

# The contribution of Brownian motion to the stress in a colloidal suspension

Duraivelan Palanisamy and Wouter K. den Otter

*Multi Scale Mechanics, Faculty of Engineering Technology and MESA+ Institute for Nanotechnology, University of Twente, Enschede, The Netherlands.*

(Dated: June 1, 2020)

The deviatoric stresses of colloidal suspensions are routinely calculated, in theoretical studies as well as in Brownian and Stokesian Dynamics simulations, using the expression introduced by Batchelor [*J. Fluid Mech.* **83**, 97–117 (1977)]. We show by example that the central feature of its derivation, the thermodynamic force representing the mean Brownian motion of colloids as flow against the density gradient, is inconsistent with the motion of the colloids on the Smoluchowski time scale. The mean Brownian motion is well-known to originate in the spatial variation of the grand mobility matrix, which therefore ought to be included in stress calculations instead. A novel expression for the stress is derived, restoring the hydrodynamic relation between the motion of suspended particles and their induced stress.

## I. INTRODUCTION

Adding rigid colloids or flexible polymers to simple fluids is well-known to affect the flow behaviour of the fluid, raising the viscosity and giving rise to visco-elastic phenomena like shear thinning; the interested reader is referred to various reviews [1–5]. Einstein [6, 7] famously derived that the viscosity of a dilute suspension of spherical colloids increases linearly with the colloidal volume fraction. Batchelor [8, 9] proposed a general expression for the deviatoric stress of non-dilute colloidal suspensions of spherical colloids subject to Brownian motion, which has become widely accepted as the standard expression in theoretical and simulation studies. It is therefore disconcerting to note that the forces entering Batchelor's stress calculation differ from those entering the equations of motion of the colloids [2, 10–12]. The problems comprise the use of a ‘thermodynamic force’ acting on the particles and the omission of a subtle correction for the configuration-dependence of the hydrodynamic matrix. A new expression for the deviatoric stress is derived by combining a number of well-known results on microhydrodynamics and Brownian motion.

Batchelor [9] derived the stress of a colloidal suspension of Brownian particles as

$$\mathbf{s}^\Sigma = -p\mathbf{1} + 2\eta_0\mathbf{E} + \frac{1}{V} \sum_i \mathbf{S}_i^E - \frac{1}{V} \sum_i (\mathbf{C}_i + \mathbf{x}_i\mathbf{1}) \cdot \mathbf{F}_i \quad (1a)$$

$$= -p\mathbf{1} + 2\eta_0\mathbf{E} + \frac{1}{V} \sum_i \mathbf{S}_i^E - \frac{k_B T}{V} \sum_i \nabla_i \cdot \mathbf{C}_i, \quad (1b)$$

where, in the first line,  $p$  is the hydrostatic pressure,  $\eta_0$  denotes the viscosity of the suspending fluid,  $\mathbf{E}$  is the imposed uniform strain rate tensor,  $V$  is the volume, the summations run over all particles,  $\mathbf{S}_i^E$  is the hydrodynamic stress on colloid  $i$  due to the strain deformation (in the absence of Brownian contributions), the product  $\mathbf{E}:\mathbf{C}_i$  yields the velocity boost of said particle in a strain deformation due to hydrodynamic interactions with the surrounding colloids,  $\mathbf{x}_i$  is the position of the

$i^{\text{th}}$  particle, and  $\mathbf{F}_i$  denotes the non-hydrodynamic force acting on the particle. Batchelor equates this force to the ‘thermodynamic force’,  $\tilde{\mathbf{F}}_i^T$ , an effective (denoted by the tilde) force describing the average result of Brownian motion as a deterministic motion against the concentration gradient; a more extensive discussion of this force is presented below. He thus arrived at the second line in the above equation, where  $k_B$  is Boltzmann's constant and  $T$  stands for the absolute temperature, as an approximate expression to the stress valid at low volume fraction only. Felderhof and Jones [13] and Nägele and Bergenholz [14] added a conservative potential-based force  $\mathbf{F}_i^\Phi = -\nabla_i\Phi$ , describing direct interactions between the colloids,  $\mathbf{F}_i = \tilde{\mathbf{F}}_i^T + \mathbf{F}_i^\Phi$ , thereby arriving at a stress expression with an inter-colloidal virial contribution,

$$\mathbf{s}^\Sigma = -p\mathbf{1} + 2\eta_\infty\mathbf{E} - \frac{k_B T}{V} \sum_i \nabla_i \cdot \mathbf{C}_i - \sum_i \mathbf{x}_i \otimes \mathbf{F}_i^\Phi - \sum_i \mathbf{C}_i \cdot \mathbf{F}_i^\Phi, \quad (2)$$

where  $\eta_\infty$  is the high frequency limiting viscosity; this expression was shown to hold true at all volume fractions. These equations and equivalent formulations are widespread in the literature, both in theoretical and numerical studies. Our objectives here are to discuss a problem underlying the inclusion of the average Brownian force in Eqs. (1b) and (2), and to provide a corrected expression for the stress.

This paper is structured as follows. Batchelor's argument for the thermodynamic force is repeated in Section II. In Section III it is shown that the thermodynamic force does not agree with the motion of colloids on the Smoluchowski time scale, and therefore ought not to be included in the derivation of a stress expression. There is a subtle Brownian contribution to the motion of colloids on the Smoluchowski time scale that must be included instead, as explained and used in Section IV to derive a novel stress expression. The new stress expression is compared against current expressions in Section V, followed by a summary of our findings in Section VI. We apolo-

gize to readers familiar with more stringent derivations of the various partial results used to refute Batchelor's argument and to derive a new stress expression: we use simple arguments to highlight the mayor points that have been systematically overlooked for the last 40 years.

## II. BACKGROUND

In a famous *Gedankenexperiment*, Einstein [6] considered the equilibrium probability distribution function  $P(\mathbf{x})$  of a dilute suspension of identical colloidal particles in an external potential  $\Phi(\mathbf{x})$ . In equilibrium, the particle flux due to the external potential is balanced by the flux due to Fickian diffusion against the concentration gradient. The macroscopic flux  $\mathbf{J}(\mathbf{x})$  then vanishes at every point  $\mathbf{x}$ , following

$$\mathbf{J} = -PM\nabla\Phi - \mathbf{D}\nabla P = \mathbf{0}, \quad (3)$$

where  $\mathbf{M}$  denotes the mobility matrix and  $\mathbf{D}$  the diffusion matrix. By inserting Boltzmann's equilibrium distribution,

$$P(\mathbf{x}) = P_0 e^{-\beta\Phi(\mathbf{x})}, \quad (4)$$

where  $P_0$  normalizes the distribution and  $\beta = 1/(k_B T)$ , Einstein showed that equilibrium implies  $\mathbf{M} = \beta\mathbf{D}$ . Using Stokes's expression for the mobility of a spherical particle of radius  $a$  in a fluid of viscosity  $\eta$ , Einstein obtained the diffusion coefficient

$$\mathbf{D} = \frac{k_B T}{6\pi\eta a} \mathbf{1}. \quad (5)$$

While the Stokes-Einstein expression was derived for colloids subject to an external force, it is equally valid for unforced colloids.

In a sequel paper, Einstein [7] likened the action of the diffusive term to a force acting on every particle, while the central equation he solved was still a flux balance. Batchelor [8] described this interpretation as follows: 'the particle flux due to Brownian migration is the same here as if a certain steady force acted on the particles (this force being equal and opposite to the external force  $\mathbf{F}^\Phi = -\nabla\Phi$  that, in the equilibrium situation, produces a convective flux which balances the diffusive flux).' The flux balance in Eq. (3) is rewritten as

$$-P \left( \mathbf{M}\nabla\Phi + \frac{1}{P}\mathbf{D}\nabla P \right) = PM \left( \mathbf{F}^\Phi + \tilde{\mathbf{F}}^T \right) = \mathbf{0}, \quad (6)$$

which, following Batchelor [8], is 'the same as if a steady force

$$\tilde{\mathbf{F}}^T(\mathbf{x}) = -k_B T \nabla \ln P(\mathbf{x}) \quad (7)$$

acted on the particle[s]. It is of course not to be supposed that the interaction of a particle with the molecules of the

surrounding medium is literally equivalent to the exertion of a steady force on the particle. When the probability density of the particle position is non-uniform, the mean Brownian velocity of a particle, conditional upon it being near a point  $\mathbf{x}$ , is non-zero simply as a consequence of the fact that the particle is more likely to have come from a direction in which the probability density increases than from one in which it decreases; and it is this bias in the statistics of particle velocities at  $\mathbf{x}$  (which is quite consistent with zero mean of the Brownian velocity of a given particle in the absence of an applied force) that is equivalent, so far as its effect on the diffusive flux is concerned, to the action of the steady force Eq. (7) on the particle.' The thermodynamic force experienced by the particles near  $\mathbf{x}$  is said to result in a thermodynamic force on colloid  $i$  given by [8, 9]

$$\tilde{\mathbf{F}}_i^T = -k_B T \frac{\partial}{\partial \mathbf{x}_i} \ln P(\mathbf{x}_1, \dots, \mathbf{x}_N), \quad (8)$$

where  $P(\mathbf{x}_1, \dots, \mathbf{x}_N)$  is the joint probability distribution function of all  $N$  particles. Batchelor [9] writes that 'these thermodynamic forces reproduce the statistical bias in the random walks of the particles which results from the non-uniformity of the joint-probability distribution function.' This use of the thermodynamic force is widespread in text books [1, 2, 5, 12, 15]. Building on the thermodynamic force, Batchelor [9] derived his widely used expression for the deviatoric stress in a suspension of spherical Brownian colloids at low strain rate, see Eq. (1b), as well as an expression for the viscosity of these suspensions up to second order in the colloidal volume fraction.

## III. THE THERMODYNAMIC FORCE REVISITED

In the reinterpretation of the flux balance as a force balance, it follows from Eq. (6) that the particles in an equilibrium suspension experience a vanishing nett force and consequently hover around an equilibrium position. But the colloids are obviously not stationary, as they are continuously subjected to rapidly fluctuating interactions with the surrounding solvent molecules in perpetual thermal motion. These fluctuating Brownian forces are not contained in the thermodynamic force, which is devoid of information on the dynamical properties of the solvent, like the viscosity, or those of the solvent molecules. Consequently, the individual colloids must also be experiencing fluctuating Brownian forces  $\mathbf{F}_i^B$ ; these forces must have a vanishing average  $\langle \mathbf{F}_i^B \rangle = \mathbf{0}$  so as not to alter the force balance of Eq. (6). As explained in the above citation of Batchelor, these Brownian forces are the origin of the diffusive flux in Eq. (3) and the thermodynamic force in Eq. (6). Randomly fluctuating Brownian forces are central to any study on the dynamics of colloids in fluids. The thermodynamic force, however, is not: the dynamics

of colloidal particles is routinely solved without reference to the thermodynamic force [2, 10–12, 16]. One obvious complication is that the probability distribution  $P$ , and hence  $\tilde{\mathbf{F}}_i^T$ , is not at hand in particle-based simulations, which require explicit expressions of the forces in terms of the particle positions – the work-around is integration by parts over configuration space to obtain a regular ensemble average, as was used in deriving Eqs. (1b) and (2). But that is not the only cause of concern.

Consider a dilute suspension of identical spherical colloids, each with the same positive excess mass  $m$  relative to the volume of fluid they displace, in a gravity field acting along the negative  $z$  direction. Using the flux  $J$  defined by the l.h.s. of Eq. (3), the evolution of the one-dimensional overall probability distribution  $P(z, t)$  follows from the conservation expression known as the Fokker-Planck-Smoluchowski equation [11, 17–19],

$$\frac{\partial P}{\partial t} = -\frac{\partial J}{\partial z} = \frac{mg}{\gamma} \frac{\partial P}{\partial z} + D \frac{\partial^2 P}{\partial z^2}, \quad (9)$$

where in the second step the effective mass  $m$ , the friction coefficient  $\gamma$ , the acceleration by gravity  $g$  and the diffusion coefficient  $D$  are assumed constant. In dilute systems, this equation applies to both the macroscopic concentration profile and the probability distribution function of an individual particle. The equilibrium solution, in the presence of a wall restricting the motion to  $z \geq 0$ , recovers the Boltzmann distribution,

$$P_{\text{eq}}(z) = P_{\text{eq}}(0)e^{-\beta mgz}, \quad (10)$$

as is readily confirmed by using Einstein’s relation  $D = k_B T / \gamma$ . We focus now on the subset of those particles that are at a specific height  $z_0$ , with  $z_0 \gg 0$ , at time 0. Their probability distribution function at a later time  $t$  is obtained as the Green’s function to the Smoluchowski equation [2, 11, 12],

$$P(z, t|z_0) = \frac{1}{\sqrt{4\pi Dt}} \exp \left[ -\frac{[z - (z_0 + v_g t)]^2}{4Dt^2} \right], \quad (11)$$

with drift velocity  $v_g = -mg/\gamma$ . This solution expresses that the colloids in an equilibrium distribution are not hovering around a constant height but have a propensity to gradually sink to the bottom of the container, where the Boltzmann probability distribution reaches its maximum. Because this subset of colloids is part of a larger system in equilibrium, it follows that the flux balance of the overall density at the macroscopic level, as in Eq. (3), does not translate into a balance of potential and thermodynamic forces at the microscopic level, as implied in the re-interpretation of the flux balance as a force balance in Eq. (6). Colloids are not subject to an effective force that drives them against the concentration gradient; instead, a concentration gradient turns the average effect of the random Brownian motion of many particles into a macroscopic flux against the concentration gradient.

The inevitable conclusion is that colloids do not experience the ‘thermodynamic force’ envisaged by Einstein and Batchelor, and expressions derived using the thermodynamic force are to be considered with care. Of particular interest here is the stress of a colloidal suspension. Batchelor [9], following his interpretation of the thermodynamic force as the average Brownian force on a particle, substituted the non-hydrodynamic force  $\mathbf{F}_i$  in Eq. (1a) with the thermodynamic force  $\tilde{\mathbf{F}}_i^T$ , see his Eq. (3.8), and after some mathematical steps arrived at Eq. (1b) as the stress expression valid for low volume fractions. Likewise, Doi and Edwards [2] emphasize in their Eq. (3.135) that the force entering their virial expression ‘must include[s] the thermodynamic force.’ This step is made by various authors – see for instance Felderhof [20], Eqs. (4.8), (5.2), (5.6), (5.7) and (7.12), Wagner [21], Eq. (6), Brady [22], Eq. (42), Strating [23], Eq. (A2), and Nägele and Bergenholtz, [14], Eq. (14) – in deriving stress expressions for colloidal suspensions. But, as observed above, *the thermodynamic force is unrelated to the motion of the colloids and should therefore not be used in deriving a stress expression* – not in the stress tensor of a given configuration, nor in an ensemble average. This is not to say that non-zero mean Brownian displacements do not contribute to the stress – Batchelor was correct to notice that they are relevant. But *the non-zero mean Brownian displacements that appear in the Langevin equation of motion, and should be used in the stress expression, result from the spatial non-homogeneity of the diffusion matrix*, rather than from the thermodynamic force. A revised expression for the stress will be derived in the next section.

For completeness, the colloids initially at  $z_0$  will eventually not sink forever. Interactions with the wall will eventually cause deviations from Eq. (11), and for long times the conditional probability converges to the equilibrium distribution,  $P(z, \infty|z_0) = P_{\text{eq}}(z)$ . This limiting behaviour is only obtained on a very long time scale, with the particles bouncing off the wall numerous times. For a macroscopic system in equilibrium, comprising many particles interacting with the wall at any moment, both the flux balance of Eq. (3) and a force balance between gravity and wall forces are obeyed nearly instantaneously. For an individual particle in this system, however, its force balance between gravity and wall forces is reached only on a very long time scale; on the far shorter Smoluchowski time scale of motion over a fraction of the colloid’s size, the persistent pull by gravity results in a downward mean motion (that is, along the concentration gradient) with super-imposed Brownian fluctuations. In summary, the thermodynamic force based on the gradient of  $k_B \ln P(\mathbf{x})$ , or on the gradient of  $\mu(\mathbf{x})/T(\mathbf{x})$  with  $\mu$  the chemical potential, provides an effective force in the phenomenological relation for the evolution of the macroscopic concentration profile [24], but this force is not to be applied to individual colloids.

#### IV. THE STRESS

We now set forth to derive an expression for the deviatoric stress in a suspension of rigid Brownian colloids in the Stokesian limit, *i.e.* for vanishing Reynolds and Stokes numbers, by combining a couple of well-known results on micro-hydrodynamics and Brownian motion. To keep the exposition focussed on the key issues, and to facilitate the comparison with earlier work [2, 9, 13, 14], we initially restrict the discussion to linear velocities only – the inclusion of angular velocities will be postponed till Section IV E. We will start by repeating a couple of well-known results, to describe the background and set the notation, before merging them into an expression for the stress.

##### A. Stokesian flow

Consider an isolated non-Brownian particle in a Newtonian fluid. The particle is described by its position  $\mathbf{x}$  and velocity  $\mathbf{U} = \dot{\mathbf{x}}$ ; the externally imposed macroscopic linear flow field is given by  $\mathbf{u}(\mathbf{r}) = \mathbf{u}_0 + \mathbf{E}\mathbf{r}$  with a constant small strain rate  $\mathbf{E}$ , *i.e.* the traceless symmetric ( $3 \times 3$ ) velocity gradient matrix. It is well-known from micro-hydrodynamics that the hydrodynamic drag force  $\mathbf{F}^H$  and stress  $\mathbf{S}^H$  experienced by the colloid are related under Stokesian flow conditions by [3, 5, 25]

$$\begin{pmatrix} \mathbf{F}^H \\ \mathbf{S}^H \end{pmatrix} = - \begin{pmatrix} \mathbf{R}_{FU} & \mathbf{R}_{FE} \\ \mathbf{R}_{SU} & \mathbf{R}_{SE} \end{pmatrix} \begin{pmatrix} \mathbf{U} - \mathbf{u}(\mathbf{x}) \\ -\mathbf{E} \end{pmatrix}, \quad (12)$$

where  $\mathbf{R}$  is the grand resistance matrix. The elements of this matrix are obtained by explicitly solving the flow and pressure fields surrounding the moving particle, followed by working out their consequences for the particle. The hydrodynamic force  $\mathbf{F}^H$  is the zeroth moment of the fluid's deviatoric stress field  $\mathbf{s}(\mathbf{r})$  (unit: N/m<sup>2</sup>) integrated over the surface of the particles, the deviatoric stress  $\mathbf{S}^H$  (unit: Nm) is obtained as the symmetric first moment of  $\mathbf{s}(\mathbf{r})$ . Analytic solutions of  $\mathbf{R}$  are available for spherical and spheroidal particles [3]; the interested reader is referred to the literature for details on numerical solvers for colloids of arbitrary shape [26–29]. Under the conditions of Stokesian flow, the total force on the particle is zero and the stress exerted by the particle on the fluid,  $\mathbf{S}$ , balances the stress by the fluid on the particle,

$$\mathbf{F}^\Phi + \mathbf{F}^H = \mathbf{0}, \quad (13a)$$

$$\mathbf{S} + \mathbf{S}^H = \mathbf{0}. \quad (13b)$$

Combining the above equations yields, by partial inversion [3, 5, 25],

$$\begin{pmatrix} \mathbf{U} - \mathbf{u} \\ \mathbf{S} \end{pmatrix} = \begin{pmatrix} \mathbf{M}_{UF} & \mathbf{M}_{UE} \\ \mathbf{M}_{SF} & \mathbf{M}_{SE} \end{pmatrix} \begin{pmatrix} \mathbf{F}^\Phi \\ \mathbf{E} \end{pmatrix}, \quad (14)$$

where the grand mobility matrix  $\mathbf{M}$  is related the grand resistance matrix by

$$\mathbf{M} = \begin{pmatrix} \mathbf{R}_{FU}^{-1} & \mathbf{R}_{FU}^{-1}\mathbf{R}_{FE} \\ \mathbf{R}_{SU}\mathbf{R}_{FU}^{-1} & \mathbf{R}_{SU}\mathbf{R}_{FU}^{-1}\mathbf{R}_{FE} - \mathbf{R}_{SE} \end{pmatrix}. \quad (15)$$

Of the two elements on the l.h.s. of Eq. (14), the velocity serves as part of an equation of motion,  $\mathbf{U} = \dot{\mathbf{x}}$ , while the stress does not.

##### B. Brownian dynamics

Consider an isolated Brownian particle in a quiescent Newtonian fluid. The evolution of its probability distribution  $P(\mathbf{x}, t)$  is described by the Smoluchowski equation

$$\frac{\partial P}{\partial t} = -\nabla \cdot \left[ \left( -\mathbf{M}_{UF}\nabla\Phi + \nabla \cdot \mathbf{D} \right) P \right] + \nabla\nabla : (\mathbf{D}P), \quad (16)$$

where, unlike in Eq. (9), it is assumed that the mobility matrix, the diffusion matrix  $\mathbf{D} = k_B T \mathbf{M}_{UF}$ , and the potential are functions of the colloidal position. In this standard form of the Smoluchowski equation, it follows from the pre-factor to  $P$  in the first term on the r.h.s., see Van Kampen [11], Eqs (IX.4.5), (IX.4.11) and (IX.4.12), and Öttinger [12], Eqs (3.78) and (3.79), that the particle experiences, in addition to the potential force  $\mathbf{F}^\Phi = -\nabla\Phi$ , an *effective* mobility-related force

$$\tilde{\mathbf{F}}^M = \mathbf{M}_{UF}^{-1} \nabla \cdot \mathbf{D} = k_B T \mathbf{M}_{UF}^{-1} \nabla \cdot \mathbf{M}_{UF}. \quad (17)$$

This term arises because a first order equation of motion is constructed to describe the dynamics resulting from a second order equation of motion including a Brownian term with a position-dependent strength. A tilde is added to emphasize that this is not a real force experienced by the particle, *i.e.* it does not enter a Newtonian equation of motion or a second order Langevin equation, but an effective force emerging in a first order Langevin equation of motion on the Smoluchowski time scale. The equation of motion in the Itô interpretation then reads as [2, 11, 12, 19, 30]

$$\dot{\mathbf{x}} = \mathbf{M}_{UF} \left( \mathbf{F}^\Phi + \tilde{\mathbf{F}}^M + \mathbf{F}^B \right), \quad (18)$$

where the Brownian force, with  $\langle \mathbf{F}^B \rangle = \mathbf{0}$ , obeys the fluctuation-dissipation theorem

$$\langle \mathbf{F}^B(t) \otimes \mathbf{F}^B(t') \rangle = 2k_B T \mathbf{R}_{FU} \delta(t - t'). \quad (19)$$

The Itô interpretation implies that all quantities appearing on the r.h.s. of Eq. (18) are evaluated using the positions  $\mathbf{x}$  before their incremental change due to the velocity  $\dot{\mathbf{x}}$ . From a physical point of view, this Langevin equation only holds true on the Smoluchowski time scale: it describes the motion on a time scale that far exceeds

the relaxation time of the velocity autocorrelation of the colloid, thereby eliminating inertia effects in the force balance, but is still short compared to motion over the colloid's size. The random force entering the dynamics then no longer consists of an infinite series of uncorrelated (Markovian) delta peaks, resulting in discontinuous jumps in the velocity, but of a well-defined time-integral over these peaks [12]. Note that merely removing the inertial term from the Newtonian equation of motion is well-known not to yield the correct Itô-form of the first-order Langevin equation of motion – inclusion of the effective force  $\tilde{\mathbf{F}}^M$ , accounting for a bias incurred by Brownian motion with a spatially varying mobility matrix, is crucial to recovering both the correct dynamics and the equilibrium Boltzmann distribution [2, 30].

The displacements of the particles over a simulation time step  $\Delta t$  are usually approximated by the forward Euler scheme [see 12], *i.e.* integrating the r.h.s. of Eq. (18) from  $t$  to  $t + \Delta t$  while keeping the coordinates fixed at their values at time  $t$ ,

$$\mathbf{x}(t + \Delta t) - \mathbf{x}(t) = \mathbf{M}_{UF} [\mathbf{F}^\Phi + \bar{\mathbf{F}}^B(t)] \Delta t + k_B T \nabla \cdot \mathbf{M}_{UF} \Delta t, \quad (20)$$

where the step-averaged Brownian force  $\bar{\mathbf{F}}^B(t) = (\Delta t)^{-1} \int_t^{t+\Delta t} \mathbf{F}^B(\tau) d\tau$  obeys the fluctuation-dissipation theorem

$$\langle \bar{\mathbf{F}}^B(t) \otimes \bar{\mathbf{F}}^B(t') \rangle = 2k_B T \Delta t \mathbf{R}_{FU} \delta_{t,t'}, \quad (21)$$

where the Kronecker delta  $\delta_{t,t'}$  equals one if  $t$  and  $t'$  refer to the same step and zero if  $t$  and  $t'$  refer to distinct steps. In practice, the displacement due to the Brownian force is readily calculated as

$$\Delta \mathbf{x}^B(t) = \mathbf{M}_{UF} \bar{\mathbf{F}}^B(t) \Delta t = \sqrt{2k_B T \mathbf{M}_{UF}^{1/2}} \boldsymbol{\theta}(t) \sqrt{\Delta t}, \quad (22)$$

where the vector  $\boldsymbol{\theta}(t)$  contains three random numbers of zero mean, unit variance and devoid of correlations; again, the forces and matrices entering these equations are evaluated at time  $t$ , before the position update. As will be discussed in more detail below, there are two Brownian-related terms affecting the displacement in Eq. (20) and hence the step-averaged velocity of the colloid; it then follows from Eq. (12) that both terms contribute to the stress. Inclusion of a slow (relative to the Smoluchowski time scale) shear flow is achieved by

$$\dot{\mathbf{x}} = \mathbf{M}_{UF} (\mathbf{F}^\Phi + \tilde{\mathbf{F}}^M + \mathbf{F}^B) + \mathbf{M}_{UE} \mathbf{E} + \mathbf{u}, \quad (23)$$

and integration from  $t$  to  $t + \Delta t$  extends Eq. (20) with the flow-related displacement terms  $-\mathbf{M}_{UE} \mathbf{E} \Delta t + \mathbf{u} \Delta t$ ; the discretized equation of motion to first order in  $\Delta t$  does not contain coupling between diffusion and strain [31, 32]. The order of strong convergence of this Euler scheme is 1/2; Mil'shtein [12, 33] method is required to reach an order of one. The above results are all well-known – they

form the starting point of theoretical developments and Brownian Dynamics simulations exploring colloidal dynamics beyond the Smoluchowski time scale. Ermak and McCammon [10] derived the above integration scheme starting from the second order Langevin equation of motion of a colloid. The implementation of this scheme for a collection of hydrodynamically interacting colloids is known as Stokesian Dynamics [16].

### C. The stress

The results of the previous two sections are now combined to obtain the deviatoric stress on an isolated colloid. Since we are dealing with a system in Stokesian flow, the hydrodynamic force and stress on the colloid are obtained by Eq. (12). The hydrodynamic stress is not part of an equation of motion, there is no related Fokker-Planck equation, nor do the hydrodynamic matrices vary with the stress. One may therefore apply Eq. (12), still in the Itô representation and keeping in mind that the equation is physically valid on the Smoluchowski time scale. Inserting the velocity derived in Eq. (23), and using the matrix relations from Eq. (15), gives

$$\mathbf{F}^H = - \left( \mathbf{F}^\Phi + \tilde{\mathbf{F}}^M + \mathbf{F}^B \right), \quad (24)$$

thereby recovering the expected force balance of a Brownian colloid in Stokesian flow on the Smoluchowski time scale. The effective mobility-related force does not feature in a second order Langevin equation of motion of a colloid, but emerges in the first order Langevin equation: it accounts for the average Brownian force on the Smoluchowski time scale being non-zero in the presence of a non-constant mobility matrix [2, 10–12, 30]. Spatial variations of the mobility matrix affect the Brownian displacements of the colloid, thereby giving rise to an effective force on the Smoluchowski time scale; the additional displacement, *i.e.* the last term in Eq. (20), contributes to the velocity of the colloid and thereby to the stress induced by the colloid.

Returning to Eq. (12) and again inserting the velocity derived in Eq. (23), one readily obtains the deviatoric hydrodynamic stress by the particle on the fluid as

$$\mathbf{S}^H = \mathbf{S}^\Phi + \tilde{\mathbf{S}}^M + \mathbf{S}^B + \mathbf{S}^E. \quad (25)$$

Using the matrix relations from Eq. (15), one finds that the stress consists of a potential term

$$\mathbf{S}^\Phi = -\mathbf{R}_{SU} \mathbf{M}_{UF} \mathbf{F}^\Phi = -\mathbf{M}_{SF} \mathbf{F}^\Phi \quad (26)$$

and a strain term

$$\mathbf{S}^E = -(\mathbf{R}_{SU} \mathbf{M}_{UE} - \mathbf{R}_{SE}) \mathbf{E} = -\mathbf{M}_{SE} \mathbf{E}, \quad (27)$$

both of which already featured in Eq. (14), as well as two

Brownian-related contributions: a fluctuating term

$$\mathbf{S}^B = -\mathbf{R}_{SU}\mathbf{M}_{UF}\mathbf{F}^B = -\mathbf{M}_{SF}\mathbf{F}^B \quad (28)$$

and a systematic term

$$\begin{aligned} \tilde{\mathbf{S}}^M &= -\mathbf{R}_{SU}\mathbf{M}_{UF}\tilde{\mathbf{F}}^M = -\mathbf{M}_{SF}\tilde{\mathbf{F}}^M \\ &= -k_B T \mathbf{R}_{SU} \nabla \cdot \mathbf{M}_{UF}. \end{aligned} \quad (29)$$

The latter two terms arise because *both the step-averaged Brownian force  $\mathbf{F}^B$ , with zero mean, and the mobility-related effective force  $\tilde{\mathbf{F}}^M$ , accounting for the non-zero mean Brownian displacement induced by spatial variations of the mobility matrix, contribute to the velocity and displacement of the colloid on the Smoluchowski time scale, see Eqs. (18) and (20), while the stress is linear in this velocity under Stokesian flow conditions.*

Combining the force balance of Eq. (24) with the stress balance of Eq. (13b), and repeating the partial inversion of Eq. (14), one arrives at

$$\begin{pmatrix} \mathbf{U} - \mathbf{u} \\ \mathbf{S} \end{pmatrix} = \begin{pmatrix} \mathbf{M}_{UF} & \mathbf{M}_{UE} \\ \mathbf{M}_{SF} & \mathbf{M}_{SE} \end{pmatrix} \begin{pmatrix} \mathbf{F}^\Phi + \tilde{\mathbf{F}}^M + \mathbf{F}^B \\ \mathbf{E} \end{pmatrix}, \quad (30)$$

again in the Itô interpretation and on the Smoluchowski time scale. Evaluating this expression recovers both the above equation of motion and the four stress contributions. For simulation purposes, forward Euler integration of Eq. (30) gives

$$\mathbf{x}(t + \Delta t) = \mathbf{x}(t) + \bar{\mathbf{U}}(\mathbf{x}(t), t)\Delta t, \quad (31a)$$

$$\bar{\mathbf{U}}(\mathbf{x}, t) = \mathbf{M}_{UF} \left[ \mathbf{F}^\Phi + \tilde{\mathbf{F}}^M + \bar{\mathbf{F}}^B(t) \right] + \mathbf{M}_{UE}\mathbf{E} + \mathbf{u}, \quad (31b)$$

$$\bar{\mathbf{S}}(\mathbf{x}, t) = \mathbf{M}_{SF} \left[ \mathbf{F}^\Phi + \tilde{\mathbf{F}}^M + \bar{\mathbf{F}}^B(t) \right] + \mathbf{M}_{SE}\mathbf{E}, \quad (31c)$$

where the step-averaged velocity  $\bar{\mathbf{U}}$  and stress  $\bar{\mathbf{S}}$  are determined to the same order in the time step  $\Delta t$ . This recovers the usual equation of motion, with a revised stress.

#### D. Fluctuating contributions to the stress

The perpetual Brownian motion of colloids affects the stress both directly and indirectly. The direct contributions are represented by the combination of  $\mathbf{S}^B$ , the hydrodynamic stress due to colloidal motions induced by the fluctuating Brownian force, and  $\tilde{\mathbf{S}}^M$ , accounting for a systematic bias in the Brownian force whenever the mobility tensor is non-uniform, *e.g.* due to hydrodynamic interactions between colloids. The indirect contribution results from the combination of Brownian motion with potential forces and imposed strain rate, collectively determining the time-evolving distribution of the colloids. Besides these two well-known contributions, the

multitude of interactions of the colloid with the solvent molecules in perpetual thermal motions gives rise to two additional stress contributions. The long-time average of the colloid-solvent interaction yields the hydrostatic stress  $-p\mathbf{1}$ , with  $p$  the hydrostatic pressure. Denoting the difference between the short-time and long-time averages as the ‘fluctuating Brownian stresslet,’  $\mathbf{S}^B$ , one finds that  $\langle \mathbf{S}^B \rangle = 0$ . Since the fluctuating Brownian force and the fluctuating Brownian stresslet are distinct projections of the same interactions of the colloid with the solvent, namely the zeroth moment and the symmetric first moment of the fluctuating stress field over the colloids surface, respectively, they are related by a generalized fluctuation-dissipation theorem [34],

$$\begin{aligned} &\left\langle \begin{pmatrix} \mathbf{F}^B(t) \\ \mathbf{S}^B(t) \end{pmatrix} \otimes \begin{pmatrix} \mathbf{F}^B(t') \\ \mathbf{S}^B(t') \end{pmatrix} \right\rangle \\ &= 2k_B T \begin{pmatrix} \mathbf{R}_{FU} & \mathbf{R}_{FE} \\ \mathbf{R}_{SU} & \mathbf{R}_{SE} \end{pmatrix} \delta(t - t'). \end{aligned} \quad (32)$$

This coupling does not alter the equation of motion, as is readily verified by noting that the usual fluctuation-dissipation theorem in Eq. (19) is a subset of the above expression, while the total deviatoric stress  $\mathbf{S}^H$  in Eq. (25) acquires the fluctuating stresslet  $\mathbf{S}^B$ .

#### E. Angular velocities

Linear flow fields may include a constant rotational component,  $\mathbf{u}(\mathbf{r}) = \mathbf{u}_0 + \boldsymbol{\omega} \times \mathbf{r} + \mathbf{E}\mathbf{r}$ , and a colloid in a flow field may acquire an angular velocity  $\boldsymbol{\Omega}$ . The corresponding extension of Eq. (30) retains the same concepts, while introducing complications that we hitherto avoided for clarity of presentation. For spherical particles, it suffices to re-interpretate the velocity vectors  $\mathbf{U}$  and  $\mathbf{u}$  as six-vectors combining the linear and angular velocities of the colloid and flow field, respectively, to re-interpret each of the forces  $\mathbf{F}^\Phi$ ,  $\tilde{\mathbf{F}}^M$  and  $\mathbf{F}^B$  as six-vectors combining a force and a torque, and to extend the grand mobility and grand resistance matrices accordingly. With these steps, the expressions for the motion and stress in Eq. (30) and the fluctuation-dissipation theorem of Eq. (32) hold true again. Only the translational equation of motion needs to be solved to explore the evolution of the system in time. If the particle is non-spherical, however, the rotational motion has to be solved as well. The complication here is that the angular velocity  $\boldsymbol{\Omega}$  is not the time derivate of a coordinate vector. One may use Euler angles or a Cartesian rotation vector to derive the corresponding mobility matrix and its divergence [35], but care must be taken to avoid the singular points of the resulting equations of motion. Furthermore, the orientation-dependence of the volume of momentum space gives rise to an additional contribution to the torque. These issues are elegantly solved by using quaternions, *i.e.* a set of four coordinates coupled by a unit-length constraint, which results in a remarkably simple equation of motion [29, 36].

## F. Multiple colloids

The above derivation is readily extended to a collection of  $N$  particles. Upon re-interpreting  $\mathbf{x}$ ,  $\mathbf{F}^B$ , etcetera, as vectors comprising all particle coordinates, all Brownian forces, etcetera, the above equations remain unaltered. One then obtains for the  $i^{\text{th}}$  particle,

$$\begin{pmatrix} \mathbf{U}_i \\ \mathbf{S}_i \end{pmatrix} = \sum_{j=1}^N \begin{pmatrix} \mathbf{M}_{UF}^{ij} & \mathbf{M}_{UE}^{ij} \\ \mathbf{M}_{SF}^{ij} & \mathbf{M}_{SE}^{ij} \end{pmatrix} \begin{pmatrix} \mathbf{F}_j^\Phi + \tilde{\mathbf{F}}_j^M + \mathbf{F}_j^B \\ \mathbf{E} \end{pmatrix} + \begin{pmatrix} \mathbf{u}(\mathbf{x}_i) \\ -\mathbf{S}_i^B \end{pmatrix}, \quad (33)$$

with fluctuation-dissipation theorem

$$\begin{aligned} & \left\langle \left( \begin{pmatrix} \mathbf{F}_i^B(t) \\ \mathbf{S}_i^B(t) \end{pmatrix} \otimes \begin{pmatrix} \mathbf{F}_j^B(t') \\ \mathbf{S}_j^B(t') \end{pmatrix} \right) \right\rangle \\ & = 2k_B T \begin{pmatrix} \mathbf{R}_{FU}^{ij} & \mathbf{R}_{FE}^{ij} \\ \mathbf{R}_{SU}^{ij} & \mathbf{R}_{SE}^{ij} \end{pmatrix} \delta(t - t'), \end{aligned} \quad (34)$$

and so on, where it should be noted that the many-particle matrices  $\mathbf{M}$  and  $\mathbf{R}$  are related by Eq. (15), whereas the two-particle matrices  $\mathbf{M}^{ij}$  and  $\mathbf{R}^{ij}$  are not. In this extension, the hydrodynamic matrices account for hydrodynamic interactions between the particles, *i.e.* a force acting on particle  $j$  contributes to the velocity and stress of particle  $i$ , and vice versa. With the potential limited to inter-particle interactions, *i.e.* in the absence of external interactions, the overall deviatoric stress exerted on the fluid,  $\mathbf{s}^\Sigma$  (unit: N/m<sup>2</sup>), at a strain rate  $\mathbf{E}$  is obtained as

$$\mathbf{s}^\Sigma = 2\eta_0 \mathbf{E} - \frac{1}{V} \sum_i \mathbf{S}_i - \frac{1}{V} \hat{\mathcal{P}} \sum_{i < j} \mathbf{x}_{ij} \otimes \mathbf{F}_{ij}^\Phi, \quad (35)$$

where the first term on the r.h.s. is the stress in the suspending fluid with viscosity  $\eta_0$ , the second term accounts for hydrodynamic interactions between the particles and the fluid, including fluid-mediated interactions between the particles, and the third term is the regular virial expression arising from direct inter-particle interactions [2, 9], where the projection

$$\hat{\mathcal{P}}\mathbf{X} = (\mathbf{X} + \mathbf{X}^T)/2 - \det(\mathbf{X})\mathbf{1} \quad (36)$$

returns the symmetric traceless part of a matrix  $\mathbf{X}$ . The total stress tensor of the suspension is obtained by adding the hydrostatic pressure  $-p\mathbf{1}$  to Eq. (35). The contribution of the colloids to this total stress, also known as the osmotic stress, is obtained from Eq. (35) by removing the bulk term  $2\eta_0 \mathbf{E}$  and the projection  $\hat{\mathcal{P}}$ , and adding the kinetic contribution  $-Nk_B T \mathbf{1}/V$ .

## V. COMPARISON OF STRESS EXPRESSIONS

Comparing the novel expression for the stress with earlier expressions, by Batchelor [9], Eqs. (2.2) and (3.10), Felderhof [20], Eq. (7.17), Brady [22], Eqs (38) through (40), and Nagele and Bergenholtz [14], Eq. (31), reveals a number of similarities and differences. The contributions due to the strain are identical, where the stress  $\mathbf{S}_i^E$  in Eq. (1b) is understood to include hydrodynamic interactions between the colloids,  $\mathbf{S}_i^E = -\sum_j \mathbf{M}_{SE}^{ij} \mathbf{E}$ . From the velocity relation under pure strain [9, 14],

$$\mathbf{U}_i = \mathbf{u}(\mathbf{x}_i) + \mathbf{E} : \mathbf{C}_i = \mathbf{u}(\mathbf{x}_i) + \sum_j \mathbf{M}_{UE}^{ij} : \mathbf{E}, \quad (37)$$

follows  $\mathbf{C}_i = \sum_j \mathbf{M}_{SF}^{ji}$ , where a symmetry rule of the grand mobility matrix [26] was used in the last step. The potential-induced hydrodynamic term in Eq. (2) then reads as  $\mathbf{S}^\Phi = -\sum_i \mathbf{C}_i \cdot \mathbf{F}_i^\Phi = -\sum_{ij} \mathbf{M}_{SF}^{ji} \mathbf{F}_i^\Phi$ , in agreement with the corresponding term in Eq. (33).

The differences are in the stress contributions by the Brownian forces. Previous derivations of stress expressions are based on the assumption that the mean contribution of the Brownian force is provided by the thermodynamic force  $\tilde{\mathbf{F}}^T$ , giving rise to a thermodynamic stress term  $\tilde{\mathbf{S}}^T$ . Here, instead, the mean contribution of the Brownian force is equated to the mobility-related effective force  $\tilde{\mathbf{F}}^M$ , see Eq. (17). These two effective forces are fundamentally different, being based on the probability distribution and the grand mobility matrix, respectively, and consequently the corresponding stresses have little in common. After rewriting the thermodynamic stress term to eliminate the probability distribution, as in Eqs. (1b) and (2), both stress terms acquire superficial similarities as divergences of segments of the hydrodynamic matrices,

$$\tilde{\mathbf{S}}^T = -\frac{k_B T}{V} \sum_i \nabla_i \cdot \mathbf{C}_i = -\frac{k_B T}{V} \sum_{ij} \nabla_i \cdot \mathbf{M}_{SF}^{ij} \quad (38)$$

and

$$\tilde{\mathbf{S}}^M = -\frac{1}{V} \sum_{ij} \mathbf{M}_{SF}^{ij} \tilde{\mathbf{F}}_j^M = -\frac{k_B T}{V} \sum_{ijk} \mathbf{R}_{SU}^{ik} \nabla_j \cdot \mathbf{M}_{UF}^{kj}, \quad (39)$$

respectively. Since  $\mathbf{M}_{SF} = \mathbf{R}_{SU} \mathbf{M}_{UF}$ , as follows from Eq. (15), the two stresses are different in general. A rare exception is a dispersion consisting of a single sphere, in which case  $\mathbf{M}_{SF} = \mathbf{R}_{SU} = \mathbf{0}$ .

A second difference is the explicit inclusion of all fluctuating Brownian contributions in Eq. (33). Their presence allows for a self-consistency test by comparing the viscosity obtained from the average stress at constant low shear rate with the viscosity extracted from the thermal stress fluctuations in equilibrium using the Green-Kubo formalism. It is not possible to conclude that a stress expression passes this test based on an analysis that bypasses the

fluctuating Brownian stress contributions and their correlations to the colloidal dynamics [14, 34]. The consistency test of the revised stress expression is a topic of ongoing research. Note that identical forces enter both the equation of motion and the novel stress expression, because the non-straining part of the hydrodynamic stress is a consequence of the motion of the colloids relative to the flow field, see Eq. (12). In applications of Batchelor’s approach, besides the omitted fluctuating Brownian contributions to the stress, the effective mobility-related force  $\tilde{\mathbf{F}}^M$  is used in the equation of motion but not in the stress, while the effective thermodynamic force  $\tilde{\mathbf{F}}^T$  features in the stress,

$$\tilde{\mathbf{S}}^T = -\frac{1}{V} \sum_{ij} \mathbf{M}_{SF}^{ij} \tilde{\mathbf{F}}_j^T, \quad (40)$$

but does not appear in the equation of motion.

## VI. DISCUSSION AND CONCLUSIONS

The ‘thermodynamic force’  $\tilde{\mathbf{F}}^T$  was presented by Batchelor [8] as ‘an alternative and much simpler method for the statistical mechanics part of the investigation which is a generalization of the argument used by Einstein and which gives the asymptotic or long-time statistical properties of the displacement of particles in terms of the thermal energy of the medium.’ Batchelor [9], and many authors since, have used the thermodynamic force as the average resultant of the Brownian force, driving the colloids against the concentration gradient, in the derivation of stress expressions for colloidal suspensions. But the average resultant of the Brownian force is well-known to derive from the divergence of the mobility matrix, referred to above as the mobility-related force  $\tilde{\mathbf{F}}^M$ . A new stress expression was derived, see Eq. (25) and Eqs. (33) through (35), in which the Brownian forces entering the stress calculation match those entering the

equation of motion of the colloids.

The interpretation of the thermodynamic force as the average contribution of Brownian motion is widespread in the literature on the stress of colloidal suspensions. Bossis and Brady [37] are a rare exception, by presenting an alternative derivation of a stress expression without making use of the thermodynamic force and including the mobility-related force instead. They replace Eq. (18) by an alternative first order Langevin equation of motion that by an averaged Mil’shtein approximation recovers Eq. (20) upon integration; the stress then follows by integration of  $\mathbf{R}_{SU}\dot{\mathbf{x}}$  over the time step, again by averaging the Mil’shtein approximation over the Brownian forces. This derivation is hampered, however, by building on the assumption that Eq. (20) is the correct expression and Eq. (18) an approximation – it is well-known that Eq. (18) is the correct expression in the Itô interpretation while Eq. (20) is an approximation to its integration over a time step [10–12].

More work is needed to establish the impact of the revised averaged Brownian contribution on the stress and viscosity calculations of the past 40 years. The good agreement between simulation results and experimental data on the viscosity of suspensions of spherical particles, see for instance the variation of viscosity with volume fraction reported by Foss and Brady [38], indicates that the mean Brownian term makes a relatively minor contribution in this particular case. A number of simulations and derivations should be repeated carefully to establish the particular consequences for other systems. Numerical results illustrating the impact on simple colloidal systems will be presented in a forthcoming publication.

## ACKNOWLEDGMENTS

This work is part of the Computational Sciences for Energy Research Industrial Partnership Programme co-financed by Shell Global Solutions B.V. and the Netherlands Organisation for Scientific Research (NWO).

- 
- [1] R. B. Bird, C. F. Curtiss, R. C. Armstrong, and O. Hassager, *Dynamics of Polymeric Liquids. Vol. 2: Kinetic Theory* (John Wiley & Sons, New York, NY, U.S.A., 1987).
  - [2] M. Doi and S. F. Edwards, *The Theory of Polymer Dynamics*, International Series of Monographs on Physics, Vol. 73 (Clarendon Press, Oxford, U.K., 1988).
  - [3] S. Kim and S. J. Karrila, *Microhydrodynamics: Principles and Selected Applications*, Butterworth-Heinemann Series in Chemical Engineering (Butterworth-Heinemann, Stoneham, MA, U.S.A., 1991).
  - [4] R. G. Larson, *The Structure and Rheology of Complex Fluids* (Oxford University Press, New York, NY, U. S. A., 1999).
  - [5] E. Guazzelli and J. F. Morris, *A Physical Introduction to Suspension Dynamics*, Cambridge Texts in Applied Mathematics (Cambridge University Press, Cambridge, U.K., 2012).
  - [6] A. Einstein, Ann. Phys. **17**, 549 (1905).
  - [7] A. Einstein, Ann. Phys. **19**, 289 (1906).
  - [8] G. K. Batchelor, J. Fluid Mech. **74**, 1 (1976).
  - [9] G. K. Batchelor, J. Fluid Mech. **83**, 97 (1977).
  - [10] D. L. Ermak and J. A. McCammon, J. Chem. Phys. **69**, 1352 (1978).
  - [11] N. G. van Kampen, *Stochastic Processes in Physics and Chemistry* (revised and enlarged edition, Elsevier, Amsterdam, The Netherlands, 1992).
  - [12] H. C. Öttinger, *Stochastic Processes in Polymeric Fluids* (Springer-Verlag, Berlin, Germany, 1996).
  - [13] B. U. Felderhof and R. B. Jones, Physica A **146**, 417 (1987).

- [14] G. Nägele and J. Bergenholtz, *J. Chem. Phys.* **108**, 9893 (1998).
- [15] J. K. G. Dhont, *An Introduction to Dynamics of Colloids*, Studies in Interface Science (Elsevier, Amsterdam, The Netherlands, 1996).
- [16] J. F. Brady and G. Bossis, *Ann. Rev. Fluid Mech.* **20**, 111 (1988).
- [17] D. A. McQuarrie, *Statistical Mechanics* (Harper & Row Publishers, New York, NY, U.S.A., 1976).
- [18] H. Risken, *The Fokker-Planck Equation. Methods of Solution and Applications*, 2nd ed., Springer Series in Synergetics, Vol. 18 (Springer Verlag, Berlin, Germany, 1989).
- [19] C. Gardiner, *Stochastic Methods. A handbook for the Natural and Social Sciences*, 4th ed., Springer Series in Synergetics (Springer-Verlag, Berlin, Germany, 2009).
- [20] B. U. Felderhof, *Physica A* **147**, 203 (1987).
- [21] N. J. Wagner, *J. Coll. Interf. Sci.* **161**, 169 (1993).
- [22] J. F. Brady, *J. Chem. Phys.* **98**, 3335 (1993).
- [23] P. Strating, *J. Chem. Phys.* **103**, 10226 (1995).
- [24] S. R. de Groot and P. Mazur, *Non-equilibrium Thermodynamics* (Dover Publications, New York, NY, U. S. A., 1984).
- [25] L. Durlofsky, J. F. Brady, and G. Bossis, *J. Fluid Mech.* **180**, 21 (1987).
- [26] M. Makino and M. Doi, *J. Phys. Soc. Japan* **73**, 2739 (2004).
- [27] J. García de la Torre, G. G. del Rio Echenique, and A. Ortega, *J. Phys. Chem. B* **111**, 955 (2007).
- [28] S. Aragon, *Methods* **54**, 101 (2011).
- [29] D. Palanisamy and W. K. den Otter, *J. Chem. Phys.* **148**, 194112 (2018).
- [30] W. J. Briels, (1998), *theory of Polymer Dynamics*, Lecture notes, University of Twente, Enschede, The Netherlands.
- [31] P. J. Dotson, *J. Chem. Phys.* **79**, 5730 (1983).
- [32] D. M. Heyes, *J. Non-Newtonian Fluid Mech.* **27**, 47 (1988).
- [33] G. N. Mil'shtein, *Theo. Probabl. Appl.* **19**, 557 (1974).
- [34] D. Palanisamy and W. K. den Otter, *J. Chem. Phys.* **152**, 074901 (2020).
- [35] T. R. Evensen, S. N. Naess, and A. Elgsaeter, *Macromol. Theory Simul.* **17**, 403 (2008).
- [36] I. M. Ilie, W. J. Briels, and W. K. den Otter, *J. Chem. Phys.* **142**, 114103 (2015).
- [37] G. Bossis and J. F. Brady, *J. Chem. Phys.* **91**, 1866 (1989).
- [38] D. R. Foss and J. F. Brady, *J. Fluid Mech.* **407**, 167 (2000).