

On information gain, Kullback-Leibler divergence, entropy production and the involution kernel

A. O. Lopes and J. K. Mengue

March 5, 2020

Abstract

It is well known that in Information Theory and in Machine Learning the Kullback-Leibler divergence, which extends the concept of Shannon entropy, plays a fundamental role. Given an *a priori* probability ν on the measurable space X and a probability π on the measurable space $X \times Y$ we are able to define the entropy of π relative to ν . Using this entropy we obtain a natural definition of information gain for general measurable spaces which coincides with the mutual information given from the K-L divergence. There is a natural way of extending the concept of information gain by considering the entropy relative to transverse functions. This extends the meaning of specific information gain of Ergodic Theory. This will be also used to extend the meaning of specific information gain for the XY model. Finally, we extend the concept of dynamical entropy production for the XY model. In this case we notice that the involution kernel is a natural tool to better understand entropy production. If the potential is symmetric the associated equilibrium probability has zero entropy production. The introduction of the *a priori* probability (indispensable here) and transverse functions is a novelty in the theory.

1 Introduction

In Information Theory and Data Compression the Shannon entropy of a probability vector $P = (p_1, \dots, p_d)$ plays an important role (see [26] and [8] chap. 5). The number¹ $S(P) = -\sum_{i=1}^d p_i \log(p_i)$, where $0 \log(0) = 0$, by

¹we will consider here $\log(x) = \ln(x)$, but any basis could be used.

convention, can be interpreted (taking basis 2 for the logarithm) as a lower bound for the average of questions of type “yes or no” which are necessary in order to analyze the statistics of a symbol picked at random - according to the probability distribution $P = (p_1, \dots, p_d)$ - on the finite alphabet $\{1, \dots, d\}$. From the sequence of answers to successive questions - of a certain type - one can introduce a binary code on the set $\{1, \dots, d\}$, where 0 corresponds to “yes” and 1 to “no” (see [8] chap. 5). We illustrate this claim through some examples in the appendix section.

In Information Theory, Machine Learning and Decision Trees it is also considered another important concept called **information gain**. Following [24] (see p. 89-90), for a probability π on $X \times Y = \{1, \dots, d\} \times \{1, \dots, r\}$ with x -marginal $P = (p_1, \dots, p_d)$, we define the information gain of π with respect to P as

$$IG(\pi, P) = \underbrace{-\sum_{x=1}^d p_x \log(p_x)}_{S(P)} - \underbrace{\sum_{y=1}^r q_y \left[-\sum_{x=1}^d \frac{\pi_{x,y}}{q_y} \log\left(\frac{\pi_{x,y}}{q_y}\right) \right]}_{H(\pi)}, \quad (1)$$

where $q_y = \sum_x \pi_{x,y}$, that is, $Q = (q_1, \dots, q_r)$ is the y -marginal of π . In this expression the number

$$-\left[\sum_{x=1}^d \frac{\pi_{x,y}}{q_y} \log\left(\frac{\pi_{x,y}}{q_y}\right) \right]$$

is the Shannon entropy of the probability obtained from the distribution of π on the line $X \times \{y\}$ and, therefore, $H(\pi)$ is just the weighted mean of these entropies according to Q . There is no explicit underline dynamics associated to the above concept. Example 52 in our appendix section illustrate a concrete interpretation of $IG(\pi, P)$.

As P and Q are the marginals of π then the number $IG(\pi, P)$ can be rewritten as

$$IG(\pi, P) = I(\pi) := \sum_{x=1}^d \sum_{y=1}^r \pi_{x,y} \log\left(\frac{\pi_{x,y}}{p_x q_y}\right),$$

which is also called of **mutual information**² of π . This last expression can be extended for measurable spaces by using the Kullback-Leibler divergence

²In the present work we avoid using explicitly the concept of random variable and we consider in our notation only its distribution.

(see [14]) of π with respect to $P \times Q$, that is, if π is absolutely continuous with respect to $P \times Q$, then, denoting by $\frac{d\pi}{dP dQ}$ the Radon-Nikodym derivative, we have

$$I(\pi) := D_{KL}(\pi | P \times Q) = \int \log \left(\frac{d\pi}{dP dQ}(x, y) \right) d\pi(x, y). \quad (2)$$

We call $J^\pi(x, y) = \frac{\pi_{x,y}}{\sum_x \pi_{x,y}}$ the **Jacobian** of the probability π (which is defined π -a.e.).

Then, we have

$$H(\pi) = - \sum_{x=1}^d \sum_{y=1}^r \pi_{x,y} \log(J^\pi(x, y)) = - \int \log(J^\pi) d\pi.$$

It is also possible to show (see for example [21], chap. 3) that

$$H(\pi) = - \sup \left\{ \sum_{x,y} f(x, y) \pi_{x,y} \mid \sum_{x \in X} e^{f(x,y)} = 1, \forall y \right\}. \quad (3)$$

For any given probability P on $X = \{1, \dots, d\}$ and any given probability $\tilde{Q} = (\tilde{q}_1, \dots, \tilde{q}_r)$ on $Y = \{1, \dots, r\}$, with $\tilde{q}_i > 0, \forall i$, consider the product measure $\pi_0 = P \times \tilde{Q}$ on $X \times Y$. Then,

1. $J^{\pi_0}(x, y) = \frac{p_x \tilde{q}_y}{\sum_x p_x \tilde{q}_y} = \frac{p_x \tilde{q}_y}{\tilde{q}_y} = p_x$,
2. $S(P) = - \sum_{x,y} (\pi_0)_{x,y} \log(p_x) = - \sum_{x,y} (\pi_0)_{x,y} \log(J^{\pi_0}(x, y)) = H(\pi_0)$,
3. If π is any probability on $X \times Y$ with x -marginal P , then we have

$$IG(\pi, P) = H(\pi_0) - H(\pi) = - \left[\int \log(J^{\pi_0}) d\pi - H(\pi) \right].$$

This allows us to extend the definition of information gain in the following way: Let π_0, π be probabilities on $X \times Y$, such that $(\pi_0)_{x,y} > 0, \forall (x, y) \in X \times Y$. We define the information gain of π with respect to π_0 as

$$IG(\pi, \pi_0) = - \left[\int \log(J^{\pi_0}) d\pi + H(\pi) \right]. \quad (4)$$

This work lies in the frontier of Information Theory and Ergodic Theory.

From a different point of view, in Ergodic Theory and Symbolic Dynamics it is usually considered the space

$$\Omega = \Omega^+ = \{1, 2, \dots, d\}^{\mathbb{N}} = \{|x_1, x_2, x_3, \dots\} \mid x_i \in \{1, 2, \dots, d\}, \forall i \in \mathbb{N}\} \quad (5)$$

and also the Kolmogorov-Sinai entropy of probabilities μ on Ω , which are invariant for the shift map $\sigma : \Omega \rightarrow \Omega$, $\sigma(|x_1, x_2, x_3, \dots\rangle) = |x_2, x_3, x_4, \dots\rangle$.

The set Ω is a compact metric space, when equipped with the metric

$$d(|x_1, x_2, x_3, \dots\rangle, |y_1, y_2, y_3, \dots\rangle) = 2^{-n},$$

where $n = \min\{i \mid x_i \neq y_i\}$, if $x \neq y$. It is a measurable space with the associated Borel σ -algebra \mathcal{B} .

For any $n \geq 1$, and any fixed symbols b_1, \dots, b_n in $\{1, 2, \dots, d\}$, we define the cylinder set $|b_1, b_2, \dots, b_n\rangle = \{|x_1, x_2, x_3, \dots\rangle \in X \mid x_1 = b_1, \dots, x_n = b_n\}$. A Borel probability μ on Ω is called shift-invariant if it satisfies $\mu(|b_1, b_2, \dots, b_n\rangle) = \sum_{i=1}^d \mu(|i, b_1, \dots, b_n\rangle)$ for any cylinder set.

The Kolmogorov-Sinai entropy of a shift-invariant Borel probability μ is given by

$$h(\mu) = \lim_{n \rightarrow \infty} -\frac{1}{n} \sum_{i_1, \dots, i_n} \mu(|i_1, \dots, i_n\rangle) \log(\mu(|i_1, \dots, i_n\rangle)). \quad (6)$$

This concept of entropy generalizes the concept of Shannon entropy in the following sense: if $P = (p_1, \dots, p_d)$ is a probability vector in $\{1, \dots, d\}$ and μ is the product (or Bernoulli) probability in $\Omega = \{1, \dots, d\}^{\mathbb{N}}$ satisfying

$$\mu(|b_1, \dots, b_n\rangle) = p_{b_1} \cdot p_{b_2} \cdots p_{b_n}$$

for any cylinder set, then $h(\mu) = S(P)$.

In Thermodynamic Formalism (see [23], [29]) it is also usual to consider the concept of pressure for a Lipschitz potential $\phi : \Omega \rightarrow \mathbb{R}$ and the associated **equilibrium probability**. We say that a shift-invariant probability μ_ϕ is the equilibrium probability for the Lipschitz function $\phi : \Omega \rightarrow \mathbb{R}$, if

$$P(\phi) := \sup_{\mu \text{ shift-invariant}} \left[\int \phi d\mu + h(\mu) \right] = \int \phi d\mu_\phi + h(\mu_\phi).$$

The number $P(\phi)$ is called the pressure of the potential ϕ .

Following [13], if μ_ϕ is the equilibrium probability for the Lipschitz function ϕ and if μ is shift-invariant, then, the **specific information gain** of μ with respect to μ_ϕ is given by

$$h(\mu, \mu_\phi) := \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{|i_1, \dots, i_n\rangle} \mu(|i_1, \dots, i_n\rangle) \log \left(\frac{\mu(|i_1, \dots, i_n\rangle)}{\mu_\phi(|i_1, \dots, i_n\rangle)} \right). \quad (7)$$

Furthermore, from Proposition 1 in [13] we get

$$h(\mu, \mu_\phi) = \underbrace{\left[\int \phi d\mu_\phi + h(\mu_\phi) \right]}_{P(\phi)} - \left[\int \phi d\mu + h(\mu) \right]. \quad (8)$$

The expression (4) is similar to the one which is given in (8), when considered $P(\phi) = 0$. Indeed, it is well known in Thermodynamic Formalism that, in the case $P(\phi) = 0$, we have

$$e^{\phi(|x_1, x_2, x_3, \dots|)} = \lim_{n \rightarrow \infty} \frac{\mu_\phi(|x_1, x_2, \dots, x_n|)}{\sum_i \mu_\phi(|i, x_2, \dots, x_n|)}$$

(see for example [23], cor. 3.2.2). We will call $J^\mu = \lim_{n \rightarrow \infty} \frac{\mu(|x_1, x_2, \dots, x_n|)}{\sum_i \mu(|i, x_2, \dots, x_n|)}$ the **Jacobian** of the shift-invariant probability μ . In this way $\log(J^{\mu_\phi}) = \phi$.

Comparing the equations (6) and (7) it is natural to interpret the specific information gain as a relative entropy. In [5] the value $h(\mu, \mu_\phi)$ is characterized by a variant of the Shannon-McMillan-Breiman theorem. Indeed, from a result on section 3.2 of [5] we get the following: consider a shift-invariant probability μ on Ω , and for a given Lipschitz function $\phi : \Omega \rightarrow \mathbb{R}$, consider the corresponding equilibrium measure μ_ϕ . Then, for μ almost every point $x = |x_1, x_2, x_3, \dots| \in \Omega$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \left(\frac{\mu(|x_1, x_2, \dots, x_n|)}{\mu_\phi(|x_1, x_2, \dots, x_n|)} \right) = h(\mu, \mu_\phi). \quad (9)$$

An interpretation of this expression in the sense of Statistical Mechanics of non equilibrium is the following: the observed system μ_ϕ is the equilibrium probability for the Lipschitz function ϕ , then, given a random point $x \in \Omega$, its time $n - 1$ orbit $\{x, \sigma(x), \dots, \sigma^{n-1}(x)\}$ describes the dynamical evolution of the system under consideration.

For each $n \in \mathbb{N}$, let $\nu_n^x = \frac{1}{n} (\delta_x + \delta_{\sigma(x)} + \dots + \delta_{\sigma^{n-1}(x)})$ the associated probability to x at time n . If x was chosen according to μ_ϕ , then, it is known that $\nu_n^x \rightarrow \mu_\phi$, as $n \rightarrow \infty$. Denote by μ another probability (which is not the equilibrium for ϕ). If x was chosen according to μ (and, so ν_n^x), then, as $n \rightarrow \infty$ we get

$$\frac{\mu(|x_1, x_2, \dots, x_n|)}{\mu_\phi(|x_1, x_2, \dots, x_n|)} \sim e^{n h(\mu, \mu_\phi)}.$$

Therefore, the value $h(\mu, \mu_\phi)$ quantifies the asymptotic exponential rate which describes how the dynamical time evolution of the system discriminates between μ_ϕ and μ , when $n \rightarrow \infty$.

In section 2 we consider measurable spaces X and Y and the product space $X \times Y$. For a fixed *a priori* probability ν on X and for any probability π on $X \times Y$ we introduce a definition of entropy of π relative to ν as

$$H^\nu(\pi) = - \sup \left\{ \int c(x, y) d\pi(x, y) \mid \int e^{c(x, y)} d\nu(x) = 1 \forall y, c \text{ measurable} \right\}. \quad (10)$$

This allows us to obtain a natural definition of information gain which coincides with the mutual information given in (2). Furthermore, by considering a generalized definition of entropy relative to a transverse function, as it was given in [15], we get a natural way of extending the concept of information gain as in (4) to measurable spaces.

We believe that the introduction of the *a priori* measure (which was indispensable in our work) and transverse functions is a novelty in the theory

We point out that what we call here transverse function is a very special and simple case of the general concept considered in the study of groupoids.

In section 3 we consider X and Y as compact metric spaces and we show that the entropy given by (10) is consistent with the definitions presented in [22] and [16] which consider a supremum taken over Lipschitz functions. We also exhibit another point of view for the concept of generalized information gain (it extends (4) and also (8)).

In section 4 we consider a dictionary which allow us to recollect the results from previous sections to the so called generalized XY model (which corresponds to replace the alphabet $\{1, \dots, d\}$ in Ω by a compact metric space). In particular we get a definition of specific information gain which generalizes the relative entropy in [16]. This definition of information gain is used later in section 5.

We will outline now some of the key elements on section 5 which considers the entropy production for the XY model. Remember that for the sake of notation (see (5)) we denote Ω by Ω^+ . The elements on Ω^+ are denoted by $x = |x_1, x_2, \dots, x_n, \dots\rangle$. Consider the space $\Omega^- = \{1, 2, \dots, d\}^{\mathbb{N}}$ (which is formally different from Ω^+). Any point in the space Ω^- is written in the form $(\dots, y_3, y_2, y_1|$ and any point in $\hat{\Omega} = \Omega^- \times \Omega^+$ will be written in the form $(\dots, y_3, y_2, y_1|x_1, x_2, x_3, \dots) = (y|x)$.

We consider on $\hat{\Omega}$ the shift map $\hat{\sigma}$ given by

$$\hat{\sigma}((\dots, y_3, y_2, y_1|x_1, x_2, x_3, \dots)) = (\dots, y_3, y_2, y_1, x_1|x_2, x_3, \dots). \quad (11)$$

The natural restriction of $\hat{\sigma}$ over $\Omega = \Omega^+$ is the shift map σ . The natural restriction of $\hat{\sigma}^{-1}$ over Ω^- is denoted by σ^- . Observe that (Ω^-, σ^-) can be identified with (Ω^+, σ) , via the conjugation $\theta : \Omega^- \rightarrow \Omega^+ = \Omega$, which is given by

$$\theta((\dots, z_3, z_2, z_1|) = |z_1, z_2, z_3, \dots). \quad (12)$$

Any σ -invariant probability μ on Ω^+ can be extended to a $\hat{\sigma}$ -invariant probability $\hat{\mu}$ on $\Omega^- \times \Omega$. The restriction of $\hat{\mu}$ to Ω^- , denoted by μ^- , is σ^- -invariant. By identifying (Ω^-, σ^-) with (Ω, σ) , via the conjugation θ and denoting by $\theta_*\mu^-$ the push forward of μ^- , we get

$$\theta_*\mu^-(|a_1, a_2, \dots, a_m|) = \mu(|a_m, \dots, a_2, a_1|). \quad (13)$$

Following [13], the **entropy production** of an equilibrium measure μ on Ω is defined by

$$e_p(\mu) := h(\mu, \theta_*\mu^-) = \lim_n \frac{1}{n} \sum_{|a_1, \dots, a_n|} \mu(|a_1, \dots, a_n|) \log \left(\frac{\mu(|a_1, \dots, a_n|)}{\mu(|a_n, \dots, a_2, a_1|)} \right).$$

It is natural in the study of the properties of the entropy production of equilibrium measures in the XY model to use as a tool the concept of involution kernel which will be defined later (for references about involution kernel see [2], [16] and [18]). In the same way as in section 4 most of the results we will consider in this section are for equilibrium probabilities of Lipschitz potentials.

We will show (see Proposition 44) that in the case the potential is symmetric the associated equilibrium probability has zero entropy production.

Results related to the role of the entropy production (the fluctuation theorem and the detailed balance condition) in problems in Physics and Dynamics can be found in [11], [13], [20], [25] and [3]. A concrete example of a system where the entropy production plays an important role is presented in [9]: a classical gas confined in a cylinder by a movable piston (see first page of [9]).

We would like to thank L. Cioletti for helpful comments during the writing of this paper.

2 Generalized Information gain

Our purpose in this section is to extend the definition of information gain $IG(\pi, P)$ and also $IG(\pi, \pi_0)$, given by (4), for the case when X and Y are measurable spaces. As we will see, a natural way of to extends (4) is by considering transverse functions and the notion of entropy given in [15]. This entropy was previously introduced in [22] for compact spaces using an a priori probability. Even in [22] it was considered for holonomic probabilities, such entropy has no dynamical content nor is related to an iterated function system. It extends the entropy previously introduced in [16] for shift-invariant probabilities when considered the generalized XY model (by identifying $Y = X^{\{2,3,4,\dots\}}$, with X compact).

From now on we consider σ -algebras \mathcal{A} on X and \mathcal{B} on Y and the product σ -algebra on $X \times Y$ (see [4] chap. 6). If $c : X \times Y \rightarrow \mathbb{R}$ is measurable then, for any fixed $y \in Y$, the function $c_y(x) := c(x, y)$ defined on X is measurable (see [4] theorem 6.7).

In order we can make an identification with the setting in [15] we consider in the space $X \times Y$ the equivalence relation $(x_1, y_1) \sim (x_2, y_2)$, if and only if, $y_1 = y_2$. So the equivalence classes are the horizontal lines of $X \times Y$.

Definition 1. We will call of an **a priori transverse function** any family $\hat{\nu} = \{\hat{\nu}^y \mid y \in Y\}$ of probabilities on $X \times Y$, such that,

- 1) $\forall y \in Y$, the probability $\hat{\nu}^y$ has support on $X_y = \{(x, y) \mid x \in X\}$,
- 2) $\forall A \subset X \times Y$ measurable, we have that $y \rightarrow \nu^y(A)$ is measurable.

The next definition is given from [15].

Definition 2. We define the entropy of any probability π on $X \times Y$ relative to the a priori transverse function $\hat{\nu}$ as

$$H^{\hat{\nu}}(\pi) = -\sup\left\{\int c(x, y) d\pi(x, y) \mid \int e^{c(x, y)} \hat{\nu}^y(dx) = 1 \forall y, c \text{ is measurable}\right\}.$$

We usually also fix a probability ν on X satisfying $\text{supp}(\nu) = X$ which we call the *a priori* probability on X . Clearly, given an a priori probability ν on X and considering the identification of X and X_y we can consider the a priori transverse function $\hat{\nu}$ given by $\hat{\nu}^y(dx) = \nu(dx)$. In this case we write $\hat{\nu} \equiv \nu$ and denote $H^{\hat{\nu}}(\pi)$ also by $H^{\nu}(\pi)$ which was given in (10).

Definition 3. We say that a measurable function $c : X \times Y \rightarrow \mathbb{R}$ is $\hat{\nu}$ -normalized if

$$\int e^{c(x, y)} \hat{\nu}^y(dx) = 1, \forall y \in Y.$$

If ν is an a priori probability on X , we say that $c : X \rightarrow \mathbb{R}$ is ν -normalized, if it is measurable and $\int e^c d\nu = 1$.

Remark 4. Following [15], a probability π on $X \times Y$ is called $\hat{\nu}$ -invariant if there is a bounded and $\hat{\nu}$ -normalized function V on $X \times Y$, such that, for any measurable function $f : X \times Y \rightarrow \mathbb{R}$, we get

$$\iint f(x, y) e^{V(x, y)} \hat{\nu}^y(dx) d\pi(x, y) = \int f(x, y) d\pi(x, y).$$

This is equivalent to say that π is a fixed point for the operator H_V^* where by definition

$$H_V(f)(x, y) = \int e^{V(x, y)} f(x, y) \hat{\nu}^y(dx).$$

The function $g = H_V(f)$ does not depend of the first coordinate, that is, g is constant on classes. Furthermore $H_V \circ H_V = H_V$ (see [15] for more results

concerning this operator). The function $J = e^V$ is called a $\hat{\nu}$ -Jacobian of π and $H^{\hat{\nu}}(\pi) = -\int V d\pi$.

It follows from Theorem 20 in [15] that for any probability Q on Y , and any bounded and $\hat{\nu}$ -normalized function V , there exists a unique $\hat{\nu}$ -invariant probability π on $X \times Y$ with $\hat{\nu}$ -Jacobian e^V and y -marginal Q . Furthermore,

$$\int f(x, y) d\pi(x, y) = \iint e^{V(x, y)} f(x, y) \hat{\nu}^y(dx) dQ(y).$$

The function $c = 0$ is $\hat{\nu}$ -normalized and therefore $H^{\hat{\nu}}(\pi) \leq 0$. If $\tilde{\nu}$ is a finite measure on X satisfying $\tilde{\nu}(X) = d$ and $d\hat{\nu} \equiv \frac{1}{d}d\tilde{\nu}$, then $H^{\hat{\nu}}(\pi) = H^{\tilde{\nu}}(\pi) + \log(d)$, where $H^{\tilde{\nu}}$ is defined in a similar way. Now, taking X and Y as finite sets and $\tilde{\nu}$ as the counting measure on X we reach the conclusion - applying equation (3) - that such definition of entropy is a natural extension of $H(\pi)$.

If P is a probability on X we also define

$$S^\nu(P) := -\sup \left\{ \int c(x) dP(x) \mid \int e^{c(x)} d\nu(x) = 1, \text{ where } c \text{ is measurable} \right\}.$$

We start by proving the next theorem which shows that the above definitions provide a variational characterization of the Kullback-Leibler divergence when considering an *a priori* probability ν .

Theorem 5. *Let P be a probability on X and π be a probability on $X \times Y$ with y -marginal Q . Let ν be an *a priori* probability on X . Then,*

$$S^\nu(P) = -D_{KL}(P \mid \nu) \quad \text{and} \quad H^\nu(\pi) = -D_{KL}(\pi \mid \nu \times Q).$$

The proof will be divided in several lemmas (until Lemma 12 which finishes the proof of Theorem 5).

Definition 6. *We say that a measurable function $J : X \times Y \rightarrow [0, +\infty)$ is a $\hat{\nu}$ -Jacobian, if $\int J(x, y) \hat{\nu}^y(dx) = 1, \forall y \in Y$.*

*Given an *a priori* probability ν on X , we say that a measurable function $J : X \rightarrow [0, +\infty)$ is a ν -Jacobian if $\int J d\nu = 1$.*

If c is $\hat{\nu}$ -normalized, then $J = e^c$ is a $\hat{\nu}$ -Jacobian. On the other hand, if J is a $\hat{\nu}$ -Jacobian and it does not assume the value zero, then $c = \log(J)$ is $\hat{\nu}$ -normalized.

Lemma 7.

$$H^{\hat{\nu}}(\pi) = -\sup \left\{ \int \log(J(x, y)) d\pi(x, y) \mid J \text{ is a } \hat{\nu}\text{-Jacobian} \right\}.$$

Proof. If c is $\hat{\nu}$ -normalized then $J = e^c$ is a $\hat{\nu}$ -Jacobian. It follows that

$$\begin{aligned} & \sup \left\{ \int \log(J(x, y)) d\pi(x, y) \mid J \text{ is a } \hat{\nu}\text{-Jacobian} \right\} \\ & \geq \sup \left\{ \int c(x, y) d\pi(x, y) \mid c \text{ is } \hat{\nu}\text{-normalized} \right\}. \end{aligned}$$

On the other hand, for any fixed $\hat{\nu}$ -Jacobian J , let $J_n = \frac{J + \frac{1}{n}}{1 + \frac{1}{n}}$. The function J_n is also a $\hat{\nu}$ -Jacobian and $\lim \log(J_n(x, y)) = \log(J(x, y))$, for any $(x, y) \in X \times Y$. Furthermore, applying Fatou's lemma, we get

$$\int \log(J) d\pi \leq \liminf_n \int \log(J + \frac{1}{n}) d\pi = \liminf_n \int \log(J_n) d\pi.$$

As the function $c_n = \log(J_n)$ is $\hat{\nu}$ -normalized we finish the proof. \square

Lemma 8. *Let P be a probability on X and π be a probability on $X \times Y$, with x -marginal P . If P is not absolutely continuous with respect to the a priori probability ν , then $S^\nu(P) = -\infty$ and $H^\nu(\pi) = -\infty$.*

Proof. If P is not absolutely continuous with respect to ν then there exists a measurable set A , such that, $\nu(A) = 0$ and $P(A) > 0$. For each $\beta > 0$, let $c_\beta : X \rightarrow \mathbb{R}$ be the measurable function defined as

$$c_\beta(x) = \begin{cases} 0 & \text{if } x \in X - A \\ \beta & \text{if } x \in A \end{cases}.$$

Then, we have $\int e^{c_\beta(x)} d\nu(x) = 1$ and $\int c_\beta(x) d\pi(x, y) = \int c_\beta(x) dP(x) = \beta P(A)$. As β is arbitrary we can take $\beta \rightarrow +\infty$, and then we get that $S^\nu(P) = -\infty$ and also $H^\nu(\pi) = -\infty$. \square

As usual, we use the notation $\mu \ll \nu$ to denote that μ is absolutely continuous with respect to ν . If $P \ll \nu$ we denote by $\frac{dP}{d\nu}$ the Radon-Nikodym derivative of P with respect to ν , which is measurable.

Observe that $J_0 := \frac{dP}{d\nu}$ is a ν -Jacobian. Let $X_0 = \{x \in X \mid J_0(x) > 0\}$. Given any measurable and bounded function $f : X \rightarrow \mathbb{R}$, we have

$$\int f dP = \int f \cdot J_0 d\nu = \int f \cdot I_{X_0} \cdot J_0 d\nu = \int_{X_0} f dP.$$

It follows that $P(X_0) = 1$ and $\int f(x) dP(x) = \int_{X_0} f(x) dP(x)$, for any measurable function f .

The next result shows that $-S^\nu(P)$ is the Kullback-Leibler divergence of P with respect to ν .

Lemma 9. *Let P be a probability on X , such that, $P \ll \nu$. Then,*

$$S^\nu(P) = -D_{KL}(P | \nu) = - \int \log\left(\frac{dP}{d\nu}\right) dP.$$

Proof. Let $J_0 := \frac{dP}{d\nu}$ and $X_0 := \{x \in X \mid J_0(x) > 0\}$. We claim that

$$\int_{X_0} \log(J_0) dP = \sup \left\{ \int_{X_0} \log(J(x)) dP(x) \mid J \text{ is a } \nu\text{-Jacobian} \right\}.$$

Indeed, for any ν -Jacobian $J : X \rightarrow [0, +\infty)$, by applying the Jensen's inequality we have

$$\int_{X_0} \log\left(\frac{J}{J_0}\right) dP \leq \log \int_{X_0} \frac{J}{J_0} dP = \log \int_{X_0} J d\nu \leq \log \int J d\nu = 0.$$

This shows that

$$\int_{X_0} \log(J) dP \leq \int_{X_0} \log(J_0) dP.$$

□

A similar result for π will be given by next result.

Lemma 10. *Assume that there exists a $\hat{\nu}$ -Jacobian J on $X \times Y$ satisfying*

$$\iint f(x, y) J_0(x, y) \hat{\nu}^y(dx) d\pi(x, y) = \int f(x, y) d\pi(x, y), \quad (14)$$

for any measurable function $f : X \times Y \rightarrow \mathbb{R}$. Then,

$$H^{\hat{\nu}}(\pi) = - \int \log(J_0) d\pi.$$

Proof. The reasoning is similar to the previous case. The set $A_0 = \{(x, y) \in X \times Y \mid J_0(x, y) > 0\}$ satisfies $\pi(A_0) = 1$. For any $\hat{\nu}$ -Jacobian $J : X \times Y \rightarrow [0, +\infty)$ we have

$$\begin{aligned} \int_{A_0} \log\left(\frac{J}{J_0}\right) d\pi &\leq \log \int_{A_0} \frac{J}{J_0} d\pi = \log \int \frac{J}{J_0} \cdot I_{A_0} \cdot J_0 \hat{\nu}^y(dx) d\pi(x, y) \\ &= \log \int J \cdot I_{A_0} \hat{\nu}^y(dx) d\pi(x, y) \leq \log \int J \hat{\nu}^y(dx) d\pi(x, y) = 0. \end{aligned}$$

□

We will say that a function J_0 satisfying (14) is a $\hat{\nu}$ -**Jacobian** of π .

Denoting by Q the y -marginal of π , the equation (14) can be rewritten as

$$\iint f(x, y) J_0(x, y) \hat{\nu}^y(dx) dQ(y) = \int f(x, y) d\pi(x, y)$$

and so $J_0(x, y) \hat{\nu}^y(dx) dQ(y)$ is the disintegration of π with respect to the horizontal lines of $X \times Y$. Supposing also that ν is an a priori probability on X and $\hat{\nu} \equiv \nu$, we get $\pi \ll \nu \times Q$ with $\frac{d\pi}{d\nu dQ} = J_0$. Then, under the hypotheses of above proposition and assuming $\hat{\nu} \equiv \nu$, we get

$$H^\nu(\pi) = -D_{KL}(\pi | \nu \times Q).$$

Lemma 11. *Let ν be an a priori probability on X and π be a probability on $X \times Y$ with y -marginal Q . Suppose that $\pi \ll \nu \times Q$. Then,*

$$H^\nu(\pi) = -D_{KL}(\pi | \nu \times Q).$$

Proof. We suppose that $\pi \ll \nu \times Q$ and we denote by $J = \frac{d\pi}{d\nu dQ}$ its Radon-Nikodym derivative. Then, for any measurable function $g : X \times Y \rightarrow \mathbb{R}$ we have

$$\iint g(x, y) J(x, y) d\nu(x) dQ(y) = \int g(x, y) d\pi(x, y).$$

As Q is the y -marginal of π , taking functions g depending just of the second coordinate, we get

$$\int g(y) \left[\int J(x, y) d\nu(x) \right] dQ(y) = \int g(y) dQ(y),$$

and, then $\int J(x, y) d\nu(x) = \frac{dQ}{dQ} = 1$, for Q -a.e. y . Replacing J by 1 in a subset of $X \times Y$ having zero measure with respect to π , we get a measurable Jacobian \tilde{J} satisfying, for any measurable function $g : X \times Y \rightarrow \mathbb{R}$,

$$\iint g(x, y) \tilde{J}(x, y) d\nu(x) dQ(y) = \int g(x, y) d\pi(x, y).$$

It follows from Lemma 10 that

$$H^\nu(\pi) = - \int \log(\tilde{J}) d\pi = - \int \log(J) d\pi = -D_{KL}(\pi | \nu \times Q).$$

□

Lemma 12. *Let ν be an a priori probability on X and π be a probability on $X \times Y$ with y -marginal Q . If π is not absolutely continuous with respect to $\nu \times Q$, then $H^\nu(\pi) = -\infty$.*

Proof. If π is not absolutely continuous with respect to $\nu \times Q$, then, there exists a measurable set $A \subset X \times Y$, such that, $(\nu \times Q)(A) = 0$ and $\pi(A) > 0$. For each $y \in Y$ let $A_y := \{x \in X \mid (x, y) \in A\}$ (it can be the empty set). The set $A_y \subset X$ is measurable and the function $\phi : Y \rightarrow [0, +\infty)$ given by $\phi(y) = \nu(A_y)$ is measurable and satisfies $\int_Y \nu(A_y) dQ(y) = (\nu \times Q)(A) = 0$ (see [4] chap. 6). It follows that $\{y \mid \nu(A_y) \neq 0\}$ is a measurable set on Y satisfying $Q(\{y \mid \nu(A_y) \neq 0\}) = 0$. The set $X \times \{y \mid \nu(A_y) \neq 0\}$ is measurable in $X \times Y$ and, as the y -marginal of π is Q , we get $\pi(X \times \{y \mid \nu(A_y) \neq 0\}) = Q(\{y \mid \nu(A_y) \neq 0\}) = 0$. Let $B = A - (X \times \{y \mid \nu(A_y) \neq 0\})$. The set B is measurable and $\pi(B) = \pi(A) > 0$, while $(\nu \times Q)(B) = 0$.

For each $\beta > 0$, let $c_\beta : X \times Y \rightarrow \mathbb{R}$ be the measurable function defined as

$$c_\beta(x, y) = \begin{cases} 0 & \text{if } (x, y) \in B^C \\ \beta & \text{if } (x, y) \in B \end{cases} .$$

For each fixed $y \in Y$ we denote $B_y = \{x \in X \mid (x, y) \in B\}$. By the construction of B we get $\nu(B_y) = 0$. Then, for fixed y we have

$$\int e^{c_\beta(x, y)} d\nu(x) = e^\beta \nu(B_y) + e^0 \nu(X - B_y) = 1.$$

This shows that c_β is ν -normalized. Furthermore, $\int c_\beta(x, y) d\pi(x, y) = \beta \pi(B)$. As β is arbitrary we can take $\beta \rightarrow +\infty$ and then we get that $H^\nu(\pi) = -\infty$. \square

The above results conclude the proof of Theorem 5.

Proposition 13. *Let P be a probability on X satisfying $P \ll \nu$. Consider any probability Q on Y and any probability π on $X \times Y$ with x -marginal P . Then, we have:*

1. $H^\nu(\pi) \leq S^\nu(P)$
2. $S^\nu(P) = H^\nu(P \times Q)$.

Proof. The proof of item 1. is a direct consequence of the definitions of S^ν and H^ν because we can consider any measurable function $c : X \rightarrow \mathbb{R}$ as a measurable function defined on $X \times Y$ which depends just on the first coordinate.

In order to prove item 2. we consider the function $J(x, y) = \frac{dP}{d\nu}(x)$. This function is a ν -Jacobian of $P \times Q$, then, applying propositions 9 and 10, we conclude the proof. \square

The proof of the next result follows the same reasoning which were used in [16] and [15] and will be left for the reader.

Proposition 14. *The entropy $H^{\hat{\nu}}(\cdot)$ has the following properties:*

1. $H^{\hat{\nu}}$ is concave
2. $H^{\hat{\nu}}$ is upper semi continuous. More precisely, if $\int f d\pi_n \rightarrow \int f d\pi$, for any measurable and bounded function f on $X \times Y$, then, $\limsup_n H^{\hat{\nu}}(\pi_n) \leq H^{\hat{\nu}}(\pi)$.

Definition 15. *We define the **information gain** of a probability π on $X \times Y$, with respect to the a priori transverse function $\hat{\nu}$, as*

$$IG(\pi, \hat{\nu}) = -H^{\hat{\nu}}(\pi).$$

If π has marginals P and Q and $\pi \ll P \times Q$, then choosing $\hat{\nu} \equiv P$ we get

$$IG(\pi, P) = -H^P(\pi) = D_{KL}(\pi | P \times Q),$$

which is the mutual information of π given by (2). This shows that the above definition is a natural extension of the usual definition of information gain. As $c = 0$ is $\hat{\nu}$ -normalized we get $IG(\pi, \hat{\nu}) \geq 0$. Furthermore, $IG(\pi, \hat{\nu}) = 0$, if $d\pi = \hat{\nu}^y(dx)dQ(y)$ for some Q .

The information gain $IG(\pi, P)$ above defined can be computed from $S^\nu(P)$ and $H^\nu(\pi)$ and it “does not depend” of the choice of an a priori probability ν on X as the following result shows.

Theorem 16. *Let P be a probability on X , π be a probability on $X \times Y$, with x -marginal P and let ν be an a priori probability on X . Assume that $S^\nu(P)$ and $H^\nu(\pi)$ are finite. Then,*

$$S^\nu(P) - H^\nu(\pi) = IG(\pi, P).$$

Proof. By hypothesis there exists $\phi : X \rightarrow [0, +\infty)$, a ν -Jacobian of P and $J : X \times Y \rightarrow [0, +\infty)$, which is a ν -Jacobian of π . Denoting by Q the y -marginal of π , we have $d\pi(x, y) = J(x, y)d\nu(x)dQ(y)$ and $dP(x) = \phi(x)d\nu(x)$. The set $A = \{x \in X | \phi(x) > 0\}$ satisfies $P(A) = 1$, and as π has x -marginal P , we finally get $\pi(A \times Y) = 1$. So we can write $d\pi(x, y) = \frac{J(x, y)}{\phi(x)}dP(x)dQ(y)$ and therefore

$$S^\nu(P) - H^\nu(\pi) = - \int \log(\phi)dP + \int \log(J)d\pi = -H^P(\pi) = IG(\pi, P).$$

□

Proposition 17. *Let $\hat{\nu}$ be an a priori transverse function and π be a probability on $X \times Y$. Given a bounded and $\hat{\nu}$ -normalized function $\phi_0 : X \times Y \rightarrow \mathbb{R}$, consider the a priori transverse function $\hat{\mu}^y(dx) = e^{\phi_0(x, y)}\hat{\nu}^y(dx)$. Then,*

$$IG(\pi, \hat{\mu}) = - \int \phi_0 d\pi + IG(\pi, \hat{\nu}).$$

Proof.

$$\begin{aligned}
IG(\pi, \hat{\mu}) &= -H^{\hat{\mu}}(\pi) = \sup\left\{\int c d\pi \mid \int e^{c(x,y)} \hat{\mu}^y(dx) = 1 \forall y\right\} \\
&= \sup\left\{\int c d\pi \mid \int e^{c+\phi_0} \hat{\nu}^y(dx) = 1 \forall y\right\} \\
&= \sup\left\{\int c - \phi_0 d\pi \mid \int e^c d\hat{\nu}^y(dx) = 1 \forall y\right\} \\
&= -\int \phi_0 d\pi - H^{\hat{\nu}}(\pi).
\end{aligned}$$

□

Corollary 18. *Let ν be an a priori probability on X and π be a probability on $X \times Y$. Given a bounded and ν -normalized function $\phi_0 : X \times Y \rightarrow \mathbb{R}$, consider the a priori transverse function $\hat{\mu}^y(dx) = e^{\phi_0(x,y)} d\nu(x)$. Then,*

$$IG(\pi, \hat{\mu}) = -\int \phi_0 d\pi - H^\nu(\pi).$$

This last result shows that the above definition of information gain, using transverse functions, is compatible with (4). Given a probability Q_0 on Y we can associate a probability π_0 on $X \times Y$ given by $d\pi_0 = e^{\phi_0(x,y)} d\nu(x) dQ_0(y)$. A natural generalization of (4) could be given by the expression

$$IG(\pi, \pi_0) = -\int \phi_0 d\pi - H^\nu(\pi). \quad (15)$$

We observe that the right hand side of this equation contains just the objects π, ϕ_0 and ν , which allow us to realize that we are not using Q_0 and the associate π_0 , but only the transverse function $\hat{\mu} = e^{\phi_0} d\nu$, which is part of a disintegration of π_0 . In this sense it is natural to define information gain by using transverse functions and to consider $IG(\pi, \hat{\mu})$ instead of trying to define $IG(\pi, \pi_0)$.

Finally we remark that (15) is not well defined. Indeed, if $Y = \{0, 1\}$ and $\pi_0 = \nu \times \delta_0$, the functions ϕ_0 and ψ_0 given by $\phi_0(x, y) = 0$ and $\psi_0(x, y) = y$, provide two different disintegrations of π_0 , which are

$$d\pi_0(x, y) = e^{\phi_0(x,y)} d\nu(x) d\delta_0(y) \quad \text{and} \quad d\pi_0(x, y) = e^{\psi_0(x,y)} d\nu(x) d\delta_0(y).$$

If $\pi = \nu \times \delta_1$, then $H^\nu(\pi) = H^\nu(\nu \times \delta_1) = S^\nu(\nu) = 0$, and so

$$-\int \psi_0 d\pi - H^\nu(\pi) = -\int \psi_0 d\pi = 1 \quad \text{while} \quad -\int \phi_0 d\pi - H^\nu(\pi) = 0.$$

3 Entropy and information gain for compact metric spaces

In this section we consider X and Y as compact metric spaces equipped with the respective Borel σ -algebras. We will prove that, in this case, H^ν coincides with the entropy considered in [22], which is defined from a supremum taken over Lipschitz functions instead measurable functions. This shows that the entropy as defined in [15], which is also considered here, extends the entropy in [16] and [22]. We also propose to consider the Information Gain by using a probability π_0 instead a transverse measure $\hat{\nu}$, as was discussed in the end of previous section. This will be coherent with the reasoning of future sections and also with (8) and (4).

Proposition 19. *Suppose that X and Y are compact metric spaces and consider the Borel sigma-algebras in X and Y . Then,*

$$H^\nu(\pi) = -\sup\left\{\int f(x, y) d\pi(x, y) \mid \int e^{f(x, y)} d\nu(x) = 1, \forall y, \text{ with } f \text{ Lipschitz}\right\}.$$

Proof. By definition

$$H^\nu(\pi) = -\sup\left\{\int f(x, y) d\pi(x, y) \mid \int e^{f(x, y)} d\nu(x) = 1, \forall y, \text{ with } f \text{ measurable}\right\}.$$

We denote

$$h^\nu(\pi) := -\sup\left\{\int f(x, y) d\pi(x, y) \mid \int e^{f(x, y)} d\nu(x) = 1, \forall y, \text{ with } f \text{ Lipschitz}\right\}.$$

It will be necessary to prove that $H^\nu = h^\nu$.

If $\psi : X \times Y \rightarrow \mathbb{R}$ is Lipschitz and ν -normalized, Q is any probability on Y and π_ψ is the probability on $X \times Y$, given by

$$\int f(x, y) d\pi_\psi(x, y) := \iint f(x, y) e^{\psi(x, y)} d\nu(x) dQ(y), \text{ for } f \text{ measurable,}$$

then, e^ψ is a ν -Jacobian of π_ψ . It follows in this case, that

$$H^\nu(\pi_\psi) = -\int \psi(x, y) d\pi_\psi(x, y) = h^\nu(\pi_\psi).$$

Suppose by contradiction there exists a probability η on $X \times Y$, such that, $-H^\nu(\eta) > -h^\nu(\eta)$. Consequently, $-h^\nu(\eta) \neq +\infty$.

First, we claim that there exists a Lipschitz function $\varphi : X \times Y \rightarrow \mathbb{R}$, such that, for any probability π on $X \times Y$,

$$\int \varphi d\eta + h^\nu(\eta) > \int \varphi d\pi + H^\nu(\pi).$$

The proof of this claim follows the same lines of the proof of Theorem 3 in [1] (see also [10] chap. 1). We consider the weak* topology on the space of finite signed-measures and we extend H^ν and h^ν as the value $-\infty$, if π is not a probability.

As $-H^\nu$ is convex, non negative and lower semi-continuous, its epigraph $\text{epi}(-H^\nu) = \{(\pi, t) \mid -H^\nu(\pi) \leq t\}$ is convex and closed. As $(\eta, -h^\nu(\eta)) \notin \text{epi}(-H^\nu)$, it follows from Hahn-Banach theorem that there is $c \in \mathbb{R}$ and a linear functional

$$(\pi, t) \rightarrow \int g d\pi + at,$$

where g is a fixed continuous function and $a \in \mathbb{R}$ is fixed, such that, for any $(\pi, t) \in \text{epi}(-H^\nu)$ we have

$$\int g d\eta - ah^\nu(\eta) < c < \int g d\pi + at.$$

Observe that necessarily $a > 0$. We denote $\varphi = -\frac{g}{a}$. If a probability π satisfies $-H^\nu(\pi) < +\infty$, then $(\pi, -H^\nu(\pi)) \in \text{epi}(-H^\nu)$, and, finally, we get

$$\int \varphi d\eta + h^\nu(\eta) > -\frac{c}{a} > \int \varphi d\pi + H^\nu(\pi).$$

Even in the case $-H^\nu(\pi) = +\infty$ these inequalities remain valid.

Finally, as the Lipschitz functions are dense in the set of continuous functions (in the uniform convergence) we can assume that for a Lipschitz function φ we have

$$\int \varphi d\eta + h^\nu(\eta) > \int \varphi d\pi + H^\nu(\pi),$$

for any probability π . This finishes the proof of the claim.

Let Q be the y -marginal of η and $\tilde{\varphi}(y) = \log(\int e^{\varphi(x,y)} d\nu(x))$. The function $\psi(x, y) = \varphi(x, y) - \tilde{\varphi}(y)$ is Lipschitz, ν -normalized and for any probability π on $X \times Y$, with y -marginal Q , we have

$$\int \psi d\eta + h^\nu(\eta) > \int \psi d\pi + H^\nu(\pi).$$

Let $\pi = \pi_\psi$ be defined by

$$\int f(x, y) d\pi_\psi(x, y) := \iint f(x, y) e^{\psi(x, y)} d\nu(x) dQ(y).$$

Then, as π_ψ has y -marginal Q and $H^\nu(\pi_\psi) = -\int \psi d\pi_\psi$, we get that

$$\int \psi d\eta + h^\nu(\eta) > \int \psi d\pi_\psi + H^\nu(\pi_\psi) = 0.$$

This is a contradiction because, by definition of h^ν , $\int \psi d\eta + h^\nu(\eta) \leq 0$. \square

The next results will be necessary in the direction of getting a different point of view for the concept of information gain.

Proposition 20. *Let X and Y be compact metric spaces and suppose that there exists a continuous/Lipschitz function $\phi : X \times Y \rightarrow \mathbb{R}$, such that, $J = e^\phi$ is a ν -Jacobian of π . Let P and Q be the marginals of π . Then, P is equivalent to ν (each one is absolutely continuous with respect to each other) and*

$$\frac{dP}{d\nu}(x) = \int e^{\phi(x, y)} dQ(y),$$

which is also continuous/Lipschitz. There exist constants $c_2 > c_1 > 0$, such that, $c_1 < \frac{dP}{d\nu} < c_2$, $\forall x \in X$. Finally, defining $\psi := \phi - \log(\frac{dP}{d\nu})$, we have that e^ψ is a P -Jacobian of π .

Proof. If a measurable and bounded function g depends only of the first coordinate, then, using Fubini's theorem, we get

$$\begin{aligned} \int g(x) dP(x) &= \int g(x) d\pi(x, y) = \iint e^{\phi(x, y)} g(x) d\nu(x) dQ(y) \\ &= \int \left[\int e^{\phi(x, y)} dQ(y) \right] g(x) d\nu(x). \end{aligned}$$

It follows that $\frac{dP}{d\nu}(x) = \int e^{\phi(x, y)} dQ(y)$ is a density. Clearly the function $\frac{dP}{d\nu}$ is continuous/Lipschitz and there are constants $c_2 > c_1 > 0$, such that, $c_1 < \frac{dP}{d\nu}(x) < c_2$, $\forall x \in X$. This shows that P and ν are equivalent and that $\log(\frac{dP}{d\nu})$ is continuous/Lipschitz.

Let $\psi(x, y) := \phi(x, y) - \log(\frac{dP}{d\nu})(x)$. Then, for any measurable and bounded function $g : X \times Y \rightarrow \mathbb{R}$ we have

$$\int e^{\psi(x, y)} g(x, y) dP(x) = \int e^{\phi(x, y)} g(x, y) d\nu(x).$$

By integrating both sides with respect to Q and using the fact that e^ϕ is a ν -Jacobian of π , we get

$$\int \left[\int e^{\psi(x,y)} g(x,y) dP(x) \right] dQ(y) = \int g(x,y) d\pi(x,y).$$

This shows that ψ is a P -Jacobian of π . \square

Proposition 21. *Suppose that X and Y are compact metric spaces. Let ν be a probability on X with $\text{supp}(\nu) = X$ and $\phi : X \times Y \rightarrow \mathbb{R}$ be a continuous function. Let π_0 be a probability on $X \times Y$ with ν -Jacobian $J = e^\phi$. If π_0 is positive on open sets of $X \times Y$, then ϕ is the unique continuous function, such that, e^ϕ is a ν -Jacobian of π_0 .*

Proof. Let Q be the y -marginal of π_0 and ϕ_2 be a continuous function, such that, e^{ϕ_2} is a ν -Jacobian of π_0 . Then,

$$d\pi_0(x,y) = e^{\phi(x,y)} d\nu(x) dQ(y) \quad \text{and} \quad d\pi_0(x,y) = e^{\phi_2(x,y)} d\nu(x) dQ(y).$$

This shows that the positive functions e^{ϕ_2} and e^ϕ satisfy $e^{\phi_2(x,y)} = e^{\phi(x,y)}$, π_0 -a.e. $(x,y) \in X \times Y$. As π_0 is positive on open sets and ϕ, ϕ_2 are also continuous, we get $\phi_2(x,y) = \phi(x,y)$, for all $(x,y) \in X \times Y$. \square

Definition 22 (Information Gain for compact spaces). *Let X and Y be compact metric spaces and π_0 be a probability on $X \times Y$ which is positive on open sets and has x -marginal P_0 . Suppose there exists a continuous function $\psi_0 : X \times Y \rightarrow \mathbb{R}$, such that, e^{ψ_0} is a P_0 -Jacobian of π_0 . For any probability π on $X \times Y$ we define the information gain of π with respect to π_0 as*

$$IG(\pi, \pi_0) = - \int \psi_0 d\pi - H^{P_0}(\pi). \quad (16)$$

Suppose that π_0 has y -marginal Q_0 and $d\pi_0(x,y) = e^{\psi_0(x,y)} dP_0(x) dQ_0(y)$. Observe that in the expression $-\int \psi_0 d\pi - H^{P_0}(\pi)$ of (16) appears ψ_0 and P_0 but not Q_0 . By considering the transverse function $\hat{\nu}^y(dx) = e^{\psi_0(x,y)} dP_0(x)$ the next result shows that this definition is coherent with Definition 15.

Proposition 23. *Under the assumptions of Definition 22, we define $\hat{\nu}^y(dx) = e^{\psi_0(x,y)} dP_0(x)$. Then,*

$$IG(\pi, \pi_0) = IG(\pi, \hat{\nu}).$$

Proof. It is a consequence of corollary 18. \square

The next proposition shows that the above interpretation of information gain does not depend, in a sense to be explained, of the choice of the *a priori* probability ν .

Proposition 24. *Let X and Y be compact metric spaces and π_0 be a probability on $X \times Y$ which is positive on open sets. Let ν be an a priori probability on X . Suppose there exist a continuous function ϕ , such that, e^ϕ is a ν -Jacobian of π_0 . Let π be any probability on $X \times Y$. Then,*

$$IG(\pi, \pi_0) = - \int \phi d\pi - H^\nu(\pi).$$

Proof. Let P and Q be the marginals of π_0 . From lemma 20, P is equivalent to ν , with $\frac{dP}{d\nu}(x) = \int e^{\phi(x,y)} dQ(y)$. Furthermore, the continuous function $\psi := \phi - \log(\frac{dP}{d\nu})$ is such that e^ψ is the P -Jacobian of π_0 . Clearly, $e^{\psi-\phi} = \frac{d\nu}{dP}$, and so a Lipschitz function c is ν -normalized, if and only if, $c + \psi - \phi$ is P -normalized.

It follows that for any probability π we get

$$H^\nu(\pi) = H^P(\pi) + \int \psi - \phi d\pi.$$

Therefore,

$$IG(\pi, \pi_0) = - \int \psi d\pi - H^P(\pi) = - \int \phi d\pi - H^\nu(\pi).$$

□

It is natural to consider that (16) generalizes (4). The next example shows that (16) also generalizes (8) in a certain sense.

Example 25. *Suppose $X = \{1, 2, \dots, d\}$, $Y = \{1, 2, \dots, d\}^{\{2,3,4,5,\dots\}}$ and identify Ω with $X \times Y$ by the homeomorphism*

$$\Omega \ni |x_1, x_2, x_3, x_4, \dots) \rightarrow (x_1, |x_2, x_3, x_4, \dots)) \in X \times Y. \quad (17)$$

When considering the a priori probability ν as the counting measure on X , we get that a ν -Jacobian of an invariant probability π is given by

$$J^\pi(x_1, x_2, x_3, \dots) = \lim_{n \rightarrow \infty} \frac{\pi(x_1, x_2, \dots, x_n)}{\pi(x_2, x_3, \dots, x_n)},$$

for π a.e. $x \in \Omega$ (it can be extended for any point of Ω by taking $J^\pi = \frac{1}{d}$ in a set of zero measure). Furthermore, the Kolmogorov-Sinai entropy of π coincides with $H^\nu(\pi)$. A measurable function $\phi : \Omega \rightarrow \mathbb{R}$ is normalized if $\sum_a e^{\phi(a, x_2, x_3, \dots)} = 1$, for all $|x_2, x_3, \dots)$. If ϕ is Lipschitz with corresponding equilibrium measure π_ϕ , then the unique continuous Jacobian of π_ϕ is e^ϕ and the pressure of ϕ is zero. It follows from (8) and Proposition 24 that for any invariant probability π we have

$$h(\pi, \pi_\phi) = IG(\pi, \pi_\phi).$$

4 Specific information gain in the generalized XY model

In this section we introduce a dictionary connecting the definitions and results of previous sections with analogous ones for the generalized XY model. This will allow us to introduce the specific information gain in this setting which will be necessary in order to introduce the concept of entropy production in the next section. First we will remember some of the main definitions and results about the generalized XY model described in [16] which will be used here.

Let (M, d) be a compact metric space and denote by $\Omega = \Omega^+$ the space $M^{\mathbb{N}}$. Elements in Ω will be written in the form $x = |x_1, x_2, x_3, \dots\rangle$, $x_i \in M$. The space Ω is compact using the metric $d(x, y) = \sum_{i=1}^{\infty} \frac{d(x_i, y_i)}{2^i}$. We also consider the Borel sigma-algebra in Ω .

The relation of the setting of this section with the previous ones can be clarified by considering $X = M$, $Y = M^{\{2,3,4,5,\dots\}}$ and identifying Ω with $X \times Y$, using the homeomorphism given in (17). Observe that Y can be also identified with $X \times Y$ using the homeomorphism $|x_2, x_3, x_4, \dots\rangle \rightarrow (x_2, |x_3, x_4, x_5, \dots\rangle)$. From this identification the shift map $\sigma : \Omega \rightarrow \Omega$ given by $\sigma(|x_1, x_2, x_3, \dots\rangle) = |x_2, x_3, \dots\rangle$ can be also interpreted as the projection on Y . We say that a probability μ on $\Omega = M^{\mathbb{N}}$ is invariant for the shift map σ (or, shift-invariant), if for any continuous function $f : \Omega \rightarrow \mathbb{R}$, we have $\int f d\mu = \int f \circ \sigma d\mu$.

Assume we fixed an *a priori* probability ν on M satisfying $\text{supp}(\nu) = M$. For each Lipschitz function $A : \Omega \rightarrow \mathbb{R}$ we consider the linear operator $\mathcal{L}_{A,\nu} : C(\Omega) \rightarrow C(\Omega)$ defined by

$$\mathcal{L}_{A,\nu}(f)(x) = \int e^{A(|a, x_1, x_2, x_3, \dots\rangle)} f(|a, x_1, x_2, x_3, \dots\rangle) d\nu(a), \quad x = |x_1, x_2, x_3, \dots\rangle.$$

We call $\mathcal{L}_{A,\nu}$ the Ruelle operator (or, transfer operator) associated to the Lipschitz potential A and the *a priori* probability ν (we refer the reader to [16] for general properties of this operator).

For this operator there exists a unique (simple) positive eigenvalue λ_A associated to a positive eigenfunction $h = h_A$. If a continuous function $h > 0$ satisfies $\mathcal{L}_{A,\nu}(h) = \lambda_A \cdot h$, then h is Lipschitz. If h_1 and h_2 are eigenfunctions associated to λ_A , then $h_2 = c \cdot h_1$ for some constant c . There exists a unique probability measure ρ_A on Ω satisfying $\mathcal{L}_{A,\nu}^*(\rho_A) = \lambda_A \cdot \rho_A$, which means that

$$\int \mathcal{L}_{A,\nu}(f) d\rho_A = \lambda_A \int f d\rho_A,$$

for any continuous function $f : \Omega \rightarrow \mathbb{R}$. For convenience we fix the eigenfunction h_A which satisfies $\int h_A d\rho_A = 1$.

We point out that in the case the space of symbols M is not countable you really need to introduce an *a priori* probability in order to get a transfer operator.

A Lipschitz function A is called ν -normalized if $\mathcal{L}_{\bar{A},\nu}(1) = 1$, that is,

$$\int e^{A(|a,x_1,x_2,x_3,\dots|)} d\nu(a) = 1, \quad \forall x = |x_1, x_2, x_3, \dots|.$$

The function

$$\bar{A} = A + \log(h_A) - \log(h_A \circ \sigma) - \log(\lambda_A) \quad (18)$$

is ν -normalized. The associated eigenprobability $\rho_{\bar{A}}$ is shift-invariant and it will be denoted also by μ_A . It also satisfies $d\mu_A = h_A d\rho_A$ and $\mathcal{L}_{\bar{A},\nu}^*(\mu_A) = \mu_A$.

The **relative entropy** of an invariant probability μ on Ω with respect to the *a priori* probability ν on M is defined in [16] as

$$h^\nu(\mu) = - \sup_{B \text{ is } \nu\text{-normalized}} \int B d\mu.$$

Considering the above identification of Ω and $X \times Y$ and applying Proposition 19 we see that this definition is consistent with the previous definition of H^ν . Let us formally enunciate this result (using also Theorem 5).

Theorem 26. *Denote by π the probability on $X \times Y$, which corresponds to the shift-invariant probability μ on Ω , we get that the relative entropy $h^\nu(\mu)$, as defined in [16], coincides with $H^\nu(\pi)$. Furthermore, if Q is the y -marginal of π (which is identified with π because μ is shift invariant), then we have*

$$h^\nu(\mu) = H^\nu(\pi) = -D_{KL}(\pi | \nu \times Q).$$

In [1] it is proved that $h^\nu(\mu)$ coincides with the so called specific entropy of Statistical Mechanics (see [12]).

For any Lipschitz function $A : \Omega \rightarrow \mathbb{R}$ we have $h^\nu(\mu_A) = - \int \bar{A} d\mu_A$. Furthermore,

$$P_\nu(A) := \sup_{\mu \text{ shift-invariant}} \int A d\mu + h^\nu(\mu) = \int A d\mu_A + h^\nu(\mu_A) = \log(\lambda_A).$$

The number $P_\nu(A)$ is called the ν -pressure of A . A probability μ attaining the supremum value $P_\nu(A)$ is called an **equilibrium probability** for A . In [1] it is proved that μ_A is the unique equilibrium probability for A .

If M is a finite set with d elements and the *a priori* probability ν is set to be the counting measure (which is not a probability), then the relative entropy $h^\nu(\mu)$ above defined coincides with the Kolmogorov-Sinai entropy $h(\mu)$ (see prop.7 in [16] and lemma 7 in [17]). If we set the *a priori* probability ν as the normalized counting measure on M (which is a probability), then $h^\nu(\mu) = h(\mu) - \log d \leq 0$.

We will call an invariant probability μ on Ω of **Gibbs probability** if there exists an a priori probability ν satisfying $\text{supp}(\nu) = M$ and a Lipschitz ν -normalized function A such that $\mathcal{L}_{(A,\nu)}^*(\mu) = \mu$. In this case the probability μ is the ν -equilibrium measure for the normalized potential A . If A is a Lipschitz function which is not normalized we can apply the construction given by (18) and we get an associated normalized potential \bar{A} (which is also Lipschitz). The probability μ is the ν -equilibrium measure for both functions A and \bar{A} .

Let $A : \Omega \rightarrow \mathbb{R}$ be a ν -normalized Lipschitz function. We will call e^A of a ν -Jacobian of the shift-invariant probability μ if for any continuous function $g : \Omega \rightarrow \mathbb{R}$ we have

$$\iint e^{A(|a,x_2,x_3,\dots\rangle)} g(|a,x_2,x_3,\dots\rangle) d\nu(a) d\mu(|x_2,x_3,\dots\rangle) = \int g(x) d\mu(x).$$

The following statements are equivalent for a ν -normalized Lipschitz function A and a shift-invariant probability μ on Ω :

- i. e^A is a ν -Jacobian of μ
- ii. μ is the ν -equilibrium measure A
- iii. $\mathcal{L}_{(A,\nu)}^*(\mu) = \mu$.

Given a Gibbs measure μ we denote by P_μ the projection of μ on the first coordinate. This means that for any continuous function $g : \Omega \rightarrow \mathbb{R}$, which depends only of the first coordinate, we have

$$\int_M g(a) dP_\mu(a) := \int_\Omega g(x_1) d\mu(|x_1,x_2,x_3,\dots\rangle).$$

The next result is a corollary of Proposition 20.

Proposition 27. *Suppose that μ is a Gibbs measure and let P_μ be the projection of μ on the first coordinate. If μ is the ν -equilibrium measure for the Lipschitz normalized potential A , then ν is equivalent to P_μ , and*

$$\frac{dP_\mu}{d\nu}(a) = \int e^{A(|a,x_1,x_2,x_3,\dots\rangle)} d\mu(|x_1,x_2,x_3,\dots\rangle),$$

which is also Lipschitz.

It is known that any Gibbs measure μ is positive on open sets of $\Omega = M^{\mathbb{N}}$ (see [7]). Therefore, the next result is a corollary of Proposition 21.

Proposition 28. *Let μ be a Gibbs measure and let ν be an a priori probability. Suppose A is a Lipschitz ν -normalized function, such that, μ is the ν -equilibrium for A , then A is the unique Lipschitz ν -Jacobian of μ .*

The next definition is inspired by (8) and (16).

Definition 29. *Let η be a shift-invariant probability and μ be a Gibbs measure. Then, we define the **specific information gain** of η with respect to μ as*

$$h(\eta, \mu) = \left[\int B d\mu + h^{P_\mu}(\mu) \right] - \left[\int B d\eta - h^{P_\mu}(\eta) \right], \quad (19)$$

where B is any Lipschitz function, such that, μ is the P_μ -equilibrium measure of B .

Observe that by the variational principle the information gain is ≥ 0 .

There exists a unique P_μ -normalized function \bar{B} , such that, μ is the P_μ -equilibrium for \bar{B} . If B is not normalized then there exists a positive function h_B and a positive number λ_B , such that,

$$\bar{B} = B + \log(h_B) - \log(h_B) \circ \sigma - \log(\lambda_B).$$

It follows that

$$\int B d\mu + h^{P_\mu}(\mu) - \int B d\eta - h^{P_\mu}(\eta) = \int \bar{B} d\mu + h^{P_\mu}(\mu) - \int \bar{B} d\eta - h^{P_\mu}(\eta).$$

This shows that $h(\eta, \mu)$ is well defined (it does not change if either B is P_μ -normalized, or not). We remark that if B is (the unique possible) normalized potential, then $\int B d\mu + h^{P_\mu}(\mu) = 0$, and, therefore, we get the following result which is a particular version of (16):

Proposition 30. *If μ is a Gibbs measure and e^B is the Lipschitz P_μ -Jacobian of μ , then*

$$h(\eta, \mu) = - \int B d\eta - h^{P_\mu}(\eta).$$

The above definition considers, for a Gibbs measure μ , the a priori probability P_μ . In this way the previous definition of specific information gain does not allow a choice of ν . The next result, which is a corollary of Proposition 24 shows that if we exchange P_μ by another a priori probability ν then it is true a similar formula for $h(\eta, \mu)$, that is, the information gain does not depend of the particular choice of ν , as long as μ is a ν -equilibrium measure.

Proposition 31. Consider any a priori probability ν and any Lipschitz function A , such that, μ is the ν -equilibrium measure for A . Let η be any invariant probability. Then, $h(\eta, \mu)$ as defined in (19), satisfies

$$h(\eta, \mu) = \left[\int A d\mu + h^\nu(\mu) \right] - \left[\int A d\eta - h^\nu(\eta) \right]. \quad (20)$$

Proof. First note that if we replace A by its normalization \bar{A} then the value on the right hand side of the above expression does not change. Then, we can suppose that A is ν -normalized. Then, it is necessary just to prove that $h(\eta, \mu) = - \left[\int A d\eta - h^\nu(\eta) \right]$. But, this follows from Proposition 24. \square

Example 32. Consider any a priori probability ν and the Lipschitz function $A = 0$. We observe that $\bar{\nu} = \nu \times \nu \times \nu \times \dots$ is the ν -equilibrium measure for $A = 0$. Given any invariant probability η , then

$$h(\eta, \bar{\nu}) = -h^\nu(\eta). \quad (21)$$

Therefore, the specific information gain generalizes the concept of relative entropy in [16].

Now we propose an interpretation of the information gain by using transfer operators defined from a priori transverse functions.

Remark 33. Let μ be a Gibbs measure and suppose that e^B is the Lipschitz ν -Jacobian of μ . Consider the identification of Ω and $X \times Y$ given by (17), and, then define an a priori transverse function on Ω by $\hat{\nu}^y(da) = e^{B(a,y)} d\nu(a)$, $y = |\cdot, x_2, x_3, x_4, \dots)$. For a fixed $\hat{\nu}$ -normalized function A let H_A be the operator acting on bounded and measurable functions $f : \Omega \rightarrow \mathbb{R}$ by

$$H_A(f)(y) = \int e^{A(a,y)} f(a, y) \hat{\nu}^y(da), \quad y = |x_1, x_2, x_3, \dots).$$

Let η be a shift-invariant probability on Ω and suppose there exists a $\hat{\nu}$ -normalized function A such that $H_A^*(\eta) = \eta$, that means, for any measurable function f we have

$$\int e^{A(a,y)} f(a, y) d\hat{\nu}^y(da) d\eta(y) = \int f(y) d\eta(y).$$

Then, $IG(\eta, \mu) = -H^{\hat{\nu}}(\eta) = \int A d\eta$. Furthermore,

$$IG(\eta, \mu) = \sup \left\{ \int c d\eta \mid c \text{ is } \hat{\nu} \text{-normalized} \right\}.$$

We finish this section by considering the exposition given in [1]. Let η, μ be two probabilities on Ω . For each $\Gamma \subset \mathbb{N}$ consider the canonical projection $\pi_\Gamma : \Omega \rightarrow M^\Gamma$ and for each $n \in \mathbb{N}$ denote by Λ_n the set $\{1, \dots, n\}$ and by \mathcal{A}_n the σ -algebra on Ω generated by the projections $\{\pi_\Gamma, \Gamma \subset \Lambda_n\}$. Denote also

$$\mathcal{H}_{\Lambda_n}(\eta | \mu) = \begin{cases} \int_{\Omega} \frac{d\eta|_{\mathcal{A}_n}}{d\mu|_{\mathcal{A}_n}} \log \left(\frac{d\eta|_{\mathcal{A}_n}}{d\mu|_{\mathcal{A}_n}} \right) d\mu, & \text{if } \eta \ll \mu \text{ on } \mathcal{A}_n \\ +\infty & \text{else} \end{cases}.$$

The next result is a consequence of Theorems 1 and 3 in [1]. From this result we get an alternative and equivalent way of extending the concept of specific information gain for the generalized XY model by considering (7) instead (8) and (16).

Proposition 34. *If μ is a Gibbs measure and η is shift-invariant on Ω , then*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \mathcal{H}_{\Lambda_n}(\eta | \mu) = h(\eta, \mu).$$

5 The Involution kernel and the Entropy production in the generalized XY model

In the same way as in last section we assume that M is a compact metric space. We denote by Ω^- the space $M^{\mathbb{N}}$ with elements written in the form $(\dots, y_3, y_2, y_1 |, y_i \in M$, and with the same metric as the one previously defined in $\Omega = \Omega^+$.

Points in $\hat{\Omega} = \Omega^- \times \Omega^+$ are written in the form $(\dots, y_3, y_2, y_1 | x_1, x_2, x_3, \dots)$. The shift map $\hat{\sigma} : \hat{\Omega} \rightarrow \hat{\Omega}$ was defined by (11). The restrictions of $\hat{\sigma}$ to Ω^+ and Ω^- are denoted, respectively, by σ and σ^- .

Observe that (Ω^-, σ^-) can be identified with (Ω, σ) from the conjugation $\theta : \Omega^- \rightarrow \Omega$, given by $\theta((\dots, z_3, z_2, z_1 |) = |z_1, z_2, z_3, \dots)$. Using this conjugation any result previously stated for (Ω, σ) has an analogous claim for (Ω^-, σ^-) .

Consider a Lipschitz function $A : \Omega^- \times \Omega \rightarrow \mathbb{R}$, which does not depend of $y \in \Omega^-$. Then, it is naturally expressed as $A(|x_1, x_2, x_3, \dots)$. One can show that there exists a (several, in fact) Lipschitz function $W : \Omega^- \times \Omega \rightarrow \mathbb{R}$, which is called an **involution kernel**, and a Lipschitz function A^- , such that

$$A^- := A \circ \hat{\sigma}^{-1} + W \circ \sigma^{-1} - W, \quad (22)$$

where the function A^- **does not depend on** $x \in \Omega$ (see [2],[16]). A^- is naturally expressed in coordinates $(\dots, y_3, y_2, y_1 |$.

All this can be written in the form:

$$A^-(\dots, y_3, y_2, y_1|) = A(|y_1, x_1, x_2, x_3, \dots) + W(\dots, y_3, y_2|y_1, x_1, x_2, x_3, \dots) - W(\dots, y_3, y_2, y_1|x_1, x_2, x_3, \dots), \quad (23)$$

for any $(\dots, y_3, y_2, y_1|x_1, x_2, x_3, \dots) \in \hat{\Omega}$.

Following [2] and [16] we state two propositions.

Proposition 35. *Let $A : \Omega^+ \rightarrow \mathbb{R}$ be a Lipschitz function and $W : \hat{\Omega} \rightarrow \mathbb{R}$ be a Lipschitz involution kernel for A . Consider the function A^- which was defined by (22). Fix an a priori probability ν on M . Then, for any $x \in \Omega^+$, $y \in \Omega^-$, and any function $f : \hat{\Omega} \rightarrow \mathbb{R}$,*

$$\mathcal{L}_{A^-, \nu}(f(\cdot|x) e^{W(\cdot|x)})(y) = \mathcal{L}_{A, \nu}(f \circ \hat{\sigma}(y|\cdot) e^{W(y|\cdot)})(x). \quad (24)$$

Proposition 36. *Let $A : \Omega^+ \rightarrow \mathbb{R}$ be a Lipschitz function and $W : \hat{\Omega} \rightarrow \mathbb{R}$ be a Lipschitz involution kernel for A . Consider the function A^- as defined by (22). Fix an a priori probability ν on M . Let ρ_A and ρ_{A^-} be the eigenmeasures for $\mathcal{L}_{A, \nu}^*$ and $\mathcal{L}_{A^-, \nu}^*$, respectively. Suppose c is such that $\iint e^{W(y|x)-c} d\rho_{A^-}(y) d\rho_A(x) = 1$ and denote $K(y|x) := e^{W(y|x)-c}$. Then,*

1. *The probability*

$$d\hat{\mu}_A = K(y|x) d\rho_{A^-}(y) d\rho_A(x)$$

is invariant for $\hat{\sigma}$ and it is an extension of the ν -equilibrium measure μ_A .

2. *The function $h_A(x) = \int K(y|x) d\rho_{A^-}(y)$ is the main eigenfunction for $\mathcal{L}_{A, \nu}$, and the function $h_{A^-}(y) = \int K(y|x) d\rho_A(x)$ is the main eigenfunction for $\mathcal{L}_{A^-, \nu}$.*

3. $\lambda_A = \lambda_{A^-}$.

Now we apply these results to the study of the entropy production. We start by refining item 1. of the last proposition.

Proposition 37. *The probability $d\hat{\mu}_A = K(y|x) d\rho_{A^-}(y) d\rho_A(x)$ is the unique $\hat{\sigma}$ -invariant extension to $\hat{\Omega}$ of the equilibrium measure μ_A on Ω .*

Proof. Let $\hat{\mu}$ be any $\hat{\sigma}$ -invariant probability on $\hat{\Omega}$ satisfying $\int g d\hat{\mu} = \int g d\mu_A$, when $g(y|x)$ does not depend of y . Consider any continuous function f on $\hat{\Omega}$. We claim that $\int f d\hat{\mu} = \int f d\hat{\mu}_A$. Indeed, as $\hat{\Omega}$ is compact, the function f is uniformly continuous. Fix any point $y_0 \in M$ and define the functions f_n on $\hat{\Omega}$, $n \in \mathbb{N}$, by $f_n(y|x) = f(y^n|x)$, where $y^n = (\dots, y_0, y_0, y_0, y_n, y_{n-1}, \dots, y_2, y_1|$.

It follows that $\{f_n\}$ converges uniformly to f , and moreover, the function $f_n((\dots, y_3, y_2, y_1|x_1, x_2, x_3, \dots))$ does not depend of y_k , for $k > n$.

From,

$$\int f_n d\hat{\mu} = \int f_n \circ \hat{\sigma}^{-n} d\hat{\mu} = \int f_n \circ \hat{\sigma}^{-n} d\mu_A = \int f_n \circ \hat{\sigma}^{-n} d\hat{\mu}_A = \int f_n d\hat{\mu}_A,$$

we conclude that $\int f d\hat{\mu} = \int f d\hat{\mu}_A$. \square

Notation 38. Let μ be a Gibbs measure on Ω^+ . We denote by $\hat{\mu}$ the unique $\hat{\sigma}$ -invariant extension to $\hat{\Omega}$ of μ and by μ^- the restriction of $\hat{\mu}$ to Ω^- .

Proposition 39. Let $A : \Omega^+ \rightarrow \mathbb{R}$ be a Lipschitz function, W be any Lipschitz involution kernel for A . Now, consider the function A^- on Ω^- as defined by (22). Fix an a priori probability ν on M . Let μ_A be the ν -equilibrium of A and let $(\mu_A)^-$ defined as above. Then, $(\mu_A)^-$ is the ν -equilibrium of A^- in Ω^- , that is

$$(\mu_A)^- = \mu_{(A^-)}.$$

Proof. From the above

$$d\hat{\mu}_A = K(y|x) d\rho_{A^-}(y) d\rho_A(x),$$

and $h_{A^-}(y) = \int K(y|x) d\rho_A(x)$ is the main eigenfunction for $\mathcal{L}_{A^-, \nu}$. Then, for any continuous function $f : \Omega^- \rightarrow \mathbb{R}$ we get

$$\begin{aligned} \int f(y) d(\mu_A)^- &= \int f(y) d\hat{\mu}_A = \iint f(y) K(y|x) d\rho_A(x) d\rho_{A^-}(y) \\ &= \int f(y) h_{A^-}(y) d\rho_{A^-}(y) = \int f(y) d\mu_{A^-}. \end{aligned}$$

\square

Definition 40. The entropy production of the Gibbs measure μ is defined as

$$e_p(\mu) = h(\mu, \theta_*\mu^-),$$

where $\theta_*\mu^-$ on Ω^+ is the push-forward of μ^- by the conjugation $\theta : \Omega^- \rightarrow \Omega^+$ given by (12).

Observe that as a consequence of the variational principle we get $e_p(\mu) \geq 0$, and it is zero, if and only if, $\mu^- = \mu$. As the specific information gain $h(\mu, \mu^-)$ “does not depend of ν ”, the above definition also “does not depend of ν ”. In fact, by definition we should have to consider the a priori probability P_{μ^-} , but, if for some a priori probability ν the measure μ is a ν -equilibrium measure, then get that the probability μ^- also is. Now, applying Proposition 31 we get an alternative formula for computing the same expression $e_p(\mu)$,

but now using the *a priori* probability ν . From now on we will exhibit other alternative ways for computing the entropy production.

Let $\hat{\theta} : \hat{\Omega} \rightarrow \hat{\Omega}$ be given by

$$\hat{\theta}(\dots, y_3, y_2, y_1 | x_1, x_2, x_3, \dots) = (\dots x_3, x_2, x_1 | y_1, y_2, y_3, \dots). \quad (25)$$

Observe that $\hat{\theta}^{-1} = \hat{\theta}$ and $\hat{\theta} \circ \hat{\sigma}^{-1} = \hat{\sigma} \circ \hat{\theta}$.

Proposition 41. *Let $A : \Omega \rightarrow \mathbb{R}$ be a Lipschitz function, $W : \hat{\Omega} \rightarrow \mathbb{R}$ be any Lipschitz involution kernel for A and let $A^- : \Omega^- \rightarrow \mathbb{R}$ be defined by (22). Let μ be any Gibbs measure on Ω and consider $\hat{\mu}$ and μ^- defined as above. Then,*

1. $\int A d\mu = \int A^- d\mu^-$
2. $\int A^- \circ \theta^{-1} d\mu = \int A \circ \theta d\mu^-$.

Proof. In order to prove item 1. we observe that

$$\int A d\mu = \int A d\hat{\mu} = \int A \circ \sigma^{-1} + W \circ \hat{\sigma}^{-1} - W d\hat{\mu} = \int A^- d\hat{\mu} = \int A^- d\mu^-.$$

Now we will prove item 2.

$$\begin{aligned} \int A^- \circ \theta^{-1} d\mu &= \int A^- \circ \hat{\theta} d\hat{\mu} = \int A \circ \hat{\sigma}^{-1} \circ \hat{\theta} + W \circ \hat{\sigma}^{-1} \circ \hat{\theta} - W \circ \hat{\theta} d\hat{\mu} \\ &= \int A \circ \hat{\theta} \circ \hat{\sigma} + W \circ \hat{\theta} \circ \hat{\sigma} - W \circ \hat{\theta} d\hat{\mu} = \int A \circ \hat{\theta} d\hat{\mu} = \int A \circ \theta d\mu^-. \end{aligned}$$

□

Proposition 42. *Let μ be a Gibbs measure and ν be an *a priori* probability. Then, $h^\nu(\mu) = h^\nu(\mu^-)$.*

Proof. For each Lipschitz function $A : \Omega^+ \rightarrow \mathbb{R}$ we can consider a Lipschitz involution kernel W , and then, we get an associated Lipschitz function $A^- : \Omega^- \rightarrow \mathbb{R}$.

For the fixed *a priori* probability ν we have $\lambda_A = \lambda_{A^-}$. Then,

$$\begin{aligned} h^\nu(\mu) &= - \sup_{A \text{ is } \nu\text{-normalized}} \int A d\mu = - \sup_{A \text{ is Lipschitz on } \Omega^+} \int A d\mu - \log(\lambda_A) \\ &= - \sup_{A^- \text{ given from some Lip. } A^+} \int A^- d\mu^- - \log(\lambda_{A^-}) \end{aligned}$$

$$\geq - \sup_{B^- \text{ is Lipschitz on } \Omega^-} \int B^- d\mu^- - \log(\lambda_{B^-}) = h^\nu(\mu^-).$$

In order to get the opposite inequality we follow a similar argument. We exchange the reasoning by $\hat{\theta}$: for each Lipschitz function $B^- : \Omega^- \rightarrow \mathbb{R}$, we take an involution kernel, and, an associated Lipschitz function $B^+ : \Omega^+ \rightarrow \mathbb{R}$. Now, we just have to proceed in the same way as before. \square

As a consequence we get the following claim for the entropy production:

Proposition 43. *Suppose that μ is a Gibbs measure and consider the associated probability μ^- . Suppose that for an a priori probability ν and for a Lipschitz function A^- we have that μ^- is the ν -equilibrium measure for A^- . Now, assume that μ^- and A^- are defined on Ω^+ via the conjugation θ . Then, the entropy production of μ satisfies*

$$e_p(\mu) = \int A^- d\mu^- - \int A^- d\mu.$$

We can take A^- such that $J^- = e^{A^-}$ is the ν -Jacobian of μ^- .

Proposition 44. *Suppose that μ is the ν -equilibrium measure for the Lipschitz function $A : \Omega^+ \rightarrow \mathbb{R}$. Let W be any Lipschitz involution kernel for A and $A^- : \Omega^- \rightarrow \mathbb{R}$ be the function defined by (22). Suppose that A^- is defined on Ω^+ using the conjugation θ . Then,*

$$e_p(\mu) = \int A - A^- d\mu.$$

Proof. It follows from the previous result and Proposition 41. \square

Proposition 45. *Suppose that μ is a Gibbs probability. Then,*

$$e_p(\mu) = h(\mu, \mu^-) = h(\mu^-, \mu) = e_p(\mu^-).$$

Proof. It follows from Proposition 36 and 37 that $(\mu^-)^- = \mu$. Consider an a priori probability ν , such that, μ is the ν -equilibrium measure for a Lipschitz function A . Let A^- defined by (22) using any involution kernel. From Proposition 43 we get

$$e_p(\mu) = h(\mu, \mu^-) = \int A^- d\mu^- - \int A^- \circ \theta^{-1} d\mu$$

and

$$e_p(\mu^-) = h(\mu^-, \mu) = \int A d\mu - \int A \circ \theta d\mu^-.$$

Now, from proposition 41 we get

$$\int A^- d\mu^- - \int A^- \circ \theta^{-1} d\mu = \int A d\mu - \int A \circ \theta d\mu^-.$$

This concludes the proof. \square

The next example consider the more simple case where $M = \{1, 2, \dots, d\}$ is a finite set.

Example 46. Take $M = \{1, 2, \dots, d\}$ and take as the a priori measure ν the counting measure on M .

Any invariant probability μ for (Ω, σ) can be extended to a $\hat{\sigma}$ -invariant probability $\hat{\mu}$ on $\hat{\Omega}$ by defining

$$\hat{\mu}([a_m, \dots, a_1 | b_1, \dots, b_n]) := \mu(|a_m, \dots, a_1, b_1, \dots, b_n]),$$

and using the extension theorem. The restriction of $\hat{\mu}$ to Ω^- satisfies

$$\mu^-([a_m, \dots, a_2, a_1 |]) = \mu(|a_m, \dots, a_2, a_1]).$$

Now, using the conjugation $\theta : \Omega^- \rightarrow \Omega$ in order to transfer μ^- to Ω^+ , we get

$$\theta_* \mu^-([a_1, a_2 \dots a_m |]) = \mu(|a_m, \dots, a_2, a_1]). \quad (26)$$

As the Kolmogorov-Sinai entropy of μ is given by

$$h(\mu) = \lim_{n \rightarrow \infty} -\frac{1}{n} \sum_{i_1, \dots, i_n} \mu(|i_1, \dots, i_n]) \log(\mu(|i_1, \dots, i_n])),$$

we conclude that $h(\mu) = h(\mu^-)$.

Suppose now, that μ is the equilibrium measure for the Lipschitz normalized potential A . Then, $e^A = J$ is the Jacobian of μ , that is,

$$e^A(|x_1, x_2, x_3, \dots>) = J(|x_1, x_2, x_3, \dots>) = \lim_{n \rightarrow \infty} \frac{\mu(|x_1, x_2, x_3, \dots, x_n])}{\mu(|x_2, x_3, \dots, x_n])}.$$

Let J^- be the Jacobian of μ^- and define $A^- := \log(J^-)$. Then, using (26),

$$e^{A^-}(\dots, y_3, y_2, y_1 |) = J^- (\dots, y_3, y_2, y_1 |) = \lim_{n \rightarrow \infty} \frac{\mu(|y_n, \dots, y_2, y_1])}{\mu(|y_n, \dots, y_2])}.$$

The next example compute the entropy production for a Markov measure μ . Our estimate is coherent with expression (1) in [13].

Example 47. Consider the line stochastic matrix $M = (p_{ij})$ and the initial probability vector $P = (\pi_i)$, such that, $PM = P$.

We denote by μ the associated Markov measure, that is, for any cylinder $[x_1, x_2, \dots, x_n]$ we set

$$\mu([x_1, x_2, \dots, x_n]) = \pi_{x_1} \cdot p_{x_1 x_2} \cdots p_{x_{n-1} x_n}.$$

Then,

$$J([i, j, x_3, \dots]) = \frac{\pi_i p_{ij}}{\pi_j}.$$

We also get

$$J^-(\dots, y_3, j, i) = \lim_{n \rightarrow \infty} \frac{\mu([y_n, \dots, y_3, j, i])}{\mu([y_n, \dots, y_3, j])} = \lim_{n \rightarrow \infty} \frac{\pi_{y_n} \cdot p_{y_n y_{n-1}} \cdots p_{y_3 j} \cdot p_{ji}}{\pi_{y_n} \cdot p_{y_n y_{n-1}} \cdots p_{y_3 j}} = p_{ji}.$$

As J^- depends only of two coordinates, μ^- is also a Markov measure.

Considering the conjugation θ , we get,

$$\mu([i, j]) = \pi_i p_{ij} \text{ and } \mu^-([i, j]) = \mu([j, i]) = \pi_j p_{ji}.$$

Taking $A = \log(J)$ and $A^- = \log(J^-)$, we also get

$$e^{A([i, j, x_3, \dots])} = \frac{\pi_i p_{ij}}{\pi_j} \text{ and } e^{A^-([i, j, z_3, z_4, \dots])} = p_{ji}.$$

Then, using the Proposition 44, we derive

$$e_p(\mu) = \int A - A^- d\mu = \sum_{i, j} \log\left(\frac{\pi_i p_{ij}}{\pi_j p_{ji}}\right) \pi_i p_{ij}.$$

We can compute $e_p(\mu)$ alternatively using Proposition 43:

$$\begin{aligned} e_p(\mu) &= \int A^- d\mu^- - \int A^- d\mu = \sum_{i, j} \log(p_{ji}) \mu^-([i, j]) - \sum_{i, j} \log(p_{ji}) \mu([i, j]) \\ &= \sum_{i, j} \log(p_{ji}) \pi_j p_{ji} - \sum_{i, j} \log(p_{ji}) \pi_i p_{ij} = \sum_{i, j} \log(p_{ij}) \pi_i p_{ij} - \sum_{i, j} \log(p_{ji}) \pi_i p_{ij} \\ &= \sum_{i, j} \log(p_{ij}) \pi_i p_{ij} - \sum_{i, j} \log(p_{ji}) \pi_i p_{ij} + \left[\sum_i \pi_i \log(\pi_i) - \sum_j \pi_j \log(\pi_j) \right] \end{aligned}$$

$$\begin{aligned}
&= \sum_{i,j} \log(p_{ij})\pi_i p_{ij} - \sum_{i,j} \log(p_{ji})\pi_i p_{ij} + \left[\sum_{i,j} \pi_i p_{ij} \log(\pi_i) - \sum_{i,j} \pi_i p_{ij} \log(\pi_j) \right] \\
&= \sum_{i,j} \log\left(\frac{\pi_i p_{ij}}{\pi_j p_{ji}}\right) \pi_i p_{ij}.
\end{aligned}$$

The case with just two symbols is quite special as we will see now.

Example 48. *Entropy production zero - Suppose $\Omega = \{1, 2\}^{\mathbb{N}}$ and assume that μ is a Markov measure (as defined above). Then, $e_p(\mu) = 0$. Indeed, as μ is invariant we get $\mu(|1, 2)) = \mu(|2, 1))$, and therefore, $\mu^-(|i, j)) = \mu(|j, i)) = \mu(|i, j)$, for any $i, j \in \{1, 2\}$. It follows that $J^- = J^+$, and therefore, $\mu^- = \mu$. Consequently,*

$$e_p(\mu) = \int \log(J) - \log(J^-) d\mu = 0.$$

That is, the entropy production is zero.

It follows from Corollary 2.3 in [28] that this result - entropy production zero - also happen for equilibrium probabilities of a more general class of functions defined on $\Omega = \{1, 2\}^{\mathbb{N}}$ (see [28]).

The probability described in [19] also has entropy production zero (see section 2 in [19]).

Markov measures on $\Omega = \{1, 2, 3\}^{\mathbb{N}}$ may have non zero entropy production.

6 Appendix: Examples in information theory

This section is intended to illustrate the theoretical results we previously described concerning Shannon entropy $S(P)$ and information gain $IG(\pi, P)$ via examples taken from Information Theory (a nice general reference on the topic is [8]). We believe that this short presentation will be helpful for mathematicians that do not have much familiarity with these concepts.

We start by considering the Shannon entropy which is called alternatively of mean information.

Example 49. *Suppose that a box has balls of 4 possible different colors. Two people will play a game with the following rules: one ball is picked of the box by one of them and the other person must discover the color of this ball by making questions of the type “yes or not”.*

If this game is repeated several times, the balls are picked randomly according with the probability $P = (p_1, p_2, p_3, p_4) = (\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4})$ and the strategy

used for the questions is optimal, what is the mean value of the number of questions which are necessary?

We will replace the colors by symbols of the set $\{1, 2, 3, 4\}$. One can consider the following strategy of questions:

Q1: is the picked symbol 1 or 2?

- with the answer "yes" it can be considered the question Q2: is the symbol 1?

- with the answer "no" it can be considered the question Q2': is the symbol 3?

Using this strategy it is necessary exactly two questions in order to discover the picked symbol (color). It coincides with the Shannon entropy (the mean information)

$$S(P) = - \sum_{i=1}^4 \frac{1}{4} \log_2\left(\frac{1}{4}\right) = 2.$$

Observe that the set of symbols $\{1, 2, 3, 4\}$ can be encoded as the answers (yy, yn, ny, nn) . Replacing y by 0 and n by 1 we can encode $\{1, 2, 3, 4\}$ as $(00, 01, 10, 11)$ in binary expansion, which is optimal.

Example 50. Proceeding as in above example, but now assuming that the colors of the balls are picked randomly according to the probability $P = (p_1, p_2, p_3, p_4) = (\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{8})$, one can use the following strategy of questions: Q1: is the symbol (color) 1? (with probability (frequency) $\frac{1}{2}$ this unique question solves the problem)

- with the answer "yes" we finish.

- with the answer "no" we consider the question Q2: is the symbol 2?

- with the answer "yes" we finish.

- with the answer "no" again, we then consider the question Q3: is the symbol 3?

If this game is repeated several times, using this strategy the mean number of questions is:

$$(1 \text{ question})\frac{1}{2} + (2 \text{ questions})\frac{1}{4} + (3 \text{ questions})\frac{1}{4} = \frac{7}{4}.$$

It coincides with the Shannon entropy (mean information)

$$S(P) = -\left[\frac{1}{2} \log_2\left(\frac{1}{2}\right) + \frac{1}{4} \log_2\left(\frac{1}{4}\right) + \frac{1}{8} \log_2\left(\frac{1}{8}\right) + \frac{1}{8} \log_2\left(\frac{1}{8}\right)\right] = \frac{7}{4}.$$

In this case $\{1, 2, 3, 4\}$ can be encoded as $\{0, 10, 110, 111\}$ in binary expansion, being this one optimal.

Example 51. Proceeding as above and supposing that there are only two colors of balls which are picked randomly according with the probability $P = (p_1, p_2) = (\frac{2}{3}, \frac{1}{3})$ one can consider the following question:

Q1: is the color (symbol) 1?

With this strategy the mean number of questions is exactly 1 which is bigger than the Shannon entropy $S(P) \approx 0,918$. In this case $\{1, 2\}$ can be encoded as $\{0, 1\}$ in binary expansion.

We refer to [8] chap. 5 for a more complete discussion of the topic. Our intention above was just to illustrate - with introductory and simple examples - the fact that the Shannon entropy is as a lower bound for the average number of questions and how one can introduce an binary code for a set of symbols $\{1, \dots, d\}$.

From now we will discuss the concept of the Information Gain $IG(\pi, P)$. We refer to [24] (see p. 89-90) for a more detailed discussion of this topic in the context of Information Theory.

Example 52. Consider - in a similar way as before - a box with a collection of 100 objects being 30 of them of the color blue and 70 of them of the color red. It's also known that:

- a. 10 of the blue objects are balls and 20 of them are cubes
- b. 45 of the red objects are balls and 25 of them are cubes.

Considering all this set of information we can construct probabilities P and π in the following way:

$$\begin{array}{l} 30 \text{ blue} \\ 70 \text{ red} \end{array} \rightarrow P = \begin{pmatrix} 0.3 \\ 0.7 \end{pmatrix} \quad \begin{array}{cc} & \begin{array}{cc} \text{balls} & \text{cubes} \end{array} \\ \begin{array}{c} \text{blue} \\ \text{red} \end{array} & \begin{pmatrix} 10 & 20 \\ 45 & 25 \end{pmatrix} \end{array} \rightarrow \pi = \begin{pmatrix} 0.10 & 0.20 \\ 0.45 & 0.25 \end{pmatrix}.$$

We consider that π is defined in a Cartesian product $X \times Y$ and has x -marginal $P = (\frac{30}{100}, \frac{70}{100})$ (adding in the lines of π) and y -marginal $Q = (\frac{55}{100}, \frac{45}{100})$ (adding in the rows of π).

We will consider two kind of different games.

Game one: One object is randomly picked of the box and we shall discover its color by asking questions of the type yes or no. In this case the Shannon's entropy, or mean information, is equal to

$$S(P) = - \left[\frac{30}{100} \log\left(\frac{30}{100}\right) + \frac{70}{100} \log\left(\frac{70}{100}\right) \right].$$

Game two: In this game - in a similar way as in game one - we have the same goal. However, in the present game, after the object was picked we receive a partial information about the result, which is: “it is a cube” or “it is a ball”.

In this game, with probability (or, frequency) $\frac{55}{100}$, the information to be received it will be that it was picked a ball. Using this information we must concentrate our attention for such class of objects and so the colors are distributed according to the probability $(\frac{10}{55}, \frac{45}{55})$. Similarly, with probability (frequency) $\frac{45}{100}$, the information received will be that a cube was picked. In this case, we consider the colors distributed according to the probability $(\frac{20}{45}, \frac{25}{45})$. Therefore, the mean information in this game is given by a weighted mean of two Shannon’s entropies, that is,

$$H(\pi) = -\frac{55}{100} \left[\frac{10}{55} \log\left(\frac{10}{55}\right) + \frac{45}{55} \log\left(\frac{45}{55}\right) \right] - \frac{45}{100} \left[\frac{20}{45} \log\left(\frac{20}{45}\right) + \frac{25}{45} \log\left(\frac{25}{45}\right) \right].$$

Finally, we observe that the information gain $IG(\pi, P)$ given in (1) is the difference between the mean information in game one and the mean information in game two,

$$IG(\pi, P) = S(P) - H(\pi).$$

A. O. Lopes partially supported by CNPq grant.

References

- [1] D. Aguiar, L. Cioletti, and R. Ruviano, A variational principle for the specific entropy for symbolic systems with uncountable alphabets, *Math. Nachr.* 291 no. 17-18, 2506-2525 (2018)
- [2] A. Baraviera, A. Lopes and Ph. Thiullen, A large deviation principle for equilibrium states of Hölder potentials: the zero temperature case, *Stochastics and Dynamics* **6** 77-96 (2006)
- [3] T. Benoist, V. Jaksic, Y. Pautrat and C-A. Pillet, On entropy production of repeated quantum measurements I. General theory, *Comm. Math. Phys.* 357, no. 1, 77-123 (2018)
- [4] M. Capinski and E. Kopp, *Measure Integral and Probability*, Springer-Verlag (2004).

- [5] J-R. Chazottes, J.-R. E. Floriani and R. Lima, Relative entropy and identification of Gibbs measures in dynamical systems, *J. Statist. Phys.* 90, no. 3-4 679-725 (1998)
- [6] J-R. Chazottes and Eric Olivier, Relative entropy, dimensions and large deviations for g-measures, *J. Phys. A* 33, no. 4, 675-689 (2000)
- [7] L. Cioletti and L. C. Melo, On the extensions of generalized transfer operators, to appear - preprint 2020, UNB
- [8] T. Cover and J. Thomas, *Elements of information theory*, 2 ed, Wiley-Interscience (2006)
- [9] G. Crooks, Entropy production fluctuation theorem and the nonequilibrium work relation for free energy differences, *Phys. Rev. E*, 60, 2721 (1999)
- [10] I. Ekeland and R. Témam, *Convex Analysis and Variational Problems*, North-Holland (1976)
- [11] G. Gallavotti and E. G. D. Cohen, Dynamical Ensembles in Nonequilibrium Statistical Mechanics, *Phys. Rev. Lett.* 74, 2694 (1995)
- [12] H-O. Georgii. *Gibbs measures and phase transitions*, Ed. Gruyter, second edition, (2011).
- [13] Da-quan Jiang, Min Qian, Min-ping Qian, Entropy Production and Information Gain in Axiom-A Systems *Commun. Math. Phys.*, 214, 389 - 409 (2000)
- [14] S. Kullback and R. A. Leibler, On Information and Sufficiency. *Ann. Math. Statist.* 22, no. 1, 79-86 (1951).
- [15] A. O. Lopes and J. K. Mengue, Thermodynamic Formalism for Haar systems in Noncommutative Integration: transverse functions and entropy of transverse measures (to appear in: *Erg. theo. Dyn. Sys.*) (arXiv:1905.09161)
- [16] A. O. Lopes, J. K. Mengue, J. Mohr and R. R. Souza, Entropy and Variational Principle for one-dimensional Lattice Systems with a general a priori probability: positive and zero temperature, *Ergodic Theory and Dynamical Systems*, 35 (6), 1925-1961 (2015)

- [17] A. Lopes, J. Mengue, J. Mohr and R. Souza, Entropy, Pressure and Duality for Gibbs plans in Ergodic transport, *Bull. Braz. Math. Soc.* Vol. 46, Issue 3, 353-389 (2015)
- [18] A. O. Lopes, E. Oliveira and Ph. Thioullien, The Dual Potential, the involution kernel and Transport in Ergodic Optimization, Dynamics, Games and Science -International Conference and Advanced School Planet Earth DGS II, Portugal (2013), Edit. J-P Bourguignon, R. Jelstch, A. Pinto and M. Viana, Springer Verlag, pp 357-398 (2015)
- [19] A. O. Lopes, J. K. Mengue, J. Mohr and C. G. Moreira, Large Deviations for Quantum Spin probabilities at temperature zero, *Stochastics and Dynamics*, Vol. 18, No. 06, 1850044 (2018)
- [20] C. Maes, The fluctuation theorem as a Gibbs property. *J. Statist. Phys.* 95 (1999), no. 1-2, 367-392
- [21] J. Mengue, Tópicos de álgebra linear e probabilidade, SBM (2016).
- [22] J. Mengue and E. Oliveira, Duality results for iterated function systems with a general family of branches, *Stochastics and Dynamics*, Vol. 17, No. 03, 1750021 (2017).
- [23] W. Parry and M. Pollicott. Zeta functions and the periodic orbit structure of hyperbolic dynamics, *Astérisque*, Vol 187-188, pp 1-268 (1990)
- [24] J. R. Quinlan, Induction of decision trees, *Machine learning*, vol 1. issue 1., 81 - 106, (1986).
- [25] D. Ruelle, A generalized detailed balance relation, *J. Stat. Phys.* 164, no. 3, 463-471 (2016)
- [26] C. E. Shannon, A Mathematical Theory of Communication, *Bell System Technical Journal* Vol. 27 Issue 3, 379-423 (1948).
- [27] M. Viana and K. Oliveira, Foundations of Ergodic Theory, Cambridge Press (2016)
- [28] P. Walters, A natural space of functions for the Ruelle operator theorem, *Ergodic Theory and Dynamical Systems*, 27, 1323–1348, (2007)
- [29] P. Walters, An introduction to Ergodic Theory, Springer Verlag, (2000)