Magnitude and significance of the peak of early embryonic mortality

Qinghua Chen¹, Zengru Di², Eduardo M. Garcia-Roger³, Hui Li⁴, Peter Richmond⁵,

Bertrand M. Roehner⁶

Version of 25 February 2020

Key-words: Embryogenesis, death rate, conception, infant death, senescence

School of Systems Science, Beijing Normal University, China.
Email: qinghuachen@bnu.edu.cn
School of Systems Science, Beijing Normal University, China.
Email: zdi@bnu.edu.cn
Institut Cavanilles de Biodiversitat I Biologia Evolutiva, University of València, Spain.
Email: eduardo.garcia@uv.es
School of Systems Science, Beijing Normal University, China.
Email: huili@bnu.edu.cn
School of Physics, Trinity College Dublin, Ireland.
Email: peter_richmond@ymail.com
Institute for Theoretical and High Energy Physics (LPTHE), Pierre and Marie Curie campus, Sorbonne University, National Center for Scientific Research (CNRS), Paris, France.
Email: roehner@lpthe.jussieu.fr

Abstract

Biologically, for any organism life does not start at birth but at fertilization of the embryo. Embryonic development is of great importance because it determines congenital anomalies and influences their severity. Whereas there is detailed qualitative knowledge of the successive steps of embryonic development, little is known about their probabilities of success or failure. Embryonic mortality as a function of post fertilization time provides a simple (albeit crude) way to identify major defects. We find that, in line with the few other species for which data are available, the embryonic mortality of zebrafish has a prominent peak shortly after fertilization. This is called the early embryonic mortality (EEM) effect. Although a number of immediate causes of death (e.g. infection, excess of carbon dioxide or of lactic acid, chromosomal defects) can be cited, the common underlying factor remains unknown.

After reviewing embryonic mortality data available for chicken and a few other farm animals, we explain that zebrafish are particularly suited for such a study because embryogenesis can be followed from its very beginning and can be observed easily thanks to transparent egg shells.

We report the following findings. (i) The mortality peak occurs in the first 15% of the 75-80 hours of embryogenesis and it is about 50 times higher than the low plateau which follows. (ii) The shape of the age-specific death rate is largely independent of the death level.

Presently, little is known about the nature of embryonic defects. However, by reviewing two special cases we show that even small initial defects, e.g. spatial cellular asymmetries or irregularities in the timing of development, carry with them lethal effects in later stages of embryogenesis.

2

Introduction

From defects in embryonic cell organization to lethality

The purpose of statistical physics is to derive the behavior of macroscopic systems from the collective properties of its microscopic components. The present investigation gives a striking illustration for it shows a massive mortality surge in early embryonic development which comes along with slight disturbances at cellular level. More specifically, investigations by two teams of researchers (Rideout et al. 2004, Cruz et al. 2012) have shown that anomalies in initial cell organization are early signals of a compromised (and eventually lethal) development. It is interesting to note that in these two cases (to be explained more fully later on) the defects are not due to individual cells but rather to defects in their interaction. Note that the defect considered in Rideout et al. is a spatial asymmetry that is fairly similar to positional defects in a crystallized solid.

In line with some of our previous papers (Berrut et al. 2016, Richmond et al. 2019a,b, Bois et al. 2019b) this study could have been entitled "The physics of embryogenesis" because all along we keep in mind possible parallels with the working and failures of physical (i.e. non living) systems; such parallels serve as guides for the exploration and suggest possible conjectures. We do not try to give a detailed description for a single species but focus on common features across species for it appears that similar peaks occur in many species; in short, we study embryonic mortality as a global effect.

So far, limited mortality data are available. Those recorded in our experiment for zebrafish embryos may be the most accurate ever produced. There are two reasons for that. The experiment was especially designed for the observation of early embryonic mortality (whereas previous experiments with farm animals had mostly an economic goal) and it involved a sample of over 1,500 embryos. It is difficult to have samples of this size for bigger animals like cows or pigs.

The early embryonic spike is not the only mortality peak to occur in the development of an organism. In order to put it in a broader perspective, in the following subsections we briefly describe the other peaks.

From fertilization to death there are three phases

In previous papers a parallel was introduced between living and technical systems (Berrut et al. 2016, Bois et al. 2019a,b). From this perspective three phases can be distinguished in human development.

(1) Embryogenesis from fertilization of the embryo to birth.




Fig. 1 Evolution of the death rate in the three phases of human development. In each phase there is a huge peak. Whereas its existence is easy to understand in phase 2 and 3, the reason of the early embryonic mortality (EEM) peak remains a mystery. In phase 2 the decrease follows a hyperbolic power law whereas in phase 3 the increase is exponential as described by Gompertz's law (1825); in phase 1 the exact shape of the curve is not yet known due to a lack of accuracy in empirical evidence, in particular in the first few weeks following fertilization. The expression "wear-out phase" is borrowed from reliability engineering; it is justified by the fact that for many functions the best performance occurs between the age of 20 and 30 and then deteriorates steadily. As examples one can mention the ability to perceive sounds of high frequency or the density of bones. In most activities of common life this wear-out remains unnoticed and becomes a hindrance only in old age. Note that the maximum death rate of the senescence curve depends upon the the oldest age group selected. A similar observation applies to the infant curve; in this case inclusion of age groups close to birth would lift up the maximum of the death rate. The data for phase 1 are for the island of Kauai in Hawaii in the late 1950s. The data for phase 2 are for the United States in 1960. The data for phase 3 are for the United States in 2016. The insets of panel 2 and 3 are (log,log) and (lin,log) respectively. Sources: Embryo phase: French et al. 1962, Léridon 1973, 1977, Jarvis 2017. Infant phase: The data from zero to 12 years are presented in two different tables (a) Under one year: Grove et al. 1968, p.210-211, (b) Over one year: Grove et al. 1968, p.318. Wear-out phase: WONDER database of the "Centers for Disease Control" (CDC), Compressed mortality, 1999-2016.

(2) From birth to 10 year old there is the infant phase¹, also known as a wear-in phase.

(3) After the age of 10 year and until senescence and death comes a phase during which the death rate increases exponentially in accordance with the well-known Gompertz's law (Gompertz 1825, Richmond et al. 2016). In the language of reliability engineering such a steady increase of the failure rate is known as a wear-out process.

Human age-specific curves for the three phases are given in Fig.1. For the sake of brevity in what follows the three phases will be referred to as phases 1, 2 and 3. Whereas there are accurate data for phases 2 and 3, embryonic death rates, and especially the early embryonic rates are subject to a number of uncertainties as empha-

¹The medical definition of infant mortality is mortality of newborns under one year. This definition is mostly motivated by statistical convenience (a more detailed discussion can be found in Berrut et al. 2016, appendix A). From a biological perspective it is more logical to include into the infant phase the whole age interval during which the death rate decreases which is 0-10 years. Note also that the "beginning" of the death rate curve has no real signification for the following reason. The first point is for age less than one year. However, if we would take as first point the mortality rate for age less than one month, or less than one day, or less than one hour, the rate would be at least 100 times higher. More details about this hyperbolic behavior can be found in one of our earlier papers (Berrut et al. 2016).

sized in the review paper by Gavin Jarvis (2017). The main reason is the difficulty of detection of fertilization in the first two weeks (e.g. early hormonal dosage is subject to several external conditions). Similarly, early fetal deaths go unrecorded. For instance, in the United States only fetal deaths at 20 or more weeks of gestation are recorded by states in the "National Vital Statistics System". This situation suggests two comments

• Despite the statistical uncertainties of human embryogenesis, because of its sheer magnitude (multiplication by 7 with respect to the base line), it is clear that there is a major mortality surge in the early phase of embryogenesis. This is called the early embryonic mortality (EEM) effect.

• This effect can be studied in much better conditions in fish than in mammals or birds because in the former fertilization occurs outside the body of the female and can be followed from the very beginning. Zebrafish eggs have the additional advantage that their shells are transparent, a feature that is not common to all fish, e.g. the egg shells of the killifish, another model fish used in laboratories, are *not* transparent.

Death rate surges in phases 2 and 3

In the embryonic and infant phase the mortality surge is at the start whereas in the senescence phase it is at the end. Coming back to humans cases 2 and 3 are fairly easy to understand.

• In phase 2 the newborns face many challenges such as breathing, regulating body temperature, being able to absorb and digest the mother's milk. An inability to perform any one of these crucial functions means that the newborn may not live very long. Defective lungs which were acceptable during pregnancy due to the oxygen supply provided by the mother may not be able to support the life of the newborn after birth. Thus, birth appears to act as a filter resulting in the loss of newborns afflicted by life threatening defects. In spite of substantial progress in neonatal assistance, present-day data still show a marked post-natal mortality surge. In fact, mortality during the first few days after birth did not change much over past decades This is because major cardiac or neurological defects can hardly be corrected.

Note that, in contrast to Gompertz's law, the post-natal fall is not an exponential but a power law (more detailed graphs can be found in Berrut et al. 2016).

• The mortality surge seen in phase 3 is not a filtering effect. In engineering language it is known as a wear-out effect. According to Gompertz's law the death rate basically doubles every 10 years of age. With a logarithmic vertical scale the graph is a straight line (as shown in the inset of Fig.1, panel 3); with a linear vertical scale it appears in the form of a sharp increase. Gompertz's law implies that the size of a cohort decreases as an exponential whose exponent (i.e. the death rate) increases

itself exponentially with age. Although it would be interesting to investigate the reasons of such a very abrupt decline, this study would bring us too far away from our main topic and it will rather be postponed to an upcoming paper.

In the following sections the paper proceeds as follows.

• In addition to the human data already discussed in the introduction, we present embryonic mortality data for the few other cases that have been studied.

• We explain the design of our zebrafish experiment and we discuss its results.

• In the conclusion we extend the 3-phase pattern given in the introduction. We explain also how embryonic mortality can be used to probe the temperature dependence of the successive biochemical processes of embryonic development. This methodology is the parallel of the injection methodology for the exploration of the DNA structure.

• Finally, appendix A discusses some basic practical aspects of the experiment (e.g. the oxygen supply issue).

Insights into the embryonic mortality surge

The literature gives us almost no real clues for understanding the EEM effect. As our only solid starting points one can recall the following facts. (i) Far from being specific to humans, such early spikes seem to exist for all species for which data are available (see below). (ii) Most often autopsies are unable to identify the immediate causes of fetal deaths. Even in those cases where specific causes are found, they appear fairly disparate (e.g. bacterial or viral infections, lack of oxygen, accumulation of ammonia or lactic acid, chromosomal anomalies) and do not reveal the common effect which may lie behind.

Even if one accepts the fairly natural idea (borrowed from the neonatal phase) that the EEM is a filtering process (which in reliability engineering is called a wear-in process) this gives little insight. Contrary to birth defects, one cannot point to failure of specific organs for in the early embryonic phase (i.e. the cleavage phase) there are only undifferentiated cells. That is why we think it may be enlightening to describe two cases in which defects in early cell organization carry lethal effects.

Effect of spatial asymmetry in the cleavage phase

Fig.2 shows position anomalies which appear at the 8-cell level that is to say, assuming a 20mn division time, some 60 mn after fertilization. This asymmetry comes together with unequal cell sizes.

The graph shows that almost all embryos affected by such a defect will die before hatching time. It would of course be of great interest to know the average time lag between the apparition of this defect and the death of the embryo. The authors of




Fig. 2 A position asymmetry in haddock embryos carries a lethal outcome. Other kinds of cellular anomalies at the same development stage are described in the same paper. It is remarkable that cellular outcrops where one or two cells protrude from the same cluster of cells do not have any negative effect in terms of hatching. The red segment corresponds to 1mm. *Source: Adapted from Rideout et al. (2004).*

Rideout et al. did not study this aspect because the paper was written for fisheries and for obvious economic reasons it was the hatching rate which was their main concern. In a subsequent paper we intend to study this aspect.

After this example of spatial asymmetry we give an example of temporal asymmetry.

How early timing irregularities affect the viability of the embryos

If all cells replicate in a synchronous way development leads successively to 1, 2, 4, 8, 16, 32, 64, \ldots cells². However, observation shows deviations from synchronicity, a fact that is not surprising actually. As they have the same DNA the daughter cells are clones of the initial zygote. However, it is known that even for clones there is a dispersion in division time (Walden et al. 2016, Bois et al. 2019b). This means that at some moments in this replication process there may be 3, 5, 6 or 7 cells (instead of 2, 4 and 8).

Will such a dispersion affect the subsequent development of the embryo? An observation based on 834 human embryos shows that the answer is yes (Cruz et al. 2012). It appears that for the 552 embryos which had a successful development³ the time spent in a 3-cell stage was on average 0.6 hour, whereas for the 282 unsuccessful developments the time spent in a 3-cell stage was one hour. All other parameters describing the timing of the divisions were the same within a $\pm 10\%$ margin. In other words, this observation suggests that a lack of synchronicity in cell divisions had a disturbing influence on embryonic development.

²In fact the divisions of the zygote (i.e. the maternal ovum after its fertilization) are not the first divisions of the maternal cell. The oocyte has already the ability to divide. Under the influence of reproductive hormones a primary oocyte completes a meiotic cell division at the end of which it splits into two separate cells: a small one which is fairly useless and a large secondary oocyte. Needless to say, this division can also be faulty.

³In this in vitro fertilization "successful" means that the embryo reached a stage (called blastocyst stage) which is just prior to its implantation in the uterus.

Here the effect is less massive than in the observation described in the previous subsection. Clearly, it would be quite important to repeat this experiment with nonhuman embryos, for instance zebra embryos. Although early embryonic developments in humans and fish are fairly different, one would expect a similar effect for the simple reason that an embryo with 5 cells has necessarily a spatial asymmetry. Even if only temporary it should affect further development.

A testable conjecture for the embryonic mortality peak

Our idea of a parallel between embryogenesis and a manufacturing process may appear somewhat speculative. However, the nice side of it is that it can be tested. Here we propose a testable conjecture. It should be noted that it is only one of several possible conjectures that can be derived from such a parallel.

In reliability engineering it is a common belief that, assuming identical technical equipment, manufacturing defects increase with the speed of the production process. The reason is simple. Control procedures take time (even if they are automatic). In successive biochemical reactions control mechanisms may be set up after each step. Such controls improve overall reliability but at the cost of slowing down the process.

It turns out that across species there are great differences in early division times: 0.5 hour for *C. elegans* (a common laboratory model organism), 0.5 hour also for zebrafish embryos, 10 hours for mice, 20 hours for cows, 24 hours for human embryos. According to the previous argument, the hatching rates of different yet "similar" species should be in proportion of their initial division times.

Clearly, the main difficulty is how to define the notion of "similar" species. For the sake of simplicity we retain the number of cells at hatching. This may be a fairly rough criterion but at least it has the advantage of being well defined; e.g. *C. elegans* has about 1,000 cells whereas zebrafish have some 25,000 cells. Thus, this argument would lead us to predict a lower hatching rate for zebra fish than for *C. elegans* larvae. More generally, the conjecture can be stated as follows.

Mortality conjecture.

For two species having approximately the same number of cells at "birth", early embryonic mortality should decrease when the initial division time increases.

Once more data become available, it will be possible to test the conjecture. If not confirmed, it may be that our criterion based on the number of cells is too crude. For instance, cell diversity may also play a key-role.

Embryonic mortality in farm animals

For humans (and more generally for mammals) a distinction is made between em-

bryonic development and fetal development. Because we will mainly be concerned with zebrafish for which there is no real need for such a distinction, we will homogenize our terminology and use "embryonic" for the whole process from fertilization to birth.

The two main sources from which data can be expected are studies of farm animals (e.g. poultry or cattle) or of organisms used in laboratories (e.g. *C. elegans* or mice).

Embryonic death in chickens

The duration of embryonic development from fertilization to hatching is about 21 days. Fig. 3a gives the graph of embryonic mortality based on data given in Peñuela et al. (2018).

In this study the number of deaths in each age interval was obtained in two ways. (i) Candling (that is to say screening the egg with the help of a light source). (ii) After hatching time the eggs which did not hatch were opened and the day of death was established. At that point the distinction between non-fertilized and fertilized eggs could be made easily: any egg which experienced embryonic development, no matter how small, could be considered fertilized.




Fig. 3a Embryonic death rate in chickens. The definition of the death rate which is used here is the same as for infant mortality that is to say number of deaths in a given age interval divided by initial number of embryos. In this study a total of 3,240 eggs were examined and there were a total of 471 embryonic deaths which gives a hatching rate of 82%. Of these deaths, 57% occurred during the first week. Older age of the female increased death rates but did not change the shape of the curve. The reasons which explain the occurrences of a maximum just prior to hatching are explained in the text. Malposition of the embryo inside the egg was a major cause of pre-hatching death. *Source: Peñuela et al. (2018, p.6505)*

Apart from the early peak there is another maximum just before hatching. It has two explanations.

• It has been observed that one half of all chick embryos which die between day 18 and 20 were in an abnormal position which did not give them access to the air cell which is on the blunt tip of the egg (Hutt 1929).

• The lungs of chicks start to work shortly before hatching; previously the oxygen absorbed into the egg through the shell was transferred to the embryo through a network of thin blood vessels constituting a kind of rudimentary lung called the allantois. If the amount of oxygen delivered is insufficient, breaking the shell becomes impossible.

Because for chicks (and more generally for all avian species whose eggs have a fairly hard shell) breaking the shell is a very challenging task, one is not surprised to see a high pre-hatching mortality peak. One would not expect a similar peak in zebrafish embryos which have a fairly soft egg membrane and this is indeed confirmed by observation. However, the same post hatching (or rather post-yolk) filtering process is expected for fish and birds likewise.

Similar results were reported for other avian species such as: pigeons, doves, ducks, grouse, pheasants, quails (Romanoff 1949) or turkeys (Fairchild et al. 2002, p. 262, Bois et al. 2019b).

Embryonic death in cattle

The widespread use of artificial insemination allows accurate determination of the time of fertilization. Fig.3b shows that the mortality peak occurs shortly after fertilization.




Fig. 3b Embryonic death rate in cows. The graph gives death rates per week of age and 1,000 successful inseminations. The study involved 63 cows and there were 35 embryonic deaths which gives a calving rate of 55%. The maximum occurred in the third week, i.e. 15 to 21 days after fertilization. The whole pregnancy lasts about 280 days. *Sources: Sreenan et al. (1986), First et al. (1988).*

In contrast with Fig.3a, there is no late maximum. This cannot be seen on Fig.3b because the graph does not extend to the end of gestation which occurs around week 40. However, in a paper by Wathes et al. (2016) one learns that from insemination to week 8 the pregnancy losses are about 50% of initial pregnancies whereas the losses between week 8 and the end of pregnancy are only 5%.

Predominance of early embryonic mortality is also observed in other farm animals. However in this case the results are less detailed than for cows as they give only the percentages of deaths which occur in the first third of the pregnancy. A constant death rate throughout embryo development would mean a share of 33% in this first third; instead the real percentages are as follows (First et al. 1988):

mare: 87%, sow: 75%, ewe: 60%

Moreover, the percentages p of total embryonic deaths as a fraction of successful inseminations were as follows:

mare: p = 40%, sow: p = 40%, ewe: p = 25%

Model organisms used in laboratories

Regrettably, almost no mortality data as a function of embryonic age could be found in the literature. The only available case concerns the European perch (Alix 2016). It reveals a huge initial peak (see Fig. 6b).

It is this paucity of data which led us to explore the embryonic mortality of zebrafish.

Exploration of the mortality of zebrafish embryos

For a number of reasons zebrafish offer ideal conditions for the exploration of embryonic mortality.

(1) As for most fish the eggs of the zebrafish are fertilized outside of the body of the female. Therefore, in contrast to many other organisms (e.g. *C. elegans*, rotifers, birds, mammals) the embryo can be observed from the very moment of fertilization to hatching.

(2) As the eggs are transparent one can see easily what is going on inside.

(3) The embryonic phase lasts about 3-4 days (depending on temperature) which is a convenient duration. Shorter, it may not give enough time to carry out the required observations. Longer, it would slow down the whole experiment without any additional benefit.

(4) After death of the embryo the eggs become opaque and therefore can be easily identified

(5) Finally, as a model organism actively studied in genetic research, zebrafish are raised in many university laboratories.

However, we will see that despite such favorable conditions, the exploration of embryonic mortality raises several questions.

The question of the black eggs

In experiments involving zebrafish embryos there are usually two preliminary steps. Firstly, after breeding, the eggs are collected and then, usually some 24 hours later, the eggs which have become opaque⁴ are removed because they are dead (see Fig.4a and panel 4 of Fig.4c). This could seem to be a simple operation but in fact it raises some questions.

• At 24 hours post fertilization (hereafter, hpf) there will be three kinds of black eggs: (i) those which were not fertilized and became black as a result, (ii) those which were fertilized but were not able to develop, (iii) those which were fertilized, developed for a few hours but died at some point before becoming black too. Most often it is assumed that all black eggs were not fertilized. We will explain in a short moment how to separate (i)+(ii) from (iii).

• A second question is what is the time-delay between the moment when a fertilized egg dies and the moment when it turns black?

Early separation of fertilized from unfertilized eggs

In order to answer the first question, one should separate the fertilized from the unfertilized embryos and do that as early as possible. A procedure is suggested by the comparison of the pictures shown in Fig.4b (panel 3) and 4c (panel 3).



Fig. 4a Live eggs (transparent) and dead eggs (opaque) with light from above. With the light from above the opaque cells appear white. On the contrary with light from below the sample, they appear black (as in the last picture of Fig.3c). *Source: Picture taken at the "Biology Institute of Paris-Seine"*.

The fertilized eggs can be easily distinguished from the non-fertilized after they have developed a second hump⁵ which occurs about one hour after the eggs were produced by the female; see Fig.4b. However in the following hours the number of humps will

12

⁴Under a stereomicroscope with light coming from below such opaque eggs appear black, whereas with light coming from above they appear white (see Fig.3a). Here, we will call them "black" because this is the color which appears on the picture.

⁵Note that at this point it cannot be strictly excluded that some fertilized eggs are nevertheless too defective to develop a second hump.



Fig. 4b Successive steps in the development of fertilized embryos. A critical step which occurs about one hour after fertilization is the apparition of a second hump (picture 3) which results from the first division of the initial embryo cell. The last picture shows the embryo shortly before hatching. *Source: Pictures (except for panel 3) taken at the "Biology Institute of Paris-Seine"*.




Fig. 4c Successive steps in the development of unfertilized embryos. The unfertilized eggs can be obtained directly from a female by a gentle massage on its belly. After that the process starts when the eggs are released in water. In fertilized as well as unfertilized eggs one sees the apparition of a first hump. Then, whereas the fertilized eggs develop a second hump as shown in the previous figure, in unfertilized eggs the single hump continues to swell while at the same time taking fairly weird shapes as shown in the third picture taken some 5 hours after release in water. In a process which starts some 8 hours after release all non fertilized eggs will turn black as shown in the last picture. *Source: Pictures taken at the Institut de Biologie de Paris-Seine by Alex Bois and at "Beijing Normal University" by Yi Zhang.*

increase to 4, 8, 16, 32 and so on. At each division the new cells become smaller because after division they are not given enough time for further growth. When there

are 32 cells (or more) they form a large hump in which individual cells can no longer be seen and which is not very different from the single humps of unfertilized eggs (see Fig.5, panel 2). In other words, the separation of unfertilized eggs should best be done between one and two hours post fertilization. As the eggs must be examined one by one a single operator can hardly treat more than 200 eggs per hour. For larger samples⁶ the solution is to increase the number of operators or/and produce the eggs in successive batches by removing the separations in the breeding boxes with time lags of about one hour. We have been using two operators and one-hour time lags between successive batches.

One way to test whether the separation was done satisfactorily is to collect the discarded eggs and to see if all of them become black after 24 hours. In the experiment presented in the next section our policy was to accept an egg as fertilized only if 2 or 4 humps could be identified clearly. Thus, when an egg was oriented in such a way that its humps were hidden (under it when using a stereo-microscope or or above it when using an inverted microscope) it was rejected even though it may have been a fertilized egg. Such a drastic selection is necessary because it is crucial not to include any unfertilized egg as this would artificially inflate initial death rates. The fact that some 95% of our discarded eggs became black after 24 hours shows that the procedure worked fairly well.

Identification of dead embryos

In following the development of the fertilized eggs the most impressive observation is that some 6 hpf the number of new black eggs starts to increase, reaching a maximum before dwindling to zero around 24 hours post fertilization. Note that there may be a time lag between the moment when an embryo dies and the moment when the egg becomes black. Our observations suggest that this time-lag increases in the course of the eggs's development. In the appendix we mention special cases for which the time-lag was of the order of two days.

With respect to the identification of deaths there are broadly speaking 3 phases:

(1) The cleavage phase which lasts about 2 hours and during which one can count the number of cells. We did not see any stoppage in development in this phase.

(2) After that the embryo assumes a shape on which changes are not easy to identify⁷. This is illustrated in Fig. 5. At this stage the most frequent defect is the lack of a thin line around the ball of yolk. Observation shows that this defect leads to death and blackening within a few hours,

⁶Say, over 400; in our experiment the largest cohort numbered 770.

⁷Of course, if one follows a single egg in the course of time even small changes may be visible but in our experiment we follow a large number of eggs and we return to the same egg only every 5-6 hours; as in such successive observations the orientation of the embryo changes, small differences may not be clearly visible.




Fig. 5 Transition from multi-hump to thin layer. The multi-hump structure results from rapid cellular division. In the first picture there are only 4 cells whereas in the second the cells are so small and their number so large that one can no longer distinguish them and count them. In the third picture the cells form a thin layer around the yolk ball. It is in this stage (from 4 hpf to 10 hpf) that the evolution of the embryo is the most difficult to follow. *Source: Pictures taken at the "Department of Life Science" of "Peking University"*.

(3) Once the embryo assumes the shape of a fish larva the development becomes easier to follow. The heart begins to beat some 25 hours post fertilization (hpf) and the flow of blood becomes gradually more visible particularly in the yolk sac and in the tail. In this phase the dead embryos can be clearly identified and counted.

In other words, it is for the age interval between 3hpf and 20hpf that one has to find appropriate criteria defining death.

In order to identify dead embryos prior to (and independently from) their blackening one can use the following two criteria.

- Further development is stopped.
- The embryo does not move.

The "no development" criterion can only be used when the development leads to changes in shape that can be clearly identified. In this respect one can make the following remarks⁸.

• As already said, in the evolution leading to 2,4,8 cells the changes in shape and in number can be clearly identified. However, in all our observations we have never seen an egg arrested in the shape of 2, 4 or 8 cells. At first sight this may suggest that in this age group the death rate is very low. It is certainly low but one must take into account that this initial cleavage phase lasts less than two hours, thus the probability of seeing a death is naturally much lower than for a phase which lasts several hours.

• Between 2 and 8 hpf the embryo has the shape of a thin border line around the yolk; this does hardly allow a clear identification of a stop in development.

• After 10 hpf the embryo takes the shape of a larva rolled up around the ball of yolk with a head and tail clearly visible; then, in the course of time, the tail becomes more and more detached from the yolk. This feature can be used to identify an

⁸In order to identify a latency in development one may try to take successive pictures of each egg in its separate well. We tried this procedure on a small sample of 40 eggs. It turns out that unless it can be done automatically, this operation is quite time consuming. Moreover the pictures are not easy to use because when the embryo has moved and shows a different side successive pictures of the same embryo do not compare easily.

arrested development.

Table 1 summarizes the successive steps of the measurement procedure.

	Time (hpf)	Device	Operation	Purpose
1	(0, 0.2)	Several breeding boxes	Production of eggs	
			(some 100-150 per batch)	
2	(0.2, 1.5)	Stereomic. (x20)	Examination of all eggs	Separation of eggs with 1 hump from those with 2 (or more)
3	(1.5, 3)	96 well plate	Repartition: 1 egg per well	To follow the development of
		Ĩ		the eggs individually
4	(3, 12h)	Light pad +	Observation of all eggs	Identification of black eggs
		x10 magnifying glass		
5	After 12h	Stereomic. (x50)	Identification of late comers	Did development stop?
6	After 24h	Stereomic. (x50)	Examination of all eggs	Identification of the embryos
				whose heart is not beating
7	After 60h	Light pad	Identification of hatched eggs	

Table 1: Operations for measuring the number of deaths

Notes: "hpf" means "hours post fertilization. Two modes of observation were alternated. The light pad surveys were fast (for a 96-well plate, it took 5mn instead of 30mn for a microscopic observation of all the eggs), and therefore they could be repeated often. However, microscopic examination was necessary to identify the eggs which were on hold and to see whether or not their heart was still beating. The identification of laggards gives a first signal that something is wrong. It can lead to death or to recovery.

Experimental results for embryonic mortality rates

The goal of our experiment was to measure the mortality rate of fertilized embryos as a function of the post fertilization age of the embryos. The curves which summarize the results are presented in Fig.6a and 6b.

Three comments are in order.

(1) In Fig.6a despite the differences in mortality levels the shapes of the curves are similar. This stability is reassuring and gives credibility to the early embryonic mortality surge. This observation comes in confirmation of previous observations made (i) for chicken (Peñuela et al. 2018, p.6505) (ii) for turkey (Fairchild et al. 2002, p.262) (iii) for European perch (Alix 2016, p.161). Based on her research at the "Institute of Marine Research" in Bergen, Dr. Maud Alix found that for cods there is also clear evidence for an early embryonic mortality peak (personal communication, results to be published shortly).

The results published for zebrafish by Yui Uchida et al. (2018) go in the same direction but the effect is less spectacular due to a small sample of embryos (n = 72).




Fig. 6a Embryonic mortality rates for fertilized zebrafish embryos. Each curve is for a separate series of observations. The numbers of fertilized embryos were as follows. (1) curve with circles: 277, (2) squares: 345, (3) triangles: 379, (4) crosses: 672 (total number=1, 673). Due probably to the young age of the parents (meaning that they have not yet been used for reproduction, but in fact the real reason of the fragility does not matter), the 4th sample involved a high percentage of "fragile" eggs which led to a huge screening process. It is to draw attention to this difference that this curve was drawn with a dotted line. However, it is interesting to observe that despite the difference in mortality levels, the shape of the curve is the same. Note that the vertical scale is logarithmic. When necessary age intervals were merged to avoid intervals with zero deaths. All experiments were performed at 23 degree Celsius. This temperature was quite stable because it was the room temperature in a laboratory with air conditioning and deprived of windows. The graph shows death rates computed with the same definition as for infant rates, i.e. number of deaths in a given age interval divided by number of initial embryos. *Source: The experiments were performed during the summer of 2019 partly at the "Department of Life Science" of "Peking University" for the initial separation of fertilized eggs from non fertilized eggs and partly at the "Institute of Physics" of the "Chinese Academy of Science" for subsequent observations of the embryos.*

(2) One may wonder what is the main factor causing variability. The comparison of curves 1,2,3 on the one hand and curve 4 on the other hand strongly suggests that the characteristics of the parents are essential. On average in our experiments some 4 male-female pairs contributed to the production of the eggs⁹. Such a low number is clearly not sufficient to achieve homogenization by mixing. In fact, what made sample 4 special is that it had 6 young pairs and only two "normal" pairs of which only one gave eggs. Thus, there was almost no homogenization. Employing more breeding boxes will improve homogenization and will likely eliminate the most spurious cases.

⁹The number of breeding boxes was usually higher, say 6 or 8, but breeding did not occur in all boxes.




Fig. 6b Average embryonic mortality rates for zebrafish and European perch. The curve with the circles is the average of the three curves 1,2,3 of Fig.6a and it corresponds to the lower horizontal scale (the whole embryonic phase lasts 3-4 days). The error bars are $\pm \sigma$ which defines the confidence intervals with a probability level of 0.66. This means that if repeated 100 times under identical conditions the experiment would give results falling within the error bars in 66 cases. The curve with the squares is for the European perch (*Perca fluviatilis*) and corresponds to the upper horizontal scale. Note that drawing a smooth line through this sharp peak was hardly possible which is why we drew straight connecting segments. *Sources: Zebrafish: same as for Fig.4a; Perca fluviatilis: Alix (2016).*

Naturally, instead of achieving homogenization by mixing one can also try to obtain it by selection, for instance by using clones. However, like real twins, clones are affected by their life path and may be identical only to some extent (more details about differences in real twins are given in the appendix of Bois et al. 2019b).

(3) How does the order of magnitude of the overall embryonic mortality compare with the mortality of the larvae after hatching? The results presented in Bois et al. (2019, Fig.4a) for a period of 50 days post hatching show an average of about 0.4 death per hour and 1,000 hatched larva. This is approximately the order of magnitude that we see on the right-hand side of the curves in Fig. 6a,b. Moreover, the fact that the amplitude of the curves is the same for the zebrafish and for the European perch suggests that embryonic death rates are fairly robust, remember that the perch is a much bigger fish than the zebrafish.

Conclusion

The measurements presented in this study show that there is a huge mortality peak

at the beginning of the embryonic phase. There can be two sources of mortality. (i) The initial elements (i.e. oocyte and sperm cell) may be defective. (ii) Subsequent defects may arise either in the process of fertilization or in the early stages of the development (of which we have given two examples).

From assisted to autonomous embryos

It is natural to interpret the death surge as a screening process which will ensure successful subsequent development of the survivors, but why should it take place at the beginning of the embryogenesis?

Taking again inspiration from the birth process in which it is the environment change which triggers the screening process, we can observe that here too there is an environment change when the embryo becomes autonomous and does no longer rely on the support of the female organism. For instance, a partially defective function may have been acceptable as long as it could be propped up through interaction with the female organism. Once released in water the fertilized embryos must rely on their own resources. The successful completion of the second hump seems to signal that the embryo has passed this first screening. Fig.7 extends the graphs of Fig.1 to the pre-embryonic phase because a knowledge of this phase should give a better understanding of the EEM.




Fig.7 Evolution of the death rate in the four phases of development. This figure illustrates a conjecture which generalizes Fig.1 in two ways. Firstly, it introduces an initial phase corresponding to the "manufacturing" of the spermatozoa and oocytes, respectively containing the male and female germ cells. Secondly, this process in four phases is presumed to apply to a broad spectrum of species from fish to mammals and birds. However, it excludes organisms whose development involves metamorphosis through successive stages (e.g. larva, puppa, imago) which follow a more complicated process.

Embryonic mortality as a probe for exploring embryogenesis

We have already mentioned the EEM measurements done by a Japanese team of the

University of Tokyo (Uchida et al. 2018) but we did not yet explain why this is truly a pioneering investigation. It can serve as a model for future studies in (at least) two respects.

• The results for zebrafish are paralleled by similar results for chicken and African frogs. Although essential to ensure a broad range of validity of the results, such comparative studies are rare. The study by Uchida et al. may convince biologists that the benefits justify the additional work required by comparative investigations.

• Although the title of the paper refers to a specific problem in evolutionary embryology, its results go beyond that. Indeed by submitting embryos to temporary shocks¹⁰ it was found that development is more "fragile" in early than in late stages of embryogenesis, in the sense that further development is more affected by early than by late shocks.

Applying temporary or permanent temperature changes means that, instead of altering the "design" (i.e. the DNA) one changes the conditions of "production" (i.e. the output of biochemical reactions). Such an analysis can be used to explore the underlying biochemical processes which are the components of embryonic development. Note that this (shock \rightarrow response) methodology is also used extensively in genetics. By observing phenotype responses to controled alterations of the DNA one can explore the DNA and RNA sequences. Because such sequences can be altered in many possible ways, these studies give an abundant harvest of data. On the other hand, due to their wide diversity, such results may be difficult to integrate into a common framework. In contrast, via Arrhenius's law¹¹, temperature is a common determining factor of biochemical reactions, and one can therefore expect that temperature effects follow fairly *basic* laws.

Finally the following objective can be mentioned for upcoming experiments.

Mathematical characterization of the shape of the embryonic spike

Does embryonic mortality follow an exponential or a power law decay? A reliable answer would require more accurate death rate measurements. As the birth peak decay is known to be hyperbolic, a similar shape for the EEM fall would give further confidence in a parallel between the two phenomena.

¹¹It can be written:

$$k = A \exp\left(-E_a/RT\right)$$

 $k = \text{reaction constant}, T = \text{Kelvin temperature}, R = \text{constant of ideal gases}, E_a = \text{activation energy}$

The formula shows that the reaction constant (which controls the speed of the reaction) increases with the temperature. At biomolecular level, the transformations undergone by the embryo consist in biochemical reactions which is why biological processes also follow Arrhenius's law.

¹⁰Several kinds of shocks were used: temperature changes, ultra violet light, chemical inhibitors, but in the subsequent discussion we focus particularly on heat shocks.

Appendix A. Some practical issues of the experiment

The purpose of this appendix is to explain some special aspects of the experiment in the hope that it may be helpful for researchers who wish to repeat this experiment with higher accuracy i.e. less background noise.

Organization of the experiment and bioethical rules

The experiment was performed in China at the Institute of Physics of the Chinese Academy of Science in Beijing. The eggs were produced by standard breeding techniques at the Life Science Department of Peking University. Independently, a number of pictures were taken at the "Biology Institute of Paris-Seine", France.

As zebrafish are vertebrates, according to European regulation experimental protocols should get prior ethical approval. However, this rule applies only to adult fish and to larva which are more than three days old post hatching. For experiments involving only embryos there are no specific rules, hence no need for prior approval.

The oxygen supply question

As explained in Table 1, after separation from the unfertilized, the fertilized eggs were distributed in a 96-well plate. Each of these wells has a diameter of 6mm and a depth of 10mm which gives a ratio $r = (\text{surface of interface})/\text{depth}=2.8\text{mm}^{-1}$. It is this ratio which rules the oxygenation process of the water (Chen et al. 2019).

With respect to oxygen supply there are two distinct phases: (i) Before the heart is working, oxygen is supplied and distributed to the embryo by diffusion. (ii) Once the heart works, the oxygen is still supplied by diffusion¹² but it can be distributed more effectively through the hemoglobin of the blood flow.

Intuitively, it seems clear that the consumption of oxygen is correlated with the activity of the embryo which suggests that it should become higher as the embryogenesis progresses. This is indeed confirmed by observation (Green 2004). While embryogenesis progresses three effects take place.

• As already said the required volume of oxygen increases

• Tail movements which appear on day 2 produce small displacements of the eggs and therefore convection through which the higher concentration of oxygen near the water surface in each well will spread to the bottom of the well where the egg is located.

• After two days there is a proliferation of bacteria (clearly visible in the vicinity of the eggs) which consume oxygen.

As it is difficult to draw a clear conclusion from these conflicting effects we set up a test. Instead of the 96-well plate we used a plate with 384 wells. These wells

¹²The gills start to work only some 20 days after hatching.

have a 3mmx3mm square section and the same height of 10mm as the 96-well plate. This gives r = 0.9mm⁻¹, i.e. three times smaller than for the 96-well plate. Even in such narrow wells the embryos did not die but had a small hatching rate of only 30%. It can also be observed that even in the 96-plate, after hatching the larva remain mostly motionless. This may be due to insufficient oxygen but it may also be due to insufficient space. It is true that the larva would have enough space for swimming around the well but at this stage their natural movements consist in straight forward leaps for which there is not enough space.

Two precautions will improve the oxygenation of the water in the wells. (1) the plate should be left open for a cover would reduce oxygen concentration particularly in the central part of the plate. (ii) The wells should be filled (and refilled) with water only to one half (or one third) of their height so as to increase the r ratio.

The temperature question

For zebrafish a temperature of 28 degree Celsius is generally considered as optimum. As a consequence of the Law of Arrhenius, the speed of the process of embryogenesis increases with temperature (at least in the interval 20-34 degrees). However, most of our experiments were done at a constant room temperature of 23 degrees. Why?

• When kept in an incubator at 28 degrees, the eggs will experience heat shocks every time they are taken out of the incubator for observation (and similarly when put back into it). We do not know how such recurrent temperature shifts will affect them.

• Beside the velocity effect, a temperature under (or over) 28 degrees may also affect the mortality and this effect may be different across age groups. The way different age groups respond may tell us something about the processes which take place.

Special episodes

We mention here two special episodes. We think that they should be seen as overextended manifestations of effects which occur also in normal cases although in less spectacular form. This is obvious for the second case and it may also be true for the first.

Latency effect In three cases we have observed a latency time in the development process¹³ in the sense that during two days there was no development nor any movement of the embryos. Then, after this delay, in two cases the development resumed normally whereas in the third case the heart stopped and shortly after the egg became

¹³One can recall that in human embryonic development, there can be a waiting time effect in which implantation of the embryo in the uterus can be delayed by several days during which the embryo remains in a quiescent stage. In natural conditions a distinction is made between diapause which is in response to adverse environmental conditions and quiescence which is rather triggered by internal factors.

black.

Effect of egg quality One set of eggs was of bad quality¹⁴ as was apparent by the high initial number of eggs in bad shape (eggs of irregular shape, broken shells, eggs containing black particles). In such a case, if the first day acts as a filter, one expects a high early mortality. This is indeed what happened. It is true that a high mortality for mediocre eggs is nothing surprising; what is more remarkable is the fact that after the first 24h their mortality rate fell as rapidly as for normal eggs.

Acknowledgments We wish to express our sincere gratitude to the following persons for their help and interest: Maud Alix of the "Institute of Marine Research" in Bergen, Norway; Alex Bois of "Sorbonne University"; Naoki Irie and Yui Uchida from the "University of Tokyo"; Xiaolong Fan, Xueri Ma and Yi Zhang from "Beijing Normal University"; Bo Zhang and Christopher Krueger from "Peking University"; Chao Jiang of the "Institute of Physics" of the "Chinese Academy of Science".

References

- Alix (M.) 2016: Etude de la variabilité de l'embryogénèse chez la perche commune: développement d'approches alternatives. [Embryogenesis of the European perch (Perca fluviatilis): alternative approaches]. PhD Thesis presented at the University of Lorraine on 15 December 2008.
- Bell (G.), Mooers (A.O.) 1997: Size and complexity among multicellular organisms. Biological Journal of the Linnean Society 60,345-363.
- Berrut (S.), Pouillard (V.), Richmond (P.), Roehner (B.M.) 2016: Deciphering infant mortality. Physica A 463,400-426.
- Bois (A.), Garcia-Roger (E.M.), Hong (E.), Hutzler (S.), Ali Irannezhad (A.), Mannioui (A.), Richmond (P.), Roehner (B.M.), Tronche (S.) 2019a: Infant mortality across species. A global probe of congenital abnormalities. Physica A 535,122308.
- Bois (A.), Garcia-Roger (E.M.), Hong (E.), Hutzler (S.), Ali Irannezhad (A.), Mannioui (A.), Richmond (P.), Roehner (B.M.), Tronche (S.) 2019b: Congenital anomalies from a physics perspective. The key role of "manufacturing" volatility. Physica A 537,122742.
- Bois (A.), Garcia-Roger (E.M.), Hong (E.), Hutzler (S.), Ali Irannezhad (A.), Mannioui (A.), Richmond (P.), Roehner (B.M.), Tronche (S.) 2020: Physical models of mortality. Implications for defects in biological systems. Preprint March 2020, Trinity College, Dublin.

Chen (Q.), Di (Z.), Richmond (P.), Roehner (B.M.), Zhou (X.), Xu (H.) 2019: Basic

¹⁴Due probably to the young age of the males and females.

rules governing the time scales of dissolution of O_2 molecules in water, with a focus on biological applications. Preprint, Beijing Normal University (November 2019).

- Cruz (M.), Garrido (N.), Herrero (J.), Pérez-Cano (I.), Meseguer (M.) 2012: Timing of cell division in human cleavage-stage embryos is linked with blastocyst formation and quality. Reproductive BioMedicine Online 25,371-381.
- Fairchild (B.D.), Christensen (V.L.), Grimes (J.L.), Wineland (M.J.), Bagley (L.G.)2002: Hen age relationship with embryonic mortality and fertility in commercial turkeys. The Journal of Applied Poultry Research 11,3,260-265.
- First (N.L.), Eyestone (W.H.) 1988: Reproductive efficiency in domestic animals. Relative efficiency of each step in reproduction and causes of variation. Annals of the New York Academy of Science 541,1,697-705.
- French (F.E.), Bierman (J.M.) 1962: Probabilities of fetal mortality. Public Health Reports 77,10,835-847.
- Glaser (O.) 1924: Temperature and forward movement of paramecium. The Journal of General Physiology 7,2,177-188.
- Gompertz (B.) 1825: On the nature of the function expressive of the law of human mortality, and on the mode of determining the value of life contingencies. Philosophical Transactions of the Royal Society 115,513-585.
- Green (B.S.) 2004: Embryogenesis and oxygen consumption in benthic egg clutches of a tropical clownfish, *Amphiprion melanopus* (Pomacentridae). Comparative Biochemistry and Physiology Part A,138,33-38.
- Grove (R.D.), Hetzel (A.M.) 1968: Vital statistics rates in the United States, 19401960. United States Printing Office, Washington, DC.
- Hutt (F.B.) 1929: Studies in embryonic mortality in the fowl. I. The frequency of various malpositions of the chick embryo and their significance. Proceedings of the Royal Society of Edinburgh 49,II,118-130.
- Jarvis (G.E.) 2017: Early embryo mortality in natural human reproduction: what the data say. FT1000Research p.1-43.
- Kirkwood (T.B.L.), Rose (M.R.) 1991: Evolution of senescence: late survival sacrificed for reproduction. Philosophical Transactions: Biological Sciences, Vol. 332, No. 1262, "The Evolution of Reproductive Strategies", pp. 15-24
- Léridon (H.) 1973: Aspects biométriques de la fécondité humaine. Presses Universitaires de France, Paris.

[An English (somewhat expanded) translation was published in 1977 under the title given in the following reference.]

Léridon (H.) and Judith Helzner (translator) 1977: Human fertility. The basic com-

ponents. Chicago University Press, Chicago. Particularly: Ch.3: Fecundability (p.22-47) and Ch.4: Intra-uterine mortality (p.48-81).

- Peñuela (A.), Hernandez (A.) 2018: Characterization of embryonic mortality in broilers. Revista MVZ (Medecina, Vetenaria, Zootecnia) Córdoba 23,1,6500-6513.
- Post (J.), Huang (C.Y.), Hoffman (J.) 1963: The replication time and pattern of the liver cell in growing rat. The Journal of Cell Biology 18,1-12.
- Richmond (P.), Roehner (B.) 2016: Predictive implications of Gompertzs law. PhysicaA 447,446454.
- Richmond (P.), Roehner (B.) 2019a: A physicist's view of the similarities and differences between tuberculosis and cancer. Physica A 534,1-18.
- Richmond (P.), Roehner (B.M.), Wang (Q-H) 2019b: The physics of large-scale food crises. Physica A 522,274293.
- Rideout (R.M.), Trippel (E.A.), Litvak (M.K.) 2004: Predicting haddock embryo viability based on early cleavage patterns. Aquaculture 230,215-228.
- Romanoff (A.) 1949: Critical periods and causes of death in avian embryonic development. The Auk 66,3,264-270.
- Sreenan (J.M.), Diskin (M.C.) 1986: The extent and timing of embryonic mortality in the cow. p.1-11 of : "Embryonic mortality in farm animals". edited by J.M. Sreenan and M.G. Diskin. Martinus Nijhoff.
- Tarkowski (A.J.K.) 1959: Experiments on the development of isolated blastomeres of mouse eggs. Nature 184,1286-1287.

[This was the first experimental proof of the notion of totipotent cells]

- Uchida (Y.), Uesaka (M.), Yamamoto (T.), Takeda (H.), Irie (N.) 2018: Embryonic lethality is not sufficient to explain hourglass-like conservation of vertebrate embryos. EvoDevo 9,7,1-11.
- Wallden (M.), Fange (D.), Lundius (E.G.), Baltekin (Ö.), Elf (J.) 2016: The synchronization of replication and division cycles in individual *E. coli* cells. Cell 166,729-739.
- Wathes (D.C.), Diskin (M.G.) 2016: Reproduction, events and management: mating management: fertility Reference Module in Food Science, 2016.