

STRONGLY GRADED GROUPOID AND DIRECTED GRAPH ALGEBRAS

LISA ORLOFF CLARK, ELLIS DAWSON AND IAIN RAEBURN

ABSTRACT. We show the reduced C^* -algebra of a graded ample groupoid is a strongly graded C^* -algebra if and only if the corresponding Steinberg algebra is a strongly graded ring. We apply this result to get a theorem about the Leavitt path algebra and C^* -algebra of an arbitrary graph.

The prototypical example of a strongly \mathbb{Z} -graded ring is the ring $K[x, x^{-1}]$ of Laurent polynomials over a field K . The homogeneous components are the sets Kx^p , and for $m, n \in \mathbb{Z}$ such that $m+n = p$, every element $rx^p \in Kx^p$ can be realised as a product $(rx^m)(1x^n)$ of elements in the homogeneous components Kx^m and Kx^n . The C^* -algebraic analogue of $K[x, x^{-1}]$ is the C^* -algebra $C(\mathbb{T})$ of continuous functions on the unit circle $\mathbb{T} \subseteq \mathbb{C}$, which by the Stone–Weierstrass theorem contains the trigonometric polynomials as a dense subspace. This is the prototype of a strongly \mathbb{Z} -graded C^* -algebra with homogeneous subspaces $C(\mathbb{T})_n = \mathbb{C}z^n$.

To be more precise, suppose that A is a C^* -algebra and Γ is a group. Following Exel [9, Definition 16.2], we say that A is a Γ -graded C^* -algebra if there are linearly independent closed subspaces $\{A_\gamma : \gamma \in \Gamma\}$ of A such that for every $\alpha, \beta \in \Gamma$, we have

- (a) $A_\alpha \cdot A_\beta := \text{span}\{ab : a \in A_\alpha, b \in A_\beta\} \subseteq A_{\alpha\beta}$,
- (b) $A_\alpha^* := \{a^* : a \in A_\alpha\} \subseteq A_{\alpha^{-1}}$ and
- (c) $\bigoplus_{\gamma \in \Gamma} A_\gamma$ is dense in A .

We say that A is a *strongly* Γ -graded C^* -algebra if in addition $A_\alpha \cdot A_\beta$ is dense in $A_{\alpha\beta}$ for all $\alpha\beta \in \Gamma$.

Our work is inspired by a recent result of Clark, Hazrat and Rigby in [7] which identifies the groupoids for which the Steinberg algebra is a strongly graded ring. Given an ample groupoid G with Hausdorff unit space and a continuous cocycle $c : G \rightarrow \Gamma$, c induces a grading on

2010 *Mathematics Subject Classification.* 16S99, 16S10, 22A22, 46L05, 46L55.

Key words and phrases. Groupoid C^* -algebra, Steinberg algebra, Leavitt path algebra.

This research was supported by a Marsden grant from the Royal Society of New Zealand.


the Steinberg R -algebra $A_R(G)$. The groupoid G is strongly graded in the sense that $c^{-1}(\alpha)c^{-1}(\beta) = c^{-1}(\alpha\beta)$ for all $\alpha, \beta \in \Gamma$ if and only if $A_R(G)$ is a strongly Γ -graded ring [7, Theorem 3.11]. Strongly graded groupoids also appear in [4]. In Proposition 2 we show that c also induces a Γ -grading on the reduced C^* -algebra $C_r^*(G)$. We then show in Theorem 3 that $C_r^*(G)$ is a strongly Γ -graded C^* -algebra if and only if the complex Steinberg algebra $A(G)$ is a strongly Γ -graded ring. Our main tool in the proof is the injective, continuous linear map j from $C_r^*(G)$ into the space of functions from G to \mathbb{C} that are bounded with respect to the uniform norm from [14, Proposition 4.4.2].

The key examples for us are the C^* -algebras $C^*(E)$ of directed graphs E , for which the gauge action $\gamma : \mathbb{T} \rightarrow \text{Aut } C^*(E)$ gives a \mathbb{Z} -grading with

$$C^*(E)_n := \{a \in C^*(E) : \gamma_z(a) = z^n a \text{ for all } z \in \mathbb{T}\}.$$

In [7], the authors identify the directed graphs E for which the Leavitt path algebras $L_R(E)$ are strongly \mathbb{Z} -graded in [7, Theorem 4.2]. (See also [10, Theorem 1.3].) We use our Theorem 3 to show $L_{\mathbb{C}}(E)$ is strongly \mathbb{Z} -graded if and only if $C^*(E)$ is strongly \mathbb{Z} -graded. Thus the graph condition of [7, Theorem 2.4] (which we will describe soon) also characterises when $C^*(E)$ is strongly \mathbb{Z} -graded.

Since we are proving a C^* -algebraic result, we use the convention that the spanning elements $\{\alpha\beta^*\}$ of $L_{\mathbb{C}}(E)$ are determined by pairs $\alpha, \beta \in E^*$ satisfying $s(\alpha) = s(\beta)$. In other words, we use the conventions of [13] rather than those of [1]. With these conventions, a directed graph E has *property (Y)* if, for every infinite path $x \in E^\infty$ and every $k \in \mathbb{N}$, there are an initial segment α of x and a finite path $\beta \in E^*$ such that $s(\alpha) = s(\beta)$ and $|\beta| - |\alpha| = k$, that is, β has k more edges than α . The reason for the name is the picture:



We can state our main graph algebra result as follows.

Theorem 1. *Suppose that E is a directed graph. Then the graph algebra $C^*(E)$ is strongly \mathbb{Z} -graded if and only if E is row-finite, has no sources, and has property (Y).*

As a corollary, we get that $C^*(E)$ is a strongly \mathbb{Z} -graded C^* -algebra if and only if $L_{\mathbb{C}}(E)$ is strongly \mathbb{Z} -graded ring. We conclude with a brief analysis of $(L(E))_0$, the core of a Leavitt path algebra which is dense inside of the core of $C^*(E)$. We show that every element of the core of a Leavitt path algebra is contained in a finite-dimensional subalgebra. Previously this was known for row-finite graphs, see for example the last line of the proof of [2, Theorem 5.3].

1. STRONGLY GRADED STEINBERG ALGEBRAS AND GROUPOID C^* -ALGEBRAS

We begin this section with some definitions. Suppose R is a ring. If A and B are subsets of R , then we write AB for the set of all finite sums of elements of the form ab where $a \in A$ and $b \in B$. Let Γ be a group. We say that R is a Γ -graded ring if there are subgroups $\{R_\gamma : \gamma \in \Gamma\}$ in R such that for every $\alpha, \beta \in \Gamma$

- (a) $R_\alpha R_\beta \subseteq R_{\alpha\beta}$ and
- (b) $\bigoplus_{\gamma \in \Gamma} R_\gamma = R$.

Further, we say R is a *strongly Γ -graded ring* if the containment in (a) above is in fact an equality.

We say a groupoid G is ample if it is a topological groupoid and has a basis of compact open bisections. When G is an ample groupoid such that $G^{(0)}$ is Hausdorff, the complex Steinberg algebra of G is the space

$$A(G) := \text{span}\{1_B : B \text{ is a compact open bisection}\}$$

where 1_X denotes the characteristic function from G to \mathbb{C} of X . Addition and scalar multiplication are defined pointwise, and the convolution product is such that

$$1_B 1_D = 1_{BD}$$

for compact open bisections B and D . If $c : G \rightarrow \Gamma$ is a continuous cocycle into a discrete group Γ , then the Steinberg algebra $A(G)$ is a Γ -graded ring such that for $\gamma \in \Gamma$

$$A(G)_\gamma := \{f \in A(G) : \text{supp } f \subseteq G_\gamma\}$$

where $G_\gamma := c^{-1}(\gamma)$. Thus

$$A(G) = \bigoplus_{\gamma \in \Gamma} A(G)_\gamma.$$

(See [8, Lemma 3.11].)

The next proposition shows that this structure also gives a grading on the groupoid C^* -algebra. For this proposition (and also the following theorem) we use the injective continuous linear map

$$j : C_r^*(G) \rightarrow \mathcal{B}(G),$$

where $\mathcal{B}(G)$ denotes the normed vector space of bounded functions from G to \mathbb{C} with respect to $\|\cdot\|_\infty$. This map restricts to the identity on $A(G)$. (See, for example, [5, page 3680].)

Proposition 2. *Let G be an ample groupoid such that $G^{(0)}$ is Hausdorff and let $c : G \rightarrow \Gamma$ be a continuous cocycle into a discrete group Γ . Then $C_r^*(G)$ is a Γ -graded C^* -algebra such that for each $\gamma \in \Gamma$ we have*

$$C_r^*(G)_\gamma := \overline{A(G)_\gamma}.$$

Proof. Fix $\alpha, \beta \in \Gamma$. Because $A(G)_\alpha \cdot A(G)_\beta$ is contained in the closed set $C_r^*(G)_{\alpha\beta}$,

$$C_r^*(G)_\alpha \cdot C_r^*(G)_\beta \subseteq C_r^*(G)_{\alpha\beta}$$

by the continuity of multiplication. Similarly

$$C_r^*(G)_\gamma^* \subseteq C_r^*(G)_{\gamma^{-1}}$$

by the continuity of the involution $*$. Next we verify that each $C_r^*(G)_\gamma$ is a linearly independent subspace. To do this, we claim that

$$j(C_r^*(G)_\gamma) \subseteq \mathcal{B}(G_\gamma).$$

We have that j is the identity map on $A(G)_\gamma \subseteq \mathcal{B}(G_\gamma)$. It is straightforward to check that $\mathcal{B}(G_\gamma)$ is closed in $\mathcal{B}(G)$ with respect to $\|\cdot\|_\infty$ and hence the claim follows from the continuity j .

Now consider a finite linear combination $\sum a_i = 0$ where each $a_i \in C_r^*(G)_{\gamma_i}$ and the γ_i are distinct elements of Γ . By linearity

$$0 = j(\sum a_i) = \sum j(a_i).$$

Since the supports of $j(a_i)$ are all disjoint, $j(a_i) = 0$ for every i . Thus $a_i = 0$ because j is an injective linear map.

Finally notice

$$A(G) = \bigoplus_{\gamma \in \Gamma} A(G)_\gamma \subseteq \bigoplus_{\gamma \in \Gamma} C_r^*(G)_\gamma \subseteq C_r^*(G).$$

Thus since $A(G)$ is dense in $C_r^*(G)$ by [17, Proposition 6.7], we have that

$$\bigoplus_{\gamma \in \Gamma} C_r^*(G)_\gamma \text{ is dense in } C_r^*(G).$$

□

Theorem 3. *Let G be an ample groupoid such that $G^{(0)}$ is Hausdorff and let $c : G \rightarrow \Gamma$ be a continuous cocycle into a discrete group Γ . The following are equivalent:*

- (a) G is strongly Γ -graded groupoid in the sense that $G_\alpha G_\beta = G_{\alpha\beta}$ for all $\alpha, \beta \in \Gamma$;
- (b) $A(G)$ is a strongly Γ -graded ring;
- (c) $C_r^*(G)$ is a strongly Γ -graded C^* -algebra.

Remark 4. Using [7, Theorem 3.11], we could replace item (b) in Theorem 3 with the equivalent statement: $A_R(G)$ is a strongly Γ -graded ring for any commutative ring R with identity.

The equivalence of (a) and (b) is established in [7, Theorem 3.11]. We establish the equivalence of (b) and (c). That (c) implies (b) is straightforward: Take $\alpha, \beta \in \Gamma$. Since $C_r^*(G)_{\alpha\beta}$ is closed and $C_r^*(G)$ is Γ -graded,

$$C_r^*(G)_\alpha \cdot C_r^*(G)_\beta$$

trivially has closure in $C_r^*(G)_{\alpha\beta}$. To see that this closure is all of $C_r^*(G)_{\alpha\beta}$, take $a \in C_r^*(G)_{\alpha\beta}$. Since $C_r^*(G)_{\alpha\beta} = \overline{A(G)_{\alpha\beta}}$, there exist $f_n \in A(G)_{\alpha\beta}$ such that $f_n \rightarrow a$ in $C_r^*(G)$. Since $A(G)$ is strongly graded, we have

$$f_n \in A(G)_\alpha \cdot A(G)_\beta \subseteq C_r^*(G)_\alpha \cdot C_r^*(G)_\beta$$

for each n and hence the limit a belongs to

$$\overline{C_r^*(G)_\alpha \cdot C_r^*(G)_\beta}.$$

For the reverse implication, we use the following lemma.

Lemma 5. *Let G be an ample groupoid such that $G^{(0)}$ is Hausdorff and let $c : G \rightarrow \Gamma$ be a continuous cocycle into a discrete group Γ . Then for $\alpha, \beta \in \Gamma$ we have*

$$A(G)_\alpha \cdot A(G)_\beta \subseteq \{f \in A(G) : f(x) \neq 0 \implies x \in G_\alpha G_\beta\}.$$

Proof. Observe that $A(G)_\alpha \cdot A(G)_\beta$ is equal to

$$\text{span}\{1_{BD} : B \subseteq G_\alpha \text{ and } D \subseteq G_\beta \text{ are compact open bisections}\}$$

(see [7, bottom of page 53]). Since the support of a sum of functions is contained in the union of the supports of the summands, the containment follows. \square

Proof of Theorem 3. To complete the proof, suppose $A(G)$ is not strongly graded. Then G is not strongly graded by [7, Theorem 3.11], that is, by (a) \implies (b) of Theorem 3. Then there exists $\gamma \in \Gamma$ and $x \in G_e \setminus G_\gamma G_{\gamma^{-1}}$

by [7, Lemma 3.1]. Since c is continuous, G_e is open so we can find $B \subseteq G_e$ a compact open bisection containing x . Since $1_B(x) = 1$,

$$1_B \in A(G)_e \setminus A(G)_\gamma A(G)_{\gamma^{-1}}$$

by Lemma 5. We show

$$1_B \notin \overline{A(G)_\gamma A(G)_{\gamma^{-1}}}$$

using an adaptation of the argument in [6, Proposition 3.3(i)]. By way of contradiction, suppose there exists a sequence

$$(f_n) \subseteq A(G)_\gamma \cdot A(G)_{\gamma^{-1}}$$

that converges to 1_B in $C_r^*(G)$. Again we apply the injective continuous linear map $j : C_r^*(G) \rightarrow \mathcal{B}(G)$. Since j restricts to the identity map on $A(G)$ we have $j(f_n) = f_n \rightarrow j(1_B) = 1_B$ in $\|\cdot\|_\infty$. Recall that $1_B(x) \neq 0$. However, $x \notin G_\gamma G_{\gamma^{-1}}$ so $f_n(x) = 0$ for all $n \in \mathbb{N}$ by Lemma 5, which is a contradiction. Therefore

$$f \notin \overline{A(G)_\gamma A(G)_{\gamma^{-1}}}$$

and $C_r^*(G)$ is not a strongly Γ -graded C^* -algebra. \square

2. LEAVITT PATH ALGEBRAS

We want to connect properties of $C^*(E)$ and properties of $L_{\mathbb{C}}(E)$, and hence we want to be able to view $L_{\mathbb{C}}(E)$ as a subalgebra of $C^*(E)$ using the homomorphism ι which takes a spanning element $\alpha\beta^*$ to the element $s_\alpha s_\beta^*$ of $C^*(E)$. That ι is an injection was established (for example) in [3, §1.3], but with an annoying hypothesis of “no sources”. So we pause to prove the following more general result.

Proposition 6. *Suppose that E is a directed graph. Then the homomorphism $\iota : L_{\mathbb{C}}(E) \rightarrow C^*(E)$ is an isomorphism of $L_{\mathbb{C}}(E)$ onto the $*$ -subalgebra*

$$A := \text{span} \{s_\alpha s_\beta^* : \alpha, \beta \in E^* \text{ and } s(\alpha) = s(\beta)\} \text{ of } C^*(E).$$

Proof. We want to apply the Graded Uniqueness Theorem for Leavitt path algebras [1, Theorem 2.2.15], but $C^*(E)$ is not \mathbb{Z} -graded in the algebraic sense, and hence that result does not apply to $\iota : L_{\mathbb{C}}(E) \rightarrow C^*(E)$. However, the $*$ -subalgebra A is \mathbb{Z} -graded, with

$$A_n = \text{span} \{s_\alpha s_\beta^* : s(\alpha) = s(\beta) \text{ and } |\alpha| - |\beta| = n\}.$$

Since we can always find Cuntz-Krieger E -families $\{S, P\}$ with $P_v \neq 0$ for every $v \in E^0$ (see the top of [13, page 8]), every p_v is non-zero. Thus [1, Theorem 2.2.15] implies that ι is an injection of $L_{\mathbb{C}}(E)$ into

A ; since every $s_\alpha s_\beta^*$ belongs to the range of ι , it is an isomorphism, as claimed. \square

Next we relate the grading spaces $C^*(E)_n$ for the C^* -algebra to those of the Leavitt path algebra. We define $\Phi_n : C^*(E) \rightarrow C^*(E)$ in terms of the gauge action $\gamma : \mathbb{T} \rightarrow \text{Aut } C^*(E)$ by

$$\Phi_n(a) = \int_{\mathbb{T}} w^{-n} \gamma_w(a) dw.$$

Left invariance of the Haar integral on \mathbb{T} implies that Φ_n is a norm-decreasing $C^*(E)^\gamma$ -bilinear map of $C^*(E)$ onto $C^*(E)_n$ such that

$$\Phi_n(s_\mu s_\nu^*) = \begin{cases} s_\mu s_\nu^* & \text{if } |\mu| - |\nu| = n, \text{ and} \\ 0 & \text{otherwise.} \end{cases}$$

The map $\Phi_n \circ \iota$ is a bounded linear map of $L_{\mathbb{C}}(E)$ onto a dense subspace of $C^*(E)_n$, and its restriction to $L_{\mathbb{C}}(E)_n$ is an injection of $L_{\mathbb{C}}(E)_n$ onto this dense subspace of $C^*(E)_n$. Hence, modulo the canonical injection ι of $L_{\mathbb{C}}(E)$ in $C^*(E)$, $\overline{C^*(E)_n}$ is the closure of $L_{\mathbb{C}}(E)_n$. We suppress ι , and write $C^*(E)_n = \overline{L_{\mathbb{C}}(E)_n}$.

Proposition 7. *Suppose that E is a directed graph. Then $C^*(E)$ is a strongly \mathbb{Z} -graded C^* -algebra if and only if $L_{\mathbb{C}}(E)$ is a strongly \mathbb{Z} -graded ring.*

With the following lemma, the proof of Proposition 7 is an immediate corollary to Theorem 3.

Lemma 8. *For any directed graph E , there is an ample Hausdorff amenable groupoid G_E and an isomorphism $\pi : C^*(E) \rightarrow C^*(G_E)$ such that $\pi(C^*(E)_n) = C^*(G_E)_n$, and such that the restriction of π to $L_{\mathbb{C}}(E)$ is a \mathbb{Z} -graded (ring) isomorphism of $L_{\mathbb{C}}(E)$ onto $A(G_E)$.*

Remark 9. Because G_E is amenable, the full and reduced C^* -algebras are equal so we drop the r subscript from the $C_r^*(G_E)$ from now on.

Proof. Let G_E be the boundary path groupoid of E as defined by Paterson on page 653 of [12]¹. Then G_E is ample and Hausdorff; see [15, Theorem 2.4] for a nice proof of this. The groupoid G_E is amenable by [12, Theorem 4.2]. We view $C^*(G_E)$ as the universal groupoid C^* -algebra. In this sense, there is a representation $\pi_{max} : C_c(G_E) \rightarrow C^*(G_E)$ that is universal for representations of $C_c(G_E)$ and $\pi_{max}(C_c(G_E))$ is dense in $C^*(G_E)$. See [16, Theorem 3.2.2] for more details. Since we view $A(G_E) \subseteq C_c(G_E)$ as subsets of $C^*(G_E)$, we identify $C_c(G_E)$ with

¹Paterson calls G_E the path groupoid

$\pi_{max}(C_c(G_E))$ inside $C^*(G_E)$, which we can do because π_{max} is injective by [16, Corollary 3.3.4].

One can show that the collection

$$P_v := 1_{Z(v)}, \quad S_e := 1_{Z(e,s(e))}$$

for $v \in E^0$ and $e \in E^1$ is a Cuntz-Krieger E -family in $C^*(G_E)$. The universal property of $C^*(E)$ gives a homomorphism $\pi : C^*(E) \rightarrow C^*(G_E)$ such that

$$\pi(p_v) = P_v \quad \text{for } v \in E^0 \quad \text{and} \quad \pi(s_e) = S_e \quad \text{for } e \in E^1.$$

(See [13, page 42].) Notice that the restriction of π to $L_{\mathbb{C}}(E)$ is precisely the \mathbb{Z} -graded ring isomorphism from $L_{\mathbb{C}}(E)$ onto $A(G_E)$ in [8, Example 3.2], and hence $\pi(L_{\mathbb{C}}(E)_n) = A(G_E)_n$.

Paterson shows in [12, Theorem 3.8] and [11, Theorem 3.1] that representations of $C^*(E)$ are in bijective correspondence with representations of $C_c(G_E)$. So the universal representation p_v, s_e of $C^*(E)$ corresponds to a representation $\phi : C_c(G_E) \rightarrow C^*(E)$: indeed, by looking at the details of the proof of [12, Theorem 3.8], we see that $\phi(P_v) = p_v$ and $\phi(S_e) = s_e$ for $v \in E^0$ and $e \in E^1$. The universal property of $C^*(G_E)$ [16, Theorem 3.2.2] gives a homomorphism $\psi : C^*(G_E) \rightarrow C^*(E)$ such that $\psi = \phi$ on $C_c(G_E)$.

Notice that $\pi \circ \phi$ and $\phi \circ \pi$ restrict to identity maps on $A(G_E)$ and $L_{\mathbb{C}}(E)$ respectively. We show that π is a bijection by showing that $\pi \circ \psi$ and $\psi \circ \pi$ are the identity maps on $C^*(G_E)$ and $C^*(E)$ respectively. Fix $a \in C^*(G_E)$. Since $A(G_E)$ is dense in $C^*(G_E)$ by [17, Proposition 6.7], $a = \lim f_n$ where $f_n \in A(G_E)$. Then using that homomorphisms between C^* -algebras are automatically continuous, we have

$$\pi(\psi(a)) = \pi(\psi(\lim f_n)) = \lim \pi(\psi(f_n)) = \lim \pi(\phi(f_n)) = \lim f_n = a.$$

For the other direction, fix $b \in C^*(E)$. Then $b = \lim b_n$ where $b_n \in L_{\mathbb{C}}(E)$, and recalling that π takes elements of $L_{\mathbb{C}}(E)$ to elements of $A(G_E)$, we deduce that

$$\psi(\pi(b)) = \psi(\pi(\lim(b_n))) = \lim(\psi(\pi(b_n))) = \lim(\phi(\pi(b_n))) = \lim b_n = b.$$

Thus π is an isomorphism with inverse ψ . Continuity of π and ψ imply $\pi(C^*(E)_n) = C_r^*(G_E)_n$, and this completes the proof. \square

Proof of Proposition 7. Because the restricted map π from Lemma 8 is a graded ring isomorphism, $L_{\mathbb{C}}(E)$ is strongly \mathbb{Z} -graded if and only if $A(G_E)$ is strongly \mathbb{Z} -graded. Now Theorem 3 says this is equivalent to $C^*(G_E)$ being strongly \mathbb{Z} -graded. Since π preserves the C^* -grading, the result follows. \square

Now we have the peices needed to prove Theorem 1. Recall that [7, Theorem 4.2] says $L_{\mathbb{C}}(E)$ is strongly \mathbb{Z} -graded if and only if E is row-finite, has no sources and has property Y. Thus Theorem 1 follows from Proposition 7 above and [7, Theorem 4.2]

In strongly graded algebras, a lot of the structure of the algebra can be seen in the core, that is, in the homogeneous component of the identity e . Thus we conclude by establishing that the core of a Leavitt path algebra is build from finite dimensional subalgebras in the following way.

Proposition 10. *Suppose that E is a directed graph and K is a field. Then every $a \in L_K(E)_0$ belongs to a finite-dimensional $*$ -subalgebra of $L_K(E)_0$.*

To prove this result, we need to understand the finite-dimensional subalgebras of $L_K(E)_0$. We start by choosing an enumeration $\{e_j : j \in \mathbb{N}\}$ of the countable set E^1 . For $J \in \mathbb{N}$, we write

$$E_J^k := \{\alpha \in E^k : \alpha_i \in \{e_j : 1 \leq j \leq J\} \text{ for all } i\},$$

and $E_J^0 := \{s(e_j) : 1 \leq j \leq J\}$. Notice that the sets E_J^k are all finite, and $E_J^k \subseteq E_{J+1}^k$. We define

$$G_{k,J} := \text{span}\{\alpha\beta^* : \alpha, \beta \in E_J^k\}.$$

For $\beta, \gamma \in E_J^k$ we have $|\beta| = |\gamma| = k$, and hence we have

$$\beta^*\gamma = \begin{cases} s(\beta) = s(\gamma) & \text{if } \beta = \gamma \\ 0 & \text{otherwise.} \end{cases}$$

It follows that

$$G_{k,J}(v) := \text{span}\{\alpha\beta^* := \alpha, \beta \in E_J^k v\}$$

is a (finite-dimensional) matrix algebra, and

$$v \neq w \implies G_{k,J}(v)G_{k,J}(w) = \{0\}.$$

Thus

$$G_{k,J} = \bigoplus_{v \in E_J^0} G_{k,J}(v)$$

is a finite direct sum of matrix algebras.

Lemma 11. *Suppose that $k, l \in \mathbb{N}$ and $k > l$. Then for every $J \in \mathbb{N}$ and $v, w \in E_J^0$, we have*

$$G_{k,J}(v)G_{l,J}(w) \subseteq G_{k,J}(v).$$

Proof. For $\alpha\beta^* \in G_{k,J}$ and $\gamma\delta^* \in G_{l,J}(w)$, we have $|\beta| = k > l = |\gamma|$, and hence

$$(\alpha\beta^*)(\gamma\delta^*) = \begin{cases} \alpha(\delta\beta')^* & \text{if } \beta = \gamma\beta' \\ 0 & \text{otherwise.} \end{cases}$$

Since $s(\delta\beta') = s(\beta) = s(\alpha) = v$, we deduce that $(\alpha\beta^*)(\gamma\delta^*) \in G_{k,J}(v)$ (it could be 0, but that is in $G_{k,J}(v)$ too). A similar argument gives $(\gamma\delta^*)(\alpha\beta^*) \in G_{k,J}(v)$. \square

Corollary 12. *For each $J \in \mathbb{N}$, the set*

$$F_{k,J} := \text{span} \left(\bigcup \{G_{l,J} : l \leq k\} \right)$$

is a finite-dimensional $$ -subalgebra of $L_K(E)_0$.*

Proof. The set $F_{k,J}$ is a vector subspace of $L_K(E)_0$ and is closed under conjugation because each $G_{l,J}$ is. The lemma implies that it is closed under multiplication. It is finite-dimensional because there are only finitely many $G_{l,J}$ in play, and their union spans $F_{k,J}$. \square

Proof of Proposition 10. We suppose that $a \in L_K(E)_0$. Then there are a finite set S of pairs $(\alpha, \beta) \in E^* \times E^*$ such that $s(\alpha) = s(\beta)$ and $|\alpha| = |\beta|$, and scalars $c_{\alpha,\beta}$ such that

$$a = \sum_{(\alpha,\beta) \in S} c_{\alpha,\beta} \alpha\beta^*.$$

Since the set S is finite, the set of vertices

$$V := \{v \in E^0 : v = s(\alpha_i) \text{ or } s(\beta_i) \text{ for some } (\alpha, \beta) \in S \text{ and } i \leq |\alpha|\}$$

is finite. Thus there exists J such that $V \subseteq E_J^0$. Then with $k := \max\{|\alpha| : (\alpha, \beta) \in S\}$, we have

$$\alpha\beta^* \in G_{|\alpha|,J} \subseteq F_{k,J} \quad \text{for all } (\alpha, \beta) \in S.$$

Thus $a = \sum c_{\alpha,\beta} \alpha\beta^*$ belongs to the subspace $F_{k,J}$, which by Corollary 12 is a finite-dimensional $*$ -subalgebra of $L_K(E)_0$. \square

REFERENCES

- [1] G. Abrams, P. Ara and M. Siles Molina, *Leavitt Path Algebras*, Lecture Notes in Math., vol. 2191, Springer-Verlag, London, 2017.
- [2] P. Ara, M.A. Moreno, E. Pardo, Nonstable K-theory for graph algebras, *Algebr. Represent. Theory* **10** (2007), 157–178.
- [3] G. Aranda Pino, F. Perera Domènech and M. Siles Molina (Editors), *Graph Algebras. Bridging the Gap between Analysis and Algebra*, Univ. of Málaga, 2007.
- [4] A. Buss and R. Meyer, Inverse semigroup actions on groupoids, *Rocky Mountain J. Math* **47**(2017), 53–159.

- [5] L.O.Clark, R. Exel, E. Pardo, A. Sims and C. Starling, Simplicity of algebras associated to non-Hausdorff groupoids, *Trans. Amer. Math. Soc.***372** (2019), 3669–3712.
- [6] L.O. Clark and J. Fletcher, Groupoid algebras as covariance algebras, <https://arxiv.org/1906.02855>
- [7] L.O. Clark, R. Hazrat and S.W. Rigby, Strongly graded groupoids and strongly graded Steinberg algebras, *J. Algebra* **530** (2019), 34–68.
- [8] L.O. Clark and A. Sims, *Equivalent groupoids have Morita equivalent Steinberg algebras*, *J. Pure Appl. Algebra* **219** (2015), 2062–2075.
- [9] R. Exel, *Partial Dynamical Systems, Fell Bundles and Applications*, Math. Surveys and Monographs, vol. 224, Amer. Math. Soc., Providence, 2017.
- [10] P. Nystedt and J. Öinert, Strongly graded Leavitt Path Algebras, <https://arxiv.org/2002.06965>.
- [11] A.L.T. Paterson, Groupoids, inverse semigroups, and their operator algebras, Birkhäuser Boston Inc., Boston, MA, 1999, xvi+274.
- [12] A.L.T. Paterson, *Graph inverse semigroups, groupoids and their C^* -algebras*, *J. Operator Theory* **48** (2002), 645–662.
- [13] I. Raeburn, *Graph Algebras*, CBMS Regional Conference Series in Math., vol. 103, Amer. Math. Soc., Providence, 2005.
- [14] J. Renault, *A groupoid approach to C^* -algebras*, volume 793 of *Lecture Notes in Mathematics*, Springer, Berlin, 1980.
- [15] S.W. Rigby, The groupoid approach to Leavitt path algebras, A. A. Ambily, R. Hazrat, B. Sury eds, *Leavitt Path Algebras and Classical K -Theory*, Indian Statistical Institute Series, Springer 2020.
- [16] A. Sims, Hausdorff étale groupoids and their C^* -algebras, *Operator algebras and dynamics: groupoids, crossed products and Rokhlin dimension*, Advanced Courses in Mathematics, CRM Barcelona, Birkhauser, to appear.
- [17] B. Steinberg, A groupoid approach to discrete inverse semigroup algebras, *Adv. Math.* **223** (2010), 689–727.

DEPARTMENT OF MATHEMATICS AND STATISTICS, VICTORIA UNIVERSITY OF WELLINGTON, PO BOX 600, WELLINGTON 6140, NEW ZEALAND