STATISTICAL STABILITY FOR DIFFEOMORPHISMS WITH MOSTLY EXPANDING AND MOSTLY CONTRACTING CENTERS

ZEYA MI AND YONGLUO CAO

Abstract. For partially hyperbolic diffeomorphisms with mostly expanding and mostly contracting centers, we establish a topological structure, called skeleton–a set consisting of finitely many hyperbolic periodic points with maximal cardinality for which there exist no heteroclinic intersections. We build the one-to-one corresponding between periodic points in any skeleton and physical measures. By making perturbations on skeletons, we study the continuity of physical measures with respect to dynamics under C^1 -topology.

1. INTRODUCTION

Given a C^1 diffeomorphism f on a smooth compact Riemannian manifold M, an invariant measure μ is called a *physical measure* if it is physical observable [\[29\]](#page-42-0), in the sense that the set of its basin

$$
B(\mu, f) = \left\{ x \in M : \frac{1}{n} \sum_{i=0}^{n-1} \delta_{f^i x} \xrightarrow{weak^*} \mu \text{ as } n \to +\infty \right\}
$$

has positive Lebesgue measure. Physical measures were discovered by Sinai, Ruelle and Bowen [\[50,](#page-43-0) [48,](#page-43-1) [11,](#page-41-0) [12\]](#page-41-1) in 1970's for uniformly hyperbolic systems, which can reflect the chaotic properties of dynamical systems from different viewpoints, such as entropies, Lyapunov exponents and equilibrium states (see [\[44,](#page-43-2) [54\]](#page-43-3)).

When the existence of physical measures is established, one expects to describe how the physical measures bifurcate as dynamics changes, which known as the subject of statistical stability. Uniformly hyperbolic systems are well-known to be statistically stable [\[49\]](#page-43-4). In recent decades, much progress has been made on the statistical stability for systems with certain weak forms of hyperbolicity (see e.g. $[1, 5, 3, 52, 8]$ $[1, 5, 3, 52, 8]$ $[1, 5, 3, 52, 8]$ $[1, 5, 3, 52, 8]$ $[1, 5, 3, 52, 8]$). In this work, we mainly study the statistical stability for a rich family of diffeomorphisms

Date: March 11, 2020.

²⁰⁰⁰ Mathematics Subject Classification. 37D30, 37D25, 37C40.

Key words and phrases. Partial hyperbolicity, Physical measures, SRB measures, Lyapunov exponent, Statistical stability.

Z.Mi was supported by NSFC 11801278 and The Startup Foundation for Introducing Talent of NUIST(Grant No. 2017r070).

Y.Cao was supported by NSFC (11771317,11790274).

beyond uniform hyperbolicity. We complete this picture by finding a geometrical combinatorial structure, which is called skeleton. By establishing the one-to-one corresponding between physical measures and elements of the skeleton, we explore the statistical properties and the stability for physical measures via skeletons in geometrical viewpoint.

A diffeomorphism f is partially hyperbolic if there exists a Df -invariant splitting $TM = E \oplus F$ of the tangent bundle such that

- $TM = E \oplus F$ is a dominated splitting: $||Df||_{E(x)}|| \cdot ||Df^{-1}|_{F(f(x))}|| <$ 1/2 for every $x \in M$, rewrite $TM = E \oplus_{\succ} F$ to indicate this domination;
- either E is uniformly expanding or F is uniformly contracting.

Throughout, we write E^u and E^s to indicate that they are uniformly expanding and uniformly contracting invariant sub-bundles, respectively. For C^{1+} C^{1+} C^{1+} partially hyperbolic diffeomorphisms¹ exhibiting E^u , Pesin and Sinai [\[46\]](#page-43-6) introduced the notion of Gibbs u-states, an invariant measure μ is a Gibbs *u*-state when its conditional measures along *strong* unstable manifolds are absolutely continuous with respect to Lebesgue measures along these manifolds. From [\[37\]](#page-42-1), we know that μ is a Gibbs u-state if and only if it satisfies the following entropy formula

(1.1)
$$
h_{\mu}(f, \mathcal{F}^{u}) = \int \log |\det Df|_{E^{u}}| d\mu,
$$

where $h_{\mu}(f, \mathcal{F}^u)$ denotes the partial entropy along strong unstable foliation \mathcal{F}^u tangent to E^u (see Definition [2.2\)](#page-8-0). It is worth noting that the above entropy formula can be defined for $C¹$ partially hyperbolic diffeomorphisms. Thus, as proposed by Croviser-Yang-Zhang [\[26\]](#page-42-2), in this paper we call μ a Gibbs u-state for f if (1.1) holds for μ . The set of Gibbs u-states plays a crucial role in the study of physical measures for partially hyperbolic diffeomorphisms (see e.g. [\[13,](#page-41-6) [15,](#page-42-3) [25,](#page-42-4) [53,](#page-43-7) [7,](#page-41-7) [8,](#page-41-5) [18\]](#page-42-5)).

In this paper, we shall mainly investigate the topological and measuretheoretic behaviors for an open class of partially hyperbolic diffeomorphisms introduced in [\[40\]](#page-43-8). Let $PH_{EC}^1(M)$ be the set of C^1 partially hyperbolic diffeomorphisms on M satisfying that for every f in it, there exists a partially hyperbolic splitting $TM = E^u \oplus_{\succ} E^{cu} \oplus_{\succ} E^{cs}$ for which

- E^{cu} is mostly expanding: μ has only positive Lyapunov exponents along E^{cu} for every Gibbs u-state μ ;
- E^{cs} is mostly contracting: μ has only negative Lyapunov exponents along E^{cs} for every Gibbs u-state μ .

Denote by $PH_{EC}^{1+}(M)$ the set of diffeomorphisms in $PH_{EC}^{1}(M)$ of class C^{1+} . Let us list some properties for this kind of partially hyperbolic diffeomorphisms:

¹We say that f is a C^{1+} diffeomorphism if it is C^{1} and Df is Hölder continuous.

- Following the result of [\[55,](#page-43-9) Theorem B], we know that $PH_{EC}^{1}(M)$ is an open subset of $C¹$ diffeomorphisms.
- $PH_{EC}^{1+}(M)$ contains both the diffeomorphisms with mostly expanding center and diffeomorphisms with mostly contracting center. Both of them have been studied from different aspects of interests recently (see e.g. [\[13,](#page-41-6) [27,](#page-42-6) [20,](#page-42-7) [21,](#page-42-8) [6,](#page-41-8) [53,](#page-43-7) [7,](#page-41-7) [55,](#page-43-9) [28,](#page-42-9) [56\]](#page-43-10)).
- By [\[40,](#page-43-8) Theorem A], any diffeomorphism of $PH_{EC}^{1+}(M)$ has finitely many ergodic physical measures, which has the basin covering prop- erty^2 erty^2 .

In [\[28\]](#page-42-9), to study the structure of physical measures for diffeomorphisms with mostly contracting center, inspired by [\[31\]](#page-42-10), Dolgopyat-Viana-Yang introduced a topological structure, which is known as skeleton. More recently, Yang [\[56\]](#page-43-10) slightly change the original definition of skeleton for diffeomorphisms with mostly expanding center. By constructing skeletons, they studied certain geometric structures of physical measures, which include the number for basins and supports of physical measures. Moreover, the study on stability of skeletons allow one can describe how physical measures bifurcate as the diffeomorphism changes under $C¹$ -topology. The question naturally arises whether one can find skeletons for diffeomorphisms in $PH_{EC}^1(M)$ and use them to study the physical measures in geometrical viewpoint?

For diffeomorphisms in $PH_{EC}^1(M)$, due to the mixture behavior occurs on centers, we propose a new definition of the skeleton which is different from those in both [\[28\]](#page-42-9) and [\[56\]](#page-43-10).

Definition 1.1. Let f be a diffeomorphism in $PH_{EC}^{1}(M)$ with dominated splitting $TM = E^u \oplus_{\succ} E^{cu} \oplus_{\succ} E^{cs}$. A set S consisting of finitely many hyperbolic periodic points of stable index $\dim E^{cs}$ is called a skeleton of f if (S1) for any C^1 disk γ transverse to E^{cs3} E^{cs3} E^{cs3} , there exists $p \in S$ such that

$$
\gamma \pitchfork W^s(\text{Orb}(p, f)) \neq \emptyset;
$$

 $(S2)$ $W^s(\text{Orb}(p, f)) \pitchfork W^u(\text{Orb}(q, f)) = \emptyset$ whenever $p, q \in S$ with $p \neq q$. In addition, we say that S is a pre-skeleton of f if it is a collection of finitely many hyperbolic periodic points of stable index $\dim E^{cs}$ and satisfies condition ([S1\)](#page-2-2).

The main difference between the present definition and those in [\[28,](#page-42-9) [56\]](#page-43-10) occurs on condition ([S1\)](#page-2-2). In mostly contracting case [\[28\]](#page-42-9), instead of ([S1\)](#page-2-2), there requires that every strong unstable manifold must intersect some stable manifold of periodic orbits of skeleton. The mostly expanding [\[56\]](#page-43-10) case requires the denseness of the union of stable manifolds of periodic orbits from skeleton, which is weaker than condition ([S1\)](#page-2-2).

In this paper, we show the existence of skeletons for diffeomorphisms in $PH_{EC}^{1}(M)$, and obtain a one-to-one correspondence between elements of a

 2 The union of the basins of all physical measures has a full Lebesgue measure.

³Meaning that $T_x\gamma \oplus E^{cs}(x) = T_xM$ for every $x \in \gamma$.

skeleton and physical measures for diffeomorphisms in $PH_{EC}^{1+}(M)$. Given a hyperbolic periodic point p of a diffeomorphism f , recall that the *homoclinic* class $H(p, f)$ of p is the closure of the set of transversal intersection points between the stable and unstable manifolds of $Orb(p, f)$.

Theorem A. Every $f \in PH_{EC}^1(M)$ admits skeletons. Moreover, if $f \in$ $PH_{EC}^{1+}(M)$, denoted by $\mathcal{P}(f) = {\mu_1, \cdots, \mu_k}$ the finitely many physical measures of f. Then for any skeleton $S(f) = \{p_1, \dots, p_\ell\}$ of f, there exists a bijective map $i \mapsto j(i)$ such that for every physical measure $\mu_i \in \mathcal{P}(f)$ and the corresponding $p_{i(i)} \in \mathcal{S}(f)$, we have

$$
supp(\mu_i) = \overline{W^u(\text{Orb}(p_{j(i)}, f))} = H(p_{j(i)}, f).
$$

In particular, we have $k = \ell$, i.e., the number of physical measures of f is equal to the cardinality of its skeleton.

Similar results were obtained for both mostly contracting case [\[28,](#page-42-9) Theorem A] and mostly expanding case [\[56,](#page-43-10) Theorem A]. Let us emphasize that the approaches of both works depend heavily on the uniform behaviors on either stable manifolds or unstable manifolds tangent to strong stable or unstable sub-bundles. In contrast to [\[28,](#page-42-9) [56\]](#page-43-10), for diffeomorphisms in $PH_{EC}^{1}(M)$, we have neither $E^{cu} = E^u$ nor $E^{cs} = E^s$, thus both E^{cu} and E^{cs} have non-uniformly behaviors. Therefore we use a different approach in the proof of Theorem [A.](#page-3-0) The following ingredients play central roles in the construction of skeletons.

- Hyperbolic periodic points in skeletons will be created near Pesin blocks admitting large weight associated to Gibbs u-states.
- We show that the forward orbit of typical point enters to small neighborhoods of periodic points obtained above at infinitely many hyperbolic times.

We mention that during the proof of Theorem [A,](#page-3-0) by utilizing structures of Gibbs cu-states established in this paper, we provide a new proof on the existence and basin covering property of physical measures, which is different from the original argument [\[40\]](#page-43-8). Let us remark that both of [\[28\]](#page-42-9) and [\[56\]](#page-43-10) stress that each closure of stable manifold of periodic orbits of skeleton coincides with the basin or essential basin of some physical measure. We do not know if it is possible to establish this property for diffeomorphisms in $PH_{EC}^{1+}(M).$

The idea of using skeletons or homoclinic classes to analyze typical measures, such as SRB measures, physical measures and maximal entropy measures appears in recent works (see e.g. [\[42,](#page-43-11) [16,](#page-42-11) [51\]](#page-43-12)).

Since $PH_{EC}^{1+}(M)$ is C^1 open, one would like to study how physical measures vary with respect to diffeomorphisms of $PH_{EC}^{1+}(M)$ under C^1 -topology. Theorem [A](#page-3-0) suggests that one can achieve this goal via analyzing the perturbed skeletons. More precisely, we have

Theorem B. Assume that $f \in PH_{EC}^{1+}(M)$ and $S(f) = \{p_1, \dots p_k\}$ is a skeleton of f. Then there exists a C^1 neighborhood V of f such that for any $g \in V$, the corresponding continuation $S(g) = \{p_1(g), \dots, p_k(g)\}\$ is a pre-skeleton of g. If $g \in V$ is of class C^{1+} , then the number of its physical measures is bounded above by the number of physical measures of f. Moreover, these two numbers coincide if and only if there exist no heteroclinic intersections between different elements of $S(g)$. For this situation, every physical measure of g is close to a physical of f , in weak^{*}-topology.

In addition, restricted to any subset of V where the number of physical measures is constant, the supports of the physical measures depends lower semi-continuously with dynamics, in the sense of Hausdorff topology.

By involving more technical details, we will prove Theorem [B](#page-4-0) in a more precisely manner in Section [7.](#page-31-0) Theorem [B](#page-4-0) gives the upper semi-continuity of the number of physical measures and the cardinality of skeleton. As a consequence, these numbers are locally constant on an open and dense subset of diffeomorphisms in $PH_{EC}^{1+}(M)$. As we mentioned, [\[28\]](#page-42-9) and [\[56\]](#page-43-10) obtained similar results for diffeomorphisms with mostly contracting and diffeomorphisms with mostly expanding center. Note that [\[8\]](#page-41-5) showed similar result for diffeomorphisms with mostly expanding center under $C^r(r > 1)$ topology, in which they didn't construct the skeletons.

Let F be a family of diffeomorphisms such that each $f \in \mathcal{F}$ admits finitely many physical measures, we say that F is *weakly statistically stable* if for every $f \in \mathcal{F}$, for every sequence $f_n \in \mathcal{F}$ that converges to f in C¹-topology, and for every sequence μ_n of physical measures of f_n , any limit measure of $\{\mu_n\}_{n\in\mathbb{N}}$ is in the convex hull of finitely many physical measures for f. In particular, if F consists of diffeomorphisms that each of which admits a unique physical measure, then we say that $\mathcal F$ is *statistically stable* if it is weakly statistically stable, or equivalently, the map $\mathcal{F} \ni f \mapsto \mu_f$ is continuous, where μ_f denotes the unique physical measure of f.

Theorem C. $PH_{EC}^{1+}(M)$ is weakly statistically stable. If $f \in PH_{EC}^{1+}(M)$ admits a unique physical measure, then there exists a C^1 neighborhood U of f such that $\mathcal{U} \cap PH_{EC}^{1+}(M)$ is statistically stable.

To the best of our knowledge, we give a partial list related to this result.

- Dolgopyat [\[27\]](#page-42-6) proved the statistical stability for some diffeomorphisms with mostly contracting center on three dimensional manifold.
- For diffeomorphisms with mostly contracting center, [\[6,](#page-41-8) [28\]](#page-42-9) achieved the (weak)statistical stability.
- Recently, [\[8\]](#page-41-5) and [\[56\]](#page-43-10) demonstrated the (weak) statistical stability for diffeomorphisms with mostly expanding center.

The method here is rather different to [\[27,](#page-42-6) [6,](#page-41-8) [8\]](#page-41-5), but with the same spirit of [\[28,](#page-42-9) [56\]](#page-43-10). Indeed, we deduce Theorem [C](#page-4-1) from Theorem [B.](#page-4-0) We point out [\[8\]](#page-41-5) used the transitivity assumption to guarantee the uniqueness of the physical measure. However, for diffeomorphisms admitting mostly contracting center, transitivity is insufficient to get the uniqueness of the physical measure, mainly due to the existence of intermingled basins^{[4](#page-5-0)}, which occurs even in robustly transitive settings. One can see $[10, \S 11.1.2], [35, 43, 24]$ for this phenomenon.

The remainder of this paper is organized as follows. Section [2](#page-5-1) consists of some notions and results used in this paper. In Section [3,](#page-9-0) we show the abundance of hyperbolic periodic points near Pesin blocks, these periodic points are candidates of the elements of skeletons. Section [4](#page-12-0) is devoted to study the structure of the set of Gibbs cu -states, such as compactness and the upper semi-continuity with respect to diffeomorphisms. Some preliminary result on properties of skeletons are given in Section [5,](#page-18-0) which will be used in next two sections. In Section [6,](#page-20-0) we introduce the set $PH_C^1(M)$ of partially hyperbolic diffeomorphims for which every diffeomorphism in $PH_{EC}^1(M)$ has its iterates in $PH_C^1(M)$. For diffeomorphisms in $PH_C^1(M)$, we obtain the existence of skeletons, and the one-to-one corresponding between physical measures and periodic points of the skeleton (Theorem [D\)](#page-20-1). In Section [7,](#page-31-0) going back to $PH_{EC}^1(M)$, we give the proofs of Theorems [A,](#page-3-0) [B](#page-4-0) and [C.](#page-4-1) We provide a detailed proof of Proposition [3.5](#page-11-0) in Appendix [A,](#page-37-0) which is stated in Section [3](#page-9-0) and used almost throughout this paper.

Acknowledgements. We are grateful to D. Yang and R. Zou for their suggestions and discussions.

2. Preliminary

Throughout, let M be a compact Riemannian manifold with metric d . Let $B(x,r)$ be the r-ball of $x \in M$ defined by $B(x,r) = \{y \in M : d(x,y) < r\}.$ Given a smooth embedded sub-manifold D, let $d_D(x, y)$ be the distance from x to y along D; and for $\delta > 0$, define $B_D(x, \delta) = \{y \in D : d_D(x, y) < \delta\}$. We use Leb_D to represent the induced Lebesgue measure on D. We say that D is an unstable disk if it is a disk inside an unstable manifold.

Denote by M the space of Borel probability measures supported on M . Let us fix a metric compatible with the weak*-topology on $\mathcal M$ as follows: for μ and ν in M, define

$$
dist(\mu, \nu) = \sum_{n \in \mathbb{N}} \frac{\left| \int \varphi_n d\mu - \int \varphi_n d\nu \right|}{2^n \sup_{x \in M} |\varphi_n(x)|},
$$

where $\{\varphi_n\}_{n\in\mathbb{N}}$ is a countable dense subset of the space of continuous functions on M. For a homoemorphism f on M, let $\mathcal{M}(f)$ be the set of finvariant measures supported on M.

⁴For two physical measures μ and ν with respect to f, we say that they are intermingled if Leb($U \cap B(\mu, f) > 0$ and Leb($U \cap B(\nu, f) > 0$ for any open set U of M.

Let f be a diffeomorphism and E a Df -invariant sub-bundle, for every $\mu \in \mathcal{M}(f)$, define

$$
\chi_{\mu}(f,E) = \int \log \|Df|_{E(x)} \|d\mu(x).
$$

2.1. Plaque families for diffeomorphisms with dominated splitting. Let f be a C^1 diffeomorphism with dominated splitting $TM = E \oplus F$. Given $\theta > 0$, define the *cone field* $\mathcal{C}_{\theta}^{E} = {\mathcal{C}_{\theta}^{E}(x) : x \in M}$ associated to E of width θ by

$$
\mathcal{C}_{\theta}^{E}(x) = \{ v = v^{E} + v^{F} \in E(x) \oplus F(x) : ||v^{F}|| \leq \theta ||v^{E}|| \}
$$

for every $x \in M$. By symmetry, we can define the cone field \mathcal{C}_{θ}^{F} associated to F. We say that a smooth embedded sub-manifold D is tangent to \mathcal{C}_{θ}^{E} if $\dim D = \dim E$ and $T_x D \subset \mathcal{C}_{\theta}^E(x)$ for every $x \in D$. In particular, one says that D is tangent to E if $T_x D = E(x)$ for every $x \in D$.

It is known that the domination is robust in the C^1 -topology. More precisely, if C^1 diffeomorphism f admits the dominated splitting $TM = E \oplus$ F, then there exists a C^1 neighborhood U of f so that every $g \in U$ admits the dominated splitting $TM = E_g \oplus F_g^5$ $TM = E_g \oplus F_g^5$. Denote by $E_g(x, \rho)$ the ρ -neighborhood of 0_x in $E_g(x)$, define $F_g(x, \rho)$ similarly. For diffeomorphisms with dominated splitting, one has the Plaque family theorem stated as follows.

Proposition 2.1. [\[32,](#page-42-14) Theorem 5.5] Let f be a C^1 diffeomorphism with dominated splitting $TM = E \oplus F$. Then for every $\theta > 0$, there exists a C^1 open neighborhood U of f and constants $0 < \delta < \rho$ such that for any $g \in \mathcal{U}$, there is a continuous family of C^1 embedded sub-manifolds $\{F^E_\rho(g,x)\}_{x\in M}$ given by

$$
\mathcal{F}^E_{\rho}(g, x) = \exp_x\{(u, \phi_x(u)) : u \in E_g(x, \rho)\}\
$$

where $\phi_x : E_g(x, \rho) \to F_g(x)$ is a C^1 map such that $\phi_x(0_x) = 0_x$ and $D\phi_x(0_x) = 0$. Similarly, one can get the continuous family of C^1 embedded $sub-manifolds \{F_{\rho}^{F}(g,x)\}_{x \in M}$. For every $*\in \{E, F\}$ and $x \in M$ one has

- $\mathcal{F}_{\rho}^*(g,x)$ is tangent to $\mathcal{C}_{\theta}^*(x)$;
- $g(\mathcal{F}_{\delta}^*(g,x)) \subset \mathcal{F}_{\rho}^*(g,g(x))$ and $g^{-1}(\mathcal{F}_{\delta}^*(g,x)) \subset \mathcal{F}_{\rho}^*(g,g^{-1}(x));$
- for $g_n \xrightarrow{C^1} g$ in U and $x_n \to x$, we have $\mathcal{F}^*_{\rho}(g_n, x_n) \to \mathcal{F}^*_{\rho}(g, x)$.

2.2. Hyperbolic time and Liao-Gan's shadowing Lemma. For $A \subset$ N, define the *lower-density* $\mathcal{D}_L(A)$ and the *upper-density* $\mathcal{D}_U(A)$ of A as follows:

$$
\mathcal{D}_L(A) := \liminf_{n \to +\infty} \frac{1}{n} \# \{ A \cap [1, n] \}, \quad \mathcal{D}_U(A) := \limsup_{n \to +\infty} \frac{1}{n} \# \{ A \cap [1, n] \}.
$$

Now we give the definition of hyperbolic times associated to some invariant sub-bundles.

 ${}^{5}E_{q}(x)$ and $F_{q}(x)$ depend continuously on q and x. In particular, $E_{f}(x) = E(x)$ and $F_f(x) = F(x)$ for every $x \in M$. Sometimes, we drop the reference to g for simplicity.

Definition 2.1. Let f be a C^1 diffeomorphism with a Df-invariant subbundle E. Given $\sigma \in (0,1)$, we call $n \in \mathbb{N}$ a (E,σ) -hyperbolic time of $x \in M$ if

$$
\prod_{i=n-k+1}^{n} \|Df^{-1}|_{E(f^{i}(x))}\| < \sigma^{k} \text{ for } k = 1, \cdots, n.
$$

For diffeomorphisms with dominated splitting, one has the backward contracting property for disks tangent to the cone field at hyperbolic times, see [\[2,](#page-41-10) Lemma 2.7] for a proof.

Lemma 2.2. Let f be a C^1 diffeomorphism admitting the dominated splitting $TM = E \oplus_{\succ} F$. Given $0 < \sigma_0 < \sigma_1 < 1$, there exist $a > 0, \delta > 0$ such that, for any C^1 embedded disk D tangent to \mathcal{C}_{a}^E that contains x, if n is a (E, σ_0) -hyperbolic time of x, then

$$
d_{f^i(D)}(f^i(y), f^i(z)) \le \sigma_1^{(n-i)} d_{f^n(D)}(f^n(y), f^n(z))
$$

for any y, z in D such that $f^{n}(y), f^{n}(z) \in B_{f^{n}(D)}(f^{n}(x), \delta)$.

As an application of Pliss Lemma (see [\[45\]](#page-43-14) and [\[2,](#page-41-10) Lemma 3.1]), one has

Proposition 2.3. [\[2,](#page-41-10) Corollary 3.2] Suppose $\alpha_0 > \alpha > 0$, and let f be a C^1 diffeomorphism admitting dominated splitting $TM = E \oplus F$, then there exists $\theta = \theta(\alpha_0, \alpha, f) \in (0, 1)$ and a C^1 neighborhood U of f such that for any $g \in \mathcal{U}$, if x is α_0 -nonuniformly expanding along E_q , i.e.,

$$
\limsup_{n \to +\infty} \frac{1}{n} \sum_{i=1}^{n} \log ||Dg^{-1}|_{E_g(g^i(x))}|| < -\alpha_0,
$$

then there exist infinitely many $(E_g, e^{-\alpha})$ -hyperbolic times $n_1 < n_2 < \cdots$ of x with lower density larger than θ .

Let us recall the Liao-Gan's shadowing lemma [\[39,](#page-43-15) [30\]](#page-42-15).

Lemma 2.4. Let f be a C^1 diffeomorphism with dominated splitting $TM =$ $E \oplus_{\succ} F$. For $\lambda \in (0,1)$, there exists $\delta_0 > 0$ and $L_0 > 0$ such that for any orbit segment $(x, f(x), \dots, f^n(x))$ with following properties:

$$
d(x, f^{n}(x)) \leq \delta_0;
$$

• for any $1 \leq k \leq n$, we have

•

$$
\prod_{i=0}^{k-1} \|Df^{-1}|_{E(f^{n-i}(x))} \| < \lambda^k, \quad \prod_{i=0}^{k-1} \|Df|_{F(f^i(x))} \| < \lambda^k,
$$

there exists a hyperbolic periodic point p with stable index dimF such that

$$
d(f^i(x), f^i(p)) \le L_0 d(x, f^n(x)) \quad \text{for any } 0 \le i \le n - 1.
$$

2.3. Entropy along unstable foliations. Let f be a $C¹$ partially hyperbolic diffeomorphism with E^u , denote by \mathcal{F}^u the unstable foliation whose leaves are strong unstable manifolds tangent to E^u .

Let μ be a Borel probability. A measurable partition ξ is μ -subordinate to \mathcal{F}^u , if for μ -almost every x, one has

- $\xi(x) \subset \mathcal{F}^u(x)$, where $\xi(x)$ is the element of ξ containing x;
- $\xi(x)$ contains an open neighborhood of x in \mathcal{F}^u ;
- $\xi(f(x)) \subset f(\xi(x)).$

Note that the existence of measurable partitions μ -subordinate to unstable foliation was given by [\[36,](#page-42-16) Lemma 3.1] and [\[55,](#page-43-9) Lemma 3.2]. Moreover, we have the following result which is contained in [\[38,](#page-43-16) Lemma 3.1.2].

Lemma 2.5. For any $\mu \in \mathcal{M}(f)$ and any measurable partitions ξ_1 and ξ_2 that are μ -subordinate to \mathcal{F}^u , we have

$$
h_{\mu}(f,\xi_1) = h_{\mu}(f,\xi_2).
$$

By Lemma [2.5,](#page-8-1) one can define the entropy along \mathcal{F}^u as follows:

Definition 2.2. For every $\mu \in \mathcal{M}(f)$, the entropy of μ along \mathcal{F}^u is defined by

$$
h_{\mu}(f, \mathcal{F}^u) = h_{\mu}(f, \xi),
$$

where ξ is any measurable partition μ -subordinated to \mathcal{F}^u .

The entropy $h_\mu(f, \mathcal{F}^u)$ has become a powerful tool in the study of partially hyperbolic diffeomorphisms in recent years, see [\[26,](#page-42-2) [33,](#page-42-17) [34,](#page-42-18) [56\]](#page-43-10) for instance.

Definition 2.3. Let f be a C^1 partially hyperbolic diffeomorphism with E^u . We call $\mu \in \mathcal{M}(f)$ a Gibbs u-state if

$$
h_{\mu}(f, \mathcal{F}^u) = \int \log |\det Df|_{E^u}|d\mu.
$$

As we have mentioned, if f is C^{1+} , then μ is a Gibbs u-state if and only its conditional measures along strong unstable manifolds are absolutely continuous w.r.t. Lebesgue measures on these manifolds. Denote by $G^u(f)$ the space of Gibbs u-states of f .

Lemma 2.6. [\[26,](#page-42-2) Corollary 2.15] Let f be a C^1 partially hyperbolic diffeomorphism with E^u . Then

- (1) $G^u(f)$ is nonempty, compact and convex subset of $\mathcal{M}(f)$.
- (2) Every ergodic component of any $\mu \in G^u(f)$ is a Gibbs u-state.

Moreover, we have the upper semi-continuity of the map $f \mapsto G^u(f)$.

Lemma 2.7. [\[34,](#page-42-18) Proposition 3.5] Let $\{f_n\}_{n\in\mathbb{N}}$ and f be C^1 partially hyperbolic diffeomorphisms exhibiting E^u , and assume that f_n converges to f in C^1 -topology, then

$$
\limsup_{n \to +\infty} G^u(f_n) \subset G^u(f).
$$

3. Abundance of hyperbolic periodic points

Let f be a C^1 diffeomorphism with dominated splitting $TM = E \oplus_{\succ} F$. For $\ell \in \mathbb{N}, \alpha > 0$, we introduce the following *Pesin block*:

$$
\Lambda_{\ell}(f, \alpha, E, F) = \{x : \frac{1}{n\ell} \sum_{i=0}^{n-1} \log ||Df^{-\ell}|_{E(f^{-i\ell}(x))}|| \leq -\alpha, \frac{1}{n\ell} \sum_{i=0}^{n-1} \log ||Df^{\ell}|_{F(f^{i\ell}(x))}|| \leq -\alpha, \quad \forall n \in \mathbb{N}\}.
$$

Remark 3.1. Note that the Pesin block defined here is different from the classical one in Pesin theory (see [\[9\]](#page-41-11)).

As usual, all points of a Pesin block admit the stable and unstable manifolds with uniform size. Moreover, this size can be made to be uniform in a neighborhood of the fixed diffeomorphism.

Lemma 3.2. Assume that f is a C^1 diffeomorphism with dominated splitting $TM = E \oplus_{\succ} F$.

Then for every $\alpha > 0$ and $\ell \in \mathbb{N}$, there exists a C^1 neighborhood U of f and constants $\tau := \tau(\alpha, \ell) \in (0, 1), C := C(\alpha, \ell) > 0, \delta := \delta(\ell, \alpha) > 0$ such that for every $g \in \mathcal{U}$, for every $x \in \Lambda_{\ell}(g, \alpha, E, F)$, there are local stable manifold $W^s_{loc}(x,g)$ and local unstable manifold $W^u_{loc}(x,g)$, which are C^1 embedded disks of radius δ centered at x with following properties:

- $W^s_{loc}(x,g)$ is tangent to E and $W^u_{loc}(x,g)$ is tangent to F;
- for every $n \in \mathbb{N}$, we have
	- $-d(g^{n}(y), g^{n}(z)) \leq C\tau^{n}d(y,z)$ for every $y, z \in W^{s}_{loc}(x,g)$, $- d(g⁻ⁿ(y), g⁻ⁿ(z)) ≤ Cτⁿd(y, z) for every y, z ∈ W_{loc}^u(x, g).$

It can be deduced from Plaque family theorem (Proposition [2.1\)](#page-6-1), note that the local stable and unstable manifolds are contained in the corresponding plaques, see [\[8,](#page-41-5) Theorem 3.5] for a detailed proof. The radius δ is also called the size of stable(unstable) manifolds. For the rest, we use $R(W^s_{loc}(x, g))$ and $R(W_{loc}^u(x,g))$ to denote the sizes of $W_{loc}^s(x,g)$ and $W_{loc}^u(x,g)$, respectively.

By Lemma [2.4,](#page-7-0) we have the following result which demonstrates the existence of hyperbolic periodic points near Pesin blocks.

Lemma 3.3. Let f be a C^1 diffeomorphism with dominated splitting $TM =$ $E \oplus_{\succ} F$. Then for any $\alpha > 0$ and $\ell \in \mathbb{N}$, there exist $\rho_{\alpha,\ell} > 0$, $\delta_{\alpha,\ell} > 0$ and $L_{\alpha,\ell} > 0$ such that if

$$
\mu(B(x,\rho) \cap \Lambda_{\ell}(f,\alpha,E,F)) > 0,
$$

for some $x \in M$, $\mu \in \mathcal{M}(f)$ and $\rho \leq \rho_{\alpha,\ell}$, then there exists a hyperbolic periodic point $p \in B(x, L_{\alpha,\ell} \cdot \rho)$ of stable index dim F such that

$$
R(W_{loc}^s(p,f)) \ge \delta_{\alpha,\ell}, \quad R(W_{loc}^u(p,f)) \ge \delta_{\alpha,\ell}.
$$

Proof. Writing $\Lambda_{\ell} := \Lambda_{\ell}(f, \alpha, E, F)$ for simplicity. Given $n \in \mathbb{N}$, assume that y and $f^{(n)}(y)$ are contained in Λ_{ℓ} , then from the definition of Λ_{ℓ} one obtains that for every $1 \leq j \leq n$,

$$
(3.2) \quad \prod_{i=0}^{j-1} \|Df^{\ell}|_{F(f^{i\ell}(y))} \| \le e^{-\alpha j\ell} \quad \text{and} \quad \prod_{i=0}^{j-1} \|Df^{-\ell}|_{E(f^{n-i\ell}(y))} \| \le e^{-\alpha j\ell}.
$$

By Lemma [2.4,](#page-7-0) there exist $\delta_0 > 0$ and $L_0 > 0$ such that when $d(y, f^n(y)) \leq$ δ_0 , then there exists a hyperbolic periodic point p of periodic n with stable index $\dim F$, which satisfies

(3.3)
$$
d(f^i(y), f^i(p)) \le L_0 d(y, f^n(y)) \text{ for every } 1 \le i \le n.
$$

Now let us take $\rho_{\alpha,\ell} = \delta_0/2$ and $L_{\alpha,\ell} = 2L_0 + 1$. For every $0 < \rho \leq \rho_{\alpha,\ell}$, assume $\mu(B(x,\rho) \cap \Lambda_{\ell}) > 0$ for some $x \in M$ and $\mu \in \mathcal{M}(f)$. By Poincare's recurrent theorem [\[57,](#page-43-17) Theorem 1.4], μ -almost every point of $B(x,\rho) \cap \Lambda_{\ell}$ is recurrent, therefore one can take $y, f^{n}(y) \in B(x, \rho) \cap \Lambda_{\ell}$ for infinitely many n, and thus we have $d(y, f^{n}(y)) < 2\rho \leq \delta_0$. Consequently, there exists a hyperbolic periodic point p with stable index dim F , which satisfies [\(3.3\)](#page-10-0) and $p \in B(x, L_{\alpha,\ell} \cdot \rho)$. Since y exhibits property [\(3.2\)](#page-10-1), one can take $\rho_{\alpha,\ell}$ small enough such that for every $j \geq 1$,

$$
\prod_{i=0}^{j-1} \left\| Df^{\ell}|_{F(f^{i\ell}(p))} \right\| \le e^{-\frac{\alpha}{2}j\ell} \quad \text{and} \quad \prod_{i=0}^{j-1} \left\| Df^{-\ell}|_{E(f^{-i\ell}(p))} \right\| \le e^{-\frac{\alpha}{2}j\ell}.
$$

Therefore, $p \in \Lambda_{\ell}(f, \alpha/2, E, F)$. According to Lemma [3.2,](#page-9-1) there exists $\delta_{\alpha,\ell}$ 0 such that the stable and unstable manifolds of p have size larger than $\delta_{\alpha,\ell}$.

When the invariant measure is ergodic, we have the following result which will be used for several times.

Lemma 3.4. Let f be a C^1 diffeomorphism with dominated splitting $TM =$ $E \oplus_{\succ} F$. Then for any $\alpha, \alpha' > 0$ and $\ell, \ell' \in \mathbb{N}$, there exist $\delta := \delta(\alpha, \alpha', \ell, \ell') >$ $0, \rho := \rho(\alpha, \alpha', \ell, \ell') > 0$ and $L := L(\alpha, \alpha', \ell, \ell') > 0$ such that if

$$
(3.4) \quad \mu\big(B(x,r)\cap \Lambda_{\ell}(f,\alpha,E,F)\big) > 0, \quad \mu\big(B(y,r)\cap \Lambda_{\ell'}(f,\alpha',E,F)\big) > 0,
$$

for some f-ergodic measure μ , points $x, y \in M$ and $r \leq \rho$, then there exists a hyperbolic periodic point $p \in B(x, Lr)$ such that $f^m(p) \in B(y, Lr)$ for some $m \in \mathbb{N}$, and both of them admit local stable and unstable manifolds of size larger than δ .

Proof. By Lemma [2.4,](#page-7-0) there exist $L_0 > 0$ and $\rho > 0$ such that for any $r \leq \rho$, for any $z \in B(x,r) \cap \Lambda_{\ell}(f,\alpha,E,F)$ with $f^{n}(z) \in B(x,r) \cap \Lambda_{\ell}(f,\alpha,E,F)$, there exists a hyperbolic periodic point p of periodic n such that

(3.5)
$$
d(f^k(p), f^k(z)) \le L_0 d(z, f^n(z)) \text{ for every } 1 \le k \le n.
$$

Given $r \leq \rho$, f-ergodic measure μ and points $x, y \in M$ satisfying [\(3.4\)](#page-10-2). By Poincare's recurrent theorem, μ -almost every point of $B(x,r) \cap \Lambda_\ell(f,\alpha,E,F)$

is recurrent. Moreover, since $\mu(B(y, r) \cap \Lambda_{\ell'}(f, \alpha', E, F)) > 0$ and μ is ergodic, the Birkhoff's ergodic theorem [\[57,](#page-43-17) Theorem 1.14] implies that for μ -almost point point, its forward orbit enters to $B(y,r) \cap \Lambda_{\ell'}(f,\alpha',E,F)$ for infinitely many times. Therefore, one can take a recurrent point z of $B(x,r) \cap \Lambda_{\ell}(f,\alpha,E,F)$ with some integers $m < n$ such that

$$
- z, f^{n}(z) \in B(x, r) \cap \Lambda_{\ell}(f, \alpha, E, F);
$$

$$
- f^{m}(z) \in B(y, r) \cap \Lambda_{\ell'}(f, \alpha', E, F).
$$

Hence, one can find a hyperbolic periodic point p of periodic n with property [\(3.5\)](#page-10-3). Furthermore, by a simple computation, one can take $L > L_0$ such that $p \in B(x, Lr)$ and $f^m(p) \in B(y, Lr)$. By definition of Pesin blocks, as showed in Lemma [3.3,](#page-9-2) by reducing ρ if necessary, one can take $\delta > 0$ such that both p and $f^m(p)$ admit local stable and unstable manifolds of size larger than δ .

We are interested in diffeomorphisms in $PH_{EC}^1(M)$, thus for every $f \in$ $PH_{EC}^{1}(M)$ with partially hyperbolic splitting $TM = E^{u} \oplus_{\succ} E^{cu} \oplus_{\succ} E^{cs}$, $\alpha > 0$ and $\ell \in \mathbb{N}$, we can rewrite $E^{wu} := E^u \oplus_{\succ} E^{cu}$ and $\Lambda_{\ell}(f, \alpha) :=$ $\Lambda_{\ell}(f, \alpha, E^{wu}, E^{cs})$ for simplicity.

The result below may be seen as a C^1 version of [\[8,](#page-41-5) Lemma 3.4], though we are dealing with a more general system than [\[8\]](#page-41-5). For completeness, a proof is provided in Appendix [A.](#page-37-0)

Proposition 3.5. For every $f \in PH_{EC}^1(M)$, there exists $\alpha > 0$ such that for every $\varepsilon > 0$, there exists $\ell \in \mathbb{N}$ and αC^1 neighborhood U of f, such that for every $g \in \mathcal{U}$, every $\mu \in G^u(f)$, it has

$$
\mu\left(\Lambda_{\ell}(g,\alpha)\right) > 1 - \varepsilon.
$$

Now we give the abundance of hyperbolic periodic points near Pesin blocks described as follows:

Theorem 3.6. For every $f \in PH_{EC}^1(M)$ with partially hyperbolic splitting $TM = E^u \oplus_{\succ} E^{cu} \oplus_{\succ} E^{cs}$, there exists $\alpha > 0$ such that for every $\varepsilon > 0$ there are $\ell \in \mathbb{N}$, $\rho_{\ell} > 0$, $\delta_{\ell} > 0$ and C^1 neighborhood U of f such that for each $q \in \mathcal{U}$, we have

$$
R(W_{loc}^s(x,g)) \ge \delta_\ell, \quad R(W_{loc}^u(x,g)) \ge \delta_\ell
$$

for all $x \in \Lambda_{\ell}(g, \alpha)$. In addition, for every $\rho \leq \rho_{\ell}$ and $\mu \in G^u(g)$, there is an open covering

$$
\mathscr{A}_{\mu}(g,\ell,\alpha,\rho) = \left\{ B(x_i,\rho) : x_i \in \text{supp}(\mu | \Lambda_{\ell}(g,\alpha)), 1 \leq i \leq k \right\}
$$

of supp $(\mu|\Lambda_{\ell}(q,\alpha))$ for some $k \in \mathbb{N}$, which admits following properties:

• each $B \in \mathscr{A}_{\mu}(g,\ell,\alpha,\rho)$ contains some hyperbolic periodic point, which we denote by p_B , so that

$$
R(W^s_{loc}(p_B, g)) \ge \delta_\ell, \quad R(W^u_{loc}(p_B, g)) \ge \delta_\ell;
$$

$$
\bullet
$$

$$
\mu\left(\bigcup_{B\in\mathscr{A}_\mu(g,\ell,\alpha,\rho)}(B\cap\Lambda_\ell(g,\alpha))\right)>1-\varepsilon.
$$

Proof. By Proposition [3.5,](#page-11-0) there exists $\alpha > 0$ such that for every $\varepsilon > 0$, there are $\ell \in \mathbb{N}$ and C^1 neighborhood $\mathcal U$ of f, such that $\mu(\Lambda_{\ell}(g, \alpha)) > 1 - \varepsilon$ for every $g \in \mathcal{U}$ and $\mu \in G^u(g)$. By Lemma [3.2](#page-9-1) and Lemma [3.3,](#page-9-2) there exist $\rho_{\ell} > 0$ and $\delta_{\ell} > 0$ such that for every $g \in \mathcal{U}$ and $\mu \in G^u(g)$, we have the following properties:

- (1) every point of $\Lambda_{\ell}(g,\alpha)$ admits local stable and unstable manifolds of size larger than δ_{ℓ} ;
- (2) the ball $B(x, \rho)$ for any $\rho \leq \rho_{\ell}$ and any $x \in \text{supp}(\mu | \Lambda_{\ell}(g, \alpha))$ contains hyperbolic periodic points, all of which admit stable and unstable manifolds of size larger than δ_{ℓ} .

Due to the compactness, for every $\rho \leq \rho_{\ell}$ there exists $k \in \mathbb{N}$, a sequence of points x_1, \dots, x_k in supp $(\mu | \Lambda_\ell(g, \alpha))$ and the family of ρ -balls

$$
\mathscr{A}_{\mu}(g,\ell,\alpha,\rho) = \left\{ B(x_i,\rho) : x_i \in \text{supp}(\mu | \Lambda_{\ell}(g,\alpha)), 1 \leq i \leq k \right\}
$$

covering supp $(\mu|\Lambda_{\ell}(g,\alpha))$. By [\(2\)](#page-12-1), for every $B \in \mathscr{A}_{\mu}(g,\ell,\alpha,\rho)$ one can fix the hyperbolic periodic point $p_B \in B$ to satisfy

$$
R(W^s_{loc}(p_B, g)) \ge \delta_\ell, \quad R(W^u_{loc}(p_B, g)) \ge \delta_\ell.
$$

Since supp $(\mu|\Lambda_{\ell}(g,\alpha))$ exhibits μ -measure larger than $1-\varepsilon$, by construction of $\mathscr{A}_{\mu}(g,\ell,\alpha,\rho)$ one gets

$$
\mu\left(\bigcup_{1\leq i\leq k} \left(B(x_i,\rho)\cap\Lambda_\ell(g,\alpha)\right)\right) > 1-\varepsilon.
$$

Now we complete the proof of Theorem [3.6.](#page-11-1)

4. Cesaro limit measures of iterates

For $x \in M$, let $\omega_{\mathcal{M}}(x, f)$ be the set of limit measures of $\{\frac{1}{n}\}$ $\frac{1}{n}\sum_{i=0}^{n-1} \delta_{f^i(x)}\}_{n\in\mathbb{N}}.$ Similarly, for every C^1 embedded sub-manifold D, denote by $\omega_{\mathcal{M}}(D, f)$ the set of limit measures of $\{\frac{1}{n}\}$ $\frac{1}{n}\sum_{j=0}^{n-1} f_*^j$ m_D}_{n∈N}, where m_D stands for the normalized Lebesgue measure on D.

Let $f \in PH_{EC}^1(M)$ with partially hyperbolic splitting $TM = E^u \oplus_{\succ}$ $E^{cu} \oplus_{\succ} E^{cs}$, writing $E^{wu} = E^u \oplus E^{cu}$ as before. Let us consider the following subspace charactered by entropy:

$$
G^{cu}(f) = \left\{ \mu \in \mathcal{M}(f) : \mu \in G^u(f), \quad h_{\mu}(f) = \int \log |\det Df|_{E^{wu}}| d\mu \right\}.
$$

We call $G^{cu}(f)$ the Gibbs cu-space of f, whose elements are called the Gibbs cu-states of f.

Remark 4.1. Since every $\mu \in G^u(f)$ has only negative Lyapunov exponents along E^{cs} , applying the Ruelle's entropy inequality [\[47\]](#page-43-18), one knows that $h_{\mu}(f) \leq \int \log |\det Df|_{E^{wu}}| d\mu$, and therefore the equality in $G^{cu}(f)$ can be replaced by $h_{\mu}(f) \geq \int \log |\det Df|_{E^{wu}}| d\mu$.

Notation. For every $f \in PH_{EC}^1(M)$, consider \mathcal{B}_f as the set of points whose Cesaro limit measures of iterates are contained in $G^{cu}(f)$, that is

 $\mathcal{B}_f := \{x \in M : \omega_{\mathcal{M}}(x, f) \subset G^{cu}(f)\}.$

The main goal of this section is to show the following result.

Theorem 4.2. Let $f \in PH_{EC}^1(M)$ with partially hyperbolic splitting $TM =$ $E^u \oplus_{\succ} E^{cu} \oplus_{\succ} E^{cs}$. If D is a C¹ disk transverse to E^{cs} , then we have

(1) the intersection of \mathcal{B}_f with D has full Leb_D-measure;

 $(2) \omega_{\mathcal{M}}(D, f) \subset G^{cu}(f).$

In addition, as a result of [\(1\)](#page-13-0), we know that \mathcal{B}_f has full Lebesgue measure on M.

Let us recall that μ is said to be an *SRB measure* of a C^{1+} diffeomorphism if it has positive Lyapunov exponents almost everywhere; and its conditional measures along Pesin unstable manifolds are absolutely continuous w.r.t. Lebesgue measures along these manifolds. As a consequence of Theorem [4.2,](#page-13-1) we have the following corollary.

Corollary 4.3. Let $f \in PH_{EC}^{1+}(M)$ with partially hyperbolic splitting $TM =$ $E^u \oplus_{\succ} E^{cu} \oplus_{\succ} E^{cs}$. If D is a C^1 disk transverse to E^{cs} , then any invariant measure of $\omega_{\mathcal{M}}(D, f)$ is an SRB measure of f. Moreover, for Leb_D-almost every point $x \in D$, any invariant measure of $\omega_{\mathcal{M}}(x, f)$ is an SRB measure.

By assuming Theorem [4.2](#page-13-1) we give the proof of Corollary [4.3](#page-13-2) immediately.

Proof of Corollary [4.3.](#page-13-2) By Theorem [4.2,](#page-13-1) we need only to show that any invariant measure of $G^{cu}(f)$ is an SRB measure. Since every $\mu \in G^{cu}(f)$ is a Gibbs u-state, it admits only positive Lyapunov exponents along E^{cu} and negative Lyapunov exponents along E^{cs} . This together with

$$
h_{\mu}(f) = \int \log |\text{det}Df|_{E^{wu}}|d\mu
$$

implies that μ must be an SRB measure of f, using the classical result [\[38,](#page-43-16) Theorem A.

4.1. Structure on Gibbs cu -spaces. We have the upper semi-continuity of metric entropy for Gibbs u-states and diffeomorphisms in $PH_{EC}^1(M)$.

Theorem 4.4. Let $\{f_n\}$ be a sequence of diffeomorphisms in $PH_{EC}^1(M)$ which converges to $f \in PH_{EC}^1(M)$ in C^1 -topology. If $\mu_n \in G^u(f_n)$ for every n and $\mu_n \to \mu$ as $n \to +\infty$, then $\mu \in G^u(f)$ and

$$
\limsup_{n \to +\infty} h_{\mu_n}(f_n) \le h_{\mu}(f).
$$

It follows from Theorem [4.4](#page-13-3) that we have the following theorem, which can be proved in a standard way as in [\[56,](#page-43-10) Proposition 5.17].

Theorem 4.5. The map $PH_{EC}^{1}(M) \ni f \mapsto G^{cu}(f)$ is upper semi-continuous. For every $f \in PH_{EC}^1(M)$, the following properties hold:

- (1) $G^{cu}(f)$ is compact and convex, all its extreme elements are ergodic;
- (2) if $f \in PH_{EC}^{1+}(M)$, then all the extreme elements of $G^{cu}(f)$ are physical measures.

Now we make effort to prove Theorem [4.4.](#page-13-3) Let us recall the definition of tail entropy.

Definition 4.1. Given $r > 0$ and $x \in M$, define the r-tail entropy^{[6](#page-14-0)} at x by $h^*(f, x, r) = h_{top}(f, B_{\pm \infty}(f, x, r)),$

where $B_{\pm\infty}(f, x, r) = \{y : d(f^{n}(x), f^{n}(y)) \le r, n \in \mathbb{Z}\}\)$. For any $\mu \in \mathcal{M}(f)$, define the r-tail entropy of μ by

$$
h^*(f, \mu, r) = \int h^*(f, x, r) d\mu(x).
$$

Lemma 4.6. [\[19,](#page-42-19) Theorem 1.2] Assume that f is a homomorphism on M with finite topological entropy. Then for any $\mu \in \mathcal{M}(f)$ and $r > 0$, we have

$$
h_{\mu}(f) - h_{\mu}(f, \mathcal{P}) \le h^*(f, \mu, r),
$$

where P is any finite measurable partition with $\text{diam}(\mathcal{P}) \leq r$.

By domination and the invariance of plaque families, one has the following result.

Lemma 4.7. [\[41,](#page-43-19) Theorem 2.1] If f is a C^1 diffeomorphism with dominated splitting $TM = E \oplus_{\succ} F$, then there exists a C^1 neighborhood U of f and constants $0 < r < \delta$ such that for every $g \in \mathcal{U}$ and any g-ergodic measure μ one has

- either $B_{\pm\infty}(g, x, r) \subset \mathcal{F}_{\delta}^{E}(g, x)$ for μ -almost every x,
- or $B_{\pm\infty}(g, x, r) \subset \mathcal{F}_{\delta}^F(g, x)$ for μ -almost every x.

Note that this result is also proved in [\[19,](#page-42-19) Lemma 2.3] for fake foliations.

Lemma 4.8. For any $f \in PH_{EC}^1(M)$, there exists $r > 0$ and a C^1 neighborhood U of f such that $h^*(g, \mu, r) = 0$ for every $g \in \mathcal{U}$ and $\mu \in G^u(g)$.

Proof. Let $f \in PH_{EC}^1(M)$ with partially hyperbolic splitting $TM = E^u \oplus_{\succ}$ $E^{cu} \oplus_{\succ} E^{cs}$. By Proposition [3.5,](#page-11-0) there exists a C^1 neighborhood $\mathcal U$ of f, and $\alpha > 0, \, \ell \in \mathbb{N}$ such that $\mu(\Lambda_{\ell}(g, \alpha)) > 0$ for every $\mu \in G^u(g)$ and $g \in \mathcal{U}$.

By definition of tail entropy, it suffices to show that there exists $r > 0$ such that for every ergodic $\mu \in G^u(g)$, one has $B_{\pm \infty}(g, x, r) = \{x\}$ for μ almost every x. Now we fix $\mu \in G^u(g)$ and $g \in \mathcal{U}$. Following Lemma [4.7,](#page-14-1)

 6 By the result of [\[17,](#page-42-20) Proposition 2.8], $h^*(f, x, r)$ is measurable and f-invariant.

without loss of generality, we assume that for μ -almost every x, $B_{\pm\infty}(q, x, r)$ is contained in the E^{wu} -direction plaque provided r is small enough. Then using the invariance property of plaques in Proposition [2.1,](#page-6-1) we have that all the forward iterates of $B_{\pm\infty}(g, x, r)$ is contained in the corresponding E^{wu} direction plaques. Since $\mu(\Lambda_{\ell}(g,\alpha)) > 0$, the Birkhoff's ergodic theorem suggests that for μ -almost every x, the forward orbit of x enters to $\Lambda_{\ell}(q,\alpha)$ for infinitely many times. Observe that the local unstable manifold of point in $\Lambda_{\ell}(g, \alpha)$ lies in the E^{wu} -direction plaque, by using the backward contracting property of local unstable manifold, one obtains that $B_{\pm\infty}(g, x, r)$ can has only one point x.

 \Box

Proof of Theorem [4.4.](#page-13-3) Assume that $f_n \to f$ in C^1 -topology and $\mu_n \to \mu$ in weak^{*}-topology, where $\{f_n\}$ and f are diffeomorphisms in $PH_{EC}^1(M)$, and $\mu_n \in G^u(f_n)$ for every $n \in \mathbb{N}$. It follows from Lemma [2.7](#page-8-2) that $\mu \in G^u(f)$. By Lemma [4.8,](#page-14-2) there exists $r > 0$ so that $h^*(f, \mu, r) = 0$ and $h^*(f_n, \mu_n, r) = 0$ for every *n* large enough. By Lemma [4.6,](#page-14-3) for any measurable partition P for which $\text{diam}(\mathcal{P}) \leq r$ and $\mu(\partial(\mathcal{P})) = 0$, we have $h_{\mu_n}(f_n) = h_{\mu_n}(f_n, \mathcal{P})$ and $h_{\mu}(f) = h_{\mu}(f, \mathcal{P})$. For this fixed \mathcal{P} , we can check by definition that

$$
\limsup_{n \to +\infty} h_{\mu_n}(f_n, \mathcal{P}) \le h_{\mu}(f, \mathcal{P}).
$$

All these together imply

$$
\limsup_{n \to +\infty} h_{\mu_n}(f_n) \le \limsup_{n \to +\infty} h_{\mu_n}(f_n, \mathcal{P}) \le h_{\mu}(f, \mathcal{P}) = h_{\mu}(f).
$$

This proves the result.

4.2. Proof of Theorem 4.2. The aim of this subsection is to complete the proof of Theorem 4.2. To begin with, let us recall the notion of *pseudophysical measure relative to compact disks*, introduced in [26,
$$
\S
$$
 2.3] as an extension of pseudo-physical measures [22, 23].

Definition 4.2. Given a compact C^1 disk D, we call $\mu \in \mathcal{M}(f)$ a pseudophysical measure relative to D if

$$
\mathrm{Leb}_D\left(\{x \in D : d(\omega_{\mathcal{M}}(x, f), \mu) < \varepsilon\}\right) > 0 \quad \text{for every } \varepsilon > 0.
$$

Lemma 4.9. [\[26,](#page-42-2) Theorem 2.3] Suppose that f is a homeomorphism on M and D is an embedded C^1 compact disk. Then the set of pseudo-physical measures relative to D is a compact non-empty set. Moreover, for Leb_D almost every $x \in D$, every invariant measure $\mu \in \omega_{\mathcal{M}}(x, f)$ is a pseudophysical measure relative to D.

Given a homeomorphism f on M, for every $\mu \in \mathcal{M}(f)$, for every $\varepsilon > 0$ and $n \in \mathbb{N}$, define the following subset of M:

$$
\mathcal{G}_{\mu}(n,\varepsilon) = \left\{ x \in M : \text{dist}\left(\frac{1}{n} \sum_{i=0}^{n-1} \delta_{f^i x}, \mu\right) < \varepsilon \right\}.
$$

Lemma 4.10. [\[26,](#page-42-2) Theorem C'] Let f be a C^1 partially hyperbolic diffeomorphism with invariant splitting $TM = E^u \oplus_{\succ} E^{cs}$. Then for any C^1 disk D transverse to E^{cs} , we have $\omega_{\mathcal{M}}(x,f) \subset G^u(f)$ for Leb_D-almost every x in D.

Lemma 4.11. [\[26,](#page-42-2) Theorem H] Let f be a C^1 diffeomorphism with dominated splitting $TM = E \oplus_{\succ} F$, then there exist $\theta > 0$, $R > 0$ such that for every $\mu \in \mathcal{M}(f)$, for any C^1 disk D tangent to \mathcal{C}_{θ}^E with diameter smaller than R, one has

$$
\lim_{\varepsilon \to 0} \limsup_{n \to +\infty} \frac{\log(\text{Leb}_D(\mathcal{G}_\mu(n,\varepsilon)))}{n} \le h_\mu(f) - \int \log|\text{det} Df|_E| d\mu.
$$

The following result is essentially contained in [\[26,](#page-42-2) § 5.2].

Lemma 4.12. Let f be a C^1 diffeomorphism with dominated splitting $TM =$ $E \oplus_{\succ} F$. Let \mathcal{M}_0 be a compact subset of $\mathcal{M}(f)$. Then for any C^1 disk D transverse to F, if $\omega_{\mathcal{M}}(x, f) \subset \mathcal{M}_0$ for Leb_D-almost every $x \in D$, then we have $\omega_{\mathcal{M}}(D, f) \subset \mathcal{M}_0$.

Now we are in position to give the proof of Theorem [4.2.](#page-13-1)

Proof of Theorem [4.2.](#page-13-1) Let θ and R be the positive constants given by Lemma [4.11,](#page-16-0) where we assume $E := E^{wu} = E^u \oplus E^{cu}$ and $F := E^{cs}$. Now we take a $C¹$ disk D transverse to E^{cs} , and prove the property [\(1\)](#page-13-0) firstly. By domination, up to considering iterates we can assume D is compact and tangent to \mathcal{C}_{θ}^{E} with diameter smaller than R. Since D can be foliated by disks transverse to $E^{cu} \oplus E^{cs}$, by Lemma [4.10](#page-16-1) we know that any measure in $\omega_{\mathcal{M}}(x, f)$ is a Gibbs u-state for Leb_D-almost every $x \in D$. Thus, according to Re-mark [4.1,](#page-13-4) it suffices to prove that for Leb_D-almost every point $x \in D$, any invariant measure $\mu \in \omega_{\mathcal{M}}(x, f)$ satisfies

(4.6)
$$
h_{\mu}(f) \geq \int \log |\det Df|_{E^{wu}}| d\mu.
$$

By Lemma [4.9,](#page-15-0) for Leb_D-almost every $x \in D$, any $\mu \in \omega_{\mathcal{M}}(x, f)$ is a pseudophysical measure relative to D, denote

$$
P_{\mu} = h_{\mu}(f) - \int \log |\det Df|_{E^{wu}}|d\mu.
$$

Now we show that μ satisfies [\(4.6\)](#page-16-2), that is $P_{\mu} \geq 0$. From Definition [4.2](#page-15-1) we know that for every $\varepsilon > 0$, there exists $\eta_{\varepsilon} > 0$ such that

$$
\text{Leb}_D\left(\{x \in D : \text{dist}(\omega_{\mathcal{M}}(x, f), \mu) < \varepsilon\}\right) > \eta_{\varepsilon},
$$

this together with the fact

$$
\{x \in D : \text{dist}(\omega_{\mathcal{M}}(x, f), \mu) < \varepsilon\} \subset \bigcap_{k=1}^{\infty} \bigcup_{n=k}^{\infty} \mathcal{G}_{\mu}(n, \varepsilon)
$$

yields the estimates

(4.7)
$$
\operatorname{Leb}_D\left(\bigcup_{n=k}^{\infty}\mathcal{G}_{\mu}(n,\varepsilon)\right) > \eta_{\varepsilon} \text{ for any } k \in \mathbb{N}.
$$

By contradiction, there is $\delta > 0$ such that $P_{\mu} < -2\delta$. Using Lemma [4.11,](#page-16-0) there are constants $\varepsilon > 0$ and $L > 0$ such that for every $n \in \mathbb{N}$,

$$
\text{Leb}_D(\mathcal{G}_\mu(n,\varepsilon)) < L \cdot \exp(n(P_\mu + \delta)) \leq L \cdot \exp(-\delta n).
$$

Consequently, for every $k \in \mathbb{N}$,

Leb_D
$$
\left(\bigcup_{n=k}^{\infty} \mathcal{G}_{\mu}(n,\varepsilon)\right) \leq L \sum_{n=k}^{\infty} \exp(-\delta n),
$$

which converges to zero as $k \to +\infty$. This a contradiction to [\(4.7\)](#page-17-0) and thus we obtain that μ satisfies [\(4.6\)](#page-16-2). We get property [\(1\)](#page-13-0) now.

Property [\(2\)](#page-13-5) is a consequence of [\(1\)](#page-13-0) and Theorem [4.5.](#page-14-4) Indeed, we have that $G^{cu}(f)$ is compact by Theorem [4.5,](#page-14-4) and we know $\omega_{\mathcal{M}}(x, f) \subset G^{cu}(f)$ for Leb_D-almost every $x \in D$ from [\(1\)](#page-13-0). Therefore, by applying Lemma [4.12](#page-16-3) to $E := E^{wu}, F := E^{cs}$ and $\mathcal{M}_0 := G^{cu}(f)$, one gets $\omega_{\mathcal{M}}(D, f) \subset G^{cu}(f)$. Observe that M can be covered by finitely many small balls, which can be foliated by disks transverse to E^{cs} , thus one can use the Fubini's theorem to show that \mathcal{B}_f has full Lebesgue measure.

The proof of Theorem [4.2](#page-13-1) is complete now.

$$
\Box
$$

4.3. Typical measurable partitions of Gibbs cu -states. Let us take $\delta_M > 0$ sufficiently small such that for any $x \in M$ the inverse of exponential map \exp_x^{-1} is well defined on the δ_M open neighborhood of x. We then identify this neighborhood with the corresponding neighborhood on tangent space.

Fix $f \in PH_{EC}^1(M)$ with partially hyperbolic splitting $TM = E^u \oplus_{\succ}$ $E^{cu} \oplus_{\succ} E^{cs}$, let $\mathcal{C}_a^{wu} := \mathcal{C}_a^{E^{wu}}$ for $a > 0$. Denote by \mathbb{D}^u , \mathbb{D}^s the compact unit disks in the Euclidean space of dimension $\dim E^{wu}$ and $\dim E^{cs}$, respectively.

Definition 4.3. A subset $C ⊂ M$ is called a cylinder if there exists a diffeomorphism $\psi : \mathbb{D}^u \times \mathbb{D}^s \to C$. We say that a smooth embedded sub-manifold γ of dimension dimE^{wu} (resp. dimE^{cs}) crosses C when $\psi^{-1}(\gamma \cap C)$ is a C¹ graph over \mathbb{D}^u (resp. \mathbb{D}^s).

By domination, we can choose $a > 0$ sufficiently small, and $0 < \rho_0 < \delta_M$ with following properties: if consider D as a compact smooth disk centered at $x \in M$ with radius $\rho \leq \rho_0$, which is tangent to \mathcal{C}_a^{wu} also, then the subset

$$
C(x, D, \rho) = \bigcup_{y \in D} \{ \exp_y(v) : v \perp T_y D, ||v|| \le \rho \}
$$

is a diffeomorphic image of $\mathbb{D}^u \times \mathbb{D}^s$, and thus a cylinder (of size ρ).

Given $\alpha > 0$ and $\ell \in \mathbb{N}$. By Lemma [3.2,](#page-9-1) one can take $\delta_{\ell,\alpha}$ such that all the points of $\Lambda_{\ell}(f,\alpha)$ admit local stable and unstable manifolds of size larger than $\delta_{\ell,\alpha}$. Thus, for any $\rho < \delta_{\ell,\alpha}$ sufficiently small, by domination and the choice of ρ_0 one can choose $r_{\ell,\alpha,\rho} \in (\rho, \min{\{\delta_{\ell,\alpha},\rho_0\}})$ such that for each $x \in M$ and disk D tangent to \mathcal{C}_a^{wu} centered at x of radius $r_{\ell,\alpha,\rho}$, we have that both $W^u_{\delta_{\ell,\alpha}}(y, f)$ and $W^s_{\delta_{\ell,\alpha}}(y, f)$ cross $C(x, D, r_{\ell,\alpha,\rho})$ whenever $y \in B(x, \rho) \cap \Lambda_{\ell}(f, \alpha)$. It follows that one can construct the measurable partition formed by local unstable disks with uniform size as follows:

$$
\mathcal{P}(x, D, \alpha, \ell, \rho) = \left\{ W^u_{\delta_{\ell, \alpha}}(y, f) \cap C(x, D, r_{\ell, \alpha, \rho}) : y \in B(x, \rho) \cap \Lambda_{\ell}(f, \alpha) \right\}.
$$

For $\mu \in \mathcal{M}(f)$, we say that $\mathcal{P}(x, D, \alpha, \ell, \rho)$ is a typical measurable partition associated to μ if $\mu(B(x,\rho) \cap \Lambda_{\ell}(f,\alpha)) > 0$.

By Proposition [3.5,](#page-11-0) we know that for every Gibbs u -state, there exists some typical measurable partition associated to it. Moreover, for Gibbs cu-states one has the following simple result.

Lemma 4.13. For every $\mu \in G^{cu}(f)$ and a typical measurable partition $\mathcal{P}(x, D, \alpha, \ell, \rho)$ associated to μ , there exists $D \in \mathcal{P}(x, D, \alpha, \ell, \rho)$ such that

- $D \subset \text{supp}(\mu)$:
- Leb_D $(\Lambda_{\ell}(f, \alpha)) > 0$.

Moreover, if μ is ergodic for f, then we can require D to satisfy that Lebesgue almost every point of D is contained in $B(\mu, f)$.

Proof. Since μ is a Gibbs cu-state, the conditional measures of μ along unstable disks of $\mathcal{P}(x, D, \alpha, \ell, \rho)$ are equivalent to the Lebesgue measures almost everywhere(see e.g. [\[9,](#page-41-11) Theorem 13.1.2]). Consequently, there exists $D \in \mathcal{P}(x, D, \alpha, \ell, \rho)$ such that $\text{Leb}_{D}(\Lambda_{\ell}(g, \alpha)) > 0$ and Lebesgue almost every point of D is contained in supp (μ) .

Now we show $D \subset \text{supp}(\mu)$. From the choice of D, we can choose $y \in D$ such that $y \in \text{supp}(\mu)$, and assume by contradiction that there is $z \in D$ which is not in supp (μ) . It follows that there exists an open neighborhood V of z such that $V \subset M \setminus \text{supp}(\mu)$, thus all the points in $V \cap D$ are not in supp (μ) , which is a contradiction to the fact that Lebesgue almost every point of D is contained in supp (μ) .

If $\mu \in G^{cu}(f)$ is ergodic, by Birkhoff's ergodic theorem one gets that $B(\mu, f)$ has full μ -measure. Thus, one can choose D to satisfy that Leb_Dalmost every point is contained in $B(\mu, f)$.

5. Certain properties of skeletons and pre-skeletons

Notation. Given a C^1 diffeomorphism f, let p, q be hyperbolic periodic points of f with the same index, denote

- $p \stackrel{f}{\prec} q$ if $W^u(\text{Orb}(p, f)) \pitchfork W^s(\text{Orb}(q, f)) \neq \emptyset$.
- $p \stackrel{f}{\sim} q$ if $p \stackrel{f}{\prec} q$ and $q \stackrel{f}{\prec} p$, or equivalently they are homoclinic related w.r.t. f .

Sometimes we drop the superscript " f " if there is no ambiguity.

In what follows, let f be a C^1 diffeomorphism in $PH_{EC}^1(M)$ with partially hyperbolic splitting $TM = E^u \oplus_{\succ} E^{cu} \oplus_{\succ} E^{cs}$ ^{[7](#page-19-0)}. By definition of preskeletons, one gets the following result directly.

Lemma 5.1. Given $n \in \mathbb{N}$, if $S = \{p_1, \dots, p_k\}$ is a pre-skeleton of f^n , then it is a pre-skeleton of f.

The proof of the following result is very similar to [\[28,](#page-42-9) Lemma 2.5], hence we omit the proof.

Lemma 5.2. Every pre-skeleton of f contains a subset to be a skeleton of f.

Lemma 5.3. If both $\{p_i : 1 \le i \le k\}$ and $\{q_i : 1 \le i \le \ell\}$ are skeletons of f, then there exists a bijective map $i \mapsto j(i)$ such that $p_i \sim q_{j(i)}$. Consequently, we must have $k = \ell$.

Proof. Consider two skeletons $S_1 = \{p_i : 1 \le i \le k\}$ and $S_2 = \{q_i : 1 \le i \le k\}$ $i \leq \ell$ of f. For each $p_i \in S_1$, we know that $W^u_{loc}(p_i, f)$ is a C^1 disk transverse to E^{cs} . It follows from the condition $(S1)$ $(S1)$ in Definition [1.1,](#page-2-3) there is $q_j \in S_2$ so that $W^u_{loc}(p_i, f)$ intersects $W^s(\text{Orb}(q_j, f))$ transversally. For this q_j , with the same reason we can find $p_l \in S_1$ such that $W^u_{loc}(q_j, f)$ intersects $W^s(\text{Orb}(p_l, f))$ transversally. As a consequence, inclination lemma [\[14,](#page-41-12) Theorem 5.7.2] leads to

$$
W^u_{loc}(p_i, f) \pitchfork W^s(\text{Orb}(p_i, f)) \neq \emptyset.
$$

Thus, condition ([S2\)](#page-2-4) in Definition [1.1](#page-2-3) gives $p_i = p_l$. Hence, we have $p_i \sim q_j$, and the choice of q_i is unique. Similarly, each element of S_2 has a unique element of S_1 such that they are homoclinically related. Consequently, we have $k = l$.

Lemma 5.4. If $\{p_i : 1 \leq i \leq k\}$ is a skeleton of f, then $H(p_i, f) =$ $W^u(\text{Orb}(p_i, f))$ for every p_i .

Proof. Since we have $H(p_i, f) \subset W^u(\text{Orb}(p_i, f))$ by definition, it suffices to show the other direction. For every point $x \in W^u(\text{Orb}(p_i, f))$, for any $\varepsilon > 0$ sufficiently small, let us take the disk $D_{\varepsilon} \subset W^u(\text{Orb}(p_i, f))$ centered at x with radius ε . By condition ([S1\)](#page-2-2) in Definition [1.1,](#page-2-3) there is $p_j \in \mathcal{S}$ such that

$$
D_{\varepsilon} \pitchfork W^s(\text{Orb}(p_j, f)) \neq \emptyset.
$$

Observe that there is no heteroclinic intersection between $W^u(\text{Orb}(p_i, f))$ and $W^s(\text{Orb}(p_j, f))$ as long as $i \neq j$. Thus, we must have $p_j = p_i$, and the arbitrariness of ε implies $x \in \overline{W^u(\text{Orb}(p_i,x))}$ \wedge $W^s(\text{Orb}(p_i,f))$. Hence, we have $\overline{W^u(\text{Orb}(p_i, f))} \subset H(p_i, f)$ to end the proof.

 7 Notice that in the proofs of lemmas in this section, we use neither the mostly expanding behavior nor the mostly contracting behavior on centers.

6. REDUCTION OF $PH_{EC}^{1}(M)$

Let $PH^1_C(M)$ be the subset of $PH^1_{EC}(M)$ for which if $f \in PH^1_C(M)$, then there exists a partially hyperbolic splitting $TM = E^u \oplus _ E^{cu} \oplus _ E^{cs}$ such that for every $\mu \in G^u(f)$, we have

$$
\chi_{\mu}(f^{-1}, E^{cu}) < 0 \quad \text{and} \quad \chi_{\mu}(f, E^{cs}) < 0.
$$

Lemma [2.6](#page-8-3) enables us to have the following result. The proof follows the same lines as the proof of [\[40,](#page-43-8) Proposition 3.3], hence omitted.

Theorem 6.1. If $f \in PH_{EC}^1(M)$, then there exists $N \in \mathbb{N}$ such that $f^N \in$ $PH^1_C(M).$

By Theorem [6.1,](#page-20-2) in order to prove Theorem [A,](#page-3-0) we can turn to prove the following parallel result for diffeomorphisms in $PH_C^1(M)$ firstly. Let $PH_C^{1+}(M)$ be the set of C^{1+} diffeomorphisms in $PH_C^1(M)$.

Theorem D. Every $f \in PH^1_C(M)$ has a skeleton. Moreover, if $f \in$ $PH_C^{1+}(M)$, then there exist finitely many physical measures of f with basin covering property, denoted by $\mathcal{P}(f) = {\mu_1, \cdots, \mu_k}.$ If $\mathcal{S}(f) = {p_1, \cdots, p_\ell}$ is a skeleton of f, then there exists a bijective map $i \mapsto j(i)$ such that

$$
supp(\mu_i) = \overline{W^u(\text{Orb}(p_{j(i)}, f))} = H(p_{j(i)}, f).
$$

for every $\mu_i \in \mathcal{P}(f)$ and the corresponding $p_{i(i)} \in \mathcal{S}(f)$. In particular, the number of physical measures of f coincides with the cardinality of its skeleton.

Theorem [D](#page-20-1) will be deduced in Subsection [6.4.](#page-29-0) Now we fix some notations for simplicity. Let $f \in PH^1_C(M)$ with partially hyperbolic splitting $TM =$ $E^u \oplus_{\succ} E^{cu} \oplus_{\succ} E^{cs}$. Given $\sigma > 0$ and $x \in M$, let $HT(x, \sigma, f)$ be the family of (E^{wu}, σ) -hyperbolic times of x w.r.t. f, where $E^{wu} = E^u \oplus E^{cu}$. For $a > 0$, rewriting $\mathcal{C}_a^{wu} := \mathcal{C}_a^{E^{wu}}$ and $\mathcal{C}_a^{cs} := \mathcal{C}_a^{E^{cs}}$.

6.1. The existence of skeletons. This subsection is dedicated to show the existence of skeletons for diffeomorphisms in $PH^1_C(M)$.

Theorem 6.2. Every $f \in PH^1_C(M)$ admits a skeleton $\mathcal{S}(f) = \{p_1, \dots, p_{k_f}\}\$ with some $k_f \in \mathbb{N}$ depending only on f.

We need the following simple observation:

Lemma 6.3. Given $\alpha > 0$, and assume that E is a Df-invariant sub-bundle of C^1 diffeomorphism f. If $x \in M$ is the point such that $\chi_{\mu}(f^{-1}, E) < -\alpha$ for every $\mu \in \omega_{\mathcal{M}}(x, f)$, then x is α -nonuniformly expanding along E.

Proposition 6.4. For every $f \in PH^1_C(M)$ with partially hyperbolic splitting $TM = E^u \oplus_{\succ} E^{cu} \oplus_{\succ} E^{cs}$, there exist $H_f > 0$, $D_f \in (0,1)$ and a C^1 neighborhood U_f of f such that for every $g \in U_f$, every point $x \in \mathcal{B}_g$ is H_f -nonuniformly expanding along E^{wu} with the density estimate

(6.8)
$$
\mathcal{D}_L\left(HT(x, e^{-H_f/2}, g)\right) \geq D_f.
$$

Proof. Since E^u is uniformly expanding and $G^u(f)$ is compact, one conclude that there exists $H_f > 0$ such that

$$
\chi_{\mu}(f^{-1}, E^{wu}) < -H_f \quad \text{for every } \mu \in G^u(f).
$$

Applying the upper semi-continuity of Gibbs u-space w.r.t. diffeomorphisms (Lemma [2.7\)](#page-8-2), there is a C^1 neighborhood \mathcal{U}_f of f such that for any $g \in \mathcal{U}_f$,

$$
\chi_{\mu}(g^{-1}, E^{wu}) < -H_f \quad \text{for every } \mu \in G^u(g).
$$

Note that for every $g \in \mathcal{U}_f$, if $x \in \mathcal{B}_g$ then $\omega_{\mathcal{M}}(x,g) \subset G^{cu}(g) \subset G^{u}(g)$, recalling definitions of \mathcal{B}_q and $G^{cu}(g)$. Thus, we have

$$
\chi_{\mu}(g^{-1}, E^{wu}) < -H_f \quad \text{for every } \mu \in \omega_{\mathcal{M}}(x, g).
$$

By Lemma [6.3,](#page-20-3) x is H_f -nonuniformly expanding along E^{wu} . Then, using Proposition [2.3,](#page-7-1) reducing \mathcal{U}_f if necessary, there exists a density $D_f \in (0,1)$ such that for every $g \in \mathcal{U}_f$, we have

$$
\mathcal{D}_L\left(HT(x, e^{-H_f/2}, g)\right) \ge D_f
$$

for every $x \in \mathcal{B}_q$.

Theorem 6.5. Let $f \in PH^1_C(M)$ with partially hyperbolic splitting $TM =$ $E^u \oplus_{\succ} E^{cu} \oplus_{\succ} E^{cs}$. There exist $\alpha > 0$, $\ell \in \mathbb{N}$, $\theta_f \in (0,1)$, $\rho_f > 0$, $H_f > 0$ and a C^1 neighborhood U of f such that, for every $\rho \leq \rho_f$, for every $g \in \mathcal{U}$, for every $x \in \mathcal{B}_g$ and every $\mu \in \omega_{\mathcal{M}}(x,g)$,

(1) the forward orbit of x enters to $\mathscr{A}_{\mu}(g,\ell,\alpha,\rho)$ for infinitely many $(E^{wu}, e^{-H_f/2})$ -hyperbolic times with upper density no less than θ_f . More precisely,

$$
\mathcal{D}_U\left(\left\{j\in HT(x, e^{-H_f/2}, g) : g^j(x)\in \bigcup_{B\in\mathscr{A}_{\mu}(g,\ell,\alpha,\rho)}B\right\}\right)\geq \theta_f;
$$

(2) there exists $B \in \mathscr{A}_{\mu}(g,\ell,\alpha,\rho)$ such that if $\gamma \ni x$ is a C^1 disk transverse to E^{cs} , then $\gamma \pitchfork W^s(\text{Orb}(p_B, g)) \neq \emptyset$. Here p_B is the hyperbolic periodic point in B given by Theorem [3.6.](#page-11-1)

Proof. Take $D_f \in (0,1)$, $H_f > 0$ and C^1 neighborhood \mathcal{U}_f of $f \in PH_C^1(M)$ as in Proposition [6.4.](#page-20-4) Choose $\varepsilon < D_f/2$, by Theorem [3.6](#page-11-1) there exist $\alpha > 0$, $\ell \in \mathbb{N}, \delta_{\ell}, \rho_{\ell} > 0$ and a C^1 neighborhood $\mathcal{U} \subset \mathcal{U}_f$ of f such that for every $\rho \leq \rho_{\ell}$, for every $g \in \mathcal{U}$ and every $\mu \in G^u(g)$, one can construct $\mathscr{A}_{\mu}(g,\ell,\alpha,\rho)$, which is consisted of ρ -balls satisfying

(6.9)
$$
\mu\left(\bigcup_{B\in\mathscr{A}_{\mu}(g,\ell,\alpha,\rho)}B\right) > 1-\varepsilon;
$$

$$
\Box
$$

• each $B \in \mathscr{A}_{\mu}(g,\ell,\alpha,\rho)$ contains a hyperbolic periodic point p_B with local stable and unstable manifolds of size larger than δ_{ℓ} .

By domination together with Lemma [2.2,](#page-7-2) one can choose $0 < \rho_f < \rho_{\ell}$, $0 < \delta_f < \delta_\ell$, and $a > 0$ such that

- (A) for any points x, y with $d(x, y) \le 2\rho_f$, if γ_x and γ_y are disks centered at x and y with radius δ_f , which are tangent to \mathcal{C}_a^{wu} and \mathcal{C}_a^{cs} , respectively, then γ_x intersects γ_y transversely;
- (B) for any C^1 embedded disk γ tangent to \mathcal{C}_a^{wu} and containing $x \in \mathcal{B}_g$, if $n \in HT(x, e^{-H_f/2}, g)$, then

$$
d_{g^i(\gamma)}(g^i(x), g^i(y)) \le e^{(-(n-i)H_f)/4} d_{g^n(\gamma)}(g^n(x), g^n(y))
$$

for any $y \in \gamma$ satisfying $d_{g^n(\gamma)}(g^n(x), g^n(y)) \leq \delta_f$.

Now let us fix $g \in \mathcal{U}$, $x \in \mathcal{B}_g$, $\mu \in \omega_{\mathcal{M}}(x,g)$, and $\rho \leq \rho_f$. For simplicity, take A_{ρ} as the union of the ρ -balls of $\mathscr{A}_{\mu}(g,\ell,\alpha,\rho)$. Due to the choice of μ , there exists a subsequence $\{n_i\}$ such that

$$
\frac{1}{n_i} \sum_{j=0}^{n_i-1} \delta_{g^j(x)} \xrightarrow{weak^*} \mu, \quad \text{as } i \to +\infty.
$$

This together with [\(6.9\)](#page-21-0) yields

$$
\liminf_{i \to +\infty} \frac{1}{n_i} \sum_{j=0}^{n_i-1} \chi_{A_\rho}(g^j(x)) \ge \mu(A_\rho) > 1 - \varepsilon,
$$

where we use the fact $\mu \in G^u(g)$. On the other hand, since x is H_f nonuniformly expanding along E^{wu} with density estimate [\(6.8\)](#page-20-5). According to the choices of ε and D_f , one has

$$
\mathcal{D}_{U}\left(\lbrace j \in HT(x, e^{-H_{f}/2}, g) : g^{j}(x) \in A_{\rho}\rbrace\right) \ge D_{f} - \varepsilon > 0.
$$

Therefore, we have complete the proof of Item [\(1\)](#page-32-0) by taking $\theta_f = D_f - \varepsilon$.

We begin to prove Item (2) . As a consequence of Item (1) , one can choose $B \in \mathscr{A}_{\mu}(g,\ell,\alpha,\rho)$ to satisfy

(6.10)
$$
\mathcal{D}_U\left(\lbrace j \in HT(x, e^{-H_f/2}, g) : g^j(x) \in B \rbrace\right) \geq \frac{\theta_f}{\#\mathscr{A}_{\mu}(g, \ell, \alpha, \rho)}.
$$

Let γ be a C^1 disk transverse to E^{cs} , containing the fixed point x in \mathcal{B}_g . By domination, we may assume that every $g^{n}(\gamma)$, $n \geq 1$ is tangent to \mathcal{C}_{a}^{wu} . It follows from [\(B\)](#page-22-0) and [\(6.10\)](#page-22-1) that there exists $n \in \mathbb{N}$ such that $g^{n}(x) \in B$ and $g^{n}(\gamma)$ contains the disk of radius δ_{f} centered at $g^{n}(x)$. Since $p, g^{n}(x) \in B$, we know $d(p, g^n(x)) < 2\rho \le 2\rho_f$, then [\(A\)](#page-22-2) gives $g^n(\gamma) \pitchfork W^s_{loc}(p_B, f) \ne \emptyset$. Consequently, we obtain $\gamma \pitchfork W^s(\text{Orb}(p_B, f)) \neq \emptyset$. Now, we finish the proof of Theorem [6.5.](#page-21-2) \Box

We are in position to give the proof of Theorem [6.2.](#page-20-6)

Proof of Theorem [6.2.](#page-20-6) Let $f \in PH^1_C(M)$ with partially hyperbolic splitting $TM = E^u \oplus_{\succ} E^{cu} \oplus_{\succ} E^{cs}$. From Lemma [5.3,](#page-19-1) it suffices to construct a skeleton of f , then it admits the cardinality depending only on f , which we denote by k_f .

By Theorem [6.5,](#page-21-2) there exist $\alpha > 0$, $\ell \in \mathbb{N}$, $\rho_f > 0$ such that, for every $x \in$ \mathcal{B}_f , when we fix $\mu \in \omega_{\mathcal{M}}(x, f)$, there is $B \in \mathscr{A}(f, \ell, \alpha, \rho_f)$ with hyperbolic periodic point $p_x := p_B$ in B so that for every C^1 disk D containing x and transverse to E^{cs} , we have that $D \pitchfork W^s(\text{Orb}(p_x, f)) \neq \emptyset$. Denote by $\mathcal{S}_0 = \{p_x : x \in \mathcal{B}_f\}$ the set consisting of all the periodic points obtained as above. By Theorem [4.2,](#page-13-1) for every C^1 disk D transverse to E^{cs} , the intersection of $\mathcal{B}_f \cap D$ admits full Lebesgue measure, thus there is $x \in \mathcal{B}_f$ for which D intersects $W^s(\text{Orb}(p_x, f))$ transversally. It follows from Theorem [3.6](#page-11-1) that every hyperbolic periodic point in S_0 admits stable and unstable manifolds of size larger than δ_{ℓ} . This implies that the number of periodic points of S_0 which are not homoclinically related to each other is finite only. Therefore, one can take $S = \{p_i : 1 \leq i \leq l\} \subset S_0$ such that different elements of S are not homoclinically related with maximal cardinality, that is, for any $p_x \in \mathcal{S}_0$ there exists $p_i \in \mathcal{S}$ so that $p_x \sim p_i$.

Next we show that S is a pre-skeleton, which implies that there exists a subset of S to be a skeleton, using Lemma [5.2.](#page-19-2) By contradiction, we suppose that there is a C^1 disk D transverse to E^{cs} which do not intersect $W^s(\text{Orb}(p, f))$ for any $p \in \mathcal{S}$. On the other hand, by the construction of $\{p_x : x \in \mathcal{B}_f\}$, one can take p_x for some $x \in \mathcal{B}_f \cap D$ such that

(6.11)
$$
D \pitchfork W^s(\text{Orb}(p_x, f)) \neq \emptyset.
$$

By the maximality of S, there exists $p \in S$ such that $p_x \sim p$, which together with [\(6.11\)](#page-23-0) implies that D intersects $W^s(Orb(p, f))$ transversally, using the inclination lemma. This is a contradiction to our assumption, thus we get the desired result and complete the proof of Theorem [6.2.](#page-20-6) \Box

The following theorem focus on the perturbations of the skeleton, which in particular gives the upper semi-continuity of $f \mapsto k_f$ for diffeomorphisms in $PH_C^1(M)$.

Theorem 6.6. Let $f \in PH^1_C(M)$ and $S(f) = \{p_i : 1 \leq i \leq k_f\}$ be a skeleton of f. Then there exists a C^1 neighborhood U of f such that for every $g \in \mathcal{U}$, the continuation $\{p_i(g) : 1 \leq i \leq k_f\}$ of $\mathcal{S}(f)$ is a pre-skeleton of g. Consequently, we have $k_q \leq k_f$ for every $g \in \mathcal{U}$.

Proof. For fixed $f \in PH^1_C(M)$ with partially hyperbolic splitting $TM =$ $E^u \oplus_{\succ} E^{cu} \oplus_{\succ} E^{cs}$, take the C^1 neighborhood $\mathcal U$ of f and size $\delta_{\ell} > 0$ as in Theorem [6.5.](#page-21-2) By Theorem [6.2,](#page-20-6) every diffeomorphism of U admits a skeleton whose periodic points exhibit local stable and unstable manifolds of size larger than δ_{ℓ} .

Now let $S(f) = \{p_i : 1 \leq i \leq k_f\}$ be any skeleton of f. For every $x \in M$, take a C^1 disk transverse to E^{cs} centered at x with radius $\delta_{\ell}/4$, and denote it by $D(x, \delta_{\ell}/4)$. By definition of skeletons, there is $p_i \in \mathcal{S}(f)$ such that $W^s(\text{Orb}(p_i, f))$ intersects $D(x, \delta_{\ell}/4)$ transversely at some point in the interior of $W^s_{\rho_x}(\text{Orb}(p_i, f))$ for some $\rho_x > 0$. Thus, one can take a small neighborhood U_x of x in M such that for every $y \in U_x$, if γ is a C^1 disk centered at y of radius $\delta_{\ell}/2$ and is transverse to E^{cs} , then it intersects $W_{\rho_x}^s(\text{Orb}(p_i, f))$ transversally. By compactness, we can take an open covering $\{U_{x_i}: 1 \leq i \leq m\}$ of M, and put $\rho = \max\{\rho_{x_i}: 1 \leq i \leq m\}$. By the continuity of the sub-bundles and the compact part of stable manifold of periodic point w.r.t. diffeomorphisms, one can reduce $\mathcal U$ small enough such that for every $g \in \mathcal{U}$, the family $\{p_i(g): 1 \leq i \leq k_f\}$ obtained as the continuation of $\{p_i : 1 \le i \le k_f\}$ possesses the following property: for every C^1 disk D transverse to E^{cs} with radius δ_{ℓ} around a point in M, there is $1 \leq i \leq k_f$ such that

$$
D \pitchfork W^s_{\rho}(\text{Orb}(p_i(g), g)) \neq \emptyset.
$$

For $g \in \mathcal{U}$, let $\mathcal{S}(g) = \{q_j : 1 \leq j \leq k_g\}$ be a skeleton of g such that every $q_j \in \mathcal{S}(g)$ has local unstable manifold of size larger than δ_{ℓ} . In particular, for every $q_j \in \mathcal{S}(g)$, there is $1 \leq i \leq k_f$ such that

$$
W^u_{loc}(q_j, g) \pitchfork W^s_{\rho}(\text{Orb}(p_i(g), g)) \neq \emptyset.
$$

Then, the inclination lemma yields $W^s(\text{Orb}(q_j, g)) \subset \overline{W^s(\text{Orb}(p_i(g), g))}$, so ${p_i(g) : 1 \le i \le k_f}$ satisfies condition ([S1\)](#page-2-2), and is a pre-skeleton. Therefore, it contains a subset to be a skeleton by Lemma [5.2,](#page-19-2) thus $k_q \leq k_f$.

 \Box

6.2. Finiteness of ergodic physical measures. For every $f \in PH_C^{1+}(M)$, use $\mathcal{P}_e(f)$ denotes the family of ergodic physical measures of f. Following Theorem [4.5,](#page-14-4) we know that $\mathcal{P}_e(f)$ is non-empty. Moreover, in terms of the definitions of physical measures and \mathcal{B}_f , we have the following observation:

Lemma 6.7. If $f \in PH_C^{1+}(M)$, then $\mu \in \mathcal{P}_e(f)$ if and only if it is an ergodic Gibbs cu-state.

Notation. Given a diffeomorphism f on M, assume that $\mu \in \mathcal{M}(f)$ and p a hyperbolic periodic point of f, writing $\mu \leftrightarrow$ p if

$$
\mathrm{supp}(\mu)=\overline{W^u(\mathrm{Orb}(p,f))}=H(p,f).
$$

We can build the following relationship between ergodic physical measures and the skeleton for any diffeomorphism in $PH_C^{1+}(M)$.

Proposition 6.8. Assume that f is a diffeomorphism in $PH_C^{1+}(M)$ with a skeleton $S = \{p_i : 1 \le i \le k_f\}$. Then there is an injective map $i \mapsto j(i)$ charactered by following property: for every $\mu_i \in \mathcal{P}_e(f)$, there is $p_{i(i)} \in \mathcal{S}$ and an unstable disk D_i such that

(1) D_i is contained in the support of μ_i , and Lebesgue almost every point of it belongs to $B(\mu_i, f)$;

(2) D_i intersects $W^s(\text{Orb}(p_{j(i)}, f))$ transversally.

Consequently, we have $\mu_i \leftrightarrow_{p_{j(i)}}$ for every $1 \leq i \leq k_f$, and $\#P_e(f) \leq k_f$.

Proof. By Lemma [5.3,](#page-19-1) it is enough to check this result for some skeleton of f. Consider the skeleton $S(f) = \{p_1, \dots, p_{k_f}\}\$ of $f \in PH_C^{1+}(M)$ that constructed as in the proof of Theorem [6.2,](#page-20-6) thus each p_i , $1 \leq i \leq k_f$ is a hyperbolic periodic point contained in $B(x_i, \rho_f) \in \mathscr{A}_{\mu}(f, \ell, \alpha, \rho_f)$ for some $x_i \in \text{supp}(\mu | \Lambda_{\ell}(f, \alpha)), \mu \in G^{cu}(f)$ and fixed $\alpha > 0, \rho_f > 0$ and $\ell \in \mathbb{N}$. Note that by the choice of ρ_f , we may assume $\rho_f \ll \min\{\rho_0, \delta_\ell\}$. Recalling the construction of cylinders and measurable partitions in Subsection [4.3,](#page-17-1) there is $r_{\ell} \in (\rho_f, \min\{\rho_0, \delta_{\ell}\})$ such that one can take disks γ_i tangent to \mathcal{C}_a^{wu} of radius r_{ℓ} centered at x_i to build the cylinders and measurable partitions as follows

$$
C_i = C(x_i, \gamma_i, r_\ell), \quad \mathcal{P}_i = \mathcal{P}(x_i, \gamma_i, \alpha, \ell, \rho_f), \quad i = 1, \cdots k_f.
$$

We observe that both $W^s_{\delta_{\ell}}(p_i, f)$ and $W^u_{\delta_{\ell}}(p_i, f)$ cross the cylinder C_i for every $1 \leq i \leq k_f$.

For every $\mu_i \in \mathcal{P}_e(f)$, since it is an ergodic Gibbs cu-state guaranteed by Lemma [6.7,](#page-24-0) using Lemma [4.13](#page-18-1) there is a C^1 unstable disk $D_i \subset \text{supp}(\mu_i)$ so that Leb_{D_i} -almost every point of D_i is contained in $B(\mu_i, f)$. Moreover, it follows from condition ([S1\)](#page-2-2) of Definition [1.1](#page-2-3) that there exists $p_{i(i)} \in \mathcal{S}(f)$ such that

(6.12)
$$
D_i \pitchfork W^s(\text{Orb}(p_{j(i)}, f)) \neq \emptyset.
$$

Hence, we have found D_i and $p_{i(i)}$ w.r.t. each ergodic physical measure μ_i to satisfy properties [\(1\)](#page-24-1) and [\(2\)](#page-25-0).

Now we show $\mu_i \leftrightarrow^f p_{j(i)}$ for every $1 \leq i \leq k_f$. Since $D_i \subset \text{supp}(\mu_i)$, and $\text{supp}(\mu_i)$ is compact and invariant, by applying the inclination lemma, one gets

(6.13)
$$
W^u(\text{Orb}(p_{j(i)}, f)) \subset \text{supp}(\mu_i)
$$

immediately. Observe that by definition each \mathcal{P}_i , $1 \leq i \leq k_f$ is a typical cylinder associated to some Gibbs cu-state, which together with Lemma [4.13](#page-18-1) implies that one can choose $D_{p_{j(i)}} \in \mathcal{P}_{j(i)}$ such that

(6.14)
$$
Leb_{D_{p_{j(i)}}}(\Lambda_{\ell}(f,\alpha)) > 0.
$$

Note that all the points in $\Lambda_{\ell}(f,\alpha)$ admit local stable manifolds of size larger than δ_{ℓ} . Using inclination lemma to [\(6.12\)](#page-25-1), we can take $n \in \mathbb{N}$ large enough such that there is a disk $\widetilde{D}_i \subset f^n(D_i)$ that is C^1 -close enough to $W^u_{\delta_\ell}(p_{j(i)},f)$ and crosses the cylinder $C_{j(i)}$. As Lebesgue almost every point of D_i is contained in $B(\mu_i, f)$, Lebesgue almost every point of D_i is contained in $B(\mu_i, f)$ as well. Observe that any stable manifold can only lie in the same

basin of some invariant measure, using [\(6.14\)](#page-25-2) and the absolute continuity of stable lamination (see e.g. [\[9,](#page-41-11) §8.6]), we get

$$
\mathrm{Leb}_{W^u_{\delta_\ell}(p_{j(i)},f)}(B(\mu_i,f))>0.
$$

This implies that one can pick a point x in $W^u_{\delta_{\ell}}(p_{j(i)}, f) \cap B(\mu_i, f)$, thus

$$
\frac{1}{n}\sum_{k=0}^{n-1}\delta_{f^k(x)} \xrightarrow{weak^*} \mu_i, \text{ as } n \to +\infty.
$$

Hence, for any $y \in \text{supp}(\mu_i)$ and any open neighborhood U of y, there exist infinitely many times k such that $f^k(x) \in U$, and thus $U \cap W^u(\text{Orb}(p_{j(i)}, g)) \neq$ \emptyset . Therefore, supp $(\mu_i) \subset \overline{W^u(\text{Orb}(p_{j(i)}, f))}$. This combines with [\(6.13\)](#page-25-3) and Lemma [5.4](#page-19-3) yields $\mu_i \leftrightarrow p_{j(i)}$.

Now we show that the map $i \mapsto j(i)$ is injective. Otherwise, there are two different ergodic physical measures ν_1 and ν_2 , $p_j \in S(f)$ for some $1 \leq j \leq k_f$ and C^1 unstable disks $\{D_i\}_{i=1,2}$ satisfying properties $(1)-(2)$ $(1)-(2)$ as follows:

- $-D_i \subset \text{supp}(\nu_i)$, and Leb_{D_i} -almost every point belongs to $B(\nu_i, f)$;
- $-D_i \pitchfork W^s(\text{Orb}(p_j, f)) \neq \emptyset.$

The second property guarantees that, there exist $n_1, n_2 \in \mathbb{N}$ and the disks $D_{n_i} \subset f^{n_i}(D_i)$, $i = 1, 2$ that are C^1 -close enough to $W^u_{\delta_{\ell}}(p_j, f)$ and cross the cylinder C_i . Then the first property ensures that Lebesgue almost every point of D_{n_i} is contained in $B(\nu_i, f)$, $i = 1, 2$. Note that $p_j \in$ $B(x_j, \rho_f)$, for the measurable partition \mathcal{P}_j , one can take $D_{p_j} \in \mathcal{P}_j$ such that $\text{Leb}_{D_{p_j}}(\Lambda_\ell(f,\alpha)) > 0$ by Lemma [4.13.](#page-18-1) Then using the absolute continuity of stable lamination again, similar to above argument we conclude that $B(\nu_1, f) \cap B(\nu_2, f) \neq \emptyset$, which is a contradiction to our assumption. Thus, we know that the map $i \mapsto j(i)$ is injective. As a direct result, we get $\#P_e(f) \leq k_f.$

6.3. Ergodicity and basin covering property of physical measures. Recall that in the previous work [\[40\]](#page-43-8), for diffeomorphisms in $PH_{EC}^{1+}(M)$, the authors have proved the finiteness and basin covering property of physical measures. In this subsection, by making more accurate analysis on Gibbs cu-states, we provide a different proof of these results. In consideration of Theorem [6.1,](#page-20-2) it suffices to prove them for diffeomorphisms in $PH_C^{1+}(M)$.

Theorem 6.9. For every $f \in PH_C^{1+}(M)$, we have

- (1) all physical measures are ergodic and coincide with the extreme elements of $G^{cu}(f)$;
- (2) Lebesque almost every point of M is contained in some basin of physical measures.

Fixed constants. Given $f \in PH_C^{1+}(M)$ with partially hyperbolic splitting $TM = E^u \oplus_{\succ} E^{cu} \oplus_{\succ} E^{cs}$. Let us fix H_f, D_f as in Proposition [6.4,](#page-20-4) and choose $\varepsilon < D_f/2$. For this $\varepsilon > 0$, one chooses $\alpha > 0$, $\ell \in \mathbb{N}$, $\rho_{\ell} > 0$, $\delta_{\ell} > 0$ as in Theorem [3.6.](#page-11-1) Furthermore, by Lemma [2.2,](#page-7-2) up to shrinking δ_{ℓ} , there exists $a > 0$ such that we have the following backward contracting property: if n is a $(E^{wu}, e^{-H_f/2})$ -hyperbolic time of x, and γ is a C^1 disk tangent to \mathcal{C}_a^{wu} , then

(6.15)
$$
d_{f^{n-i}\gamma}(f^{n-i}(y), f^{n-i}(z)) \le e^{-iH_f/4} d_{f^n\gamma}(f^n(y), f^n(z))
$$

whenever $y, z \in \gamma$ within $f^{n}(x), f^{n}(y)$ contained in $B_{f^{n}(\gamma)}(f^{n}(x), \delta_{\ell})$. Let us mention that from the choices of ρ_{ℓ} and δ_{ℓ} , we can and will assume $\rho_{\ell} \ll \delta_{\ell}$.

Following [\[2,](#page-41-10) Corollary 4.2], for any C^2 disk D transverse to E^{cs} , the iterate $f^{n}(D)$ admits the *bounded Hölder curvature* for all *n* sufficiently large, see [\[2,](#page-41-10) § 2.1] for more details. Therefore, without loss of generality, we assume that any C^2 disk transverse to E^{cs} considered in the rest of this subsection exhibits the bounded Hölder curvature.

By Theorem [4.5,](#page-14-4) all the extreme elements of $G^{cu}(f)$ are ergodic, they are also ergodic physical measures. Hence, applying Proposition [6.8](#page-24-2) we know that there are at most k_f extreme elements in $G^{cu}(f)$. Denote them by μ_1, \cdots, μ_k for some $k \leq k_f$. Rewriting

$$
\mathscr{A}_i := \mathscr{A}_{\mu_i}(f, \ell, \alpha, \rho_\ell), \quad i = 1, \cdots, k,
$$

and

$$
\mathbb{B}:=\bigcup_{1\leq i\leq k}\bigcup_{B\in\mathscr{A}_i}B.
$$

For $x \in \mathcal{B}_f$, let us denote

(6.16)
$$
HT_{\mathbb{B}}(x, e^{-H_f/2}, f) = \left\{ n : n \in HT(x, e^{-H_f/2}, f) \text{ and } f^n(x) \in \mathbb{B} \right\}.
$$

Lemma 6.10. For every $x \in \mathcal{B}_f$, we have $\mathcal{D}_U(HT_{\mathbb{B}}(x, e^{-H_f/2}, f)) \ge D_f/2$.

Proof. Take $\mu \in \omega_{\mathcal{M}}(x, f)$, which belongs to $G^{cu}(f)$ by definition. Therefore, it can be written as the convex combination of the extreme elements of $G^{cu}(f)$, thus there exists a sequence of real numbers $\xi_i \in [0,1]$ for $1 \leq i \leq k$ satisfying $\xi_1 + \cdots + \xi_k = 1$ so that $\mu = \xi_1 \mu_1 + \cdots + \xi_k \mu_k$. Due to the construction of \mathscr{A}_i in Theorem [3.6](#page-11-1) we know $\mu_i(\cup_{B \in \mathscr{A}_i} B) > 1 - \varepsilon$ for every $1 \leq i \leq k$. Consequently,

$$
\mu(\mathbb{B}) > 1 - \varepsilon.
$$

As $\mu \in \omega_{\mathcal{M}}(x, f)$, we can choose subsequence $\{n_l\}$ tends to infinity such that

$$
\frac{1}{n_l} \sum_{i=0}^{n_l-1} \delta_{f^i(x)} \xrightarrow{weak^*} \mu, \quad \text{as } l \to +\infty.
$$

Combined with [\(6.17\)](#page-27-0) it follows that the times n for which $f^{n}(x)$ enters to B admits upper density estimate as follows:

$$
\mathcal{D}_U\left(\{n \in \mathbb{N} : f^n(x) \in \mathbb{B}\}\right) \ge 1 - \varepsilon.
$$

This together with the choices of ε and D_f yields

$$
\mathcal{D}_U\left(HT_{\mathbb{B}}(x, e^{-H_f/2}, f)\right) \ge D_f/2.
$$

By Lemma [6.10,](#page-27-1) we can deduce the following result which is a variation of [\[4,](#page-41-13) Lemma 5.4], thus we omit the proof.

Lemma 6.11. Assume that D is a C^2 disk transverse to E^{cs} . Let A be a subset of \mathcal{B}_f and U an open subset in D satisfying $\text{Leb}_D(A \cap U) > 0$. For any $\eta > 0$, there exists $n \in HT_{\mathbb{B}}(x, e^{-H_f/2}, f)$ for some $x \in A$, such that $f^{n}(D)$ contains a ball B_n of radius $\delta_{\ell}/4$ centered at $f^{n}(x)$, and we have

$$
\frac{\operatorname{Leb}_{f^n(D)}(f^n(A) \cap B_n)}{\operatorname{Leb}_{f^n(D)}(B_n)} > 1 - \eta.
$$

Now we can give the proof of Theorem [6.9.](#page-26-0)

Proof of Theorem [6.9.](#page-26-0) Let μ_1, \dots, μ_k be the extreme elements of $G^{cu}(f)$, and take constants $\alpha > 0$, $\ell \in \mathbb{N}$, $\rho_{\ell} > 0$, $a > 0$ as fixed after the statement of Theorem [6.9.](#page-26-0) Recall that \mathcal{A}_i is the collection of finitely many ρ_{ℓ} -balls that covers supp $(\mu_i | \Lambda_{\ell}(f, \alpha))$, and the union of ρ_{ℓ} -balls from \mathscr{A}_i , $1 \leq i \leq k$ is denoted by $\mathbb B$. Note also $HT_{\mathbb B}(x, e^{-H_f/2}, f)$ is given by [\(6.16\)](#page-27-2). For this ρ_{ℓ} and δ_{ℓ} , by domination there exists $r_{\ell} \in (\rho_{\ell}, \delta_{\ell})$ such that for each $B :=$ $B(x,\rho_\ell) \in \mathscr{A}_i, 1 \leq i \leq k,$

(I) one can take a C^1 disk D tangent to \mathcal{C}_a^{wu} centered at x of radius r_ℓ , with which we build cylinder and measurable partition as follows:

$$
C_B := C(x, D, r_\ell), \quad \mathcal{P}_B = \mathcal{P}(x, D, \alpha, \ell, \rho_\ell);
$$

Recalling their constructions in Subsection [4.3.](#page-17-1)

(II) If γ is a C^1 sub-manifold tangent to \mathcal{C}_a^{wu} which contains a disk of radius $\delta_{\ell}/4$ centered at a point of B, then this disk must cross C_B .

For each $B \in \mathscr{A}_i$, since we have $\mu_i(B \cap \Lambda_\ell(f, \alpha)) > 0$, \mathcal{P}_B is a typical measurable partition associated to μ_i , so there exists $\gamma_B \in \mathcal{P}_B$ such that

$$
\mathrm{Leb}_{\gamma_B}(B(\mu_i, f) \cap \Lambda_\ell(f, \alpha)) > 0.
$$

Note that all points of $\Lambda_{\ell}(f,\alpha)$ admit local stable manifolds of size larger than δ_{ℓ} , and the points of a stable manifold can only lie in the same basin. Then, in view of [\(II\)](#page-28-0), by using the absolute continuity of stable lamination, we can take $L_0 > 0$ so that

(6.18)
$$
\frac{\text{Leb}_{\gamma}(B_{\gamma}(x,\delta_{\ell}/4) \cap B(\mu_i,f))}{\text{Leb}_{\gamma}(B_{\gamma}(x,\delta_{\ell}/4))} \ge L_0
$$

whenever $x \in B \in \mathcal{A}_i$ and γ is a C^2 sub-manifold containing the ball of radius $\delta_{\ell}/4$ centered at x. Here we use the finiteness of balls in \mathscr{A}_i , $1 \leq i \leq k$.

We are ready to give the proof of Item [\(1\)](#page-31-1) firstly. Let μ be a physical measure of f, then $B(\mu, f)$ exhibits positive Lebesgue measure, and thus $\mathcal{B}_f \cap B(\mu, f)$ has positive Lebesgue measure. Therefore, we know $\mu \in G^{cu}(f)$, which also implies $B(\mu, f) \subset \mathcal{B}_f$ by definition. Next we want to prove $\mu = \mu_i$ for some $1 \leq i \leq k$.

Pick a density point of $B(\mu, f)$ and foliate the small neighborhood of it with smooth sub-manifolds transverse to E^{cs} ; then using the Fubini's theorem, one can find a C^2 disk D lie in some sub-manifold such that Leb_D($O \cap B(\mu, f) > 0$ for some open set O in D. By taking $\eta < L_0$ and $A := B(\mu, f)$, Lemma [6.11](#page-28-1) implies that there exist $x \in B(\mu, f)$ and $n \in HT_{\mathbb{B}}(x, e^{-H_f/2}, f)$ sufficiently large such that

$$
\frac{\text{Leb}_{f^{n}(D)} (B_{f^{n}(D)}(f^{n}(x), \delta_{\ell}/4) \cap B(\mu, f))}{\text{Leb}_{f^{n}(D)} (B_{f^{n}(D)}(f^{n}(x), \delta_{\ell}/4))} > 1 - \eta,
$$

where we use the fact that $B(\mu, f)$ is f-invariant. Note also that $f^{n}(x)$ is contained in some ball of \mathscr{A}_i for some $1 \leq i \leq k$, we have the estimate [\(6.18\)](#page-28-2) by taking $x := fⁿ(x)$ and $\gamma := fⁿ(D)$. By the choices of η and L_0 , one must have

$$
\operatorname{Leb}_{f^n(D)}\left(B_{f^n(D)}(f^n(x), \delta_{\ell}/4) \cap B(\mu_i, f) \cap B(\mu, f)\right) > 0,
$$

which implies $B(\mu, f) \cap B(\mu_i, f) \neq \emptyset$, and thus $\mu = \mu_i$. This shows that μ is ergodic and is an extreme element of $G^{cu}(f)$, which is an ergodic physical measure following from Theorem [4.5.](#page-14-4) This completes the proof of Item [\(1\)](#page-26-1).

From Item [\(1\)](#page-26-1), we know that μ_1, \cdots, μ_k are all the physical measures of f. Because \mathcal{B}_f has full Lebesgue measure by Theorem [4.2,](#page-13-1) to see Item [\(2\)](#page-31-2), it suffices to prove that Lebesgue almost every point of \mathcal{B}_f is contained in $B(\mu_i, f)$ for some $1 \leq i \leq k$. Assume by contradiction that $\mathcal{N} = \mathcal{B}_f \setminus \mathcal{S}_i$ $\cup_{1\leq i\leq k} B(\mu_i, f)$ exhibits positive Lebesgue measure, note that N is an finvariant subset of \mathcal{B}_f . Applying Lemma [6.11](#page-28-1) and [\(6.18\)](#page-28-2) to $A := \mathcal{N}$ and $\eta < L_0$, with the same argument as in the proof of Item [\(1\)](#page-26-1), one obtains $B(\mu_i, f) \cap \mathcal{N} \neq \emptyset$ for some $1 \leq i \leq k$. This is a contradiction to the definition of $\mathcal N$, thus we complete the proof of Item [\(2\)](#page-26-2).

6.4. Skeleton generated by physical measures: Proof of Theorem **[D.](#page-20-1)** We have given the existence of skeletons for diffeomorphisms in $PH_C^1(M)$ by Theorem [6.2.](#page-20-6) Moreover, among $PH_C^{1+}(M)$, Proposition [6.8](#page-24-2) and Theorem [6.9](#page-26-0) tell us that each physical measure μ_i is attached with a periodic point $p_{j(i)}$ in the skeleton, and $i \mapsto j(i)$ is injective. Thus, for proving Theorem [D,](#page-20-1) it remains to show the map $i \mapsto j(i)$ is indeed bijective. To this end, we will establish the skeleton in a finer way than those in Theorem [6.2.](#page-20-6)

Proof of Theorem [D.](#page-20-1) By Theorem [6.9,](#page-26-0) consider μ_1, \dots, μ_k for some $k \leq \kappa_f$ as the family of physical measures of f , which are also the extreme elements

of $G^{cu}(f)$. Fixing $H_f > 0$, $\alpha > 0$, $0 < \varepsilon < D_f/2$, $\ell \in \mathbb{N}$, $0 < \rho_\ell \ll \delta_\ell$ as in Subsection [6.3.](#page-26-3) As before, for each $1 \leq i \leq k$, let \mathscr{A}_i be the family of finitely many ρ_{ℓ} -balls associated to μ_i . Let $\mathbb B$ be the union of ρ_{ℓ} -balls from \mathscr{A}_i , $1 \leq i \leq k$, and the notion $HT_{\mathbb{B}}(x, e^{-H_f/2}, f)$ is given by [\(6.16\)](#page-27-2). For the sake of more detailed argument, putting

- $\#(\mathscr{A}_i) = L_i;$
- $\mathscr{A}_i = \{B_{i,j} : 1 \leq j \leq L_i\}, \quad i = 1, \cdots, k;$
- Let $p_{i,j} := p_{B_{i,j}}$ be the hyperbolic periodic point in $B_{i,j}$ given by Theorem [3.6.](#page-11-1)

As established in the proof of Theorem [6.9,](#page-26-0) consider $C_{i,j} := C_{B_{i,j}}$ and $\mathcal{P}_{i,j} :=$ $\mathcal{P}_{B_{i,j}}$ as the cylinder and measurable partition around $B_{i,j}$, respectively.

Lemma 6.12. $S_0 = \{p_{i,j} : 1 \le i \le k, 1 \le j \le L_i\}$ is a pre-skeleton of f.

Proof. Let us take any C^1 disk D transverse to E^{cs} , we need to find $p_{i,j} \in \mathcal{S}_0$ such that $D \pitchfork W^s(\text{Orb}(p_{i,j}, f)) \neq \emptyset$. From [\(1\)](#page-13-0) of Theorem [4.2,](#page-13-1) one can pick $x \in D \cap \mathcal{B}_f$, using Lemma [6.10](#page-27-1) we have

$$
\mathcal{D}_U\left(HT_{\mathbb{B}}(x, e^{-H_f/2}, f)\right) \ge D_f/2.
$$

According to [\(6.15\)](#page-27-3), there is $n \in HT_{\mathbb{B}}(x, e^{-H_f/2}, f)$ such that $f^{(n)}(D)$ contains a disk of radius δ_{ℓ} centered at $f^{n}(x) \in \mathbb{B}$. Therefore, there is some $B_{i,j} \in \mathscr{A}_{i}$ such that $f^{n}(D)$ crosses the cylinder $C_{i,j}$. In particular, we have

$$
f^{n}(D) \pitchfork W_{\delta_{\ell}}^{s}(p_{i,j},f) \neq \emptyset.
$$

Hence, we get $D \pitchfork W^s(\text{Orb}(p_{i,j}, f)) \neq \emptyset$ and complete the proof of Lemma $6.12.$

Lemma [5.2](#page-19-2) guarantees that S_0 contains a subset to be a skeleton. Furthermore, we have:

Lemma 6.13. If $S_f \subset S_0$ is a skeleton of f, then for each $1 \leq i \leq k$, there exists a unique $1 \leq j_i \leq L_i$ such that $p_{i,j_i} \in \mathcal{S}_f$.

Proof. Let $S_f \subset S_0$ be a skeleton of f. Now we show firstly that for each $1 \leq i \leq k$, different elements of $\{p_{i,j} : 1 \leq j \leq L_i\}$ are homoclinically related to each other. Given $1 \leq i \leq k$, let us take $1 \leq m, n \leq L_i$ with $m \neq n$. Recalling the construction of balls in \mathscr{A}_i , suppose

$$
B_{i,m} := B(x_m, \rho_\ell), \quad B_{i,n} := B(x_n, \rho_\ell)
$$

for some $x_m, x_n \in \text{supp}(\mu_i | \Lambda_\ell(f, \alpha))$. Therefore, for any $\rho > 0$, we have

$$
\mu_i(B(x_m,\rho)\cap\Lambda_\ell(f,\alpha))>0,\quad \mu_i(B(x_n,\rho)\cap\Lambda_\ell(f,\alpha))>0.
$$

By Lemma [3.4,](#page-10-4) one can find integers $K < N$ and a hyperbolic periodic point p of periodic n with properties as follows:

 $-p \in B_{i,m}$ and $f^K(p) \in B_{i,n};$

– both p and $f^K(p)$ have stable and unstable manifolds of size larger than δ_{ℓ} .

Since $\rho_{\ell} \ll \delta_{\ell}$, the above properties imply $p \sim p_{i,m}$ and $f^{K}(p) \sim p_{i,n}$, thus $p_{i,n} \sim p_{i,m}$. Now, to complete the proof it suffices to show that for each $1 \leq i \leq k$, there exists at least one $1 \leq j \leq L_i$ such that $p_{i,j} \in \mathcal{S}_f$. Suppose by contradiction that there is $1 \leq i_0 \leq k$ such that $p_{i_0,j} \notin S_f$ for any $1 \leq j \leq L_{i_0}$. From Proposition [6.8,](#page-24-2) we know that for μ_{i_0} , there exists some $p_{i,j} \in \mathcal{S}_f$ with $i \neq i_0$ and an unstable disk D_{i_0} satisfying

- (1) Lebesgue almost every point of D_{i_0} belongs to $B(\mu_{i_0}, f)$;
- (2) D_{i_0} intersects $W^s(\text{Orb}(p_{i,j}, f))$ transversally.

On the other hand, by Lemma [4.13,](#page-18-1) there exists $D_{i,j} \in \mathcal{P}_{i,j}$ such that Leb $D_{i,j}(\Lambda_\ell(f,\alpha)\cap B(\mu_i,f))>0$. Applying inclination lemma to [\(2\)](#page-31-2), there is $n \in \mathbb{N}$ so that $f^n(D_{i_0})$ contains a disk \widetilde{D}_{i_0} which is C^1 -close to $W^s_{\delta_{\ell}}(p_{i,j}, f)$ and crosses $C_{i,j}$. Thus, in view of the choice of $D_{i,j}$, using the absolute continuity of stable lamination one must have $\text{Leb}_{D_{i_0}}(B(\mu_i, f)) > 0$, which is a contradiction to [\(1\)](#page-31-1).

By Lemma [6.13,](#page-30-1) we know that for each μ_i , there exists a unique $p_{i,j_i} \in \mathcal{S}_f$ and an unstable disk D_i satisfying

- $-D_i$ is contained in the support of μ_i , and Lebesgue almost every point of D_i belongs to $B(\mu_i, f)$;
- D_i intersects $W^s(\text{Orb}(p_{i,j_i}, f))$ transversally.

Therefore, we have $\mu_i \leftrightarrow p_{i,j_i}$, as showed in Proposition [6.8.](#page-24-2) Then, using Lemma [5.3](#page-19-1) to get the desired result.

7. Proofs of the main results

In this section we will provide the proofs of Theorems [A,](#page-3-0) [B](#page-4-0) and [C.](#page-4-1) Now we show firstly the existence of skeletons and the upper semi-continuity of the cardinality of skeleton w.r.t. diffeomorphisms in $PH_{EC}^1(M)$.

Theorem 7.1. Every $f \in PH_{EC}^1(M)$ admits a skeleton. If $\mathcal{S}(f) = \{p_i :$ $1 \leq i \leq k_f$ is a skeleton of f, then there exists a C^1 neighborhood U of f such that for every $g \in \mathcal{U}$, the continuation $\{p_i(g): 1 \leq i \leq k_f\}$ of $\mathcal{S}(f)$ is a pre-skeleton of g. Consequently, we have $k_g \leq k_f$ for every $g \in \mathcal{U}$.

Proof. By Theorem [6.1,](#page-20-2) there is $n_0 \in \mathbb{N}$ such that $f^{n_0} \in PH^1_C(M)$. Then by the construction of skeletons in Theorem [6.2,](#page-20-6) there exists a C^1 neighborhood \mathcal{U}_0 of f^{n_0} such that any diffeomorphism in it admits a skeleton, whose periodic points exhibit local stable and unstable manifolds with uniform size. Take a C^1 neighborhood $\mathcal U$ of f such that $g^{n_0} \in \mathcal U_0$ whenever $g \in \mathcal U$. By applying Lemmas [5.1](#page-19-4) and [5.2,](#page-19-2) we obtain that every diffeomorphism in U admits a skeleton that formed by periodic points admitting local stable and unstable manifolds with uniform size. Using the similar argument as in the proof of Theorem [6.6,](#page-23-1) mainly the continuity of the compact part of stable manifolds of periodic points w.r.t. diffeomorphisms, one can conclude the desired result.

Notation. Let f be a homeomorphism on M and μ a probability on M, define

$$
\mu^{(n)} = \frac{1}{n} \sum_{i=0}^{n-1} f_*^i \mu, \quad n \in \mathbb{N}.
$$

By definition, if μ is f^n -invariant, then $\mu^{(n)}$ is f -invariant. Moreover, one has the next result.

Lemma 7.2. Let f be a homeomorphism on M, assume that μ and ν are f^n -ergodic measures. Then we have that $\mu^{(n)} = \nu^{(n)}$ if and only if $\mu = f^i_* \nu$ *for some* $i \in \{0, 1 \cdots, n-1\}.$

Proposition 7.3. Given $N \in \mathbb{N}$ and a C^{1+} diffeomorphism f on M with $f^N \in PH_C^{1+}(M)$. Let μ_1, \cdots, μ_k be physical measures of f^N . Assume that $\mathcal{S}(f^N) = \{p_1, \cdots, p_k\}$ is a skeleton of f^N satisfying $\mu_i \stackrel{f^N}{\longleftrightarrow} p_i$ for every $1 \leq i \leq k$. Then the following properties hold:

(1)
$$
\mu_i^{(n)} = \mu_j^{(n)}
$$
 implies $p_i \stackrel{f}{\sim} p_j$.
\n(2) $p_i \stackrel{f}{\sim} p_j$ implies $\mu_i^{(n)} = \mu_j^{(n)}$.

Proof. We prove Item [\(1\)](#page-32-0) firstly. By Lemma [7.2,](#page-32-1) $\mu_i^{(n)} = \mu_j^{(n)}$ $j^{(n)}$ implies $\mu_i = f_*^{\kappa} \mu_j$ for some $\kappa \in \{0, 1, \dots, N-1\}$. Let us fix $\ell, \alpha, \delta_{\ell}, \rho_{\ell}$ for the diffeomorphism f^N as in Subsection [6.3.](#page-26-3) By our definition of Pesin blocks, one can verify that there exist $\ell_0 := \ell_0(\ell, \alpha, \kappa), \ \alpha_0 := \alpha_0(\ell, \alpha, \kappa)$ such that $f^{\kappa}(x) \in \Lambda_{\ell_0}(f^N, \alpha_0)$ whenever $x \in \Lambda_{\ell}(f^N, \alpha)$, that is,

(7.19)
$$
f^{\kappa}(\Lambda_{\ell}(f^N,\alpha)) \subset \Lambda_{\ell_0}(f^N,\alpha_0).
$$

For any small $\rho \leq \rho_{\ell}$, one can create skeleton $\mathcal{S}_{f^N} = \{P_1, \cdots, P_k\}$ as in Lemmas [6.12](#page-30-0) and [6.13,](#page-30-1) with properties as follows:

 $- \mu_i \stackrel{f^N}{\longleftrightarrow} P_i$ for every $1 \leq i \leq k$;

– up to shrinking δ_{ℓ} , we assume each $f^j(P_i)$, $1 \leq i \leq k$, $0 \leq j \leq \kappa$ has stable and unstable manifolds of size larger than δ_{ℓ} .

Moreover, we assume that each P_i is contained in $B(x_i, \rho) \in \mathscr{A}_{\mu_i}(f^N, \ell, \alpha, \rho)$ for some $x_i \in \text{supp}(\mu_i | \Lambda_\ell(f^N, \alpha))$. As before, let C_i be the cylinder around $B(x_i, \rho)$ for every $1 \leq i \leq k$. By continuity of f^{κ} , one can take $\delta := \delta(\rho) > 0$ so that

(7.20)
$$
f^{\kappa}B(x,\rho) \subset B(f^{\kappa}(x),\delta) \text{ for every } x \in M.
$$

Using [\(7.19\)](#page-32-2), [\(7.20\)](#page-32-3) together with the fact $\mu_i = f^{\kappa}_* \mu_j$, we obtain

$$
\mu_i\left(B(f^{\kappa}(x_j),\delta) \cap \Lambda_{\ell_0}(f^N,\alpha_0)\right) \geq \mu_i\left(f^{\kappa}(B(x_j,\rho)) \cap f^{\kappa}(\Lambda_{\ell}(f^N,\alpha))\right) \n= \mu_j\left(B(x_j,\rho) \cap \Lambda_{\ell}(f^N,\alpha)\right) \n> 0.
$$

Note that one can make δ small by choosing ρ sufficiently small.

Recall that we have $\mu_i(B(x_i, \rho) \cap \Lambda_\ell(f^N, \alpha)) > 0$ by construction. With the ergodicity of μ_i , by applying Lemma [3.4,](#page-10-4) we know that up to reducing δ_{ℓ} if necessary, by taking ρ small enough, there exist $r \ll \delta_{\ell}$ and $m \in \mathbb{N}$ such that

- there exists a hyperbolic periodic point P, which is r-close to $f^{\kappa}(P_i)$ and exhibits stable and unstable manifolds of size larger than δ_{ℓ} .
- $-f^m(P)$ is r-close to P_i , with local stable and unstable manifolds of size larger than δ_{ℓ} .

These two facts imply that $f^{\kappa}(P_j) \stackrel{f^N}{\sim} P_i$, so $f^{\kappa}(P_j) \stackrel{f}{\sim} P_i$. Therefore, we get $P_j \stackrel{f}{\sim} P_i$ immediately. On the other hand, following from Lemma [5.3](#page-19-1) and Proposition [6.8,](#page-24-2) we have $p_i \stackrel{f^N}{\sim} P_i$ and $p_j \stackrel{f^N}{\sim} P_j$, thus $p_i \stackrel{f}{\sim} P_i$ and $p_j \stackrel{f}{\sim} P_j$. Altogether, we obtain $p_i \stackrel{f}{\sim} p_j$.

To prove Item [\(2\)](#page-32-4), consider skeleton S_{fN} as above, by Proposition [6.8,](#page-24-2) we can choose C^1 unstable disks D_i, D_j such that for every $* \in \{i, j\}$

- $-D_*$ intersects $W^s(\text{Orb}(P_*, f^N))$ transversely;
- Lebesgue almost every point of D_* is contained in $B(\mu_*, f^N)$.

Following from $p_i \stackrel{f}{\prec} p_j$, one gets $P_i \stackrel{f}{\prec} P_j$ immediately. Therefore, there is $L \in \mathbb{N}$ such that $W^u(f^L(P_i), f)$ intersects $W^s(P_j, f)$ transversely. Note that Lebesgue almost every point of $f^L(D_i)$ is contained in $B(f^L\mu_i, f^N)$. By using inclination lemma, there exists m as a multiple of N and a disk D_1 inside $f^{L+m}(D)$ and a disk D_2 inside $f^m(D_j)$ such that, both D_1 and D_2 cross the cylinder C_j around P_j . Then Lemma [4.13](#page-18-1) gives a disk D_0 crossing C_i that admits a subset with positive Lebesgue measure, which possess stable manifolds of size larger than δ_{ℓ} . With the absolute continuity of stable lamination, one deduce that $\mu_j = f_*^L \mu_i$. Hence, we get $\mu_i^{(n)} = \mu_j^{(n)}$ j by Lemma [7.2.](#page-32-1) This completes the proof of Proposition [7.3.](#page-32-5)

Now we can give the proof of Theorem of [A](#page-3-0) by applying Proposition [7.3](#page-32-5) and the previous results.

Proof of Theorem [A.](#page-3-0) Let $f \in PH_{EC}^{1+}(M)$ with partially hyperbolic splitting $TM = E^u \oplus_{\succ} E^{cu} \oplus_{\succ} E^{cs}$. By Theorem [6.1,](#page-20-2) we have $f^N \in PH_C^{1+}(M)$ for some $N \in \mathbb{N}$. By Theorem [6.9](#page-26-0) and Theorem [D,](#page-20-1) there exists $s \in \mathbb{N}$ with physical measures μ_1, \dots, μ_s for f^N and skeleton $\mathcal{S}(f^N) = \{p_1, \dots, p_s\}$ for f^N such that

- (1) $\mu_i \stackrel{f^N}{\longleftrightarrow} p_i$ for every $1 \leq i \leq s$;
- (2) the union of $B(\mu_1, f^N), \cdots B(\mu_s, f^N)$ has full Lebesgue measure.

Claim 7.4. Rearrange if necessary, there is $1 \leq k \leq s$ such that $\{p_1, \dots, p_k\}$ is a skeleton of f, and $\mu_i^{(N)}$ $i^{(N)}$, $1 \leq i \leq k$ are (ergodic) physical measures with basin covering property. In addition, $\mu_i^{(N)}$ $i^{(N)} \leftrightarrow p_i$ for every $1 \leq i \leq k$.

By assuming Claim [7.4](#page-34-0) and using Lemma [5.3,](#page-19-1) we complete the proof of Theorem [A.](#page-3-0)

Now it remains to give the proof of the claim.

Proof of Claim [7.4.](#page-34-0) We obtain the ergodicity of $\mu_i^{(N)}$ $i^{(N)}$, $1 \leq i \leq s$ by construction. Moreover, we have $B(\mu_i, f^N) \subset B(\mu_i^{(N)})$ $\binom{y}{i}$, f) for every $1 \leq i \leq s$, hence ${\rm Leb}(B(\mu_i^{(N)}))$ $\mathcal{L}^{(N)}_i(f)) \geq \text{Leb}(B(\mu_i, f^N)) > 0$, which means that $\{\mu_i^{(N)}\}$ $i_i^{(N)}: 1 \leq i \leq s$ are physical measures for f ; from (2) one gets that the union of basins of $\{\mu_i^{(N)}\}$ $i^{(N)}: 1 \leq i \leq s$ w.r.t. f covers a full Lebesgue measure subset. Note that it may be happened that $\mu_i^{(N)} = \mu_j^{(N)}$ $j_j^{(N)}$ for some $i, j \in \{1, \cdots, s\}$ with $i \neq j$.

It follows from [\(1\)](#page-34-1) and the construction that we have

(7.21)
$$
\mu_i^{(N)} \xleftarrow{f} p_i \quad \text{for every } 1 \le i \le s.
$$

Since $\mathcal{S}(f^N) = \{p_1, \cdots, p_s\}$ is a skeleton for f^N , by applying Lemma [5.1](#page-19-4) we know that $\{p_1, \cdots, p_s\}$ is a pre-skeleton for f . Taking $\mathcal{P} \subset \{\mu_i^{(N)}\}$ $i_i^{(N)}: 1 \leq i \leq s$ such that elements of P are different to each other and exhibits the basin covering property. By reordering if necessary, we may assume

$$
\mathcal{P} = {\mu_1^{(N)}, \cdots, \mu_k^{(N)}} \quad \text{for some } 1 \le k \le s.
$$

It suffices to prove that $S = \{p_1, \dots, p_k\}$ is a skeleton of f. We show that S satisfies ([S1\)](#page-2-2) firstly. For every C^1 disk D transverse to E^{cs} , since $\mathcal{S}(f^N)$ is a pre-skeleton of f, there is $p_i \in \mathcal{S}(f^N)$ such that D intersects $W^s(\text{Orb}(p_i, f))$ transversely. In addition, one can make i to satisfy $i \leq k$. Indeed, if k $i \leq s$, then by the choice of P , there is $j \leq k$ so that $\mu_j^{(N)} = \mu_i^{(N)}$ $i^{(IV)}$. By using [\(1\)](#page-32-0) of Proposition [7.3,](#page-32-5) we get $p_i \stackrel{f}{\sim} p_j$. This together with inclination lemma implies that D must intersect $W^s(\text{Orb}(p_j, f))$ transversely also. This says that S satisfies condition ([S1\)](#page-2-2). Now we show that S satisfies condition ([S2\)](#page-2-4). By absurd we have $p_i \stackrel{f}{\prec} p_j$ for some $i, j \in \{1, \cdots, k\}$, then by applying [\(2\)](#page-32-4) of Proposition [7.3](#page-32-5) we get $\mu_i^{(N)} = \mu_j^{(N)}$ $j^{(N)}$, this is a contradiction. Now, we can use (7.21) and Lemma [5.4](#page-19-3) to complete the proof.

We are in ready to give the proof of Theorem [B.](#page-4-0)

Proof of Theorem [B.](#page-4-0) By Theorem [7.1](#page-31-3) and Theorem [A,](#page-3-0) we get the upper semi-continuity of the number of physical measures and the cardinality of skeleton in C^1 -topology among $PH_{EC}^{1+}(M)$. Indeed, as showed in Theorem [7.1,](#page-31-3) for each diffeomorphism $f \in PH_{EC}^{1+}(M)$, there exists a C^1 neighborhood V of f such that $k_g \leq k_f$ for every $g \in V$. Furthermore, there exists a C^1 open and dense subset V_0 of V on which the cardinality of skeleton is locally constant. To this end, we consider a sequence of subsets of V as follows:

 $\mathcal{U}_0 = \emptyset; \quad \mathcal{U}_i = \{g \in \mathcal{V} : k_g \leq i\} \quad \text{when} \quad 1 \leq i \leq k_f.$

By definition, it suffices to take $V_0 = \bigcup_{1 \leq i \leq k_f} (\mathcal{U}_i \setminus \overline{\mathcal{U}_{i-1}})$. Using Theorem [A,](#page-3-0) we know that the number of the physical measures for C^{1+} diffeomorphisms of V_0 is also locally constant.

Now we verify the continuity of the physical measures w.r.t. C^{1+} diffeomorphisms of V_0 in following precise way:

Lemma 7.5. Let g be a C^{1+} diffeomorphism in V_0 . Assume that $S(g)$ = $\{p_1, \dots, p_{k_g}\}\$ is a skeleton of g and $\{\mu_i : 1 \leq i \leq k_g\}$ are physical measures of g satisfying $\mu_i \leftrightarrow_{\mathcal{P}}^g$ for each $1 \leq i \leq k_g$. If g_n is a sequence of C^{1+} diffeomorphisms in V_0 that converges to g in C^1 -topology, then for every $n \in \mathbb{N}$ sufficiently large, we have

- the continuation of $S(g)$, denoted by $S(g_n) = \{p_1(g_n), \cdots, p_{k_g}(g_n)\},\$ is a skeleton of q_n .
- if we let $\{\mu_{i,n}: 1 \leq i \leq k_q\}$ be the set of physical measures of g_n satisfying $\mu_{i,n} \xleftarrow{g_n} p_i(g_n)$ for each $1 \leq i \leq k_g$, then $\mu_{i,n} \xrightarrow{weak^*} \mu_i$ as $n \to +\infty$ whenever $i \in \{1, \cdots, k_g\}.$

Proof. As a direct consequence of Theorem [7.1](#page-31-3) and the definition of V_0 , one knows that the first item is true. Now we make effort to show the second item. By Theorem [6.1,](#page-20-2) there is $N \in \mathbb{N}$ such that $f^N \in PH_C^{1+}(M)$. Consequently, shrinking V if necessary, we may assume $g^N \in PH_C^{1+}(M)$ with dominated splitting $TM = E^u \oplus E^{cu} \oplus E^{cs}$ for every $g \in V$. In view of Claim [7.4,](#page-34-0) without loss of generality we can assume $\mathcal{V} \subset PH^1_C(M)$.

If the conclusion is not true, then there exists $\varkappa \in \{1, \cdots, k_g\}$ such that $\mu_{\varkappa,n}$ does not converge to μ_{\varkappa} when n goes to infinite. Note that all physical measures are Gibbs cu-states, by the upper semi-continuity of Gibbs cuspace w.r.t. diffeomorphisms as stated in Theorem [4.5,](#page-14-4) up to considering subsequence we assume $\mu_{\varkappa,n}$ converges to $\mu \in G^{cu}(q)$ different to μ_{\varkappa} . It follows from [\(1\)](#page-26-1) of Theorem [6.9](#page-26-0) that μ is a convex combination of physical measures of g, thus there exists $\xi_i \in [0,1]$ such that

$$
\mu = \sum_{i=1}^{k_g} \xi_i \mu_i, \quad \text{where } \sum_{i=1}^{k_g} \xi_i = 1.
$$

Since $\mu \neq \mu_{\varkappa}$, there exists $\iota \in \{1, \cdots, k_q\}$ with $\iota \neq \varkappa$ such that $\xi_{\iota} > 0$.

Choosing $\varepsilon < \xi_{\iota}/(1+\xi_{\iota})$, Proposition [3.5](#page-11-0) applies to g and ε , we know that there exist $\alpha > 0$, $\ell \in \mathbb{N}$ and a C^1 neighborhood $\mathcal{W} \subset \mathcal{V}$ of g such that for every $h \in \mathcal{W}$ we have

(7.22)
$$
\mu(\Lambda_{\ell}(h,\alpha)) > 1 - \varepsilon, \quad \forall \mu \in G^{u}(h).
$$

By Lemma [3.3,](#page-9-2) there exist $\rho_{\ell} > 0$ and $\delta_{\ell} > 0$ such that for every $h \in \mathcal{W}$, for every $\mu \in G^u(h)$ and $x \in \text{supp}(\mu | \Lambda_\ell(h, \alpha))$, any ball $B(x, \rho)(\rho \le \rho_\ell)$ contains hyperbolic periodic points with stable and unstable manifolds of size larger than δ_{ℓ} .

For fixed μ_{ι} , which is a Gibbs u-state of g, take $\rho \leq \rho_{\ell}$ within $\rho \ll \delta_{\ell}$, recalling the definition of $\mathscr{A}_{\mu}\left(g,\ell,\alpha,\rho\right)$ in Theorem [3.6,](#page-11-1) and put

$$
A_{\rho} = \bigcup_{B \in \mathscr{A}_{\mu_{\iota}}(g,\ell,\alpha,\rho)} B.
$$

For every $B \in \mathscr{A}_{\mu_{\iota}}(g,\ell,\alpha,\rho)$, let $P_{B,\iota}$ be a fixed hyperbolic periodic point in it which admits stable and unstable manifolds of size larger than δ_{ℓ} . Moreover, In view of the proof of Lemma [6.13,](#page-30-1) we know

$$
(7.23) \t\t P_{B,\iota} \stackrel{g}{\sim} p_{\iota}.
$$

Using [\(7.22\)](#page-36-0) to g and μ_{ι} , we have $\mu_{\iota}(A_{\rho}) > 1 - \varepsilon$, which implies $\mu(A_{\rho}) >$ $\xi_{\iota}(1-\varepsilon)$ by representation of μ . Since $\mu_{\varkappa,n} \xrightarrow{weak^*} \mu$ as $n \to +\infty$, and A_{ρ} is open, we get

$$
\mu_{\varkappa,n}(A_{\rho}) > \xi_{\iota}(1 - \varepsilon)
$$

for any n large enough. On the other hand, note that $\mu_{\varkappa,n}$ is a Gibbs u-state of g_n for every n, use [\(7.22\)](#page-36-0) again we obtain $\mu_{\varkappa,n}(\Lambda_\ell(g_n,\alpha)) > 1-\varepsilon$ for every n sufficiently large. Combining this with [\(7.24\)](#page-36-1) and our choice of ε , we see that for every large n ,

(7.25)
$$
\mu_{\varkappa,n}(\Lambda_{\ell}(g_n,\alpha)\cap A_{\rho})>0.
$$

Observe that $\mathscr{A}_{\mu_{\iota}}(g,\ell,\alpha,\rho)$ is consisted of finitely many ρ -balls. By [\(7.25\)](#page-36-2), up to taking subsequence, one may fix $B \in \mathscr{A}_{\mu_{\iota}}(g, \ell, \alpha, \rho)$ such that

$$
\mu_{\varkappa,n}(\Lambda_{\ell}(g_n,\alpha)\cap B)>0
$$
 for every large *n*.

This implies that for every n large enough, one can take some point $x_n \in$ $\text{supp}(\mu_{\varkappa,n}|\Lambda_{\ell}(g_n,\alpha))\cap B$. From the choices of ρ and δ_{ℓ} , there exists a hyperbolic periodic point $p_{\varkappa,n} \in B(x_n, \rho)$, which has stable and unstable manifolds of size larger than δ_{ℓ} . According Lemma [6.13,](#page-30-1) for each n, one can take a hyperbolic periodic point $p_{\varkappa}^{(n)}$ in a ball $B_{\varkappa,n} \in \mathscr{A}_{\mu_{\varkappa,n}}(g_n,\ell,\alpha,\rho)$ such that $\mu_{\varkappa,n} \leftrightarrow p_{\varkappa}^{(n)}$. Since we have also $\mu_{\varkappa,n} \leftrightarrow p_{\varkappa}(g_n)$ from assumption, we get $\left(\ldots \right)$ a_{n}

$$
(7.26) \t\t\t p_{\varkappa}^{(n)} \stackrel{g_n}{\sim} p_{\varkappa}(g_n).
$$

By construction, we have

$$
\mu_{\varkappa,n}(B_{\varkappa,n}\cap\Lambda_{\ell}(g_n,\alpha))>0,\quad \mu_{\varkappa,n}(B(x_n,\rho)\cap\Lambda_{\ell}(g_n,\alpha))>0.
$$

As $\mu_{\varkappa,n}$ is g_n -ergodic, by Lemma [3.4,](#page-10-4) similar to the argument in the proof of Lemma [6.13,](#page-30-1) we get $p_{\varkappa}^{(n)} \stackrel{g_n}{\sim} p_{\varkappa,n}$, choosing ρ smaller if necessary. This together with [\(7.26\)](#page-36-3) yields

$$
(7.27) \t\t p_{\varkappa,n} \stackrel{g_n}{\sim} p_{\varkappa}(g_n).
$$

For every n large enough, denote by $P_{B,\iota,n}$ the continuation of $P_{B,\iota}$ w.r.t. g_n . Using [\(7.23\)](#page-36-4) and the fact $d(P_{B,\iota}, p_{\kappa,n}) < 2\rho$, $\rho \ll \delta_{\ell}$, it follows that one can fix n_0 large enough such that both $p_{\iota}(g_{n_0})$ and p_{\varkappa,n_0} are homoclinically related to P_{B,ι,n_0} , thus $p_{\iota}(g_{n_0}) \stackrel{g_{n_0}}{\sim} p_{\varkappa,n_0}$. This together with [\(7.27\)](#page-37-1) by taking $n = n_0$ yields $p_{\varkappa}(g_{n_0}) \stackrel{g_{n_0}}{\sim} p_{\iota}(g_{n_0})$, this is a contradiction to the definition of skeletons.

The proof of Lemma [7.5](#page-35-0) is complete now.

Observe that for each fixed $L > 0$, by applying the unstable manifold theorem of hyperbolic periodic point, the compact part $W_L^u(\text{Orb}(p_i(g), g))$ depends continuously on diffeomorphism g , which implies that the closure of $W^u(\text{Orb}(p_i(g), g))$ vary lower semi-continuously on g with the Hausdorff topology. By applying Theorem [A,](#page-3-0) we have the lower semi-continuity of the

 \Box

We close this section by finishing the proof of Theorem [C.](#page-4-1)

Proof of Theorem [C.](#page-4-1) We show the weak statistical stability firstly. We start by taking a sequence of diffeomorphisms $f_n \in PH_{EC}^{1+}(M)$ that converges to $f \in PH_{EC}^{1+}(M)$ in C¹-topology, and let μ_n be the physical measure of f_n respectively. Up to considering subsequences, we assume that μ_n converges to some $\mu \in \mathcal{M}(f)$. Since all ergodic Gibbs cu-states are physical measures of diffeomorphisms in $PH_{EC}^{1+}(M)$, by applying Theorem [4.5](#page-14-4) we know that μ is a Gibbs cu-state of f, which is a convex combination of the physical measures of f .

supports of physical measures. Now we have complete the proof of Theorem

Assume that $f \in PH_{EC}^{1+}(M)$ admits a unique physical measure μ_f . By Theorem [B](#page-4-0) there exists a C^1 neighborhood V of f such that every $g \in$ $V \cap PH_{EC}^{1+}(M)$ has a unique physical measure μ_g also, and one must have $\mu_{f_n} \mapsto \mu_f$ whenever $f_n \in \mathcal{V} \cap PH_{EC}^{1+}(M)$ converges to f in C¹-topology. \Box

Appendix A. Proof of Proposition [3.5](#page-11-0)

The following lemma given by [\[18,](#page-42-5) Lemma 5.8] is a variation version of the Pliss-Like Lemma founded by Andersson-Vasquez [\[8,](#page-41-5) Lemma A].

Lemma A.1. Given constants $C_1 < C_2 \le \max\{0, C_2\} < C$, for any $\varepsilon > 0$, there is $\rho = \rho(C_1, C_2, C, \varepsilon) > 0$ such that for any any sequence $\{a_n\}_{n \in \mathbb{N}} \subset \mathbb{R}$ satisfying:

• $|a_n| \leq C, \forall n \in \mathbb{N},$

• there is a subset $P \subset \mathbb{N}$ such that $\mathcal{D}_L(P) > 1 - \rho$ and $a_n \leq C_1$ for any $n \in P$,

then there is a subset $Q \subset \mathbb{N}$ with $\mathcal{D}_U(Q) > 1 - \varepsilon$ such that for any $j \in Q$, one has that

$$
\sum_{i=0}^{n-1} a_{i+j} \le nC_2, \ \forall n \in \mathbb{N}.
$$

The following lemma plays a key role in the proof of Proposition [3.5.](#page-11-0)

Lemma A.2. Let f be a C^1 diffeomorphism with dominated splitting $TM =$ $E \oplus F$. Given $0 < \alpha < \alpha_0$, if μ is an f-invariant measure such that

(A.28)
$$
\lim_{n \to +\infty} \frac{1}{n} \log \|Df^{-n}|_{E(x)} \| < -\alpha_0, \quad \mu - a.e. \ x \in M.
$$

(A.29)
$$
\lim_{n \to +\infty} \frac{1}{n} \log \|Df^n|_{F(x)} \| < -\alpha_0, \quad \mu - a.e. \ x \in M.
$$

Then, for every $\varepsilon > 0$ there exists $\ell \in \mathbb{N}$ and a neighborhood U of f such that for every $q \in \mathcal{U}$ and g-invariant measure ν , we have

$$
\nu\left(\Lambda_{\ell}(g,\alpha,E,F)\right) > 1 - \varepsilon.
$$

Proof. For simplicity, for every g that is C^1 -close to f, we introduce the two blocks $\Lambda_{\ell}^-(g, \alpha)$ and $\Lambda_{\ell}^+(g, \alpha)$ defined as follows:

$$
\Lambda_{\ell}^-(g,\alpha) = \left\{ x : \frac{1}{n\ell} \sum_{i=0}^{n-1} \log \|Dg^{-\ell}|_{E(f^{-i\ell}(x))} \| \leq -\alpha, \quad \forall n \in \mathbb{N} \right\};
$$

$$
\Lambda_{\ell}^+(g,\alpha) = \left\{ x : \frac{1}{n\ell} \sum_{i=0}^{n-1} \log \|Dg^{\ell}|_{F(f^{i\ell}(x))} \| \leq -\alpha, \quad \forall n \in \mathbb{N} \right\}.
$$

Hence, $\Lambda_{\ell}(g, \alpha) = \Lambda_{\ell}^{-}(g, \alpha) \cap \Lambda_{\ell}^{+}$ $\ell^+(g, \alpha)$. Given $n \in \mathbb{N}$, let us define

$$
\Lambda_{g,n}^E = \left\{ x : \frac{1}{n} \log \|Dg^{-n}|_{E(x)} \| < -\alpha_0 \right\};
$$
\n
$$
\Lambda_{g,n}^F = \left\{ x : \frac{1}{n} \log \|Dg^n|_{F(x)} \| < -\alpha_0 \right\}.
$$

Take $\Lambda_{g,n} = \Lambda_{g,n}^E \cap \Lambda_{g,n}^F$.

Choose $\varepsilon' > 0$ satisfying $(1 - \varepsilon')^2 > 1 - \varepsilon/2$ and fix constants

$$
C_1 = -\alpha_0
$$
, $C_2 = -\alpha$, $C = \max_{x \in M} |\log ||Df^{\pm 1}|_{E(x)}|| + 1$.

Then take $\rho = \rho(C_1, C_2, C, \varepsilon)$ as in Lemma [A.1.](#page-37-2) It follows from [\(A.28\)](#page-38-0) and [\(A.29\)](#page-38-1) that for $0 < \rho \varepsilon' < 1$, there exists $\ell \in \mathbb{N}$ such that $\mu(\Lambda_{f,\ell}) > 1 - \rho \varepsilon'$. By the continuity of diffeomorphisms and [\[8,](#page-41-5) Lemma 3.2], there exists an open neighborhood $\mathbb U$ of (f, μ) such that

$$
(A.30) \t\t\t \nu(\Lambda_{g,\ell}) > 1 - \rho \varepsilon'
$$

and $\max_{x \in M} |\log ||Dg^{\pm 1}|_{E(x)}|| \leq C$ for every $(g, \nu) \in \mathbb{U}$. Consequently, we have $\nu(\Lambda_{g,\ell}^E) > 1 - \rho \varepsilon'$, $\nu(\Lambda_{g,\ell}^F) > 1 - \rho \varepsilon'$ for every $(g,\nu) \in \mathbb{U}$. This implies that there exists a C^1 open neighborhood U of f satisfying $(g, \nu) \in \mathbb{U}$ for every $g \in \mathcal{U}$ and g-invariant measure ν .

Since ν is g^{ℓ} -invariant, by Birkhoff's ergodic theorem we get that for ν -almost every $x \in M$, there exists the limit

$$
\phi(x) = \lim_{n \to +\infty} \frac{1}{n} \# \left\{ i : 0 \le i \le n - 1, g^{-\ell i}(x) \in \Lambda_{g,\ell}^E \right\}.
$$

Furthermore, applying [\(A.30\)](#page-38-2) we have

(A.31)
$$
\int \phi(x) d\nu = \nu(\Lambda_{g,\ell}^E) > 1 - \rho \varepsilon'.
$$

Let $G = \{x : 1 - \rho < \phi(x) \leq 1\}$, we claim that $\nu(G) > 1 - \varepsilon'$. Indeed, observe that $M \setminus G = \{x : \rho \leq 1 - \phi(x) \leq 1\}$, this together with [\(A.31\)](#page-39-0) imply

$$
\rho \varepsilon' > \int (1 - \phi) d\nu \ge \int (1 - \phi) \chi_{M \setminus G} d\nu \ge \rho \nu(M \setminus G).
$$

Thus, $\nu(G) > 1 - \varepsilon'$. For any $x \in G$, we define

$$
a_i = \frac{1}{\ell} \log ||Dg^{-\ell}|_{E(g^{-i\ell}x)}||, \quad \forall i \ge 0,
$$

and

$$
P = \{ i \ge 0 : g^{-i\ell}(x) \in \Lambda_{g,\ell}^E \}.
$$

Then, by properties of $\Lambda_{g,\ell}^E$ and G, one has that $a_i < -\alpha$ for every $i \in P$, and $\mathcal{D}_L(P) > 1 - \rho$. Thus, by applying Lemma [A.1,](#page-37-2) there exists a subset $Q \subset \mathbb{N}$ with density estimate

(A.32) D^U (Q) > 1 − ε ′ .

Moreover, for every $j \in Q$, we have

$$
\sum_{i=0}^{n-1} a_{i+j} \le -n\alpha, \quad \forall n \in \mathbb{N},
$$

which can be rewrite as

$$
\prod_{i=0}^{n-1} \|Dg^{-\ell}|_{E(g^{-i+j\ell}(x))}\| \le e^{-n\ell\alpha}, \quad \forall n \in \mathbb{N}.
$$

Therefore, by combining [\(A.32\)](#page-39-1) with Birkhoff's ergodic theorem we know that for ν -almost every $x \in G$,

$$
\lim_{n \to +\infty} \frac{1}{n} \# \left\{ i \in [0, n-1] : g^{-i\ell}(x) \in \Lambda_{\ell}^-(g, \alpha) \right\} > 1 - \varepsilon'.
$$

Therefore, there exists $n_0 \in \mathbb{N}$ and a subset

$$
G_0 = \left\{ x \in G : \frac{1}{n_0} \# \left\{ i \in [0, n_0 - 1] : g^{-i\ell}(x) \in \Lambda_{\ell}^-(g, \alpha) \right\} > 1 - \varepsilon' \right\}
$$

satisfying $\nu(G_0) > 1 - \varepsilon'$. Then, we have

$$
\nu(\Lambda_{\ell}^{-}(g,\alpha)) = \int \frac{1}{n_0} \sum_{i=0}^{n_0-1} \chi_{\Lambda_{\ell}^{-}(g,\alpha)}(g^{-i\ell}x) d\nu
$$

\n
$$
\geq \int_{G_0} \frac{1}{n_0} \sum_{i=0}^{n_0-1} \chi_{\Lambda_{\ell}^{-}(g,\alpha)}(g^{-i\ell}x) d\nu
$$

\n
$$
= \int_{G_0} \frac{1}{n_0} \# \left\{ i \in [0,n_0-1] : g^{-i\ell}(x) \in \Lambda_{\ell}^{-}(g,\alpha) \right\} d\nu
$$

\n
$$
\geq (1-\varepsilon')\nu(G_0)
$$

\n
$$
> (1-\varepsilon')^2 > 1-\varepsilon/2.
$$

Similarly, for F direction one can get $\nu(\Lambda_{\ell}^-(g, \alpha)) > 1 - \varepsilon/2$. Therefore,

$$
\nu(\Lambda_{\ell}(g,\alpha)) > 1 - \varepsilon.
$$

Now the proof of Lemma [A.2](#page-38-3) is completed. \square

By Lemma [2.6,](#page-8-3) one can conclude the following result. It can be seen as a combination of [\[40,](#page-43-8) Lemma 3.4] and [\[40,](#page-43-8) Lemma 3.5], though the definition of Gibbs u -state is different from that in [\[40\]](#page-43-8).

Lemma A.3. If $f \in PH_{EC}^1(M)$, then there exist $N \in \mathbb{N}$ and $\alpha > 0$ such that for every $\mu \in G^u(f)$, one has

$$
\lim_{n \to +\infty} \frac{1}{n} \sum_{i=0}^{n-1} \log \|Df^{-N}|_{E^{cu}(f^{-iN}(x))}\| < -\alpha
$$

for μ -almost every x.

Now we can finish the proof of Proposition [3.5.](#page-11-0)

Proof of Proposition [3.5.](#page-11-0) By Lemma [A.3,](#page-40-0) there exist $N \in \mathbb{N}$ and $\alpha > 0$ such that for every $\mu \in G^u(f)$ and μ -almost every x we have

$$
\lim_{n \to +\infty} \frac{1}{n} \sum_{i=0}^{n-1} \log \|Df^{-N}|_{E^{cu}(f^{-iN}(x))}\| < -\alpha,
$$

by taking $\alpha_0 = \alpha/N$, we get that for μ -almost every x,

$$
\lim_{n \to +\infty} \frac{1}{n} \log ||Df^{-n}|_{E^{cu}(x)}|| < -\alpha_0.
$$

Similarly, we have that

$$
\lim_{n \to +\infty} \frac{1}{n} \log ||Df^n|_{E^{cs}(x)}|| < -\alpha_0.
$$

for μ -almost every x. Now fix $\alpha \in (0, \alpha_0)$. By Lemma [A.2,](#page-38-3) for every $\varepsilon > 0$, for any $\mu \in G^u(f)$, there exists $\ell_\mu \in \mathbb{N}$ and neighborhood $\mathcal{U}_\mu \times \mathcal{V}_\mu$ of (f, μ)

in $G^u(PH^1_{EC}(M))^8$ $G^u(PH^1_{EC}(M))^8$ such that

 $(A.33)$ $\nu(\Lambda_{\ell_\mu}(g, \alpha)) > 1 - \varepsilon$

for every $(g, \nu) \in \mathcal{U}_{\mu} \times \mathcal{V}_{\mu}$. By compactness of $G^u(f)$ as stated in Item (1) of Lemma [2.6,](#page-8-3) there exist finitely many Gibbs u-states μ_1, \dots, μ_j such that the corresponding neighborhoods $V_i := V_{\mu_i}, 1 \leq i \leq j$ form a covering of $G^u(f)$. Writting $\mathcal{U}_i := \mathcal{U}_{\mu_i}$ and $\ell_i := \ell_{\mu_i}$ for every $1 \leq i \leq j$. Let

$$
\mathcal{U} = \bigcap_{1 \leq i \leq j} \mathcal{U}_i; \quad \mathcal{V} = \bigcup_{1 \leq i \leq j} \mathcal{V}_i; \quad \ell = \prod_{i=1}^j \ell_j.
$$

Note that by definition of ℓ , we have $\Lambda_{\ell_i}(g, \alpha) \subset \Lambda_{\ell}(g, \alpha)$ for every $1 \leq i \leq j$. By Lemma [2.7,](#page-8-2) up to shrinking U we assume $G^u(g) \subset V$ for every $g \in U$. Therefore, for every $(g, \nu) \in \mathcal{U} \times \mathcal{V}$, there is some $1 \leq i \leq j$ such that $(g, \nu) \in \mathcal{U}_i \times \mathcal{V}_i$, and thus $(A.33)$ implies $\nu(\Lambda_\ell(g, \alpha)) > 1 - \varepsilon$.

REFERENCES

- [1] J. Alves. SRB measures for non-hyperbolic systems with multidimensional expansion, Ann. Sci. Ecole Norm. Sup. (4), 33(1) (2000), 1-32.
- [2] J. Alves, C. Bonatti and M. Viana, SRB measures for partially hyperbolic systems whose central direction is mostly expanding, Invent. Math., 140 (2000), 351-398.
- [3] J. Alves. Strong statistical stability of non-uniformly expanding maps, Nonlinearity, $17(4)$ (2004), 1193-1215.
- [4] J. Alves, V. Pinheiro, Topological structure of (partially) hyperbolic sets with positive volume, Trans. Amer. Math. Soc., 360 (2008), 5551-5569.
- [5] J. Alves and Marcelo Viana. Statistical stability for robust classes of maps with non-uniform expansion, Ergod. Th.& Dynam. Sys., $22(1)$ (2002) , 1-32.
- [6] M. Andersson. Robust ergodic properties in partially hyperbolic dynamics, Trans. Amer. Math. Soc., 362 (2010), 1831-1867.
- [7] M. Andersson and C. Vasquez, On mostly expanding diffeomorphisms, Ergod. Th.& Dynam. Sys., 38 (2018), 2838-2859.
- [8] M. Andersson and C. Vasquez, Statistical stability of mostly expanding diffeomorphisms, [arXiv:1710.07970V](http://arxiv.org/abs/1710.07970)3.
- [9] L. Barreira, Y. Pesin, Nonuniform Hyperbolicity: Dynamics of Systems with Nonzero Lyapunov Exponents, Encyclopedia of Mathematics and its Applications. Cambridge University Press, 2007.
- [10] C. Bonatti, L. Diaz and M. Viana, Dynamics beyond uniform hyperbolicity, A global geometric and probabilistic perspective, Encyclopaedia of Mathematical Sciences, 102. Mathematical Physics, 2005. III. Springer-Verlag, Berlin.
- [11] R. Bowen, Equilibrium states and the ergodic theory of Anosov diffeomorphisms, Springer Lectures Notes in Math. (1975).
- [12] R. Bowen, D. Ruelle, The ergodic theory of Axiom A flows, Invent. Math., 29 (1975), 181-202.
- [13] C. Bonatti, M. Viana, SRB measures for partially hyperbolic systems whose central direction is mostly contracting, Israel J. Math., 115 (2000), 157-193.
- [14] M. Brin, S. Garrett, Introduction to dynamical systems, Cambridge university press, 2002.

 ${}^8G^u(PH_{EC}^1(M)) = \{(f,\mu) : \mu \in G^u(f), f \in PH_{EC}^1(M)\}.$

- [15] K. Burns, D. Dolgopyat, Y. Pesin, Pollicott M., Stable ergodicity for partially hyperbolic attractors with negative central exponents, J. Mod. Dyn., 2 (2008), 63-81.
- [16] J. Buzzi, S. Crovisier, O. Sarig. Measures of maximal entropy for surface diffeomorphisms, [arXiv:1811.02240.](http://arxiv.org/abs/1811.02240)
- [17] Y. Cao and D. Yang, On Pesins entropy formula for dominated splittings without mixed behavior, *J. Differential Equations.*, **261**(2016), 39643986.
- [18] Y. Cao, Z. Mi and D. Yang, On the abundance of SRB measures, [arXiv:1804.03328.](http://arxiv.org/abs/1804.03328)
- [19] Y. Cao, G. Liao and Z. You, Upper bounds on measure theoretic tail entropy for dominated splittings, Ergod Th.& Dynam. Sys., DOI: 10.1017/etds.2019.10.
- [20] A. A. Castro, Backward inducing and exponential decay of correlations for partially hyperbolic attractors with mostly contracting central direction, Israel J. *Math.*, **130** (2002), 29-75.
- [21] A. A. Castro, Fast mixing for attractors with a mostly contracting central direction, Ergod Th.& Dynam. Sys., 24 (2004),1744.
- [22] E. Catsigeras and H. Enrich, SRB-like measures for C^0 dynamics, *Bull. Pol.* Acad. Sci. Math., 59 (2011), 151-164.
- [23] E. Catsigeras, M. Cerminara and H. Enrich, The Pesin entropy formula for C1 diffeomorphisms with dominated splitting, Ergod Th.& Dynam. Sys., 35 (2015), 737-761.
- [24] C. Cheng, S. Gan and Y. Shi, A robustly transitive diffeomorphism of Kan's type, Discrete Contin. Dyn. Syst., 38(2) (2018), 867-888.
- [25] W. Cowieson , L.-S. Young, SRB measures as zero-noise limits, Ergod. Th. & Dynam. Sys., 25(4) (2005), 1115-1138.
- [26] S. Crovisier, D. Yang, J. Zhang, Empirical measures of partially hyperbolic attractors, Comm. Math. Phys., (2020). https://doi.org/10.1007/s00220-019- 03668-1
- [27] D. Dolgopyat, On dynamics of mostly contracting diffeomorphisms, Comm. Math. Phys., 213 (2000), 181-201.
- [28] D. Dolgopyat, M. Viana and J. Yang, Geometric and measure-theoretical structures of maps with mostly contracting center, Comm. Math. Phys., 341 2016, 991-1014.
- [29] J.-P. Eckmann and D. Ruelle, Ergodic theory of chaos and strange attractors, Rev. Modern Phys., 57 (1985), 617-656.
- [30] S. Gan. A generalized shadowing lemma, Discrete Contin. Dyn. Syst., 8(3) (2002), 627-632.
- [31] Rodriguez Hertz, M. A. Rodriguez Hertz, A. Tahzibi, and R. Ures. New criteria for ergodicity and nonuniform hyperbolicity, Duke Math. J., 160 (2011), 599- 629.
- [32] M. W. Hirsch, C. Pugh and M. Shub, Invariant manifolds, Lecture Notes in Mathmatics, Springer-Verlag, Vol. 583 (1977).
- [33] H. Hu, Y. Hua and W. Wu, Unstable Entropies and Variational Principle for Partially Hyperbolic Diffeomorphisms, Adv. Math., 321 (2017), 31-68.
- [34] Y. Hua, F. Yang and J. Yang, A new criterion of physical measures for partially hyperbolic diffeomorphisms, *Trans. Amer. Math. Soc.*, **373(1)** (2020), 385-417.
- [35] I. Kan, Open sets of diffeomorphisms having two attractors, each with an everywhere dense basin, *Bull. Am. Math. Soc.*, $31(1)$ (1994), 68-74.
- [36] F. Ledrappier and J. Strelcyn, A proof of the estimation from below in Pesin's entropy formula. Ergod. Th. & Dynam. Sys., 2 (1982), 203-219.
- [37] F. Ledrappier. Propriétés ergodiques des mesures de Sinaï, Publ. Math. I.H.E.S., 59 (1984), 163-188.
- [38] F. Ledrappier, L.-S. Young, The metric entropy of difeomorphisms Part I: Characterization of measures satisfying Pesin's entropy formula, Ann. Math., 122 (1985), 509-539.
- [39] S. Liao. On (η, d) -contractible orbits of vector fields, *Systems Science and Math*ematica Sciences, 2 (1989), 193-227.
- [40] Z. Mi, Y. Cao and D. Yang, A note on partially hyperbolic systems with mostly expanding centers, *Proc. Amer. Math. Soc.*, $145(12)$ (2017), 5299-5313.
- [41] Z. Mi, Radnom entropy expansiveness for diffeomorphism with dominated splittings, *Stoch. Dynam*, $20(1)$ (2020), 2050014 (25 pages).
- [42] P. Mehdipour and A. Tahzibi, SRB measures and homoclinic relation for endomorphisms, *J. Statist. Phys.*, **163(1)** (2016), 139-155.
- [43] A. Okunev, Milnor Attractors of Skew Products with the Fiber a Circle, J. Dyn. Control. Syst., 23 (2017), 421-433.
- [44] J. Palis, A global view of Dynamics and a conjecture on the denseness of finitude of attractors, $Astérisque$, 261 (1999), 339-351.
- [45] V. Pliss, On a conjecture due to Smale, Diff. Uravenenija, 8 (1972), 262-268.
- [46] Y. Pesin, Y. Sinai, Gibbs measures for partially hyperbolic attractors, Ergod. Th. & Dynam. Sys., 2(1982), 417-438.
- [47] D. Ruelle, An inequality of the entropy of differentiable maps, Bol. Sc. Bra. Math., 9 (1978), 83-87.
- [48] D. Ruelle, A measure associated with Axiom A attractors, Amer. J. Math., 98(1976), 619-654.
- [49] D. Ruelle, Differentiation of SRB states, Comm. Math. Phys.,187 (1997), 227- 241.
- [50] Y. Sinai, Gibbs measures in ergodic theory, Russ. Math., 27(4) (1972), 21-69.
- [51] R. Ures, M.Viana and J.Yang, Maximal entropy measures of diffeomorphisms of circle fiber bundles, [arXiv:1909.00219.](http://arxiv.org/abs/1909.00219)
- [52] C. Vásquez, Statistical stability for diffeomorphisms with dominated splitting, Ergod Th.& Dynam. Sys., 27 (2007), 253-283.
- [53] M. Viana and J. Yang, Physical measures and absolute continuity for onedimensional center direction, Ann. Inst. H. Poincaré Anal. Non Linéaire, 30(2013), 845-877.
- [54] L.-S. Young, What are SRB measures, and which dynamical systems have them?, J. Statist. Phys., 108(2002), 733-754.
- [55] J. Yang, Entropy along expanding foliations, [arXiv:1601.05504v](http://arxiv.org/abs/1601.05504)1.
- [56] J.Yang, Geometrical and measure-theoretic structures of maps with mostly expanding center, [arXiv:1904.10880v](http://arxiv.org/abs/1904.10880)1.
- [57] P. Walters, An introduction to ergodic theory, Springer Verlag, 1982.

School of Mathematics and Statistics, Nanjing University of Information Science and Technology, Nanjing 210044, Jiangsu, P.R. China. E-mail address: mizeya@163.com

Department of Mathematics, Shanghai key laboratory of PMMP, East China Normal University, Shanghai 200062, P.R. China.

Department of Mathematics, Center for Dynamical Systems and Differential Equations, Soochow University, Suzhou 215006, Jiangsu, P.R. China.

E-mail address: ylcao@suda.edu.cn