Cell cycle heritability and localization phase transition in growing populations

Takashi Nozoe¹ and Edo Kussell^{1,2}
¹Department of Biology, New York University, 12 Waverly Place, New York, NY 10003.
²Department of Physics, New York University, 726 Broadway, New York, NY 10003.

(Dated: March 12, 2020)

The cell cycle duration is a variable cellular phenotype that underlies long-term population growth and age structures. By analyzing the stationary solutions of a branching process with heritable cell division times, we demonstrate existence of a phase transition, which can be continuous or first-order, by which a non-zero fraction of the population becomes localized at a minimal division time. Just below the transition, we demonstrate coexistence of localized and delocalized age-structure phases, and power law decay of correlation functions. Above it, we observe self-synchronization of cell cycles, collective divisions, and slow 'aging' of population growth rates.

The duration of a cell-cycle, or inter-division time (IDT), is a fluctuating quantity in cellular populations, and its statistical properties are thought to result from biological mechanisms that regulate cell growth and division [1–3]. For single cells that grow in size exponentially, as in bacteria [1] and yeast [4], models of cell-size regulation typically predict negative mother-daughter correlation of IDTs [5]. Yet in a subset of bacterial experiments and in most observations on mammalian cells. positive IDT correlations have been measured (see Table S1 in [6]), indicating that the cell cycle duration is a heritable, fluctuating single-cell phenotype. Such heritability enables selection to act on the distribution of IDTs in a population to increase the long-term population growth rate. Here we take up the question of how selection and cell cycle heritability interact to determine long-term population dynamics, a fundamental step toward understanding how evolution has shaped cell cycle control mechanisms.

To investigate the effect of IDT heritability on population growth, we considered a model with heritable IDTs that was first introduced in [7], which we refer to as the Lebowitz-Rubinow model below. We noticed the existence of a heritability threshold above which the population's distribution of IDTs would localize at the minimal IDT corresponding to the fastest possible single-cell growth rate. We identify and characterize a localization phase transition in this model. We show the ancestral mean IDT provides an order parameter of the transition, and above the heritability threshold predict the emergence of arbitrarily long single cell lineages that maintain perfect inheritance of a minimal IDT. This prediction is confirmed in numerical simulations of finite populations. We observe slow punctuated dynamics above the heritability threshold, characteristic of systems that exhibit aging, as well as cell-cycle synchronization in finite populations. Our work provides fundamental connections between dynamics of age-structured populations [8] and well-studied error-threshold phenomena of evolutionary theory [9, 10], both of which are described by the unifying framework of statistical physics of phase transitions.

Model and stationary solutions. We model a proliferating population by a branching process in which $K(\tau,\tau')$ is the transition probability density from τ' to τ , where τ' and τ are the parent and offspring IDTs, respectively. To study the effect of heritability of IDTs, we focus on the analytically tractable Lebowitz-Rubinow model [7], which uses the transition kernel

$$K(\tau, \tau') = \beta \delta(\tau - \tau') + (1 - \beta) k(\tau) , \qquad (1)$$

where $\delta\left(\cdot\right)$ is the Dirac delta function, β represents the heritability of IDTs in the model $(0 \leq \beta < 1)$, and $k\left(\tau\right)$ is a probability density function on the interval $[\tau_0, \infty)$, with $0 < \tau_0 < \infty$. A cell produces \hat{z} offsprings at every division, where \hat{z} is independently drawn from a fixed probability distribution and its average is denoted by z > 0. For example, z = 2 for binary division organisms and z = 1 for the isolated single cells.

The dynamics of cell divisions in the population are governed by

$$n_{\rm div}(\tau;t+\tau) = z \int_0^\infty K(\tau,\tau') n_{\rm div}(\tau';t) d\tau'.$$
 (2)

where $n_{\rm div}\left(\tau;t\right)d\tau dt$ is the expected number of dividing cells (cells at the termination of cell-cycle) between times t and t+dt with IDT between τ and $\tau+d\tau$ [11]. The expected number of divisions occurring between t and t+dt is given by $N_{\rm div}\left(t\right)dt:=dt\int_0^\infty n_{\rm div}\left(\tau;t\right)d\tau$. The time-dependent population growth rate is defined by $\Lambda_t:=(z-1)\,N_{\rm div}\left(t\right)/N\left(t\right)$, where $N\left(t\right)$ is the expected population size at time t (see [6] for detailed derivations).

The steady-state, exponentially growing solution of (2) can be found by substituting the form $n_{\rm div}(\tau;t) = N_{\rm div}(t) \cdot p_{\rm div}(\tau)$, where $p_{\rm div}(\tau)$ is the stationary probability density of IDTs of dividing cells, yielding

$$p_{\text{div}}(\tau) = ze^{-\Lambda\tau} \int_0^\infty K(\tau, \tau') p_{\text{div}}(\tau') d\tau' , \qquad (3)$$

Using (1) above, and solving the equation one obtains

$$p_{\text{div}}(\tau) = q_{\beta,z,\Lambda}(\tau) := \frac{(1-\beta)k(\tau)ze^{-\Lambda\tau}}{1-\beta ze^{-\Lambda\tau}}$$
(4)

where the steady-state growth rate Λ is determined by the normalization condition

$$Q(\beta, z, \Lambda) := \int_0^\infty q_{\beta, z, \Lambda}(\tau) d\tau = 1.$$
 (5)

If z=1, for example, then $\Lambda=0$ and $p_{\rm div}\left(\tau\right)=k\left(\tau\right)$ for any $0\leq\beta<1$. Namely, the stationary IDT distribution of isolated single cells is invariant of β . We remark that β is the correlation coefficient of IDTs between parent and offspring for isolated, single cell growth at steady-state [6].

In the absence of IDT correlations $(\beta = 0)$, one recovers the well-known result $p_{\text{div}}(\tau) = ze^{-\Lambda\tau}k(\tau)$ where Λ is the unique real root of the integral equation $z\int_0^\infty e^{-\Lambda\tau}k(\tau)\,d\tau = 1$ [3, 12]. For $\beta > 0$, one additionally must have $1 - \beta ze^{-\Lambda\tau} > 0$ for all τ in the support of $k(\tau)$ to ensure $q_{\beta,z,\Lambda}(\tau) \geq 0$. If $k(\tau) > 0$ for all $\tau > \tau_0$, we find

$$\Lambda \ge \omega_0(\beta, z) := \sup_{\tau > \tau_0} \frac{\ln(\beta z)}{\tau} = \max\left(0, \frac{\ln(\beta z)}{\tau_0}\right); \quad (6)$$

(see [6] for $k(\tau)$ with bounded support). Since $Q(\beta, z, \Lambda)$ is a monotonically decreasing function of Λ tending to zero as $\Lambda \to \infty$, Eq. 5 has a unique root $\Lambda_{\beta,z}$ provided that $Q(\beta, z, \omega_0(\beta, z)) \geq 1$. This condition holds for $\beta < \beta_c$, where β_c is a critical heritability threshold defined by

$$Q(\beta_c, z, \omega_0(\beta_c, z)) = 1, \qquad (7)$$

such that for $\beta > \beta_c$, $Q(\beta, z, \omega_0(\beta, z)) < 1$ and Eq. 5 does not admit a real solution $\Lambda_{\beta,z}$. For $\beta > \beta_c$, the solution $p_{\text{div}}(\tau)$ given in (4) is incomplete, as there is missing probability 1 - Q. For z > 1, the full solution is

$$p_{\text{div}}(\tau) = q_{\beta,z,\Lambda_{\beta,z}}(\tau) + (1 - Q(\beta, z, \Lambda_{\beta,z})) \delta(\tau - \tau_0),$$
(8)

and substitution into (3) yields $\Lambda_{\beta,z} = \tau_0^{-1} \ln(\beta z)$ for the steady-state growth rate when $\beta > \beta_c$ [6].

Further analysis of (7) shows that a heritability threshold $\beta_c < 1$ exists if and only if $\int_0^\infty d\tau k(\tau)/(\tau - \tau_0)$ converges; e.g. if $k(\tau) \sim (\tau - \tau_0)^{\gamma}$ near τ_0 for some $\gamma > 0$. The steady state population growth rate $\Lambda_{\beta,z}$ qualitatively changes as β crosses the threshold: it depends on $k(\tau)$ for $\beta < \beta_c$, and becomes independent of it for $\beta > \beta_c$. The expected number of offspring having the same IDT as their parent is βz , and only parents with $\tau' = \tau_0$ can generate offspring with $\tau = \tau_0$. Thus, the fraction of the population with IDT τ_0 grows with rate $\tau_0^{-1} \ln{(\beta z)}$. If we add a small fraction of τ_0 cells to a population, they go extinct if $\beta z < 1$, while if $\beta z > 1$, they can constitute a giant cluster in the population's genealogy. The subpopulation localized at $\tau = \tau_0$ will be outcompeted by the rest of the population for $z^{-1} < \beta < \beta_c$, with $\Lambda_{\beta,z}$ determined by (5); or it will dominate the population for $\beta > \beta_c$, and thus dictate its growth rate to be $\Lambda_{\beta,z} = \tau_0^{-1} \ln{(\beta z)}$. In Fig. 1A, we show a range of




FIG. 1. Stationary solutions and their dependence on the heritability parameter β . (A) The IDT distribution, $k\left(\tau\right)$, is chosen to be a gamma distribution, shifted by $\tau_{0}=0.2$, with shape parameters $\alpha=1.5$, 4, and 20; $k\left(\tau\right)=\Gamma\left(\alpha\right)^{-1}\theta^{-\alpha}\left(\tau-\tau_{0}\right)^{\alpha-1}e^{-\left(\tau-\tau_{0}\right)/\theta}$ for $\tau\geq\tau_{0}$. For the mean IDT to be 1, the scale parameter θ is chosen as $\tau_{0}+\alpha\theta=1$. (B-D) Dependence of key quantities of the stationary solutions on β using z=2; all other parameters are the same as in A. $\beta_{c}=0.68$, 0.60 and 0.58 respectively for $\alpha=1.5$, 4, and 20. (B) Population growth rate, $\Lambda_{\beta,z}$. (C) Reciprocal mean block size, $\overline{m}_{\beta,z}^{-1}$. (D) Reciprocal ancestral mean IDT, $\overline{\tau}_{\beta,z}^{-1}$.

examples $k(\tau)$ which admit a threshold β_c . Increasing β from 0 to 1, $\Lambda_{\beta,z}$ increases monotonically with a shallow slope, while past β_c , the slope of the growth rate changes markedly (Fig. 1B).

Localization phase transition of population age structure. We now show that the threshold behavior identified above constitutes a phase transition in the strict sense. We map the population to a statistical mechanical ensemble, as follows. From the viewpoint of single cell lineages – i.e. the history of an individual and all of its ancestors – an age-structured population constitutes an ensemble of trajectories: the sequence of IDTs along a lineage $(\ldots, \tau_{i-1}, \tau_i, \tau_{i+1}, \ldots)$ is analogous to a microscopic state of a large system (e.g. configuration of spins on a lattice, conformation of a polymer in space, etc.); while a single ancestral cell division τ_i specifies the state of a single component (e.g a spin, or a monomer). The population is an ensemble of lineages, and the steady-state population growth rate is its free energy [6, 13].

To analyze the structure of lineages observed above and below β_c , we consider the number of generations with which the same IDT is consecutively inherited, which we denote by m and call the "block size". The probability distribution of m over lineages is analogous to a correlation function, and its mean measures the typical correlation length. To compute these quantities, we let (τ', τ) denote a pair of IDTs, where τ' and τ are parent and offspring IDTs, respectively. From (3) one can infer

that the stationary probability density to find (τ',τ) is $ze^{-\Lambda\tau}K(\tau,\tau')\,p_{\rm div}(\tau')$. In particular, the probability of observing (τ,τ) is $\beta ze^{-\Lambda\tau}p_{\rm div}(\tau)$. Repeatedly multiplying $\beta ze^{-\Lambda_{\beta,z}\tau}$, the probability for τ drawn from $p_{\rm div}(\tau)$ to be inherited at least m-1 times is $(\beta ze^{-\Lambda_{\beta,z}\tau})^{m-1}$, hence the joint probability distribution of block size m and IDT τ is

$$p_{\text{lin}}(m,\tau) := \left(1 - \beta z e^{-\Lambda_{\beta,z}\tau}\right) \left(\beta z e^{-\Lambda_{\beta,z}\tau}\right)^{m-1} p_{\text{div}}(\tau) . \tag{9}$$

For $\beta < \beta_c$, the mean block size on lineage is

$$\overline{m}_{\beta,z} := \sum_{m>1} \int_0^\infty m \cdot p_{\text{lin}}(m,\tau) \, d\tau < \infty , \qquad (10)$$

and the probability distribution of IDT on lineage, also known as the ancestral distribution [10], is

$$p_{\text{lin}}(\tau) := \frac{\sum_{m \ge 1} m \cdot p_{\text{lin}}(m, \tau)}{\overline{m}_{\beta, z}}, \qquad (11)$$

and represents the probability distribution of IDTs of ancestors in the infinite past [6]. For $\beta > \beta_c$, $\overline{m}_{\beta,z}$ diverges and $p_{\text{lin}}(\tau)$ must be computed as an appropriate limit [6]; we obtain

$$p_{\text{lin}}(\tau) = \begin{cases} \frac{(1-\beta)k(\tau)ze^{-\Lambda_{\beta,z}\tau}}{\overline{m}_{\beta,z}(1-\beta ze^{-\tau\Lambda_{\beta,z}})^2}, & 0 \le \beta < \beta_c\\ \delta(\tau - \tau_0), & \beta_c < \beta < 1 \end{cases} . (12)$$

The expression indicates that the IDT distribution on lineages in the population is localized entirely at $\tau = \tau_0$ above β_c , despite the fact that the IDT distribution of isolated lineages remains $k(\tau)$.

The mean block size $\overline{m}_{\beta,z}$ serves as an order parameter that diverges for $\beta > \beta_c$, and the continuity of the phase transition can be characterized by its behavior near β_c . If $\lim_{\beta \uparrow \beta_c} \overline{m}_{\beta,z} = \infty$, the transition is a continuous phase transition, while if $\lim_{\beta \uparrow \beta_c} \overline{m}_{\beta,z} < \infty$ the transition is referred to as discontinuous, or 'first-order'. Analogously, the lineage mean IDT, $\overline{\tau}_{\beta,z} := \int_0^\infty \tau p_{\text{lin}}(\tau) d\tau$, which is given by $\partial \Lambda_{\beta,z}/\partial \log z$ (i.e. a first-order derivative of the free energy [6]), exhibits the same type of phase transition. Examples of the β dependence of $\overline{m}_{\beta,z}$ and $\overline{\tau}_{\beta,z}$ are shown in Fig. 1C and D. Assuming the law of large numbers, $\overline{\tau}_{\beta,z}^{-1} \simeq D/t$ and $\overline{m}_{\beta,z}^{-1} \simeq S/D$ hold where D and S denote the number of divisions and the number of switches to different values of τ on lineage, respectively [6].

Using the form of $k(\tau)$ as in Fig. 1, where $k(\tau) \sim (\tau - \tau_0)^{\gamma}$ for τ near τ_0 , and computing $\overline{m}_{\beta,z}$ we find that the phase transition is continuous for $0 < \gamma \le 1$, and becomes discontinuous for $\gamma > 1$. For the latter type of transition, one expects to observe coexistence of two phases at the transition point, which is seen in numerical simulations of finite populations shown below (Fig. 2B). We can also compute the marginal probability distribution $p_{\text{lin}}^{\text{M}}(m)$ of block size m, which decays exponentially




FIG. 2. Aging of growth rates in a finite population. The parameters are as in Fig. 1 using $\alpha = 4$, which exhibits a first-order localization transition. In simulations, some of the cells in the population may be removed when newborn cells appear in order to keep the population size N = 100 constant. See [6] for simulation details. $\beta_{c,N} = 0.73$. (A-D) Time courses of reciprocal empirical minimal IDT $\hat{\tau}_0^{-1}$ (top, orange, solid) compared with the minimal IDT τ_0^{-1} (gray, dashed line) and growth rates (bottom, blue, solid) compared with predicted stationary value (gray, dashed line). (A) is below the transition ($\beta = 0.5$) and the initial state is sampled from the delocalized state. (B) shows coexistence of delocalized and localized states ($\beta = 0.7$, starting from the delocalized state). Time courses of growth rates are compared to $\hat{\tau}_0^{-1} \log (\beta z)$ (orange dot). (C) Time courses above the transition showing aging dynamics ($\beta = 0.8$, starting from the delocalized state). (D) For $\beta = 0.8$, when the initial population is sampled from predicted stationary solution, the growth rates fluctuate around the theoretically predicted stationary growth rate. (E, F) Growth rate vs β . The initial condition was chosen as delocalized state for (E) and predicted stationary for (F). For each β and stationary conditions, 10 simulations were run, and the mean is plotted with error bars showing the standard deviation.

below the transition, and follows power law statistics, $p_{\text{lin}}^{\text{M}}(m) \sim m^{-\gamma-1}$ for large m, in the limit $\beta \uparrow \beta_c$ as expected for correlation functions in the vicinity of a phase transition [6]. Such an appearance of long memory of IDT inheritance at the transition point is likewise observed in simulations (Fig. 2B).

Aging of long-term growth rate in finite populations. In finite sized populations, the stationary distribution (8) for $\beta > \beta_c$ is not achievable because any parent cell with IDT $\tau' > \tau_0$ has probability zero of generating offspring with IDT τ_0 . Additionally, the distribution (8) cannot be

maintained as a steady-state because cells with IDT τ_0 can be lost from the population within finite time due to coalescence. To observe dynamics in finite populations, we conducted exact stochastic simulations in which cells are randomly removed to maintain a fixed population size [6]. We sampled the initial population with size N=100 independently from the stationary probability distribution with $\beta=0$, which we refer to as the delocalized state, and simulated the population forward in time for a given value of $\beta>0$.

In simulations with $\beta < \beta_c$, growth rates Λ_t fluctuate around the expected steady-state growth rate (Fig. 2A). For β slightly above β_c , however, Λ_t exhibits sudden transitions between two distinct, long-lived states (Fig 2B), which is expected for systems near a first-order phase transition. One of these states represents localization at the empirical minimum IDT, denoted by $\hat{\tau}_0$, which is the minimum IDT among all the dividing cells within each time bin [6]. In this state, the growth rate fluctuates around $\hat{\tau}_0^{-1} \ln (\beta z)$ over a sufficient period during which $\hat{\tau}_0$ is constant. The other phase represents delocalization, where $\hat{\tau}_0$ exhibits large fluctuations. For higher values of β , the observed growth rate increases stepwise and fluctuates around $\hat{\tau}_0^{-1} \ln (\beta z)$ (Fig. 2C). In this case, a fraction of the population localized at $\hat{\tau}_0$ is maintained over a significant time interval until a new value of $\hat{\tau}_0$ replaces the current empirical minimum IDT. The time intervals between these replacement events become increasingly long as $\hat{\tau}_0$ approaches τ_0 , because IDTs that are shorter than the current minimum become increasingly rare. When the simulation starts with the population sampled from the predicted stationary probability distribution with the same β used in Fig. 2C, Λ_t fluctuates around $\hat{\tau}_0^{-1} \ln (\beta z)$ and $\hat{\tau}_0 = \tau_0$ is preserved over the entire simulation time (Fig. 2D). Values of Λ_t at different simulation times are plotted as function of β initialized at either the delocalized state (Fig. 2E) or the predicted stationary solution (Fig. 2F). For both initial conditions, the curve at $t=10^4$ possesses an inflection point $\beta_{c,N}$ slightly greater than β_c , which we refer to as the effective transition point at fixed population size N (see [6] for detailed definition). For both of these initial conditions, the population would reach the steady-state where $\hat{\tau}_0$ fluctuates around some specific $\tau_{\rm opt}$ (> τ_0) but it was not observed in reasonable simulation time due to the observed aging behavior of the dynamics. Despite this challenge in numerically observing the true steady-state, due to the slow aging of the dynamics, our finite population simulations demonstrate the existence of the transition point above which the localized phase emerges in the dynamics, and that the dependence of the observed growth rate on β is quantitatively predicted by theoretical analysis of the stationary solution.

Collective divisions and cell cycle synchronization. In addition to aging dynamics, we observed collective divisions within the population above the transition (Fig.

3). The number of divisions that occurred, binned in short intervals, is plotted over time. Compared to Figs. 3A and C, Fig. 3E shows division events occurring collectively and periodically. Its auto-correlation function does not oscillate below the transition (3B), but exhibits decaying oscillations near the transition point (3D) and sustained oscillations above the transition (3F). The fact that the period of the autocorrelation function is close to τ_0 reflects localization at $\hat{\tau}_0$ close to τ_0 . We also tested the robustness of the transition properties to noise in the inheritance of IDTs by allowing small fluctuations of the offspring's IDT when it inherits its parent's IDT with probability β . To do so, we modified the transition kernel to be

$$K(\tau, \tau') = \beta p_{\text{norm}}(\tau - \tau', \sigma) + (1 - \beta) k(\tau)$$
 (13)

where $p_{\text{norm}}(x,\sigma)$ is a normal distribution density function with standard deviation σ , truncated at τ_0 . As a result, the population can reach equilibrium over a reasonable time scale, that is, the long-term behaviors coincide between the two distinct initial conditions (Figs. S6 and S7 in [6]). Even without exact inheritance of parental IDTs, signatures of the phase transition are still observed (Figs. S8-S15 in [6]). Collective divisions are weaker but still detectable through the decaying oscillations of the autocorrelation function of divisions in population (Figs. S16 and S17 in [6]). Division rate oscillations have also been predicted to arise in cell size control models, due to negative IDT correlations [14].

Discussion. We analyzed how the strength of the inter-division time heritability affects age-structured population dynamics. In a model with heritable cell cycle durations first introduced in [7], we demonstrated the existence of a localization phase transition. While the existence of a heritability threshold β_c was not explicitly suggested in [7], the potential limit on the validity of the stationary solution was pointed out. A similar localization transition in exponentially growing populations is known in mutation-selection models of theoretical population genetics [10, 15, 16]. For example, Kingman's house-of-cards model [16] describes population dynamics on a specific type of fitness landscape, which exhibits localization at maximal fitness below a critical mutation rate. The localization phase transition as a stationary state in the house-of-cards model can be generalized to include cases where fitness is correlated between parent and offspring, and the existence and the uniqueness of the stationary solutions has been proven using the theory of positive linear operators [17].

We showed that the signature of localization phase transition can be observed even in finite populations as small as 100 cells, which is comparable to the capacity allowed by typical microfluidic experiments [3, 18, 19], indicating that this transition could in principle be experimentally observed. Above the transition, population




FIG. 3. Synchronization of cell-cycles above the transition. The simulation is initialized by the delocalized state. The parameters are as in Fig. 2 using $\alpha = 4$ and N = 100. $\beta_{c,N} = 0.73$. Number of divisions binned over 0.1 doubling time (A, C, E) and their auto-correlations (B, D, F) are shown. Simulations are run over 10⁴ time units and the last 10 time units of the series are shown for A, C and E, where time point 0 indicates the simulation end. Auto-correlations are computed over entire time series. The correlation coefficient is normalized to equal 1 at zero lag. Dotted line indicates zero correlation. (A,B) Below the transition ($\beta = 0.6$) timing of divisions are not synchronized. (C, D) Near the transition point ($\beta = 0.7$), timing of divisions are partially synchronized, observed as decaying oscillation of the auto-correlation function. (E, F) Above the transition ($\beta > \beta_{c,N}$), collective divisions are observed indicating self-synchronization of cell cycles.

growth rates exhibit aging dynamics. Similar phenomena are seen in evolution in random unbounded fitness landscape with rare mutation rate in finite population, known as the "diluted record process" [20] or "successional mutation regime" [21]. While the distribution and auto-correlation of individual fitness are difficult to measure in reality, measuring those of IDTs is highly feasible using in vivo single cell tracking in microfluidics-based experiments. The aging dynamics we observed in agestructured populations are closely related to the emergence of self-synchronized cell-cycles (Fig. 3). Future work on the spectral structure of the linear semigroup of age-structured population dynamics [22, 23] may further elucidate the emergence of synchronized growth above the transition.

While aging dynamics above the transition are specific to the noiseless inheritance of IDT, we found that the signature of localization and self-synchronization can still be observable in "noisier" inheritance systems. This result indicates that our analytical results are applicable to real biological systems in which IDT heritability is noisy. The Lebowitz-Rubinow model does not include cell-cell

interactions which are thought to be a major mechanism for cell-cycle synchronization. Remarkably, our analysis predicts strong but imperfect correlation may be enough to cause self-synchronization of cell cycles in finite populations in the absence of cell-cell interactions. Our findings could provide a basis for the design of new types of synthetic biological oscillators which leverage population-level selective forces to establish robust cell cycle synchronization and to support sustained oscillations.

We thank Yuichi Wakamoto for discussions. This work was supported by NIH grant R01-120231 to E.K.

- S. Taheri-Araghi, S. Bradde, J. T. Sauls, N. S. Hill, P. A. Levin, J. Paulsson, M. Vergassola, and S. Jun, Curr. Biol. 25, 385 (2015).
- [2] L. Susman, M. Kohram, H. Vashistha, J. T. Nechleba, H. Salman, and N. Brenner, Proc. Natl. Acad. Sci. U. S. A. 115, 201615526 (2018), arXiv:1805.05058.
- [3] M. Hashimoto, T. Nozoe, H. Nakaoka, R. Okura, S. Akiyoshi, K. Kaneko, E. Kussell, and Y. Wakamoto, Proc. Natl. Acad. Sci. 113, 3251 (2016).
- [4] H. Nakaoka and Y. Wakamoto, PLoS Biol. 15, e2001109 (2017).
- [5] A. Amir, Phys. Rev. Lett. 112, 208102 (2014).
- [6] Supplemental Material at [URL will be inserted by publisher].
- [7] J. L. Lebowitz and S. I. Rubinow, J. Math. Biol. 1, 17 (1974).
- [8] B. Charlesworth, Evolution in Age-Structured Populations, 2nd ed., Cambridge Studies in Mathematical Biology (Cambridge University Press, 1994).
- [9] M. Eigen, J. McCaskill, and P. Schuster, Adv. Chem. Phys 75, 149 (1989).
- [10] J. Hermisson, O. Redner, H. Wagner, and E. Baake, Theor. Popul. Biol. 62, 9 (2002).
- [11] IDTs of sibling cells may be correlated, which does not change Eq. (2), as long as the number of offspring is independent of the parent's IDT.
- [12] R. Fisher, The Genetical Theory of Natural Selection (Clarendon, 1930).
- [13] Y. Wakamoto, A. Y. Grosberg, and E. Kussell, Evolution (N. Y). 66, 115 (2012).
- [14] F. Jafarpour, Phys. Rev. Lett. 122, 118101 (2019), arXiv:1809.10217.
- [15] M. Eigen, Naturwissenschaften **58**, 465 (1971).
- [16] J. F. C. Kingman, J. Appl. Probab. **15**, 1 (1978).
- [17] R. Bürger and I. M. Bomze, Adv. Appl. Probab. **28**, 227 (1996).
- [18] M. Wallden and J. Elf, Curr. Opin. Biotechnol. **22**, 81 (2011).
- [19] G. Lambert and E. Kussell, PLoS Genet. **10**, e1004556 (2014).
- [20] S. C. Park and J. Krug, J. Stat. Mech. Theory Exp. 2008, 10.1088/1742-5468/2008/04/P04014 (2008), arXiv:0711.1989.
- [21] T. Brotto, G. Bunin, and J. Kurchan, J. Stat. Mech. Theory Exp. 2016, 10.1088/1742-5468/2016/03/033302 (2016), arXiv:1407.4669.

- [22] G. F. Webb, J. Math. Biol. 23, 269 (1986).[23] M. Boulanouar, J. Evol. Equations 11, 531 (2011).

Supplemental Material: Cell cycle heritability and localization phase transition in growing populations

Takashi Nozoe¹ and Edo Kussell^{1,2}

¹Department of Biology, New York University, 12 Waverly Place, New York, NY 10003.

²Department of Physics, New York University, 726 Broadway, New York, NY 10003.

(Dated: March 12, 2020)

CONTENTS			Classification heritability threshold of LR model	16
List of Tables	1			
List of Figures	1		LIST OF FIGURES	
I.General stationary solutions	2	S1	Synchronization of cell-cycles without inheritance noise	17
$I\beta$ as correlation coefficient	3	S2 S3	Finite N effects, initial delocalized state	
Intationary solution in the Lebowitz-Rubinow model Alteritability threshold	3 3		Phase transition at various N , initial delocalized state	18
Case II: $z > 1$ Case II: $z = 1$	4 5	S4	Finite N effects, initial predicted stationary state	18
Case III: $0 < z < 1$ Necessary and sufficient condition for	5	S5	Phase transition at various N , initial predicted stationary state	19
$Q_{0}\left(eta,z ight) <\infty$	6	S6		19
Estationary solution for $\beta > \beta_c$ CSmall variation limit of IDT distribution	6 7	S7	Disappearing aging dynamics with 10% inheritance noise	20
Independent divisions: $\beta = 0$	7	S8	Finite N effects with 1% inheritance noise, initial delocalized state	20
Intatistics of Lineages Allean block size for $\beta > \beta_c$ in finite time	8	S9	Phase transition at various N with 1% inheritance noise, initial delocalized state	21
BLineage IDT distribution for $\beta > \beta_c$ COn-lineage mean IDT	9	S10	Finite N effects with 1% inheritance noise, initial predicted stationary state	21
DBlock size distribution	9	S11	Phase transition at various N with 1% inheritance noise, initial predicted station-	
VAncestral distribution	9	S12	ary state	22
VIDerivatives of growth rate function	10		noise, initial delocalized state Phase transition at various N with 10%	22
VThermodynamic analogy for lineage structures in the Lebowitz-Rubinow model	10			23
Wilmerical simulations	12			23
AIDT distribution in isolation BGrowth rate	12 12	510	inheritance noise, initial predicted stationary state	24
Cinitial conditions Dimulation	12 13	S16	Synchronization of cell-cycles with 1% in-	24
IXX-dependence of population growth and definition of		S17	Synchronization of cell-cycles with 10% in-	
$eta_{c,N}$	13		heritance noise	20
References	13			

LIST OF TABLES

S1	List of mother-daughter correlation coeffi-	
	cients of IDTs	15

I. GENERAL STATIONARY SOLUTIONS

Let $n(a, \tau; t) da d\tau$ be the expected number of cells with age a and IDT τ at time t. a is the time since division, which is reset to 0 just after division $(0 \le a < \tau)$. In particular, $n_{\text{div}}(\tau; t) d\tau dt := \lim_{a \to \tau} n(a, \tau; t) d\tau dt$ is the expected number of dividing cells at age τ during the infinitesimal interval [t, t + dt]. For fixed τ , $n(a, \tau; t)$ satisfies the partial differential equation:

$$\frac{\partial n\left(a,\tau;t\right)}{\partial a} + \frac{\partial n\left(a,\tau;t\right)}{\partial t} = 0 \tag{S1}$$

We denote the initial distribution of a and τ by $n(a,\tau;0) = n_0(a,\tau)$. Then the solution of Eq. (S1) satisfies

$$n\left(a,\tau;t\right) = \begin{cases} n\left(0,\tau;t-a\right) & t > a \\ n_0\left(a-t,\tau\right) & t \le a \end{cases}$$
 (S2)

Let $K(\tau, \tau')$ denote the transition probability density from τ' to τ . The boundary condition is given by

$$n(0,\tau;t) = z \int_{0}^{\infty} K(\tau,\tau') n_{\text{div}}(\tau';t) d\tau'$$
 (S3)

Let

$$N_t := \int_0^\infty \int_0^\tau n(a, \tau; t) \, da \, d\tau \tag{S4}$$

denote the expected population size at time t (relative to the initial population size) and

$$p(a,\tau;t) := n(a,\tau;t) \cdot N_t^{-1}$$
 (S5)

denote the probability density of (a, τ) at time t. The marginal distribution of age is denoted by

$$p_{\text{age}}(a;t) := \int_{a}^{\infty} p(a,\tau;t) d\tau . \tag{S6}$$

By integration of Eq. (S3), we have

$$\int_{0}^{\infty} n(0,\tau;t) d\tau = z \int_{0}^{\infty} n_{\text{div}}(\tau;t) d\tau \qquad (S7)$$

This equation represents the balance between numbers of newly formed cells and dividing cells. Thus the exponential growth rate at time t, denoted by Λ_t , is

$$\Lambda_{t} := \frac{d}{dt} \ln N_{t} = N_{t}^{-1} \int_{0}^{\infty} \left(\int_{0}^{\tau} \frac{\partial n\left(a, \tau; t\right)}{\partial t} da \right) d\tau$$

$$= -N_{t}^{-1} \int_{0}^{\infty} \left(\int_{0}^{\tau} \frac{\partial n\left(a, \tau; t\right)}{\partial a} da \right) d\tau$$

$$= N_{t}^{-1} \int_{0}^{\infty} \left(n\left(0, \tau; t\right) - n_{\text{div}}\left(\tau; t\right) \right) d\tau$$

$$= \left(1 - z^{-1} \right) p_{\text{age}}\left(0; t\right) \tag{S8}$$

This definition of time-dependent growth rate is equivalent to its definition in Main Text $(\Lambda_t := (z-1) N_{\text{div}}(t)/N(t))$ because $p_{\text{age}}(0;t) = z N_{\text{div}}(t)/N(t)$.

Three interelated distributions of IDTs can be defined, as follows: (i) the offspring's IDT distribution,

$$p_{\text{birth}}(\tau;t) := \frac{p(0,\tau;t)}{p_{\text{age}}(0;t)} = \frac{1-z^{-1}}{\Lambda_t} p(0,\tau;t) ,$$
 (S9)

which is the probability distribution of τ conditioned on a=0; (ii) the parent's IDT distribution,

$$p_{\text{div}}(\tau;t) := \frac{n_{\text{div}}(\tau;t)}{\int_{0}^{\infty} n_{\text{div}}(\tau;t) d\tau} = \frac{z \cdot n_{\text{div}}(\tau;t)}{N_{t} p_{\text{age}}(0;t)}$$

$$= \begin{cases} z \frac{\Lambda_{t-\tau}}{\Lambda_{t}} e^{-\int_{t-\tau}^{t} \Lambda_{s} ds} p_{\text{birth}}(\tau;t-\tau) & \tau < t \\ z \frac{\Lambda_{0}}{\Lambda_{t}} e^{-\int_{0}^{t} \Lambda_{s} ds} p(\tau-t,\tau;0) & \tau \ge t \end{cases},$$
(S10)

which is the probability distribution of age among dividing cells at time t; and (iii) the population's IDT distribution,

$$p_{\mathrm{idt}}\left(\tau;t\right) := \int_{0}^{\tau} p\left(a,\tau;t\right) da , \qquad (S11)$$

i.e. the marginal of $p(a, \tau; t)$ with respect to τ , which is the probability distribution of τ at time t.

The joint distribution of τ and a at time t can be expressed as

$$p\left(a, \tau; t\right) = \begin{cases} \frac{\Lambda_{t} e^{-\int_{t-a}^{t} \Lambda_{s} ds}}{1 - z^{-1}} p_{\text{birth}}\left(\tau; t - a\right) & a < t \\ e^{-\int_{0}^{t} \Lambda_{s} ds} p\left(a - t, \tau; 0\right) & a \ge t \end{cases}$$
(S12)

Stationary distributions are found by substituting $p\left(a,\tau;t\right)=p\left(a,\tau\right)$, which is independent of t. Accordingly, we denote $p_{\rm age}\left(a\right):=\int_a^\infty p\left(a,\tau\right)d\tau$, $p_{\rm idt}\left(\tau\right):=\int_0^\tau p\left(a,\tau\right)da$, $p_{\rm birth}\left(\tau\right):=p\left(0,\tau\right)/p_{\rm age}\left(0\right)$ and $p_{\rm div}\left(\tau\right):=p\left(\tau,\tau\right)/\int_0^\infty p\left(\tau',\tau'\right)d\tau'$. Let $\Lambda:=\left(1-z^{-1}\right)p_{\rm age}\left(0\right)$ denote the steady-state population growth rate. Then we formally obtain

$$p(a,\tau) = e^{-\Lambda a} p(0,\tau)$$

$$= \frac{\Lambda}{1 - z^{-1}} e^{-\Lambda a} p_{\text{birth}}(\tau)$$
 (S13)

$$p_{\text{age}}(a) = \frac{\Lambda}{1 - z^{-1}} e^{-\Lambda a} \int_{a}^{\infty} p_{\text{birth}}(\tau) d\tau \qquad (S14)$$

$$p_{\text{div}}(\tau) = ze^{-\Lambda \tau} p_{\text{birth}}(\tau)$$
 (S15)

and

$$p_{\rm idt}(\tau) = \frac{1 - e^{-\Lambda \tau}}{1 - z^{-1}} p_{\rm birth}(\tau) , \qquad (S16)$$

where Λ and $p_{\text{birth}}(\tau)$ are determined as the solutions of

$$p_{\text{birth}}(\tau) = z \int_0^\infty K(\tau, \tau') e^{-\Lambda \tau'} p_{\text{birth}}(\tau') d\tau' , \quad (S17)$$

or equivalently,

$$p_{\rm div}(\tau) = ze^{-\Lambda\tau} \int_0^\infty K(\tau, \tau') p_{\rm div}(\tau') d\tau' .$$
 (S18)

Using the above, we obtain the relation

$$p_{\text{birth}}(\tau) = \frac{z - 1}{z} p_{\text{idt}}(\tau) + \frac{1}{z} p_{\text{div}}(\tau) . \qquad (S19)$$

Eq. (S3) becomes at stationary state

$$p_{\text{birth}}(\tau) = \int_0^\infty K(\tau, \tau') p_{\text{div}}(\tau') d\tau' . \qquad (S20)$$

II. β AS CORRELATION COEFFICIENT

The covariance between mother and daughter IDTs of isolated single cells at steady state is given by

$$\int_{0}^{\infty} d\tau \int_{\tau_{0}}^{\infty} d\tau' \tau \tau' K(\tau, \tau') k(\tau') - \left(\int_{0}^{\infty} \tau k(\tau) d\tau\right)^{2}$$

$$= \beta \left(\int_{0}^{\infty} \tau^{2} k(\tau) d\tau - \left(\int_{0}^{\infty} \tau k(\tau) d\tau\right)^{2}\right)$$
(S21)

Thus, we find that β is the Pearson correlation coefficient.

III. STATIONARY SOLUTION IN THE LEBOWITZ-RUBINOW MODEL

In this section, we derive the stationary distribution for p_{div} given in the main text Eq. (8), and determine the conditions in which a heritability threshold exists.

Let $[\tau_0, \tau_1]$ be the support of the IDT distribution $k(\tau)$

$$\operatorname{supp}(k) := \overline{\{\tau \in \mathbb{R} | k(\tau) \neq 0\}} = [\tau_0, \tau_1]. \tag{S22}$$

Here, we consider the general case $0 \le \tau_0 < \tau_1 \le \infty$, while in the main text and throughout the other sections of the Supplement, we present results for $\tau_0 > 0$ and $\tau_1 = \infty$. We recall that the function $k\left(\tau\right)$ is non-negative and normalized such that $\int_{\tau_0}^{\tau_1} k\left(\tau\right) d\tau = 1$; the parameter z denotes the average number of offspring $(0 < z < \infty)$; and β denotes the IDT heritability $(0 \le \beta < 1)$.

The stationary distribution $p_{\rm div}$ is a probability measure on the interval $[\tau_0, \tau_1]$ (see Sec. III B for details on the measure-theoretic formulation). We wish to determine the range of β for which $p_{\rm div}$ can be expressed by a probability density function $p_{\rm div}(\tau)$; in such cases substituting Eq. (1) in Eq. (3), we obtain

$$p_{\text{div}}(\tau) = \frac{(1-\beta)zk(\tau)}{e^{\Lambda\tau} - \beta z},$$
 (S23)

and Λ is determined by the normalization condition

$$\int_{\tau_0}^{\tau_1} p_{\text{div}}(\tau) d\tau = \int_{\tau_0}^{\tau_1} \frac{(1-\beta) z k(\tau)}{e^{\Lambda \tau} - \beta z} d\tau = 1 . \quad (S24)$$

If the above normalization condition does not admit a solution $\Lambda < \infty$ such that the integrand is a non-negative function of τ , the measure $p_{\rm div}$ is not equivalent to a density function (formally, it is not absolutely continuous with respect to Lebesgue measure), and, as we will show below, the measure contains a singular portion corresponding to a delta function (i.e. a Dirac measure).

To determine the conditions such that $p_{\rm div}$ is given by a density, we define

$$q_{\beta,z,\lambda}(\tau) := \frac{(1-\beta)zk(\tau)}{e^{\lambda\tau} - \beta z},$$
 (S25)

and its integral on $[\tau_0, \tau_1]$,

$$Q(\beta, z, \lambda) := \int_{\tau_0}^{\tau_1} q_{\beta, z, \lambda}(\tau) d\tau.$$
 (S26)

For fixed β , z, and $k(\tau)$, $p_{\rm div}$ is given by a density if and only if there exists $\lambda < \infty$ for which the following conditions both hold:

Condition 1 $q_{\beta,z,\lambda}(\tau) \geq 0$ for any $\tau \in [\tau_0, \tau_1]$.

Condition 2 $Q(\beta, z, \lambda) = 1$.

As we will see below, condition 2 may not be satisfied even though condition 1 holds for a range of λ .

We characterize the range of β for which $p_{\rm div}$ has a density by the value β_c , which is defined as the supremum over the set of β for which both conditions can be satisfied for some $\lambda < \infty$, given z and $k(\tau)$ are fixed; that is,

$$\beta_c := \sup\{\beta : 0 < \beta < 1 \text{ and } \exists \lambda < \infty \\
\text{s.t. conditions 1 and 2 both hold}\},$$

or $\beta_c := 0$ if the above set is empty.

When $\beta_c = 1$, $p_{\rm div}$ is given by a density for all values of β . If $\beta_c < 1$, then the system undergoes a phase transition as β increases past β_c , such that for $\beta > \beta_c$, $p_{\rm div}$ is not given by a density, and contains a singular portion that is localized at a single point. We refer to a system for which $\beta_c < 1$ as one that exhibits a heritability threshold; and refer to $p_{\rm div}$ as being localized if $\beta > \beta_c$ or delocalized if $\beta < \beta_c$. Below, we show that the behavior of $p_{\rm div}$ at precisely $\beta = \beta_c$ depends on additional details including the form of $k(\tau)$.

In Sec. III A, we determine the general conditions in which there exists a heritability threshold, i.e. when $\beta_c < 1$. In Sec. III B, we derive the form of $p_{\rm div}$ in the localized state. In Sec. III C we provide an explicit calculation of β_c in a special case, and in Sec. III D we discuss the case $\beta = 0$.

A. Heritability threshold

Throughout this subsection, we assume $0 < \beta < 1$ unless otherwise noted. We consider the definition of β_c

given in (S27), and observe that condition 1 is satisfied if and only if

$$\omega_0(\beta, z) \le \lambda < \infty$$
 (S28)

where

$$\omega_0(\beta, z) := \begin{cases} \ln (\beta z) / \tau_1, & 0 < \beta z < 1 \\ 0, & \beta z = 1 \\ \ln (\beta z) / \tau_0, & \beta z > 1 \end{cases}$$
 (S29)

If $\tau_0 > 0$, then ω_0 is finite and there exists a non-empty range of λ satisfying (S28). If $\tau_0 = 0$, then the range (S28) is non-empty if and only if $\beta z \leq 1$ (since for $\beta z > 1$ we have $\omega_0 = \infty$).

To additionally satisfy condition 2, we first identify the values of λ such that $Q(\beta, z, \lambda) < \infty$. For any λ in the range

$$\omega_0(\beta, z) < \lambda < \infty ,$$
 (S30)

we can bound the integral in (S26) yielding

$$Q(\beta, z, \lambda) \le \max\left(\frac{(1-\beta)z}{e^{\lambda \tau_0} - \beta z}, \frac{(1-\beta)z}{e^{\lambda \tau_1} - \beta z}\right)$$
(S31)

and the right-hand side is finite if and only if $\beta z \neq 1$ or $\tau_0 \neq 0$. Thus, we have $Q(\beta, z, \lambda) < \infty$ for all λ satisfying (S30) except in the special case in which both $\beta z = 1$ and $\tau_0 = 0$. We will treat that case separately when needed below.

Given that $Q(\beta, z, \lambda) < \infty$, we have that $Q(\beta, z, \lambda)$ monotonically decreases as λ increases because

$$\frac{\partial}{\partial \lambda} Q(\beta, z, \lambda) = -\int_{\tau_0}^{\tau_1} \frac{(1 - \beta) z \tau e^{\lambda \tau} k(\tau)}{(e^{\lambda \tau} - \beta z)^2} d\tau < 0.$$
(S32)

For sufficiently large λ ,

$$Q(\beta, z, \lambda) \le \frac{(1 - \beta)z}{e^{\lambda \tau_0} - \beta z} , \qquad (S33)$$

and thus $\lim_{\lambda\to\infty} Q(\beta, z, \lambda) = 0$ if $\tau_0 > 0$. If $\tau_0 = 0$ (and thus $\beta z \leq 1$ by condition 1), $\lim_{\lambda\to\infty} Q(\beta, z, \lambda) = 0$

likewise holds because for any $\epsilon \in (0, \tau_1)$,

$$Q(\beta, z, \lambda) \le \frac{(1 - \beta)z}{1 - \beta z} \int_0^{\epsilon} k(\tau)d\tau + \frac{(1 - \beta)z}{e^{\lambda \epsilon} - \beta z} \int_{\epsilon}^{\tau_1} k(\tau)d\tau .$$
(S34)

Therefore, over the range of λ satisfying condition 1, we have shown that $Q(\beta, z, \lambda)$ monotonically decreases to 0 with increasing λ . We conclude that condition 2 is satisfied if and only if

$$Q_0(\beta, z) := \lim_{\lambda \downarrow \omega_0(\beta, z)} Q(\beta, z, \lambda) \ge 1.$$
 (S35)

In this case, p_{div} is delocalized and the long-term growth rate $\Lambda_{\beta,z}$ is given by the unique solution of $Q(\beta, z, \Lambda_{\beta,z}) = 1$, where $\omega_0(\beta, z) \leq \Lambda_{\beta,z} < \infty$.

To determine when (S35) is satisfied, we analyze separately the cases z > 1, z = 1 and 0 < z < 1 below. We treat the case $\beta = 0$ separately in Sec.III D, while for the cases below we will always have $0 < \beta < 1$.

Case I:
$$z > 1$$

Given that $0 < \beta < 1$, we consider three sub-cases below, consisting of $\beta z < 1$, $\beta z > 1$, and $\beta z = 1$.

Sub-case I-1: $\beta z < 1$. In this case, we have $\omega_0(\beta,z) = \ln(\beta z)/\tau_1 \le 0$. Since $Q(\beta,z,\lambda)$ increases for decreasing λ , we have by the definition in (S35) that $Q_0(\beta,z) \ge Q(\beta,z,0)$. Since $Q(\beta,z,0) = \frac{(1-\beta)z}{(1-\beta z)}$, and given that we have assumed z>1, we have $Q(\beta,z,0)>1$. Hence, $Q_0(\beta,z)>1$ and condition 2 holds for $\beta< z^{-1}$. We conclude that $\beta_c \ge z^{-1}$.

Sub-case I-2: $\beta z > 1$. We have already seen that for $\beta z > 1$ and $\tau_0 = 0$, condition 1 cannot be satisfied implying that $\beta_c \leq z^{-1}$. Since sub-case I-1 above yielded $\beta_c \geq z^{-1}$, which holds in general, we conclude that if $\tau_0 = 0$ then $\beta_c = z^{-1}$.

We now consider $\tau_0 > 0$. Below, we show that for $\beta z > 1$, $Q_0(\beta, z) < \infty$ if and only if

$$\int_{\tau_0}^{\tau_1} \frac{k(\tau) d\tau}{\tau - \tau_0} < \infty . \tag{S36}$$

When $Q_0(\beta, z) = \infty$, Eq. S35 is satisfied, hence condition 2 is satisfied for all $\beta > z^{-1}$; this implies that $\beta_c = 1$. We thus turn to cases when $Q_0(\beta, z) < \infty$. We have $\omega_0(\beta, z) = \ln(\beta z)/\tau_0$, from which

$$Q_0(\beta, z) = \frac{1 - \beta}{\beta} \int_{\tau_0}^{\tau_1} \frac{k(\tau)}{(\beta z)^{\tau/\tau_0 - 1} - 1} d\tau$$
 (S37)

and

$$\frac{\partial}{\partial \beta} Q_0(\beta, z) = -\frac{1}{\beta^2} \int_{\tau_0}^{\tau_1} \frac{\left((1 + (1 - \beta) (\tau/\tau_0 - 1)) (\beta z)^{\tau/\tau_0 - 1} - 1 \right) k(\tau)}{\left((\beta z)^{\tau/\tau_0 - 1} - 1 \right)^2} d\tau < 0, \tag{S38}$$

since $(1+(1-\beta)(\tau/\tau_0-1))(\beta z)^{\tau/\tau_0-1} \ge 1$ holds for $\tau \in [\tau_0,\tau_1]$ with equality only when $\tau=\tau_0$. That is, $Q_0(\beta,z)$ is monotonically decreasing as β increases over the range $z^{-1} < \beta < 1$. Since relation (S36) holds, Eq. S37 yields $\lim_{\beta \to z^{-1}} Q_0(\beta,z) = \infty$ and $\lim_{\beta \to 1} Q_0(\beta,z) = 0$; therefore, there exists β_c such that $Q_0(\beta_c,z) = 1$ with $z^{-1} < \beta_c < 1$, i.e. a heritability threshold exists.

In summary, if $\tau_0 > 0$ and z > 1, we conclude that $z^{-1} < \beta_c < 1$ if $\int_{\tau_0}^{\tau_1} d\tau^k (\tau)/(\tau - \tau_0) < \infty$, and $\beta_c = 1$ otherwise. We know that $p_{\rm div}$ is localized (delocalized) if $\beta > \beta_c$ ($\beta < \beta_c$). Moreover, since $Q_0(\beta_c, z) = 1$ satisfies (S35), we have that at $\beta = \beta_c$, $p_{\rm div}$ is delocalized. If $\tau_0 = 0$, we have $\beta_c = z^{-1}$; the behavior of $p_{\rm div}$ at β_c corresponds to the unique special case in which the bound given in (S31) is not finite, and $Q(\beta, z, \lambda) < \infty$ may or may not hold. This special case is analyzed next.

Sub-case I-3: $\beta = z^{-1}$. We only need to analyze the case $\tau_0 = 0$, since for $\tau_0 > 0$, we showed above that $\beta_c > z^{-1}$, hence we know that p_{div} is delocalized for $\beta = z^{-1}$. For $\tau_0 = 0$, we have

$$Q\left(\beta,z,\lambda\right) = Q\left(z^{-1},z,\lambda\right) = \int_{0}^{\tau_{1}} \frac{\left(z-1\right)k\left(\tau\right)}{e^{\lambda\tau} - 1} d\tau ,$$
(S39)

where the integral may diverge or converge for $\lambda > \omega_0(z^{-1},z) = 0$. If $\int_0^{\tau_1} \tau^{-1} k(\tau) d\tau = \infty$, $Q(z^{-1},z,\lambda) = \infty$ for any $\lambda > 0$ and condition 2 cannot be satisfied. If $\int_0^{\tau_1} \tau^{-1} k(\tau) d\tau < \infty$, $Q(z^{-1},z,\lambda) < \infty$ for all $\lambda > 0$, and the argument leading to relation (S35) applies.

In summary, for $\tau_0 = 0$ and z > 1, the system exhibits a heritability threshold, given by $\beta_c = z^{-1}$. Convergence or divergence of $\int_0^{\tau_1} \tau^{-1} k(\tau) d\tau$ affects the uniqueness of the stationary solution at $\beta = \beta_c$ (see Sec. III B).

In conclusion, for z > 1 we have shown that

$$\beta_c \begin{cases} = z^{-1}, & \text{if } \tau_0 = 0, \\ > z^{-1}, & \text{if } \tau_0 > 0 \text{ and } \int_{\tau_0}^{\tau_1} \frac{k(\tau)}{(\tau - \tau_0)} < \infty, \\ = 1, & \text{if } \tau_0 > 0 \text{ and } \int_{\tau_0}^{\tau_1} \frac{k(\tau)}{(\tau - \tau_0)} = \infty. \end{cases}$$
(S40)

Case II:
$$z = 1$$

Since $Q(\beta, 1, 0) = 1$, $\Lambda = 0$ is the unique solution of $Q(\beta, 1, \Lambda) = 1$. Thus $\beta_c = 1$ for z = 1.

Case III:
$$0 < z < 1$$

For z < 1, we have $0 < \beta z < 1$, $\omega_0(\beta, z) = \ln(\beta z)/\tau_1 \le 0$, and

$$Q_0(\beta, z) = \frac{1 - \beta}{\beta} \int_{\tau_0}^{\tau_1} \frac{k(\tau)}{(\beta z)^{\tau/\tau_1 - 1} - 1} d\tau .$$
 (S41)

If $\tau_1 = \infty$, $Q_0(\beta, z) = \frac{(1-\beta)z}{(1-\beta z)} < 1$ and thus by relation (S35) condition 2 cannot be satisfied for any value of β , implying $\beta_c = 0$.

If $\tau_1 < \infty$, we show below that for $\beta z < 1$, $Q_0(\beta, z) < \infty$ if and only if

$$\int_{\tau_0}^{\tau_1} \frac{k(\tau) d\tau}{\tau_1 - \tau} < \infty . \tag{S42}$$

If $Q_0(\beta, z) = \infty$, then relation (S35) implies that $\beta_c = 1$. For $Q_0(\beta, z) < \infty$, we can show that

$$\frac{\partial}{\partial \beta} Q_0(\beta, z) = -\frac{1}{\beta^2} \int_{\tau_0}^{\tau_1} \frac{\left((1 + (1 - \beta) (\tau/\tau_1 - 1)) (\beta z)^{\tau/\tau_1 - 1} - 1 \right) k(\tau)}{\left((\beta z)^{\tau/\tau_1 - 1} - 1 \right)^2} d\tau < 0.$$
 (S43)

To prove this inequality, it suffices to prove that $((1-\beta)x-1)(\beta z)^{-x}+1>0$ for $x\in(0,1)$; since $(\beta z)^x<\beta^x$ for x>0, the result will follow by showing that $\beta^x<1-(1-\beta)x$ for $x\in(0,1)$. To see this, we define $f(x):=1-(1-\beta)x-\beta^x$, which has a unique maximum at $x^*:=\ln[(\beta-1)/\ln\beta]/\ln\beta$. $0<1-\beta<-\ln\beta$ implies $x^*>0$ and $\beta^{-1}-1>-\ln\beta$ implies $x^*<1$. Together with f(0)=f(1)=0 and continuity of f(x) and f'(x), f(x)>0 is true for $x\in(0,1)$.

Next, we show that $\lim_{\beta \uparrow 1} Q_0(\beta, z) = 0$ and $\lim_{\beta \downarrow 0} Q_0(\beta, z) = \infty$. We have

$$\lim_{\beta \uparrow 1} Q_0(\beta, z) = 0 \cdot \int_{\tau_0}^{\tau_1} \frac{k(\tau)}{z^{\tau/\tau_1 - 1} - 1} d\tau = 0, \quad (S44)$$

since the integral converges by relation (S42) which holds. For $\tau_0 > 0$, we can bound the integral in (S41) to obtain

$$Q_0(\beta, z) \ge \frac{(1 - \beta) z}{(\beta z)^{\tau_0/\tau_1} - \beta z}, \tag{S45}$$

while for $\tau_0 = 0$, given $\epsilon \in (0, \tau_1)$ we have

$$Q_{0}(\beta, z) \ge (1 - \beta) z \left((\beta z)^{\epsilon/\tau_{1}} - \beta z \right)^{-1} \int_{\epsilon}^{\tau_{1}} k(\tau) d\tau;$$
(S46)

hence $\lim_{\beta \downarrow 0} Q_0(\beta, z) = \infty$. Therefore, there exists $0 < \beta_c < 1$ such that $Q_0(\beta_c, z) = 1$.

In conclusion, for 0 < z < 1 we have shown that

$$\beta_c \begin{cases} = 0, & \text{if } \tau_1 = \infty, \\ < 1, & \text{if } \tau_1 < \infty \text{ and } \int_{\tau_0}^{\tau_1} \frac{k(\tau)}{|(\tau_1 - \tau)|} < \infty, \\ = 1, & \text{if } \tau_1 < \infty \text{ and } \int_{\tau_0}^{\tau_1} \frac{k(\tau)}{|(\tau_1 - \tau)|} = \infty. \end{cases}$$
 (S47)

Table S2 provides a summary of the results obtained in this subsection.

Necessary and sufficient condition for $Q_0(\beta, z) < \infty$

It suffices to prove that $\int_{\tau_0}^{\tau_1} k(\tau) d\tau/(\tau - \tau_0) < \infty$ is equivalent to $Q_0(\beta, z) < \infty$ when $\tau_0 > 0$, z > 1 and $\beta z > 1$. In the same way one can show that for $\tau_1 < \infty$, 0 < z < 1, and $\beta z < 1$, $\int_{\tau_0}^{\tau_1} \frac{k(\tau)}{(\tau_1 - \tau)} < \infty$ is the equivalent condition.

We write

$$Q_{0}(\beta, z) \cdot \beta / (1 - \beta) = \int_{\tau_{0}}^{\tau_{1}} \frac{k(\tau)}{(\beta z)^{\tau/\tau_{0} - 1} - 1} d\tau$$
 (S48)
$$= \int_{\tau_{0}}^{\tau_{0} + \epsilon} \frac{k(\tau)}{(\beta z)^{\tau/\tau_{0} - 1} - 1} d\tau + \int_{\tau_{0} + \epsilon}^{\tau_{1}} \frac{k(\tau)}{(\beta z)^{\tau/\tau_{0} - 1} - 1} d\tau$$

where $0 < \epsilon \ll 1$. The second term of RHS is finite, while the integrand of the first term can be expanded with respect to $\tau - \tau_0$, yielding

$$\int_{\tau_0}^{\tau_0 + \epsilon} \frac{k(\tau)}{\omega_0} \left(\frac{1}{\tau - \tau_0} + \mathcal{O}(1) \right) d\tau$$

$$= \frac{1}{\omega_0} \int_{\tau_0}^{\tau_0 + \epsilon} \frac{k(\tau)}{\tau - \tau_0} d\tau + \mathcal{O}(\epsilon)$$
 (S49)

where $\omega_0 = \tau_0^{-1} \ln{(\beta z)}$. Thus, for any β satisfying $z^{-1} < \beta < 1$, $Q_0(\beta, z)$ is finite if and only if $\int_{\tau_0}^{\tau_0 + \epsilon} {}_k(\tau) d\tau/(\tau - \tau_0)$ is finite for small $\epsilon > 0$. This is equivalent to $\int_{\tau_0}^{\tau_1} {}_k(\tau) d\tau/(\tau - \tau_0) < \infty$.

For example, if $k\left(\tau\right)$ behaves power of $\tau-\tau_0$ near τ_0 , that is, there exists a real number γ and $\lim_{\tau\downarrow\tau_0}\frac{k(\tau)}{(\tau-\tau_0)^{\gamma}}=A$ where A is a positive finite constant, then the dominant term of $\int_{\tau_0}^{\tau_0+\epsilon}\frac{k(\tau)}{\tau-\tau_0}d\tau$ will be $A\int_0^\epsilon x^{\gamma-1}dx$. Thus $\int_{\tau_0}^{\tau_0+\epsilon}\frac{k(\tau)}{\tau-\tau_0}d\tau<\infty$ if and only if $\gamma>0$.

B. Stationary solution for $\beta > \beta_c$

For $\beta > \beta_c$, the measure $p_{\rm div}$ does not have a density relative to Lebesgue measure μ on the interval $[\tau_0, \tau_1]$. We find that in this regime $p_{\rm div}$ includes a Dirac delta function, and we obtain the exact form of the solution, which we will show below to be unique.

We seek a solution of the form

$$p_{\text{div}}(\tau) = \widetilde{p}(\tau) + C_1 \delta(\tau - \tau_*) \tag{S50}$$

where $\tau_* \in [\tau_0, \tau_1]$, $C_1 > 0$, $\delta(\cdot)$ denotes the Dirac delta function, and $\widetilde{p}(\tau) \geq 0$ is a Lebesgue integrable function

on $[\tau_0, \tau_1]$, i.e. $\widetilde{p} \in L^1(\mu)$. The normalization is given by $\int_{\tau_0}^{\tau_1} p_{\text{div}}(\tau) d\tau = 1$. Substituting Eq. (S50) in Eq. (3)we have

$$(1 - \beta z e^{-\Lambda \tau}) \widetilde{p}(\tau) = (1 - \beta) z e^{-\Lambda \tau} k(\tau), \quad (S51a)$$

$$\left(1 - \beta z e^{-\Lambda \tau_*}\right) C_1 = 0. \tag{S51b}$$

Since $\widetilde{p}(\tau) \geq 0$ for $\tau \in [\tau_0, \tau_1]$, $\Lambda \geq \omega_0(\beta, z)$; and $C_1 > 0$ implies $\Lambda = \ln(\beta z) / \tau_* \leq \omega_0(\beta, z)$ for $\tau_* \in [\tau_0, \tau_1]$. Thus

$$\Lambda = \omega_0(\beta, z) = \ln(\beta z) / \tau_* . \tag{S52}$$

Using $\beta > \beta_c \ge z^{-1}$ for z > 1, we have $\beta z > 1$ hence $\omega_0 = \ln(\beta z)/\tau_0$; while for z < 1, $\beta z < 1$ yields $\omega_0 = \ln(\beta z)/\tau_1$. These relations together with (S52) imply

$$\tau_* = \begin{cases} \tau_0, & z > 1\\ \tau_1, & z < 1 \end{cases}$$
 (S53)

Due to the normalization $\int_{\tau_0}^{\tau_1} p_{\text{div}}(\tau) d\tau = 1$,

$$C_{1} = 1 - Q(\beta, z, \omega_{0}(\beta, z))$$

$$= 1 - \int_{\tau_{0}}^{\tau_{1}} \frac{(1 - \beta) zk(\tau')}{(\beta z)^{\tau'/\tau_{*} - 1} - \beta z} d\tau', \qquad (S54)$$

and $0 < C_1 \le 1$ holds because $Q(\beta, z, \omega_0(\beta, z)) = Q_0(\beta, z) < 1$ for $\beta > \beta_c$.

To summarize, for $\tau_0 > 0$, z > 1 and $\beta > \beta_c$,

$$\Lambda = \frac{\ln(\beta z)}{\tau_0},\tag{S55a}$$

$$p_{\text{div}}(\tau) = q_{\beta,z,\Lambda}(\tau) + (1 - Q(\beta,z,\Lambda)) \delta(\tau - \tau_0),$$
(S55b)

and for $\tau_1 < \infty$, z < 1 and $\beta > \beta_c$,

$$\Lambda = \frac{\ln(\beta z)}{\tau_1},\tag{S56a}$$

$$p_{\text{div}}\left(\tau\right) = q_{\beta,z,\Lambda}\left(\tau\right) + \left(1 - Q\left(\beta,z,\Lambda\right)\right)\delta\left(\tau - \tau_{1}\right) . \tag{S56b}$$

We note that in the limits $\tau_0 \to 0$ and $\tau_1 \to \infty$, the limiting forms of the solution above are likewise valid. The case in which z > 1 and $\tau_0 \to 0$ is unique in that we have $\Lambda = \infty$ and $Q(\beta, z, \Lambda) = 0$, hence $p_{\rm div} = \delta(\tau - \tau_0)$ for all $\beta > \beta_c = z^{-1}$, and the entire population is localized at τ_0 above the transition. Moreover, in this special case, uniqueness of the solution at $\beta = \beta_c$ is dependent on whether or not $\int_0^{\tau_1} \tau^{-1} k(\tau) d\tau$ converges, as discussed below.

For cases in which z < 1, a finite population will eventually go extinct, and in practice the stationary solution Eq. (S56) may or may not be realized before extinction occurs. Lastly, we remark that Eqs. (S55b) and (S56b) also hold for $\beta < \beta_c$ as $Q(\beta, z, \Lambda) = 1$.

Uniqueness of the solution. We establish existence and uniqueness of the stationary solution by the same approach used in [1] to study Kingman's "house of cards" model in evolutionary dynamics.

We set $X = [\tau_0, \tau_1]$, and let $\mathcal{B}(X)$ be the collection of Borel sets in X, let \mathcal{P} denote the set of all Borel probability measures on X, and let μ denote the Lebesgue measure on X. Eq. (3) is rewritten for a stationary probability measure $P \in \mathcal{P}$ as

$$P(A) = \int_{A} \beta z e^{-\Lambda \tau} P(d\tau) + \int_{A} (1 - \beta) z e^{-\Lambda \tau} k(\tau) \mu(d\tau)$$
(S57)

for any set $A \subset \mathcal{B}(X)$, and where P(X) = 1.

Given that $P \in \mathcal{P}$ of Eq. (S57) exists, then by the Lebesgue decomposition theorem there exist unique $P_a, P_s \in \mathcal{P}$ and a number $C_1 \in [0, 1]$ such that

$$P = (1 - C_1) P_a + C_1 P_s \tag{S58}$$

where P_a is absolutely continuous with respect to μ and P_s is singular with respect to μ ; i.e. there exists $f \geq 0 \in L^1(\mu)$ such that $P_a(A) = \int_A f(\tau)\mu(d\tau)$ and P_s is supported on a μ -null set. Decomposing Eq. (S57) into absolutely continuous and singular parts with respect to μ , we have

$$(1 - C_1) \int_A (1 - \beta z e^{-\Lambda \tau}) P_a(d\tau)$$

$$= \int_A (1 - \beta) z e^{-\Lambda \tau} k(\tau) \mu(d\tau)$$

$$C_1 \int_A (1 - \beta z e^{-\Lambda \tau}) P_s(d\tau) = 0$$
(S59a)

From Eq. (S59a),

$$\widetilde{p}(\tau) = \frac{(1-\beta)ze^{-\Lambda\tau}k(\tau)}{1-\beta ze^{-\Lambda\tau}}, \, \mu\text{-a.s. on } [\tau_0, \tau_1] \quad (S60)$$

where $\widetilde{p}(\tau)$ is a non-negative Lebesgue integrable function such that $(1-C_1)P_a(A)=\int_A \widetilde{p}(\tau)\,\mu\,(d\tau)$ for any measurable set $A\in [\tau_0,\tau_1]$, denoted by $(1-C_1)P_a=\widetilde{p}\,\mu$ in short. $\widetilde{p}(\tau)\geq 0,\,\mu$ -a.s. on $[\tau_0,\tau_1]$ implies $\Lambda\geq \omega_0\,(\beta,z)$. By the normalization $P_a\,([\tau_0,\tau_1])=1$, we have

$$C_1 = 1 - \int_{\tau_0}^{\tau_1} \widetilde{p}(\tau) \mu(d\tau) = 1 - Q(\beta, z, \Lambda) , \quad (S61)$$

for $Q(\beta, z, \Lambda) < 1$. From Eq. (S59b), $C_1 = 0$ or

$$e^{\Lambda \tau} = \beta z, P_s$$
-a.s. on $[\tau_0, \tau_1]$. (S62)

Therefore, if $C_1 > 0$, P_s is the Dirac measure concentrated on τ_* , denoted by δ_{τ_*} , with $\tau_* = \frac{\ln{(\beta z)}}{\Lambda} \in [\tau_0, \tau_1]$. This implies that if $C_1 > 0$ then $\Lambda \leq \omega_0(\beta, z)$ and thus $\Lambda = \omega_0(\beta, z)$.

From the discussion in Sec. III A, for $\tau_0 > 0$ and $0 < \beta \le \beta_c$, there exists $\Lambda \ge \omega_0(\beta, z)$ such that $Q(\beta, z, \Lambda) = 1$ and $P = \widetilde{p}\mu$. For $\beta > \beta_c$, on the other hand, $\Lambda = 1$

 $\omega_0(\beta, z)$ and $P = \widetilde{p} \mu + (1 - Q(\beta, z, \omega_0(\beta, z))) \delta_{\tau_*}$ where τ_* is given by Eq. (S53).

In the special case of $\tau_0 = 0$, for $\beta > \beta_c = z^{-1}$, $\Lambda = \ln(\beta z)/\tau_0 = \infty$, which yields $\tilde{p}(\tau) = 0$, $Q(\beta, z, \Lambda) = 0$, and $C_1 = 1$; hence the stationary solution P is given by Dirac measure δ_{τ_0} concentrated at $\tau_0 = 0$. At $\beta = \beta_c = z^{-1}$, if $\int_0^{\tau_1} \tau^{-1} k(\tau) d\tau = \infty$, we have $Q(\beta, z, \Lambda) = \infty$, and it is not possible to normalize $\tilde{p}(\tau)$ for any $\Lambda < \infty$; in this case, the only possible solution of Eq. S59a can be understood as the limiting case $\beta z \downarrow 1$, which yields $C_1 = 1$, $\Lambda = \infty$, and $P = \delta_{\tau_0}$. If on the other hand $\int_0^{\tau_1} \tau^{-1} k(\tau) d\tau < \infty$, then for all $0 \le C_1 \le 1$, there exists $\Lambda(C_1)$ such that $Q(z^{-1}, z, \Lambda(C_1)) = 1 - C_1$, or

$$1 - C_1 = \int_0^{\tau_1} \frac{(z - 1)k(\tau)}{e^{\Lambda(C_1)\tau} - 1} \mu(d\tau) , \qquad (S63)$$

and we obtain solutions at all growth rates $\Lambda \geq \Lambda(0)$.

C. Small variation limit of IDT distribution

We consider the limit in which $k(\tau)$ has vanishingly small variation. We fix $\bar{\tau} \in [\tau_0, \tau_1]$, and taking the limit $k(\tau) \to \delta(\tau - \bar{\tau})$,

$$Q(\beta, z, \lambda) = \frac{(1 - \beta)z}{e^{\lambda \bar{\tau}} - \beta z}$$
 (S64)

and

$$Q_0(\beta, z) = \begin{cases} \frac{(1-\beta)z}{(\beta z)^{\bar{\tau}/\tau_0} - \beta z}, & 0 < \beta z < 1\\ \frac{(1-\beta)z}{(\beta z)^{\bar{\tau}/\tau_1} - \beta z}, & \beta z > 1 \end{cases}$$
 (S65)

For z>1, we have $\beta_c z>1$, and $Q_0\left(\beta_c,z\right)=1$ implies $\beta_c=z^{\tau_0/\bar{\tau}-1};$ while for z<1, we find $\beta_c=z^{\tau_1/\bar{\tau}-1}.$ Therefore

$$\beta_c = \begin{cases} z^{\tau_1/\bar{\tau}-1}, & 0 < z < 1\\ 1, & z = 1\\ z^{\tau_0/\bar{\tau}-1}, & z > 1 \end{cases}$$
 (S66)

We also observe $\beta_c \downarrow 0$ as $\tau_1 \uparrow \infty$ for z < 1 and $\beta_c \downarrow z^{-1}$ as $\tau_0 \downarrow 0$ for z > 1. This limiting case is also included in Table S2.

D. Independent divisions: $\beta = 0$

For $\beta=0$, condition 1 is automatically satisfied for any λ , since $q_{0,z,\lambda}(\tau)=ze^{-\lambda\tau}k(\tau)$; here we determine when condition 2 is satisfied.

For z>1, since Q(0,z,0)=z>1, it suffices to show that $Q(0,z,\lambda)$ is monotonically decreasing (which follows from (S32)) to 0 for $\lambda \geq 0$. If $\tau_0>0$, $Q(0,z,\lambda)$ is finite and $\lim_{\lambda\to\infty}Q(0,z,\lambda)=0$ because $0\leq Q(0,z,\lambda)\leq ze^{-\lambda\tau_0}\to 0$ as $\lambda\to\infty$. If $\tau_0=0$, $\lim_{\lambda\to\infty}Q(0,z,\lambda)=0$ holds since the argument given at Eq. (S34) holds for

 $\beta=0.$ We thus have a unique $\lambda>0$ satisfying conditions 1 and 2.

For z=1, $Q(0,1,\lambda)$ is monotonically decreasing to 0 for $\lambda \geq 0$. Since $Q(0,1,\lambda) > 1$ for any $\lambda < 0$, $\lambda = 0$ is the unique root of $Q(0,1,\lambda) = 1$.

Finally, let 0 < z < 1. If $\tau_1 < \infty$, then $Q(0, z, \lambda) \le ze^{-\lambda \tau_1} < \infty$ for any $\lambda \le 0$; in particular, $Q(\beta, z, 0) < 1$. For some $\epsilon \in (0, \tau_1)$,

$$Q\left(0, z, \lambda\right) \ge z \int_{\epsilon}^{\tau_{1}} e^{-\lambda \tau} k\left(\tau\right) \ge z e^{-\lambda \epsilon} \int_{\epsilon}^{\tau_{1}} k\left(\tau\right) \quad (S67)$$

Thus $\lim_{\lambda \to -\infty} Q\left(0,z,\lambda\right) = \infty$. Since $Q\left(0,z,\lambda\right)$ decreases monotonically with respect to λ , there is a unique $\lambda < 0$ satisfying conditions 1 and 2. If $\tau_1 = \infty$, λ may be bounded below by $\lambda_c \in (-\infty,0)$ for $Q\left(0,z,\lambda\right)$ to be finite. In this case, a unique $\lambda < 0$ satisfying conditions 1 and 2 exists as long as $\lim_{\lambda \downarrow \lambda_c} Q\left(0,z,\lambda\right) > 1$ holds. For example, consider $k\left(\tau\right) = \tau^{-2}e^{-\tau}/\int_1^\infty \tau^{-2}e^{-\tau}d\tau$ for $\tau \geq 1$ $(\tau_0 = 1, \tau_1 = \infty)$. Then $Q\left(0,z,\lambda\right) < \infty$ for $\lambda \geq -1$ and $Q\left(0,z,\lambda\right) = \infty$ for $\lambda < -1$. In particular, $Q\left(0,z,-1\right) = z/\int_1^\infty \tau^{-2}e^{-\tau}d\tau$. Thus if $z < \int_1^\infty \tau^{-2}e^{-\tau}d\tau(<1)$, there is no λ such that $Q\left(0,z,\lambda\right) = 1$.

IV. STATISTICS OF LINEAGES

A. Mean block size for $\beta > \beta_c$ in finite time

We consider the case in which a heritability threshold β_c exists, and analyze $\beta > \beta_c$. Then $\Lambda_{\beta,z} = \omega_0 = \tau_0^{-1} \ln{(\beta z)}$. We choose t > 0 large enough so that $\int_t^\infty k\left(\tau\right) d\tau \ll 1$ and let $m_t = m_t(m,\tau)$ denote the block size with cutoff $\lfloor t/\tau \rfloor$; we define $m_t = m$ if $m < \lfloor t/\tau \rfloor$ and otherwise $m_t = \lfloor t/\tau \rfloor$, where $\lfloor x \rfloor$ denotes the integer part of a real x. Let $\rho(\tau) := \beta z e^{-\omega_0 \tau}$. Note that $\rho(\tau) < 1$ for $\tau > \tau_0$ and $\rho(\tau) = 1$ for $\tau = \tau_0$. Using Eq. (9), the joint probability distribution of m_t and τ is

$$\widetilde{p}_{\text{lin}}(m_t, \tau) = \widetilde{p}_{\text{lin}}(m_t | \tau) \, p_{\text{div}}(\tau),$$
 (S68)

where

$$\widetilde{p}_{\text{lin}}(m_t|\tau) = \begin{cases} (1 - \rho(\tau))\rho(\tau)^{m_t - 1}, & 1 \le m_t < \lfloor t/\tau \rfloor \\ \rho(\tau)^{m_t - 1}, & m_t = \lfloor t/\tau \rfloor \end{cases}$$
(S69)

Noting

$$\sum_{m_{t}=1}^{\lfloor t/\tau \rfloor} m_{t} \cdot \widetilde{p}_{\text{lin}} (m_{t}|\tau) = \sum_{m_{t}=1}^{\lfloor t/\tau \rfloor - 1} m_{t} (1 - \rho(\tau)) \rho(\tau)^{m_{t} - 1}$$

$$+ \lfloor t/\tau \rfloor \rho(\tau)^{\lfloor t/\tau \rfloor - 1}$$

$$= \sum_{m_{t}=1}^{\lfloor t/\tau \rfloor} \rho(\tau)^{m_{t} - 1}$$

$$= \begin{cases} \frac{1 - \rho(\tau)^{\lfloor t/\tau \rfloor}}{1 - \rho(\tau)}, & \tau > \tau_{0} \\ \lfloor t/\tau_{0} \rfloor, & \tau = \tau_{0}, \end{cases}$$
 (S70)

the mean block size with cutoff is defined as

$$\overline{m_t} := \int_0^t d\tau \sum_{m_t=1}^{\lfloor t/\tau \rfloor} m_t \cdot \widetilde{p}_{\text{lin}}(m_t, \tau)$$

$$= \int_0^t q_{\beta, z, \omega_0}(\tau) \frac{1 - \rho(\tau)^{\lfloor t/\tau \rfloor}}{1 - \rho(\tau)} d\tau$$

$$+ (1 - Q(\beta, z, \omega_0)) \lfloor t/\tau_0 \rfloor. \tag{S71}$$

If $\int_0^\infty q_{\beta,z,\omega_0}(\tau) \left(1 - \beta z e^{-\omega_0 \tau}\right)^{-1} d\tau < \infty$,

$$\overline{m_t} = (1 - Q(\beta, z, \omega_0)) \left| \frac{t}{\tau_0} \right| + \mathcal{O}(1)$$
 (S72)

If $k(\tau) \simeq A(\tau - \tau_0)^{\alpha - 1}$ (A is a positive constant and $\alpha > 1$) near τ_0 , we can show that the leading term of $\overline{m_t}$ is $(1 - Q(\beta, z, \omega_0)) \lfloor t/\tau_0 \rfloor$ even when $\int_0^\infty q_{\beta,z,\omega_0}(\tau) \left(1 - \beta z e^{-\omega_0 \tau}\right)^{-1} d\tau = \infty$. Let $1 < \alpha \le 2$. We show $\int_0^t q_{\beta,z,\omega_0}(\tau) \frac{1 - \left(\beta z e^{-\omega_0 \tau}\right)^{\lfloor t/\tau \rfloor}}{1 - \beta z e^{-\omega_0 \tau}} d\tau$ is bounded by $\mathcal{O}\left(t^{2-\alpha}\right)$. Let $\tau_c := \tau_0 + \tau_0/(\omega_0 t)$. Then

$$\frac{1 - (\beta z e^{-\omega_0 \tau})^{\lfloor t/\tau \rfloor}}{1 - \beta z e^{-\omega_0 \tau}} \le \begin{cases} \left\lfloor \frac{t}{\tau_0} \right\rfloor, & \tau_0 \le \tau < \tau_c \\ \left(1 - \beta z e^{-\omega_0 \tau}\right)^{-1}, & \tau \ge \tau_c \end{cases}$$
(S73)

holds. Thus

$$\int_{0}^{t} q_{\beta,z,\omega_{0}}(\tau) \frac{1 - (\beta z e^{-\omega_{0}\tau})^{\lfloor t/\tau \rfloor}}{1 - \beta z e^{-\omega_{0}\tau}} d\tau \leq (S74)$$

$$\left\lfloor \frac{t}{\tau_{0}} \right\rfloor \int_{\tau_{0}}^{\tau_{c}} q_{\beta,z,\omega_{0}}(\tau) d\tau + \int_{\tau_{c}}^{\tau_{m}} \frac{q_{\beta,z,\omega_{0}}(\tau)}{1 - \beta z e^{-\omega_{0}\tau}} d\tau$$

$$+ \int_{\tau}^{\infty} \frac{q_{\beta,z,\omega_{0}}(\tau)}{1 - \beta z e^{-\omega_{0}\tau}} d\tau ,$$

where $\tau_m > \tau_c$ is a constant independent of t. For large enough t, τ_m can be close enough to τ_0 such that $k(\tau_m) \simeq A(\tau_m - \tau_0)^{\alpha - 1}$ and $1 - \beta z e^{-\omega_0 \tau_m} \simeq \omega_0(\tau_m - \tau_0)$. Note that the third term of RHS is constant. The first and second terms are evaluated as

$$\int_{\tau_0}^{\tau_c} q_{\beta,z,\omega_0}(\tau) d\tau = \mathcal{O}\left(\left(\frac{\tau_0}{t}\right)^{\alpha-1}\right)$$
 (S75)

and

$$\int_{\tau_c}^{\tau_m} \frac{q_{\beta,z,\omega_0}(\tau)}{1 - \beta z e^{-\omega_0 \tau}} d\tau = \begin{cases} \mathcal{O}\left(\left(\frac{t}{\tau_0}\right)^{2 - \alpha}\right), & 1 < \alpha < 2\\ \mathcal{O}\left(\ln\left(\frac{t}{\tau_0}\right)\right), & \alpha = 2 \end{cases}$$
(S76)

Therefore, $\int_0^t q_{\beta,z,\omega_0}(\tau) \frac{1-\left(\beta z e^{-\omega_0 \tau}\right)^{\lfloor t/\tau \rfloor}}{1-\beta z e^{-\omega_0 \tau}} d\tau$ is increasing slower than $\mathcal{O}(t)$. Thus, for $\beta > \beta_c$, the leading term of $\overline{m_t}$ is $(1-Q(\beta,z,\omega_0))\lfloor t/\tau \rfloor$ not only when $\alpha > 2$ but also when $1 < \alpha \le 2$.

B. Lineage IDT distribution for $\beta > \beta_c$

Using the same approach as in the calculations above, we can compute $p_{\text{lin}}(\tau)$ for $\beta > \beta_c$ as the $t \to \infty$ limit of the finite time version of expression Eq. (11); that is, we define

$$\widetilde{p}_{\text{lin}}\left(\tau\right) := \frac{\sum_{m_t=1}^{\lfloor t/\tau \rfloor} m_t \cdot \widetilde{p}_{\text{lin}}\left(m_t, \tau\right)}{\overline{m}_t} \ . \tag{S77}$$

We integrate the above distribution over a small interval $(\tau, \tau + \delta t)$, and using Eqs. S68 and S70, we find

$$\int_{\tau}^{\tau+\delta t} \widetilde{p}_{\text{lin}}(\tau') d\tau' = \begin{cases} \frac{q_{\beta,z,\omega_{0}}(\tau)}{\overline{m}_{t}} \cdot \frac{1-\rho(\tau)^{\lfloor t/\tau \rfloor}}{1-\rho(\tau)} \delta t, & \tau > \tau_{0} \\ \frac{1-Q(\beta,z,\omega_{0})}{\overline{m}_{t}} \lfloor t/\tau_{0} \rfloor + \mathcal{O}(\delta t), & \tau = \tau_{0}. \end{cases}$$
(S78)

Using Eq. S72, we see that as $t \to \infty$, $\widetilde{p}_{lin}(\tau) \to \delta(\tau - \tau_0)$.

C. On-lineage mean IDT

For $\beta<\beta_c, \overline{m}_{\beta,z}<\infty$ always holds, while for $\beta>\beta_c$, $\overline{m}_{\beta,z}=\infty$, but $\lim_{\beta\uparrow\beta_c}\overline{m}_{\beta,z}$ may diverge or converge depending on k (τ). The distribution $p_{\text{lin}}(\tau)$ converges to a density as $\beta\uparrow\beta_c$ when $\lim_{\beta\uparrow\beta_c}\overline{m}_{\beta,z}<\infty$. In contrast, when $\lim_{\beta\uparrow\beta_c}\overline{m}_{\beta,z}=\infty$, $p_{\text{lin}}(\tau)$ weakly converges to δ ($\tau-\tau_0$) as $\beta\uparrow\beta_c$ because $\lim_{\beta\uparrow\beta_c}\int_{\tau_0+\epsilon}^\infty p_{\text{lin}}(\tau)\,d\tau=0$ for any $\epsilon>0$. Thus $\lim_{\beta\uparrow\beta_c}\overline{m}_{\beta,z}^{-1}$ is an important quantity to distinguish continuous or discontinuous transition; $\lim_{\beta\uparrow\beta_c}\overline{m}_{\beta,z}^{-1}>0$ indicates a discontinuous transition and $\lim_{\beta\uparrow\beta_c}\overline{m}_{\beta,z}^{-1}=0$ a continuous one. Under the condition $\int_{\tau_0}^\infty d\tau^k(\tau)/(\tau-\tau_0)<\infty$,

$$\int_{\tau_0}^{\infty} \frac{(\tau - \tau_0) q_{\beta_c, z, \omega_0}(\tau)}{1 - \beta_c z e^{-\omega_0 \tau}} d\tau$$

$$= \frac{1 - \beta_c}{\beta_c} \int_{\tau_0}^{\infty} \frac{(\tau - \tau_0) (\beta_c z)^{1 - \tau/\tau_0} k(\tau)}{\left(1 - (\beta_c z)^{1 - \tau/\tau_0}\right)^2} d\tau < \infty \quad (S79)$$

holds. Therefore, the on-lineage mean IDT, $\overline{\tau}_{\beta,z} = \int_0^\infty \tau p_{\text{lin}}(\tau) d\tau$ shows continuous/discontinuous phase transition depending on convergence/divergence of $\lim_{\beta \uparrow \beta,z} \overline{m}_{\beta,z}$:

$$\lim_{\beta \uparrow \beta_c} \overline{\tau}_{\beta,z} - \tau_0 = \frac{1}{\lim_{\beta \uparrow \beta_c} \overline{m}_{\beta,z}} \int_{\tau_0}^{\infty} \frac{(\tau - \tau_0) \, q_{\beta_c,z,\omega_0}(\tau)}{1 - \beta_c z e^{-\omega_0 \tau}} d\tau$$

$$\begin{cases}
> 0, & \lim_{\beta \uparrow \beta_c} \overline{m}_{\beta,z} < \infty \\
= 0, & \lim_{\beta \uparrow \beta_c} \overline{m}_{\beta,z} = \infty
\end{cases}$$
(S80)

Applying this result to Eq. (S94), $\partial_{\ln \beta} \Lambda(\beta, z)$ is discontinuous at $\beta = \beta_c$ if $\lim_{\beta \uparrow \beta_c} \overline{m}_{\beta,z} < \infty$ and otherwise continuous. If $\lim_{\tau \downarrow \tau_0} \frac{k(\tau)}{(\tau - \tau_0)^{\alpha - 1}}$ converges a positive finite constant where $\alpha > 1$,

$$\lim_{\beta \uparrow \beta_c} \overline{m}_{\beta,z} \begin{cases} < \infty, & \alpha > 2 \\ = \infty, & 1 < \alpha \le 2 \end{cases}$$
 (S81)

and therefore

$$\lim_{\beta \uparrow \beta_c} \overline{\tau}_{\beta,z} - \tau_0 \begin{cases} > 0, & \alpha > 2 \\ = 0, & 1 < \alpha \le 2 \end{cases}$$
 (S82)

D. Block size distribution

The lineage distribution of m for $\beta < \beta_c$ is

$$\begin{split} p_{\text{lin}}^{\text{M}}\left(m\right) &:= \int_{0}^{\infty} p_{\text{lin}}\left(m,\tau\right) d\tau \\ &= \frac{1-\beta}{\beta} \left(\beta z e^{-\Lambda_{\beta,z}\tau_{0}}\right)^{m} \!\! \int_{0}^{\infty} \!\! k\left(\tau\right) e^{-m\Lambda_{\beta,z}\left(\tau-\tau_{0}\right)} d\tau \end{split} \tag{S83}$$

To obtain an analytical expression for the integral $\int_0^\infty k\left(\tau\right)e^{-m\Lambda_{\beta,z}(\tau-\tau_0)}d\tau$, we use a gamma distribution $k\left(\tau\right)=\Gamma\left(\alpha\right)^{-1}\theta^{-\alpha}\left(\tau-\tau_0\right)^{\alpha-1}e^{-(\tau-\tau_0)/\theta}$. Then $\int_0^\infty k\left(\tau\right)e^{-m\Lambda_{\beta,z}(\tau-\tau_0)}d\tau=\left(1+m\Lambda_{\beta,z}\theta\right)^{-\alpha}$. Thus

$$p_{\text{lin}}^{\text{M}}(m) = \frac{1-\beta}{\beta} \left(\beta z e^{-\Lambda_{\beta,z}\tau_0}\right)^m \left(1 + m\Lambda_{\beta,z}\theta\right)^{-\alpha}. (S84)$$

For $\beta < \beta_c$, $p_{\text{lin}}^{\text{M}}(m)$ exponentially decays with typical length $(\Lambda_{\beta,z}\tau_0 - \ln \beta z)^{-1}$. As β approaches β_c , $\beta z e^{-\Lambda_{\beta,z}\tau_0}$ approaches 1 and thereby $(1 - \beta z e^{-\Lambda_{\beta,z}\tau_0})^{-1}$ diverges, meaning that the typical block size becomes much larger as β is approaching β_c . In the limit $\beta \uparrow \beta_c$, $p_{\text{lin}}^{\text{M}}(m)$ decays with $m^{-\alpha}$ for large m. For general forms of $k(\tau)$, the integral $\int_{\tau_0}^{\infty} k(\tau) e^{-m\Lambda_{\beta,z}(\tau-\tau_0)} d\tau$, for large enough m, is contributed dominantly by the values of $k(\tau)$ for τ near τ_0 . Thus if $k(\tau) \simeq A(\tau-\tau_0)^{\alpha-1}$ (A is positive constant and $\alpha > 1$), a power law $p_{\text{lin}}^{\text{M}}(m) \sim m^{-\alpha}$ holds in the limit $\beta \uparrow \beta_c$.

V. ANCESTRAL DISTRIBUTION

Let $\hat{N}(\tau; t')$ denote the expected number of cells after time t' has passed since a cell completed a cell-cycle with IDT τ . This satisfies

$$\hat{N}(\tau;t') = z \int_0^{t'} \hat{N}(\tau';t'-\tau') K(\tau',\tau) d\tau'.$$
 (S85)

In a stationary growing population, it is natural to assume $\hat{N}(\tau;t') = \phi(\tau) e^{\Lambda_{\beta,z}t'}$, where $\phi(\tau) \geq 0$. Then Eq. (S85) is rewritten as

$$\phi(\tau) = z \int_0^\infty \phi(\tau') e^{-\Lambda_{\beta,z}\tau'} K(\tau',\tau) d\tau'.$$
 (S86)

where the limit $t'\to\infty$ is taken. Since $\int_0^\infty p_{\mathrm{div}}(\tau)\,\hat{N}(\tau;t)d\tau=e^{\Lambda_{\beta,z}t}$ holds, we have

$$\int_0^\infty p_{\text{div}}(\tau) \,\phi(\tau) d\tau = 1 \,\,, \tag{S87}$$

where the quantity $p_{\rm div}\left(\tau\right)\phi(\tau)$ is the distribution of τ of the ancestor in the infinite past in a steadily growing population, which is also called the ancestral distribution [2]; this requires that $\phi(\tau)$ is an integrable function with respect to $p_{\rm div}(\tau)d\tau$ (i.e. $\phi\in L^1(p_{\rm div})$, where $p_{\rm div}$ is a measure on $[\tau_0,\infty)$). For example, if $K\left(\tau,\tau'\right)=k\left(\tau\right),\,p_{\rm div}\left(\tau\right)=ze^{-\Lambda_{\beta,z}\tau}k\left(\tau\right)$ and $\phi\left(\tau\right)=1$, therefore $p_{\rm lin}\left(\tau\right)=p_{\rm div}\left(\tau\right)$. The empirical distribution of IDT on lineages in a population converges to the ancestral distribution as $t\to\infty$ (Sec. VII). In the main text, we derived $p_{\rm lin}\left(\tau\right)$ as Eq. (12). Here, we directly compute $p_{\rm lin}\left(\tau\right)=p_{\rm div}\left(\tau\right)\phi(\tau)$.

In the Lebowitz-Rubinow model, Eq. (S86) yields

$$\phi(\tau) = \beta z e^{-\Lambda_{\beta,z}\tau} \phi(\tau) + (1 - \beta) z \int_0^\infty \phi(\tau') e^{-\Lambda_{\beta,z}\tau'} k(\tau') d\tau'.$$
 (S88)

This implies, there exists a constant $C_2 \in \mathbb{R}$

$$\phi(\tau) = \frac{C_2}{1 - \beta z e^{-\Lambda_{\beta,z}\tau}} \tag{S89}$$

for $\tau \in [\tau_0, \tau_1]$ if $\beta < \beta_c$ and for $\tau \neq \tau_*$ otherwise. Substituting Eq. (S89) to Eq. (S88), we have

$$C_2 \left(1 - Q \left(\beta, z, \Lambda_{\beta, z} \right) \right) = 0. \tag{S90}$$

If $\beta < \beta_c$, $Q(\beta, z, \Lambda_{\beta,z}) = 1$ and we find $C_2 = \overline{m}_{\beta,z}^{-1}$ by the normalization Eq. (S87).

If $\beta=\beta_c,\ Q\left(\beta,z,\Lambda_{\beta,z}\right)=1$ holds too and thus $C_2=1/\lim_{\beta\uparrow\beta_c}\overline{m}_{\beta,z}$ if $\lim_{\beta\uparrow\beta_c}\overline{m}_{\beta,z}<\infty$. If $\lim_{\beta\uparrow\beta_c}\overline{m}_{\beta,z}=\infty$, no strictly positive C_2 satisfy the normalization Eq. (S87). Additionally, $C_2=0$ leads $\int_0^\infty p_{\mathrm{div}}\left(\tau\right)\phi(\tau)d\tau=0$. Thus there is no ϕ satisfying the normalization Eq. (S87) if $\lim_{\beta\uparrow\beta_c}\overline{m}_{\beta,z}=\infty$.

If $\beta > \beta_c$, $Q(\beta, z, \Lambda_{\beta, z}) < 1$ implies $C_2 = 0$; i.e. $\phi(\tau) = 0$ for $\tau \neq \tau_*$ (τ_* is defined by Eq. (S53)). Noting for $\phi \in L^1(p_{\text{div}})$ that p_{div} includes an atomic measure at τ_0 (see Eq. S50), by the normalization condition (S87) we have $\phi(\tau_0) = C_1^{-1} = (1 - Q_0(\beta, z))^{-1}$ and $\phi(\tau > \tau_0) = 0$; equivalently, $\phi(\tau) = C_1^{-1}\delta_{\tau_0,\tau}$, where we define the Kronecker delta function $\delta_{x,y} = 1$ for x = y, and $\delta_{x,y} = 0$ for $x \neq y$.

In summary,

$$\phi(\tau) = \begin{cases} \overline{m}_{\beta,z}^{-1} \cdot \left(1 - \beta z e^{-\Lambda_{\beta,z}\tau}\right)^{-1}, & 0 \le \beta \le \beta_c \\ \left(1 - Q(\beta, z, \Lambda_{\beta,z})\right)^{-1} \cdot \delta_{\tau_0,\tau}, & \beta_c < \beta < 1 \end{cases}$$
(S91)

Multiplying $\phi(\tau)$ by $p_{\text{div}}(\tau)$, we have

$$p_{\text{lin}}(\tau) = \begin{cases} \overline{m}_{\beta,z}^{-1} \cdot \frac{(1-\beta)k(\tau)ze^{-\Lambda_{\beta,z}\tau}}{(1-\beta ze^{-\Lambda_{\beta,z}\tau})^2}, & 0 \le \beta \le \beta_c \\ \delta(\tau - \tau_0), & \beta_c < \beta < 1 \end{cases}$$

Note that ϕ and p_{lin} exist at $\beta = \beta_c$ if $\lim_{\beta \uparrow \beta_c} \overline{m}_{\beta,z} < \infty$.

VI. DERIVATIVES OF GROWTH RATE FUNCTION

For $\beta < \beta_c$, $Q(\beta, z, \Lambda_{\beta,z}) = 1$ determines $\Lambda_{\beta,z}$. Taking the first derivatives of both side of $Q(\beta, z, \Lambda_{\beta,z}) = 1$ with respect to $\ln z$, we obtain

$$\frac{\partial \Lambda_{\beta,z}}{\partial \ln z} = \overline{\tau}_{\beta,z}^{-1}.$$
 (S93)

Namely, the inverse of ancestral mean IDT $\overline{\tau}_{\beta,z}^{-1}$ is equal to the first derivative of $\Lambda_{\beta,z}$ with respect to $\ln z$. For $\beta > \beta_c$, Eq. (S93) is clear because $\overline{\tau}_{\beta,z} = \tau_0$. On the other hand, the derivative of $\Lambda_{\beta,z}$ with respect to $\ln \beta$ is

$$\frac{\partial \Lambda_{\beta,z}}{\partial \ln \beta} = \frac{1}{\overline{\tau}_{\beta,z}} \left(1 - (1 - \beta)^{-1} \overline{m}_{\beta,z}^{-1} \right). \tag{S94}$$

Eq. (S94) is also derived by taking the first derivatives of both side of $Q(\beta, z, \Lambda_{\beta,z}) = 1$ and holds as well for $\beta > \beta_c$. $\frac{\partial \Lambda_{\beta,z}}{\partial \ln \beta} > 0$ holds because for $\beta < \beta_c$,

$$\overline{m}_{\beta,z} - (1-\beta)^{-1}$$

$$= \int_{\tau_0}^{\infty} \frac{(1-\beta)ze^{-\Lambda_{\beta,z}\tau}k(\tau)}{(1-\beta ze^{-\Lambda\tau})^2} d\tau - \int_{\tau_0}^{\infty} \frac{k(\tau)}{1-\beta} d\tau$$

$$= \int_{\tau_0}^{\infty} \left(\frac{(1-\beta)ze^{-\Lambda_{\beta,z}\tau}}{1-\beta ze^{-\Lambda_{\beta,z}\tau}} - 1\right) \frac{1-\beta^2 ze^{-\Lambda_{\beta,z}\tau}}{1-\beta ze^{-\Lambda_{\beta,z}\tau}} \frac{k(\tau)}{1-\beta} d\tau$$

$$> \int_{\tau_0}^{\infty} \left(\frac{(1-\beta)ze^{-\Lambda_{\beta,z}\tau}}{1-\beta ze^{-\Lambda_{\beta,z}\tau}} - 1\right) \frac{k(\tau)}{1-\beta} d\tau = 0. \tag{S95}$$

Together, $\Lambda_{\beta,z}$ increases as β or z increases. Since $\overline{m}_{\beta,1} = (1-\beta)^{-1}$ holds, this inequality implies that the mean cluster size on lineage is always greater in growing population (z > 1) rather than in isolation.

VII. THERMODYNAMIC ANALOGY FOR LINEAGE STRUCTURES IN THE LEBOWITZ-RUBINOW MODEL

Here we show that the population structure of lineages in the Lebowitz-Rubinow model is mapped to a statistical mechanical ensemble, which yields the thermodynamic interpretation of population growth rate and other quantities. The results hold for general forms of the transition kernel $K(\tau, \tau')$.

Let $\Omega := \{(a,\tau) \mid 0 \le a \le \tau; \tau_0 \le \tau < \infty\}$ be the domain for age a and IDT τ . The initial condition is given by $n(a,\tau;0) = N_0 p_0(a,\tau)$ where N_0 is the initial population size and p_0 is a probability distribution on Ω . Let $n^{(D)}(a,\tau;t) da d\tau$ denote the expected number of offspring in the interval $(a,a+da) \times (\tau,\tau+d\tau)$ at time t after D divisions. Recursively using Eq. (S2) and Eq. (S3), we have

$$n(a, \tau; t) = N_0 \sum_{D=0}^{\infty} z^D P^{(D)}(a, \tau, t)$$
 (S96)

where $P^{(D)}\left(a,\tau;t\right)$ denotes the probability distribution of (a,τ) at time t after D divisions in isolation and the expected population size at time t is

$$N_t = \iint_{\Omega} n\left(a, \tau; t\right) da \, d\tau = N_0 \sum_{D=0}^{\infty} z^D P_{\text{iso}}\left(D; t\right) \quad (S97)$$

where

$$P_{\text{iso}}(D;t) := \iint_{\Omega} P^{(D)}(a,\tau,t) \, da \, d\tau \tag{S98}$$

denotes the probability distribution of number of divisions along lineages in isolation. The probability distribution of number of divisions along lineage in population is

$$P_{\text{pop}}(D;t) := \frac{\iint_{\Omega} n^{(D)}(a,\tau;t) \, da \, d\tau}{N_t} = \frac{N_0}{N_t} z^D P_{\text{iso}}(D;t)$$
(S99)

 P_{iso} and P_{pop} are referred to as "chronological" and "retrospective" distribution respectively in [3].

A lineage can be identified as the sequences (m_0, \dots, m_S) and $(x_0, \dots, x_{S-1}, \tau)$ where m_j denotes the j-th block size with IDT x_j $(x_S = \tau)$. S denotes the number of switching to the other IDT and $D = m_0 + \dots + m_S - 1$ denotes the number of divisions on the lineage. Then, the joint probability distribution of number of divisions D and number of switches S on a lineage in isolation can be defined as

$$P_{\rm iso}\left(D,S;t\right) := \iint_{\Omega} P^{(D,S)}\left(a,\tau,t\right) da \, d\tau \qquad (S100)$$

where $P^{(D,S)}\left(a,\tau;t\right)$ denotes probability density of (a,τ) at time t after D divisions and S switches in isolation. Therefore

$$\frac{N_t}{N_0} = \sum_{D=0}^{\infty} \sum_{S=0}^{D} z^D P_{\text{iso}}(D, S; t)$$
 (S101)

holds and it is reasonable to define

$$P_{\text{pop}}(D, S; t) := \frac{N_0}{N_t} z^D P_{\text{iso}}(D, S; t)$$
 (S102)

It is noticeable that $P_{\text{iso}}\left(D,S;t\right)$ depends on β in the form of $\left(1-\beta\right)^{S}\beta^{D-S}$.

In this lineage formulation, the long-term growth rate $\Lambda_{\beta,z}$ can be interpreted as the cumulant generating function scaled by time. Let D denote the number of divisions on lineage and S denote the number of switching to the other τ on lineage. Let $\Phi_{\beta}(\xi, \eta; t)$ be the cumulant generating function of D and S:

$$\Phi_{\beta}\left(\xi, \eta; t\right) := \ln \sum_{D, S} P_{\text{iso}}\left(D, S; t\right) e^{\xi D + \eta S}. \tag{S103}$$

 $\Phi_{\beta}(\xi, \eta)$ depends on β via $(1 - \beta)^S \beta^{D-S}$ in $P_{\text{iso}}(D, S; t)$. The biased probability distribution of (D, S) is defined by

$$P_{\xi,\eta}(D,S;t) := P_{\text{iso}}(D,S;t) e^{\xi D + \eta S} e^{-\Phi_{\beta}(\xi,\eta;t)}, \text{ (S104)}$$

Note that $\Phi_{\beta}(\ln z, 0) = \ln^{N_t}/N_0$ and $P_{\ln z, 0}(D, S; t) = P_{\text{pop}}(D, S; t)$. In particular, $\Phi_{\beta}(\ln z, 0) = t\Lambda_{\beta, z}$ holds at steady state. Below we assume steady state, that is, the initial distribution p_0 the stationary solution. Let \hat{D}_t and \hat{S}_t denote the random variables taking values D and S along lineage with time length t and let $\langle \cdot \rangle_{\text{pop}}$ denote the expectation with respect to $P_{\text{pop}}(D, S; t)$. For any positive integer j,

$$\frac{\partial^{j} \Lambda_{\beta,z}}{(\partial \ln z)^{j}} = \left. \frac{1}{t} \frac{\partial^{j} \Phi_{\beta} (\xi, 0)}{\partial \xi^{j}} \right|_{\xi = \ln z}$$
 (S105)

holds. In particular, for j = 1,

$$\frac{\partial \Lambda_{\beta,z}}{\partial \ln z} = \frac{1}{t} \frac{\partial \Phi_{\beta}(\xi,0)}{\partial \xi} \bigg|_{\xi = \ln z} = \left\langle \frac{\hat{D}_t}{t} \right\rangle_{\text{pop}} \tag{S106}$$

Together with Eq. (S93), we have

$$\left\langle \frac{\hat{D}_t}{t} \right\rangle_{\text{pop}} = \left\langle \tau \right\rangle_{\text{anc}}^{-1}$$
 (S107)

For j=2,

$$\frac{\partial^{2} \Lambda \left(\beta, z\right)}{\left(\partial \ln z\right)^{2}} = \frac{1}{t} \frac{\partial^{2} \Phi_{\beta} \left(\xi, 0\right)}{\partial \xi^{2}} \bigg|_{\xi = \ln z} = \frac{1}{t} \operatorname{Var} \left[\hat{D}_{t}\right]_{\text{pop}} \tag{S108}$$

holds, where $\operatorname{Var}\left[\hat{D}_{t}\right]_{\operatorname{pop}} = \left\langle \hat{D}_{t}^{2}\right\rangle_{\operatorname{pop}} - \left\langle \hat{D}_{t}\right\rangle_{\operatorname{pop}}^{2}$. If $\beta > \beta_{c}$, $\Lambda_{\beta,z} = \tau_{0}^{-1} \ln{(\beta z)}$ implies $t^{-1}\operatorname{Var}\left[\hat{D}_{t}\right]_{\operatorname{pop}}^{2} = 0$ holds. In terms of law of large number, a random variable \hat{D}_{t}/t converges to $\left\langle \tau \right\rangle_{\operatorname{anc}}^{-1}$ as $t \to \infty$ with respect to $P_{\operatorname{pop}}\left(D,S;t\right)$. We mean this convergence in probability P_{pop} by $\operatorname{plim}_{t\to\infty}\hat{D}_{t}/t = \left\langle \tau \right\rangle_{\operatorname{anc}}^{-1}$ ("plim" for probability limit). In addition, the derivative of $\Lambda_{\beta,z}$ with respect to $\ln \beta$ can be expressed by derivatives of $\Phi_{\beta}\left(\xi,\eta\right)$ with respect to ξ and η . For a small $\epsilon > 0$,

$$(1 - \beta e^{\epsilon})^{S} (\beta e^{\epsilon})^{D-S} = (1 - \beta)^{S} \beta^{D-S} e^{\epsilon D - \frac{\epsilon S}{1-\beta}} + \mathcal{O}(\epsilon^{2}).$$
(S109)

That is, $\Phi_{\beta e^{\epsilon}}(\xi, \eta) = \Phi_{\beta}\left(\xi + \epsilon, \eta - \frac{\epsilon}{1-\beta}\right) + \mathcal{O}\left(\epsilon^{2}\right)$ and thereby

$$\frac{\partial \Phi_{\beta}\left(\xi,\eta\right)}{\partial \ln \beta} = \left(\frac{\partial}{\partial \xi} - \frac{1}{1-\beta} \frac{\partial}{\partial \eta}\right) \Phi_{\beta}\left(\xi,\eta\right)$$
 (S110)

holds. Therefore, using the relation

$$\left. \frac{\partial \Phi_{\beta} \left(\ln z, \eta \right)}{\partial \eta} \right|_{\eta=0} = \left\langle \frac{\hat{S}_t}{t} \right\rangle_{\text{pop}} \tag{S111}$$

we have

$$\frac{\partial \Lambda (\beta, z)}{\partial \ln \beta} = \frac{\partial \Phi_{\beta} (\xi, 0)}{\partial \xi} \bigg|_{\xi = \ln z} - \frac{1}{1 - \beta} \frac{\partial \Phi_{\beta} (\ln z, \eta)}{\partial \eta} \bigg|_{\eta = 0}$$

$$= \left\langle \frac{\hat{D}_t}{t} \right\rangle_{\text{pop}} - \frac{1}{1 - \beta} \left\langle \frac{\hat{S}_t}{t} \right\rangle_{\text{pop}} \tag{S112}$$

Comparing it to Eq. (S94), we have

$$\left\langle \frac{\hat{S}_t}{t} \right\rangle_{\text{pop}} = \overline{m}_{\beta,z}^{-1} \overline{\tau}_{\beta,z}^{-1} \tag{S113}$$

In terms of law of large number,

$$\left\langle \frac{\hat{S}_t}{t} \right\rangle_{\text{pop}} = \text{plim}_{t \to \infty} \frac{\hat{S}_t}{t}$$

$$= \text{plim}_{t \to \infty} \frac{\hat{S}_t}{\hat{D}_t} \cdot \text{plim}_{t \to \infty} \frac{\hat{D}_t}{t}$$

$$= \overline{m}_{\beta,z}^{-1} \cdot \overline{\tau}_{\beta,z}^{-1}$$
(S114)

implies $\operatorname{plim}_{t\to\infty} \hat{S}_t/\hat{D}_t = \overline{m}_{\beta,z}^{-1}$, which is reasonable in that $\overline{m}_{\beta,z}$ represents the mean cluster size on lineage in population.

The cumulant generating function $t \cdot \Phi_{\beta}(\xi, \eta)$ plays the role of free energy in thermodynamics. A lineage is analogous to a specific configuration of a macroscopic system. The total time t, denoting the lineage length, corresponds to the physical size of the system, i.e. its volume. $\langle \hat{D}_t \rangle_{\rm pop}$ and $\langle \hat{S}_t \rangle_{\rm pop}$ are extensive variables, like energy and entropy in thermodynamics, as they are typically proportional to t. $\xi (= \ln z)$ and η are the conjugate variables to $\langle \hat{D}_t \rangle_{\rm pop}$ and $\langle \hat{S}_t \rangle_{\rm pop}$; and as in thermodynamics, the first derivative of free energy with respect to an intensive variable becomes its conjugate extensive variable. In this point of view, a discontinuous transition of $\overline{\tau}_{\beta,z}^{-1}$ at $\beta = \beta_c$ is a first-order phase transition.

VIII. NUMERICAL SIMULATIONS

A. IDT distribution in isolation

In simulations, $k(\tau)$ is chosen as

$$k(\tau) = \begin{cases} \frac{(\tau - \tau_0)^{\alpha - 1}}{\Gamma(\alpha)\theta^{\alpha}} e^{-(\tau - \tau_0)/\theta}, & \tau \ge \tau_0\\ 0, & otherwise \end{cases}$$
(S115)

To fix division time scale, we always set the mean interdivision time in isolation to 1: $\int_0^\infty \tau k(\tau) d\tau = \tau_0 + \alpha \theta =$ 1.

B. Growth rate

To sample (a, τ) for initial conditions, we need to compute $\Lambda_{\beta,z}$. For $\beta=0$, $\Lambda_{0,z}$ is computed using Lambert W function W(x): W(x) satisfies W(x) $e^{W(x)}=x$ and $W(x) \geq -1$ with its domain $[-e^{-1}, \infty)$. Since $\Lambda_{0,z}$ sat-

isfies

$$1 = \int_{0}^{\infty} z e^{-\lambda \tau} k(\tau) d\tau = \frac{z e^{-\tau_0 \Lambda_{0,z}}}{(1 + \theta \Lambda_{0,z})^{\alpha}}$$
$$= \left(\frac{\frac{\tau_0}{\alpha \theta} e^{\tau_0/(\alpha \theta)} z^{1/\alpha}}{\frac{\tau_0}{\alpha} (\Lambda_{0,z} + \theta^{-1}) \exp\left(\frac{\tau_0}{\alpha} (\Lambda_{0,z} + \theta^{-1})\right)}\right)^{\alpha}$$
(S116)

Thus we get

$$\Lambda_{0,z} = \frac{\alpha}{\tau_0} W \left(\frac{\tau_0}{\alpha \theta} e^{\tau_0/(\alpha \theta)} z^{1/\alpha} \right) - \frac{1}{\theta}$$
 (S117)

When $\beta > 0$, we numerically solve the integral equation

$$\int_{0}^{\infty} \frac{(1-\beta)ze^{-\lambda\tau}k(\tau)}{1-\beta ze^{-\lambda\tau}}d\tau = 1$$
 (S118)

in the range of $\lambda > \max(0, \tau_0^{-1} \ln(\beta z))$ to get $\Lambda_{\beta,z}$, and if there is no solution in that range, we set $\Lambda_{\beta,z} = \tau_0^{-1} \ln(\beta z)$.

C. Initial conditions

Let $\Omega := \{(a, \tau) | 0 \le a \le \tau; \tau_0 \le \tau < \infty\}$ be the domain for age a and IDT τ . The stationary joint probability distribution of (a, τ) is given by Eq (S13). For LR model, this becomes

$$\varphi_{\beta,z}(a,\tau) := \frac{z}{z-1} \Lambda_{\beta,z} e^{-\Lambda_{\beta,z}a} \times ((1 - G_{\beta,z}) \delta(\tau - \tau_0) + g_{\beta,z}(\tau)) \quad (S119)$$

where

$$g_{\beta,z}(\tau) = \frac{(1-\beta)k(\tau)}{1-\beta z e^{-\Lambda_{\beta,z}\tau}}$$
 (S120)

and

$$G_{\beta,z} = \int_{0}^{\infty} g_{\beta,z}(\tau) d\tau \qquad (S121)$$

Noting that $\varphi_{\beta,z}(a,\tau)$ is proportional to the product of $\Lambda_{\beta,z}e^{-\Lambda_{\beta,z}a}$ and of $(1-G_{\beta,z})\,\delta\,(\tau-\tau_0)+g_{\beta,z}\,(\tau)$, each of which is a univariate probability distribution, sampling (a,τ) from $\varphi_{\beta,z}$ is done as follows. First, generate a random number \hat{a} , which is exponentially distributed with mean $\Lambda_{0,z}^{-1}$. Next generate a random number τ , which follows $(1-G_{\beta,z})\,\delta\,(\tau-\tau_0)+g_{\beta,z}\,(\tau)$. To do so, set $\tau=\tau_0$ with probability $1-G_{\beta,z}$. Otherwise, generate uniformly distributed (x,y) in sufficiently large rectangular region in \mathbb{R}^2 and set $\tau=x$ if $g_{\beta,z}(x)>y$.

In the numerical simulations described in the main text, we set 2 different initial conditions. We mean by delocalized state that (a,τ) is sampled from $\varphi_{0,z}(a,\tau)$ at time 0. Each (a,τ) among N cells is independent of each other. The other condition is predicted stationary, for which (a,τ) is sampled from $\varphi_{\beta,z}(a,\tau)$ at time 0, where β is the same number as the one used in time-evolution.

D. Simulation

Fix an integer N>0 for population size. Fix a real number T>0 for the simulation time. The offspring number per cell is set to z=2. Let \hat{t}_i denote the time of i-th division in the population and let B_i denote the number of divisions at time \hat{t}_i . To keep the population size constant, B_i cells among $N+B_i$ are randomly excluded. Let τ_{ij} denote the IDT of the j-th dividing cell among B_i cells. Let Δt be the length of the time window within which the growth rate is averaged and the locally minimum generation time is determined. Let $t_n:=n\Delta t$, $n=0,1,\ldots$ Then the empirical growth rate

$$\Lambda^{(n)} := \frac{1}{\Delta t} \sum_{i:t_{n-1} < \hat{t}_i < t_n} \ln\left(1 + \frac{B_i}{N}\right)$$
 (S122)

and the empirical minimum IDT

$$\hat{\tau}_0^{(n)} := \min_{i: t_{n-1} \le \hat{t}_i < t_n} \min_{1 \le j \le B_i} \tau_{ij}$$
 (S123)

are computed. We also record all the lineages that remain by the end of the simulation. We validated that N individuals at time T coalesce to a single ancestor. In other words, the simulation time T is taken much longer than the coalescence time for N individuals.

- [1] R. Bürger and I. M. Bomze, Adv. Appl. Probab. **28**, 227 (1996).
- [2] J. Hermisson, O. Redner, H. Wagner, and E. Baake, Theor. Popul. Biol. 62, 9 (2002).
- [3] T. Nozoe, E. Kussell, and Y. Wakamoto, PLOS Genet. 13, e1006653 (2017).
- [4] N. Mosheiff, B. M. C. Martins, S. Pearl-Mizrahi, A. Grünberger, S. Helfrich, I. Mihalcescu, D. Kohlheyer, J. C. W. Locke, L. Glass, and N. Q. Balaban, Phys. Rev. X 8, 021035 (2018).
- [5] H. E. Kubitschek, Exp. Cell Res. 26, 439 (1962).
- [6] M. Hashimoto, T. Nozoe, H. Nakaoka, R. Okura, S. Akiyoshi, K. Kaneko, E. Kussell, and Y. Wakamoto, Proc. Natl. Acad. Sci. 113, 3251 (2016).
- [7] F. B. Yu, L. Willis, R. M. W. Chau, A. Zambon, M. Horowitz, D. Bhaya, K. C. Huang, and S. R. Quake, BMC Biol. 15, 1 (2017).
- [8] E. D. Hawkins, J. F. Markham, L. P. McGuinness, and P. D. Hodgkin, Proc. Natl. Acad. Sci. 106, 13457 (2009).
- [9] H. Sasaki, H. Yoshinaga, and K. Kawano,

IX. N-DEPENDENCE OF POPULATION GROWTH AND DEFINITION OF $\beta_{c,N}$

To observe the population size dependence of population growth and lineage quantities, we computed empirical counterparts for $\Lambda_{\beta,z}$, $\overline{\tau}_{\beta,z}^{-1}$ and $\overline{m}_{\beta,z}^{-1}$ with varied population size N and with fixed simulation time T. As the counterpart of $\Lambda_{\beta,z}$ in a finite population, time-average of $\Lambda^{(n)}$ is taken, denoted by $\Lambda_{\beta,z,N}$. To compute the counterparts of $\overline{\tau}_{\beta,z}^{-1}$ and $\overline{m}_{\beta,z}$, a single-cell lineage is randomly chosen and we let D and S be the number of divisions and number of switching events to a different IDT, respectively. As the counterpart of $\overline{\tau}_{\beta,z,N}^{-1}$ in finite population, D/T is taken, denoted by $\overline{\tau}_{\beta,z,N}^{-1}$. As the counterpart of $\overline{m}_{\beta,z,N}^{-1}$ in finite population, S/D is taken, denoted by $\overline{m}_{\beta,z,N}^{-1}$. For each empirical evaluation, the mean and standard deviation are calculated over 10 simulations

We denote by $\beta_{c,N}$ the critical heritability of IDT at finite population size N. This is defined as β which maximizes

$$\overline{\tau}_{\beta,z,N}^{-1} \ln z - \Lambda_{\beta,z,N} . \tag{S124}$$

If $N = \infty$, this is discontinuous at β_c and

$$\lim_{\beta \downarrow \beta_c} \left(\tau_{\beta,z}^{-1} \ln z - \Lambda_{\beta,z} \right) = -\tau_0^{-1} \ln \beta_c$$
 (S125)

is the supreme of $\tau_{\beta,z}^{-1} \ln z - \Lambda_{\beta,z}$ for $\beta \in [0,1)$. For finite N, this function becomes smoother around β_c and have a peak at $\beta_{c,N} > \beta_c$. Still, the result can depend on initial state due to aging dynamics. Thus introducing small noise in inheritance will allow convergence over a feasible time-scale (Figs S6 and S7).

- Radiat. Res. **72**, 364 (1977).
- [10] R. G. Staudte, M. Guiguet, and M. C. D'Hooghe, J. Theor. Biol. 109, 127 (1984).
- [11] M. Hola and P. A. Riley, J. Cell Sci. 88 (Pt 1), 73 (1987).
- [12] S. Chakrabarti, A. L. Paek, J. Reyes, K. A. Lasick, G. Lahav, and F. Michor, Nat. Commun. 9, 1 (2018).
- [13] G. Froese, Exp. Cell Res. **35**, 415 (1964).
- [14] R. Van Wijk and K. W. Van De Poll, Cell Prolif. 12, 659 (1979).
- [15] M. Absher, D. Sylwester, and B. A. Hart, Environ. Res. 30, 34 (1983).
- [16] M. Absher and V. J. Cristofalo, J. Cell. Physiol. 119, 315 (1984).
- [17] O. Sandler, S. P. Mizrahi, N. Weiss, O. Agam, I. Simon, and N. Q. Balaban, Nature 519, 468 (2015).
- [18] H. Miyamoto, E. Zeuthen, and L. Rasmussen, J. Cell Sci. 13, 879 (1973).
- [19] A. van Meeteren, R. van Wijk, J. Stap, and B. F. Deys, Cell Prolif. 17, 105 (1984).
- [20] J. Grasman, Bull. Math. Biol. 52, 535 (1990).

- [21] R. Shields and J. A. Smith, J. Cell. Physiol. **91**, 345 (1977).
- [22] R. G. Staudte, R. M. Huggins, J. Zhang, D. E. Axelrod, and M. Kimmel, Math. Biosci. 143, 103 (1997).
- [23] R. Sennerstam and J. O. Strömberg, J. Theor. Biol. 131, 151 (1988).
- [24] K. B. Dawson, H. Madoc-Jones, and E. O. Field, Exp. Cell Res. 38, 75 (1965).

TABLE S1. List of mother-daughter correlation coefficients of IDTs. For the same cell type in the same reference, brief summary of the acquisition conditions of the data is described in the "Data set" column. "Correlation type" indicates Spearman rank correlation or Pearson correlation. "*" denotes the definition of correlation coefficient is not explicitly described in the reference. "N" denotes the sample size. "M-D corr." denotes the mother-daughter correlation of IDTs.

	Cell type	Data set	Correlation type	N	M-D corr.	Error range	Ref
	Cornebacterium	n/a	Spearman	51	-0.21	-0.21 ± 0.12	[4]
	glutamicum	,	G	6.0	0.00	0.001.005	F . 3
	E.coli B Rel606	n/a	Spearman	60	-0.06	-0.06 ± 0.25	[4]
	E.coli B/r	n/a	Pearson	160	0.11	n/a	[5]
	E.coli B/r ΔfimAΔflu	M9 glucose, 37C	Pearson	n/a	-0.18	-0.18 ± 0.03	[6]
	$^{ m rpsL-gfp}_{,,}$	M9 glycerol, 37C	Pearson	n/a	0.02	0.02 ± 0.02	[6]
	E.coli W3110	M9 glucose, 37C	Pearson	n/a	0.02 0.11	0.02 ± 0.02 0.11 ± 0.03	[6]
Bacteria	Δ fliC Δ fim $A\Delta$ flu LVS	WIS glucose, STC	1 carson	11/α	0.11	0.11±0.05	ران
	E.coli W3110	M9 cas. Acids, 37C	Pearson	n/a	-0.06	-0.06 ± 0.03	[6]
	Δ fliC Δ fim $A\Delta$ flu rpsL-gfp			/	0.00	0.00_0.00	[-]
	,,	M9 glucose, 30C	Pearson	n/a	0.13	0.13 ± 0.03	[6]
	"	M9 glucose, 37C	Pearson	n/a	0.26	$0.26 {\pm} 0.02$	[6]
	"	M9 glycerol, 30C	Pearson	n/a	-0.08	-0.08 ± 0.03	[6]
	"	M9 glycerol, 37C	Pearson	n/a	0.08	0.08 ± 0.03	[6]
	"	M9 LB, 37C	Pearson	n/a	-0.06	-0.06 ± 0.04	[6]
	E.coli W3110	M9 glucose, 30C	Pearson	n/a	0.02	0.02 ± 0.03	[6]
	Δ fliC Δ fim $A\Delta$ flu T7-venus	3.60 3		,	0		F = 7
	"	M9 glucose, 37C	Pearson	n/a	0.08	0.08 ± 0.02	[6]
	Synechococcus elongatus	n/a	Spearman	65	-0.16	-0.16 ± 0.08	[4]
	WT	m / a	C	74	0.17	0.1710.00	[4]
	Synechococcus elongatus Δ kaiBC	n/a	Spearman	74	-0.17	-0.17 ± 0.08	[4]
	Synechocystis sp. PCC6803	n/a	Pearson*	n/a	-0.1	n/a	[7]
	B lymphocyte	Fam2	Spearman	n/a	0.66	n/a	[8]
	n iy in phocy te	Fam3	Spearman	n/a	0.658	n/a n/a	[8]
	СЗН	n/a	Pearson*	144	-0.09	n/a	[9]
	EMT6	1st generation	Spearman	66	0.46	n/a	[10]
	"	2nd generation	Spearman	106	0.63	n/a	[10]
	"	3rd generation	Spearman	131	0.64	n/a	[10]
	"	4th generation	Spearman	37	0.14	n/a	[10]
	GPK	n/a	Pearson*	40	0.27	n/a	[11]
	HCT116 p53V-KI	n/a	Pearson	71	-0.03	[-0.26, 0.16] (95%CI)	[12]
	HeLa	series case II, $\tau_d < 15$	Pearson*	n/a	0.57	n/a	[13]
	"	series case II,	Pearson*	n/a	-0.79	n/a	[13]
	"	$15 < \tau_d < 25$ series case II, $\tau_d > 25$	Pearson*	n/a	0.19	'	[13]
	HelaS	series case ii, $\tau_d > 25$	Pearson*	$_{ m n/a}$	$0.19 \\ 0.12$	$_{ m n/a}$	[13]
Mammalian	HTC	n/a	Pearson*	n/a	$0.12 \\ 0.55$	n/a n/a	[14]
	human embryonic lung	n/a n/a	Pearson*	28	-0.245	n/a n/a	[15]
	fibroblasts	/			5.210	/ 😅	[[-]
	IMR-90	young	Pearson*	41	0.1	n/a	[16]
	"	$\stackrel{\circ}{\operatorname{aged}}$	Pearson*	14	0.31	n/a	[16]
	L1210	closed system	Spearman	432	0.04	0.04 ± 0.08	[17]
	"	constant media flow	Spearman	381	0.3	0.3 ± 0.03	[17]
	mouse L-cells	n/a	Pearson*	100	0.648	n/a	[18]
	mouse osteosarcoma	1st generation	Pearson*	133	0.33	n/a	[19]
	"	2nd generation	Pearson*	122	0.28	n/a	[19]
	"	n/a	Pearson*	117	0.52	n/a	[20]
	neuroblastoma	n/a	Pearson*	60	0.61	n/a	[20]
	NIH3T3	n/a	Pearson*	n/a	0.06	n/a	[21]
	"	rich medium	Pearson*	141	-0.418	n/a	[22]
		poor medium	Pearson*	166	-0.241 0.41	n/a	[22]
	PCC3 PCC4 azal	$ \frac{n/a}{n/a} $	Pearson*	100 84	$0.41 \\ 0.07$	n/a	[23]
	rat S6 sarcoma	n/a n/a	Pearson* Pearson*	n/a	0.07 0.194	n/a n/a	[23] [24]
<u> </u>	Tat DU SatCOIIIà	п/а	1 carson	11/a	0.134	11/а	[24]

TABLE S2. Classification heritability threshold of LR model. First column denotes the cases explained in Sec. III A. Second column shows the type of heritability threshold. Third to fifth columns show each limiting case. n/a implies it is the same as the one in the second column.

	$0 < \tau_0 < \tau_1 < \infty$	ω_0	$\tau_0 \downarrow 0$	$\tau_1 \uparrow \infty$	$k(\tau) \to \delta(\tau - \bar{\tau})$
z > 1 (case I)	Localize to τ_0 above $\beta_c (> z^{-1})^a$	$\ln(\beta z)/\tau_0$	$\beta_c \downarrow z^{-1}$	n/a	$\beta_c = z^{\tau_0/\bar{\tau} - 1}$
z = 1 (case II)	No threshold $(\beta_c = 1)$	0	n/a	n/a	$\beta_c = 1$
0 < z < 1 (case III)	Localize to τ_1 above $\beta_c (> 0)$ b	$\ln(\beta z)/\tau_1$	n/a	$\beta_c \downarrow 0$	$\beta_c = z^{\tau_1/\bar{\tau} - 1}$

a $\beta_c < 1$ when $\int_{\tau_0}^{\tau_1} d\tau^k (\tau) / (\tau - \tau_0) < \infty$ b $\beta_c < 1$ when $\int_{\tau_0}^{\tau_1} d\tau^k (\tau) / (\tau_1 - \tau) < \infty$




FIG. S1. Synchronization of cell-cycles without inheritance noise ($\sigma = 0$ in Eq. (13)), starting from predicted stationary. Except for the initial condition, parameters used for the simulation are the same as Fig. 3 in the main text. Captions for each figure are the same with those in Fig. 3.




FIG. S2. The effect of finite population size N on population growth rate and lineage quantities, starting from delocalized. Simulation time t=10000. The parameters are the same with $\alpha=4$ in Fig. 1. (A) Population growth rate. (B) The frequency of divisions per unit time on lineage. (C) The frequency of switching to the other τ per generation on lineage. (D) Results in (B) multiplied by $\ln 2$ subtracted by results in (A).




FIG. S3. Phase transition across population size N, starting from delocalized state. The same results with Fig. S2 are used for these figures. (A) Threshold of β in finite population of fixed size, computed as the position of peak in Fig S2. (B-D) Rearrangements of Fig S2A-C against N with the certain range of β around transition point.




FIG. S4. The effect of finite population size N on population growth rate and lineage quantities, starting from predicted stationary. Captions for each figure are the same with those in Fig. S2.




FIG. S5. Phase transition across population size N, starting from predicted stationary. The same results with Fig. S4 are used for these figures. Captions for each figure are the same with those in Fig. S3.




FIG. S6. Disappearing aging dynamics with 1% inheritance noise ($\sigma = 0.01\tau_0$ in Eq. (13)). The parameters are as in Fig. 1 using $\alpha = 4$, which exhibits a first-order localization transition. See the caption of Fig. 2 for the description of figures.




FIG. S7. Disappearing aging dynamics with 10% inheritance noise ($\sigma = 0.1\tau_0$ in Eq. (13)). The parameters are as in Fig. 1 using $\alpha = 4$, which exhibits a first-order localization transition. See the caption of Fig. 2 for the description of figures.




FIG. S8. The effect of finite population size N on population growth rate and lineage quantities, with 1% inheritance noise $(\sigma = 0.01\tau_0)$ in Eq. (13), starting from delocalized. Simulation time t = 10000. The parameters are the same with $\alpha = 4$ in Fig. 1. Captions for each figure are the same with those in Fig. S2.




FIG. S9. Phase transition across population size N, with 1% inheritance noise, starting from delocalized. The same results with Fig. S8 are used for these figures. Captions for each figure are the same with those in Fig. S3.




FIG. S10. The effect of finite population size N on population growth rate and lineage quantities, with 1% inheritance noise $(\sigma = 0.01\tau_0)$ in Eq. (13), starting from predicted stationary. Simulation time t = 10000. The parameters are the same with $\alpha = 4$ in Fig. 1. Captions for each figure are the same with those in Fig. S2.




FIG. S11. Phase transition across population size N, with 1% inheritance noise, starting from predicted stationary. The same results with Fig. S10 are used for these figures. Captions for each figure are the same with those in Fig. S3.




FIG. S12. The effect of finite population size N on population growth rate and lineage quantities, with 10% inheritance noise $(\sigma = 0.1\tau_0 \text{ in Eq. (13)})$, starting from delocalized. Simulation time t = 10000. The parameters are the same with $\alpha = 4$ in Fig. 1. Captions for each figure are the same with those in Fig. S2.




FIG. S13. Phase transition across population size N, with 10% inheritance noise, starting from delocalized. The same results with Fig. S12 are used for these figures. Captions for each figure are the same with those in Fig. S3.




FIG. S14. The effect of finite population size N on population growth rate and lineage quantities, with 10% inheritance noise $(\sigma = 0.1\tau_0)$ in Eq. (13)), starting from predicted stationary. Simulation time t = 10000. The parameters are the same with $\alpha = 4$ in Fig. 1. Captions for each figure are the same with those in Fig. S2.




FIG. S15. Phase transition across population size N, with 10% inheritance noise, starting from predicted stationary. The same results with Fig. S14 are used for these figures. Captions for each figure are the same with those in Fig. S3.




FIG. S16. Synchronization of cell-cycles with 1% inheritance noise ($\sigma = 0.01\tau_0$ in Eq. (13)), starting from delocalized. The parameters are the same with $\alpha = 4$ in Fig. 1. N = 100. Simulation time t = 10000. Captions for each figure are the same with those in Fig. 3.




FIG. S17. Synchronization of cell-cycles with 10% inheritance noise ($\sigma=0.1\tau_0$ in Eq. (13)), starting from delocalized. The parameters are the same with $\alpha=4$ in Fig. 1. N=100. Simulation time t=10000. Captions for each figure are the same with those in Fig. 3.