

Boundedness of Singular Integral Operators on Weak Herz Type Spaces with Variable Exponent

Hongbin WANG* and Zongguang LIU

Abstract In this paper, the authors define the weak Herz spaces and the weak Herz-type Hardy spaces with variable exponent. As applications, the authors establish the boundedness for a large class of singular integral operators including some critical cases.

Key Words weak Herz spaces; weak Herz-type Hardy spaces; variable exponent; boundedness; singular integral operators.

MR(2010)Subject Classification 42B35; 42B20; 46E30.

1 Introduction

The main purpose of this article is to introduce the weak Herz type spaces with variable exponent and give the boundedness for a large class of singular integral operators. The classical weak Herz type spaces can be traced to the work of Hu, Lu and Yang ([7,8]) on the study of the boundedness for some operators. The classical weak Herz type spaces mainly include the weak Herz spaces and the weak Herz-type Hardy spaces. As well known, many important operators in harmonic analysis, such as Calderón-Zygmund operators, Ricci-Stein oscillatory singular integral operators, are not bounded on $L^1(\mathbb{R}^n)$, but they map $L^1(\mathbb{R}^n)$ to weak $L^1(\mathbb{R}^n)$. So the weak Lebesgue space $WL^p(\mathbb{R}^n)$ are very important in operator theory. The weak Herz spaces are the analog of $WL^p(\mathbb{R}^n)$ in the setting of Herz spaces. Meanwhile, the weak Herz-type Hardy spaces are the suitable local version of the weak Hardy spaces $WH^p(\mathbb{R}^n)$. The classical weak Herz type spaces and related function spaces have interesting applications in studying boundedness of many operators and the theory of partial differential equations; see, for example [5,10,12,14,16].

On the other hand, due to their wide applications in partial differential equations with non-standard growth ([6]), electrorheological fluids ([13]) and image processing ([1]), the theory of function spaces with variable exponents have attracted a lot of attentions in recent years. Particularly, such theory have achieved great progresses after the notable work of Kováčik and Rákosník [11] in 1991; see [3], [4] and the references therein. In 2010, Izuki [9] studied the Herz spaces with variable exponent $\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ ($K_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$) and proved the boundedness of some sublinear operators on the spaces. In addition, Wang and Liu [15] introduced a certain Herz-type Hardy spaces with variable exponent $H\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ ($HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$) in 2012. Recently, Yan et al. introduced the variable weak Hardy spaces $WH^{p(\cdot)}(\mathbb{R}^n)$ and established some characterizations and the boundedness of some operators in [17].

Motivated by [8,9,15,17], we aim to develop the weak Herz spaces and the weak Herz-type Hardy spaces in the setting of variable exponents. In Section 2, we first briefly recall some standard notations and lemmas in variable function spaces. Then we will define the weak Herz spaces

*Corresponding author.

and the weak Herz-type Hardy spaces with variable exponent. In Section 3, firstly we will prove the boundedness for a class of sublinear operators from $\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ (or $K_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$) into the weak Herz spaces with variable exponent $W\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ (or $WK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$). Then we will establish the boundedness of local Calderón-Zygmund type operators from $H\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ (or $HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$) into $\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ (or $K_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$) and from $H\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ (or $HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$) into $W\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ (or $WK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$) at the critical index. Subsequently, we will show the standard Calderón-Zygmund operators is bounded on $H\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ (or $HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$) and from $H\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ (or $HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$) into the weak Herz-type Hardy spaces with variable exponent $WH\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ (or $WHK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$) at the endpoint case.

In addition, we denote the Lebesgue measure and the characteristic function of a measurable set $A \subset \mathbb{R}^n$ by $|A|$ and χ_A respectively. The notation $f \approx g$ means that there exist constants $C_1, C_2 > 0$ such that $C_1g \leq f \leq C_2g$.

2 Weak Herz type spaces with variable exponent

In this section we first recall some basic definitions and properties of variable function spaces, and then introduce weak Herz spaces and weak Herz-type Hardy spaces with variable exponent.

Given an open set $E \subset \mathbb{R}^n$, and a measurable function $p(\cdot) : E \rightarrow [1, \infty)$, $L^{p(\cdot)}(E)$ denotes the set of measurable functions f on E such that for some $\eta > 0$,

$$\int_E \left(\frac{|f(x)|}{\eta} \right)^{p(x)} dx < \infty.$$

This set becomes a Banach function space when equipped with the Luxemburg-Nakano norm

$$\|f\|_{L^{p(\cdot)}(E)} = \inf \left\{ \eta > 0 : \int_E \left(\frac{|f(x)|}{\eta} \right)^{p(x)} dx \leq 1 \right\}.$$

These spaces are referred as variable L^p spaces, since they generalized the standard L^p spaces: if $p(x) = p$ is constant, then $L^{p(\cdot)}(E)$ is isometrically isomorphic to $L^p(E)$.

The space $L_{\text{loc}}^{p(\cdot)}(E)$ is defined by

$$L_{\text{loc}}^{p(\cdot)}(E) := \{f : f \in L^{p(\cdot)}(F) \text{ for all compact subsets } F \subset E\}.$$

Define $\mathcal{P}(E)$ to be set of measurable functions $p(\cdot) : E \rightarrow [1, \infty)$ such that

$$p^- = \text{ess inf}\{p(x) : x \in E\} > 1, \quad p^+ = \text{ess sup}\{p(x) : x \in E\} < \infty.$$

Denote $p'(x) = p(x)/(p(x) - 1)$.

Let f be a locally integrable function. The Hardy-Littlewood maximal operator is defined by

$$Mf(x) = \sup_{B \ni x} \frac{1}{|B|} \int_{B \cap E} |f(y)| dy,$$

where the supremum is taken over all balls B containing x . Let $\mathcal{B}(E)$ be the set of $p(\cdot) \in \mathcal{P}(E)$ such that M is bounded on $L^{p(\cdot)}(E)$.

In variable L^p spaces there are some important lemmas as follows.

Lemma 2.1^[2] *Given an open set $E \subset \mathbb{R}^n$. If $p(\cdot) \in \mathcal{P}(E)$ and satisfies*

$$|p(x) - p(y)| \leq \frac{C}{-\log(|x - y|)}, \quad |x - y| \leq 1/2 \quad (2.1)$$

and

$$|p(x) - p(y)| \leq \frac{C}{\log(|x| + e)}, \quad |y| \geq |x|, \quad (2.2)$$

then $p(\cdot) \in \mathcal{B}(E)$, that is the Hardy-Littlewood maximal operator M is bounded on $L^{p(\cdot)}(E)$.

Lemma 2.2^[11] (Generalized Hölder's inequality) *Given an open set $E \subset \mathbb{R}^n$ and let $p(\cdot) \in \mathcal{P}(E)$. If $f \in L^{p(\cdot)}(E)$ and $g \in L^{p'(\cdot)}(E)$, then fg is integrable on E and*

$$\int_E |f(x)g(x)|dx \leq r_p \|f\|_{L^{p(\cdot)}(E)} \|g\|_{L^{p'(\cdot)}(E)},$$

where

$$r_p = 1 + 1/p^- - 1/p^+.$$

Lemma 2.3^[9] *Let $p(\cdot) \in \mathcal{B}(\mathbb{R}^n)$. Then there exists a positive constant C such that for all balls B in \mathbb{R}^n and all measurable subsets $S \subset B$,*

$$\frac{\|\chi_B\|_{L^{p(\cdot)}(\mathbb{R}^n)}}{\|\chi_S\|_{L^{p(\cdot)}(\mathbb{R}^n)}} \leq C \frac{|B|}{|S|}, \quad \frac{\|\chi_S\|_{L^{p(\cdot)}(\mathbb{R}^n)}}{\|\chi_B\|_{L^{p(\cdot)}(\mathbb{R}^n)}} \leq C \left(\frac{|S|}{|B|} \right)^{\delta_1} \quad \text{and} \quad \frac{\|\chi_S\|_{L^{p'(\cdot)}(\mathbb{R}^n)}}{\|\chi_B\|_{L^{p'(\cdot)}(\mathbb{R}^n)}} \leq C \left(\frac{|S|}{|B|} \right)^{\delta_2},$$

where $0 < \delta_1, \delta_2 < 1$ depend on $p(\cdot)$.

Throughout this paper δ_2 is the same as in Lemma 2.3.

Lemma 2.4^[9] *Suppose $p(\cdot) \in \mathcal{B}(\mathbb{R}^n)$. Then there exists a positive constant C such that for all balls B in \mathbb{R}^n ,*

$$\frac{1}{|B|} \|\chi_B\|_{L^{p(\cdot)}(\mathbb{R}^n)} \|\chi_B\|_{L^{p'(\cdot)}(\mathbb{R}^n)} \leq C.$$

Lemma 2.5^[3] *Given E and $p(\cdot) \in \mathcal{P}(E)$, let $f : E \times E \rightarrow \mathbb{R}$ be a measurable function (with respect to product measure) such that for almost every $y \in E$, $f(\cdot, y) \in L^{p(\cdot)}(E)$. Then*

$$\left\| \int_E f(\cdot, y) dy \right\|_{L^{p(\cdot)}(E)} \leq C \int_E \|f(\cdot, y)\|_{L^{p(\cdot)}(E)} dy.$$

Now we recall the definition of the Herz-type spaces with variable exponent. Let $B_k = \{x \in \mathbb{R}^n : |x| \leq 2^k\}$ and $A_k = B_k \setminus B_{k-1}$ for $k \in \mathbb{Z}$. Denote \mathbb{Z}_+ and \mathbb{N} as the sets of all positive and non-negative integers, $\chi_k = \chi_{A_k}$ for $k \in \mathbb{Z}$, $\tilde{\chi}_k = \chi_k$ if $k \in \mathbb{Z}_+$ and $\tilde{\chi}_0 = \chi_{B_0}$.

Definition 2.1^[9] Let $\alpha \in \mathbb{R}$, $0 < p \leq \infty$ and $q(\cdot) \in \mathcal{P}(\mathbb{R}^n)$. The homogeneous Herz space with variable exponent $\dot{K}_{q(\cdot)}^{\alpha, p}(\mathbb{R}^n)$ is defined by

$$\dot{K}_{q(\cdot)}^{\alpha, p}(\mathbb{R}^n) = \{f \in L_{\text{loc}}^{q(\cdot)}(\mathbb{R}^n \setminus \{0\}) : \|f\|_{\dot{K}_{q(\cdot)}^{\alpha, p}(\mathbb{R}^n)} < \infty\},$$

where

$$\|f\|_{\dot{K}_{q(\cdot)}^{\alpha, p}(\mathbb{R}^n)} = \left\{ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \|f \chi_k\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p}.$$

The non-homogeneous Herz space with variable exponent $K_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ is defined by

$$K_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n) = \{f \in L_{\text{loc}}^{q(\cdot)}(\mathbb{R}^n) : \|f\|_{K_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)} < \infty\},$$

where

$$\|f\|_{K_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)} = \left\{ \sum_{k=0}^{\infty} 2^{k\alpha p} \|f \tilde{\chi}_k\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p}.$$

In [15], the authors gave the definition of Herz-type Hardy space with variable exponent $HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ and the atomic decomposition characterizations. $\mathcal{S}(\mathbb{R}^n)$ denotes the space of Schwartz functions, and $\mathcal{S}'(\mathbb{R}^n)$ denotes the dual space of $\mathcal{S}(\mathbb{R}^n)$. Let $G_N(f)(x)$ be the grand maximal function of $f(x)$ defined by

$$G_N(f)(x) = \sup_{\phi \in \mathcal{A}_N} |\phi_{\nabla}^*(f)(x)|,$$

where $\mathcal{A}_N = \{\phi \in \mathcal{S}(\mathbb{R}^n) : \sup_{|\alpha|, |\beta| \leq N} |x^\alpha D^\beta \phi(x)| \leq 1\}$ and $N > n + 1$, ϕ_{∇}^* is the nontangential maximal operator defined by

$$\phi_{\nabla}^*(f)(x) = \sup_{|y-x| < t} |\phi_t * f(y)|$$

with $\phi_t(x) = t^{-n} \phi(x/t)$.

Definition 2.2^[15] Let $\alpha \in \mathbb{R}$, $0 < p < \infty$, $q(\cdot) \in \mathcal{P}(\mathbb{R}^n)$ and $N > n + 1$.

(i) The homogeneous Herz-type Hardy space with variable exponent $HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ is defined by

$$HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n) = \left\{ f \in \mathcal{S}'(\mathbb{R}^n) : G_N(f)(x) \in \dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n) \right\}$$

and $\|f\|_{HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)} = \|G_N(f)\|_{\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)}$.

(ii) The non-homogeneous Herz-type Hardy space with variable exponent $HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ is defined by

$$HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n) = \left\{ f \in \mathcal{S}'(\mathbb{R}^n) : G_N(f)(x) \in K_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n) \right\}$$

and $\|f\|_{HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)} = \|G_N(f)\|_{K_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)}$.

For $x \in \mathbb{R}$ we denote by $[x]$ the largest integer less than or equal to x .

Definition 2.3^[15] Let $n\delta_2 \leq \alpha < \infty$, $q(\cdot) \in \mathcal{P}(\mathbb{R}^n)$, and non-negative integer $s \geq [\alpha - n\delta_2]$.

(i) A function $a(x)$ on \mathbb{R}^n is said to be a central $(\alpha, q(\cdot))$ -atom, if it satisfies

$$(1) \text{supp } a \subset B(0, r) = \{x \in \mathbb{R}^n : |x| < r\}.$$

$$(2) \|a\|_{L^{q(\cdot)}(\mathbb{R}^n)} \leq |B(0, r)|^{-\alpha/n}.$$

$$(3) \int_{\mathbb{R}^n} a(x) x^\beta dx = 0, |\beta| \leq s.$$

(ii) A function $a(x)$ on \mathbb{R}^n is said to be a central $(\alpha, q(\cdot))$ -atom of restricted type, if it satisfies the conditions (2), (3) above and

$$(1)' \text{supp } a \subset B(0, r), r \geq 1.$$

If $r = 2^k$ for some $k \in \mathbb{Z}$ in Definition 1.3, then the corresponding central $(\alpha, q(\cdot))$ -atom is called a dyadic central $(\alpha, q(\cdot))$ -atom.

Lemma 2.6^[15] Let $n\delta_2 \leq \alpha < \infty$, $0 < p < \infty$ and $q(\cdot) \in \mathcal{B}(\mathbb{R}^n)$. Then $f \in \dot{H}K_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ (or $HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$) if and only if

$$f = \sum_{k=-\infty}^{\infty} \lambda_k a_k \left(\text{or } \sum_{k=0}^{\infty} \lambda_k a_k \right), \quad \text{in the sense of } \mathcal{S}'(\mathbb{R}^n),$$

where each a_k is a central $(\alpha, q(\cdot))$ -atom (or central $(\alpha, q(\cdot))$ -atom of restricted type) with support contained in B_k and $\sum_{k=-\infty}^{\infty} |\lambda_k|^p < \infty$ (or $\sum_{k=0}^{\infty} |\lambda_k|^p < \infty$). Moreover,

$$\|f\|_{\dot{H}K_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)} \approx \inf \left(\sum_{k=-\infty}^{\infty} |\lambda_k|^p \right)^{1/p} \left(\text{or } \|f\|_{HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)} \approx \inf \left(\sum_{k=0}^{\infty} |\lambda_k|^p \right)^{1/p} \right),$$

where the infimum is taken over all above decompositions of f .

Next we recall the definition of the weak Herz spaces and the weak Herz-type Hardy spaces. For $k \in \mathbb{Z}$, let $m_k(\sigma, f) = |\{x \in A_k : |f(x)| > \sigma\}|$; for $k \in \mathbb{N}$, let $\tilde{m}_k(\sigma, f) = m_k(\sigma, f)$ and $\tilde{m}_0(\sigma, f) = |\{x \in B(0, 1) : |f(x)| > \sigma\}|$.

Definition 2.4^[7] Let $\alpha \in \mathbb{R}$, $0 < p \leq \infty$ and $0 < q < \infty$. A measurable function $f(x)$ on \mathbb{R}^n is said to belong to the homogeneous weak Herz space $WK_q^{\alpha,p}(\mathbb{R}^n)$, if

$$\|f\|_{WK_q^{\alpha,p}(\mathbb{R}^n)} = \sup_{\lambda > 0} \lambda \left\{ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} m_k(\lambda, f)^{p/q} \right\}^{1/p} < \infty,$$

where the usual modification is made when $p = \infty$.

A measurable function $f(x)$ on \mathbb{R}^n is said to belong to the non-homogeneous weak Herz space $WK_q^{\alpha,p}(\mathbb{R}^n)$, if

$$\|f\|_{WK_q^{\alpha,p}(\mathbb{R}^n)} = \sup_{\lambda > 0} \lambda \left\{ \sum_{k=0}^{\infty} 2^{k\alpha p} \tilde{m}_k(\lambda, f)^{p/q} \right\}^{1/p} < \infty,$$

where the usual modification is made when $p = \infty$.

Definition 2.5^[8] Let $\alpha \in \mathbb{R}$, $0 < p, q \leq \infty$ and N be sufficiently large. We define the spaces

$$WH\dot{K}_q^{\alpha,p}(\mathbb{R}^n) = \{f \in \mathcal{S}'(\mathbb{R}^n) : G_N(f)(x) \in WK_q^{\alpha,p}(\mathbb{R}^n)\}$$

and

$$WHK_q^{\alpha,p}(\mathbb{R}^n) = \{f \in \mathcal{S}'(\mathbb{R}^n) : G_N(f)(x) \in WK_q^{\alpha,p}(\mathbb{R}^n)\}.$$

Moreover, we define that

$$\|f\|_{WH\dot{K}_q^{\alpha,p}(\mathbb{R}^n)} = \|G_N(f)\|_{WK_q^{\alpha,p}(\mathbb{R}^n)} \quad \text{and} \quad \|f\|_{WHK_q^{\alpha,p}(\mathbb{R}^n)} = \|G_N(f)\|_{WK_q^{\alpha,p}(\mathbb{R}^n)}.$$

Now we extend the Definitions 2.4 and 2.5 to the case of function spaces with variable exponent.

Definition 2.6 Let $\alpha \in \mathbb{R}$, $0 < p \leq \infty$ and $q(\cdot) \in \mathcal{P}(\mathbb{R}^n)$. A measurable function $f(x)$ on \mathbb{R}^n is said to belong to the homogeneous weak Herz space with variable exponent $WK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$, if

$$\|f\|_{WK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)} = \sup_{\lambda > 0} \lambda \left\{ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \|\chi_{\{x \in A_k : |f(x)| > \lambda\}}\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} < \infty,$$

where the usual modification is made when $p = \infty$.

A measurable function $f(x)$ on \mathbb{R}^n is said to belong to the non-homogeneous weak Herz space with variable exponent $WK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$, if

$$\|f\|_{WK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)} = \sup_{\lambda>0} \lambda \left\{ \sum_{k=0}^{\infty} 2^{k\alpha p} \|\chi_{\{x \in A_k: |f(x)|>\lambda\}}\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} < \infty,$$

where the usual modification is made when $p = \infty$.

Definition 2.7 Let $\alpha \in \mathbb{R}$, $0 < p < \infty$, $q(\cdot) \in \mathcal{P}(\mathbb{R}^n)$ and $N > n + 1$. We define the spaces

$$WH\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n) = \{f \in \mathcal{S}'(\mathbb{R}^n) : G_N(f)(x) \in W\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)\}$$

and

$$WHK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n) = \{f \in \mathcal{S}'(\mathbb{R}^n) : G_N(f)(x) \in WK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)\}.$$

Moreover, we define that

$$\|f\|_{WH\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)} = \|G_N(f)\|_{W\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)} \text{ and } \|f\|_{WHK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)} = \|G_N(f)\|_{WK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)}.$$

Remark 2.1 If $q(\cdot) = q$ is constant in Definition 2.6 and Definition 2.7, then we can easily get the Definition 2.4 and Definition 2.5 respectively.

Remark 2.2 If $\alpha = 0$, then $WL^{q(\cdot)}(\mathbb{R}^n) \subset W\dot{K}_{q(\cdot)}^{0,p}(\mathbb{R}^n)$, where $WL^{q(\cdot)}(\mathbb{R}^n)$ is the weak Lebesgue space with variable exponent and

$$\|f\|_{WL^{q(\cdot)}(\mathbb{R}^n)} = \sup_{\lambda>0} \lambda \|\chi_{\{x \in A_k: |f(x)|>\lambda\}}\|_{L^{q(\cdot)}(\mathbb{R}^n)} < \infty.$$

3 Boundedness of some singular integral operators

Firstly we establish the boundedness for a large class of operators on Herz type spaces with variable exponent.

Theorem 3.1 Let $0 < \alpha < n\delta_2$, $q(\cdot) \in \mathcal{B}(\mathbb{R}^n)$, $p \in (0, 1]$. If T be a sublinear operator and bounded on $WL^{q(\cdot)}(\mathbb{R}^n)$ satisfying

$$|Tf(x)| \leq C|x|^{-n} \int_{\mathbb{R}^n} |f(y)|dy \quad (3.1)$$

for any $f \in L_{\text{loc}}^1(\mathbb{R}^n)$, $\text{supp } f \subset B(0, r)$ and $x \notin B(0, 2r)$. Then T maps continuously $\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ (or $K_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$) into $W\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ (or $WK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$).

Proof We only prove the homogeneous case. The non-homogeneous case can be proved in the same way. For any $k \in \mathbb{Z}$, we decompose f into

$$f(x) = f(x)\chi_{\{|x| \leq 2^{k-3}\}}(x) + f(x)\chi_{\{|x| > 2^{k-3}\}}(x) = f_1(x) + f_2(x).$$

Then $|Tf(x)| \leq |Tf_1(x)| + |Tf_2(x)|$, and

$$\begin{aligned} \|Tf\|_{W\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)} &= \sup_{\lambda>0} \lambda \left\{ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \|\chi_{\{x \in A_k: |Tf(x)|>\lambda\}}\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} \\ &\leq C \sup_{\lambda>0} \lambda \left\{ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \|\chi_{\{x \in A_k: |Tf_1(x)|>\frac{\lambda}{2}\}}\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} \end{aligned}$$

$$\begin{aligned}
& + C \sup_{\lambda > 0} \lambda \left\{ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \|\chi_{\{x \in A_k: |Tf_2(x)| > \frac{\lambda}{2}\}}\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} \\
& = I_1 + I_2.
\end{aligned}$$

Using the $WL^{q(\cdot)}(\mathbb{R}^n)$ -boundedness of T and $0 < p \leq 1$, we have

$$\begin{aligned}
I_2 & \leq C \left\{ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \|Tf_2\chi_k\|_{WL^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} \\
& \leq C \left\{ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \left(\sum_{l=k-2}^{\infty} \|f\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)} \right)^p \right\}^{1/p} \\
& \leq C \left\{ \sum_{l=-\infty}^{\infty} \|f\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \left(\sum_{k=-\infty}^{l+2} 2^{k\alpha p} \right) \right\}^{1/p} \\
& \leq C \left\{ \sum_{l=-\infty}^{\infty} 2^{l\alpha p} \|f\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} \\
& = C \|f\|_{\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)}.
\end{aligned}$$

For I_1 , noting that $x \in A_k$ and $\text{supp } f_1 \subset \{x \in \mathbb{R}^n : |x| \leq 2^{k-3}\}$, by (3.1) and the generalized Hölder inequality we have

$$\begin{aligned}
|Tf_1(x)| & \leq C|x|^{-n} \|f_1\|_{L^1(\mathbb{R}^n)} \\
& \leq C2^{-kn} \sum_{l=-\infty}^{k-3} \|f\chi_l\|_{L^1(\mathbb{R}^n)} \\
& \leq C2^{-kn} \sum_{l=-\infty}^{k-3} \|f\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)} \|\chi_l\|_{L^{q'(\cdot)}(\mathbb{R}^n)}.
\end{aligned}$$

So by Lemmas 2.3 and 2.4 we have

$$\begin{aligned}
& \left\| \chi_{\{x \in A_k: |Tf_1(x)| > \frac{\lambda}{2}\}} \right\|_{L^{q(\cdot)}(\mathbb{R}^n)} \\
& \leq \left\| \chi_{\{x \in A_k: C2^{-kn} \sum_{l=-\infty}^{k-3} \|f\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)} \|\chi_l\|_{L^{q'(\cdot)}(\mathbb{R}^n)} > \frac{\lambda}{2}\}} \right\|_{L^{q(\cdot)}(\mathbb{R}^n)} \\
& \leq C2^{-kn+1} \lambda^{-1} \sum_{l=-\infty}^{k-3} \|f\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)} \|\chi_l\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \|\chi_k\|_{L^{q(\cdot)}(\mathbb{R}^n)} \\
& \leq C2^{-kn} \lambda^{-1} \sum_{l=-\infty}^{k-3} \|f\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)} \|\chi_{B_l}\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \|\chi_{B_k}\|_{L^{q(\cdot)}(\mathbb{R}^n)}^{-1} |B_k| \\
& \leq C\lambda^{-1} \sum_{l=-\infty}^{k-3} \|f\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)} 2^{(l-k)n\delta_2}.
\end{aligned}$$

Thus by $0 < p \leq 1$ and $0 < \alpha < n\delta_2$ we obtain

$$\begin{aligned}
I_1 & = C \sup_{\lambda > 0} \lambda \left\{ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \|\chi_{\{x \in A_k: |Tf_1(x)| > \frac{\lambda}{2}\}}\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} \\
& \leq C \left\{ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \sum_{l=-\infty}^{k-3} \|f\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p 2^{(l-k)n\delta_2 p} \right\}^{1/p}
\end{aligned}$$

$$\begin{aligned}
&\leq C \left\{ \sum_{l=-\infty}^{\infty} 2^{l\alpha p} \|f\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \left(\sum_{k=l+3}^{\infty} 2^{(l-k)(n\delta_2-\alpha)p} \right) \right\}^{1/p} \\
&\leq C \left\{ \sum_{l=-\infty}^{\infty} 2^{l\alpha p} \|f\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} \\
&= C \|f\|_{\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)}.
\end{aligned}$$

This completes the proof of Theorem 3.1.

Remark 3.1 If the condition (3.1) is substituted by the following condition

$$|Tf(x)| \leq C \int_{\mathbb{R}^n} \frac{|f(y)|}{|x-y|^n} dy, \quad x \notin \text{supp} f, \quad (3.2)$$

then the conclusions of Theorem 3.1 are still true.

Remark 3.2 The condition (3.1) is very weak and many classical operators satisfy (3.1), such as Calderón-Zygmund operators, multipliers and oscillatory singular integrals, Bochner-Riesz operators at the critical index and so on.

Now we turn our attention to the behaviours of local Calderón-Zygmund type operators on the Herz-type Hardy spaces with variable exponent.

Theorem 3.2 *Let $T : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$ be a linear and continuous operator. Assume that the distribution kernel of T coincides in the complement of the diagonal with a locally integrable function $k(x, y)$ which satisfies*

$$\sup_{y \in B_k} \|[k(\cdot, y) - k(\cdot, 0)]\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)} \leq C 2^{(k-l)\delta - ln} \|\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)} \quad (3.3)$$

for $k, l \in \mathbb{Z}$, some $\delta \in (0, 1]$ and some $q(\cdot) \in \mathcal{B}(\mathbb{R}^n)$. Suppose that T can be extended to a continuous operator on $L^{q(\cdot)}(\mathbb{R}^n)$. If $n\delta_2 \leq \alpha < n\delta_2 + \delta$ and $0 < p \leq \infty$, then T maps continuously $H\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ into $\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$.

Remark 3.3 If the condition (3.3) is true only for $k \in \mathbb{N}$ and $l \in \mathbb{Z}_+$, then the operator T in Theorem 3.2 maps continuously $H\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ into $\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$.

Remark 3.4 Assuming more regularity on the kernel $k(x, y)$, we can extend Theorem 3.2 to larger range of α .

Proof of Theorem 3.2 Firstly we suppose that $0 < p \leq 1$. In this case, we only need to prove $\|Ta_k\|_{\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)} \leq C$, where a_k is a dyadic central $(\alpha, q(\cdot))$ -atom with the support B_k and C is independent of k . Write

$$\begin{aligned}
\|Ta_k\|_{\dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)} &\leq C \left\{ \sum_{l=-\infty}^{k+3} 2^{l\alpha p} \|(Ta_k)\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} \\
&\quad + C \left\{ \sum_{l=k+4}^{\infty} 2^{l\alpha p} \|(Ta_k)\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} \\
&= J_1 + J_2.
\end{aligned}$$

Since T is bounded on $L^{q(\cdot)}(\mathbb{R}^n)$, by Definition 2.3 we have

$$J_1 \leq C \left\{ \sum_{l=-\infty}^{k+3} 2^{l\alpha p} \|Ta_k\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p}$$

$$\begin{aligned}
&\leq C \left\{ \sum_{l=-\infty}^{k+3} 2^{l\alpha p} \|a_k\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} \\
&\leq C 2^{-k\alpha} \left\{ \sum_{l=-\infty}^{k+3} 2^{l\alpha p} \right\}^{1/p} \leq C,
\end{aligned}$$

where C is independent of k .

For J_2 , using (3.3), Lemmas 2.3, 2.4, 2.5 and the generalized Hölder inequality we have

$$\begin{aligned}
\|(Ta_k)\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)} &\leq C \int_{B_k} \|[k(\cdot, y) - k(\cdot, 0)]\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)} |a_k(y)| dy \\
&\leq C 2^{(k-l)\delta - ln} \|\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)} \|a_k\|_{L^{q(\cdot)}(\mathbb{R}^n)} \|\chi_k\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \\
&\leq C 2^{(k-l)\delta - k\alpha} \|\chi_{B_l}\|_{L^{q'(\cdot)}(\mathbb{R}^n)}^{-1} \|\chi_{B_k}\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \\
&\leq C 2^{(k-l)(\delta + n\delta_2) - k\alpha}.
\end{aligned}$$

Thus by $\alpha < n\delta_2 + \delta$ we obtain

$$J_2 \leq C \left\{ \sum_{l=k+4}^{\infty} 2^{(l-k)(\alpha - \delta - n\delta_2)p} \right\}^{1/p} \leq C,$$

where C is independent of k . This finishes the proof for the case $0 < p \leq 1$. Now let $1 < p < \infty$ and

$f \in H\dot{K}_{q(\cdot)}^{\alpha, p}(\mathbb{R}^n)$. By Lemma 2.6 we get $f = \sum_{k=-\infty}^{\infty} \lambda_k a_k$, where $\|f\|_{H\dot{K}_{q(\cdot)}^{\alpha, p}(\mathbb{R}^n)} \approx \inf \left(\sum_{k=-\infty}^{\infty} |\lambda_k|^p \right)^{1/p}$

(the infimum is taken over above decompositions of f), and a_k is a dyadic central $(\alpha, q(\cdot))$ -atom with the support B_k . Write

$$\begin{aligned}
\|Tf\|_{\dot{K}_{q(\cdot)}^{\alpha, p}(\mathbb{R}^n)} &\leq C \left\{ \sum_{l=-\infty}^{\infty} 2^{l\alpha p} \left(\sum_{k=-\infty}^{l-4} |\lambda_k| \|(Ta_k)\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)} \right)^p \right\}^{1/p} \\
&\quad + C \left\{ \sum_{l=-\infty}^{\infty} 2^{l\alpha p} \left(\sum_{k=l-3}^{\infty} |\lambda_k| \|(Ta_k)\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)} \right)^p \right\}^{1/p} \\
&= U_1 + U_2.
\end{aligned}$$

For U_2 , by the Hölder inequality and the fact that T is bounded on $L^{q(\cdot)}(\mathbb{R}^n)$, we have

$$\begin{aligned}
U_2 &\leq C \left\{ \sum_{l=-\infty}^{\infty} 2^{l\alpha p} \left(\sum_{k=l-3}^{\infty} |\lambda_k| \|(Ta_k)\|_{L^{q(\cdot)}(\mathbb{R}^n)} \right)^p \right\}^{1/p} \\
&\leq C \left\{ \sum_{l=-\infty}^{\infty} 2^{l\alpha p} \left(\sum_{k=l-3}^{\infty} |\lambda_k| 2^{-k\alpha} \right)^p \right\}^{1/p} \\
&\leq C \left\{ \sum_{l=-\infty}^{\infty} 2^{l\alpha p/2} \left(\sum_{k=l-3}^{\infty} |\lambda_k|^p 2^{-k\alpha p/2} \right) \right\}^{1/p} \\
&\leq C \left\{ \sum_{k=-\infty}^{\infty} |\lambda_k|^p \right\}^{1/p} \leq C \|f\|_{H\dot{K}_{q(\cdot)}^{\alpha, p}(\mathbb{R}^n)}.
\end{aligned}$$

On the other hand, noting that $\alpha < n\delta_2 + \delta$. So by (3.3), Lemmas 2.3, 2.4, 2.5 and the generalized Hölder inequality we have

$$\begin{aligned}
U_1 &\leq C \left\{ \sum_{l=-\infty}^{\infty} 2^{l\alpha p} \left(\sum_{k=-\infty}^{l-4} |\lambda_k| \|a_k\|_{L^1(\mathbb{R}^n)} 2^{(k-l)\delta - ln} \|\chi_l\|_{L^{q(\cdot)}(\mathbb{R}^n)} \right)^p \right\}^{1/p} \\
&\leq C \left\{ \sum_{l=-\infty}^{\infty} \left(\sum_{k=-\infty}^{l-4} |\lambda_k| 2^{(l-k)(\alpha - \delta - n\delta_2)} \right)^p \right\}^{1/p} \\
&\leq C \left\{ \sum_{l=-\infty}^{\infty} \left(\sum_{k=-\infty}^{l-4} |\lambda_k|^p 2^{(l-k)(\alpha - \delta - n\delta_2)p/2} \right) \right\}^{1/p} \\
&\leq C \left\{ \sum_{k=-\infty}^{\infty} |\lambda_k|^p \left(\sum_{l=k+4}^{\infty} 2^{(l-k)(\alpha - \delta - n\delta_2)p/2} \right) \right\}^{1/p} \\
&\leq C \left\{ \sum_{k=-\infty}^{\infty} |\lambda_k|^p \right\}^{1/p} \leq C \|f\|_{HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)}.
\end{aligned}$$

Therefore, we complete the proof of Theorem 3.2.

When $\alpha = n\delta_2 + \delta$, we have the following result.

Theorem 3.3 *Let $T : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$ be a linear operator. Suppose that the distribution kernel of T coincides in the complement of the diagonal with a locally integrable function $k(x, y)$ satisfying*

$$|k(x, y) - k(x, 0)| \leq C \frac{|y|^\delta}{|x|^{n+\delta}}, \quad (3.4)$$

if $2|y| < |x|$, for some $\delta \in (0, 1]$. Suppose that T is bounded on $L^{q(\cdot)}(\mathbb{R}^n)$ for some $q(\cdot) \in \mathcal{B}(\mathbb{R}^n)$, $\alpha = n\delta_2 + \delta$ and $0 < p \leq 1$, then T maps continuously $HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ (or $HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$) into $WK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ (or $WK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$).

Proof We only prove the homogeneous case. Let $\alpha = n\delta_2 + \delta$ and $f \in HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$. By Lemma 2.6

we get $f = \sum_{l=-\infty}^{\infty} \lambda_l a_l$, where $\|f\|_{HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)} \approx \inf \left(\sum_{l=-\infty}^{\infty} |\lambda_l|^p \right)^{1/p}$ (the infimum is taken over above decompositions of f), and a_l is a dyadic central $(\alpha, q(\cdot))$ -atom with the support B_l . Given $\lambda > 0$, we can write

$$\begin{aligned}
&\lambda \left\{ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \|\chi_{\{x \in A_k : |Tf(x)| > \lambda\}}\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} \\
&\leq C\lambda \left\{ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \|\chi_{\{x \in A_k : |\sum_{l=-\infty}^{k-4} \lambda_l T a_l(x)| > \frac{\lambda}{2}\}}\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} \\
&\quad + C\lambda \left\{ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \|\chi_{\{x \in A_k : |\sum_{l=k-3}^{\infty} \lambda_l T a_l(x)| > \frac{\lambda}{2}\}}\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} \\
&= CV_1 + CV_2.
\end{aligned}$$

For V_2 , by the $L^{q(\cdot)}(\mathbb{R}^n)$ -boundedness of T , we have

$$\begin{aligned}
V_2 &\leq C \left\{ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \left\| \sum_{l=k-3}^{\infty} |\lambda_l T a_l| \chi_k \right\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} \\
&\leq C \left\{ \sum_{k=-\infty}^{\infty} 2^{k\alpha p} \left(\sum_{l=k-3}^{\infty} |\lambda_l| \|a_l\|_{L^{q(\cdot)}(\mathbb{R}^n)} \right)^p \right\}^{1/p} \\
&\leq C \left\{ \sum_{k=-\infty}^{\infty} \sum_{l=k-3}^{\infty} |\lambda_l|^p 2^{(k-l)\alpha p} \right\}^{1/p} \\
&\leq C \left\{ \sum_{l=-\infty}^{\infty} |\lambda_l|^p \sum_{k=-\infty}^{l+3} 2^{(k-l)\alpha p} \right\}^{1/p} \\
&\leq C \left\{ \sum_{l=-\infty}^{\infty} |\lambda_l|^p \right\}^{1/p} \leq C \|f\|_{HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)}.
\end{aligned}$$

To estimate V_1 , we notice that if $x \in A_k$ and $k \geq l + 4$, then by (3.4), $\alpha = n\delta_2 + \delta$, Lemma 2.3 and the generalized Hölder inequality we have

$$\begin{aligned}
|T a_l(x)| &\leq \int_{\mathbb{R}^n} |k(x, y) - k(x, 0)| |a_l(y)| dy \\
&\leq C 2^{l\delta - k(n+\delta)} \int_{\mathbb{R}^n} |a_l(y)| dy \\
&\leq C 2^{l\delta - k(n+\delta)} \|a_l\|_{L^{q(\cdot)}(\mathbb{R}^n)} \|\chi_l\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \\
&\leq C 2^{l(\delta - \alpha) - k(n+\delta)} \|\chi_l\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \\
&\leq C 2^{l(\delta - \alpha) - k(n+\delta)} 2^{(l-k)n\delta_2} \|\chi_k\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \\
&= C 2^{-k(n+\delta+n\delta_2)} \|\chi_k\|_{L^{q'(\cdot)}(\mathbb{R}^n)}.
\end{aligned}$$

So by $0 < p \leq 1$ we have

$$\begin{aligned}
\left| \sum_{l=-\infty}^{k-4} \lambda_l T a_l(x) \right| &\leq C 2^{-k(n+\delta+n\delta_2)} \|\chi_k\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \sum_{l=-\infty}^{k-4} |\lambda_l| \\
&\leq C 2^{-k(n+\delta+n\delta_2)} \|\chi_k\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \left(\sum_{l=-\infty}^{k-4} |\lambda_l|^p \right)^{1/p} \\
&\leq C 2^{-k(n+\delta+n\delta_2)} \|\chi_k\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \|f\|_{HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)}.
\end{aligned}$$

If $|\{x \in A_k : |\sum_{l=-\infty}^{k-4} \lambda_l T a_l(x)| > \frac{\lambda}{2}\}| \neq 0$, then

$$\lambda \leq 2C_0 2^{-k(n+\delta+n\delta_2)} \|\chi_k\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \|f\|_{HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)}.$$

For any given $\lambda > 0$, let k_λ be the maximal positive integer such that

$$2^{k_\lambda(n+\delta+n\delta_2)} \|\chi_{k_\lambda}\|_{L^{q'(\cdot)}(\mathbb{R}^n)}^{-1} \leq 2C_0 \lambda^{-1} \|f\|_{HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)}.$$

Thus by $\alpha = n\delta_2 + \delta$ and Lemma 2.4 we obtain

$$\begin{aligned}
V_1 &\leq C\lambda \left\{ \sum_{k=-\infty}^{k_\lambda} 2^{k\alpha p} \|\chi_{\{x \in A_k : |\sum_{l=-\infty}^{k-4} \lambda_l T a_l(x)| > \frac{\lambda}{2}\}}\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} \\
&\leq C\lambda \left\{ \sum_{k=-\infty}^{k_\lambda} 2^{k\alpha p} \|\chi_k\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} \\
&\leq C\lambda \left\{ \sum_{k=-\infty}^{k_\lambda} 2^{k\alpha p} 2^{kn p} \|\chi_k\|_{L^{q'(\cdot)}(\mathbb{R}^n)}^{-p} \right\}^{1/p} \\
&= C\lambda \left\{ \sum_{k=-\infty}^{k_\lambda} 2^{k(n+\delta+n\delta_2)p} \|\chi_k\|_{L^{q'(\cdot)}(\mathbb{R}^n)}^{-p} \right\}^{1/p} \\
&\leq C\|f\|_{HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)}.
\end{aligned}$$

This completes the proof of Theorem 3.3.

Remark 3.5 Similar to the method in the proof of Theorem 3.3, we can extend the result of Theorem 3.1 to the case $\alpha = n\delta_2$. Here we omit the details.

If the operator T in Theorem 3.2 is of convolution type, then we can obtain the following stronger conclusion.

Theorem 3.4 *Let $Tf(x) = \text{p.v.}(k * f)(x)$ be a bounded operator on $L^{q(\cdot)}(\mathbb{R}^n)$ for some $q(\cdot) \in \mathcal{B}(\mathbb{R}^n)$, and for some $\delta \in (0, 1]$, the kernel k satisfy*

$$|k(x-y) - k(x)| \leq C \frac{|y|^\delta}{|x|^{n+\delta}}, \quad \text{if } |x| > 2|y|. \quad (3.5)$$

If $n\delta_2 \leq \alpha < n\delta_2 + \delta$ and $0 < p < \infty$, then T can be extended to a bounded operator on $HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ (or $HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$).

Using the atomic and molecular theory for the spaces $HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ and $HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$, we can prove Theorem 3.4 by a standard procedure, see [15]. Here we omit the details.

For the operator T in Theorem 3.4, we have the following result which is stronger than Theorem 3.3 with the end case $\alpha = n\delta_2 + \delta$.

Theorem 3.5 *Let T be the same as in Theorem 3.4 with $\delta \in (0, 1)$. If $\alpha = n\delta_2 + \delta$, $q(\cdot) \in \mathcal{B}(\mathbb{R}^n)$ and $0 < p < \infty$, then T maps continuously $HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ (or $HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$) into $WHK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ (or $WHK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$).*

Proof We only prove the homogeneous case. Let $\alpha = n\delta_2 + \delta$ and $f \in HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$. By Lemma

2.6 we get $f = \sum_{k=-\infty}^{\infty} \lambda_k a_k$, where $\|f\|_{HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)} \approx \inf(\sum_{k=-\infty}^{\infty} |\lambda_k|^p)^{1/p}$ (the infimum is taken over above decompositions of f), and a_k is a dyadic central $(\alpha, q(\cdot))$ -atom with the support B_k . For a fixed $\lambda > 0$, we can write

$$\lambda \left\{ \sum_{j=-\infty}^{\infty} 2^{j\alpha p} \|\chi_{\{x \in A_j : G_N(Tf)(x) > \lambda\}}\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p}$$

$$\begin{aligned}
&\leq C\lambda \left\{ \sum_{j=-\infty}^{\infty} 2^{j\alpha p} \|\chi_{\{x \in A_j: \sum_{k=-\infty}^{j-4} |\lambda_k| G_N(Ta_k)(x) > \lambda\}}\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} \\
&\quad + C\lambda \left\{ \sum_{j=-\infty}^{\infty} 2^{j\alpha p} \|\chi_{\{x \in A_j: \sum_{k=j-3}^{\infty} |\lambda_k| G_N(Ta_k)(x) > \lambda\}}\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} \\
&= E_1 + E_2.
\end{aligned}$$

For E_2 , by the $L^{q(\cdot)}(\mathbb{R}^n)$ -boundedness of G_N and T , we have

$$\begin{aligned}
E_2 &\leq C \left\{ \sum_{j=-\infty}^{\infty} 2^{j\alpha p} \left\| \sum_{k=j-3}^{\infty} |\lambda_k| G_N(Ta_k) \right\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right\}^{1/p} \\
&\leq C \left\{ \sum_{j=-\infty}^{\infty} 2^{j\alpha p} \left(\sum_{k=j-3}^{\infty} |\lambda_k|^p \|G_N(Ta_k)\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right) \right\}^{1/p} \\
&\leq C \left\{ \sum_{j=-\infty}^{\infty} 2^{j\alpha p} \left(\sum_{k=j-3}^{\infty} |\lambda_k|^p \|Ta_k\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right) \right\}^{1/p} \\
&\leq C \left\{ \sum_{j=-\infty}^{\infty} 2^{j\alpha p} \left(\sum_{k=j-3}^{\infty} 2^{-k\alpha p} |\lambda_k|^p \right) \right\}^{1/p} \\
&\leq C \left\{ \sum_{k=-\infty}^{\infty} |\lambda_k|^p \right\}^{1/p} \leq C \|f\|_{HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)}.
\end{aligned}$$

Now we consider the term E_1 . We want to obtain the pointwise estimate for $G_N(Ta_k)(x)$ with $x \in A_j$ and $k \leq j-4$. Note that $\int_{\mathbb{R}^n} Ta_k(x) dx = 0$. Let $|x-y| < t$ and write

$$\begin{aligned}
|(Ta_k * \phi_t)(y)| &\leq \int_{\mathbb{R}^n} \left| \int_{B_k} a_k(v) k(w-v) dv \right| t^{-n} \left| \phi\left(\frac{y-w}{t}\right) - \phi\left(\frac{y}{t}\right) \right| dw \\
&\leq \int_{|w| < 2^{k+1}} \left| \int_{B_k} a_k(v) k(w-v) dv \right| t^{-n} \left| \phi\left(\frac{y-w}{t}\right) - \phi\left(\frac{y}{t}\right) \right| dw \\
&\quad + \int_{2^{k+1} \leq |w| < |x|/2} \left| \int_{B_k} a_k(v) k(w-v) dv \right| t^{-n} \left| \phi\left(\frac{y-w}{t}\right) - \phi\left(\frac{y}{t}\right) \right| dw \\
&\quad + \int_{|w| \geq |x|/2} \left| \int_{B_k} a_k(v) k(w-v) dv \right| t^{-n} \left| \phi\left(\frac{y-w}{t}\right) - \phi\left(\frac{y}{t}\right) \right| dw \\
&=: F_1 + F_2 + F_3.
\end{aligned}$$

For F_1 , by Lemma 2.3, the generalized Hölder inequality and the mean value theorem, we obtain

$$\begin{aligned}
F_1 &\leq C \|Ta_k\|_{L^{q(\cdot)}(\mathbb{R}^n)} t^{-n} \left\| \left(\phi\left(\frac{y-\cdot}{t}\right) - \phi\left(\frac{y}{t}\right) \right) \chi_{B_{k+1}} \right\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \\
&\leq C \|a_k\|_{L^{q(\cdot)}(\mathbb{R}^n)} t^{-n} \left\| \sup_{|\beta|=1} \left| D^\beta \phi\left(\frac{y-\theta \cdot}{t}\right) \right| \frac{|\cdot|}{t} \chi_{B_k} \right\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \\
&\leq C \|a_k\|_{L^{q(\cdot)}(\mathbb{R}^n)} \left\| \frac{|\cdot| \chi_{B_k}}{(|x-y| + |y-\theta \cdot|)^{n+1}} \right\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \\
&\leq C 2^{-k\alpha} \frac{1}{|x|^{n+1}} \| |\cdot| \chi_{B_k} \|_{L^{q'(\cdot)}(\mathbb{R}^n)}
\end{aligned}$$

$$\begin{aligned}
&\leq C2^{-k\alpha+k} \frac{1}{|x|^{n+1}} \|\chi_{B_k}\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \\
&\leq C2^{k\delta-k\alpha} \frac{1}{|x|^{n+\delta}} \|\chi_{B_k}\|_{L^{q'(\cdot)}(\mathbb{R}^n)}
\end{aligned}$$

where $0 < \theta < 1$. For F_2 , by (3.5) and the vanishing condition of a_k we have

$$\begin{aligned}
F_2 &= \int_{2^{k+1} \leq |w| < |x|/2} \left| \int_{B_k} a_k(v)(k(w-v) - k(w)) dv \right| t^{-n} \left| \phi\left(\frac{y-w}{t}\right) - \phi\left(\frac{y}{t}\right) \right| dw \\
&\leq \int_{2^{k+1} \leq |w| < |x|/2} \left(\int_{B_k} |a_k(v)| |k(w-v) - k(w)| dv \right) t^{-n} \sup_{|\beta|=1} \left| D^\beta \phi\left(\frac{y-\theta w}{t}\right) \right| \frac{|w|}{t} dw \\
&\leq C \int_{2^{k+1} \leq |w| < |x|/2} \left(\int_{B_k} |a_k(v)| |v|^\delta dv \right) \frac{|w|}{|x|^{n+1}} \frac{1}{|w|^{n+\delta}} dw \\
&\leq C2^{k\delta} \|a_k\|_{L^{q(\cdot)}(\mathbb{R}^n)} \|\chi_{B_k}\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \frac{1}{|x|^{n+1}} \int_{2^{k+1} \leq |w| < |x|/2} |w|^{1-n-\delta} dw \\
&\leq C2^{k\delta-k\alpha} \|\chi_{B_k}\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \frac{1}{|x|^{n+1}} |x|^{1-\delta} \\
&\leq C2^{k\delta-k\alpha} \frac{1}{|x|^{n+\delta}} \|\chi_{B_k}\|_{L^{q'(\cdot)}(\mathbb{R}^n)},
\end{aligned}$$

where $0 < \theta < 1$. By (3.5) and the vanishing condition of a_k we can obtain

$$\begin{aligned}
F_3 &\leq \int_{|w| \geq |x|/2} \left| \int_{B_k} a_k(v)(k(w-v) - k(w)) dv \right| t^{-n} \left(\left| \phi\left(\frac{y-w}{t}\right) \right| + \left| \phi\left(\frac{y}{t}\right) \right| \right) dw \\
&\leq C2^{k\delta-k\alpha} \|\chi_{B_k}\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \int_{|w| \geq |x|/2} |w|^{-n-\delta} t^{-n} \left(\left| \phi\left(\frac{y-w}{t}\right) \right| + \left| \phi\left(\frac{y}{t}\right) \right| \right) dw \\
&\leq C2^{k\delta-k\alpha} \|\chi_{B_k}\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \left(|x|^{-n-\delta} + |x|^{-n} \int_{|w| \geq |x|/2} |w|^{-n-\delta} dw \right) \\
&\leq C2^{k\delta-k\alpha} \frac{1}{|x|^{n+\delta}} \|\chi_{B_k}\|_{L^{q'(\cdot)}(\mathbb{R}^n)},
\end{aligned}$$

where we have invoked the fact that $t + |y| > |x - y| + |y| > |x|$. Thus, if $x \in A_j$ and $k \leq j - 4$, then we have

$$G_N(Ta_k)(x) \leq C2^{k\delta-k\alpha} \frac{1}{|x|^{n+\delta}} \|\chi_{B_k}\|_{L^{q'(\cdot)}(\mathbb{R}^n)}.$$

Therefore, we obtain

$$\sum_{k=-\infty}^{j-4} |\lambda_k| G_N(Ta_k)(x) \leq C \sum_{k=-\infty}^{j-4} |\lambda_k| 2^{k\delta-k\alpha} \frac{1}{|x|^{n+\delta}} \|\chi_{B_k}\|_{L^{q'(\cdot)}(\mathbb{R}^n)}.$$

If $|\{x \in A_j : \sum_{k=-\infty}^{j-4} |\lambda_k| G_N(Ta_k)(x) > \lambda\}| \neq 0$, then by Lemma 2.3 and $\alpha = n\delta_2 + \delta$ we have

$$\begin{aligned}
\lambda &< C \sum_{k=-\infty}^{j-4} |\lambda_k| 2^{k\delta-k\alpha} \frac{1}{|x|^{n+\delta}} \|\chi_{B_k}\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \\
&\leq C \sum_{k=-\infty}^{j-4} |\lambda_k| 2^{k\delta-k\alpha} 2^{-j(n+\delta)} 2^{(k-j)n\delta_2} \|\chi_{B_j}\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \\
&\leq C2^{-j(n+\alpha)} \|\chi_{B_j}\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \sum_{k=-\infty}^{\infty} |\lambda_k| \\
&\leq C2^{-j(n+\alpha)} \|\chi_{B_j}\|_{L^{q'(\cdot)}(\mathbb{R}^n)} \|f\|_{HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)}.
\end{aligned}$$

Let j_λ be the maximal integer such that the above inequality holds. Then by Lemma 2.4 we have

$$\begin{aligned} E_1 &\leq C\lambda \left(\sum_{j=-\infty}^{j_\lambda} 2^{j\alpha p} \|\chi_j\|_{L^{q(\cdot)}(\mathbb{R}^n)}^p \right)^{1/p} \\ &\leq C\lambda \left(\sum_{j=-\infty}^{j_\lambda} 2^{j\alpha p} 2^{jn p} \|\chi_{B_j}\|_{L^{q'(\cdot)}(\mathbb{R}^n)}^{-p} \right)^{1/p} \\ &\leq C\|f\|_{HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)}. \end{aligned}$$

This finishes the proof of Theorem 3.5.

In general, we have the following theorems. The proofs are similar. Here we omit the details.

Theorem 3.6 *Let $q(\cdot) \in \mathcal{B}(\mathbb{R}^n)$, $\alpha \geq n\delta_2$, $s = [\alpha + n\delta_2]$ and $\varepsilon_0 = \alpha + n\delta_2 - s$. Suppose $Tf(x) = \text{p.v.}(k * f)(x)$ is bounded on $L^{q(\cdot)}(\mathbb{R}^n)$ and the kernel k satisfies*

$$|D^J k(x-y) - D^J k(x)| \leq C|y|^\varepsilon |x|^{-n-\varepsilon}$$

for all multi-index J with $|J| = s$, some $\varepsilon > \varepsilon_0$ and $|x| > 2|y|$. If $0 < p < \infty$, then T can be extended to a bounded operator on $HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ (or $HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$).

Theorem 3.7 *Let T be the same as in Theorem 3.6 with $\varepsilon = \varepsilon_0$ and $\alpha > n\delta_2$. If $0 < p \leq 1$, then T maps continuously $HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ (or $HK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$) into $WHK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$ (or $WHK_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$).*

Acknowledgements The authors are very grateful to the referees for their valuable comments. This work was supported by National Natural Science Foundation of China (Grant Nos. 11926343, 11926342, 11761026 and 11671397), Shandong Provincial Natural Science Foundation (Grant No. ZR2017MA041) and Project of Shandong Province Higher Educational Science and Technology Program (Grant No. J18KA225).

References

- [1] Chen Y, Levin S and Rao M, Variable exponent, linear growth functionals in image restoration, *SIAM J. Appl. Math.*, **66**(2006)1383-1406.
- [2] Cruz-Uribe D, Fiorenza A and Neugebauer C, The maximal function on variable L^p spaces, *Ann. Acad. Sci. Fenn. Math.*, **28**(2003)223-238.
- [3] Cruz-Uribe D and Fiorenza A, Variable Lebesgue Spaces: Foundations and Harmonic Analysis (Applied and Numerical Harmonic Analysis), Springer, Heidelberg, 2013.
- [4] Diening L, Harjulehto P, Hästö P and Růžička M, *Lebesgue and Sobolev spaces with variable exponents*, Lecture Notes in Math., vol. **2017**, Springer, Heidelberg, 2011.
- [5] Ferreira L and Pérez-López J, Besov-weak-Herz spaces and global solutions for Navier-Stokes equations, *Pacific J. Math.*, **296**(2018)57-77.
- [6] Harjulehto P, Hästö P, Lê U V and Nuortio M, Overview of differential equations with non-standard growth, *Nonlinear Anal.*, **72**(2010)4551-4574.
- [7] Hu G, Lu S and Yang D, The weak Herz spaces. *J. Beijing Normal Univ. (Natur. Sci.)*, **33**(1997)27-34.
- [8] Hu G, Lu S and Yang D, The applications of weak Herz spaces. *Adv. Math. (China)*, **26**(1997)417-428.
- [9] Izuki M, Boundedness of sublinear operators on Herz spaces with variable exponent and application to wavelet characterization, *Anal. Math.*, **36**(2010)33-50.

- [10] Komori Y, Weak type estimates for Caldern-Zygmund operators on Herz spaces at critical indexes, *Math. Nachr.*, **259**(2003)42-50.
- [11] Kováčik O and Rákosník J, On spaces $L^{p(x)}$ and $W^{k,p(x)}$, *Czechoslovak Math. J.*, **41**(1991)592-618.
- [12] Liu L, The inequalities of commutators on weak Herz spaces, *J. Korean Math. Soc.*, **39**(2002)899-912.
- [13] Růžička M, Electrorheological fluids: modeling and mathematical theory, Springer, Berlin, 2000.
- [14] Tsutsui Y, The Navier-Stokes equations and weak Herz spaces, *Adv. Differential Equations*, **16**(2011)1049-1085.
- [15] Wang H and Liu Z, The Herz-type Hardy spaces with variable exponent and their applications, *Taiwanese J. Math.*, **16**(2012)1363-1389.
- [16] Wang H, Some estimates of intrinsic square functions on weighted Herz-type Hardy spaces, *J. Inequal. Appl.*, **2015**(2015)62: 22 pages.
- [17] Yan X, Yang D, Yuan W and Zhuo C, Variable weak Hardy spaces and their applications, *J. Funct. Anal.*, **271**(2016)2822-2887.

Hongbin WANG

School of Mathematics and Statistics, Shandong University of Technology, Zibo, Shandong, 255049, China

E-mail: wanghb@sdut.edu.cn

Zongguang LIU

Department of Mathematics, China University of Mining and Technology(Beijing), Beijing, 100083, China

E-mail: liuzg@cumtb.edu.cn