

ARCMINUTE MICROKELVIN IMAGER OBSERVATIONS AT 15 GHz OF THE 2020 FEBRUARY OUTBURST OF CYGNUS X-3

David A. Green¹ and Patrick Elwood¹

¹*Cavendish Laboratory,
19 J. J. Thomson Ave., Cambridge, CB3 0HE, UK*

Keywords: High mass x-ray binary stars (733) — Galactic radio sources (571) — Radio continuum emission (1340) — Variable radiation sources (1759)

The high mass X-ray binary Cygnus X-3 shows occasional giant radio flares, with flux densities up to ~ 10 Jy. The first such burst was observed in 1972 (Gregory & Kronberg 1972), and subsequently there have been several similar flares, e.g. see Johnston et al. (1986); Waltman et al. (1995); Fender et al. (1997); Mioduszewski et al. (2001); Corbel et al. (2012); Zdziarski et al. (2016); Egron et al. (2017) for some examples.

Here we report observations of a recent giant radio flare from Cygnus X-3 in 2020 February, made with the Arcminute MicroKelvin Imager (AMI, Zwart et al. 2008; Hickish et al. 2018). These observations were triggered by the detection of an increase in the emission from Cygnus X-3 seen at 37 GHz with the Metsähovi Radio Observatory (Karri Koljonen, private communication). The observations were made with the AMI ‘Small Array’ which is a radio interferometer consisting of ten 3.7-m diameter antennas. A single linear polarisation, Stokes parameter $I + Q$, was observed over a frequency range of 13.0 to 18.0 GHz. However, in practice the ends of this frequency range were not used (either due to poor sensitivity or, at the lower frequencies, satellite interference), and the observed band was 14.2 to 17.4 GHz.

The observations consisted of 1-hour integrations on Cygnus X-3, interleaved with 400-s observations of a nearby, compact calibrator source J2052+3635. The first observation was made on 2020 Feb 7th, for ≈ 6.4 hours. Long observations (up to ≈ 9.7 hours, after flagging) were made on all but one of the following days up to and including

2020 Feb 15th; the exception was Feb 9th, when only a short observation (≈ 1.5 hours) was possible, due to extremely high winds, which meant the array had to be stowed. The number of antennas available during these observations was usually 8 (except on Feb 9th and 13th when the number was, respectively, 7 and 9).

The data were processed using standard procedures, with the flux density scale established from short observations of the standard calibrator source 3C286 which were made most days, together with the ‘rain gauge’ measurements made during the observations which were used to correct for varying atmospheric conditions (see Zwart et al.). The data were flagged: (i) automatically to eliminate bad data due to various technical problems, interference, and when some antennas were shadowed at the end of the observations (at low elevations); (ii) manually, to eliminate remaining interference, and some periods with very heavy rain. The interleaved observations of J2052+3635 provided the initial phase calibration of each antenna in the array throughout each observation. The amplitudes of the J2052+3635 observations were used to check the consistency of the flux density scale during the observations. It was found that the r.m.s. deviation of the J2052+3635 flux densities was 1.8 per cent. Subsequently the observations were phase self-calibrated on a timescale of 10 min. Flux densities were derived for 10-min averages, for 5 broad frequency channels covering 14.2 to 17.4 GHz, and then a power law fit was made to obtain a flux density at 15 GHz.

Figure 1 shows the 15-GHz light curve of Cygnus X-3 from these observations. This shows a rapid increase on Feb 7th, rising from ≈ 2 Jy to a peak of ≈ 10 Jy in just over 4 hours. During this period the radio spectrum across the AMI




Figure 1. AMI Small Array observations of Cygnus X-3 during the period 2020 February 7th to 15th, at 15 GHz (Stokes $I + Q$). Each data point is a 10-min average. Statistical error bars are plotted, although these are usually smaller than the size of symbols.

band changed from optically thick (flux density rising with increasing frequency) to optically thin (flux density falling with increasing frequency) synchrotron emission. In subsequent days Cygnus X-3 showed other peaks at ≈ 11 Jy on Feb 8th, and ≈ 14 Jy on Feb 11th, after which it faded, approximately exponentially (with a characteristic time of 1.7 days).

We thank the staff of the Mullard Radio Astronomy Observatory, University of Cambridge, for their support in the maintenance, and operation of AMI. We acknowledge support from the European Research Council under grant ERC-2012-StG-307215 LODESTONE.

REFERENCES

- Corbel, S., Dubus, G., Tomsick, J. A., et al. 2012, *MNRAS*, 421, 2947
- Egron, E., Pellizzoni, A., Giroletti, M., et al. 2017, *MNRAS*, 471, 2703
- Fender, R. P., Bell Burnell, S. J., Waltman, E. B., et al. 1997, *MNRAS*, 288, 849
- Gregory, P. C., & Kronberg, P. P. 1972, *Nature*, 239, 440
- Hickish, J., Razavi-Ghods, N., Perrott, Y. C., et al. 2018, *MNRAS*, 475, 5677
- Johnston, K. J., Spencer, J. H., Simon, R. S., et al. 1986, *ApJ*, 309, 707
- Mioduszewski, A. J., Rupen, M. P., Hjellming, R. M., et al. 2001, *ApJ*, 553, 766
- Waltman, E. B., Ghigo, F. D., Johnston, K. J., et al. 1995, *AJ*, 110, 290
- Zdziarski, A. A., Segreto, A., & Pooley, G. G. 2016, *MNRAS*, 456, 775
- Zwart, J. T. L., Barker, R. W., Biddulph, P., et al. 2008, *MNRAS*, 391, 1545