A critical branching process with immigration in random environment^{*}

Afanasyev V.I.

Steklov Mathematical Institute of Russian Academy of Sciences, 8 Gubkin St., Moscow, 119991, Russia Email: viafan@mi.ras.ru

Abstract

A Galton-Watson branching process with immigration evolving in a random environment is considered. Its associated random walk is assumed to be oscillating. We prove a functional limit theorem in which the process under consideration is normalized by a random coefficient depending on the random environment only. The distribution of the limiting process is described in terms of a strictly stable Levy process and a sequence of independent and identically distributed random variables which is independent of this process.

Keywords: Branching process in random environment, branching process with immigration, functional limit theorem

1. Introduction and statement of main result

Let $(\Omega, \mathcal{F}, \mathbf{P})$ be a probability space and Δ be the space of probability measures on $N_0 := \{0, 1, \ldots\}$ equipped with the metric of total variation. *A random environment* is a sequence of random elements Q_1, Q_2, \ldots , mapping the space $(\Omega, \mathcal{F}, \mathbf{P})$ into Δ^2 . Thus, Q_n for each $n \in \mathbf{N}$ has the form (F_n, G_n) , where F_n, G_n are probability measures on N0. *A branching process with immigration in random environment* ((BPIRE)) is a stochastic process possessing the following properties. For a fixed random environment $\{Q_n, n \in \mathbb{N}\}\)$ this is an inhomogeneous branching Galton-Watson process with immigration (see [\[1\]](#page-31-0), Chapter 6, § 7). Here, for each $n \in \mathbb{N}$, the number of immigrants joining the $(n-1)$ th gen-

[✩]This work is supported by the Russian Science Foundation under grant 19-11-00111.

eration has the distribution G_n and the offspring reproduction law of particles of the $(n-1)$ th generation is F_n .

Let Z_n be the size of nth generation without the immigrants which joined this generation (we assume that $Z_0 = 0$), η_n be the number of immigrants which joined the *n*th generation. Let $f_n(\cdot)$ and $g_n(\cdot)$ be generating functions of distributions F_n and G_n respectively.

We consider this model under the assumption that the random elements Q_1, Q_2, \ldots are independent and identically distributed. A more detailed definition of the BPIRE can be found in [\[2\]](#page-32-0).

Set for $i \in \mathbb{N}$

$$
X_i = \ln f'_i(1), \qquad \mu_i = g'_i(1)
$$

(suppose that $0 < f'_1(1) < +\infty$, $0 < g'_i(1) < +\infty$ a.s.). Introduce the so-called *associated random walk*:

$$
S_0 = 0,
$$
 $S_n = \sum_{i=1}^n X_i, n \in \mathbb{N}.$

It is clear that the random vectors $(X_1, \mu_1), (X_2, \mu_2), \ldots$ are independent and identically distributed under our assumptions.

We impose the following restriction on the distribution of X_1 .

Hypothesis A. The distribution of X_1 belongs without centering to the domain of attraction of some stable law with index $\alpha \in (0, 2]$ and the limit law is not a one-sided stable law.

Under Hypothesis A the Skorokhod functional limit theorem is valid (see, for instance, [\[3\]](#page-32-1), Chapter 16): there are such positive normalizing constants C_n that, as $n \to \infty$,

$$
W_n \stackrel{D}{\to} W,\tag{1}
$$

where $W_n = \left\{C_n^{-1}S_{\lfloor nt \rfloor}, t \ge 0\right\}$, the process $W = \left\{W(t), t \ge 0\right\}$ is a strictly stable Levy process with index $\alpha \in (0, 2]$ and the symbol $\stackrel{D}{\rightarrow}$ means convergence in distribution in the space $D[0, +\infty)$ with Skorokhod topology. Moreover,

$$
C_n = n^{1/\alpha} l(n),
$$

where $\{l(n), n \in \mathbb{N}\}\$ is a slowly varying sequence. It is known that the finitedimensional distributions of the process W are absolutely continuous. Note that $\rho := \mathbf{P}(W(1) > 0) \in (0, 1)$ given Hypothesis A. Thus, the Spitzer-Doney condition is satisfied:

$$
\lim_{n \to \infty} \mathbf{P}(S_n > 0) = \rho \in (0, 1). \tag{2}
$$

The Spitzer-Doney condition means that the random walk $\{S_n\}$ is oscillating. As result, the absolute values of its strict descending ladder heights constitute a renewal process with the corresponding renewal function $v(x)$, $x \geq 0$ (see [\[4\]](#page-32-2) for a detailed definition of the function $v(\cdot)$). Similarly, weak ascending ladder heights of the random walk $\{S_n\}$ generate a renewal process with the corresponding renewal function $u(x)$, $x \geq 0$.

The aim of this paper is to prove a functional limit theorem for the process $\{Z_{\lfloor nt \rfloor}, t \ge 0\}$, as $n \to \infty$ (see Theorem 1).

We need some notation and definitions to formulate the theorem. Let for $n\in\mathbf{N}$

$$
M_n = \max_{1 \le i \le n} S_i, \qquad L_n = \min_{0 \le i \le n} S_i.
$$

It is known (see, for instance, [\[4](#page-32-2)], Lemma 2.5) that, if the Spitzer-Doney condition (2) is satisfied, then, as $n \to \infty$,

$$
\left\{ (Q_i, S_i, \mu_i), i \in \mathbf{N} \mid L_n \ge 0 \right\} \stackrel{D}{\to} \left\{ \left(Q_i^+, S_i^+, \mu_i^+ \right), i \in \mathbf{N} \right\},\tag{3}
$$

$$
\left\{ (Q_i, S_i, \mu_i), i \in \mathbf{N} | M_n < 0 \right\} \stackrel{D}{\to} \left\{ \left(Q_i^-, S_i^-, \mu_i^- \right), i \in \mathbf{N} \right\},\tag{4}
$$

where $\{(Q_i^+,S_i^+,\mu_i^+)\},\{(Q_i^-,S_i^-, \mu_i^-)\}$ are some random sequences. Moreover: a) the sequences $\{Q_i^+, i \in \mathbb{N}\}, \{Q_i^-, i \in \mathbb{N}\}$ can be viewed as some random environments; b) the sequences $\{S_i^+, i \in \mathbb{N}\}, \{S_i^-, i \in \mathbb{N}\}$ are the corresponding associated random walks $(S_0^+ = S_0^- = 0)$; c) the sequences $\{\mu_i^+, i \in \mathbb{N}\}\$ and $\left\{\mu_i^-, \, i \in \mathbf{N}\right\}$ are positive and constructed by $\left\{Q_i^+, \, i \in \mathbf{N}\right\}$ and $\left\{Q_i^-, \, i \in \mathbf{N}\right\}$, respectively, the same as the sequence $\{\mu_i, i \in \mathbf{N}\}\)$ is constructed by $\{Q_i, i \in \mathbf{N}\}\$. Suppose that the sequences $\{Q_i^+, i \in \mathbb{N}\}, \{Q_i^-, i \in \mathbb{N}\}$ are defined on the same probability space $(\Omega^*, \mathcal{F}^*, \mathbf{P}^*)$ and are independent (below we denote the expectation on this probability space by \mathbf{E}^*).

We now come back to our initial BPIRE. Set $N_i = \{i, i+1, ...\}$ for $i \in \mathbb{Z}$. Fix $i \in \mathbb{N}_0$ and, for $n \in \mathbb{N}_i$, denote by $Z_{i,n}$ the total number of particles in the nth generation which are the descendants of the immigrants joined the ith generation (we assume that $Z_{i,n} = 0$ for $i \geq n$ and $i < 0$). Note that the random sequence $\{\eta_i; Z_{i,n}, n \in \mathbf{N}_{i+1}\}\$ is a usual (without immigration) branching process in the random environment $\{G_{i+1}; F_n, n \in \mathbb{N}_{i+1}\}$. In particular, if the random environment is fixed, then G_{i+1} is the distribution of the random variable η_i which should be interpreted as the number of particles in the initial generation. Set for $n \in \mathbf{N}_i$

$$
a_{i,n} = e^{-(S_n - S_i)}.
$$

The sequence $\{\eta_i; a_{i,n}\mathbb{Z}_{i,n}, n \in \mathbb{N}_{i+1}\}\$ is a nonnegative martingale if the random environment $\{G_{i+1}; F_n, n \in \mathbb{N}_{i+1}\}$ is fixed. Hence (without assuming that the random environment is fixed), there is a finite limit $\lim_{n\to\infty} a_{i,n} Z_{i,n}$ **P**-a.s.

Set

$$
\begin{aligned} Q_i^* &= \left\{ \begin{array}{cl} Q_i^+, \quad & i \in \mathbf{N}, \\ Q_{-i+1}^-, \quad & i \in \mathbf{Z} \setminus \mathbf{N}, \\ S_i^* &= \left\{ \begin{array}{cl} S_i^+, \quad & i \in \mathbf{N}_0, \\ -S_{-i}^-, \quad & i \in \mathbf{Z} \setminus \mathbf{N}_0, \\ \mu_i^* &= \left\{ \begin{array}{cl} \mu_i^+, \quad & i \in \mathbf{N}, \\ \mu_{-i+1}^-, \quad & i \in \mathbf{Z} \setminus \mathbf{N}. \end{array} \right. \end{aligned} \right. \end{aligned}
$$

The sequence $\mathcal{E}^* := \{Q_k^*, k \in \mathbf{Z}\}\$ can be considered as a random environment (we denote the components of Q_k^* by G_k^* and F_k^*). We assume that the probability space $(\Omega^*, \mathcal{F}^*, \mathbf{P}^*)$ is reach enough for we are able to define on it a branching process with immigration in the random environment \mathcal{E}^* . Fix $i \in \mathbf{Z}$ and, for $j \in \mathbf{N}_i$, denote by $Z^*_{i,j}$ the total number of particles in the jth generation being descendants of immigrants which joined the ith generation (we denote the number of such immigrants as η_i^*). Note that the sequence $\{\eta_i^*; Z_{i,j}^*, j \in \mathbf{N}_{i+1}\}\$ is a branching process in the random environment $\{G_{i+1}^*; F_j^*, j \in \mathbb{N}_{i+1}\}\$ with the initial value η_i^* . The sequence $\{S_j^* - S_i^*, j \in \mathbf{N}_i\}$ is the associated random walk and the random variable μ_i^* is under fixed environment the mean of the random variable η_i^* . Set

$$
a_{i,j}^* = e^{-\left(S_j^* - S_i^*\right)}.
$$

In accordance with the above the limit

$$
\lim_{j \to \infty} a_{i,j}^* Z_{i,j}^* =: \zeta_i^* \tag{5}
$$

exists \mathbf{P}^* -a.s. and \mathbf{P}^* $(\zeta_i^* > 0) > 0$ for $i \in \mathbf{N}_0$ (see [\[4\]](#page-32-2), Proposition 3.1).

Introduce the following random series:

$$
\Sigma_1:=\sum_{i\in{\mathbf{Z}}}\mu^*_{i+1}e^{-S^*_i},\qquad \Sigma_2:=\sum_{i\in{\mathbf{Z}}}\zeta^*_i e^{-S^*_i}
$$

It is clear that $\Sigma_1 > 0$ **P**^{*}-a.s. and **P**^{*} ($\Sigma_2 > 0$) > 0. Both series converge P[∗] -a.s. under certain restrictions (see Lemma 4).

Let W be a strictly stable Levy process with index α (in the sequel we call W simply *the Levy process*). By the Levy process we specify *the (lower) level* $L = \{L(t), t \geq 0\}$ *of the Levy process* as

$$
L(t) = \inf_{s \in [0,t]} W(s).
$$

Let, further, $\gamma_1, \gamma_2, \ldots$ be an independent of W sequence of independent random variables distributed as the random variable Σ_2/Σ_1 .

By these ingredients we define finite-dimensional distributions of a random process $Y = \{Y(t), t \ge 0\}$ which plays an important role in the sequel. First we set $Y(0) = 0$. Consider an arbitrary $m \in \mathbb{N}$ and arbitrary moments t_1, t_2, \ldots, t_m : $0 = t_0 < t_1 < t_2 < \ldots < t_m$. The random vector $\{Y(t_1),\ldots,Y(t_m)\}$ coincides in distribution with the following vector $\widehat{Y}\;:=\;$ $\{\hat{Y}_1,\ldots,\hat{Y}_m\}$. We describe at first the possible values of the vector \hat{Y} . Its first several coordinates coincide with γ_1 , the next several coordinates coincide with γ_2 and so on up to the mth coordinate. The coordinates of the vector \hat{Y} are specified according to the level L of the Levy process W . The first coordinate \widehat{Y}_1 is equal to γ_1 . Let the coordinate \widehat{Y}_k for some $k < m$ be known. For instance, $\widehat{Y}_k = \gamma_l$ for some $l \in \mathbb{N}$. If the level of the Levy process at the moment t_{k+1}

remains the same as at moment t_k , i.e. $L(t_{k+1}) = L(t_k)$, then $Y_{k+1} = \gamma_l$. If the level of the Levy process at the moment t_{k+1} is changed, i.e. $L(t_{k+1}) < L(t_k)$, then $\hat{Y}_{k+1} = \gamma_{l+1}$.

Set for $n \in \mathbb{N}_0$

$$
a_n = e^{-S_n}
$$
, $b_n = \sum_{i=0}^{n-1} \mu_{i+1} e^{-S_i}$ $(b_0 = 0)$.

Introduce for each $n \in \mathbb{N}$ the random process $Y_n = \{Y_n(t), t \geq 0\}$, where

$$
Y_n(0) = 0, \qquad Y_n(t) = \frac{a_{\lfloor nt \rfloor}}{b_{\lfloor nt \rfloor}} Z_{\lfloor nt \rfloor}.
$$

Note that for $k \in \mathbb{N}$ the ratio b_k/a_k is equal to the mean of Z_k for a fixed random environment.

Let the symbol \Rightarrow means convergence of random processes in the sense of finite-dimensional distributions and $\ln^+ x = \max(0, \ln x)$ for $x > 0$.

Theorem 1. If Hypothesis A is valid and $\mathbf{E} (\ln^+ \mu_1)^{\alpha+\varepsilon} < +\infty$ for some $\varepsilon > 0$, then, as $n \to \infty$,

$$
Y_n \Rightarrow Y.
$$

A detailed description of the theory of critical (when Hypothesis A is valid) branching processes in random environment is available in [\[4\]](#page-32-2) and [\[5](#page-32-3)].

A particular case of a subcritical BPIRE (when the offspring generating function $f_n(\cdot)$ is fractional-linear and $g_n(s) \equiv s$ for each $n \in \mathbb{N}$) was considered in [\[6\]](#page-32-4). The main attention there was paid to obtaining an exponential estimate for the tail distribution of the so-called life period of this process (i.e., the time until the first extinction). A more general class of subcritical BPIRE was analyzed in [\[7](#page-32-5)] where a limit theorem describing the population size at a distant moment was proved and an exponential estimate for the tail distribution of the life period was established. A strong law of large numbers and a central limit theorem for a wide class of subcritical BPIRE were proved in [\[8](#page-32-6)].

A critical BPIRE was considered in [\[9\]](#page-32-7) where sufficient conditions of transience and recurrence were obtained. The author of [\[10\]](#page-32-8), studying a random walk in random environment, proved a particular case of Theorem 1 (when the offspring generating function $f_n(\cdot)$ is fractional-linear and $g_n(s) \equiv s$ for each $n \in \mathbb{N}$). We would like to stress that the proof used in the present paper differs significantly from that one given in [\[10](#page-32-8)]. We also mention the papers [\[11\]](#page-32-9), [\[12\]](#page-33-0) and [\[13](#page-33-1)] in which critical and supercritical processes (with stopped immigration) are considered under some restrictions on their lifetime.

Recent papers [\[2\]](#page-32-0) and [\[14\]](#page-33-2) contain exact asymptotic formulae for the tail distribution of the life period for critical and subcritical BPIRE.

2. Auxiliary statements

Let τ_n be the first moment when the minimum of the random walk S_0, \ldots, S_n is attained:

$$
\tau_n = \min \{ i : S_i = L_n, \, 0 \le i \le n \} \, .
$$

Set for $n\in{\bf N}$

$$
S'_{i,n} = \left\{ \begin{array}{cl} S_{\tau_n+i} - S_{\tau_n}, & i \in \mathbf{N}_{(-\tau_n)}, \\ 0, & i \in \mathbf{Z} \setminus \mathbf{N}_{(-\tau_n)}. \end{array} \right.
$$

For positive integers numbers $n_1 < n_2$ set

$$
L_{n_1,n_2} = \min_{n_1 \le i \le n_2} S_i.
$$

Lemma 1. *If the Spitzer-Doney condition* (2) *is satisfied, then, as* $n \to \infty$,

$$
\left\{S_{i,n}^{\prime}, i \in \mathbf{Z}\right\} \stackrel{D}{\to} \left\{S_{i}^{*}, i \in \mathbf{Z}\right\}.
$$
 (6)

Proof. We demonstrate for simplicity only convergence of one-dimensional distributions. Fix $i \in \mathbb{N}_0$. Let A be a one-dimensional S_i^* -continuous (relative to the measure \mathbf{P}^*) Borel set. Then for $n \geq i$

$$
\mathbf{P}\left(S'_{i,n} \in A, \, \tau_n + i \leq n\right)
$$
\n
$$
= \sum_{k=0}^{n-i} \mathbf{P}\left(S'_{i,n} \in A, \, \tau_n = k\right)
$$

$$
= \sum_{k=0}^{n-i} \mathbf{P}(S_{k+i} - S_k \in A, S_k < L_{k-1}, S_k \le L_{k+1,n})
$$

and by the Markov property of random walks we have that

$$
\mathbf{P}(S_{k+i} - S_k \in A, S_k < L_{k-1}, S_k \le L_{k+1,n})
$$
\n
$$
= \mathbf{P}(S_k < L_{k-1}) \mathbf{P}(S_{k+i} - S_k \in A, S_k \le L_{k+1,n})
$$
\n
$$
= \mathbf{P}(S_k < L_{k-1}) \mathbf{P}(S_i \in A, L_{n-k} \ge 0)
$$
\n
$$
= \mathbf{P}(S_i \in A | L_{n-k} \ge 0) \mathbf{P}(S_k < L_{k-1}) \mathbf{P}(L_{n-k} \ge 0)
$$
\n
$$
= \mathbf{P}(S_i \in A | L_{n-k} \ge 0) \mathbf{P}(\tau_n = k).
$$

Thus,

$$
\mathbf{P}\left(S'_{i,n} \in A, \, \tau_n + i \leq n\right) = \sum_{k=0}^{n-i} \mathbf{P}\left(S_i \in A \,|\, L_{n-k} \geq 0\right) \mathbf{P}\left(\tau_n = k\right). \tag{7}
$$

If the Spitzer-Doney condition is satisfied, then the following generalized arcsine law is valid (see, for instance, [\[15](#page-33-3)], Chapter 8, Theorem 8.9.9): for $x \in [0,1]$

$$
\lim_{n \to \infty} \mathbf{P}\left(\frac{\tau_n}{n} \le x\right) = \frac{\sin(\pi \rho)}{\pi} \int_{0}^{x} u^{\rho - 1} \left(1 - u\right)^{-\rho} du. \tag{8}
$$

We pass to the limit in formula (7), as $n \to \infty$. Due to (8)

$$
\lim_{n \to \infty} \mathbf{P}(\tau_n + i \le n) = \lim_{n \to \infty} \mathbf{P}(\tau_n/n \le 1 - i/n) = 1.
$$

Therefore the limit of the left-hand side of (7) coincides with the limit of probability $\mathbf{P}(S'_{i,n} \in A)$, as $n \to \infty$, if at least one of these limits exists.

If $\varepsilon \in (0,1)$ and n is large enough, then by (7)

$$
\mathbf{P}\left(S'_{i,n}\in A,\,\tau_n+i\leq n\right)=P_1\left(n,\varepsilon\right)+P_2\left(n,\varepsilon\right),\tag{9}
$$

where

$$
P_1(n,\varepsilon) = \sum_{k=0}^{\lfloor (1-\varepsilon)n \rfloor} \mathbf{P}\left(S_i \in A \mid L_{n-k} \ge 0\right) \mathbf{P}\left(\tau_n = k\right),
$$

$$
P_2(n,\varepsilon) = \sum_{k=\lfloor (1-\varepsilon)n \rfloor+1}^{n-i} \mathbf{P}\left(S_i \in A \mid L_{n-k} \ge 0\right) \mathbf{P}\left(\tau_n = k\right).
$$

Clearly,

$$
P_2(n,\varepsilon) \leq \sum_{k=\lfloor(1-\varepsilon)n\rfloor+1}^n \mathbf{P}(\tau_n = k) = \mathbf{P}(\tau_n > \lfloor(1-\varepsilon)n\rfloor)
$$

$$
\xrightarrow{n\to\infty} 1 - \frac{\sin(\pi\rho)}{\pi} \int_0^{1-\varepsilon} u^{\rho-1} (1-u)^{-\rho} du \xrightarrow{\varepsilon \to 0} 0.
$$

Therefore

$$
\lim_{\varepsilon \to 0} \limsup_{n \to \infty} P_2(n, \varepsilon) = 0.
$$
\n(10)

In view of (3) the probability $\mathbf{P}\left(S_i \in A \mid L_{n-k} \geq 0\right)$ tends, as $n \to \infty$, to \mathbf{P}^* $(S_i^* \in A) = \mathbf{P}^*$ $(S_i^* \in A)$ uniformly over $0 \le k \le \lfloor (1 - \varepsilon) n \rfloor$. Consequently,

$$
\lim_{n \to \infty} P_1(n, \varepsilon) = \mathbf{P}^* (S_i^* \in A) \lim_{n \to \infty} \sum_{k=0}^{\lfloor (1-\varepsilon)n \rfloor} \mathbf{P} (\tau_n = k)
$$

$$
= \mathbf{P}^* (S_i^* \in A) \lim_{n \to \infty} \mathbf{P} (\tau_n \le \lfloor (1-\varepsilon)n \rfloor)
$$

$$
= \mathbf{P}^* (S_i^* \in A) \frac{\sin(\pi \rho)}{\pi} \int_{0}^{1-\varepsilon} u^{\rho-1} (1-u)^{-\rho} du
$$

$$
\xrightarrow{\varepsilon \to 0} \mathbf{P}^* (S_i^* \in A)
$$

implying

$$
\lim_{\varepsilon \to 0} \lim_{n \to \infty} P_1(n, \varepsilon) = \mathbf{P}^* \left(S_i^* \in A \right). \tag{11}
$$

It follows from relations (9)-(11) that for $i \in \mathbb{N}_0$

$$
\lim_{n \to \infty} \mathbf{P}\left(S'_{i,n} \in A, \, \tau_n + i \leq n\right) = \mathbf{P}^*\left(S_i^* \in A\right).
$$

Thus,

$$
\lim_{n \to \infty} \mathbf{P} \left(S'_{i,n} \in A \right) = \mathbf{P}^* \left(S_i^* \in A \right). \tag{12}
$$

We now fix $i \in \mathbb{N}$. Let A be a one-dimensional S^*_{-i} -continuous (relative to the measure \mathbf{P}^*) Borel set. Then for $n \geq i$

$$
\mathbf{P}\left(S'_{-i,n} \in A, \tau_n - i \ge 0\right)
$$

$$
= \sum_{k=i}^{n} \mathbf{P}\left(S'_{-i,n} \in A, \tau_n = k\right)
$$

$$
= \sum_{k=i}^{n} \mathbf{P}(S_{k-i} - S_k \in A, S_k < L_{k-1}, S_k \le L_{k+1,n})
$$

and by the Markov property and the duality property of random walks we have that

$$
\mathbf{P}(S_{k-i} - S_k \in A, S_k < L_{k-1}, S_k \le L_{k+1,n})
$$
\n
$$
= \mathbf{P}(S_{k-i} - S_k \in A, S_k < L_{k-1}) \mathbf{P}(S_k \le L_{k+1,n})
$$
\n
$$
= \mathbf{P}(-S_i \in A, M_k < 0) \mathbf{P}(S_k \le L_{k+1,n})
$$
\n
$$
= \mathbf{P}(-S_i \in A | M_k < 0) \mathbf{P}(M_k < 0) \mathbf{P}(S_k \le L_{k+1,n})
$$
\n
$$
= \mathbf{P}(-S_i \in A | M_k < 0) \mathbf{P}(\tau_n = k).
$$

Thus,

$$
\mathbf{P}\left(S'_{-i,n} \in A, \, \tau_n - i \ge 0\right) = \sum_{k=i}^{n} \mathbf{P}\left(-S_i \in A \,|\, M_k < 0\right) \mathbf{P}\left(\tau_n = k\right),\tag{13}
$$

therefore, if $\varepsilon \in (0,1)$ and n is large enough, then

$$
\mathbf{P}\left(S'_{-i,n} \in A, \, \tau_n - i \geq 0\right) = P_3\left(n, \varepsilon\right) + P_4\left(n, \varepsilon\right),\,
$$

where

$$
P_3(n,\varepsilon) = \sum_{k=i}^{\lfloor \varepsilon n \rfloor} \mathbf{P}(-S_i \in A \mid M_k < 0) \mathbf{P}(\tau_n = k),
$$
\n
$$
P_4(n,\varepsilon) = \sum_{k=\lfloor \varepsilon n \rfloor + 1}^n \mathbf{P}(-S_i \in A \mid M_k < 0) \mathbf{P}(\tau_n = k).
$$

It is not difficult to show (see our proof of relation (10)) that

$$
\lim_{\varepsilon \to 0} \limsup_{n \to \infty} P_3(n, \varepsilon) = 0.
$$

Due to (4) the probability $\mathbf{P}(-S_{-i} \in A | M_k < 0)$ tends, as $n \to \infty$, to $\mathbf{P}^*(-S_{-i}^- \in A) = \mathbf{P}^* (S_{-i}^* \in A)$ uniformly over $\lfloor \varepsilon n \rfloor < k \leq n$. Therefore

$$
\lim_{\varepsilon \to 0} \lim_{n \to \infty} P_4(n, \varepsilon) = \mathbf{P}^* \left(S_{-i}^* \in A \right).
$$

As result, we obtain that

$$
\lim_{n \to \infty} \mathbf{P} \left(S'_{-i,n} \in A, \, \tau_n - i \ge 0 \right) = \mathbf{P}^* \left(S^*_{-i} \in A \right)
$$

proving (12) for $i \in \mathbf{Z} \backslash \mathbf{N}_0$. Thus, convergence of one-dimensional distributions in (6) is established.

The lemma is proved.

Remark 1. It is not difficult to verify (see [\[16\]](#page-33-4), Lemma 1) that relation (6) admits the following generalization: for any $a \leq 0$ and $b > 0$, as $n \to \infty$,

$$
\left\{S'_{i,n}, i \in \mathbf{Z} \middle| \frac{L_n}{C_n} \le a, \frac{S_n - L_n}{C_n} \le b \right\} \stackrel{D}{\to} \left\{S_i^*, i \in \mathbf{Z} \right\}.
$$

Recall that (Ω, \mathcal{F}, P) is the underlying probability space. Set

$$
I_n^{(2)} := \{(i, j) : i, j \in \{0, \dots, n\} \text{ and } i \leq j\}.
$$

Let \mathcal{F}_n , $n \in \mathbb{N}$, denote the σ -algebra generated by the segment of the random environment Q_1, \ldots, Q_n and the random variables $Z_{i,j}$ for $(i,j) \in I_n^{(2)}$. We now introduce a probability measure \mathbf{P}^+ on the σ -algebra $\mathcal{F}_{\infty} := \sigma(\bigcup_{n=1}^{\infty} \mathcal{F}_n)$, defined for each $n\in \mathbf{N}_0$ and each $\mathcal{F}_n\text{-measurable nonnegative random variable}$ β by the formula

$$
\mathbf{E}^{+}\beta = \mathbf{E}\left(\beta v\left(S_{n}\right); L_{n} \geq 0\right). \tag{14}
$$

This may require a change of the underlying probability space (see [\[4\]](#page-32-2) for more details). Similarly, we also introduce a probability measure \mathbf{P}^- on the σ -algebra \mathcal{F}_{∞} , defined for each $n \in \mathbb{N}_0$ and each \mathcal{F}_n -measurable nonnegative random variable β by the formula

$$
\mathbf{E}^{-}\beta = \mathbf{E}\left(\beta u\left(-S_{n}\right); M_{n} < 0\right). \tag{15}
$$

Recall that the functions $v(\cdot)$ and $u(\cdot)$ in formulae (14) and (15) are defined after relation (2). Thus, three measures P, P^+, P^- are defined on one and the same measurable space $(\Omega, \mathcal{F}_{\infty})$. To explicitly indicate the measure on $(\Omega, \mathcal{F}_{\infty})$ according to which we consider this or those random elements we use the measure symbol as a lower index.

For instance, it is shown in Lemma 2.5 from [\[4\]](#page-32-2) that

$$
\{(Q_i^+, S_i^+, \mu_i^+), i \in \mathbf{N}\} \stackrel{D}{=} \{(Q_i, S_i, \mu_i), i \in \mathbf{N}\}_{\mathbf{P}^+}
$$
(16)

(the lower index \mathbf{P}^+ for a random sequence shows here that the measure \mathbf{P}^+ is used on the space $(\Omega, \mathcal{F}_{\infty})$). Similarly,

$$
\{(Q_i^-, S_i^-, \mu_i^-), i \in \mathbf{N}\} \stackrel{D}{=} \{(Q_i, S_i, \mu_i), i \in \mathbf{N}\}_{\mathbf{P}^-}.
$$
 (17)

Due to (16) , (17) and our assumption about the independence of the lefthand sides of these relations, the product of probability spaces $(\Omega, \mathcal{F}_{\infty}, \mathbf{P}^+)$ and $(\Omega, \mathcal{F}_{\infty}, \mathbf{P}^{-})$ may be considered as a probability space $(\Omega^{*}, \mathcal{F}^{*}, \mathbf{P}^{*})$ and, consequently, the direct product of the measures \mathbf{P}^+ and \mathbf{P}^- may be treated as the measure P[∗] .

Remark 2. If a random element ξ is given on the space $(\Omega, \mathcal{F}_{\infty}, \mathbf{P}^+)$ we can define the random element ξ^+ , specified on the product of the spaces $(\Omega, \mathcal{F}_{\infty}, \mathbf{P}^+)$ and $(\Omega, \mathcal{F}_{\infty}, \mathbf{P}^-)$ by means of the formula $\xi^+(\omega_1, \omega_2) = \xi(\omega_1)$ for $(\omega_1, \omega_2) \in \Omega \times \Omega$. It is clear that $\mathbf{P}^*(\xi^+ \in A) = \mathbf{P}^+(\xi \in A)$ for an arbitrary one-dimensional Borel set A. Similarly, if a random element ξ is given on the space $(\Omega, \mathcal{F}_{\infty}, \mathbf{P}^-)$ we can define the random element ξ^- , specified on the product of the spaces $(\Omega, \mathcal{F}_\infty, \mathbf{P}^+)$ and $(\Omega, \mathcal{F}_\infty, \mathbf{P}^-)$ by means of the formula $\xi^-(\omega_1,\omega_2)=\xi(\omega_2)$ for $(\omega_1,\omega_2)\in\Omega\times\Omega$, and $\mathbf{P}^*(\xi^-\in A)=\mathbf{P}^-(\xi\in A)$ for an arbitrary one-dimensional Borel set A. In accordance with the agreement we can consider the random elements standing in the left-hand sides of formulae (16) and (17) as generated by the random elements $\{(Q_i, S_i, \mu_i), i \in \mathbf{N}\}_{\mathbf{P}^+}$ and $\{(Q_i, S_i, \mu_i), i \in \mathbf{N}\}_{\mathbf{P}^-}$ respectively.

Lemma 2. If the Spitzer-Doney condition (2) is satisfied, then, as $n \to \infty$,

$$
\{a_{i,n}Z_{i,n}, i \in \mathbf{N}_0 \mid L_n \ge 0\} \stackrel{D}{\to} \{\zeta_i^*, i \in \mathbf{N}_0\},\tag{18}
$$

where $\{\zeta_i^*, i \in \mathbb{N}_0\}$ *is the random sequence defined by relation* (5).

Proof. By virtue of the first part of Lemma 2.5 from [\[4](#page-32-2)] for $k \in \mathbb{N}$, as $n \to \infty$,

.

$$
\left\{ (a_{i,j}, Z_{i,j}), (i,j) \in I_k^{(2)} \middle| L_n \geq 0 \right\} \stackrel{D}{\to} \left\{ (a_{i,j}, Z_{i,j}), (i,j) \in I_k^{(2)} \right\}_{\mathbf{P}^+}
$$

Note (see [\[4](#page-32-2)], Section 3) that in view of (14) for a fixed $i \in \mathbb{N}_0$ the random sequence $\{\eta_i; Z_{i,j}, j \in \mathbb{N}_{i+1}\}_{\mathbf{P}^+}$ given on the probability space $(\Omega, \mathcal{F}_{\infty}, \mathbf{P}^+)$ is a branching process in the random environment $\{G_{i+1}; F_n, n \in \mathbb{N}_{i+1}\}_{\mathbb{P}^+}$. Hence, if the random environment is fixed, the sequence $\{\eta_i; a_{i,j}Z_{i,j}, j \in \mathbf{N}_{i+1}\}_{\mathbf{P}^+}$ is a non-negative martingale. Because of this (without assuming that the random environment is fixed) there is \mathbf{P}^+ -a.s. the finite limit

$$
\lim_{n \to \infty} a_{i,n} Z_{i,n} =: \zeta_i.
$$

It means, in view of the second part of Lemma 2.5 from [\[4\]](#page-32-2), that, as $n \to \infty$,

$$
\{a_{i,n}Z_{i,n}, i \in \mathbf{N}_0 \mid L_n \geq 0\} \stackrel{D}{\rightarrow} \{\zeta_i, i \in \mathbf{N}_0\}_{\mathbf{P}^+}.
$$

To prove relation (18), it remains to note that in view of Remark 2

$$
\{\zeta_i, i \in \mathbf{N}_0\}_{\mathbf{P}^+} \stackrel{D}{=} \{\zeta_i^*, i \in \mathbf{N}_0\}.
$$

The lemma is proved.

Set for $n\in{\bf N}$

$$
\begin{aligned} Z'_{i,n} &= \left\{ \begin{array}{ll} Z_{\tau_n+i,n}, & i \in \mathbf{N}_{(-\tau_n)}, \\ 0, & i \in \mathbf{Z} \setminus \mathbf{N}_{(-\tau_n)}, \end{array} \right. \\ a'_{i,n} &= \left\{ \begin{array}{ll} a_n/a_{\tau_n+i}, & i \in \mathbf{N}_{(-\tau_n)}, \\ 1, & i \in \mathbf{Z} \setminus \mathbf{N}_{(-\tau_n)}. \end{array} \right. \end{aligned}
$$

Lemma 3. *If the Spitzer-Doney condition* (2) *is satisfied, then, as* $n \to \infty$,

$$
\left\{a'_{i,n}Z'_{i,n}, i \in \mathbf{Z}\right\} \stackrel{D}{\to} \left\{\zeta_i^*, i \in \mathbf{Z}\right\},\tag{19}
$$

where $\{\zeta_i^*, i \in \mathbb{N}_0\}$ *is the random sequence defined by relation* (5).

Proof. We demonstrate for simplicity only convergence of one-dimensional distributions. Fix $i \in \mathbb{N}_0$. Let A be an arbitrary one-dimensional ζ_i^* -continuous (relative to the measure \mathbf{P}^*) Borel set. Then for $n \geq i$

$$
\mathbf{P}\left(a'_{i,n}Z'_{i,n} \in A, \, \tau_n + i \leq n\right)
$$
\n
$$
= \sum_{k=0}^{n-i} \mathbf{P}\left(a'_{i,n}Z'_{i,n} \in A, \, \tau_n = k\right)
$$
\n
$$
= \sum_{k=0}^{n-i} \mathbf{P}\left(\frac{a_n}{a_{k+i}}Z_{k+i,n} \in A, \, S_k < L_{k-1}, \, S_k \leq L_{k+1,n}\right)
$$

and

$$
\mathbf{P}\left(\frac{a_n}{a_{k+i}}Z_{k+i,n} \in A, S_k < L_{k-1}, S_k \le L_{k+1,n}\right)
$$
\n
$$
= \mathbf{P}\left(S_k < L_{k-1}\right) \mathbf{P}\left(\frac{a_n}{a_{k+i}}Z_{k+i,n} \in A, S_k \le L_{k+1,n}\right)
$$
\n
$$
= \mathbf{P}\left(S_k < L_{k-1}\right) \mathbf{P}\left(\frac{a_{n-k}}{a_i}Z_{i,n-k} \in A, L_{n-k} \ge 0\right)
$$
\n
$$
= \mathbf{P}\left(a_{i,n-k}Z_{i,n-k} \in A \mid L_{n-k} \ge 0\right) \mathbf{P}\left(S_k < L_{k-1}\right) \mathbf{P}\left(L_{n-k} \ge 0\right)
$$
\n
$$
= \mathbf{P}\left(a_{i,n-k}Z_{i,n-k} \in A \mid L_{n-k} \ge 0\right) \mathbf{P}\left(\tau_n = k\right).
$$

Thus,

$$
\mathbf{P}\left(a'_{i,n}Z'_{i,n} \in A, \, \tau_n + i \leq n\right) \n= \sum_{k=0}^{n-i} \mathbf{P}\left(a_{i,n-k}Z_{i,n-k} \in A \mid L_{n-k} \geq 0\right) \mathbf{P}\left(\tau_n = k\right).
$$

Therefore, if $\varepsilon \in (0,1)$ and n is large enough, then

$$
\mathbf{P}\left(a'_{i,n}Z'_{i,n}\in A,\,\tau_n+i\leq n\right)=P_1\left(n,\varepsilon\right)+P_2\left(n,\varepsilon\right),
$$

where

$$
P_1(n,\varepsilon) = \sum_{k=0}^{\lfloor (1-\varepsilon)n \rfloor} \mathbf{P}(a_{i,n-k} Z_{i,n-k} \in A | L_{n-k} \ge 0) \mathbf{P}(\tau_n = k),
$$

$$
P_2(n,\varepsilon) = \sum_{k=\lfloor (1-\varepsilon)n \rfloor+1}^{n-i} \mathbf{P}(a_{i,n-k} Z_{i,n-k} \in A | L_{n-k} \ge 0) \mathbf{P}(\tau_n = k).
$$

It is easy to show (see the proof of relation (10)) that

$$
\lim_{\varepsilon \to 0} \limsup_{n \to \infty} P_2(n, \varepsilon) = 0.
$$

By Lemma 2 the probability $\mathbf{P}(a_{i,n-k}Z_{i,n-k}\in A\mid L_{n-k}\geq 0)$ tends, as $n\to\infty$, to \mathbf{P}^* ($\zeta_i^* \in A$) uniformly over $0 \leq k \leq \lfloor (1 - \varepsilon) n \rfloor$. Therefore

$$
\lim_{\varepsilon \to 0} \lim_{n \to \infty} P_1(n, \varepsilon) = \mathbf{P}^* \left(\zeta_i^* \in A \right).
$$

As result, we obtain that

$$
\lim_{n \to \infty} \mathbf{P} \left(a'_{i,n} Z'_{i,n} \in A, \, \tau_n + i \leq n \right) = \mathbf{P}^* \left(\zeta_i^* \in A \right).
$$

This justifies the one-dimensional convergence in (19) for $i \in \mathbb{N}_0$.

Now fix $i \in \mathbb{N}$. Then for $x \ge 0$ and $n \ge i$

$$
\mathbf{P}\left(a'_{-i,n}Z'_{-i,n} \leq x, \tau_n - i \geq 0\right)
$$
\n
$$
= \sum_{k=i}^{n} \mathbf{P}\left(a'_{-i,n}Z'_{-i,n} \leq x, \tau_n = k\right)
$$
\n
$$
= \sum_{k=i}^{n} \mathbf{P}\left(\frac{a_n}{a_{k-i}}Z_{k-i,n} \leq x, S_k < L_{k-1}, S_k \leq L_{k+1,n}\right). \tag{20}
$$

Note that the random sequence $\{(Z_{k-i,n}, a_{k-i,n})\,,\,n\in{\bf N}_{k-i}\}$ is Markovian. Denote by $Z_{k,n}$ (l) the number of particles of nth generation being descendants of l particles of kth generation. Since $Z_{k-i,n} \stackrel{D}{=} Z_{k,n}(l)$ given $Z_{k-i,k} = l$, it follows that

$$
\mathbf{P}\left(\frac{a_n}{a_{k-i}}Z_{k-i,n} \le x, S_k < L_{k-1}, S_k \le L_{k+1,n}\right)
$$
\n
$$
= \mathbf{E}\left(U\left(Z_{k-i,k}, a_{k-i,k}\right); S_k < L_{k-1}\right),
$$

where

$$
U(l, y) = \mathbf{P}\left(a_{k,n}Z_{k,n}(l) \leq \frac{x}{y}, S_k \leq L_{k+1,n}\right).
$$

Clearly,

$$
U(l, y) = \mathbf{P}\left(a_{0,n-k}Z_{0,n-k}(l) \leq \frac{x}{y}, L_{n-k} \geq 0\right).
$$

As result, we obtain that

$$
\mathbf{P}\left(\frac{a_n}{a_{k-i}}Z_{k-i,n} \leq x, S_k < L_{k-1}, S_k \leq L_{k+1,n}\right)
$$
\n
$$
= \mathbf{E}\left(H_{n-k}\left(Z_{k-i,k}, \frac{x}{a_{k-i,k}}\right) \middle| S_k < L_{k-1}\right) \mathbf{P}\left(S_k < L_{k-1}\right) \mathbf{P}\left(L_{n-k} \geq 0\right)
$$
\n
$$
= \mathbf{E}\left(H_{n-k}\left(Z_{k-i,k}, x/a_{k-i,k}\right) \middle| S_k < L_{k-1}\right) \mathbf{P}\left(\tau_n = k\right), \tag{21}
$$

where

$$
H_n(l, x) = \mathbf{P}(a_{0,n}Z_{0,n}(l) \le x | L_n \ge 0)
$$

for $l \in \mathbb{N}_0$ and $x \geq 0$.

Set $Q_{k,l} = (Q_k, \ldots, Q_l)$ for $k, l \in \mathbb{N}$. In the sequel, we will to explicitly include a random environment in the notation. For example, we will write $Z_{k-i,k} \langle Q_{k-i+1,k} \rangle$ instead of $Z_{k-i,k}$. Set

$$
Q_k = \widetilde{Q}_1, \ldots, Q_{k-i+1} = \widetilde{Q}_i, \ldots, Q_1 = \widetilde{Q}_k
$$

and consider a branching process with immigration in the random environment $\widetilde{Q}_1, \ldots, \widetilde{Q}_k$. Then

$$
\mathbf{E}\left(H_{n-k}\left(Z_{k-i,k}\left\langle Q_{k-i+1,k}\right\rangle\right),x/a_{k-i,k}\right)|S_k < L_{k-1}\right)
$$
\n
$$
= \mathbf{E}\left(H_{n-k}\left(Z_{0,i}\left\langle\widetilde{Q}_{i,1}\right\rangle\right),x/\widetilde{a}_{0,i}\right)\middle|\widetilde{M}_k < 0\right),
$$

where the symbols $\widetilde{a}_{0,i}, \widetilde{M}_k, \widetilde{Q}_{i,1}$ have the same meaning for the random environment $\widetilde{Q}_1, \ldots, \widetilde{Q}_k$ as the symbols $a_{0,i}, M_k, Q_{i,1}$ mean for the random environment Q_1, \ldots, Q_k . Further, the random environments $\widetilde{Q}_1, \ldots, \widetilde{Q}_k$ and $Q_1, , \ldots, Q_k$ are identically distributed. Therefore

$$
\mathbf{E}\left(H_{n-k}\left(Z_{0,i}\left\langle\widetilde{Q}_{i,1}\right\rangle,x/\widetilde{a}_{0,i}\right)\middle|\widetilde{M}_{k}<0\right) \n= \mathbf{E}\left(H_{n-k}\left(Z_{0,i}\left\langle Q_{i,1}\right\rangle,x/a_{0,i}\right)\middle|\ M_{k}<0\right).
$$

As result, we obtain that

$$
\mathbf{E} (H_{n-k} (Z_{k-i,k} \langle Q_{k-i+1,k} \rangle, x/a_{k-i,k}) | S_k < L_{k-1})
$$

=
$$
\mathbf{E} (H_{n-k} (Z_{0,i} \langle Q_{i,1} \rangle, x/a_{0,i}) | M_k < 0).
$$
 (22)

Set $\psi_i = Z_{0,i} \langle Q_{i,1} \rangle$. We have from (20)-(22) that

$$
\mathbf{P}\left(a'_{-i,n}Z'_{-i,n} \leq x, \, \tau_n - i \geq 0\right) \n= \sum_{k=i}^{n} \mathbf{E}\left(H_{n-k}\left(\psi_i, x/a_{0,i}\right) | M_k < 0\right) \mathbf{P}\left(\tau_n = k\right).
$$

Therefore, if $\varepsilon \in (0,1)$ and n is large enough, then

 \mathbf{L}

$$
\mathbf{P}\left(a'_{-i,n}Z'_{-i,n} \leq x, \, \tau_n - i \geq 0\right) = P_3\left(n, \varepsilon\right) + P_4\left(n, \varepsilon\right) + P_5\left(n, \varepsilon\right),\tag{23}
$$

where

$$
P_3(n,\varepsilon) = \sum_{k=i}^{\lfloor \varepsilon n \rfloor} \mathbf{E} \left(H_{n-k} \left(\psi_i, x/a_{0,i} \right) | M_k < 0 \right) \mathbf{P} \left(\tau_n = k \right),
$$
\n
$$
P_4(n,\varepsilon) = \sum_{k=\lfloor (1-\varepsilon)n \rfloor+1}^n \mathbf{E} \left(H_{n-k} \left(\psi_i, x/a_{0,i} \right) | M_k < 0 \right) \mathbf{P} \left(\tau_n = k \right),
$$
\n
$$
P_5(n,\varepsilon) = \sum_{k=\lfloor \varepsilon n \rfloor+1}^{\lfloor (1-\varepsilon)n \rfloor} \mathbf{E} \left(H_{n-k} \left(\psi_i, x/a_{0,i} \right) | M_k < 0 \right) \mathbf{P} \left(\tau_n = k \right).
$$

Similar to relation (10) we conclude that

$$
\lim_{\varepsilon \to 0} \limsup_{n \to \infty} P_3(n, \varepsilon) = 0,
$$
\n(24)

$$
\lim_{\varepsilon \to 0} \limsup_{n \to \infty} P_4(n, \varepsilon) = 0.
$$
\n(25)

Let $l\in\mathbf{N}_0$ be fixed. It is not difficult to demonstrate that

$$
\lim_{n \to \infty} a_{0,n} Z_{0,n} (l) =: \zeta_0 (l)
$$
\n(26)

exists a.s. on the probability space $(\Omega, \mathcal{F}_{\infty}, \mathbf{P}^{+})$. By the arguments to those used in Lemma 2 one can show, as $n \to \infty$,

$$
\{a_{0,n}Z_{0,n}(l), i \in \mathbf{N}_0 \mid L_n \ge 0\} \stackrel{D}{\to} \{\zeta_0(l), i \in \mathbf{N}_0\}_{\mathbf{P}^+}.
$$
 (27)

For $x\geq 0$ set

$$
H(l,x) = \mathbf{P}^{+}(\zeta_0(l) \leq x).
$$

It follows from (27) that

$$
\lim_{n \to \infty} H_n(l, x) = H(l, x) \tag{28}
$$

if $x \geq 0$ belongs to the set of continuity points of $H(l, \cdot)$ (with respect to the second argument). By Lemma 2.5 in [\[4\]](#page-32-2)

$$
\{Z_{0,i}\langle Q_{i,1}\rangle,a_{0,i}\,|\,M_n\langle0\rangle\stackrel{D}{\to}\langle Z_{0,i}\langle Q_{i,1}\rangle,a_{0,i}\rangle_{\mathbf{P}}.
$$

as $n \to \infty$. Therefore

$$
\{\psi_i, a_{0,i} \mid M_n < 0\} \stackrel{D}{\to} \left(Z^*_{-i,0}, a^*_{-i,0}\right). \tag{29}
$$

We show that, for fixed $l \in \mathbf{N}_0$ and $K > 0$

$$
\lim_{n \to \infty} \mathbf{E} \left(H_{n-k} \left(l, x/a_{0,i} \right) I_{\{\psi_i = l, x/a_{0,i} \le K\}} \middle| M_k < 0 \right)
$$
\n
$$
= \mathbf{E}^* \left(H \left(l, x/a_{-i,0}^* \right) ; Z_{-i,0}^* = l, x/a_{-i,0}^* \le K \right) \tag{30}
$$

uniformly over $\lfloor \varepsilon n \rfloor < k \leq \lfloor (1 - \varepsilon) n \rfloor$ (here I_A is the indicator of the event A). Let $0 = x_0 < x_1 < \ldots < x_m = K$ for some $m \in \mathbb{N}$. The monotonicity of the function $H(l, \cdot)$ with respect to the second argument gives

$$
\mathbf{E}\left(H_{n-k}\left(l, x/a_{0,i}\right)I_{\{\psi_i=l, x/a_{0,i}\leq K\}}\middle| M_k < 0\right)
$$
\n
$$
= \sum_{j=1}^m \mathbf{E}\left(H_{n-k}\left(l, x/a_{0,i}\right)I_{\{\psi_i=l, x_{j-1}< x/a_{0,i}\leq x_j\}}\middle| M_k < 0\right)
$$
\n
$$
\leq \sum_{j=1}^m H_{n-k}\left(l, x_j\right) \mathbf{P}\left(\psi_i=l, x_{j-1}< x/a_{0,i}\leq x_j\middle| M_k < 0\right). \tag{31}
$$

In view of (28) and (29) the right-hand side of (31) converges, as $n \to \infty$, to

$$
\sum_{j=1}^{m} H(l, x_j) \mathbf{P}^* \left(Z^*_{-i,0} = l, x_{j-1} < x/a^*_{-i,0} \le x_j \right)
$$

uniformly over $\lfloor \varepsilon n \rfloor < k \leq \lfloor (1 - \varepsilon) n \rfloor$, if the selected x_1, \ldots, x_m are simultaneously the continuity points of $H(l, \cdot)$ with respect to the second argument and of \mathbf{P}^* $(Z^*_{-i,0} = l, x/a^*_{-i,0} \leq y)$ with respect to y. Thus, if $\delta > 0$ and n is large enough, the following inequality holds

$$
\mathbf{E}\left(H_{n-k}\left(l,x/a_{0,i}\right)I_{\{\psi_i=l,\,x/a_{0,i}\leq K\}}\,\big|\,M_k<0\right)
$$
\n
$$
\leq \sum_{j=1}^m H\left(l,x_j\right)\mathbf{P}^*\left(Z_{-i,0}^*=l,\,x_{j-1}<\left(a_{-i,0}^*\right)^{-1}x\leq x_j\right)+\delta\qquad(32)
$$

for $\lfloor \varepsilon n \rfloor < k \leq \lfloor (1 - \varepsilon) n \rfloor$. Similarly, if $\delta > 0$ and n is large enough, then

$$
\mathbf{E}\left(H_{n-k}\left(l,x/a_{0,i}\right)I_{\{\psi_i=l,\,x/a_{0,i}\leq K\}}\Big|\ M_k<0\right)
$$
\n
$$
\geq\sum_{j=1}^m H\left(l,x_{j-1}\right)\mathbf{P}^*\left(Z_{-i,0}^*=l,\,x_{j-1}
$$

for $\lfloor \varepsilon n \rfloor < k \leq \lfloor (1 - \varepsilon) n \rfloor$. Since $0 \leq H(l, x) \leq 1$ for $x \geq 0$, the sums in the right-hand sides of (32) and (33) converge, as $\max_{1 \leq j \leq m} (x_j - x_{j-1}) \to 0$, to (see [\[17\]](#page-33-5), Chapter 2, § 6, Section 11)

$$
\mathbf{E}^* \left[H\left(l, x/a_{-i,0}^*\right) ;\, Z_{-i,0}^* = l,\, x/a_{-i,0}^* \leq K \right].
$$

Hence, if $\delta > 0$ and n is large enough, then

$$
\mathbf{E}^* \left(H \left(l, x/a_{-i,0}^* \right); Z_{-i,0}^* = l, x/a_{-i,0}^* \le K \right) - \delta
$$

\n
$$
\le \mathbf{E} \left(H_{n-k} \left(l, x/a_{0,i} \right) I_{\{\psi_i = l, x/a_{0,i} \le K\}} \middle| M_k < 0 \right)
$$

\n
$$
\le \mathbf{E}^* \left(H \left(l, x/a_{-i,0}^* \right); Z_{-i,0}^* = l, x/a_{-i,0}^* \le K \right) + \delta
$$

for $\lfloor \varepsilon n \rfloor < k \leq \lfloor (1 - \varepsilon) n \rfloor$. Since $\delta > 0$ is arbitrary, we obtain the required relation (30).

Now we show that

$$
\lim_{n \to \infty} \mathbf{E} \left(H_{n-k} \left(\psi_i, x/a_{0,i} \right) \mid M_k < 0 \right) = \mathbf{E}^* H \left(Z_{-i,0}^*, x/a_{-i,0}^* \right) \tag{34}
$$

uniformly over $\lfloor \varepsilon n \rfloor < k \leq \lfloor (1-\varepsilon) \, n \rfloor.$ For $N \in \mathbf N$ and $K > 0$ we write

$$
\mathbf{E}\left(H_{n-k}\left(\psi_{i},x/a_{0,i}\right) | M_{k}<0\right)=E_{1}\left(k,n,N,K\right)+E_{2}\left(k,n,N,K\right),\qquad(35)
$$

where

$$
E_1(k, n, N, K) = \mathbf{E} \left(H_{n-k} \left(\psi_i, x/a_{0,i} \right) I_{\{\psi_i \le N, x/a_{0,i} \le K\}} \middle| M_k < 0 \right),
$$

$$
E_2(k, n, N, K) = \mathbf{E} \left(H_{n-k} \left(\psi_i, x/a_{0,i} \right) I_{\{\psi_i > N\} \cup \{x/a_{0,i} > K\}} \middle| M_k < 0 \right).
$$

Since

$$
E_2 (k, n, N, K) \le \mathbf{P} \left(\{ \psi_i > N \} \cup \{ x/a_{0,i} > K \} \mid M_k < 0 \right),
$$

it follows by (29) that

$$
\lim_{K \to \infty} \lim_{N \to \infty} \limsup_{n \to \infty} E_2(k, n, N, K) = 0
$$
\n(36)

uniformly over $\lfloor \varepsilon n \rfloor < k \leq \lfloor (1 - \varepsilon) n \rfloor$. Clearly,

$$
E_1(k, n, N, K) = \sum_{l=0}^{N} \mathbf{E} \left(H_{n-k} (l, x/a_{0,i}) I_{\{\psi_i = l, x/a_{0,i} \leq K\}} \middle| M_k < 0 \right).
$$

Hence, using (30) we conclude that

$$
\lim_{K \to \infty} \lim_{N \to \infty} \lim_{n \to \infty} E_1(k, n, N, K) = \mathbf{E}^* H\left(Z^*_{-i,0}, x/a^*_{-i,0}\right). \tag{37}
$$

uniformly over $\lfloor \varepsilon n \rfloor < k \leq \lfloor (1 - \varepsilon) n \rfloor$. Combining (35)-(37) we obtain the desired relation (34).

It follows from (34) that

$$
\lim_{\varepsilon \to 0} \lim_{n \to \infty} P_5(n, \varepsilon) = \mathbf{E}^* H\left(Z^*_{-i,0}, x/a^*_{-i,0}\right). \tag{38}
$$

Now (23)-(25) and (38) imply

$$
\lim_{n \to \infty} \mathbf{P} \left(a'_{-i,n} Z'_{-i,n} \leq x, \, \tau_n - i \geq 0 \right) = \mathbf{E}^* H \left(Z^*_{-i,0}, x/a^*_{-i,0} \right).
$$

Hence,

$$
\lim_{n \to \infty} \mathbf{P} \left(a'_{-i,n} Z'_{-i,n} \le x \right) = \mathbf{E}^* H \left(Z^*_{-i,0}, x/a^*_{-i,0} \right). \tag{39}
$$

We now analyze a branching process with immigration in the random environment $\{Q_k^*, k \in \mathbf{Z}\}$. The random sequence $\{(Z_{-i,n}^*, a_{-i,n}^*)$, $n \in \mathbf{N}_{-i}\}$ is Markovian. Denote by $Z_{k,n}^*(l)$ the number of particles in nth generation which are descendants of l particles of the kth generation. Note that

$$
\left\{ \left(Z_{0,n}^{*}\left(l\right) ,a_{0,n}^{*}\right) ,\, n\in\mathbf{N}_{0}\right\} \stackrel{D}{=} \left\{ \left(Z_{0,n}\left(l\right) ,a_{0,n}\right) ,\, n\in\mathbf{N}_{0}\right\} _{\mathbf{P}^{+}}.\tag{40}
$$

Since $Z^*_{-i,n} \stackrel{D}{=} Z^*_{0,n}(l)$ given $Z^*_{-i,0} = l$, it follows that, for any bounded and continuous function $f: \mathbf{R} \to \mathbf{R}$ and $n \in \mathbf{N}_{-i}$,

$$
\mathbf{E}^* f\left(a_{-i,n}^* Z_{-i,n}^*\right) = \mathbf{E}^* V_n\left(Z_{-i,0}^*, 1/a_{-i,0}^*\right),\tag{41}
$$

where

$$
V_n(l, y) = \mathbf{E}^* f\left(a_{0,n}^* Z_{0,n}^*(l) / y\right).
$$

By (40)

$$
V_n(l, y) = \mathbf{E}^+ f(a_{0,n} Z_{0,n}(l) / y).
$$
 (42)

In view of (5), as $n \to \infty$,

$$
a_{-i,n}^* Z_{-i,n}^* \stackrel{D}{\to} \zeta_{-i}^*,\tag{43}
$$

and in view of (26)

$$
(a_{0,n}Z_{0,n}(l))_{\mathbf{P}^+}\stackrel{D}{\to}(\zeta_0(l))_{\mathbf{P}^+}.
$$
\n(44)

Using (42), (44) and applying the dominated convergence theorem we see that

$$
\lim_{n \to \infty} V_n(l, y) = V(l, y), \qquad (45)
$$

where

$$
V(l, y) = \mathbf{E}^{+} f(\zeta_0(l)/y).
$$

Applying the dominated convergence theorem again we obtain from (41), (43) and (45) that

$$
\mathbf{E}^* f\left(\zeta_{-i}^*\right) = \mathbf{E}^* V\left(Z_{-i,0}^*, 1/a_{-i,0}^*\right). \tag{46}
$$

Fix $x \geq 0$. As relation (46) is valid for any bounded and continuous function f, it is valid, even when a function f is the indicator of the semi-axis $(-\infty, x]$. It means that

$$
\mathbf{P}^* \left(\zeta_{-i}^* \le x \right) = \mathbf{E}^* H \left(Z_{-i,0}^*, x/a_{-i,0}^* \right) \tag{47}
$$

(we take into account that $V(l, y) = H(l, xy)$ for the specified function f).

Equalities (39) and (47) imply the one-dimensional convergence in relation (19) for $i \in \mathbf{Z} \setminus \mathbf{N}_0$.

The lemma is proved.

Remark 3. It is not difficult to verify that (19) admits the following generalization: for any $a \leq 0$ and $b > 0$, as $n \to \infty$,

$$
\left\{a'_{i,n}Z'_{i,n},\, i\in\mathbf{Z}\,\middle|\, \frac{L_n}{C_n}\leq a,\, \frac{S_n-L_n}{C_n}\leq b\right\}\stackrel{D}{\to}\left\{\zeta_i^*,\, i\in\mathbf{Z}\right\}.
$$

Lemma 4. *If the conditions of Theorem 1 are satisfied*, *then* P[∗] -*a.s.*

$$
\Sigma_1<+\infty, \qquad \Sigma_2<+\infty.
$$

Proof. It is shown in Lemma 2.7 from [\[4\]](#page-32-2) that, if the conditions of Theorem 1 are satisfied, then the series $\sum_{i=0}^{\infty} \mu_{i+1}e^{-S_i}$ converges \mathbf{P}^+ -a.s. Hence, the series $\sum_{i=0}^{\infty} \mu_{i+1}^+ e^{-S_i^+}$ converges **P**^{*}-a.s. Similarly we can prove that the series $\sum_{i=1}^{\infty} \mu_i^- e^{S_i^-}$ converges \mathbf{P}^* -a.s. As result, we obtain that the series $\sum_{i\in\mathbf{Z}} \mu_{i+1}^* e^{-S_i^*}$ converges \mathbf{P}^* -a.s. Thus, $\Sigma_1 < +\infty$ \mathbf{P}^* -a.s.

Fix $i \in \mathbb{Z}$. If the random environment \mathcal{E}^* is fixed, the random sequence $\{\eta_i^*; a_{i,j}^* Z_{i,j}^*, j \in \mathbf{N}_{i+1}\}\$ is a martingale. Therefore

$$
\mathbf{E}^* \left(a_{i,j}^* Z_{i,j}^* \mid \mathcal{E}^* \right) = \mu_{i+1}^* \tag{48}
$$

for $j \in \mathbf{N}_{i+1}$. By (5), (48) using Fatou's lemma we obtain that

$$
\mathbf{E}^* \left(\zeta_i^* \mid \mathcal{E}^* \right) \le \liminf_{j \to \infty} \mathbf{E}^* \left(a_{i,j}^* Z_{i,j}^* \mid \mathcal{E}^* \right) = \mu_{i+1}^*
$$

and, consequently,

$$
\mathbf{E}^* \left(\zeta_i^* e^{-S_i^*} \middle| \mathcal{E}^* \right) = e^{-S_i^*} \mathbf{E}^* \left(\zeta_i^* \middle| \mathcal{E}^* \right) \le \mu_{i+1}^* e^{-S_i^*}. \tag{49}
$$

We have proved that the series $\sum_{i\in\mathbf{Z}}\mu_{i+1}^*e^{-S_i^*}$ converges \mathbf{P}^* -a.s. This fact combined with (49) implies convergence of the series $\sum_{i\in \mathbf{Z}} \mathbf{E}^* \left(\zeta_{i+1}^* e^{-S_i^*} \mid \mathcal{E}^* \right)$ **P**^{*}-a.s. Since the random variables $\zeta_{i+1}^*e^{-S_i^*}$ are nonnegative, it follows that the series $\sum_{i\in \mathbf{Z}} \zeta_i^* e^{-S_i^*}$ converges a.s. for any fixed environment \mathcal{E}^* . Hence, Σ_2 < + ∞ **P**^{*}-a.s.

The lemma is proved.

Set

$$
\begin{split} \Sigma_1^{(1)} &= \sum_{i=0}^\infty \mu^+_{i+1} e^{-S^+_i} = \sum_{i \in \mathbf{N}_0} \mu^*_{i+1} e^{-S^*_i}, \\ \Sigma_1^{(2)} &= \sum_{i=1}^\infty \mu^-_i e^{S^-_i} = \sum_{i \in \mathbf{Z} \backslash \mathbf{N}_0} \mu^*_{i+1} e^{-S^*_i}. \end{split}
$$

Clearly,

$$
\Sigma_1 = \Sigma_1^{(1)} + \Sigma_1^{(2)} \tag{50}
$$

and by virtue of Lemma $4P^*$ -a.s.

$$
\Sigma_1^{(1)} < +\infty, \qquad \Sigma_1^{(2)} < +\infty. \tag{51}
$$

Lemma 5. *If the conditions of Theorem 1 are satisfied*, *then* P[∗] -*a.s.*, *as* $n \to \infty$, $\left(\sum_{n=1}^{n-1} \right)$

$$
\left\{ \sum_{i=0}^{n-1} \mu_{i+1} e^{-S_i} \middle| L_n \ge 0 \right\} \xrightarrow{D} \Sigma_1^{(1)}, \tag{52}
$$

$$
\left\{ \sum_{i=1}^{n-1} \mu_i e^{S_i} \middle| M_n < 0 \right\} \xrightarrow{D} \Sigma_1^{(2)}.
$$
\n
$$
\tag{53}
$$

Proof. Let $f: \mathbf{R} \to \mathbf{R}$ be a bounded and continuous function. By virtue of (3) for fixed $k \in \mathbf{N}$

$$
\left\{ f\left(\sum_{i=0}^k \mu_{i+1} e^{-S_i}\right) \middle| L_n \ge 0 \right\} \stackrel{D}{\to} f\left(\sum_{i=0}^k \mu_{i+1}^+ e^{-S_i^+}\right)
$$

as $n \to \infty$. Recalling (51) we conclude that

$$
\lim_{k \to \infty} f\left(\sum_{i=0}^k \mu_{i+1}^+ e^{-S_i^+}\right) = f\left(\Sigma_1^{(1)}\right)
$$

P[∗]-a.s. From these two facts, in view of Lemma 2.5 of [\[4\]](#page-32-2), it follows that

$$
\left\{ f\left(\sum_{i=0}^{n-1} \mu_{i+1} e^{-S_i}\right) \middle| L_n \ge 0 \right\} \xrightarrow{D} f\left(\Sigma_1^{(1)}\right).
$$

Thus, relation (52) is true. Relation (53) can be proved by similar arguments.

The lemma is proved.

Remark 4. It is not difficult to verify that if we combine the left-hand sides of relations (3) and (52) (or (4) and (53)), then the respective statements concerning convergence in distribution of the four dimensional tuples of the random elements given $L_n \geq 0$ (or $M_n < 0$) are still force.

Set for $n\in\mathbf{N}$

$$
\mu'_{i,n} = \left\{ \begin{array}{ll} \mu_{\tau_n+i}, & i \in \mathbf{N}_{(-\tau_n)}, \\[1ex] 0, & i \in \mathbf{Z} \setminus \mathbf{N}_{(-\tau_n)}. \end{array} \right.
$$

Let

$$
\Sigma_1^{(1)}(n) = \sum_{j=0}^{n-1-\tau_n} \mu'_{j+1,n} e^{-S'_{j,n}}, \qquad \Sigma_1^{(2)}(n) = \sum_{j=1}^{\tau_n} \mu'_{-j+1,n} e^{-S'_{j,n}}.
$$

Lemma 6. *If the conditions of Theorem 1 are satisfied*, *then* P[∗] -*a.s.*, *as* $n \to \infty$,

$$
\left(\left\{\left(\mu'_{i,n}, S'_{i,n}\right), i \in \mathbf{N}_0\right\}, \, \Sigma_1^{(1)}\left(n\right)\right) \stackrel{D}{\to} \left(\left\{\left(\mu_i^*, S_i^*\right), i \in \mathbf{N}_0\right\}, \, \Sigma_1^{(1)}\right),\tag{54}
$$

$$
\left(\left\{\left(\mu'_{-i,n}, S'_{-i,n}\right), i \in \mathbf{N}\right\}, \, \Sigma_1^{(2)}\left(n\right)\right) \stackrel{D}{\rightarrow} \left(\left\{\left(\mu^*_{-i}, S^*_{-i}\right), i \in \mathbf{N}\right\}, \, \Sigma_1^{(2)}\right). \tag{55}
$$

Moreover, the left-hand sides of these relations are asymptotically independent.

Proof. We prove for simplicity only convergence in distribution (for a fixed *i*) of the random sequences $(\mu'_{i,n}, S'_{i,n}, \Sigma_1^{(1)}(n))$ and $(\mu'_{-i,n}, S'_{-i,n}, \Sigma_1^{(2)}(n))$, as $n \to \infty$.

Fix $i \in \mathbb{N}_0$. Similarly to relation (7), we can show that, for any bounded and continuous function $f: \mathbb{R}^3 \to \mathbb{R}$,

$$
\mathbf{E}\left[f\left(\mu'_{i,n}, S'_{i,n}, \sum_{j=0}^{n-1-\tau_n} \mu'_{j+1,n} e^{-S'_{j,n}}\right); \tau_n + i \leq n\right]
$$
\n
$$
= \sum_{k=0}^{n-i} \mathbf{E}\left(f\left(\mu_i, S_i, \sum_{j=0}^{n-1-k} \mu_{j+1} e^{-S_j}\right) \middle| L_{n-k} \geq 0\right) \mathbf{P}\left(\tau_n = k\right) (56)
$$

for $n \geq i$. Repeating the arguments of Lemma 1 and using Lemma 5 and Remark 4, we can deduce from (56) that

$$
\lim_{n \to \infty} \mathbf{E} f\left(\mu'_{i,n}, S'_{i,n}, \sum_{j=0}^{n-1-\tau_n} \mu'_{j+1,n} e^{-S'_{j,n}}\right)
$$
\n
$$
= \mathbf{E}^* f\left(\mu_i^+, S_i^+, \sum_{j \in \mathbf{N}_0} \mu_{j+1}^+ e^{-S_j^+}\right) = \mathbf{E}^* f\left(\mu_i^*, S_i^*, \Sigma_1^{(1)}\right).
$$

Thus, relation (54) is proved.

Now fix $i \in \mathbb{N}$. It is easy to show (see the proof of relation (13)) that for $n\geq i$

$$
\mathbf{E}\left[f\left(\mu'_{-i,n}, S'_{-i,n}, \sum_{j=1}^{\tau_n} \mu'_{-j+1,n} e^{-S'_{-j,n}}\right); \tau_n - i \ge 0\right]
$$

$$
= \sum_{k=i}^n \mathbf{E}\left(f\left(\mu_{i+1}, -S_i, \sum_{j=1}^k \mu_j e^{S_j}\right) \middle| M_k < 0\right) \mathbf{P}(\tau_n = k)
$$

and therefore (see Lemma 5 and Remark 4)

$$
\lim_{n \to \infty} \mathbf{E} f\left(\mu'_{-i,n}, S'_{-i,n}, \sum_{j=1}^{\tau_n} \mu'_{-j+1,n} e^{-S'_{-j,n}}\right)
$$
\n
$$
= \mathbf{E}^* f\left(\mu_{i+1}^-, -S_i^-, \sum_{j=1}^\infty \mu_j^- e^{S_j^-}\right) = \mathbf{E}^* f\left(\mu_{-i}^*, S_{-i}^*, \Sigma_1^{(2)}\right)
$$

.

,

This proves (55). The asymptotic independence of the left-hand sides of relations (54) and (55) is obvious.

The lemma is proved.

Remark 5. It is not difficult to verify that statement (54) admits the following generalization: for any $a \leq 0$ and $b > 0$, as $n \to \infty$,

$$
\left(\left\{ \left(\mu'_{i,n}, S'_{i,n} \right), i \in \mathbb{N}_0 \right\}, \, \Sigma_1^{(1)}(n) \, \middle| \, \frac{L_n}{C_n} \le a, \, \frac{S_n - L_n}{C_n} \le b \right)
$$

$$
\xrightarrow{D} \left(\left\{ \left(\mu_i^*, S_i^* \right), i \in \mathbb{N}_0 \right\}, \, \Sigma_1^{(1)} \right).
$$

Statement (55) allows for a similar generalization.

Lemma 7. *If the conditions of Theorem 1 are satisfied, then, as* $n \rightarrow \infty$,

$$
\left\{\frac{b_n - b_{\tau_n + i}}{b_n}, i \in \mathbb{N}_0\right\} \xrightarrow{D} \left\{\frac{\sum_{j=i}^{\infty} \mu_{j+1}^+ \exp\left(-S_j^+\right)}{\Sigma_1}, i \in \mathbb{N}_0\right\}
$$
\n
$$
\left\{\frac{b_{\tau_n - i}}{b_n}, i \in \mathbb{N}\right\} \xrightarrow{D} \left\{\frac{\sum_{j=i+1}^{\infty} \mu_j^- \exp\left(S_j^-\right)}{\Sigma_1}, i \in \mathbb{N}\right\},\
$$
\n
$$
\left\{\frac{a_{\tau_n + i}}{b_n}, i \in \mathbb{N}_0\right\} \xrightarrow{D} \left\{\frac{\exp\left(-S_i^+\right)}{\Sigma_1}, i \in \mathbb{N}_0\right\},\
$$
\n
$$
\left\{\frac{a_{\tau_n - i}}{b_n}, i \in \mathbb{N}\right\} \xrightarrow{D} \left\{\frac{\exp\left(S_i^-\right)}{\Sigma_1}, i \in \mathbb{N}\right\}.
$$

Proof. To simplify the presentation we check the first statement only. Moreover, we prove only convergence of one-dimensional distributions. Fix $i \in \mathbb{N}_0$. Note that for $\tau_n + i \leq n$

$$
\frac{b_{\tau_n+i}}{b_n} = \frac{\sum_{j=0}^{\tau_n+i-1} \mu_{j+1} \exp(-S_j)}{\sum_{j=0}^{n-1} \mu_{j+1} \exp(-S_j)} = \frac{\sum_{j=0}^{\tau_n+i-1} \mu_{j+1} \exp(-(S_j - S_{\tau_n}))}{\sum_{j=0}^{n-1} \mu_{j+1} \exp(-(S_j - S_{\tau_n}))}
$$

$$
= \frac{\sum_{j=0}^{i-1} \mu'_{j+1,n} \exp\left(-S'_{j,n}\right) + \Sigma_1^{(2)}(n)}{\Sigma_1^{(1)}(n) + \Sigma_1^{(2)}(n)}.
$$

Since the last expression is a bounded continuous function of the random element mentioned in Lemma 6, it follows that

$$
\frac{b_{\tau_n+i}}{b_n} \stackrel{D}{\to} \frac{\sum_{j=0}^{i-1} \mu_{j+1}^+ \exp\left(-S_j^+\right) + \Sigma_1^{(2)}}{\Sigma_1^{(1)} + \Sigma_1^{(2)}}
$$

as $n \to \infty$. Whence, taking into account (50) we obtain the required relation. The remaining three statements may be proved by similar arguments.

The lemma is proved.

Remark 6. We can construct a new random element by combining the left-hand sides of all the relations included in Lemmas 3 and 7. It is not difficult to prove convergence in distribution of the sequence of these random elements to a random element constructed by the right-hand sides of the corresponding relations of Lemmas 3 and 7. Moreover, a random element constructed by the left-hand sides is asymptotically independent, as $n \to \infty$, of the random event

$$
\left\{C_n^{-1}L_n \le a, \, C_n^{-1} \left(S_n - L_n\right) \le b\right\}
$$

for any $a \leq 0$ and $b > 0$.

3. Proof of the main result

First part. We establish convergence of one-dimensional distributions: if $t > 0$, then, as $n \to \infty$,

$$
\frac{a_{\lfloor nt \rfloor}}{b_{\lfloor nt \rfloor}} Z_{\lfloor nt \rfloor} \stackrel{D}{\to} \frac{\Sigma_2}{\Sigma_1}.
$$
\n
$$
(57)
$$

Set for $r \in \mathbf{N}$

$$
U_r^{(i)} = \sum_{j=\tau_r-i}^{\tau_r+i-1} Z_{j,r},
$$

$$
V_r^{(i)} = \sum_{j=0}^{\tau_r-i-1} Z_{j,r} + \sum_{j=\tau_r+i}^{r-1} Z_{j,r}.
$$

It is clear that for $i \in {\bf N}$

$$
Z_{\lfloor nt \rfloor} = \sum_{j=0}^{\lfloor nt \rfloor - 1} Z_{j, \lfloor nt \rfloor} = U_{\lfloor nt \rfloor}^{(i)} + V_{\lfloor nt \rfloor}^{(i)}.
$$
 (58)

Note that

$$
\mathbf{E}\left(a_{j,\lfloor nt\rfloor}Z_{j,\lfloor nt\rfloor} \;|\; Q_{1,\lfloor nt\rfloor}\right) = \mu_{j+1},\tag{59}
$$

if $1 \leq j < \lfloor nt \rfloor$. Observing that $a_{\lfloor nt \rfloor} = a_j a_{j, \lfloor nt \rfloor}$ for $1 \leq j < \lfloor nt \rfloor$ we obtain by (59) that

$$
\mathbf{E} \left(\frac{a_{\lfloor nt \rfloor}}{b_{\lfloor nt \rfloor}} V_{\lfloor nt \rfloor}^{(i)} \right)
$$
\n
$$
= \mathbf{E} b_{\lfloor nt \rfloor}^{-1} \left(\sum_{j=0}^{\tau_{\lfloor nt \rfloor - i - 1}} a_j a_{j, \lfloor nt \rfloor} Z_{j, \lfloor nt \rfloor} + \sum_{j=\tau_{\lfloor nt \rfloor + i}}^{\lfloor nt \rfloor - 1} a_j a_{j, \lfloor nt \rfloor} Z_{j, \lfloor nt \rfloor} \right)
$$
\n
$$
= \mathbf{E} b_{\lfloor nt \rfloor}^{-1} \left(\sum_{j=0}^{\tau_{\lfloor nt \rfloor - i - 1}} \mu_{j+1} a_j + \sum_{j=\tau_{\lfloor nt \rfloor + i}}^{\lfloor nt \rfloor - 1} \mu_{j+1} a_j \right)
$$
\n
$$
= \mathbf{E} \frac{b_{\tau_{\lfloor nt \rfloor - i}} + (b_{\lfloor nt \rfloor - b_{\tau_{\lfloor nt \rfloor + i}})})}{b_{\lfloor nt \rfloor}}.
$$
\n(60)

Applying Lemma 7 to the right-hand side of (60), we conclude that

$$
\lim_{n \to \infty} \mathbf{E}\left(\frac{a_{\lfloor nt \rfloor}}{b_{\lfloor nt \rfloor}} V_{\lfloor nt \rfloor}^{(i)}\right) = \frac{\sum_{j=i}^{\infty} \mu_{j+1}^{+} \exp\left(-S_{j}^{+}\right) + \sum_{j=i+1}^{\infty} \mu_{j}^{-} \exp\left(S_{j}^{-}\right)}{\Sigma_{1}}
$$

and, therefore (see Lemma 4),

$$
\lim_{i \to \infty} \lim_{n \to \infty} \mathbf{E}\left(\frac{a_{\lfloor nt \rfloor}}{b_{\lfloor nt \rfloor}} V_{\lfloor nt \rfloor}^{(i)}\right) = 0. \tag{61}
$$

By Markov inequality for any $\varepsilon>0$

$$
\mathbf{P}\left(\frac{a_{\lfloor nt \rfloor}}{b_{\lfloor nt \rfloor}}V_{\lfloor nt \rfloor}^{(i)} \geq \varepsilon\right) \leq \varepsilon^{-1}\mathbf{E}\left(\frac{a_{\lfloor nt \rfloor}}{b_{\lfloor nt \rfloor}}V_{\lfloor nt \rfloor}^{(i)}\right).
$$

Hence, taking into account (61) we obtain that

$$
\lim_{i \to \infty} \lim_{n \to \infty} \mathbf{P}\left(\frac{a_{\lfloor nt \rfloor}}{b_{\lfloor nt \rfloor}} V_{\lfloor nt \rfloor}^{(i)} \ge \varepsilon\right) = 0. \tag{62}
$$

Observe that we may assume in the sequel that $i\leq \tau_{\lfloor nt\rfloor}<\lfloor nt\rfloor-i$ (see the proof of Lemma 1). Note that

$$
U_{\lfloor nt \rfloor}^{(i)} = \sum_{j=-i}^{i-1} Z_{\tau_{\lfloor nt \rfloor} + j, \lfloor nt \rfloor} = \sum_{j=-i}^{i-1} Z'_{j, \lfloor nt \rfloor}
$$

and, therefore,

$$
\frac{a_{\lfloor nt \rfloor}}{b_{\lfloor nt \rfloor}} U_{\lfloor nt \rfloor}^{(i)} = \sum_{j=-i}^{i-1} \frac{a_{\tau_{\lfloor nt \rfloor} + j}}{b_{\lfloor nt \rfloor}} a'_{j, \lfloor nt \rfloor} Z'_{j, \lfloor nt \rfloor}.
$$
\n(63)

Applying Lemmas 3, 7 and Remark 6 to relation (63), we obtain that, as $n \to \infty$,

$$
\frac{a_{\lfloor nt \rfloor}}{b_{\lfloor nt \rfloor}} U_{\lfloor nt \rfloor}^{(i)} \stackrel{D}{\to} \frac{1}{\sum_{1}} \sum_{j=-i}^{i-1} \zeta_j^* e^{-S_j^*}.
$$
 (64)

Hence, for all but a countable set of $x\geq 0$

$$
\lim_{n \to \infty} \mathbf{P}\left(\frac{a_{\lfloor nt \rfloor}}{b_{\lfloor nt \rfloor}} U_{\lfloor nt \rfloor}^{(i)} \le x\right) = \mathbf{P}\left(\frac{1}{\Sigma_1} \sum_{j=-i}^{i-1} \zeta_j^* e^{-S_j^*} \le x\right). \tag{65}
$$

In view of Lemma 4

$$
\lim_{i \to \infty} \mathbf{P}\left(\frac{1}{\Sigma_1} \sum_{j=-i}^{i-1} \zeta_i^* e^{-S_i^*} \le x\right) = \mathbf{P}\left(\frac{\Sigma_2}{\Sigma_1} \le x\right). \tag{66}
$$

We obtain by (65) and (66) that

$$
\lim_{i \to \infty} \lim_{n \to \infty} \mathbf{P}\left(\frac{a_{\lfloor nt \rfloor}}{b_{\lfloor nt \rfloor}} U_{\lfloor nt \rfloor}^{(i)} \le x\right) = \mathbf{P}\left(\frac{\Sigma_2}{\Sigma_1} \le x\right). \tag{67}
$$

It follows from (58), (62) and (67) that for all but a countable set of $x \ge 0$

$$
\lim_{n \to \infty} \mathbf{P}\left(\frac{a_{\lfloor nt \rfloor}}{b_{\lfloor nt \rfloor}} Z_{\lfloor nt \rfloor} \leq x\right) = \mathbf{P}\left(\frac{\Sigma_2}{\Sigma_1} \leq x\right).
$$

This proves (57).

Remark 7. It is not difficult to verify that relation (64) admits the following generalization: for any $a\leq 0$ and $b>0,$ as $n\to\infty,$

$$
\left\{\frac{a_{\lfloor nt \rfloor}}{b_{\lfloor nt \rfloor}}U_{\lfloor nt \rfloor}^{(i)} \mid \frac{L_{\lfloor nt \rfloor}}{C_n} \le a, \frac{S_{\lfloor nt \rfloor} - L_{\lfloor nt \rfloor}}{C_n} \le b \right\} \xrightarrow{D} \frac{1}{\sum_{1}} \sum_{j=-i}^{i+1} \zeta_j^* e^{-S_j^*}.
$$

Second part. Now we establish convergence of two-dimensional distributions. Select $0 < t_1 < t_2,$ fix an $\varepsilon > 0$ and introduce the following random events:

$$
A_{n,\varepsilon} = \left\{ L_{\lfloor nt_1 \rfloor} > L_{\lfloor nt_1 \rfloor, \lfloor nt_2 \rfloor} + \varepsilon C_n \right\},
$$

\n
$$
B_{n,\varepsilon} = \left\{ L_{\lfloor nt_1 \rfloor} < L_{\lfloor nt_1 \rfloor, \lfloor nt_2 \rfloor} - \varepsilon C_n \right\},
$$

\n
$$
D_{n,\varepsilon} = \left\{ \left| L_{\lfloor nt_1 \rfloor} - L_{\lfloor nt_1 \rfloor, \lfloor nt_2 \rfloor} \right| \leq \varepsilon C_n \right\}.
$$

We show that, as $n \to \infty$,

$$
\left\{ \frac{a_{\lfloor nt_1 \rfloor}}{b_{\lfloor nt_1 \rfloor}} Z_{\lfloor nt_1 \rfloor}, \frac{a_{\lfloor nt_2 \rfloor}}{b_{\lfloor nt_2 \rfloor}} Z_{\lfloor nt_2 \rfloor} \middle| A_{n,\varepsilon} \right\} \xrightarrow{D} (\gamma_1, \gamma_2), \tag{68}
$$

$$
\left\{ \frac{a_{\lfloor nt_1 \rfloor}}{b_{\lfloor nt_1 \rfloor}} Z_{\lfloor nt_1 \rfloor}, \frac{a_{\lfloor nt_2 \rfloor}}{b_{\lfloor nt_2 \rfloor}} Z_{\lfloor nt_2 \rfloor} \middle| B_{n,\varepsilon} \right\} \xrightarrow{D} (\gamma_1, \gamma_1), \tag{69}
$$

where γ_1, γ_2 are independent random variables and $\gamma_1 \stackrel{D}{=} \gamma_2 \stackrel{D}{=} \Sigma_2/\Sigma_1$.

First we establish (68). To this aim we prove that, for any fixed $i \in \mathbb{N}$ and for all but a countable set of (x_1, x_2) with $x_1, x_2 \geq 0$,

$$
\lim_{n \to \infty} \mathbf{P} \left(\frac{a_{\lfloor nt_1 \rfloor}}{b_{\lfloor nt_1 \rfloor}} U_{\lfloor nt_1 \rfloor}^{(i)} \le x_1, \frac{b_{\lfloor nt_2 \rfloor}}{a_{\lfloor nt_2 \rfloor}} U_{\lfloor nt_2 \rfloor}^{(i)} \le x_2 \middle| A_{n,\varepsilon} \right)
$$
\n
$$
= \mathbf{P} \left(\frac{1}{\Sigma_1} \sum_{j=-i}^{i+1} \zeta_j^* e^{-S_j^*} \le x_1 \right) \mathbf{P} \left(\frac{1}{\Sigma_1} \sum_{j=-i}^{i+1} \zeta_j^* e^{-S_j^*} \le x_2 \right). \tag{70}
$$

Provided the random event $A_{n,\varepsilon}$ occurred, it follows that, as $n \to \infty$,

$$
\frac{b_{\lfloor nt_2 \rfloor}}{a_{\lfloor nt_2 \rfloor}} \sim \frac{b_{\lfloor nt_2 \rfloor} - b_{\lfloor nt_1 \rfloor}}{a_{\lfloor nt_2 \rfloor}} = \frac{\widetilde{b}_{\lfloor nt_2 \rfloor - \lfloor nt_1 \rfloor}}{\widetilde{a}_{\lfloor nt_2 \rfloor - \lfloor nt_1 \rfloor}},
$$

where the values $\tilde{a}_{\lfloor nt_2\rfloor-\lfloor nt_1\rfloor}$ and $\tilde{b}_{\lfloor nt_2\rfloor-\lfloor nt_1\rfloor}$ are constructed by the random environment $Q_i := Q_{\lfloor nt_1 \rfloor + i}, i = 1, \ldots, \lfloor nt_2 \rfloor - \lfloor nt_1 \rfloor$, just as the values $a_{\lfloor nt_2 \rfloor - \lfloor nt_1 \rfloor}$ and $b_{\lfloor nt_2\rfloor - \lfloor nt_1\rfloor}$ are constructed by the random environment $Q_{1,\lfloor nt_2\rfloor - \lfloor nt_1\rfloor}$.

Further, given $A_{n,\varepsilon}$, the inequality $\tau_{\lfloor nt_2\rfloor} > \tau_{\lfloor nt_1\rfloor}$ is true (we may assume that $\lfloor nt_2 \rfloor - i > \tau_{\lfloor nt_2 \rfloor} > \tau_{\lfloor nt_1 \rfloor} + i$. Thus, if the random environment $\{Q_n, n \in \mathbf{N}\}\$ is fixed, the distribution of the random variable $U_{\lfloor nt \rfloor}^{(i)}$ $\begin{bmatrix} u \\ nt_1 \end{bmatrix}$ is completely determined by the random environment $Q_{1, \lfloor nt_1 \rfloor}$ and the distribution of the random variable $U_{\parallel n i}^{(i)}$ $\binom{v}{n}$ is completely determined by the random environment $Q_{\lfloor nt_1\rfloor+1,\lfloor nt_2\rfloor}$. Moreover, $U_{\lfloor nt_2\rfloor}^{(i)} = \widetilde{U}_{\lfloor nt_2\rfloor-\lfloor nt_1\rfloor}^{(i)}$, where $\widetilde{U}_{\lfloor nt_2\rfloor-\lfloor nt_1\rfloor}^{(i)}$ has the same meaning for the environment Q_i , $i = 1, ..., \lfloor nt_2 \rfloor - \lfloor nt_1 \rfloor$, as $U^{(i)}_{\perp m}$ $\binom{v}{n}$ _{[nt₂]− $\lfloor nt_1 \rfloor$ has for the environment $Q_{1,\lfloor nt_2 \rfloor - \lfloor nt_1 \rfloor}$.}

Summarizing the arguments above, we see that to prove (70) it is sufficient to show that

$$
\lim_{n\to\infty} \mathbf{P}\left(\frac{a_{\lfloor nt_1 \rfloor}}{b_{\lfloor nt_1 \rfloor}}U_{\lfloor nt_1 \rfloor}^{(i)} \leq x_1, \frac{\widetilde{a}_{\lfloor nt_2 \rfloor - \lfloor nt_1 \rfloor}}{\widetilde{b}_{\lfloor nt_2 \rfloor - \lfloor nt_1 \rfloor}}\widetilde{U}_{\lfloor nt_2 \rfloor - \lfloor nt_1 \rfloor}^{(i)} \leq x_2 \middle| A_{n,\varepsilon} \right)
$$

$$
= \mathbf{P}\left(\frac{1}{\Sigma_{1}}\sum_{j=-i}^{i+1} \zeta_{j}^{*} e^{-S_{j}^{*}} \leq x_{1}\right) \mathbf{P}\left(\frac{1}{\Sigma_{1}}\sum_{j=-i}^{i+1} \zeta_{j}^{*} e^{-S_{j}^{*}} \leq x_{2}\right).
$$
 (71)

Note that

$$
\mathbf{P}\left(\frac{a_{\lfloor nt_1 \rfloor}}{b_{\lfloor nt_1 \rfloor}}U_{\lfloor nt_1 \rfloor}^{(i)} \leq x_1, \frac{\widetilde{a}_{\lfloor nt_2 \rfloor - \lfloor nt_1 \rfloor}}{\widetilde{b}_{\lfloor nt_2 \rfloor - \lfloor nt_1 \rfloor}}\widetilde{U}_{\lfloor nt_2 \rfloor - \lfloor nt_1 \rfloor}^{(i)} \leq x_2, A_{n,\varepsilon}\right)
$$
\n
$$
= \int_{-\infty}^{0} \int_{0}^{+\infty} \mathbf{P}\left(\frac{a_{\lfloor nt_1 \rfloor}}{b_{\lfloor nt_1 \rfloor}}U_{\lfloor nt_1 \rfloor}^{(i)} \leq x_1, \frac{L_{\lfloor nt_1 \rfloor}}{C_n} \in da, \frac{S_{\lfloor nt_1 \rfloor} - L_{\lfloor nt_1 \rfloor}}{C_n} \in db\right)
$$
\n
$$
\times \mathbf{P}\left(\frac{a_{\lfloor nt_2 \rfloor - \lfloor nt_1 \rfloor}}{b_{\lfloor nt_2 \rfloor - \lfloor nt_1 \rfloor}}U_{\lfloor nt_2 \rfloor - \lfloor nt_1 \rfloor}^{(i)} \leq x_2, \frac{L_{\lfloor nt_2 \rfloor - \lfloor nt_1 \rfloor}}{C_n} < b - a - \varepsilon\right).
$$

Hence, taking into account Remark 7 we deduce that, as $n \to \infty$,

$$
\mathbf{P}\left(\frac{a_{\lfloor nt_1 \rfloor}}{b_{\lfloor nt_1 \rfloor}}U_{\lfloor nt_1 \rfloor}^{(i)} \leq x_1, \frac{\widetilde{a}_{\lfloor nt_2 \rfloor - \lfloor nt_1 \rfloor}}{\widetilde{b}_{\lfloor nt_2 \rfloor - \lfloor nt_1 \rfloor}}U_{\lfloor nt_2 \rfloor}^{(i)} \leq x_2, A_{n,\varepsilon}\right)
$$
\n
$$
\sim \mathbf{P}\left(\frac{1}{\Sigma_1}\sum_{j=-i}^{i+1} \zeta_j^* e^{-S_j^*} \leq x_1\right) \mathbf{P}\left(\frac{1}{\Sigma_1}\sum_{j=-i}^{i+1} \zeta_j^* e^{-S_j^*} \leq x_2\right)
$$
\n
$$
\times \int_{-\infty}^{0} \int_{0}^{+\infty} \mathbf{P}\left(\frac{L_{\lfloor nt_1 \rfloor}}{C_n} \in da, \frac{S_{\lfloor nt_1 \rfloor} - L_{\lfloor nt_1 \rfloor}}{C_n} \in db\right)
$$
\n
$$
\times \mathbf{P}\left(\frac{L_{\lfloor nt_2 \rfloor - \lfloor nt_1 \rfloor}}{C_n} < b - a - \varepsilon\right).
$$

Since the last integral is equal to $\mathbf{P}(A_{n,\varepsilon})$, we obtain (71) and, as result, the required relation (70).

It follows from (70) that (see (67))

$$
\lim_{i \to \infty} \lim_{n \to \infty} \mathbf{P}\left(\frac{a_{\lfloor nt_1 \rfloor}}{b_{\lfloor nt_1 \rfloor}} U_{\lfloor nt_1 \rfloor}^{(i)} \le x_1, \frac{b_{\lfloor nt_2 \rfloor}}{a_{\lfloor nt_k \rfloor}} U_{\lfloor nt_2 \rfloor}^{(i)} \le x_2 \middle| A_{n,\varepsilon} \right)
$$
\n
$$
= \mathbf{P}\left(\frac{\Sigma_2}{\Sigma_1} \le x_1\right) \mathbf{P}\left(\frac{\Sigma_2}{\Sigma_1} \le x_2\right). \tag{72}
$$

Applying now the same arguments which we have used in First part of the proof to establish (57) from (67) , we obtain (68) from (72) .

We now prove (69). To this aim we check that, for any fixed $i \in \mathbb{N}$ and for all but a countable set of (x_1, x_2) with $x_1, x_2 \ge 0$,

$$
\lim_{n \to \infty} \mathbf{P}\left(\frac{a_{\lfloor nt_1 \rfloor}}{b_{\lfloor nt_1 \rfloor}} U_{\lfloor nt_1 \rfloor}^{(i)} \le x_1, \frac{b_{\lfloor nt_2 \rfloor}}{a_{\lfloor nt_k \rfloor}} U_{\lfloor nt_2 \rfloor}^{(i)} \le x_2 \middle| B_{n,\varepsilon}\right)
$$

$$
= \mathbf{P}\left(\frac{1}{\Sigma_{1}}\sum_{j=-i}^{i+1} \zeta_{j}^{*} e^{-S_{j}^{*}} \le \min(x_{1}, x_{2})\right). \tag{73}
$$

Set

$$
Z'_{i,n}(m) = Z_{\tau_n + i, m},
$$

$$
U_n^{(i)}(m) = \sum_{j=-i}^{i+1} Z'_{j,n}(m).
$$

Given that the random event $B_{n,\varepsilon}$ occurred, $\tau_{\lfloor nt_2\rfloor} = \tau_{\lfloor nt_1\rfloor}$ and

$$
\frac{b_{\lfloor nt_2 \rfloor}}{a_{\lfloor nt_2 \rfloor}} \sim \frac{b_{\lfloor nt_1 \rfloor}}{a_{\lfloor nt_2 \rfloor}}
$$

as $n \to \infty$. Therefore

$$
U_{\lfloor nt_1 \rfloor}^{(i)} = \sum_{j=-i}^{i+1} Z'_{i,\lfloor nt_1 \rfloor} (\lfloor nt_1 \rfloor) = U_{\lfloor nt_1 \rfloor}^{(i)} (\lfloor nt_1 \rfloor),
$$

$$
U_{\lfloor nt_2 \rfloor}^{(i)} = \sum_{j=-i}^{i+1} Z'_{i,\lfloor nt_1 \rfloor} (\lfloor nt_2 \rfloor) = U_{\lfloor nt_1 \rfloor}^{(i)} (\lfloor nt_2 \rfloor).
$$

Thus, to prove (73) it is sufficient to show that

$$
\lim_{n\to\infty} \mathbf{P}\left(\frac{a_{\lfloor nt_1 \rfloor}}{b_{\lfloor nt_1 \rfloor}} U_{\lfloor nt_1 \rfloor}^{(i)}\left(\lfloor nt_1 \rfloor\right) \leq x_1, \frac{a_{\lfloor nt_2 \rfloor}}{b_{\lfloor nt_1 \rfloor}} U_{\lfloor nt_1 \rfloor}^{(i)}\left(\lfloor nt_2 \rfloor\right) \leq x_2 \middle| B_{n,\varepsilon}\right)
$$
\n
$$
= \mathbf{P}\left(\frac{1}{\Sigma_1} \sum_{j=-i}^{i+1} \zeta_j^* e^{-S_j^*} \leq \min\left(x_1, x_2\right)\right). \tag{74}
$$

Applying the arguments similar to those used to establish relation (19), we can show that

$$
\left\{a'_{i,m}Z'_{i,n}\left(m\right),\,i\in\mathbf{Z}\right\}\stackrel{D}{\rightarrow}\left\{\zeta_{i}^{*},\,i\in\mathbf{Z}\right\},\
$$

as $m \geq n \to \infty$. Moreover,

$$
\left\{ \left(a'_{i,n} Z'_{i,n}(n), a'_{i,m} Z'_{i,n}(m) \right), i \in \mathbf{Z} \right\} \xrightarrow{D} \left\{ \left(\zeta_i^*, \zeta_i^* \right), i \in \mathbf{Z} \right\} \tag{75}
$$

and the left-hand side of this relation is asymptotically independent from the random event $\{C_n^{-1}L_n \le a, C_n^{-1}(S_n - L_n) \le b\}$ for any $a \le 0$ and $b > 0$. It follows from (75) that (see the proof of (64))

$$
\left(\frac{a_n}{b_n}U_n^{(i)}(n), \frac{a_m}{b_n}U_n^{(i)}(m)\right) \xrightarrow{D} \frac{1}{\Sigma_1} \left(\sum_{j=-i}^{i+1} \zeta_j^* e^{-S_j^*}, \sum_{j=-i}^{i+1} \zeta_j^* e^{-S_j^*}\right),\tag{76}
$$

as $m \ge n \to \infty$. From (76) we obtain the desired relation (74) and, as result, (73). Now statement (69) follows from (73) in a standard way.

Finally, according to the Skorokhod functional limit theorem (see (1))

$$
\lim_{\varepsilon \to 0} \lim_{n \to \infty} \mathbf{P}(A_{n,\varepsilon}) = \mathbf{P}(L(t_1) > L(t_1, t_2)) = \mathbf{P}(L(t_1) > L(t_2)), \quad (77)
$$

$$
\lim_{\varepsilon \to 0} \lim_{n \to \infty} \mathbf{P}\left(B_{n,\varepsilon}\right) = \mathbf{P}\left(L\left(t_1\right) < L\left(t_1, t_2\right)\right) = \mathbf{P}\left(L\left(t_1\right) = L\left(t_2\right)\right),\tag{78}
$$

where $L(t_1, t_2) = \inf_{t \in [t_1, t_2]} W(t)$, and

$$
\lim_{\varepsilon \to 0} \lim_{n \to \infty} \mathbf{P}(D_{n,\varepsilon}) = 0.
$$
\n(79)

By the total probability formula

$$
\mathbf{P}\left(\frac{a_{\lfloor nt_1 \rfloor}}{b_{\lfloor nt_1 \rfloor}} Z_{\lfloor nt_1 \rfloor} \leq x_1, \frac{a_{\lfloor nt_2 \rfloor}}{b_{\lfloor nt_2 \rfloor}} Z_{\lfloor nt_2 \rfloor} \leq x_2\right)
$$
\n
$$
= \mathbf{P}\left(\frac{a_{\lfloor nt_1 \rfloor}}{b_{\lfloor nt_1 \rfloor}} Z_{\lfloor nt_1 \rfloor} \leq x_1, \frac{a_{\lfloor nt_2 \rfloor}}{b_{\lfloor nt_2 \rfloor}} Z_{\lfloor nt_2 \rfloor} \leq x_2 \middle| A_{n,\varepsilon}\right) \mathbf{P}(A_{n,\varepsilon})
$$
\n
$$
+ \mathbf{P}\left(\frac{a_{\lfloor nt_1 \rfloor}}{b_{\lfloor nt_1 \rfloor}} Z_{\lfloor nt_1 \rfloor} \leq x_1, \frac{a_{\lfloor nt_2 \rfloor}}{b_{\lfloor nt_2 \rfloor}} Z_{\lfloor nt_2 \rfloor} \leq x_2 \middle| B_{n,\varepsilon}\right) \mathbf{P}(B_{n,\varepsilon})
$$
\n
$$
+ \mathbf{P}\left(\frac{a_{\lfloor nt_1 \rfloor}}{b_{\lfloor nt_1 \rfloor}} Z_{\lfloor nt_1 \rfloor} \leq x_1, \frac{a_{\lfloor nt_2 \rfloor}}{b_{\lfloor nt_2 \rfloor}} Z_{\lfloor nt_2 \rfloor} \leq x_2 \middle| D_{n,\varepsilon}\right) \mathbf{P}(D_{n,\varepsilon}). \quad (80)
$$

Combining (68) , (69) and $(77)-(80)$ we deduce that

$$
\lim_{n \to \infty} \mathbf{P}\left(\frac{a_{\lfloor nt_1 \rfloor}}{b_{\lfloor nt_1 \rfloor}} Z_{\lfloor nt_1 \rfloor} \le x_1, \frac{a_{\lfloor nt_2 \rfloor}}{b_{\lfloor nt_2 \rfloor}} Z_{\lfloor nt_2 \rfloor} \le x_2\right)
$$
\n
$$
= \mathbf{P}(\gamma_1 \le x_1, \gamma_2 \le x_2) \mathbf{P}(L(t_1) > L(t_2))
$$
\n
$$
+ \mathbf{P}(\gamma_1 \le x_1, \gamma_1 \le x_2) \mathbf{P}(L(t_1) = L(t_2)),
$$

This gives the desired convergence of two-dimensional distributions.

Third part. The proof of convergence of multidimensional distributions (for dimensions exceeding two) is carried out by induction using the reasonings of Second part of the proof.

References

[1] K. Athreya, P. Ney, Branching processes, Springer-Verlag, Berlin-Heidelberg-New-York, 1972.

- [2] E. Dyakonova, L. Doudou, V. Vatutin, M. Zhang, Branching processes in a random environment with immigration stopped at zero (in print).
- [3] O. Kallenberg, Foundations of modern probability. 2nd Edition, Springer-Verlag, Berlin-Heidelberg-New-York, 2002.
- [4] V. Afanasyev, J. Geiger, G. Kersting, V. Vatutin, A limit theorem for a critical branching process in random environment, Ann. Probab. 33 (2) (2005) 645–673. [doi:10.1214/009117904000000928](http://dx.doi.org/10.1214/009117904000000928).
- [5] G. Kersting, V. Vatutin, Discrete time branching processes in random environment, John Viley and Sons, Inc., London, 2017.
- [6] H. Kesten, M. Kozlov, F. Spitzer, A limit law for random walk in a random environment, Compositio mathematica 30 (1975) 145–168.
- [7] E. Key, Limiting distributions and regeneration times for multitype branching processes with immigration in a random environment, Ann. Probab. 15 (1) (1987) 344–353.
- [8] A. Roitershtein, A note on multitype branching processes with immigration in a random environment, Ann. Probab. 35 (4) (2007) 1573–1592. [doi:10.1214/009117906000001015](http://dx.doi.org/10.1214/009117906000001015).
- [9] E. Bauernschubert, Recurrence and transience of critical branching processes in random environment with immigration and an application to excited random walks, Adv. in Appl. Probab. 46 (3) (2014) 683–703. [doi:10.1239/aap/1409319555](http://dx.doi.org/10.1239/aap/1409319555).
- [10] V. Afanasyev, About time of reaching a high level by a random walk in a random environment, Theory Probab. Appl. 57 (4) (2013) 547–567. [doi:10.1137/S0040585X97986175](http://dx.doi.org/10.1137/S0040585X97986175).
- [11] V. Afanasyev, Conditional limit theorem for maximum of random walk in a random environment, Theory Probab. Appl. 58 (4) (2014) 525–545. [doi:10.1137/S0040585X97986746](http://dx.doi.org/10.1137/S0040585X97986746).
- [12] V. Afanasyev, On the time of attaining a high level by a transient random walk in a random environment, Theory Probab. Appl. 61 (2) (2017) 178– 207. [doi:10.1137/S0040585X97T988101](http://dx.doi.org/10.1137/S0040585X97T988101).
- [13] V. Afanasyev, On the non-recurrent random walk in a random environment, Discrete Math. Appl. 28 (3) (2018) 139–156. [doi:10.1515/dma-2018-0014](http://dx.doi.org/10.1515/dma-2018-0014).
- [14] D. Li, V. Vatutin, M. Zhang, Subcritical branching processes in random environment with immigration stopped at zero, J. Theor. Probability (2020) 1–23[doi:10.1007/s10959-020-00991-5](http://dx.doi.org/10.1007/s10959-020-00991-5).
- [15] N. Bingham, C. Goldie, J.L.Teugels, Regular variation, Cambridge Univ. Press, Cambridge, 1987.
- [16] V. Afanasyev, A limit theorem for a critical branching process in random environment, Diskretnaya Matematika 5 (1) (1993) 45–58.
- [17] A. Shiryaev, Probability-1, Springer, New York, 2018.