

IMPROVED APPROXIMATIONS OF HEDGES' g^*

XIAOHUAN XUE

ABSTRACT. Hedges' unbiased estimator g^* has been broadly used in statistics. We propose a sequence of polynomials to better approximate the multiplicative correction factor of g^* by incorporating analytic estimations to the ratio of gamma functions.

1. INTRODUCTION

Hedges proposed the widely used unbiased estimator, g^* , of standardized mean differences [1, 2, 3]. Suppose group i with sample size n_i is y_{i1}, \dots, y_{in_i} , and group j with sample size n_j are y_{j1}, \dots, y_{jn_j} , and we assume the samples in two groups are normally distributed with the same variance, that is,

$$(1.1) \quad y_{i1}, \dots, y_{in_i} \sim N(\mu_i, \sigma^2)$$

$$(1.2) \quad y_{j1}, \dots, y_{jn_j} \sim N(\mu_j, \sigma^2)$$

Hedges' g^* is given by the sample mean difference divided by a multiple of the pooled sample standard deviation as following

$$(1.3) \quad g^* = J(n_i + n_j - 2) \cdot \frac{\bar{y}_j - \bar{y}_i}{s_p}$$

where

$$(1.4) \quad J(m) = \frac{\Gamma(\frac{m}{2})}{\sqrt{\frac{m}{2}} \Gamma(\frac{m-1}{2})}$$

and s_p is the pooled sample standard deviation

$$(1.5) \quad s_p = \sqrt{\frac{(n_j - 1)s_j^2 + (n_i - 1)s_i^2}{n_j + n_i - 2}}$$

The multiplicative correction term $J(\cdot)$ in Hedges' g^* is not easy to calculate in practice, and one commonly used approximation is given by Hedges [1, 2]

$$(1.6) \quad J(m) = \frac{\Gamma(\frac{m}{2})}{\sqrt{\frac{m}{2}} \Gamma(\frac{m-1}{2})} \approx 1 - \frac{3}{4m - 1}$$

and let us denote Hedges' estimation as $H(m)$:

$$(1.7) \quad H(m) = 1 - \frac{3}{4m - 1}$$

2010 *Mathematics Subject Classification*. Primary 41A10; Secondary 33B15, 41A60, 62H12, 62P10.

Key words and phrases. Gamma function, polynomial approximation, standardized mean difference.

We now propose a sequence of polynomials to give more accurate approximations of $J(\cdot)$ and thus improve the accuracy of Hedges' g^* .

2. WALLIS RATIO AND APPROXIMATION OF HEDGE'S ESTIMATOR

The Wallis ratio [4, 5, 6] is defined by

$$(2.1) \quad \frac{\Gamma(x+1)}{\Gamma(x+\frac{1}{2})}$$

and we can see

$$(2.2) \quad \sqrt{\frac{m}{2}} J(m) = \frac{\Gamma(\frac{m}{2})}{\Gamma(\frac{m-1}{2})}$$

is a special case of Wallis ratio when $x = \frac{m}{2} - 1$. Thus the approximations of Wallis ratio as a result of properties of gamma function can be applied to estimating $J(\cdot)$.

For Wallis ratio, there are Mortici's approximations [5, 6, 7]

$$(2.3) \quad \frac{\Gamma(x+1)}{\Gamma(x+\frac{1}{2})} \approx \sqrt{x+\frac{1}{4}}$$

$$(2.4) \quad \frac{\Gamma(x+1)}{\Gamma(x+\frac{1}{2})} \approx \sqrt[4]{x^2+\frac{1}{2}x+\frac{1}{8}}$$

$$(2.5) \quad \frac{\Gamma(x+1)}{\Gamma(x+\frac{1}{2})} \approx \sqrt[6]{x^3+\frac{3}{4}x^2+\frac{9}{32}x+\frac{5}{128}}$$

$$(2.6) \quad \frac{\Gamma(x+1)}{\Gamma(x+\frac{1}{2})} \approx \sqrt[8]{x^4+x^3+\frac{1}{2}x^2+\frac{1}{8}x}$$

$$(2.7) \quad \frac{\Gamma(x+1)}{\Gamma(x+\frac{1}{2})} \approx \sqrt[10]{x^5+\frac{5}{4}x^4+\frac{25}{32}x^3+\frac{35}{128}x^2+\frac{75}{2048}x+\frac{3}{8192}}$$

$$(2.8) \quad \frac{\Gamma(x+1)}{\Gamma(x+\frac{1}{2})} \approx \sqrt[12]{x^6+\frac{3}{2}x^5+\frac{9}{8}x^4+\frac{1}{2}x^3+\frac{15}{128}x^2+\frac{3}{256}x+\frac{11}{1024}}$$

By letting $x = \frac{m}{2} - 1$, we can have an approximation of $J(m)$

$$(2.9) \quad J(m) \approx P_1(m) = \sqrt{1-\frac{3}{2m}}$$

$$(2.10) \quad \begin{aligned} J(m) \approx P_2(m) &= \sqrt{\frac{2}{m} \sqrt[4]{(\frac{m}{2}-1)^2 + \frac{1}{2}(\frac{m}{2}-1) + \frac{1}{8}}} \\ &= \sqrt[4]{1-\frac{3}{m} + \frac{5}{2m^2}} \end{aligned}$$

$$(2.11) \quad \begin{aligned} J(m) \approx P_3(m) &= \sqrt{\frac{2}{m} \sqrt[6]{\left(\frac{m}{2} - 1\right)^3 + \frac{3}{4}\left(\frac{m}{2} - 1\right)^2 + \frac{9}{32}\left(\frac{m}{2} - 1\right) + \frac{5}{128}}} \\ &= \sqrt[6]{1 - \frac{9}{2m} + \frac{57}{8m^2} - \frac{63}{16m^3}} \end{aligned}$$

$$(2.12) \quad \begin{aligned} J(m) \approx P_4(m) &= \sqrt{\frac{2}{m} \sqrt[8]{\left(\frac{m}{2} - 1\right)^4 + \left(\frac{m}{2} - 1\right)^3 + \frac{1}{2}\left(\frac{m}{2} - 1\right)^2 + \frac{1}{8}\left(\frac{m}{2} - 1\right)}} \\ &= \sqrt[8]{1 - \frac{6}{m} + \frac{14}{m^2} - \frac{15}{m^3} + \frac{6}{m^4}} \end{aligned}$$

(2.13)

$$J(m) \approx P_5(m)$$

$$\begin{aligned} &= \sqrt{\frac{2}{m} \sqrt[10]{\left(\frac{m}{2} - 1\right)^5 + \frac{5}{4}\left(\frac{m}{2} - 1\right)^4 + \frac{25}{32}\left(\frac{m}{2} - 1\right)^3 + \frac{35}{128}\left(\frac{m}{2} - 1\right)^2 + \frac{75}{2048}\left(\frac{m}{2} - 1\right) + \frac{3}{8192}}} \\ &= \sqrt[10]{1 - \frac{15}{2m} + \frac{185}{8m^2} - \frac{585}{16m^3} + \frac{3755}{128m^4} - \frac{2409}{256m^5}} \end{aligned}$$

$$(2.14) \quad J(m) \approx P_6(m) = \sqrt[12]{1 - \frac{9}{m} + \frac{69}{2m^2} - \frac{72}{m^3} + \frac{687}{8m^4} - \frac{441}{8m^5} + \frac{247}{16m^6}}$$

3. ACCURACY OF DIFFERENT APPROXIMATIONS

In this section, we compare the accuracy of all the given approximations by measuring the absolute values of their errors to the real value with a broad range of m . For convenience, we introduce the notations for the absolute errors

$$(3.1) \quad \delta_0(m) = |H(m) - J(m)|$$

$$(3.2) \quad \delta_i(m) = |P_i(m) - J(m)|; i = 1, \dots, 6.$$

Figure 1 shows the performance of approximations and we can see Hedges' approximation $H(m)$, which is the dashed red line, has less accuracy compared to $P_2(m), P_3(m), \dots, P_m(m)$. In terms of absolute errors, we can see

$$(3.3) \quad \delta_6(m) < \delta_5(m) < \delta_4(m) < \delta_3(m) < \delta_2(m) < \delta_0(m) < \delta_1(m).$$

This can also be verified by performing numerical analysis demonstrated by Table 1, from which the order of accuracy is more straightforward.

TABLE 1. Numerical accuracy of different approximations

m	δ_0	δ_1	δ_2	δ_3	δ_4	δ_5	δ_6
10	0.00033	0.00079	$5.34e-06$	$4.66e-06$	$1.43e-07$	$1.35e-07$	$8.47e-09$
30	$3.55e-05$	$7.49e-05$	$4.60e-08$	$4.03e-08$	$1.13e-10$	$1.067e-10$	$6.24e-13$
50	$1.27e-05$	$2.62e-05$	$5.56e-09$	$4.86e-09$	$4.71e-12$	$4.45e-12$	$1.79e-14$
70	$6.44e-06$	$1.32e-05$	$1.40e-09$	$1.23e-09$	$5.93e-13$	$5.60e-13$	$5.66e-15$
100	$3.15e-06$	$6.39e-06$	$3.29e-10$	$2.88e-10$	$4.49e-14$	$4.11e-14$	$2.31e-14$
200	$7.84e-07$	$1.58e-06$	$2.00e-11$	$1.75e-11$	$4.29e-14$	$4.29e-14$	$4.19e-14$




FIGURE 1. The approximation accuracy of H and P_n .




FIGURE 2. The approximation accuracy of P_5 and P_6 when $m > 200$.

4. DISCUSSION

In this paper, we have proposed a sequence of more accurate approximations to the multiplicative correction factor, $J(m)$, in Hedges' unbiased estimator of standardized mean difference.

It is also worth mentioning that the difference between $P_5(m)$ and $P_6(m)$ are small when m is over 100, and there is almost no difference when m is over 200. We can also see this from Figure 2, both $P_5(m)$ and $P_6(m)$ are oscillating around 0 within the magnitude of 2×10^{-13} .

More accurate and efficient approximations to Hedges' g^* would be available with deeper understandings of the properties of gamma functions, which has always been an appealing topic in both mathematics itself and our future work.

REFERENCES

1. Hedges, L. V. (1981) *Distribution Theory for Glass Estimator of Effect size and Related Estimators*, Journal of Educational Statistics, 6(2), pp. 107128. doi: 10.3102/10769986006002107.
2. Hedges, L. V., and Olkin I. (2014) *Statistical Methods for Meta-Analysis*. Academic Press.
3. Borenstein, Michael, Larry V. Hedges, Julian P. T. Higgins, and Hannah Rothstein. (2009) *Introduction to Meta-Analysis*. Chichester, U.K. John Wiley and Sons.
4. J. T. Chu, *A modified Wallis product and some applications*, Amer. Math. Monthly 69 (1962), no. 5, 402-404.
5. C. Mortici, *New approximation formulas for evaluating the ratio of gamma functions*, Math. Comp. Modelling 52 (2010) 425433.
6. Dumitrescu, S., *Estimates for the ratio of gamma functions by using higher order roots*, Studia Universitatis Babeş-Bolyai, Mathematica . Jun2015, Vol. 60 Issue 2, p173-181. 9p.
7. Paris, R. B. (2011). *Asymptotic approximations for $n!$* , Applied Mathematical Sciences.
8. Xue, X. (2020) *Estimation of within-study covariances in multivariate meta-analysis*, arXiv:2003.05092.

DEPARTMENT OF MATHEMATICS AND STATISTICS, WAKE FOREST UNIVERSITY, WINSTON-SALEM,
NORTH CAROLINA 27109

Current address: Department of Psychology, University of California, Los Angeles, California
90095

E-mail address: xhxue@g.ucla.edu