

THE RECONSTRUCTION CONJECTURE FOR FINITE SIMPLE GRAPHS AND ASSOCIATED DIRECTED GRAPHS

TETSUYA HOSAKA

ABSTRACT. In this paper, we study the Reconstruction Conjecture for finite simple graphs. Let Γ and Γ' be finite simple graphs with at least three vertices such that there exists a bijective map $f : V(\Gamma) \rightarrow V(\Gamma')$ and for any $v \in V(\Gamma)$, there exists an isomorphism $\phi_v : \Gamma - v \rightarrow \Gamma' - f(v)$. Then we define the associated directed graph $\tilde{\Gamma} = \tilde{\Gamma}(\Gamma, \Gamma', f, \{\phi_v\}_{v \in V(\Gamma)})$ with two kinds of arrows from the graphs Γ and Γ' , the bijective map f and the isomorphisms $\{\phi_v\}_{v \in V(\Gamma)}$. By investigating the associated directed graph $\tilde{\Gamma}$, we study when are the two graphs Γ and Γ' isomorphic.

1. INTRODUCTION AND PRELIMINARIES

A finite simple graph Γ is said to be *reconstructible*, if any simple graph Γ' with the following property (*) is isomorphic to Γ .

- (*) There exists a bijective map $f : V(\Gamma) \rightarrow V(\Gamma')$ such that for any $v \in V(\Gamma)$, there exists an isomorphism $\phi_v : \Gamma - v \rightarrow \Gamma' - f(v)$.

Here $V(\Gamma)$ and $V(\Gamma')$ are the vertex sets of Γ and Γ' respectively. Also $\Gamma - v$ and $\Gamma' - f(v)$ are the full subgraphs of Γ and Γ' whose vertex sets are $V(\Gamma) - \{v\}$ and $V(\Gamma') - \{f(v)\}$ respectively.

The Reconstruction Conjecture (also often called Ulam Conjecture or Kelly-Ulam Conjecture) introduced by P.J. Kelly and S.M. Ulam in 1942 is one of the very famous open problems in Graph Theory and in Mathematics.

Reconstruction Conjecture. Every finite simple graph with at least three vertices will be reconstructible.

Some classes of reconstructible graphs are known (see [1], [2], [3], [4], [5], [6], [7], [8], [9], [10], [11], [12] and [13]) as follows: Let Γ be a finite simple graph with at least three vertices.

- (i) If Γ is a regular graph, then it is reconstructible.
- (ii) If Γ is a tree, then it is reconstructible.
- (iii) If Γ is not connected, then it is reconstructible.
- (iv) If Γ has at most 11 vertices, then it is reconstructible.
- (v) If Γ is a maximal planar graph, then it is reconstructible.
- (vi) If Γ is a finite graph that is the 1-skeleton of some simplicial flag complex that is a homology manifold of dimension $n \geq 1$, then it is reconstructible.

Date: June 18, 2021.

2010 Mathematics Subject Classification. 05C60.

Key words and phrases. the reconstruction conjecture; reconstructible graphs.

Also it is known that if all 2-connected graphs are reconstructible, then every finite simple graph is reconstructible [14].

Let Γ and Γ' be finite simple graphs with at least three vertices satisfying the property (*) as above.

Then we define the directed graph

$$\tilde{\Gamma} = \tilde{\Gamma}(\Gamma, \Gamma', f, \{\phi_v\}_{v \in V(\Gamma)})$$

with two kinds of arrows (called *the associated directed graph*) as follows:

- (1) The vertex set $V(\tilde{\Gamma})$ of $\tilde{\Gamma}$ is $V(\Gamma)$.
- (2) For $v_1, v_2 \in V(\tilde{\Gamma})$, there is a normal-arrow from v_1 to v_2 in $\tilde{\Gamma}$, denoted by $v_1 \longrightarrow v_2$, if and only if $[v_1, v_2] \notin E(\Gamma)$ and $[f(v_1), \phi_{v_1}(v_2)] \in E(\Gamma')$.
- (3) For $v_1, v_2 \in V(\tilde{\Gamma})$, there is a dashed-arrow from v_1 to v_2 in $\tilde{\Gamma}$, denoted by $v_1 \dashrightarrow v_2$, if and only if $[f(v_1), f(v_2)] \notin E(\Gamma')$ and $[v_1, \phi_{v_1}^{-1}(f(v_2))] \in E(\Gamma)$

(see Figure 1).




FIGURE 1. Definition of arrows in $\tilde{\Gamma}$

The purpose of this paper is to consider when are the two graphs Γ and Γ' isomorphic by investigating the associated directed graph $\tilde{\Gamma}$, and we study the problem of when are finite simple graphs reconstructible.

We give some examples of $\tilde{\Gamma}$ in Section 2 and we investigate some properties of $\tilde{\Gamma}$ in Section 3.

Proposition 3.2. *For all $v \in V(\tilde{\Gamma})$, the number of normal-arrows arising from v and the number of dashed-arrows arising from v are equal in $\tilde{\Gamma}$.*

We show the following theorem in Section 3.

Theorem 3.3. *For any vertex $v \in V(\Gamma) = V(\tilde{\Gamma})$, the following two statements are equivalent:*

- (i) *There are no normal-arrows arising from v (and hence no dashed-arrows arising from v) in the associated directed graph $\tilde{\Gamma}$.*
- (ii) *The isomorphism $\phi_v : \Gamma - v \rightarrow \Gamma' - f(v)$ extends to the isomorphism $\overline{\phi_v} : \Gamma \rightarrow \Gamma'$ as $\overline{\phi_v}|_{\Gamma - v} = \phi_v$ and $\overline{\phi_v}(v) = f(v)$.*

By Theorem 3.3, if for some vertex $v \in V(\tilde{\Gamma})$,

- (i) there are no normal-arrows arising from v (and no dashed-arrows arising from v) in the associated directed graph $V(\tilde{\Gamma})$,

then $\Gamma \cong \Gamma'$.

Now we state the otherwise case. Suppose that for every vertex $v \in V(\tilde{\Gamma})$ the statement (i) above does not hold. Then there exist a normal-arrow and a dashed-arrow arising from each vertex $v \in V(\tilde{\Gamma})$ by Proposition 3.2. Since the directed graph $\tilde{\Gamma}$ is finite, for some positive number $k \in \mathbb{N}$ and some vertices $v_1, \dots, v_{2k} \in V(\tilde{\Gamma})$, there exist arrows $v_{2i-1} \rightarrow v_{2i}$ and $v_{2i} \dashrightarrow v_{2i+1}$ in $\tilde{\Gamma}$ for any $i = 1, \dots, k$, where $v_{2k+1} := v_1$ (see Figure 2). Here the vertices v_1, \dots, v_{2k} need not be different all together.




FIGURE 2. Examples of cycles with alternate normal-arrows and dashed-arrows

We investigate on cycles with alternate normal-arrows and dashed-arrows in $\tilde{\Gamma}$ in Section 4. For such a cycle, we define “ α -type” and “ β -type”. We show that if such a cycle is α -type then there exist normal-arrows $v_1 \rightarrow v_2$ and $v_2 \rightarrow v_1$ in $\tilde{\Gamma}$ in Section 4.

Proposition 4.2. *If for some positive number $k \in \mathbb{N}$ and some sequence $v_1, \dots, v_{2k} \in V(\tilde{\Gamma})$, there exist arrows $v_{2i-1} \rightarrow v_{2i}$ and $v_{2i} \dashrightarrow v_{2i+1}$ in $\tilde{\Gamma}$ for any $i = 1, \dots, k$ (where $v_{2k+1} := v_1$) and if the cycle v_1, \dots, v_{2k} is α -type then there exist normal-arrows $v_1 \rightarrow v_2$ and $v_2 \rightarrow v_1$ in $\tilde{\Gamma}$.*

By Proposition 4.2, we obtain the following.

Corollary 4.3. *Let Γ and Γ' be finite simple graphs with at least three vertices satisfying the property (*). Then for the associated directed graph $\tilde{\Gamma}$,*

- (I) *there exists a vertex $v \in V(\tilde{\Gamma})$ with no normal-arrow arising from v in $\tilde{\Gamma}$ (then $\Gamma \cong \Gamma'$ by Theorem 3.3),*
- (II) *for some $v_1, v_2 \in V(\tilde{\Gamma})$, there exist normal-arrows $v_1 \rightarrow v_2$ and $v_2 \rightarrow v_1$ in $\tilde{\Gamma}$, or*
- (III) *some and all cycles with alternate normal-arrows and dashed-arrows in $\tilde{\Gamma}$ are β -type.*

In the case that (II) holds and the cycle v_1, v_2 is α -type, we investigate structures of Γ and Γ' in Section 5. As a preparation, we define a finite simple graph $A = A(n; B, C)$.

Definition 1.1. For a number $n \in \mathbb{N}$ at least 3 and subsets B and C of the set $\{1, \dots, n-1\}$, we define the finite simple graph $A = A(n; B, C)$ as follows:

- (1) The graph A has n vertices denoted by $V(A) = \{a_1, a_2, \dots, a_n\}$.
- (2) $b \in B$ if and only if $[a_{2t-1}, a_{2t-1+b}] \in E(A)$ for any number $t \in \mathbb{N}$ as $1 \leq 2t-1 < 2t-1+b \leq n$.

- (3) $c \in C$ if and only if $[a_{2t}, a_{2t+c}] \in E(A)$ for any number $t \in \mathbb{N}$ as $2 \leq 2t < 2t + c \leq n$.

We give some examples of $A(n; B, C)$ in Figure 3.




FIGURE 3. Examples of graphs $A(n; B, C)$ for $n = 7, 8$

Then we have the following by the definition of $A(n; B, C)$.

Lemma 1.2. *Let $n \in \mathbb{N}$ be a number at least 3 and let B and C be subsets of the set $\{1, \dots, n-1\}$. Let $A := A(n; B, C)$ and $A' := A(n; C, B)$, where $V(A) = \{a_1, a_2, \dots, a_n\}$ and $V(A') = \{a'_1, a'_2, \dots, a'_n\}$ are the numbering of vertices as in the definition of $A = A(n; B, C)$ and $A' = A(n; C, B)$. Let $a'_i := a_{n-i+1}$ for $i = 1, \dots, n$. (Hence $V(A') = \{a'_1, a'_2, \dots, a'_n\}$.) Then the two graphs A and A' satisfy the following:*

- (1) *There exists the isomorphism $\Psi : A - \{a_1, a_2\} \rightarrow A - \{a_{n-1}, a_n\}$ such that $\Psi(a_i) = a_{i-2}$ for any $i = 3, \dots, n$.*
- (2) *There exists the isomorphism $\Psi' : A' - \{a'_1, a'_2\} \rightarrow A' - \{a'_{n-1}, a'_n\}$ such that $\Psi'(a'_i) = a'_{i-2}$ for any $i = 3, \dots, n$.*
- (3) *There exists the isomorphism $\phi_1 : A - a_1 \rightarrow A' - a'_1$ such that $\phi_1(a_i) = a'_{n-i+2}$ for any $i = 2, \dots, n$.*
- (4) *There exists the isomorphism $\phi_2 : A - a_n \rightarrow A' - a'_n$ such that $\phi_2(a_i) = a'_{n-i}$ for any $i = 1, \dots, n-1$.*

Definition 1.3. Let $A := A(n; B, C)$ where n is a number at least 3 and B and C are subsets of the set $\{1, \dots, n-1\}$. Then we define the numbers $\beta(A)$ and $\gamma(A)$ as follows;

$$\beta(A) := \#\{b \in B \mid n - b \text{ is odd}\} \text{ and}$$

$$\gamma(A) := \#\{c \in C \mid n - c \text{ is odd}\}.$$

Here we note that $[n - b \text{ is odd}]$ if and only if $[[b \text{ is even if } n \text{ is odd}] \text{ and } [b \text{ is odd if } n \text{ is even}]]$.

For a vertex v of a graph Γ , we denote $\deg_{\Gamma} v$ as the degree of the vertex v in the graph Γ . If Γ and Γ' are finite simple graphs with at least three vertices satisfying the property $(*)$, then it is well known that $\deg_{\Gamma} v = \deg_{\Gamma'} f(v)$ for all $v \in V(\Gamma)$.

In Sections 5 and 6, we give some remarks and examples of $A = A(n; B, C)$ and $A' = A(n; C, B)$. We obtain the following in Section 5.

Lemma 5.7. *Let $A = A(n; B, C)$, $A' = A(n; C, B)$, $V(A) = \{a_1, a_n, \dots, a_n\}$ and $V(A') = \{a'_1, a'_2, \dots, a'_n\}$ as in Lemma 1.2. Then the following two statements are equivalent:*

- (i) $\beta(A) = \gamma(A)$.
- (ii) $|E(A)| = |E(A')|$, $\deg_A a_1 = \deg_{A'} a'_1$ and $\deg_A a_n = \deg_{A'} a'_n$.

In Theorem 5.8, we show that for finite simple graphs Γ and Γ' with at least three vertices satisfying the property $(*)$, if in the associated directed graph $\tilde{\Gamma}$,

- (II) for some $v_1, v_2 \in V(\tilde{\Gamma})$, there exist normal-arrows $v_1 \rightarrow v_2$ and $v_2 \rightarrow v_1$ in $\tilde{\Gamma}$ and if the cycle v_1, v_2 is α -type, then Γ and Γ' have some structure (\mathcal{F}) .

An example of Γ and Γ' with the structure (\mathcal{F}) is given in Figure 4. In this example, $A = A(6; \{1\}, \{1\})$ and $A' = A(6; \{1\}, \{1\})$.




FIGURE 4. An example of graphs Γ and Γ' with the structure (\mathcal{F})

We introduce some examples and remarks on $A = A(n; B, C)$ and $A' = A(n; C, B)$ in Section 6 and finally we give an example and a remark on β -type cycles in Section 7.

2. EXAMPLES ON THE ASSOCIATED DIRECTED GRAPHS

We give some examples of the associated directed graphs.

Example 2.1. Let Γ and Γ' be the graphs as Figure 5 (I-1). We define the map $f : V(\Gamma) \rightarrow V(\Gamma')$ as $f(v_i) = v'_i$ for $i = 1, 2, 3, 4, 5$. For each $v_i \in V(\Gamma)$, we define the isomorphism $\phi_{v_i} : \Gamma - v_i \rightarrow \Gamma' - v'_i$ as Figure 5 (I-2). Then the associated directed graph $\tilde{\Gamma} = \tilde{\Gamma}(\Gamma, \Gamma', f, \{\phi_v\}_{v \in V(\Gamma)})$ becomes as Figure 5 (I-3).

Here we note that for each vertex $v \in V(\tilde{\Gamma})$, the normal-arrows and the dashed-arrows arising from v in $\tilde{\Gamma}$ are determined by the isomorphism $\phi_v : \Gamma - v \rightarrow \Gamma' - f(v)$.

In this example, there are no normal-arrows arising from v_2 (and no dashed-arrows arising from v_2). Then there exists the isomorphism $\overline{\phi_{v_2}} : \Gamma \rightarrow \Gamma'$ as $\overline{\phi_{v_2}}|_{\Gamma - v_2} = \phi_{v_2}$ and $\overline{\phi_{v_2}}(v_2) = f(v_2) = v'_2$ (as Theorem 3.3).

Example 2.2. Let Γ and Γ' be the graphs as Figure 6 (II-1). We define the map $f : V(\Gamma) \rightarrow V(\Gamma')$ as $f(v_i) = v'_i$ for $i = 1, 2, 3, 4$. For each $v_i \in V(\Gamma)$, we define the isomorphism $\phi_{v_i} : \Gamma - v_i \rightarrow \Gamma' - v'_i$ as Figure 6 (II-2). Then the associated directed graph $\tilde{\Gamma} = \tilde{\Gamma}(\Gamma, \Gamma', f, \{\phi_v\}_{v \in V(\Gamma)})$ becomes as Figure 6 (II-3).

In this example, there are no normal-arrows (and no dashed-arrows) arising from v_3 and this implies that there exists the isomorphism $\overline{\phi_{v_3}} : \Gamma \rightarrow \Gamma'$ as $\overline{\phi_{v_3}}|_{\Gamma - v_3} = \phi_{v_3}$ and $\overline{\phi_{v_3}}(v_3) = f(v_3) = v'_3$.

Example 2.3. Let Γ and Γ' be the graphs as Figure 7 (III-1) that are the same graphs in Example 2.2. We define the map $f : V(\Gamma) \rightarrow V(\Gamma')$ as $f(v_i) = v'_i$ for $i = 1, 2, 3, 4$. For each $v_i \in V(\Gamma)$, we define the isomorphism $\phi_{v_i} : \Gamma - v_i \rightarrow \Gamma' - v'_i$ as Figure 7 (III-2). Then the associated directed graph $\tilde{\Gamma} = \tilde{\Gamma}(\Gamma, \Gamma', f, \{\phi_v\}_{v \in V(\Gamma)})$ becomes as Figure 7 (III-3).

In this example, for every vertex $v \in V(\tilde{\Gamma})$, there exists a normal-arrow (and a dashed-arrow) arising from v in $\tilde{\Gamma}$. Also there are arrows $v_1 \rightarrow v_4$, $v_4 \rightarrow v_1$, $v_1 \dashrightarrow v_4$ and $v_4 \dashrightarrow v_1$ in $\tilde{\Gamma}$.

3. ON THE ASSOCIATED DIRECTED GRAPHS

Let Γ and Γ' be finite simple graphs with at least three vertices satisfying the property (*) and let $\tilde{\Gamma}$ be the associated directed graph. We investigate some properties of $\tilde{\Gamma}$. The following lemma is obtained immediately from the definition of $\tilde{\Gamma}$.

Lemma 3.1. (1) For $v_1, v_2 \in V(\tilde{\Gamma})$, if there exists a normal-arrow $v_1 \rightarrow v_2$ in $\tilde{\Gamma}$ then $[v_1, v_2] \notin E(\Gamma)$.
(2) For $v_1, v_2 \in V(\tilde{\Gamma})$, if there exists a dashed-arrow $v_1 \dashrightarrow v_2$ in $\tilde{\Gamma}$ then $[f(v_1), f(v_2)] \notin E(\Gamma')$.

We show a proposition.

Proposition 3.2. For all $v \in V(\tilde{\Gamma})$, the number of normal-arrows arising from v and the number of dashed-arrows arising from v are equal in $\tilde{\Gamma}$.

Proof. For all $v \in V(\Gamma)$, $\deg_{\Gamma} v = \deg_{\Gamma'} f(v)$ holds.

Let v be a fixed vertex of Γ and let $v' := f(v) \in V(\Gamma')$. Then the degree $\deg_{\Gamma} v$ is the sum of the number P_v of edges $[v, w]$ as $[v', \phi_v(w)] \in E(\Gamma')$ and the number Q_v of edges $[v, w]$ as $[v', \phi_v(w)] \notin E(\Gamma')$. Also, the degree $\deg_{\Gamma'} v'$ is the sum of the






FIGURE 5. Example 2.1




FIGURE 6. Example 2.2

number P'_v of edges $[v', u']$ as $[v, \phi_v^{-1}(u')] \in E(\Gamma)$ and the number Q'_v of edges $[v', u']$ as $[v, \phi_v^{-1}(u')] \notin E(\Gamma)$. Hence

$$\begin{aligned}
P_v &= \#\{w \in V(\Gamma - v) \mid [v, w] \in E(\Gamma) \text{ and } [v', \phi_v(w)] \in E(\Gamma')\}, \\
Q_v &= \#\{w \in V(\Gamma - v) \mid [v, w] \in E(\Gamma) \text{ and } [v', \phi_v(w)] \notin E(\Gamma')\}, \\
P'_{v'} &= \#\{u' \in V(\Gamma' - v') \mid [v', u'] \in E(\Gamma') \text{ and } [v, \phi_v^{-1}(u')] \in E(\Gamma)\}, \\
Q'_{v'} &= \#\{u' \in V(\Gamma' - v') \mid [v', u'] \in E(\Gamma') \text{ and } [v, \phi_v^{-1}(u')] \notin E(\Gamma)\}, \\
\deg_{\Gamma} v &= P_v + Q_v \text{ and} \\
\deg_{\Gamma'} v' &= P'_{v'} + Q'_{v'}.
\end{aligned}$$

Let $w \in V(\Gamma - v)$ and let $u' := \phi_v(w)$. Here $u' \in V(\Gamma' - v')$. Then $[v', \phi_v(w)] \in E(\Gamma')$ if and only if $[v', u'] \in E(\Gamma')$. Also $[v, w] \in E(\Gamma)$ if and only if $[v, \phi_v^{-1}(u')] \in E(\Gamma)$.

Hence $P_v = P'_{v'}$ and

$$Q_v = \deg_{\Gamma} v - P_v = \deg_{\Gamma'} v' - P'_{v'} = Q'_{v'}.$$




FIGURE 7. Example 2.3

Also

$$\begin{aligned} Q_v &= \#\{u' \in V(\Gamma' - v') \mid [v, \phi_v^{-1}(u')] \in E(\Gamma) \text{ and } [v', u'] \notin E(\Gamma')\} \\ &= \#\{u \in V(\Gamma - v) \mid v \dashrightarrow u \text{ in } \tilde{\Gamma}\} \end{aligned}$$

and

$$\begin{aligned} Q'_{v'} &= \#\{w \in V(\Gamma - v) \mid [v', \phi_v(w)] \in E(\Gamma') \text{ and } [v, w] \notin E(\Gamma)\} \\ &= \#\{w \in V(\Gamma - v) \mid v \rightarrow w \text{ in } \tilde{\Gamma}\}. \end{aligned}$$

Thus, the number of normal-arrows arising from v and the number of dashed-arrows arising from v are equal in $\tilde{\Gamma}$. \square

We show the following theorem. This is the first motivation for considering the associated directed graph $\tilde{\Gamma}$.

Theorem 3.3. *For any vertex $v \in V(\Gamma) = V(\tilde{\Gamma})$, the following two statements are equivalent:*

- (i) *There are no normal-arrows arising from v (and no dashed-arrows arising from v) in the associated directed graph $\tilde{\Gamma}$.*

(ii) The isomorphism $\phi_v : \Gamma - v \rightarrow \Gamma' - f(v)$ extends to the isomorphism $\overline{\phi}_v : \Gamma \rightarrow \Gamma'$ as $\overline{\phi}_v|_{\Gamma-v} = \phi_v$ and $\overline{\phi}_v(v) = f(v)$.

Proof. Let $v' := f(v)$. We suppose that (i) holds. Then by Proposition 3.2, there are no dashed-arrows arising from v , $Q_v = Q_{v'} = 0$ and $\deg_{\Gamma} v = P_v = P_{v'} = \deg_{\Gamma'} v'$. Hence for $w \in V(\Gamma - v)$, $[v, w] \in E(\Gamma)$ if and only if $[v', \phi_v(w)] \in E(\Gamma')$. Thus we can obtain the isomorphism $\overline{\phi}_v : \Gamma \rightarrow \Gamma'$ such that $\overline{\phi}_v|_{\Gamma-v} = \phi_v$ and $\overline{\phi}_v(v) = f(v) = v'$.

We suppose that (ii) holds. Then for any $w \in V(\Gamma - v)$, $[v, w] \in E(\Gamma)$ if and only if $[v', \phi_v(w)] \in E(\Gamma)$. By the definition of $\tilde{\Gamma}$, the statement (i) holds. \square

4. ON CYCLES WITH ALTERNATE NORMAL-ARROWS AND DASHED-ARROWS IN $\tilde{\Gamma}$

We suppose that there does not exist a vertex $v \in V(\tilde{\Gamma})$ with no normal-arrows arising from v in the associated directed graph $\tilde{\Gamma}$. Then by the argument in Section 1, there exist a positive number $k \in \mathbb{N}$ and a sequence $v_1, \dots, v_{2k} \in V(\tilde{\Gamma})$ such that there exist arrows $v_{2i-1} \rightarrow v_{2i}$ and $v_{2i} \dashrightarrow v_{2i+1}$ in $\tilde{\Gamma}$ for any $i = 1, \dots, k$, where $v_{2k+1} := v_1$ by Proposition 3.2 (see Figure 2), because $\tilde{\Gamma}$ is finite.

We define α -type and β -type for such a cycle v_1, \dots, v_{2k} .

Definition 4.1. We consider a sequence $\{b'_i\}$ in $V(\Gamma')$ as

$$\begin{aligned}
b'_1 &:= v'_1, \\
b'_2 &:= \phi_{v_1}(v_2), \\
b'_3 &:= \phi_{v_1} \circ \phi_{v_2}^{-1}(v'_3), \\
b'_4 &:= \phi_{v_1} \circ \phi_{v_2}^{-1} \circ \phi_{v_3}(v_4), \\
b'_5 &:= \phi_{v_1} \circ \phi_{v_2}^{-1} \circ \phi_{v_3} \circ \phi_{v_4}^{-1}(v'_5), \\
&\dots \\
b'_{2k} &:= \phi_{v_1} \circ \phi_{v_2}^{-1} \circ \phi_{v_3} \circ \phi_{v_4}^{-1} \circ \dots \circ \phi_{v_{2k-1}}(v_{2k}) = \overline{\phi}_{2k-1}(v_{2k}), \\
b'_{2k+1} &:= \overline{\phi}_{2k-1} \circ \phi_{v_{2k}}^{-1}(v'_1) = \Psi_0(b'_1), \\
b'_{2k+2} &:= \Psi_0(b'_2), \\
b'_{2k+3} &:= \Psi_0(b'_3), \\
b'_{2k+4} &:= \Psi_0(b'_4), \\
&\dots \\
b'_{4k} &:= \Psi_0(b'_{2k}), \\
b'_{4k+1} &:= (\Psi_0)^2(b'_1), \\
b'_{4k+2} &:= (\Psi_0)^2(b'_2), \\
b'_{4k+3} &:= (\Psi_0)^2(b'_3), \\
b'_{4k+4} &:= (\Psi_0)^2(b'_4), \\
&\dots,
\end{aligned}$$

where $\overline{\phi}_{2k-1} := \phi_{v_1} \circ \phi_{v_2}^{-1} \circ \phi_{v_3} \circ \phi_{v_4}^{-1} \circ \dots \circ \phi_{v_{2k-1}}$ and $\Psi_0 := \overline{\phi}_{2k-1} \circ \phi_{v_{2k}}^{-1} = \phi_{v_1} \circ \phi_{v_2}^{-1} \circ \phi_{v_3} \circ \phi_{v_4}^{-1} \circ \dots \circ \phi_{v_{2k-1}} \circ \phi_{v_{2k}}^{-1}$.

Here if b'_i is not defined, then we consider b'_i is blank and we denote $b'_i = \square$.

For example, $b'_5 = \phi_{v_1} \circ \phi_{v_2}^{-1} \circ \phi_{v_3} \circ \phi_{v_4}^{-1}(v'_5)$ is defined if and only if $v'_5 \neq v'_4$, $\phi_{v_4}^{-1}(v'_5) \neq v_3$, $\phi_{v_3} \circ \phi_{v_4}^{-1}(v'_5) \neq v'_2$ and $\phi_{v_2}^{-1} \circ \phi_{v_3} \circ \phi_{v_4}^{-1}(v'_5) \neq v_1$. Here we note that $v'_5 \neq v'_4$ and $\phi_{v_4}^{-1}(v'_5) \neq v_3$ always hold. Hence in this case, b'_5 is blank if and only if $\phi_{v_3} \circ \phi_{v_4}^{-1}(v'_5) = v'_2$ or $\phi_{v_2}^{-1} \circ \phi_{v_3} \circ \phi_{v_4}^{-1}(v'_5) = v_1$.

Let $b_i^{(0)} := b'_i$ ($i = 1, 2, \dots$) and let $A'_0 := \{b_i^{(0)} \mid i = 1, 2, \dots\}$.

Similarly, we consider a sequence $\{b_i\}$ in $V(\Gamma)$ as

$$\begin{aligned}
b_1 &:= v_2, \\
b_2 &:= \phi_{v_2}^{-1}(v'_3), \\
b_3 &:= \phi_{v_2}^{-1} \circ \phi_{v_3}(v_4), \\
b_4 &:= \phi_{v_2}^{-1} \circ \phi_{v_3} \circ \phi_{v_4}^{-1}(v'_5), \\
b_5 &:= \phi_{v_2}^{-1} \circ \phi_{v_3} \circ \phi_{v_4}^{-1} \circ \phi_{v_5}(v_6), \\
&\dots \\
b_{2k-1} &:= \phi_{v_2}^{-1} \circ \phi_{v_3} \circ \phi_{v_4}^{-1} \circ \dots \circ \phi_{v_{2k-1}}(v_{2k}), \\
b_{2k} &:= \phi_{v_2}^{-1} \circ \phi_{v_3} \circ \phi_{v_4}^{-1} \circ \dots \circ \phi_{v_{2k-1}} \circ \phi_{v_{2k}}^{-1}(v'_1) = \bar{\phi}_{2k}(v'_1), \\
b_{2k+1} &:= \bar{\phi}_{2k} \circ \phi_{v_1}(v_2) = \Psi'_0(b_1), \\
b_{2k+2} &:= \Psi'_0(b_2), \\
b_{2k+3} &:= \Psi'_0(b_3), \\
b_{2k+4} &:= \Psi'_0(b_4), \\
&\dots \\
b_{4k} &:= \Psi'_0(b_{2k}), \\
b_{4k+1} &:= (\Psi'_0)^2(b_1), \\
b_{4k+2} &:= (\Psi'_0)^2(b_2), \\
b_{4k+3} &:= (\Psi'_0)^2(b_3), \\
b_{4k+4} &:= (\Psi'_0)^2(b_4), \\
&\dots,
\end{aligned}$$

where $\bar{\phi}_{2k} := \phi_{v_2}^{-1} \circ \phi_{v_3} \circ \phi_{v_4}^{-1} \circ \dots \circ \phi_{v_{2k-1}} \circ \phi_{v_{2k}}^{-1}$ and $\Psi'_0 := \bar{\phi}_{2k} \circ \phi_{v_1} = \phi_{v_2}^{-1} \circ \phi_{v_3} \circ \phi_{v_4}^{-1} \circ \dots \circ \phi_{v_{2k-1}} \circ \phi_{v_{2k}}^{-1} \circ \phi_{v_1}$.

Let $b_i^{(1)} := b_i$ ($i = 1, 2, \dots$) and let $A_1 := \{b_i^{(1)} \mid i = 1, 2, \dots\}$.

Since v_1, \dots, v_{2k} is a cycle, similarly we can define the sequences $A'_{2t} = \{b_i^{(2t)} \mid i = 1, 2, \dots\}$ and $A_{2t+1} = \{b_i^{(2t+1)} \mid i = 1, 2, \dots\}$ for $t = 0, 1, \dots, k-1$ (where if $b_i^{(2t)}$ (and $b_i^{(2t+1)}$) is not defined then we consider $b_i^{(2t)}$ (and $b_i^{(2t+1)}$) is blank). Here if $b = \square$ then $\phi_{v_{2t}}(b) = \square$ and $\phi_{v_{2t+1}}^{-1}(b) = \square$.

We say that the cycle v_1, \dots, v_{2k} is α -type, if there exists a positive number n_0 such that for any $t = 0, 1, \dots, k-1$,

$$\cdot b_i^{(2t)} \neq \square \text{ and } b_i^{(2t+1)} \neq \square \text{ for all } 1 \leq i \leq n_0,$$




FIGURE 8. A'_{2i} and A_{2i+1}

- $b'_i^{(2t)} = \square$ and $b_i^{(2t+1)} = \square$ for all $i \geq n_0 + 1$ and
- $b'_{n_0}^{(2t)} = v'_{2t}$ and $b_{n_0}^{(2t+1)} = v_{2t+1}$,

where $v'_0 = v'_{2k}$ and $A'_0 = A'_{2k}$. In the otherwise case, if there is no such a positive number n_0 , then the cycle v_1, \dots, v_{2k} is said to be β -type.

If the cycle v_1, \dots, v_{2k} is α -type then for all $t = 1, \dots, k$, $A'_{2t} = \phi_{v_{2t}}(A_{2t-1} - \{v_{2t}\}) \cup \{v'_{2t}\}$ and $A_{2t+1} = \phi_{v_{2t+1}}^{-1}(A'_{2t} - \{v'_{2t+1}\}) \cup \{v_{2t+1}\}$ hold (see Figure 8).

We also consider a cycle $v_1, v_2 \in V(\tilde{\Gamma})$ such that normal-arrows $v_1 \rightarrow v_2$ and $v_2 \rightarrow v_1$ are in $\tilde{\Gamma}$. Then similarly we define α -type and β -type for the cycle v_1, v_2 by the above corresponding sequences $\{b'_i\}$ and $\{b_i\}$.

We obtain the following proposition.

Proposition 4.2. *If for some positive number $k \in \mathbb{N}$ and some sequence $v_1, \dots, v_{2k} \in V(\tilde{\Gamma})$, there exist arrows $v_{2i-1} \rightarrow v_{2i}$ and $v_{2i} \dashrightarrow v_{2i+1}$ in $\tilde{\Gamma}$ for any $i = 1, \dots, k$ (where $v_{2k+1} := v_1$) and if the cycle v_1, \dots, v_{2k} is α -type, then there exist normal-arrows $v_1 \rightarrow v_2$ and $v_2 \rightarrow v_1$ in $\tilde{\Gamma}$.*

Proof. We consider the sequence $\{b'_i\}$ in $V(\Gamma')$ as in Definition 4.1. Here the cycle v_1, \dots, v_{2k} is α -type. Let n_0 be the positive number such that $b'_i \neq \square$ for $1 \leq i \leq n_0$ and $b'_i = \square$ for $i \geq n_0 + 1$. Let $A'_0 := \{b'_i \mid i = 1, \dots, n_0\}$. Here $b'_i \neq b'_j$ if $i \neq j$.

Then there exist the following edges in Γ' .

- (1) Since $v_1 \rightarrow v_2$ is in $\tilde{\Gamma}$, we have $[v'_1, \phi_{v_1}(v_2)] = [b'_1, b'_2] \in E(\Gamma')$.

(2) Since $v_2 \dashrightarrow v_3$ is in $\tilde{\Gamma}$, we have $[v_2, \phi_{v_2}^{-1}(v'_3)] \in E(\Gamma)$ and $[\phi_{v_1}(v_2), \phi_{v_1} \circ \phi_{v_2}^{-1}(v'_3)] = [b'_2, b'_3] \in E(\Gamma')$. (Here $\phi_{v_2}^{-1}(v'_3) \neq v_1$. Indeed, $[v_2, \phi_{v_2}^{-1}(v'_3)] \in E(\Gamma)$ and $[v_1, v_2] \notin E(\Gamma)$, since $v_1 \rightarrow v_2$ is in $\tilde{\Gamma}$.)

(3) Since $v_3 \rightarrow v_4$ is in $\tilde{\Gamma}$, we have $[v'_3, \phi_{v_3}(v_4)] \in E(\Gamma')$ and $[\phi_{v_1} \circ \phi_{v_2}^{-1}(v'_3), \phi_{v_1} \circ \phi_{v_2}^{-1} \circ \phi_{v_3}(v_4)] = [b'_3, b'_4] \in E(\Gamma')$.

(4) Since $v_4 \dashrightarrow v_5$ is in $\tilde{\Gamma}$, we have $[v_4, \phi_{v_4}^{-1}(v'_5)] \in E(\Gamma)$ and $[\phi_{v_1} \circ \phi_{v_2}^{-1} \circ \phi_{v_3}(v_4), \phi_{v_1} \circ \phi_{v_2}^{-1} \circ \phi_{v_3} \circ \phi_{v_4}^{-1}(v'_5)] = [b'_4, b'_5] \in E(\Gamma')$.

(2k-1) Since $v_{2k-1} \rightarrow v_{2k}$ is in $\tilde{\Gamma}$, we have $[v'_{2k-1}, \phi_{v_{2k-1}}(v_{2k})] \in E(\Gamma')$ and for

$$\bar{\phi}_{2k-2} := \phi_{v_1} \circ \phi_{v_2}^{-1} \circ \phi_{v_3} \circ \phi_{v_4}^{-1} \circ \cdots \circ \phi_{v_{2k-3}} \circ \phi_{v_{2k-2}}^{-1},$$

$$[\bar{\phi}_{2k-2}(v'_{2k-1}), \bar{\phi}_{2k-2} \circ \phi_{v_{2k-1}}(v_{2k})] = [b'_{2k-1}, b'_{2k}] \in E(\Gamma').$$

(2k) Since $v_{2k} \dashrightarrow v_1$ is in $\tilde{\Gamma}$, we have $[v_{2k}, \phi_{v_{2k}}^{-1}(v'_1)] \in E(\Gamma)$ and for

$$\bar{\phi}_{2k-1} := \phi_{v_1} \circ \phi_{v_2}^{-1} \circ \phi_{v_3} \circ \phi_{v_4}^{-1} \circ \cdots \circ \phi_{v_{2k-3}} \circ \phi_{v_{2k-2}}^{-1} \circ \phi_{v_{2k-1}},$$

$$[\bar{\phi}_{2k-1}(v_{2k}), \bar{\phi}_{2k-1} \circ \phi_{v_{2k}}^{-1}(v'_1)] = [b'_{2k}, b'_{2k+1}] \in E(\Gamma').$$

Thus $[b'_i, b'_{i+1}] \in E(\Gamma')$ for any $i = 1, \dots, n_0 - 1$.

Let $A_{2i-1} := \phi_{v_{2i-1}}^{-1}(A'_{2i-2} - \{v'_{2i-1}\}) \cup \{v_{2i-1}\}$ and let $A'_{2i} := \phi_{v_{2i}}(A_{2i-1} - \{v_{2i}\}) \cup \{v'_{2i}\}$ for each $i = 1, \dots, k$. Here $A'_{2k} = A'_0$. We can also denote $A_{2i-1} = \{b_1^{(2i-1)}, \dots, b_{n_0}^{(2i-1)}\}$ and $A'_{2i} = \{b_1^{(2i)}, \dots, b_{n_0}^{(2i)}\}$ for $i = 1, \dots, k$ as in Definition 4.1 (see Figure 8).

By the definition of α -type, $b_{n_0}^{(1)} = v_1$ on A_1 and $b_{n_0}^{(2)} = v'_2$ on A'_2 . Hence $[\phi_{v_2}(v_1), v'_2] \in E(\Gamma')$ on A'_2 . Also $[v_1, v_2] \notin E(\Gamma)$. Thus there exists a normal-arrow $v_2 \rightarrow v_1$ in $\tilde{\Gamma}$. (Here by the same argument, we can obtain that there exist arrows $v_{2i} \rightarrow v_{2i-1}$ and $v_{2i+1} \dashrightarrow v_{2i}$ in $\tilde{\Gamma}$ for all $i = 1, \dots, k$ where $v_{2k+1} = v_1$.) \square

From Proposition 4.2, we obtain the following.

Corollary 4.3. *Let Γ and Γ' be finite simple graphs with at least three vertices satisfying the property (*). Then for the associated directed graph $\tilde{\Gamma}$,*

- (I) *there exists a vertex $v \in V(\tilde{\Gamma})$ with no normal-arrow arising from v in $\tilde{\Gamma}$ (then $\Gamma \cong \Gamma'$ by Theorem 3.3),*
- (II) *for some $v_1, v_2 \in V(\tilde{\Gamma})$, there exist normal-arrows $v_1 \rightarrow v_2$ and $v_2 \rightarrow v_1$ in $\tilde{\Gamma}$, or*
- (III) *some and all cycles with alternate normal-arrows and dashed-arrows in $\tilde{\Gamma}$ are β -type.*

5. IN THE CASE THAT $v_1 \rightarrow v_2$ AND $v_2 \rightarrow v_1$ ARE IN $\tilde{\Gamma}$ WHOSE CYCLE IS α -TYPE

We suppose that there exist normal-arrows $v_1 \rightarrow v_2$ and $v_2 \rightarrow v_1$ in $\tilde{\Gamma}$ for some vertices $v_1, v_2 \in V(\tilde{\Gamma})$ and suppose that the cycle v_1, v_2 is α -type.

We consider the isomorphism

$$\Psi := \phi_{v_2}^{-1} \circ \phi_{v_1} : \Gamma - v_1 - \phi_{v_1}^{-1}(v'_2) \xrightarrow[\cong]{\phi_{v_1}|} \Gamma' - v'_1 - v'_2 \xrightarrow[\cong]{\phi_{v_2}^{-1}|} \Gamma - v_2 - \phi_{v_2}^{-1}(v'_1).$$

Let A , X and Y be the full-subgraphs of Γ induced by

$$\begin{aligned} V(A) &= \{a \in V(\Gamma) \mid \Psi^i(a) \in \{v_1, \phi_{v_1}^{-1}(v'_2)\} \text{ for some } i \geq 0\}, \\ V(X) &= \left\{ x \in V(\Gamma) - \{v_1, \phi_{v_1}^{-1}(v'_2)\} \mid \begin{array}{l} \Psi(x) \neq x \text{ and} \\ \Psi^i(x) \notin \{v_1, \phi_{v_1}^{-1}(v'_2)\} \text{ for any } i \geq 0 \end{array} \right\} \text{ and} \\ V(Y) &= \{y \in V(\Gamma) - \{v_1, \phi_{v_1}^{-1}(v'_2)\} \mid \Psi(y) = y\}. \end{aligned}$$

Here

$$V(\Gamma) = V(X) \cup V(Y) \cup V(A)$$

that is a disjoint union.

Let $X \cup Y$ be the full-subgraph of Γ with $V(X \cup Y) = V(X) \cup V(Y)$. Then $\Psi : X \cup Y \rightarrow X \cup Y$ is a graph-automorphism. Here X is the moving-part and Y is the fixed-part by Ψ . If X is non-empty then the graph-automorphism group G of $X \cup Y$ generated by Ψ is

$$G = \{\Psi^i \mid i \in \mathbb{Z}\} \cong \mathbb{Z}_m$$

for some number $m \geq 2$, since Γ is a finite graph.

Also we consider the isomorphism

$$\Psi' := \phi_{v_2} \circ \phi_{v_1}^{-1} : \Gamma' - v'_1 - \phi_{v_1}(v_2) \xrightarrow[\cong]{\phi_{v_1}^{-1}|} \Gamma - v_1 - v_2 \xrightarrow[\cong]{\phi_{v_2}|} \Gamma' - v'_2 - \phi_{v_2}(v_1).$$

Let A' , X' and Y' be the full-subgraphs of Γ' induced by

$$\begin{aligned} V(A') &= \{a' \in V(\Gamma') \mid (\Psi')^i(a') \in \{v'_1, \phi_{v_1}(v_2)\} \text{ for some } i \geq 0\}, \\ V(X') &= \left\{ x' \in V(\Gamma') - \{v'_1, \phi_{v_1}(v_2)\} \mid \begin{array}{l} \Psi'(x') \neq x' \text{ and} \\ (\Psi')^i(x') \notin \{v'_1, \phi_{v_1}(v_2)\} \text{ for any } i \geq 0 \end{array} \right\} \text{ and} \\ V(Y') &= \{y' \in V(\Gamma') - \{v'_1, \phi_{v_1}(v_2)\} \mid \Psi'(y') = y'\}. \end{aligned}$$

Here

$$V(\Gamma') = V(X') \cup V(Y') \cup V(A')$$

that is a disjoint union.

We consider a sequence $\{a_i\}$ in $V(\Gamma)$ as

$$\begin{aligned} a_1 &:= v_1, \\ a_2 &:= \phi_{v_1}^{-1}(v'_2), \\ a_3 &:= \phi_{v_1}^{-1} \circ \phi_{v_2}(v_1), \\ a_4 &:= \phi_{v_1}^{-1} \circ \phi_{v_2} \circ \phi_{v_1}^{-1}(v'_2), \\ a_5 &:= \phi_{v_1}^{-1} \circ \phi_{v_2} \circ \phi_{v_1}^{-1} \circ \phi_{v_2}(v_1), \\ &\dots \end{aligned}$$

Here the cycle v_1, v_2 is α -type by assumption. Let n be the positive number such that $a_i \neq \square$ for all $1 \leq i \leq n$ and $a_i = \square$ for all $i \geq n + 1$. Let $A_0 := \{a_1, a_2, \dots, a_n\}$. Here $a_i \neq a_j$ if $i \neq j$.

We also define a sequence $\{a'_i \mid i = 1, \dots, n\}$ in $V(\Gamma')$ as $a'_i := \phi_{v_1}(a_{n-i+2})$ for any $i = 2, \dots, n$ and $a'_1 := v'_1$. Let $A'_0 := \{a'_1, a'_2, \dots, a'_n\}$. Here $A'_0 = \phi_{v_1}(A_0 - \{v_1\}) \cup \{v'_1\}$.

Then by the argument in Section 4 (where $k = 1$ and $n_0 = n$), we obtain that




FIGURE 9. A_0 and A'_0

- (1) $V(A) = A_0$,
- (2) $V(A') = A'_0$,
- (3) $\phi_{v_1}(A - a_1) = A' - a'_1$ where $a_1 = v_1$ and $a'_1 = v'_1$,
- (4) $\phi_{v_1}(a_i) = a'_{n-i+2}$ for any $i = 2, \dots, n$,
- (5) $\phi_{v_2}(A - a_n) = A' - a'_n$ where $a_n = v_2$ and $a'_n = v'_2$, and
- (6) $\phi_{v_2}(a_i) = a'_{n-i}$ for any $i = 1, \dots, n-1$

(see Figure 9). Here each vertices a_i, a_{i+1} do not have to span an edge in Γ .

Then $A'_0 = \phi_{v_1}(A_0 - \{v_1\}) \cup \{v'_1\}$ and $A'_0 = \phi_{v_2}(A_0 - \{v_2\}) \cup \{v'_2\}$ both hold. Here

$$a_1 = v_1, \quad a_2 = \phi_{v_1}^{-1}(v'_2), \quad a_{n-1} = \phi_{v_2}^{-1}(v'_1), \quad a_n = v_2,$$

$$a'_1 = v'_1, \quad a'_2 = \phi_{v_1}(v_2), \quad a'_{n-1} = \phi_{v_2}(v_1) \quad \text{and} \quad a'_n = v'_2.$$

Also

- (7) $\Psi = \phi_{v_2}^{-1} \circ \phi_{v_1} : \Gamma - \{a_1, a_2\} \rightarrow \Gamma - \{a_{n-1}, a_n\}$ is an isomorphism such that $\Psi(a_i) = a_{i-2}$ for any $i = 3, \dots, n$,
- (8) $\Psi' = \phi_{v_2} \circ \phi_{v_1}^{-1} : \Gamma' - \{a'_1, a'_2\} \rightarrow \Gamma' - \{a'_{n-1}, a'_n\}$ is an isomorphism such that $\Psi'(a'_i) = a'_{i-2}$ for any $i = 3, \dots, n$,

- (9) $\Psi|_{A-\{a_1, a_2\}} : A-\{a_1, a_2\} \rightarrow A-\{a_{n-1}, a_n\}$ is an isomorphism such that $\Psi(a_i) = a_{i-2}$ for any $i = 3, \dots, n$, and
- (10) $\Psi'|_{A'-\{a'_1, a'_2\}} : A'-\{a'_1, a'_2\} \rightarrow A'-\{a'_{n-1}, a'_n\}$ is an isomorphism such that $\Psi'(a'_i) = a'_{i-2}$ for any $i = 3, \dots, n$.

We obtain the following.

Proposition 5.1. *Then $\phi_{v_i}(X \cup Y) = X' \cup Y'$, $\phi_{v_i}(X) = X'$ and $\phi_{v_i}(Y) = Y'$ for each $i = 1, 2$. Hence*

$$X \cup Y \cong X' \cup Y', \quad X \cong X' \text{ and } Y \cong Y'.$$

Their isomorphisms are obtained by the restrictions of ϕ_{v_1} and ϕ_{v_2} both.

Proof. Since $\phi_{v_1}(A - v_1) = A' - v'_1$ and $\phi_{v_2}(A - v_2) = A' - v'_2$ by (3) and (5) above, we have that $\phi_{v_i}(X \cup Y) = X' \cup Y'$ for each $i = 1, 2$.

We show that $\phi_{v_1}(X) = X'$. Let $x_0 \in V(X)$. We consider $x'_1 := \phi_{v_1}(x_0)$, $x_2 := \phi_{v_2}^{-1}(x'_1) = \Psi(x_0)$ and $x'_3 := \phi_{v_1}(x_2) = (\Psi')^{-1}(x'_1)$. Then $x_0 \neq x_2$, since $x_0 \in V(X)$. Hence $x'_1 = \phi_{v_1}(x_0) \neq \phi_{v_1}(x_2) = x'_3$. Thus $\Psi'(x'_1) \neq x'_1$ and $x'_1 \in V(X')$. This follows that $\phi_{v_1}(x_0) \in V(X')$ for all $x_0 \in V(X)$. Thus $X \cong X'$ by ϕ_{v_1} .

Also since $V(Y) = V(X \cup Y) - V(X)$ and $V(Y') = V(X' \cup Y') - V(X')$, we obtain that $\phi_{v_1}(y) \in V(Y')$ for all $y \in V(Y)$. Thus $Y \cong Y'$ by ϕ_{v_1} .

By the same argument, we can also show that $X \cong X'$ and $Y \cong Y'$ by ϕ_{v_2} . \square

By (1)–(10) above, we can denote $A = A(n; B, C)$ and $A' = A(n; C, B)$ for some subsets B and C of the set $\{1, \dots, n-1\}$. Here $V(A) = \{a_1, a_n, \dots, a_n\}$ is the numbering of vertices in the definition of $A = A(n; B, C)$. Also if $V(A') = \{b_1, b_2, \dots, b_n\}$ is the numbering of vertices in the definition of $A' = A(n; C, B)$, then $a'_i = b_{n-i+1}$ for all $i = 1, \dots, n$ and $V(A') = \{a'_1, a'_2, \dots, a'_n\}$.

Then

- (a) since $v_1 \rightarrow v_2$ is in $\tilde{\Gamma}$, we have that $[a'_1, a'_2] = [v'_1, \phi_{v_1}(v_2)] \in E(\Gamma')$ holds, and
- (b) since $v_2 \rightarrow v_1$ is in $\tilde{\Gamma}$, we have that $[a'_n, a'_{n-1}] = [v'_2, \phi_{v_2}(v_1)] \in E(\Gamma')$ holds.

Here $1 \in C$, because $A' = A(n; C, B)$ and $[a'_n, a'_{n-1}] = [b_1, b_2] \in E(A')$.

Also

- (c) $v_1 \dashrightarrow v_2$ is in $\tilde{\Gamma}$ if and only if $[a_1, a_2] = [v_1, \phi_{v_1}^{-1}(v'_2)] \in E(\Gamma)$, and
- (d) $v_2 \dashrightarrow v_1$ is in $\tilde{\Gamma}$ if and only if $[a_n, a_{n-1}] = [v_2, \phi_{v_2}^{-1}(v'_1)] \in E(\Gamma)$.

Thus we obtain the following.

Lemma 5.2. *We can denote $A = A(n; B, C)$ and $A' = A(n; C, B)$ for some subsets B and C of the set $\{1, \dots, n-1\}$. Here $1 \in C$.*

Then we show some lemmas.

Lemma 5.3. *The following statements are equivalent:*

- (1) *The sequence a_1, \dots, a_n is a Hamilton-path of A .*
- (2) *The sequence a'_1, \dots, a'_n is a Hamilton-path of A' .*
- (3) *There exist dashed-arrows $v_1 \dashrightarrow v_2$ and $v_2 \dashrightarrow v_1$ in $\tilde{\Gamma}$.*

Proof. (1) \iff (2): For $A = A(n; B, C)$ and $A' = A(n; C, B)$, the sequence a_1, \dots, a_n is a Hamilton-path of A if and only if $[a_1, a_2]$ and $[a_2, a_3]$ are in $E(A)$; that is, $1 \in B$ and $1 \in C$. This is equivalent that the sequence b_1, \dots, b_n is a Hamilton-path of $A' = A(n; C, B)$ where $a'_i = b_{n-i+1}$ for $i = 1, \dots, n$. Hence (1) and (2) are equivalent.

[(1) and (2)] \iff (3): Now a dashed-arrow $v_1 \dashrightarrow v_2$ is in $\tilde{\Gamma}$ if and only if $[a_1, a_2] \in E(A)$. Also a dashed-arrow $v_2 \dashrightarrow v_1$ is in $\tilde{\Gamma}$ if and only if $[a_n, a_{n-1}] \in E(A)$.

Thus, if (1) and (2) hold then we obtain (3) holds.

Suppose that (3) holds. Then $[a_1, a_2] \in E(A)$ by the above. Hence $1 \in B$ in $A = A(n; B, C)$. Also $1 \in C$ by Lemma 5.2. Thus (1) and (2) hold. \square

Lemma 5.4. *If n is odd, then the sequence a_1, \dots, a_n is a Hamilton-path of A , the sequence a'_1, \dots, a'_n is a Hamilton-path of A' and there exist dashed-arrows $v_1 \dashrightarrow v_2$ and $v_2 \dashrightarrow v_1$ in $\tilde{\Gamma}$.*

Proof. Suppose that n is odd. By the assumption, a normal-arrow $v_1 \rightarrow v_2$ is in $\tilde{\Gamma}$ and $[a'_1, a'_2] = [b_n, b_{n-1}] \in E(A')$. Then since n is odd, $[a'_{n-1}, a'_{n-2}] = [b_2, b_3] \in E(A')$. Hence $1 \in B$, because $A' = A(n; C, B)$. Thus, $[a_1, a_2] \in E(A)$. Also $1 \in C$ and $[a_2, a_3] \in E(A)$ by the assumption. Hence the sequence a_1, \dots, a_n is a Hamilton-path of A and there exist dashed-arrows $v_1 \dashrightarrow v_2$ and $v_2 \dashrightarrow v_1$ in $\tilde{\Gamma}$ by Lemma 5.3. \square

We investigate some properties of $A(n; B, C)$ and $A(n; C, B)$ in general case.

Definition 5.5. For $b, c \in \{1, \dots, n-1\}$, let $A_1(b)$ and $A_2(c)$ be the simple graphs as $V(A_1(b)) = V(A_2(c)) = V(A)$,

$$E(A_1(b)) = \{ [a_{2t-1}, a_{2t-1+b}] \mid t \in \mathbb{N} \text{ as } 1 \leq 2t-1 < 2t-1+b \leq n \} \text{ and}$$

$$E(A_2(c)) = \{ [a_{2t}, a_{2t+c}] \mid t \in \mathbb{N} \text{ as } 2 \leq 2t < 2t+c \leq n \}.$$

Then $A_1(b)$ is a subgraph of $A = A(n; B, C)$ if and only if $b \in B$. Also $A_2(c)$ is a subgraph of $A = A(n; B, C)$ if and only if $c \in C$. Hence $E(A) = \bigcup_{b \in B} E(A_1(b)) \cup \bigcup_{c \in C} E(A_2(c))$.

Definition 5.6. Let $A := A(n; B, C)$ where $n \in \mathbb{N}$ is a number at least 3 and B and C are subsets of the set $\{1, \dots, n-1\}$. Then we define the numbers $\beta(A)$, $\beta'(A)$, $\gamma(A)$ and $\gamma'(A)$ as follows;

$$\begin{aligned} \beta(A) &:= \#\{b \in B \mid n-b \text{ is odd}\}, \\ \beta'(A) &:= |B| - \beta(A), \\ \gamma(A) &:= \#\{c \in C \mid n-c \text{ is odd}\} \text{ and} \\ \gamma'(A) &:= |C| - \gamma(A). \end{aligned}$$

Here we note that $[n-b \text{ is odd}]$ if and only if $[[b \text{ is even if } n \text{ is odd}] \text{ and } [b \text{ is odd if } n \text{ is even}]]$.

Then we have the following.

Lemma 5.7. *Let $A = A(n; B, C)$, $A' = A(n; C, B)$, $V(A) = \{a_1, a_n, \dots, a_n\}$ and $V(A') = \{a'_1, a'_2, \dots, a'_n\}$ where $V(A) = \{a_1, a_n, \dots, a_n\}$ and $V(A') = \{b_1, b_2, \dots, b_n\}$ are the numbering of vertices as in the definition of $A = A(n; B, C)$ and $A' =$*

$A(n; C, B)$. Let $a'_i := b_{n-i+1}$ for $i = 1, \dots, n$. (Hence $V(A') = \{a'_1, a'_2, \dots, a'_n\}$.) Then the following two statements are equivalent:

- (i) $\beta(A) = \gamma(A)$.
- (ii) $|E(A)| = |E(A')|$, $\deg_A a_1 = \deg_{A'} a'_1$ and $\deg_A a_n = \deg_{A'} a'_n$.

Proof. Let $b \in \{1, \dots, n-1\}$. In the case that n is even, $n-b$ is odd if and only if b is odd. Also in the case that n is odd, $n-b$ is odd if and only if b is even.

Here $n-b$ is odd if and only if for each $j = 1, 2$, $A_j(b)$ is symmetric and there exists the automorphism $\tau : A_j(b) \rightarrow A_j(b)$ as $\tau(a_i) = a_{n-i+1}$ for any $i = 1, \dots, n$. Also $n-b$ is even if and only if for each $j = 1, 2$, $A_j(b) - a_1$ and $A_j(b) - a_n$ are symmetric and there exist the automorphisms $\tau' : A_j(b) - a_1 \rightarrow A_j(b) - a_1$ and $\tau'' : A_j(b) - a_n \rightarrow A_j(b) - a_n$ such that $\tau'(a_i) = a_{n-i+2}$ for any $i = 2, \dots, n$ and $\tau''(a_i) = a_{n-i}$ for any $i = 1, \dots, n-1$ (see Figure 3).

Here if $n-b$ is even then $|E(A_1(b))| = |E(A_2(b))|$ holds. Also if $n-b$ is odd then $|E(A_1(b))| = |E(A_2(b))| + 1$ holds. Thus, for $A = A(n; B, C)$ and $A' = A(n; C, B)$, $|E(A)| = |E(A')|$ if and only if $\beta(A) = \beta(A') = \gamma(A)$. Here $\beta(A') = \gamma(A)$ since $A = A(n; B, C)$ and $A' = A(n; C, B)$.

We also note that $\deg_A a_1 = |B|$, $\deg_A a_n = \beta(A) + \gamma'(A)$, $\deg_{A'} b_1 = \deg_{A'} a'_n = |C|$ and $\deg_{A'} b_n = \deg_{A'} a'_1 = \beta(A') + \gamma'(A') = \gamma(A) + \beta'(A)$. Then

$$\begin{aligned} \deg_A a_1 = \deg_{A'} a'_1 &\iff |B| = \gamma(A) + \beta'(A) \\ &\iff |B| - \beta'(A) = \gamma(A) \\ &\iff \beta(A) = \gamma(A). \end{aligned}$$

Also

$$\begin{aligned} \deg_A a_n = \deg_{A'} a'_n &\iff \beta(A) + \gamma'(A) = |C| \\ &\iff \beta(A) = |C| - \gamma'(A) \\ &\iff \beta(A) = \gamma(A). \end{aligned}$$

□

From the above argument, we obtain the following theorem.

Theorem 5.8. *Let Γ and Γ' be finite simple graphs with at least three vertices satisfying the property (*). If in the associated directed graph $\tilde{\Gamma}$,*

(II) *for some $v_1, v_2 \in V(\tilde{\Gamma})$, there exist normal-arrows $v_1 \rightarrow v_2$ and $v_2 \rightarrow v_1$ in $\tilde{\Gamma}$ and if the cycle v_1, v_2 is α -type, then the graphs Γ and Γ' have the following structure (\mathcal{F}):*

- (\mathcal{F}) (1) *There exists a positive number $n \geq 3$.*
- (2) *There exists a full-subgraph A of Γ with $V(A) = \{a_1, a_2, \dots, a_{n-1}, a_n\}$ and $|V(A)| = n$.*
- (3) *There exists a full-subgraph A' of Γ' with $V(A') = \{a'_1, a'_2, \dots, a'_{n-1}, a'_n\}$ and $|V(A')| = n$.*
- (4) *$a_1 = v_1$, $a_n = v_2$, $a'_1 = v'_1$ and $a'_n = v'_2$, where $v'_1 := f(v_1)$ and $v'_2 := f(v_2)$.*
- (5) *$A = A(n; B, C)$ and $A' = A(n; C, B)$ for some subsets B and C of the set $\{1, \dots, n-1\}$.*

- (6) $1 \in C$ and $\beta(A) = \gamma(A)$ hold.
- (7) If n is odd or if there exist dashed-arrows $v_1 \dashrightarrow v_2$ and $v_2 \dashrightarrow v_1$ in $\tilde{\Gamma}$, then the sequence a_1, \dots, a_n is a Hamilton-path of A , the sequence a'_1, \dots, a'_n is a Hamilton-path of A' and $1 \in B$.
- (8) There exists an isomorphism $\Psi : \Gamma - \{a_1, a_2\} \rightarrow \Gamma - \{a_{n-1}, a_n\}$ such that $\Psi(a_i) = a_{i-2}$ for any $i = 3, \dots, n$.
- (9) There exists an isomorphism $\Psi' : \Gamma' - \{a'_1, a'_2\} \rightarrow \Gamma' - \{a'_{n-1}, a'_n\}$ such that $\Psi'(a'_i) = a'_{i-2}$ for any $i = 3, \dots, n$.
- (10) The full-subgraph $X \cup Y$ of Γ with $V(X \cup Y) = V(\Gamma) - V(A)$ is Ψ -invariant.
- (11) The full-subgraph $X' \cup Y'$ of Γ' with $V(X' \cup Y') = V(\Gamma') - V(A')$ is Ψ' -invariant.
- (12) $X \cup Y$ is isomorphic to $X' \cup Y'$ by $\phi_{v_1}|_{X \cup Y}$ and $\phi_{v_2}|_{X \cup Y}$ both.
- (13) The full-subgraph X of Γ with $V(X) = \{x \in V(X \cup Y) \mid \Psi(x) \neq x\}$ is isomorphic to the full-subgraph X' of Γ' with $V(X') = \{x' \in V(X' \cup Y') \mid \Psi'(x') \neq x'\}$ by $\phi_{v_1}|_X$ and $\phi_{v_2}|_X$ both. (Here X and X' are the moving-parts by Ψ and Ψ' respectively.)
- (14) The full-subgraph Y of Γ with $V(Y) = \{y \in V(X \cup Y) \mid \Psi(y) = y\}$ is isomorphic to the full-subgraph Y' of Γ' with $V(Y') = \{y' \in V(X' \cup Y') \mid \Psi'(y') = y'\}$ by $\phi_{v_1}|_Y$ and $\phi_{v_2}|_Y$ both. (Here Y and Y' are the fixed-parts by Ψ and Ψ' respectively.)
- (15) $\phi_{v_1}(a_i) = a'_{n-i+2}$ for any $i = 2, \dots, n$.
- (16) $\phi_{v_2}(a_i) = a'_{n-i}$ for any $i = 1, \dots, n-1$.

6. EXAMPLES AND REMARKS ON $A = A(n; B, C)$ AND $A' = A(n; C, B)$

We give some examples and remarks on $A = A(n; B, C)$ and $A' = A(n; C, B)$.

Let $A = A(n; B, C)$ and $A' = A(n; C, B)$ for a number $n \in \mathbb{N}$ at least 3 and subsets B and C of the set $\{1, \dots, n-1\}$. Here $V(A) = \{a_1, a_n, \dots, a_n\}$ is the numbering of vertices in the definition of $A = A(n; B, C)$. Also if $V(A') = \{b_1, b_2, \dots, b_n\}$ is the numbering of vertices in the definition of $A' = A(n; C, B)$, then $a'_i := b_{n-i+1}$ for all $i = 1, \dots, n$ and $V(A') = \{a'_1, a'_2, \dots, a'_n\}$.

We first show a proposition.

Proposition 6.1. *Let*

$$B_0 := \{b \in B \mid n - b \text{ is odd}\} \text{ and}$$

$$C_0 := \{c \in C \mid n - c \text{ is odd}\}.$$

Then $B_0 = C_0$ if and only if the following all statements hold.

- (1) $A - a_1 = A - v_1$ is symmetric and there exists the automorphism $\tau_0 : A - a_1 \rightarrow A - a_1$ such that $\tau_0(a_i) = a_{n-i+2}$ for any $i = 2, \dots, n$.
- (2) $A - a_n = A - v_2$ is symmetric and there exists the automorphism $\tau_1 : A - a_n \rightarrow A - a_n$ such that $\tau_1(a_i) = a_{n-i}$ for any $i = 1, \dots, n-1$.
- (3) There exists the isomorphism $\bar{\varphi} : A \rightarrow A'$ such that $\bar{\varphi}(a_i) = a'_i$ for any $i = 1, \dots, n$. Hence $A \cong A'$.
- (4) The isomorphisms $\varphi_0 := \phi_{v_1} \circ \tau_0 : A - v_1 \rightarrow A' - v'_1$ and $\varphi_1 := \phi_{v_2} \circ \tau_1 : A - v_2 \rightarrow A' - v'_2$ both extend to the isomorphism $\bar{\varphi} : A \rightarrow A'$ in (3).

Proof. Suppose that $B_0 = C_0$.

Then for any $b \in B_0$, since $b \in C_0$, the graphs $A_1(b)$ and $A_2(b)$ (in Definition 5.5) both are subgraphs of A . Let $A_1(b) \cup A_2(b)$ be the simple graph as $V(A_1(b) \cup A_2(b)) = V(A)$ and $E(A_1(b) \cup A_2(b)) = E(A_1(b)) \cup E(A_2(b))$. Then $A_1(b) \cup A_2(b)$, $A_1(b) \cup A_2(b) - a_1$ and $A_1(b) \cup A_2(b) - a_n$ are symmetric in the above sense.

Also by the argument in the proof of Lemma 5.7, for any $b \in B - B_0$ ($n - b$ is even) and for any $j = 1, 2$, $A_j(b) - a_1$ and $A_j(b) - a_n$ are symmetric in the above sense. Hence the statements (1) and (2) hold.

Now we show that the statements (3) and (4) hold. For any $i = 2, \dots, n$,

$$\varphi_0(a_i) = \phi_{v_1}(\tau_0(a_i)) = \phi_{v_1}(a_{n-i+2}) = a'_i = \bar{\varphi}(a_i).$$

Also for any $i = 1, \dots, n - 1$,

$$\varphi_1(a_i) = \phi_{v_2}(\tau_1(a_i)) = \phi_{v_2}(a_{n-i}) = a'_i = \bar{\varphi}(a_i).$$

Here $a_1 = v_1$ and $a_n = v_2$ (and $a'_1 = v'_1$ and $a'_n = v'_2$) do not span an edge of A (and A' respectively).

Hence for any $i, j \in \{1, \dots, n\}$, $[a_i, a_j] \in E(A)$ if and only if $[a'_i, a'_j] \in E(A')$, because φ_0 and φ_1 are isomorphisms. Thus $\bar{\varphi} : A \rightarrow A'$ as $\bar{\varphi}(a_i) = a'_i$ for $i = 1, \dots, n$ is an isomorphism and the statements (3) and (4) hold.

Conversely we show that if the statements (1) and (2) hold then $B_0 = C_0$. Suppose that $B_0 \neq C_0$. Then there exists $b_0 \in B_0 - C_0$ or $c_0 \in C_0 - B_0$. If there is $b_0 \in B_0 - C_0$, then $A_1(b_0)$ is a subgraph of A and $A_2(b_0)$ is not a subgraph of A . If there is $c_0 \in C_0 - B_0$, then $A_2(c_0)$ is a subgraph of A and $A_1(c_0)$ is not a subgraph of A . Hence $A - a_1$ and $A - a_n$ are not symmetric in the above sense. Thus (1) and (2) do not hold. \square

We remark that any $A = A(n; B, C)$ and $A' = A(n; C, B)$ with $1 \in C$ and $B_0 = C_0$ as in Proposition 6.1 are isomorphic and they can be realized as the full-subgraphs A and A' of Γ and Γ' respectively in the structure (\mathcal{F}) . On the other hand, if $B_0 \neq C_0$ then in general there is a possibility that $A = A(n; B, C)$ and $A' = A(n; C, B)$ can not become finite simple graphs satisfying the condition (*). We introduce this in Example 6.6. Also in the case that $B_0 \neq C_0$, even if $A = A(n; B, C)$ and $A' = A(n; C, B)$ are finite simple graphs satisfying the property (*), it is not obvious that $A \cong A'$. This can be seen from Examples 6.2–6.5.

Example 6.2. We consider $A = A(8; \{3\}, \{1\})$ and $A' = A(8; \{1\}, \{3\})$ (see Figure 10). Here $n = 8$ is even. Then A and A' have two connected components and they are isomorphic.




FIGURE 10. Example 6.2

In this example, since $A \cong A'$, we can construct a bijective map $f : V(A) \rightarrow V(A')$ and an isomorphism $\phi_v : A - v \rightarrow A' - f(v)$ for any $v \in V(A)$. Here for $v_1 = a_1$, $v_2 = a_8$, $v'_1 = f(v_1) = a'_1$ and $v'_2 = f(v_2) = a'_8$, we can take the isomorphisms $\phi_{v_1} : A - v_1 \rightarrow A' - v'_1$ and $\phi_{v_2} : A - v_2 \rightarrow A' - v'_2$ as (14) and (15) in Theorem 5.8 and for $\Gamma := A$ and $\Gamma' := A'$, there exist normal-arrows $v_1 \rightarrow v_2$ and $v_2 \rightarrow v_1$ in $\tilde{\Gamma}$. Then there do not exist dashed-arrows $v_1 \dashrightarrow v_2$ and $v_2 \dashrightarrow v_1$ in $\tilde{\Gamma}$, since the sequence a_1, a_2, \dots, a_8 is not a Hamilton-path of A .

Example 6.3. We consider $A = A(8; \{1, 3\}, \{1, 5\})$ and $A' = A(8; \{1, 5\}, \{1, 3\})$ (see Figure 11). Here $n = 8$ is even. Then A and A' are isomorphic as in Figure 11.




FIGURE 11. Example 6.3

In this example, since $A \cong A'$, we can construct a bijective map $f : V(A) \rightarrow V(A')$ and an isomorphism $\phi_v : A - v \rightarrow A' - f(v)$ for any $v \in V(A)$ such that for $v_1 = a_1$, $v_2 = a_8$, $v'_1 = f(v_1) = a'_1$, $v'_2 = f(v_2) = a'_8$ and for $\Gamma := A$ and $\Gamma' := A'$, there exist arrows $v_1 \rightarrow v_2$, $v_2 \rightarrow v_1$, $v_1 \dashrightarrow v_2$ and $v_2 \dashrightarrow v_1$ in $\tilde{\Gamma}$.

Example 6.4. We consider $A = A(8; \{3, 4\}, \{1\})$ and $A' = A(8; \{1\}, \{3, 4\})$ (see Figure 12). Here $n = 8$ is even and there are an even number and an odd number both in $B = \{3, 4\}$. Then A and A' have two connected components and they are isomorphic as in Figure 12.

In this example, since $A \cong A'$, we can construct a bijective map $f : V(A) \rightarrow V(A')$ and an isomorphism $\phi_v : A - v \rightarrow A' - f(v)$ for any $v \in V(A)$ such that for $v_1 = a_1$, $v_2 = a_8$, $v'_1 = f(v_1) = a'_1$, $v'_2 = f(v_2) = a'_8$ and for $\Gamma := A$ and $\Gamma' := A'$, there exist normal-arrows $v_1 \rightarrow v_2$ and $v_2 \rightarrow v_1$ in $\tilde{\Gamma}$. Here there do not exist dashed-arrows $v_1 \dashrightarrow v_2$ and $v_2 \dashrightarrow v_1$ in $\tilde{\Gamma}$.

Example 6.5. We consider $A = A(7; \{1, 2, 3\}, \{1, 4\})$ and $A' = A(7; \{1, 4\}, \{1, 2, 3\})$ (see Figure 13). Here $n = 7$ is odd and there are an even number and an odd number both in $B - C = \{2, 3\}$. Then A and A' are isomorphic as in Figure 13.




FIGURE 12. Example 6.4




FIGURE 13. Example 6.5

In this example, since $A \cong A'$, we can construct a bijective map $f : V(A) \rightarrow V(A')$ and an isomorphism $\phi_v : A - v \rightarrow A' - v'$ for any $v \in V(A)$ such that for $v_1 = a_1$, $v_2 = a_7$, $v'_1 = f(v_1) = a'_1$, $v'_2 = f(v_2) = a'_7$ and for $\Gamma := A$ and $\Gamma' := A'$, there exist arrows $v_1 \rightarrow v_2$, $v_2 \rightarrow v_1$, $v_1 \dashrightarrow v_2$ and $v_2 \dashrightarrow v_1$ in $\tilde{\Gamma}$.

Example 6.6. We consider $A = A(8; \{1, 3, 4\}, \{1, 5\})$ and $A' = A(8; \{1, 5\}, \{1, 3, 4\})$ (see Figure 14). Here $\beta(A) = \gamma(A)$ holds. Also we note that there exists a bijective map $g : V(A) \rightarrow V(A')$ such that $\deg_A v = \deg_{A'} g(v)$ for all $v \in V(A)$.




FIGURE 14. Example 6.6

Then A and A' are not isomorphic. Indeed, there is a triangle a'_3, a'_7, a'_8 in A' such that $\deg_{A'} a'_3 = 4$, $\deg_{A'} a'_7 = 4$ and $\deg_{A'} a'_8 = 2$. On the other hand, there is not such a triangle in A . In the graph A , a_8 is the unique vertex with degree 2 and a_5 and a_7 are the vertices connecting to a_8 with degree 4. Here a_5 and a_7 do not span an edge in A .

It is known that every simple graph Γ with at least 3 vertices and at most 11 vertices is reconstructible [11]. Hence the two graphs $A = A(8; \{1, 3, 4\}, \{1, 5\})$ and $A' = A(8; \{1, 5\}, \{1, 3, 4\})$ can not satisfy the property (*).

By Example 6.6, in general, there is a possibility that $A = A(n; B, C)$ and $A' = A(n; C, B)$ (still if they satisfy $\beta(A) = \gamma(A)$ and still if there is a bijective map $g : V(A) \rightarrow V(A')$ such that $\deg_A v = \deg_{A'} g(v)$ for all $v \in V(A)$) can not become finite simple graphs satisfying the property (*).

Here the following problem arises.

Problem. The full-subgraphs $A = A(n; B, C)$ and $A' = A(n; C, B)$ of Γ and Γ' respectively in Theorem 5.8 will be isomorphic?

7. EXAMPLES AND REMARKS ON β -TYPE CYCLES

We give an example and a remark on β -type cycles.

Example 7.1. We consider two graphs Γ and Γ' as in Figure 15. Here Γ and Γ' are isomorphic. We define isomorphisms $\phi_{v_1} : \Gamma - v_1 \rightarrow \Gamma' - v'_1$ and $\phi_{v_2} : \Gamma - v_2 \rightarrow \Gamma' - v'_2$ as in Figure 16.

We consider a sequence $\{b'_i\}$ in $V(\Gamma)$ as

$$\begin{aligned} b'_1 &:= v'_1, \\ b'_2 &:= \phi_{v_1}(v_2), \\ b'_3 &:= \phi_{v_1} \circ \phi_{v_2}^{-1}(v'_1), \\ b'_4 &:= \phi_{v_1} \circ \phi_{v_2}^{-1} \circ \phi_{v_1}(v_2), \\ b'_5 &:= \phi_{v_1} \circ \phi_{v_2}^{-1} \circ \phi_{v_1} \circ \phi_{v_2}^{-1}(v'_1), \\ &\dots \end{aligned}$$

Here b'_4, b'_6 and b'_i ($i \geq 8$) are blanks.




FIGURE 15. Example 7.1

Also consider a sequence $\{b_i\}$ in $V(\Gamma)$ as

$$\begin{aligned}
b_1 &:= v_2, \\
b_2 &:= \phi_{v_2}^{-1}(v'_1), \\
b_3 &:= \phi_{v_2}^{-1} \circ \phi_{v_1}(v_2), \\
b_4 &:= \phi_{v_2}^{-1} \circ \phi_{v_1} \circ \phi_{v_2}^{-1}(v'_1), \\
b_5 &:= \phi_{v_2}^{-1} \circ \phi_{v_1} \circ \phi_{v_2}^{-1} \circ \phi_{v_1}(v_2), \\
&\dots
\end{aligned}$$

Here b_5, b_7 and b_i ($i \geq 8$) are blanks.

Then $[v'_1, \phi_{v_1}(v_2)] = [b'_1, b'_2] \in E(\Gamma')$, $[v'_2, \phi_{v_2}(v_1)] = [b'_7, b'_2] \in E(\Gamma')$, $[v_1, \phi_{v_1}^{-1}(v'_2)] = [b_3, b_6] \notin E(\Gamma)$ and $[v_2, \phi_{v_2}^{-1}(v'_1)] = [b_1, b_2] \in E(\Gamma)$. Hence there exist arrows $v_1 \rightarrow v_2$, $v_2 \rightarrow v_1$ and $v_2 \dashrightarrow v_1$ in $\tilde{\Gamma}$. The cycle v_1, v_2 is β -type.

The isomorphism

$$\phi_{v_2} \circ \phi_{v_1}^{-1} : \Gamma' - \{b'_1, b'_2\} \xrightarrow[\cong]{\phi_{v_1}^{-1}|} \Gamma - \{v_1, v_2\} \xrightarrow[\cong]{\phi_{v_2}|} \Gamma' - \{b'_2, b'_7\}$$

is as in Figure 16, where $b'_1 = v'_1$, $b'_2 = \phi_{v_1}(v_2) = \phi_{v_2}(v_1)$ and $b'_7 = v'_2$.

Remark. We define a subgraph $\bar{A}(n, p; B, C)$ of $A(n; B, C)$.

Let $n \in \mathbb{N}$ be a number at least 3, let $p \in \mathbb{N}$ with $p < n$ and let B and C be subsets of the set $\{1, \dots, n-1\}$. Let $\{a_1, \dots, a_n\}$ be the vertex set of $A(n; B, C)$ whose numbering is as in the definition of $A(n; B, C)$. Let $P := \{a_i \mid i = p + 2t \ (0 \leq t \in \mathbb{Z})\}$.

Then we define $\bar{A}(n, p; B, C)$ as the full-subgraph of $A(n; B, C)$ whose vertex set is $\{a_1, \dots, a_n\} - P$; that is, a_i is blank if $a_i \in P$.




FIGURE 16. Isomorphisms ϕ_{v_1} and ϕ_{v_2}

In Example 7.1, the graphs Γ and Γ' with the vertex sets $\{b_1, b_2, b_3, b_4, b_5, b_6, b_7\}$ and $\{b'_1, b'_2, b'_3, b'_4, b'_5, b'_6, b'_7\}$ respectively can be denoted by $\Gamma = \overline{A}(7, 5; \{1, 5\}, \{1, 4\})$ and $\Gamma' = \overline{A}(7, 4; \{1, 4\}, \{1, 5\})$.

REFERENCES

- [1] M. Bilinski, Y.S. Kwon and X. Yu, On the reconstruction of planar graphs, *J. Combin. Theory Ser. B* 97 (2007), 745–756.
- [2] J.A. Bondy and R.L. Hemminger, Graph reconstruction—a survey, *J. Graph Theory* 1 (1977), 227–268.
- [3] S. Fiorini and J. Lauri, The reconstruction of maximal planar graphs. I. Recognition, *J. Combin. Theory Ser. B* 30 (1981), 188–195.
- [4] T. Hosaka, Reconstructible graphs, simplicial flag complexes of homology manifolds and associated right-angled Coxeter groups, *Osaka J. Math.* 52 (2015), 1173–1180.
- [5] T. Hosaka and Y. Xiao, A new class of reconstructible graphs from some neighbourhood conditions, *Hokkaido Math. J.* 44 (2015), 327–340.
- [6] P.J. Kelly, On isomorphic transformations, PhD thesis, University of Wisconsin, 1942.
- [7] P.J. Kelly, A congruence theorem for trees, *Pacific J. Math.* 7 (1957), 961–968.
- [8] J. Lauri, The reconstruction of maximal planar graphs. II. Reconstruction, *J. Combin. Theory Ser. B* 30 (1981), 196–214.
- [9] B. Manvel, On reconstruction of graphs, Ph.D. thesis, University of Michigan, 1988.
- [10] B.D. McKay, Computer reconstruction of small graphs, *J. Graph Theory* 1 (1977), 281–283.
- [11] B.D. McKay, Small graphs are reconstructible, *Australas. J. Combin.* 15 (1997), 123–126.
- [12] A. Nijenhuis, Note on the unique determination of graphs by proper subgraphs, *Notices Amer. Math. Soc.* 24 (1977), A-290.

- [13] S.M. Ulam, A Collection of Mathematical Problems, Wiley (Interscience), New York, 1960, p. 29.
- [14] Y. Yongzhi, The reconstruction conjecture is true if all 2-connected graphs are reconstructible, J. Graph theory 12 (1988), 237–243.

Email address: `hosaka.tetsuya@shizuoka.ac.jp`

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, SHIZUOKA UNIVERSITY, 836 OHYA,
SURUGA-KU, SHIZUOKA, 422-8529, JAPAN