The long-time behaviour of a stochastic SIR epidemic model with distributed delay and multidimensional Lévy jumps

Driss Kiouach^{*} and Yassine Sabbar

LPAIS Laboratory, Faculty of Sciences Dhar El Mahraz, Sidi Mohamed Ben Abdellah University, Fez, Morocco.

Abstract

arXiv:2003.08219v2 [math.DS] 23 Jan 2021

Recently, emerging epidemics like COVID-19 and its variants require predictive mathematical models to implement suitable responses in order to limit their negative and profound impact on society. The SIR (Susceptible-Infected-Removed) system is a straightforward mathematical formulation to model the dissemination of many infectious diseases. The present paper reports novel theoretical and analytical results for a perturbed version of an SIR model with Gamma-distributed delay. Notably, our epidemic model is represented by Itô-Lévy stochastic differential equations in order to simulate sudden and unexpected external phenomena. By using some new and ameliorated mathematical approaches, we study the long-run characteristics of the perturbed delayed model. Within this scope, we give sufficient conditions for two interesting asymptotic proprieties: extinction and persistence of the epidemic. One of the most interesting results is that the dynamics of the stochastic model are closely related to the intensities of white noises and Lévy jumps, which can give us a good insight into the evolution of the epidemic in some unexpected situations. Our work complements the results of some previous investigations and provides a new approach to predict and analyze the dynamic behavior of epidemics with distributed delay. For illustrative purposes, numerical examples are presented for checking the theoretical study.

Keywords: Distributed delay; Epidemic model; White noise; Lévy jumps; Extinction; Persistence in the mean. Mathematics Subject Classification 2020: 60H10; 34A12; 34A26; 37C10; 60H30; 92D30.

1. Introduction

Epidemic models under various formal frameworks are an auxiliary tool to acquire information about the dynamics of epidemic transmission and the impact of different intervention strategies [17]. Recently, the employment of these models to generate long-term epidemic forecasts is increased with the rising number of emerging and re-emerging epidemic outbreaks. For example, COVID-19 and its new variant identified in England present an urgent and serious challenge due to their contagious nature and frequently changing characteristics [20, 46, 19, 9, 2]. A mathematical model for measuring and preventing the continued spread of COVID-19 is firmly required to understand the mechanisms of its dynamic and predict its future. Considerable attention has been paid to the analysis of susceptibleinfectious-removed (SIR) type model, which is proposed to describe the dissemination of COVID-19 before suggesting more adapted and complex epidemic models. In the SIR model, we often assume that recovered individuals can get continuous immunity [3]. Many studies have paid close attention to the characteristics of the long-term epidemics immune response [12, 31, 37, 50]. To confer the realistic aspect of the epidemic model and make it biologically reasonable, numerous scholars considered the SIR epidemic model with time delay because an individual may not be infectious until some time after becoming infected [23, 3]. In the above-mentioned works, the time delay is assumed to be single-valued. The constant delay may be considered if the variation of the time is known exactly, which is not real for many biological reasons [28]. Considering the variable infectivity in the time interval yields a model with a distributed delay [4]. Therefore, it is more realistic to introduce a continuously distributed delay in the biological modeling [45, 42]. Analyzing the characteristics of the SIR model with a distributed time delay still a rich subject that may deliver new comprehension of the epidemics propagation which motivates our work. According to the approach of Muroya et al. [32], we consider the delay kernel $\mathcal{G}: [0,\infty) \to [0,\infty)$ as a normalized L¹-function, i.e, $\int_0^\infty \mathcal{G}(s) ds = 1$.

*Corresponding author.

E-mail addresses: d.kiouach@uiz.ac.ma (D. Kiouach), yassine.sabbar@usmba.ac.ma (Y. Sabbar).

The average delay for the kernel \mathcal{G} can be presented by the following quantity $\int_0^\infty s\mathcal{G}(s)ds < \infty$. Hence, the incidence rate at time τ can be presented as the following form: $\beta S(\tau) \int_{-\infty}^{\tau} \mathcal{G}(\tau - s)I(s)ds$, where β denotes the transmission rate, S(t) and I(t) represent the fractions of susceptible and infective individuals at time t. The SIR epidemic model with distributed delay can be expressed as follows [26]:

$$\begin{cases} dS(t) = \left(A - \mu_1 S(t) - \beta S(t) \int_{-\infty}^t \mathcal{G}(t-s) I(s) ds\right) dt, \\ dI(t) = \left(\beta S(t) \int_{-\infty}^t \mathcal{G}(t-s) I(s) ds - (\mu_2 + \gamma) I(t)\right) dt, \\ dR(t) = \left(\gamma I(t) - \mu_3 R(t)\right) dt, \end{cases}$$
(1)

where R(t) is the fraction number of recovered populations at time t. The remaining parameters appearing in this system are described as follows:

- A is the recruitment rate of susceptible individuals corresponding to births and immigration.
- μ_1 , μ_3 are the natural death rates associated respectively to the susceptible and recovered populations, μ_2 is a general mortality rate including the effect of the disease fatality.
- γ is the rate of individuals leaving I to R (recovered rate).

The threshold number of the deterministic system (1) is $\mathcal{T}^{\star} = \frac{\beta A}{\mu_1(\mu_2 + \gamma)}$ which determines the persistence $(\mathcal{T}^{\star} > 1)$ or the extinction $(\mathcal{T}^{\star} < 1)$ of the epidemic. Many studies showed that the deterministic epidemic model (1) is suitable to describe the transmission process of some known epidemics such as Rubella, Whooping cough, Measles and Smallpox. Due to many biological and mathematical considerations [26], in this paper, we consider the delay kernel with Gamma distribution $\mathcal{G}(s) = \frac{s^n \eta^{n+1} e^{-\eta s}}{n!}$, $s \in (0, \infty)$, where the constant $\eta > 0$ is the rate of exponential fading memory, which means the retrogradation of the past memories effect. In this paper, we consider the low kernel function \mathcal{G} with n = 0. By letting $\mathcal{D}(t) = \int_{-\infty}^{t} \eta e^{-\eta(t-s)} I(s) ds$ and using the linear chain approach, system (1) can be transformed into the following equivalent system:

$$\begin{cases} dS(t) = (A - \mu_1 S(t) - \beta S(t) \mathcal{D}(t)) dt, \\ dI(t) = (\beta S(t) \mathcal{D}(t) - (\mu_2 + \gamma) I(t)) dt, \\ dR(t) = (\gamma I(t) - \mu_3 R(t)) dt, \\ d\mathcal{D}(t) = \eta (I(t) - \mathcal{D}(t)) dt. \end{cases}$$

$$(2)$$

Although the use of deterministic models can explain and simulate some phenomena in real life, such models do not actually consider the effect of the natural stochasticity, and we plainly wish to learn how randomness affects our epidemic models [49, 52, 51, 39, 1, 33, 36]. Generally, one of the ordinary extensions from the deterministic SIR model to the stochastic version is to incorporate environmental white noises, which appear from an almost continuous series of small variations on the model parameters [6, 58, 56, 40, 41, 34, 38, 30]. Therefore, the stochastic delayed SIR epidemic can be an accurate tool to predict the long-run dynamics of infectious epidemics [25, 27, 47, 16, 14, 15, 24, 18, 13]. In [26], the authors inserted the stochastic perturbation in the model (1) by assuming that the white noise is directly proportional to the variable S and they obtained the following stochastic system:

$$\begin{cases} dS(t) = \left(A - \mu_1 S(t) - \beta S(t) \mathcal{D}(t)\right) dt + \sigma S(t) d\mathcal{W}(t), \\ dI(t) = \left(\beta S(t) \mathcal{D}(t) - (\mu_2 + \gamma) I(t)\right) dt, \\ dR(t) = \left(\gamma I(t) - \mu_3 R(t)\right) dt, \\ d\mathcal{D}(t) = \eta \left(I(t) - \mathcal{D}(t)\right) dt. \end{cases}$$

$$(3)$$

where W(t) is a standard Brownian motion with associated intensity $\sigma > 0$. Specifically, they proved the existence and uniqueness of an ergodic stationary distribution to the model (3). Then, they established sufficient conditions for the extinction of a disease that spreads according to this model. On the basis of these findings, one question was catches our attention. It is possible to develop and generalize the stochastic model proposed in [26]?. So, the objective of this work is to expound on this problem and provide a suitable analytical context. Specifically, we aim to describe the strong fluctuations by considering a general version of the dynamical model (3). It is clear that the population systems may suffer certain sudden environmental catastrophes, such as earthquakes, floods, droughts, etc [54, 55, 57]. For example, the recent massive explosion in the port city of Beirut. The impact of this unexpected disaster has been extremely devastating, especially when it has occurred simultaneously with the COVID-19 pandemic. This led to a sudden worsening of the health situation and a jump increase in the number of deaths. Mathematically, we use the Lévy process to describe the phenomena that cause a big jump to occur occasionally [11, 43, 44, 53, 61]. By considering this type of random perturbations, the model (2) becomes the following system of stochastic differential equations with Lévy jumps (SDE-Js for short):

$$\begin{cases} dS(t) = \left(A - \mu_1 S(t) - \beta S(t) \mathcal{D}(t)\right) dt + \sigma_1 S(t) d\mathcal{W}_1(t) + \int_{\mathcal{U}} \lambda_1(u) S(t^-) \widetilde{\mathcal{N}}(dt, du), \\ dI(t) = \left(\beta S(t) \mathcal{D}(t) - (\mu_2 + \gamma) I(t)\right) dt + \sigma_2 I(t) d\mathcal{W}_2(t) + \int_{\mathcal{U}} \lambda_2(u) I(t^-) \widetilde{\mathcal{N}}(dt, du), \\ dR(t) = \left(\gamma I(t) - \mu_3 R(t)\right) dt + \sigma_3 R(t) d\mathcal{W}_3(t) + \int_{\mathcal{U}} \lambda_3(u) R(t^-) \widetilde{\mathcal{N}}(dt, du), \\ d\mathcal{D}(t) = \eta \left(I(t) - \mathcal{D}(t)\right) dt + \sigma_4 \mathcal{D}(t) d\mathcal{W}_4(t) + \int_{\mathcal{U}} \lambda_4(u) \mathcal{D}(t^-) \widetilde{\mathcal{N}}(dt, du), \end{cases}$$
(4)

where $S(t^-)$, $I(t^-)$, $R(t^-)$ and $\mathcal{D}(t^-)$ are the left limits of S(t), I(t), R(t) and $\mathcal{D}(t)$, respectively. $\mathcal{W}_i(t)$ (i = 1, 2, 3, 4)are independent Brownian motions and $\sigma_i > 0$ (i = 1, 2, 3, 4) are their intensities. \mathcal{N} is a Poisson counting measure with compensating martingale $\tilde{\mathcal{N}}$ and characteristic measure ν on a measurable subset \mathcal{U} of $(0, \infty)$ satisfying $\nu(\mathcal{U}) < \infty$. $\mathcal{W}_i(t)$ (i = 1, 2, 3, 4) are independent of \mathcal{N} . We assumed that ν is a Lévy measure such that $\tilde{\mathcal{N}}(dt, du) = \mathcal{N}(dt, du) - \nu(du)dt$ and we suppose that the function $\lambda_i : Z \times \Omega \to \mathbb{R}$ is bounded and continuous. Throughout this paper, we let $(\Omega, \mathcal{F}, \mathbb{P})$ denotes a complete probability space with a filtration $\{\mathcal{F}_t\}_{t\geq 0}$ satisfying these conditions: right continuous and \mathcal{F}_0 contains all \mathbb{P} -null sets. We also assume that $\mathcal{W}_i(t)$ is defined on this probability space. Since the compartment R(t) does not appear in the equations of S(t) I(t) and $\mathcal{D}(t)$, it is sufficient to analyze the dynamic behavior of the following SDE-J model:

$$\begin{cases} dS(t) = \left(A - \mu_1 S(t) - \beta S(t) \mathcal{D}(t)\right) dt + \sigma_1 S(t) d\mathcal{W}_1(t) + \int_{\mathcal{U}} \lambda_1(u) S(t^-) \widetilde{\mathcal{N}}(dt, du), \\ dI(t) = \left(\beta S(t) \mathcal{D}(t) - (\mu_2 + \gamma) I(t)\right) dt + \sigma_2 I(t) d\mathcal{W}_2(t) + \int_{\mathcal{U}} \lambda_2(u) I(t^-) \widetilde{\mathcal{N}}(dt, du), \\ d\mathcal{D}(t) = \eta \left(I(t) - \mathcal{D}(t)\right) dt + \sigma_4 \mathcal{D}(t) d\mathcal{W}_4(t) + \int_{\mathcal{U}} \lambda_4(u) \mathcal{D}(t^-) \widetilde{\mathcal{N}}(dt, du). \end{cases}$$
(5)

In this study, we develop a new analysis to deal with stochastic models with jumps in epidemiology. Our main goal is to investigate sufficient conditions of the stochastic extinction and persistence in the mean. These two important properties are sufficient to predict and analyze the dynamics of a given epidemic. We apply a new approach to estimate the values of the averages $t^{-1} \int_0^t \psi(s) ds$ and $t^{-1} \int_0^t \psi^2(s) ds$, where $\psi(t)$ is the positive solution of the following subsystem:

$$\begin{cases} \mathrm{d}\psi(t) = \left(A - \mu_1\psi(t)\right)\mathrm{d}t + \sigma_1\psi(t)\mathrm{d}\mathcal{W}_1(t) + \int_{\mathcal{U}}\lambda_1(u)\psi(t^-)\tilde{\mathcal{N}}(\mathrm{d}t,\mathrm{d}u), \quad \forall t > 0, \\ \psi(0) = S(0) > 0. \end{cases}$$
(6)

Our approach allows us to close the gap left by using the classical method presented for example in [60]. Furthermore, we give an optimal sufficient condition for the stochastic extinction. For the purpose of well understanding the dynamics of the delayed model (5), we give a sufficient condition of the disease persistence. The analysis in this paper seems to be promising to investigate other related stochastic delayed models with Lévy noises in epidemiology and even in biology.

This work is organized as follows. In section 2, we verify the well-posedness of the stochastic model (5). In section 3, we give sufficient conditions for the extinction and the persistence in the mean of the disease. Finally, in section 4, numerical simulations are carried out to confirm the theoretical study.

2. Existence and uniqueness of the global positive solution

To study the long-term properties of an infectious disease system, the prime concern is whether the solution is unique, positive and global in time. In this short section, motivated by the approach presented in [21], we show the

well-posedness of the stochastic model (5). Mainly, the primary key to treat the said problem is to construct a suitable Lyapunov function. According to some analytical and mathematical reasons, it is necessary that we make the following two hypotheses:

- (\mathcal{H}_1): We assume that the jump coefficients $\lambda_i(u)$ in (5) satisfy $\int_{\mathcal{H}} \lambda_i^2(u)\nu(\mathrm{d}u) < \infty$ (i = 1, 2, 4).
- (\mathcal{H}_2): For all i = 1, 2, 4, we assume that $1 + \lambda_i(u) > 0$ and $\int_{\mathcal{U}} \left(\lambda_i(u) \ln\left(1 + \lambda_i(u)\right)\right) \nu(\mathrm{d}u) < \infty$.

By assumption (\mathcal{H}_1) , the coefficients of the system (5) are locally Lipschitz continuous, then for any initial value $(S(0), I(0), \mathcal{D}(0)) \in \mathbb{R}^3_+$ there is a unique local solution $(S(t), I(t), \mathcal{D}(t))$ on $t \in [0, \tau_e)$, herein, τ_e represents the explosion time. In the following theorem, our goal is to show that the solution is positive and global.

Theorem 2.1. Let hypotheses (\mathcal{H}_1) and (\mathcal{H}_2) hold. For any initial value $(S(0), I(0), \mathcal{D}(0)) \in \mathbb{R}^3_+$, there exists a unique positive solution $(S(t), I(t), \mathcal{D}(t))$ of the SDE-J (5) on $t \ge 0$ and the solution will remain in \mathbb{R}^3_+ with probability one, namely $(S(t), I(t), \mathcal{D}(t)) \in \mathbb{R}^3_+$ for all $t \ge 0$ almost surely (a.s.).

Proof. We only need to prove that $\tau_e = \infty$ almost surely. Let $\epsilon_0 > 0$ be sufficiently large such that each component of $(S(0), I(0), \mathcal{D}(0))$ all lies in the interval $\left(\frac{1}{\epsilon_0}, \epsilon_0\right)$. For each integer $\epsilon \ge \epsilon_0$, we define the following stopping time

$$\tau_{\epsilon} = \inf\left\{ t \in [0, \tau_{e}) | \ S(t) \notin \left(\frac{1}{\epsilon}, \epsilon\right), \text{ or } I(t) \notin \left(\frac{1}{\epsilon}, \epsilon\right), \text{ or } \mathcal{D}(t) \notin \left(\frac{1}{\epsilon}, \epsilon\right) \right\}$$

Set inf $\emptyset = \infty$ (\emptyset denotes the empty set) and let $\tau_{\infty} = \lim_{\epsilon \to \infty} \tau_{\epsilon}$. Evidently, τ_{ϵ} is increasing as $\epsilon \to \infty$. Moreover, $\tau_{\infty} \leq \tau_{\epsilon}$. If we can prove that $\tau_{\infty} = \infty$ a.s., then $\tau_{e} = \infty$ and $(S(t), I(t), \mathcal{D}(t)) \in \mathbb{R}^{3}_{+}$ for all $t \geq 0$ almost surely. Specifically, we need to show that $\tau_{\infty} = \infty$ a.s. By supposing the opposite, we can consider a pair of positive constants T > 0 and $k \in (0, 1)$ such that $\mathbb{P}\{\tau_{\infty} \leq T\} > k$. Hence, there exists an integer $\epsilon_{1} \geq \epsilon_{0}$ such that

$$\mathbb{P}\{\tau_{\epsilon} \le T\} \ge k \quad \text{for all} \quad \epsilon > \epsilon_1. \tag{7}$$

Construct a \mathcal{C}^2 -function $\mathcal{V}: \mathbb{R}^3_+ \to [0, +\infty)$ by

$$\mathcal{V}(S, I, \mathcal{D}) = \left(S - m - m\ln\frac{S}{m}\right) + (I - 1 - \ln I) + \frac{\mu_2 + \gamma}{\eta}(\mathcal{D} - 1 - \ln \mathcal{D})$$

where m > 0 is a positive constant to be determined later. Obviously, this function is non-negative which can be seen from $x - 1 - \ln x \ge 0$ for all x > 0. According to the general Itô's formula [8], we obtain for all $0 \le t < \tau_{\epsilon}$,

$$d\mathcal{V}(S, I, \mathcal{D}) = \mathcal{L}\mathcal{V}(S, I, \mathcal{D})dt + \left(1 - \frac{m}{S}\right)\sigma_1 S d\mathcal{W}_1(t) + \left(1 - \frac{1}{I}\right)\sigma_2 I d\mathcal{W}_2(t) + \frac{\mu_2 + \gamma}{\eta}\left(1 - \frac{1}{\mathcal{D}}\right)\sigma_4 \mathcal{D} d\mathcal{W}_4(t) + \int_{\mathcal{U}} \lambda_1(u)S - m\ln(1 + \lambda_1(u))\widetilde{\mathcal{N}}(dt, du) + \int_{\mathcal{U}} \lambda_2(u)I - \ln(1 + \lambda_2(u))\widetilde{\mathcal{N}}(dt, du) + \int_{\mathcal{U}} \frac{\mu_2 + \gamma}{\eta} \left(\lambda_4(u)\mathcal{D} - \ln(1 + \lambda_4(u))\right)\widetilde{\mathcal{N}}(dt, du),$$

where,

$$\begin{aligned} \mathcal{LV}(S, I, \mathcal{D}) &= A - \mu_1 S - \frac{mA}{S} + m\beta \mathcal{D} + m\mu_1 - (\mu_2 + \gamma)I - \frac{\beta S\mathcal{D}}{I} \\ &+ (\mu_2 + \gamma) + (\mu_2 + \gamma)I - (\mu_2 + \gamma)\mathcal{D} - \frac{(\mu_2 + \gamma)I}{\mathcal{D}} + (\mu_2 + \gamma) \\ &+ \frac{m\sigma_1^2}{2} + \frac{\sigma_2^2}{2} + \frac{\mu_2 + \gamma}{\eta} \frac{\sigma_4^2}{2} + \int_{\mathcal{U}} m\lambda_1(u) - m\ln(1 + \lambda_1(u))\nu(\mathrm{d}u) \\ &+ \int_{\mathcal{U}} \lambda_2(u) - \ln(1 + \lambda_2(u))\nu(\mathrm{d}u) + \int_{\mathcal{U}} \frac{\mu_2 + \gamma}{\eta} \Big(\lambda_4(u) - \ln(1 + \lambda_4(u))\Big)\nu(\mathrm{d}u) \end{aligned}$$

Then

$$\mathcal{LV}(S, I, \mathcal{D}) \leq A + 2(\mu_2 + \gamma) + m\mu_1 + \left(m\beta - (\mu_2 + \gamma)\right)\mathcal{D} + \frac{m\sigma_1^2}{2} + \frac{\sigma_2^2}{2} + \frac{\mu_2 + \gamma}{\eta} \frac{\sigma_4^2}{2} + \int_{\mathcal{U}} m\lambda_1(u) - m\ln(1 + \lambda_1(u))\nu(\mathrm{d}u) + \int_{\mathcal{U}} \lambda_2(u) - \ln(1 + \lambda_2(u))\nu(\mathrm{d}u) + \int_{\mathcal{U}} \frac{\mu_2 + \gamma}{\eta} \left(\lambda_4(u) - \ln(1 + \lambda_4(u))\right)\nu(\mathrm{d}u).$$

Given the fact that $x - \ln(1 + x) \ge 0$ for all x > 1 and the hypothesis (\mathcal{H}_2) , we define

$$\mathcal{J}_{1} \equiv \int_{\mathcal{U}} m\lambda_{1}(u) - m\ln(1+\lambda_{1}(u))\nu(\mathrm{d}u) + \int_{\mathcal{U}} \lambda_{2}(u) - \ln(1+\lambda_{2}(u))\nu(\mathrm{d}tu) + \int_{\mathcal{U}} \frac{\mu_{2}+\gamma}{\eta} \Big(\lambda_{4}(u) - \ln(1+\lambda_{4}(u))\Big)\nu(\mathrm{d}u).$$

To simplify, we choose $m = \frac{\mu_2 + \gamma}{\beta}$. Then, we obtain

$$\mathcal{LV}(S, I, \mathcal{D}) \le A + 2(\mu_2 + \gamma) + m\mu_1 + \frac{m\sigma_1^2}{2} + \frac{\sigma_2^2}{2} + \frac{\mu_2 + \gamma}{\eta} \frac{\sigma_4^2}{2} + \mathcal{J}_1 \equiv \mathcal{J}_2$$

The proof of the remainder is similar to the proof of Theorem 2 in [21], so we omitted it.

3. Conditions of stochastic extinction and permanence in the mean of the epidemic

In mathematical epidemiology, we are generally interested in two things, the first is to know when the epidemic will die out, and the second is when it will continue and persist. In this section, we will try our best to find sufficient conditions for these two interesting asymptotic proprieties in terms of model parameters and intensities of noises. For the sake of notational simplicity, we define

- $\bar{\sigma} \triangleq \max\{\sigma_1^2, \sigma_2^2, \sigma_4^2\}$ and $\vartheta \triangleq \min\{\mu_1, \mu_2 + \gamma \eta, \eta\}.$
- $\bar{\lambda}(u) \triangleq \max\{\lambda_1(u), \lambda_2(u), \lambda_4(u)\}$ and $\underline{\lambda}(u) \triangleq \min\{\lambda_1(u), \lambda_2(u), \lambda_4(u)\}.$

•
$$\hat{\zeta}_p(u) \triangleq \left[1 + \bar{\lambda}(u)\right]^p - 1 - p\bar{\lambda}(u)$$
 and $\check{\zeta}_p(u) \triangleq \left[1 + \underline{\lambda}(u)\right]^p - 1 - p\underline{\lambda}(u).$

•
$$\xi(u) \triangleq \max \{\hat{\zeta}_p(u), \check{\zeta}_p(u)\}$$
 and $\ell_p \triangleq \int_{\mathcal{U}} \xi(u)\nu(\mathrm{d}u)$

To properly study the long-term of the perturbed model (5), we have the following additionally hypotheses on the jump-diffusion coefficients:

• (\mathcal{H}_3) : For i = 1, 2, 4, we assume that $\int_{\mathcal{U}} \left[\ln(1 + \lambda_i(u)) \right]^2 \nu(\mathrm{d}u) < \infty$.

• (
$$\mathcal{H}_4$$
): For $i = 1, 2, 4$, we assume that $\int_{\mathcal{U}} \left[\left(1 + \bar{\lambda}(u) \right)^2 - 1 \right]^2 \nu(\mathrm{d}u) < \infty$.

• (\mathcal{H}_5): We suppose that there exists some real number p > 2 such that $\chi_{1,p} = \vartheta - \frac{(p-1)}{2}\bar{\sigma} - \frac{1}{p}\ell_p > 0$.

For the convenience of discussion in the stochastic model (5), we introduce two lemmas which will be used in our analysis.

Lemma 3.1 ([22]). We assume that the conditions (\mathcal{H}_4) and (\mathcal{H}_5) hold. Let $(S(t), I(t), \mathcal{D}(t))$ be the positive solution of the system (5) with any given initial condition $(S(0), I(0), \mathcal{D}(0)) \in \mathbb{R}^3_+$. Let also $\psi(t) \in \mathbb{R}_+$ be the solution of the equation (6) with any given initial value $\psi(0) = S(0) \in \mathbb{R}_+$. Then

• $\lim_{t \to \infty} t^{-1} \psi(t) = 0$, $\lim_{t \to \infty} t^{-1} \psi^2(t) = 0$, $\lim_{t \to \infty} t^{-1} S(t) = 0$, $\lim_{t \to \infty} t^{-1} I(t) = 0$, and $\lim_{t \to \infty} t^{-1} \mathcal{D}(t) = 0$ a.s.

$$\begin{array}{l} & \lim_{t \to \infty} t^{-1} \int_{0}^{t} \psi(s) \mathrm{d}\mathcal{W}_{1}(s) = 0, \quad \lim_{t \to \infty} t^{-1} \int_{0}^{t} \psi^{2}(s) \mathrm{d}\mathcal{W}_{1}(s) = 0, \quad \lim_{t \to \infty} t^{-1} \int_{0}^{t} S(s) \mathrm{d}\mathcal{W}_{1}(s) = 0, \\ & \lim_{t \to \infty} t^{-1} \int_{0}^{t} I(s) \mathrm{d}\mathcal{W}_{2}(s) = 0, \quad and \quad \lim_{t \to \infty} t^{-1} \int_{0}^{t} \mathcal{D}(s) \mathrm{d}\mathcal{W}_{4}(s) = 0 \quad a.s. \end{array}$$

$$\begin{array}{l} & \lim_{t \to \infty} t^{-1} \int_{0}^{t} \int_{\mathcal{U}} \lambda_{1}(u) \psi(s^{-}) \widetilde{\mathcal{N}}(\mathrm{d}s, \mathrm{d}u) = 0, \quad \lim_{t \to \infty} t^{-1} \int_{0}^{t} \int_{\mathcal{U}} \left((1 + \lambda_{1}(u))^{2} - 1 \right) \psi^{2}(s^{-}) \widetilde{\mathcal{N}}(\mathrm{d}s, \mathrm{d}u) = 0 \\ & \lim_{t \to \infty} t^{-1} \int_{0}^{t} \int_{\mathcal{U}} \lambda_{1}(u) S(s^{-}) \widetilde{\mathcal{N}}(\mathrm{d}s, \mathrm{d}u) = 0, \quad \lim_{t \to \infty} t^{-1} \int_{0}^{t} \int_{\mathcal{U}} \lambda_{2}(u) I(s^{-}) \widetilde{\mathcal{N}}(\mathrm{d}s, \mathrm{d}u) = 0, \\ & and \quad \lim_{t \to \infty} t^{-1} \int_{0}^{t} \int_{\mathcal{U}} \lambda_{4}(u) \mathcal{D}(s^{-}) \widetilde{\mathcal{N}}(\mathrm{d}s, \mathrm{d}u) = 0 \quad a.s. \end{array}$$

Remark 3.2. By using the same approach adopted in Lemma 2.5 of [22], we can easily prove the last result. Note that the hypothesis (\mathcal{H}_5) is an ameliorated version of it corresponding hypothesis frequently used in many previous works, for example, [8, 7, 10]. Therefore, the adoption of $\chi_{1,p}$ in our paper raises the optimality of our calculus and results.

Remark 3.3. In the absence of Lévy jumps (see for example [48]), the stationary distribution expression is used to estimate the time averages of the auxiliary process solution by employing the ergodic theorem [29]. Unluckily, the said expression is still unknown in the case of the Lévy noise. This issue is implicitly mentioned in [60, 59] as an open question, and the authors presented the threshold analysis of their model with an unknown stationary distribution formula. In this article, we propose an alternative method to establish the exact expression of the threshold parameter without having recourse to the use of ergodic theorem. This new idea that we propose is presented in the following lemma.

Lemma 3.4. Assume that the conditions (\mathcal{H}_4) and (\mathcal{H}_5) hold. Let $\psi(t)$ be the solution of (6) with an initial value $\psi(0) \in \mathbb{R}_+$. Then $\chi_2 = 2\mu_1 - \sigma_1^2 - \int_{\mathcal{U}} \lambda_1^2(u)\nu(\mathrm{d}u) > 0$, and

•
$$\lim_{t \to \infty} t^{-1} \int_0^t \psi(s) ds = \frac{A}{\mu_1} \quad a.s.$$

•
$$\lim_{t \to \infty} t^{-1} \int_0^t \psi^2(s) ds = \frac{2A^2}{\mu_1 \chi_2} \quad a.s.$$

Proof. Integrating from 0 to t on both sides of (6) yields

$$\frac{\psi(t) - \psi(0)}{t} = A - \frac{\mu_1}{t} \int_0^t \psi(s) \mathrm{d}s + \frac{\sigma_1}{t} \int_0^t \psi(s) \mathrm{d}\mathcal{W}_1(s) + t^{-1} \int_0^t \int_Z \lambda_1(u)\psi(s^-)\widetilde{\mathcal{N}}(\mathrm{d}s,\mathrm{d}u).$$

Clearly, we can derive that

$$t^{-1} \int_0^t \psi(s) \mathrm{d}s = \frac{A}{\mu_1} - \frac{\psi(t) - \psi(0)}{\mu_1 t} + \frac{\sigma_1}{\mu_1 t} \int_0^t \psi(s) \mathrm{d}\mathcal{W}_1(s) + \frac{1}{\mu_1 t} \int_0^t \int_Z \lambda_1(u) \psi(s^-) \widetilde{\mathcal{N}}(\mathrm{d}s, \mathrm{d}u).$$

By Lemma 3.1, we get

$$\lim_{t \to \infty} t^{-1} \int_0^t \psi(s) \mathrm{d}s = \frac{A}{\mu_1} \text{ a.s.}$$

Now, applying the generalized Itô's formula to model (6) leads to

$$d\psi^{2}(t) = \left(2\psi(t)\left(A - \mu_{1}\psi(t)\right) + \sigma_{1}^{2}\psi^{2}(t) + \int_{\mathcal{U}}\psi^{2}(t)\left((1 + \lambda_{1}(u))^{2} - 1 - 2\lambda_{1}(u)\right)\nu(\mathrm{d}u)\right)\mathrm{d}t + 2\sigma_{1}\psi^{2}(t)\mathrm{d}\mathcal{W}_{1}(t) + \int_{\mathcal{U}}\psi^{2}(t^{-})\left((1 + \lambda_{1}(u))^{2} - 1\right)\widetilde{\mathcal{N}}(\mathrm{d}t, \mathrm{d}u).$$

Integrating both sides of the last expression from 0 to t and then dividing by t, yields

$$\frac{\psi^2(t) - \psi^2(0)}{t} = 2A \times \frac{1}{t} \int_0^t \psi(s) ds - \overbrace{\left(2\mu_1 - \sigma_1^2 - \int_{\mathcal{U}} \lambda_1^2(u)\nu(du)\right)}^{\chi^2} \times \frac{1}{t} \int_0^t \psi^2(s) ds + 2\sigma_1 \times \frac{1}{t} \int_0^t \psi^2(s) d\mathcal{W}_1(s) + \frac{1}{t} \int_0^t \int_Z \psi^2(s^-) \left((1 + \lambda_1(u))^2 - 1\right) \widetilde{\mathcal{N}}(ds, du).$$

Therefore

$$\frac{\chi_2}{t} \int_0^t \psi^2(s) \mathrm{d}s = \frac{2A}{t} \int_0^t \psi(s) \mathrm{d}s + \frac{\left(\psi^2(0) - \psi^2(t)\right)}{t} + \frac{2\sigma_1}{t} \int_0^t \psi^2(s) \mathrm{d}\mathcal{W}_1(s) \tag{8}$$

$$+\frac{1}{t}\int_0^t \int_{\mathcal{U}} \psi^2(s^-) \Big((1+\lambda_1(u))^2 - 1\Big) \widetilde{\mathcal{N}}(\mathrm{d} s, \mathrm{d} u).$$
(9)

Clearly, $\chi_2 \neq 0$, because if it is not the case, we will obtain by letting t go to infinity in (8) $\frac{2A^2}{\mu} = 0$, which is obviously impossible. So, and using Lemma 3.1, we can easily verify that

$$\lim_{t \to \infty} t^{-1} \int_0^t \psi^2(s) ds = \frac{2A^2}{\mu_1 \chi_2} \text{ a.s.}$$

and since $t^{-1} \int_0^t \psi^2(s) ds > 0$ for all t > 0, we can conclude also that $\frac{2A^2}{\mu_1 \chi_2} > 0$ and then $\chi_2 > 0$. Hence the proof is completed.

We are now in the position to state and prove the main results of this paper. In the following, we always presume that the hypotheses (\mathcal{H}_1) - (\mathcal{H}_5) hold.

3.1. Stochastic extinction of the epidemic

In this subsection, we give a sufficient condition for the stochastic extinction of the disease in the system (5), but before stating the main result, we shall first recall the concept of the stochastic extinction.

Definition 3.5. Let $\mathcal{X}(t)$ be a stochastic process that describes the evolution of an infectious disease under a host population.

- The disease is said to be exponentially extinct if $\limsup_{t\to\infty} t^{-1} \ln \mathcal{X}(t) < 0$ a.s.
- The disease is said to be stochastically extinct, or extinctive, if $\lim_{t\to\infty} \mathcal{X}(t) = 0$ a.s.

Remark 3.6. Obviously, it can be seen from the above definitions that the exponential extinction implies the extinction.

For brevity and simplicity in writing the next result, we adopt the following notations:

•
$$\Upsilon \triangleq \min\{\mu_2 + \gamma, \eta\}(\sqrt{\mathcal{T}^{\star}} - 1)\mathbb{1}_{\{\mathcal{T}^{\star} \le 1\}} + \max\{\mu_2 + \gamma, \eta\}(\sqrt{\mathcal{T}^{\star}} - 1)\mathbb{1}_{\{\mathcal{T}^{\star} > 1\}}$$

•
$$\Lambda \triangleq \sigma_1^2 + \int_{\mathcal{U}} \lambda_1^2(u) \nu(\mathrm{d}u)$$
 and $\Sigma \triangleq \left(2\left(\sigma_2^{-2} + \sigma_4^{-2}\right)\right)^{-1}$.

- $\bar{\aleph}(u) \triangleq \left(\ln \left(1 + \lambda_2(u) \land \lambda_4(u) \right) \lambda_2(u) \land \lambda_4(u) \right) \times \mathbb{1}_{\{\lambda_2(u) \land \lambda_4(u) > 0\}}.$
- $\underline{\aleph}(u) \triangleq \left(\ln \left(1 + \lambda_2(u) \lor \lambda_4(u) \right) \lambda_2(u) \lor \lambda_4(u) \right) \times \mathbb{1}_{\{\lambda_2(u) \lor \lambda_4(u) \le 0\}}.$

•
$$\aleph(u) \triangleq \overline{\aleph}(u) + \underline{\aleph}(u)$$
 and $\Pi \triangleq \int_{\mathcal{U}} \aleph(u)\nu(\mathrm{d}u).$

• $\Theta \triangleq \Upsilon + \Pi - \Sigma + \eta \left(\frac{\mathcal{T}^*\Lambda}{\chi_2}\right)^{\frac{1}{2}}.$

• For any vector $v \in \mathbb{R}^n$, we denote its transpose by v^T .

Theorem 3.7. Let us denote by $(S(t), I(t), \mathcal{D}(t))$ the solution of the stochastic system (5) that starts from a given initial data $(S(0), I(0), \mathcal{D}(0)) \in \mathbb{R}^3_+$. Under the hypotheses (\mathcal{H}_1) - (\mathcal{H}_5) , we have

$$\limsup_{t \to \infty} t^{-1} \ln \left(\frac{1}{\mu_2 + \gamma} I(t) + \frac{\sqrt{\mathcal{T}^{\star}}}{\eta} \mathcal{D}(t) \right) \le \Theta \quad a.s$$

Notably, if $\Theta < 0$, then the epidemic will go to zero exponentially with probability one. Consequently,

$$\lim_{t \to \infty} I(t) = 0 \quad and \quad \lim_{t \to \infty} \mathcal{D}(t) = 0 \quad a.s.$$

Proof. Our proof starts with the use of Theorem 1.4 in [5] to establish that there is a left eigenvector of the following matrix $\beta A = \beta A$

$$\mathfrak{M}_0 = \left(\begin{array}{cc} 0 & \frac{\beta A}{\mu_1(\mu_2 + \gamma)} \\ 1 & 0 \end{array}\right)$$

corresponding to $\sqrt{\mathcal{T}^{\star}}$. This vector will be denoted by $(e_1, e_2) = (1, \sqrt{\mathcal{T}^{\star}})$. Then, $\sqrt{\mathcal{T}^{\star}}(e_1, e_2) = (e_1, e_2)\mathfrak{M}_0$. On the other hand, we define a \mathcal{C}^2 -function $\mathcal{M} : \mathbb{R}^2_+ \to \mathbb{R}_+$ by

$$\mathcal{M}(I(t), \mathcal{D}(t)) = \omega_1 I(t) + \omega_2 \mathcal{D}(t),$$

where $\omega_1 = \frac{e_1}{\mu_2 + \gamma}$ and $\omega_2 = \frac{e_2}{\eta}$. By applying the generalized Itô's formula with Lévy jumps we obtain

$$d\ln \mathcal{M}(I(t), \mathcal{D}(t)) = \mathcal{L} \ln \mathcal{M}(I(t), \mathcal{D}(t)) dt + \frac{1}{\omega_1 I(t) + \omega_2 D(t)} \Big\{ \omega_1 \sigma_2 I(t) d\mathcal{W}_2(t) + \omega_2 \sigma_4 D(t) d\mathcal{W}_3(t) \Big\} \\ + \int_{\mathcal{U}} \ln \left(1 + \frac{\omega_1 \lambda_2(u) I(t) + \omega_2 \lambda_4(u) \mathcal{D}(t)}{\omega_1 I(t) + \omega_2 D(t)} \right) \widetilde{\mathcal{N}}(dt, du),$$

where

$$\mathcal{L}\ln\mathcal{M}(I(t),\mathcal{D}(t)) = \frac{1}{\omega_1 I(t) + \omega_2 D(t)} \Big\{ \omega_1 \Big(\beta S(t) \mathcal{D}(t) - (\mu_2 + \gamma) I(t) \Big) + \omega_2 \eta \Big(I(t) - \mathcal{D}(t) \Big) \Big\} \\ - \frac{1}{2(\omega_1 I(t) + \omega_2 D(t))^2} \Big\{ \omega_1^2 \sigma_2^2 I^2(t) + \omega_2^2 \sigma_4^2 D^2(t) \Big\} \\ + \int_{\mathcal{U}} \left[\ln\left(1 + \frac{\omega_1 \lambda_2(u) I(t) + \omega_2 \lambda_4(u) \mathcal{D}(t)}{\omega_1 I(t) + \omega_2 D(t)} \right) - \frac{\omega_1 \lambda_2(u) I(t) + \omega_2 \lambda_4(u) \mathcal{D}(t)}{\omega_1 I(t) + \omega_2 D(t)} \right] \nu(du).$$

Moreover, it is easy to show the following inequality

$$\left(\frac{1}{\sigma_2^2} + \frac{1}{\sigma_4^2}\right) \times \left(\omega_1^2 \sigma_2^2 I^2(t) + \omega_2^2 \sigma_4^2 \mathcal{D}^2(t)\right) \ge \left(\frac{1}{\sigma_2} \omega_1 \sigma_2 I(t) + \frac{1}{\sigma_4} \omega_2 \sigma_4 \mathcal{D}(t)\right)^2.$$

In order to find an optimal and good majorization, we adopt the fact that

$$\int_{\mathcal{U}} \left[\ln \left(1 + \frac{\omega_1 \lambda_2(u) I(t) + \omega_2 \lambda_4(u) \mathcal{D}(t)}{\omega_1 I(t) + \omega_2 D(t)} \right) - \frac{\omega_1 \lambda_2(u) I(t) + \omega_2 \lambda_4(u) \mathcal{D}(t)}{\omega_1 I(t) + \omega_2 D(t)} \right] \nu(\mathrm{d}u) \le \Pi.$$
(10)

By using the last two results, we get

$$\mathcal{L}\ln\mathcal{M}(I(t),\mathcal{D}(t)) \leq \frac{\omega_1\beta\mathcal{D}(t)}{\omega_1I(t)+\omega_2D(t)} \Big(S(t) - \frac{A}{\mu_1}\Big) + \Pi - \Sigma \\ + \frac{1}{\omega_1I(t)+\omega_2D(t)} \Bigg\{\omega_1\Big(\frac{\beta A}{\mu_1}\mathcal{D}(t) - (\mu_2 + \gamma)I(t)\Big) + \omega_2\eta\Big(I(t) - \mathcal{D}(t)\Big)\Bigg\}.$$

By the stochastic comparison theorem, we have

$$\mathcal{L}\ln\mathcal{M}(I(t),\mathcal{D}(t)) \leq \frac{\omega_1\beta\mathcal{D}(t)}{\omega_1I(t)+\omega_2D(t)} \Big(\psi(t) - \frac{A}{\mu_1}\Big) + \Pi - \Sigma \\ + \frac{1}{\omega_1I(t)+\omega_2D(t)} \Bigg\{ \frac{e_1}{\mu_2+\gamma} \left(\frac{\beta A}{\mu_1}\mathcal{D}(t) - (\mu_2+\gamma)I(t)\right) + \frac{e_2}{\eta} \Big(\eta I(t) - \eta \mathcal{D}(t)\Big) \Bigg\}.$$

Then, we obtain that

$$\begin{aligned} \mathcal{L}\ln\mathcal{M}(I(t),\mathcal{D}(t)) &\leq \frac{\omega_1\beta}{\omega_2} \Big| \psi(t) - \frac{A}{\mu_1} \Big| + \Pi - \Sigma + \frac{1}{\omega_1 I(t) + \omega_2 D(t)} (e_1, e_2) \Big(\mathfrak{M}_0(I(t), \mathcal{D}(t))^T - (I(t), \mathcal{D}(t))^T \Big) \\ &= \frac{\omega_1\beta}{\omega_2} \Big| \psi(t) - \frac{A}{\mu_1} \Big| + \Pi - \Sigma + \frac{1}{\omega_1 I(t) + \omega_2 D(t)} \Big(\sqrt{\mathcal{T}^{\star}} - 1 \Big) \Big(e_1 I(t) + e_2 D(t) \Big) \\ &= \frac{\omega_1\beta}{\omega_2} \Big| \psi(t) - \frac{A}{\mu_1} \Big| + \Pi - \Sigma + \frac{1}{\omega_1 I(t) + \omega_2 D(t)} \Big(\sqrt{\mathcal{T}^{\star}} - 1 \Big) \Big(\omega_1(\mu_2 + \gamma) I(t) + \eta \omega_2 D(t) \Big) \\ &\leq \Upsilon + \Pi - \Sigma + \frac{\omega_1\beta}{\omega_2} \Big| \psi(t) - \frac{A}{\mu_1} \Big|. \end{aligned}$$

Hence, we deduce that

$$d\ln \mathcal{M}(I(t), \mathcal{D}(t)) \leq \left(\Upsilon + \Pi - \Sigma\right) dt + \frac{\omega_1 \beta}{\omega_2} \left| \psi(t) - \frac{A}{\mu_1} \right| dt + \frac{1}{\omega_1 I(t) + \omega_2 D(t)} \left\{ \omega_1 \sigma_2 I(t) d\mathcal{W}_2(t) + \omega_2 \sigma_4 D(t) d\mathcal{W}_4(t) \right\} + \int_{\mathcal{U}} \ln \left(1 + \lambda(u)\right) \widetilde{\mathcal{N}}(dt, du),$$

where $\lambda(u) = \max{\{\lambda_2(u), \lambda_4(u)\}}$. Now, by integrating both sides of the last inequality and dividing by t, we find immediately that

$$t^{-1} \ln \mathcal{M}(I(t), \mathcal{D}(t)) \leq t^{-1} \ln \mathcal{M}(I(0), \mathcal{D}(0)) + \Upsilon + \Pi - \Sigma + \frac{\omega_1 \beta}{\omega_2 t} \int_0^t \left| \psi(s) - \frac{A}{\mu_1} \right| ds + t^{-1} \mathcal{J}_3(t) + t^{-1} \mathcal{J}_4(t),$$
(11)

where

$$\mathcal{J}_{3}(t) = \int_{0}^{t} \frac{\sigma_{2}\omega_{1}I(s)}{\omega_{1}I(s) + \omega_{2}D(s)} \mathrm{d}\mathcal{W}_{2}(s) + \int_{0}^{t} \frac{\sigma_{4}\omega_{2}D(s)}{\omega_{1}I(s) + \omega_{2}D(s)} \mathrm{d}\mathcal{W}_{4}(s),$$
$$\mathcal{J}_{4}(t) = \int_{0}^{t} \int_{\mathcal{U}} \ln(1 + \lambda(u))\widetilde{\mathcal{N}}(\mathrm{d}s, \mathrm{d}u).$$

It is easy to check that $\mathcal{J}_3(t)$ is a local martingale with finite quadratic variation, and by the hypothesis (\mathcal{H}_3) we can affirm that $\mathcal{J}_4(t)$ is also a local martingale with finite quadratic variation. By the strong law of large numbers for local martingales [29], we get

$$\lim_{t \to \infty} t^{-1} \mathcal{J}_3(t) = 0 \quad \text{a.s} \quad \text{and} \quad \lim_{t \to \infty} t^{-1} \mathcal{J}_4(t) = 0 \quad \text{a.s.}$$

Now, by using the Hölder's inequality, we deduce that

$$t^{-1} \int_0^t \left| \psi(s) - \frac{A}{\mu_1} \right| \mathrm{d}s \le t^{-\frac{1}{2}} \left(\int_0^t \left(\psi(s) - \frac{A}{\mu_1} \right)^2 \mathrm{d}s \right)^{\frac{1}{2}} = \left(t^{-1} \int_0^t \left(\psi^2(s) - \frac{2A}{\mu_1} \psi(s) + \left(\frac{A}{\mu_1} \right)^2 \right) \mathrm{d}s \right)^{\frac{1}{2}}.$$

It follows from Lemme 3.4 that

$$\lim_{t \to \infty} t^{-1} \int_0^t \left| \psi(s) - \frac{A}{\mu_1} \right| \mathrm{d}s \le \left(\frac{2A^2}{\mu_1 \chi_2} - \frac{2A^2}{\mu_1^2} + \frac{A^2}{\mu_1^2} \right)^{\frac{1}{2}} = \left(\frac{A^2 \left(\sigma_1^2 + \int_{\mathcal{U}} \lambda_1^2(u) \nu(\mathrm{d}u) \right)}{\mu_1^2 \chi_2} \right)^{\frac{1}{2}}.$$

Taking the superior limit on both sides of (11) leads to

$$\limsup_{t \to \infty} t^{-1} \ln \mathcal{M}(I(t), \mathcal{D}(t)) \leq \Upsilon + \Pi - \Sigma + \frac{\omega_1 \beta}{\omega_2} \left(\frac{A^2 \left(\sigma_1^2 + \int_{\mathcal{U}} \lambda_1^2(u) \nu(\mathrm{d}u) \right)}{\mu_1^2 \chi_2} \right)^{\frac{1}{2}} \quad \text{a.s.}$$

Which implies,

$$\limsup_{t \to \infty} t^{-1} \ln \mathcal{M}(I(t), \mathcal{D}(t)) \le \Upsilon + \Pi - \Sigma + \eta \left(\frac{\mathcal{T}^*\Lambda}{\chi_2}\right)^{\frac{1}{2}} = \Theta \quad \text{a.s}$$

That is to say, if $\Theta < 0$, then $\limsup_{t \to \infty} t^{-1} \ln I(t) < 0$, and $\limsup_{t \to \infty} t^{-1} \ln \mathcal{D}(t) < 0$ a.s., which implies in turn that the disease will die out with probability one and this completes the proof.

Remark 3.8. When the jumps coefficients $\lambda_i(u)$ (i = 1, 2, 4) and the white intensities σ_i (i = 2, 4) degenerate to zero, our results in Theorem 3.7 coincide with Theorem 2.3 in [26]. Therefore, our results generalize the consequence of the mentioned paper.

3.2. Persistence in mean of the epidemic

The study of the persistence in the mean is a significant characteristic to know more about epidemic dynamics. For this reason, in this section, we will give the condition for the disease persistence, but before stating the main result, we shall first recall the concept of persistence in the mean.

Definition 3.9. An infected population $\mathcal{I}(t)$ is said to be be strongly persistent in the mean, or just persistent in the mean, if $\liminf_{t\to\infty} t^{-1} \int_0^t \mathcal{I}(s) ds > 0$ almost surely.

For simplicity of notation, we define the following quantity

$$\widetilde{\mathcal{T}^{\star}} = \beta \left(\frac{A}{\mu_1 + \bar{\sigma}_1} \right) \left((\mu_2 + \gamma + \bar{\sigma}_2) + \beta \left(\frac{A}{\mu_1 + \bar{\sigma}_1} \right) \frac{\bar{\sigma}_4}{\eta} \right)^{-1}$$

where $\bar{\sigma}_i \triangleq 0.5\sigma_i^2 + \int_{\mathcal{U}} \left(\lambda_i(u) - \ln(1 + \lambda_i(u))\right) \nu(\mathrm{d}u), i = 1, 2, 4.$

Theorem 3.10. Let $(S(t), I(t), \mathcal{D}(t))$ be the solution of (5) with any initial data $(S(0), I(0), \mathcal{D}(0)) \in \mathbb{R}^3_+$. The stochastic model (5) has the following property: if $\widetilde{\mathcal{T}^*} > 1$ holds, then the disease I(t) persists in the mean almost surely.

Proof. Begin by considering the following function

$$\mathcal{Z}(S(t), I(t), \mathcal{D}(t)) = -\mathfrak{c}_1 \ln S(t) - \ln I(t) - \mathfrak{c}_2 \ln \mathcal{D}(t) + \mathfrak{c}_3 \mathcal{D}(t).$$

where c_i , (i = 1, 2, 3) are positive constants to be determined in the following. From Itô's formula and system (5), we have

$$\begin{split} \mathrm{d}\mathcal{Z}(S(t),I(t),\mathcal{D}(t)) &= \mathcal{L}\mathcal{Z}(S(t),I(t),\mathcal{D}(t))\mathrm{d}t - \mathfrak{c}_{1}\sigma_{1}\mathrm{d}\mathcal{W}_{1}(t) - \sigma_{2}\mathrm{d}\mathcal{W}_{2}(t) - \mathfrak{c}_{2}\sigma_{4}\mathrm{d}\mathcal{W}_{4}(t) \\ &+ \mathfrak{c}_{3}\sigma_{4}D(t)\mathrm{d}\mathcal{W}_{4}(t) - \int_{\mathcal{U}}\mathfrak{c}_{1}\ln\left(1 + \lambda_{1}(u)\right)\widetilde{\mathcal{N}}(\mathrm{d}t,\mathrm{d}u) - \int_{\mathcal{U}}\ln\left(1 + \lambda_{2}(u)\right)\widetilde{\mathcal{N}}(\mathrm{d}t,\mathrm{d}u) \\ &- \int_{\mathcal{U}}\mathfrak{c}_{2}\ln\left(1 + \lambda_{4}(u)\right)\widetilde{\mathcal{N}}(\mathrm{d}t,\mathrm{d}u) + \int_{\mathcal{U}}\mathfrak{c}_{3}\lambda_{4}(u)\mathcal{D}(t^{-})\widetilde{\mathcal{N}}(\mathrm{d}t,\mathrm{d}u), \end{split}$$

where

$$\begin{aligned} \mathcal{LZ}(S(t), I(t), \mathcal{D}(t)) &= -\frac{\mathfrak{c}_1}{S(t)} (A - \mu_1 S(t) - \beta S(t) \mathcal{D}(t)) + \frac{\mathfrak{c}_1 \sigma_1^2}{2} - \frac{1}{I(t)} \left(\beta S(t) \mathcal{D}(t) - (\mu_2 + \gamma) I(t) \right) \\ &+ \frac{\sigma_2^2}{2} - \frac{\mathfrak{c}_2 \eta}{\mathcal{D}(t)} (I(t) - \mathcal{D}(t)) + \frac{\mathfrak{c}_2 \sigma_4^2}{2} + \mathfrak{c}_3 \eta \left(I(t) - \mathcal{D}(t) \right) \\ &+ \int_{\mathcal{U}} \mathfrak{c}_1 \left(\lambda_1(u) - \ln(1 + \lambda_1(u)) \nu(\mathrm{d}u) + \int_{\mathcal{U}} \left(\lambda_2(u) - \ln(1 + \lambda_2(u)) \nu(\mathrm{d}u) \right) \\ &+ \int_{\mathcal{U}} \mathfrak{c}_2 \left(\lambda_4(u) - \ln(1 + \lambda_4(u)) \nu(\mathrm{d}u) \right). \end{aligned}$$

We then find that

$$\begin{aligned} \mathcal{LZ}(S(t), I(t), \mathcal{D}(t)) &= -\frac{\beta S(t) \mathcal{D}(t)}{I(t)} - \frac{\mathfrak{c}_1 A}{S(t)} - \frac{\mathfrak{c}_2 \eta I(t)}{\mathcal{D}(t)} + (\mathfrak{c}_1 \beta - \mathfrak{c}_3 \eta) \mathcal{D}(t) \\ &+ \mathfrak{c}_1(\mu_1 + \bar{\sigma}_1) + \mathfrak{c}_2(\eta + \bar{\sigma}_4) + (\mu_2 + \gamma + \bar{\sigma}_2) + \mathfrak{c}_3 \eta I(t) \\ &\leq -3 \big(\beta A \eta \mathfrak{c}_1 \mathfrak{c}_2\big)^{\frac{1}{3}} + (\mathfrak{c}_1 \beta - \mathfrak{c}_3 \eta) \mathcal{D}(t) + \mathfrak{c}_1(\mu_1 + \bar{\sigma}_1) \\ &+ \mathfrak{c}_2(\eta + \bar{\sigma}_4) + (\mu_2 + \gamma + \bar{\sigma}_2) + \mathfrak{c}_3 \eta I(t). \end{aligned}$$

By choosing

$$\begin{aligned} \mathbf{c}_1 &= \beta \left(\frac{A}{\mu_1 + \bar{\sigma}_1} \right)^2 (\eta + \bar{\sigma}_4) / A\eta, \\ \mathbf{c}_2 &= \beta \left(\frac{A}{\mu_1 + \bar{\sigma}_1} \right) / (\eta + \bar{\sigma}_4), \\ \mathbf{c}_3 &= \mathbf{c}_1 \beta / \eta, \end{aligned}$$

we may actually obtain that

$$\mathcal{LZ}(S(t), I(t), \mathcal{D}(t)) \leq -\beta \left(\frac{A}{\mu_1 + \bar{\sigma}_1}\right) + (\mu_2 + \gamma + \bar{\sigma}_2) + \beta \left(\frac{A}{\mu_1 + \bar{\sigma}_1}\right) \frac{\bar{\sigma}_4}{\eta} + \mathfrak{c}_1 \beta I(t)$$
$$= -\beta \left(\frac{A}{\mu_1 + \bar{\sigma}_1}\right) \left(1 - \frac{1}{\widetilde{\mathcal{T}^{\star}}}\right) + \mathfrak{c}_1 \beta I(t).$$

Hence, we get

$$\begin{split} \mathrm{d}\mathcal{Z}(S(t),I(t),\mathcal{D}(t)) &\leq \left(-\beta\Big(\frac{A}{\mu_1+\bar{\sigma}_1}\Big)\left(1-\frac{1}{\widetilde{\mathcal{T}^{\star}}}\right) + \mathfrak{c}_1\beta I(t)\right)\mathrm{d}t - \mathfrak{c}_1\sigma_1\mathrm{d}\mathcal{W}_1(t) - \sigma_2\mathrm{d}\mathcal{W}_2(t) \\ &-\mathfrak{c}_2\sigma_4\mathrm{d}\mathcal{W}_4(t) + \mathfrak{c}_3\sigma_4D(t)\mathrm{d}\mathcal{W}_4(t) - \int_{\mathcal{U}}\mathfrak{c}_1\ln\big(1+\lambda_1(u)\big)\widetilde{\mathcal{N}}(\mathrm{d}t,\mathrm{d}u) \\ &-\int_{\mathcal{U}}\ln\big(1+\lambda_2(u)\big)\widetilde{\mathcal{N}}(\mathrm{d}t,\mathrm{d}u) - \int_{\mathcal{U}}\mathfrak{c}_2\ln\big(1+\lambda_4(u)\big)\widetilde{\mathcal{N}}(\mathrm{d}t,\mathrm{d}u) \\ &+\int_{\mathcal{U}}\mathfrak{c}_3\lambda_4(u)\mathcal{D}(t^-)\widetilde{\mathcal{N}}(\mathrm{d}t,\mathrm{d}u). \end{split}$$

Integrating from 0 to t and dividing by t on both sides of the last inequality, yields

$$t^{-1} \mathcal{Z}(S(t), I(t), \mathcal{D}(t)) \leq t^{-1} \mathcal{Z}(S(0), I(0), \mathcal{D}(0)) - \beta \left(\frac{A}{\mu_1 + \bar{\sigma}_1}\right) \left(1 - \frac{1}{\widetilde{\mathcal{T}^{\star}}}\right) + \mathfrak{c}_1 \beta t^{-1} \int_0^t I(s) \mathrm{d}s + t^{-1} \mathcal{J}_5(t) + t^{-1} \mathcal{J}_6(t),$$

where

$$\mathcal{J}_{5}(t) = -\left(\mathfrak{c}_{1}\sigma_{1}\mathcal{W}_{1}(t) + \sigma_{2}\mathcal{W}_{2}(t) + \mathfrak{c}_{2}\sigma_{4}\mathcal{W}_{4}(t)\right) + \mathfrak{c}_{3}\sigma_{4}\int_{0}^{t}D(s)\mathrm{d}\mathcal{W}_{4}(s),$$
$$\mathcal{J}_{6}(t) = -\int_{0}^{t}\int_{\mathcal{U}}\mathfrak{c}_{1}\ln\left(1 + \lambda_{1}(u)\right)\widetilde{\mathcal{N}}(\mathrm{d}t,\mathrm{d}u) - \int_{0}^{t}\int_{\mathcal{U}}\ln\left(1 + \lambda_{2}(u)\right)\widetilde{\mathcal{N}}(\mathrm{d}t,\mathrm{d}u) - \int_{0}^{t}\int_{\mathcal{U}}\ln\left(1 + \lambda_{2}(u)\right)\widetilde{\mathcal{N}}(\mathrm{d}t,\mathrm{d}u) - \int_{0}^{t}\int_{\mathcal{U}}\ln\left(1 + \lambda_{2}(u)\right)\widetilde{\mathcal{N}}(\mathrm{d}t,\mathrm{d}u) + \int_{0}^{t}\int_{\mathcal{U}}\mathfrak{c}_{3}\lambda_{4}(u)\mathcal{D}(t^{-})\widetilde{\mathcal{N}}(\mathrm{d}t,\mathrm{d}u).$$

By using, the strong law of large numbers for local martingales and Lemma 3.1, we can obtain

$$\lim_{t \to \infty} t^{-1} \mathcal{J}_5(t) = 0 \quad \text{a.s} \quad \text{and} \quad \lim_{t \to \infty} t^{-1} \mathcal{J}_6(t) = 0 \quad \text{a.s}$$

Therefore

$$\liminf_{t \to \infty} t^{-1} \int_0^t I(s) \mathrm{d}s \ge \frac{1}{\mathfrak{c}_1} \left(\frac{A}{\mu_1 + \bar{\sigma}_1} \right) \left(1 - \frac{1}{\widetilde{\mathcal{T}^{\star}}} \right) > 0 \quad \text{a.s.}$$

This shows that the disease persists in the mean as claimed.

Remark 3.11. Persistence in the mean is an important concept in mathematical epidemiology. It captures the long-term survival of the disease even when the population size is quite low at t = 0. Moreover, the persistence of the model refers to a situation where the disease is endemic in a population.

4. Numerical simulations

This section is devoted to illustrate our theoretical results by employing numerical simulations. In the three following examples, we apply the algorithm presented in [35] to discretize the disturbed system (5). Using the software Matlab2015b and the parameter values listed in Table 1, we numerically simulate the solution of the system (5) with the initial value $(S(0), I(0), \mathcal{D}(0)) = (0.6, 0.3, 0.05)$.

Parameters	Description	Numerical values			
A	Recruitment rate	0.9	0.3	0.6	0.6
μ_1	Natural mortality rate of S	0.3	0.3	0.4	0.4
β	Transmission rate	0.07	1.3	0.35	0.8
γ	Recovered rate	0.05	0.05	0.2	0.3
μ_2	General mortality of I	0.5	0.5	0.3	0.3
η	Exponentially fading memory rate	0.09	0.09	0.7	0.2
σ_1	Intensity of $\mathcal{W}_1(t)$	0.15	0.15	0.2	0.169
σ_2	Intensity of $\mathcal{W}_2(t)$	0.25	0.25	0.15	0.15
σ_4	Intensity of $\mathcal{W}_4(t)$	0.27	0.27	0.13	0.13
λ_1	Jump intensity of S	0.2	0.2	0.5	0.5
λ_2	Jump intensity of I	0.23	0.23	0.3	0.3
λ_4	Jump intensity of \mathcal{D}	0.1	0.1	0.7	0.7
		Figure 1	Figure 2	Figure 3	Figure 4

Table 1: Nominal values of the system parameters and disturbances intensities adopted in the different simulation examples .

4.1. The stochastic extinction case

In order to exhibit the strong random fluctuations effect on epidemic dynamics, we present in Figure 1, the trajectories of the stochastic solution $(S(t), I(t), \mathcal{D}(t))$. We assume that $\mathcal{U} = (0, \infty)$ and $\nu(\mathcal{U}) = 1$, then by using the parameters listed in Table 1, we must check the existence of p such that $\chi_{1,p} > 0$. By simple calculation, we easily get $\chi_{1,p} = 0.0206$ for p = 2.1. Then, the condition (\mathcal{H}_5) is satisfied. With the chosen parameters, we can obtain the following values:

Expression	Value
$\mathcal{T}^{\star} \triangleq \beta A(\mu_1(\mu_2 + \gamma))^{-1}$	0.3818
$\Upsilon \triangleq \min\{\mu_2 + \gamma, \eta\}(\sqrt{\mathcal{T}^{\star}} - 1)$	-0.0344
$\Pi \triangleq \int_{\mathcal{U}} \aleph(u) \nu(\mathrm{d}u)$	-0.0047
$\Sigma \triangleq \left(2 \left(\sigma_2^{-2} + \sigma_4^{-2} \right) \right)^{-1}$	0.0168
$\Lambda \triangleq \sigma_1^2 + \int_{\mathcal{U}} \lambda_1^2(u) \nu(\mathrm{d}u)$	0.0625
$\chi_2 = 2\mu_1 - \sigma_1^2 - \int_{\mathcal{U}} \lambda_1^2(u)\nu(\mathrm{d}u)$	0.5375
$\Theta \triangleq \Upsilon + \Pi - \Sigma + \eta \sqrt{\mathcal{T}^{\star}} \Lambda^{\frac{1}{2}} \chi_2^{-\frac{1}{2}}$	-0.0369

Table 2: Some expressions and their corresponding values.

From Table 2, we have $\Theta < 0$, then the condition of Theorem 3.7 is verified. That is to say that the epidemic dies out exponentially almost surely which is exactly illustrated in Figure 1.

4.2. The stochastic persistence case

Consider the system (5) with parameters appearing in Table 1. Then, we obtain the following values:



Figure 1: The paths of S(t), I(t) and $\mathcal{D}(t)$ for the stochastic model (5) when $\Theta = -0.0369 < 0$.

Expression		
$\bar{\sigma}_1 \triangleq 0.5\sigma_1^2 + \int_{\mathcal{U}} \left(\lambda_1(u) - \ln(1 + \lambda_1(u)) \right) \nu(\mathrm{d}u)$	0.0289	
$\bar{\sigma}_2 \triangleq 0.5\sigma_2^2 + \int_{\mathcal{U}} \left(\lambda_2(u) - \ln(1 + \lambda_2(u)) \right) \nu(\mathrm{d}u)$	0.0542	
$\bar{\sigma}_4 \triangleq 0.5\sigma_4^2 + \int_{\mathcal{U}} \left(\lambda_4(u) - \ln(1 + \lambda_4(u)) \right) \nu(\mathrm{d}u)$	0.0411	
$\widetilde{\mathcal{T}^{\star}} \triangleq \beta \left(\frac{A}{\mu_1 + \bar{\sigma}_1} \right) \left((\mu_2 + \gamma + \bar{\sigma}_2) + \beta \left(\frac{A}{\mu_1 + \bar{\sigma}_1} \right) \frac{\bar{\sigma}_4}{\eta} \right)^{-1}$	1.0344	

 Table 3: Some expressions and their corresponding values.

Therefore, $\widetilde{\mathcal{T}^{\star}} > 1$. From Figure 2, we observe the persistence of the epidemic I(t) in this case, which agree well with Theorem 3.10. Furthermore, the solutions S(t) and D(t) are persistent which implies the non-extinction of the stochastic model (5).

4.3. The Lévy jumps effect on the epidemic dynamics

To find out the effect of white noise and jumps intensities on epidemic dynamics, in this example, we will compare the trajectories of the following systems:

- The deterministic model (3) $(\sigma_i = 0 \text{ and } \lambda_i = 0, i = 1, 2, 4).$
- The stochastic version of (3) with degenerate diffusion [26] ($\sigma_2 = \sigma_4 = 0$ and $\lambda_i = 0$, i = 1, 2, 4).



Figure 2: The paths of S(t), I(t) and $\mathcal{D}(t)$ for the stochastic model (5) when $\widetilde{\mathcal{T}^{\star}} = 1.0344 > 1$.

• The SDE-J system (5) ($\sigma_i \neq 0$ and $\lambda_i \neq 0$, i = 1, 2, 4).

We take the values appearing in Table 1 which are the same as those used in [26]. For the rest of parameters, we choose $\sigma_1 = 0.2$, $\sigma_2 = 0.15$, $\sigma_4 = 0.13$, $\lambda_1 = 0.5$, $\lambda_2 = 0.3$, and $\lambda_4 = 0.7$. For the sake of a comparison, we choose the following initial value used in $(S(0), I(0), \mathcal{D}(0)) = (0.2, 0.3, 0.4)$ used in [26]. We see from Figure 3 that the effects of Lévy jumps lead to the extinction of the disease while the deterministic model (3) and the perturbed model driven by degenerate diffusion both predict persistence. Therefore, we say that the jumps have negative effects on the prevalence of epidemics. This means that jumps can change the asymptotic behavior of the epidemic model significantly. To examine the effect of jumps intensities on dynamical system (3) in the case of persistence, we shall decrease the intensity σ_1 to 0.169 and take other parameter as in the last column of Table 1. From Figure 4, we observe the persistence of the epidemic in all cases with a greater variation in the case of Lévy jumps.

5. Conclusion and discussion

Environmental factors and unexpected phenomena have significant impacts on the spread of epidemics. This paper takes into account these two factors. Specifically, we have analyzed a delayed SIR epidemic model that incorporates proportional Lévy jumps. For analytical reasons, we have employed the linear chain approach to transform the model with a weak kernel case (1) into the equivalent system (2). After proving the well-posedness of this perturbed model, we have analyzed its long-term behavior. Under some hypotheses, the main epidemiological findings of our study are presented as follows:



Figure 3: The paths of S(t), I(t) and $\mathcal{D}(t)$ for the deterministic model (2), the model (3) with degenerate diffusion, and the SDE-J (5).

- 1. We have given sufficient condition for the extinction of the epidemic.
- 2. We have established sufficient condition for the persistence in the mean of the epidemic.

Compared to the existing literature, the novelty of our work lies in new mathematical analysis techniques and improvements which are summarized in the following items:

- 1. Our work is distinguished from previous works [8, 7, 10] by the use of the expression $\chi_{1,p}$ which boosts the optimality of our calculus and results.
- 2. Our study offers an alternative method to the gap mentioned in (Theorem 2.2, [60]). Without using the explicit formula of the distribution stationary $\pi^*(\cdot)$ of ψ (which still up to now unknown), we calculate the following time averages:

$$\lim_{t \to \infty} t^{-1} \int_0^t \psi(s) \mathrm{d}s \text{ and } \lim_{t \to \infty} t^{-1} \int_0^t \psi^2(s) \mathrm{d}s \text{ a.s.}$$

3. In order to find an optimal and good majorization, we have considered the inequality (10) in our analysis without eliminated it (since $\ln(1 + x) - x \le 0$ for all x > -1) which differs from the calculus presented in [7].

Generally speaking, our theoretical results indicate that the conditions of extinction and persistence are mainly depending on the magnitude of the noise intensities as well as the system parameters. From numerical simulations, we remark that Lévy jumps affect significantly the long-run behavior of an epidemic. Eventually, we point out that this paper extends the study presented in [26] to the case of Lévy jumps and delivers some new insights for understanding the propagation of diseases with distributed delay. Furthermore, the method developed in this paper can be used to investigate a class of related stochastic models driven by Lévy noise.

Data Availability

The theoretical data used to support the findings of this study are included in the article.

Conflicts of Interest

On behalf of all authors, the corresponding author states that there is no conflict of interest.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Authors' Contributions

The authors declare that the study was conducted in collaboration with the same responsibility. All authors read and approved the final manuscript.

References

- S. Bentout, Y. Chen, and S. Djilali. Global dynamics of an SEIR model with two age structures and a nonlinear incidence. Acta Applicandae Mathematicae, 171(7), 2021.
- [2] S. Bentout, A. Tridane, S. Djilali, and T. M. Touaoula. Age-structured modeling of COVID-19 epidemic in the USA, UAE and Algeria. Alexandria Engineering Journal, 60(1):401–411, 2021.
- [3] E. Beretta, T. Hara, and W. Ma. Global asymptotic stability of an SIR epidemic model with distributed time delay. *Nonlinear Analysis*, 47:4107–4115, 2001.
- [4] E. Beretta and Y. Takeuchi. Global stability of an SIR epidemic model with time delay. Journal of Mathematical Biology, 33:250–260, 1995.
- [5] A. Berman and R.J. Plemmons. Nonnegative matrices in the mathematical sciences. Academic Press, New York, 1979.
- [6] G. Chen and T. Li. Stability of stochastic delayed SIR model. Stochastics and Dynamics, 9(2):231–252, 2009.
- [7] Y. Cheng, M. Li, and F. Zhang. A dynamics stochastic model with HIV infection of CD4 T cells driven by Levy noise. *Chaos, Solitons and Fractals*, 129:62–70, 2019.
- [8] Y. Cheng, F. Zhang, and M. Zhao. A stochastic model of HIV infection incorporating combined therapy of haart driven by Levy jumps. Advance in Difference Equations, 321, 2019.
- [9] S. Djilali, L. Benahmadi, A. Tridane, and K. Niri. Modeling the impact of unreported cases of the COVID-19 in the north african countries. *Biology*, 9(11):373, 2020.
- [10] M. Gao, D. Jiang, T. Hayat, and A. Alsaedi. Threshold behavior of a stochastic lotka volterra food chain chemostat model with jumps. *Chaos, Solitons and Fractals*, 523:191–203, 2019.
- [11] I. I. Gihman and A. V. Skorohod. Stochastic Differential Equations. Springer-Verlag Berlin Heidelberg, 1972.
- [12] H. Guo, M.Y. Li, and Z. S. Shuai. Global stability of the endemic equilibrium of multigroup SIR epidemic models. Canadian Applied Mathematics Quarterly, 14:259–284, 2006.
- [13] G. Hussain, A. Khan, M. Zahri, and G. Zaman. Stochastic permanence of an epidemic model with a saturated incidence rate. *Chaos Solitons and Fractals*, 139:110005, 2020.



Figure 4: The paths of S(t), I(t) and $\mathcal{D}(t)$ associated respectively to the models (2),(3) and (5).

- [14] C. Ji and D. Jiang. Threshold behaviour of a stochastic SIR model. Applied Mathematical Modelling, 38:5067– 5079, 2014.
- [15] C. Ji, D. Jiang, and N. Shi. Asymptotic behavior of global positive solution to a stochastic SIR model. Applied Mathematical Modelling, 45:221–232, 2011.
- [16] C. Ji, D. Jiang, and N. Shi. The behavior of an SIR epidemic model with stochastic perturbation. Stochastic analysis and applications, 30:755–773, 2012.
- [17] W. O. Kermack and A. G. McKendrick. A contribution to the mathematical theory of epidemics. Proceedings of The Royal Society A Mathematical Physical and Engineering Sciences, 115(772):700–721, 1927.
- [18] A. Khan, G. Hussain, M. Zahri, and G. Zaman. A stochastic SACR epidemic model for HBV transmission. Journal of Biological Dynamics, 14(1):788–801, 2020.
- [19] M. A. Khan, A. Atangana, E. Alzahrani, and Fatmawati. The dynamics of COVID-19 with quarantined and isolation. Advances in Difference Equations, 425, 2020.
- [20] D. Kiouach, S. E. A. El-idrissi, and Y. Sabbar. Advanced and comprehensive research on the dynamics of COVID-19 under mass communication outlets intervention and quarantine strategy: a deterministic and probabilistic approach. ArXiv preprint, 2101.00517, 2021.
- [21] D. Kiouach and Y. Sabbar. Stability and threshold of a stochastic SIRS epidemic model with vertical transmission and transfer from infectious to susceptible individuals. *Discrete Dynamics in Nature and Society*, (7570296), 2018.

- [22] D. Kiouach, Y. Sabbar, and S. E. A. El-idrissi. New results on the asymptotic behavior of an SIS epidemiological model with quarantine strategy, stochastic transmission, and Levy disturbance. arXiv preprint, 2012.00875, 2020.
- [23] Y. Kyrychko and K. Blyuss. Global properties of a delayed SIR model with temporary immunity and nonlinear incidence rate. *Nonlinear Analysis*, 6:495–507, 2005.
- [24] Y. Lin, D. Jiang, and P. Xia. Long-time behavior of a stochastic SIR model. Applied Mathematics and Computation, 236:1–9, 2014.
- [25] Q. Liu, Q. Chen, and D. Jiang. The threshold of a stochastic delayed SIR epidemic model with temporary immunity. *Physica A*, 450:115–125, 2016.
- [26] Q. Liu, D. Jiang, T. Hayat, and A. Alsaedi. Dynamics of a stochastic SIR epidemic model with distributed delay and degenerate diffusion. *Journal of the Franklin Institute*, 356:7347–7370, 2019.
- [27] Q. Liu, D. Jiang, N. Shi, T. Hayat, and A. Alsaedi. Asymptotic behaviors of a stochastic delayed sir epidemic model with nonlinear incidence. *Communications in Nonlinear Science and Numerical Simulation*, 40:89–99, 2016.
- [28] W. Ma, Y. Takeuchi, T. Hara, and E. Beretta. Epermanence of an SIR epidemic model with distributed time delays. *Tohoku Mathematical Journal*, 54:581–591, 2002.
- [29] X. Mao. Stochastic Differential Equations and Applications. Horwoodl, Chichester, 1997.
- [30] S. B. Mendrazitsky and L. Shaikhet. Stability analysis of delayed tumor-antigen-activated immune response in combined BCG and IL-2 immunotherapy of bladder cancer. MDPI, Processes, 8(12):1–17, 2020.
- [31] X. Z. Meng and L. S. Chen. The dynamics of a new SIR epidemic model concerning pulse vaccination strategy. Applied Mathematics and Computation, 197:528–597, 2008.
- [32] Y. Muroya, T. Kuniya, and J. Wang. Stability analysis of a delayed multi-group SIS epidemic model with nonlinear incidence rates and patch structure. *Journal of Mathematical Analysis and Applications*, 425:415–439, 2015.
- [33] M. Pitchaimani and D. M. Brasanna. Stochastic dynamical probes in a triple delayed SICR model with general incidence rate and immunization strategies. *Chaos, Solitons and Fractals*, 143:110540, 2021.
- [34] S. P. Rajasekar M. Pitchaimani. Ergodic stationary distribution and extinction of a stochastic sirs epidemic model with logistic growth and nonlinear incidence. *Physica A*, 482:125143, 2020.
- [35] P. Portter and D. Talay. The euler scheme for Levy driven stochastic differential equations. Ann. Probab., 25:393-423, 1997.
- [36] S. P. Rajasekar, M. Pitchaimani, and Q. Zhu. Dynamic threshold probe of stochastic SIR model with saturated incidence rate and saturated treatment function. *Physica A*, 535:122300, 2019.
- [37] M. Roy and R. D. Holt. Effects of predation on host-pathogen dynamics in SIR models. Theoretical Population Biology, 73:319–331, 2008.
- [38] L. Shaikhet. Improving stability conditions for equilibria of SIR epidemic model with delay under stochastic perturbations. *Mathematics*, 8(8):1302, 2020.
- [39] L. Shaikhet. Stability of stochastic differential equations with distributed and state-dependent delays. Journal of Applied Mathematics and Computation, 4(4):181–188, 2020.
- [40] L. Shaikhet. Behavior of solution of stochastic delay differential equation with additive fading perturbations. Applied Mathematics Letters, 111(2021)(106640), 2021.
- [41] L. Shaikhet and T. Caraballo. Stability of delay evolution equations with fading stochastic perturbations. International Journal of Control, pages 1–7, 2020.
- [42] H. Shu, D. Fan, and J. Wei. Global stability of multi-group seir epidemic models with distributed delays and nonlinear transmission. NNonlinear Analysis: Real World Applications, 13:1581–1591, 2012.

- [43] F. Sun. Dynamics of an imprecise stochastic Holling ii one predator two prey system with jumps. arXiv preprint, 2006.14943, 2020.
- [44] F. Sun. Dynamics of an imprecise stochastic multimolecular biochemical reaction model with Levy jumps. arXiv preprint, 2004.14163, 2020.
- [45] Y. Takeuchi, W. Ma, and E. Beretta. Global asymptotic properties of a delay SIR epidemic model with finite incubation times. *Nonlinear Analysis*, 42:931–947, 2000.
- [46] B. Tang, F. Xia, S. Tang, N. L. Bragazzi, Q. Li, X. Sun, J. Liang, Y. Xiao, and J. Wu. The effectiveness of quarantine and isolation determine the trend of the COVID-19 epidemics in the final phase of the current outbreak in china. *International Journal of Infectious Diseases*, 95:288–293, 2020.
- [47] E. Tornatore, S. Buccellato, and P. Vetro. Stability of a stochastic SIR system. *Physica A*, 354:111–126, 2005.
- [48] Y. Wang and D. Jiang. Stationary distribution and extinction of a stochastic viral infection model. Discrete Dynamics in Nature and Society, 2017, ID 6027509, 2017.
- [49] C. Xu, S. Yuan, and T. Zhang. Competitive exclusion in a general multi-species Chemostat model with stochastic perturbations. Bulletin of Mathematical Biology, 83(4), 2021.
- [50] S. Yan and S. Yuan. Critical value in a SIR network model with heterogeneous infectiousness and susceptibility. Mathematical Biosciences and Engineering, 17(5):5802–5811, 2020.
- [51] A. Yang, B. Song, and S. Yuan. Noise-induced transitions in a non-smooth SIS epidemic model with media alert. Mathematical Biosciences and Engineering, 18(1):745–763, 2020.
- [52] S. Zhang, T. Zhang, and S. Yuan. Dynamics of a stochastic predator-prey model with habitat complexity and prey aggregation. *Ecological Complexity*, 45:100889, 2021.
- [53] X. Zhang and K. Wang. Stochastic SIR model with jumps. Applied Mathematics letters, 826:867–874, 2013.
- [54] X. B. Zhang, H. Huo, H. Xiang, and X. Meng. Dynamics of the deterministic and stochastic SIQS epidemic model with nonlinear incidence. *Applied Mathematics and Computation*, 243:546–558, 2014.
- [55] X. B. Zhang, H. Huo, H. Xiang, Q. Shi, and D. Li. The threshold of a stochastic SIQS epidemic model. *Physica* A, 482:362–374, 2017.
- [56] X. B. Zhang and R. J. Liu. The stationary distribution of a stochastic SIQS epidemic model with varying total population size. Applied Mathematics Letters, 116(2021)(106974), 2021.
- [57] X. B. Zhang, Q. Shi, S. Ma, H. Huo, and D. Li. Dynamic behavior of a stochastic SIQS epidemic model with Levy jumps. *Nonlinear Dynamics*, 93:1481–1493, 2018.
- [58] X. B. Zhang and X. H. Zhang. The threshold of a deterministic and a stochastic SIQS epidemic model with varying total population size. *Applied Mathematical Modelling*, 91:749–767, 2021.
- [59] D. Zhao and S. Yuan. Sharp conditions for the existence of a stationary distribution in one classical stochastic chemostat. Applied Mathematics and Computation, 339:199–205, 2018.
- [60] D. Zhao, S. Yuan, and H. Liu. Stochastic dynamics of the delayed chemostat with Levy noises. International Journal of Biomathematics, 12(5), 2019.
- [61] Y. Zhou and W. Zhang. Threshold of a stochastic SIR epidemic model with Levy jumps. Physica A, 446:204–2016, 2016.