

# Forecasting and evaluating intervention of Covid-19 in the World

Zixin Hu, Ph.D<sup>1,2</sup>; Qiyang Ge, MS<sup>3</sup>; Shudi Li, MS<sup>4</sup>; Eric Boerwinkle<sup>4</sup>, Ph.D<sup>4</sup>, Li Jin, Ph.D<sup>1,2</sup>  
and Momiao Xiong, Ph.D<sup>4,\*</sup>

<sup>1</sup> State Key Laboratory of Genetic Engineering and Innovation Center of Genetics and Development, School of Life Sciences, Fudan University, Shanghai, China.

<sup>2</sup> Human Phenome Institute, Fudan University, Shanghai, China.

<sup>3</sup> The School of Mathematic Sciences, Fudan University, Shanghai, China.

<sup>4</sup> School of Public Health, The University of Texas Health Science Center at Houston, Houston, TX 77030, USA.

**Running Title:** Real Time Forecasting of Covid-19 in the world

**Keywords:** Cov-19, artificial intelligence, transmission dynamics, forecasting, time series, auto-encoder.

\*Address for correspondence and reprints: Dr. Momiao Xiong, Department of Biostatistics and Data Science, School of Public Health, The University of Texas Health Science Center at Houston, P.O. Box 20186, Houston, Texas 77225, (Phone): 713-500-9894, (Fax): 713-500-0900, E-mail: [Momiao.Xiong@uth.tmc.edu](mailto:Momiao.Xiong@uth.tmc.edu).

## **Key Points**

**Question** As the Covid-19 pandemic surges around the world, what are the number of cases worldwide at its peak, the length of the pandemic before receding, the fatality rate and the timing of public health interventions to significantly stop the spread of Covid-19?

**Findings** We estimated the maximum number of cumulative cases under immediate intervention to be 1,530,276; under later intervention the number increased to a frightening 255,392,154 and the case ending time was delayed to January 10, 2021.

**Meaning** If there is not immediate aggressive action to intervene, we will face serious consequences.

## **Abstract**

**Importance** When the Covid-19 pandemic enters dangerous new phase, whether and when to take aggressive public health interventions to slow down the spread of COVID-19.

**Objective** To develop the artificial intelligence (AI) inspired methods for real-time forecasting and evaluating intervention strategies to curb the spread of Covid-19 in the World.

**Design, Setting, and Participants** A modified auto-encoder for modeling the transmission dynamics of the epidemics is developed and applied to the surveillance data of cumulative and new Covid-19 cases and deaths from WHO, as of March 16, 2020.

**Interventions and Exposures** Public health interventions and documented novel coronavirus (2019-nCoV)-infected pneumonia (NCIP).

**Main Outcomes and Measures** Data on the number of lab confirmed cumulative, new and death cases of Covid-19 from January 20, 2020 to March 16 were obtained from WHO.

**Results** The average errors of 5-step forecasting were 2.5%. The total peak number of cumulative cases and new cases, and the maximum number of cumulative cases in the world with later intervention (comprehensive public health intervention is implemented 4 weeks later) could reach 75,249,909, 10,086,085, and 255,392,154, respectively. The case ending time was January 10, 2021. However, the total peak number of cumulative cases and new cases and the maximum number of cumulative cases in the world with one week later intervention were reduced to 951,799, 108,853 and 1,530,276, respectively. Duration time of the Covid-19 spread would be reduced from 356 days to 232 days. The case ending time was September 8, 2020. We

observed that delaying intervention for one month caused the maximum number of cumulative cases to increase 166.89 times, and the number of deaths increase from 53,560 to 8,938,725.

**Conclusions and Relevance** Complimentary to other approaches to model the transmission dynamics of the virus outbreaks, the AI-based method provides forecasting tools for tracking, estimating the trajectory of epidemics, assessing their severity, and predicting the lengths of epidemics with various interventions. We estimated the maximum number of cumulative cases under immediate intervention to be 1,530,276; under later intervention the number increased to a frightening 255,392,154 and the case ending time was delayed to January 10, 2021.

## **Introduction**

As of March 16, 2020, global confirmed cases of COVID-19 passed 170,568 and has spread to more than 152 countries. As this coronavirus passes into a pandemic,<sup>1</sup> a number of questions arise among the public and government and business leaders: How many cases will there be worldwide? How many deaths can we expect? When will we see a peak in the number of cases? When will this pandemic end? And, how will recommended immediate actions slow the spread?

A number of the statistical and dynamic models of the Covid-19 outbreak have been applied to analyze its transmission dynamics.<sup>2-8</sup> These epidemiological models are useful for estimating the dynamics of transmission, targeting resources and evaluating the impact of intervention strategies, but the models require values for unknown parameters and depend on many assumptions,<sup>8-10</sup> leading to low accuracy and unsure prediction.

## **Methods**

### **Modified Auto-encoder for Modeling Time Series**

To overcome these limitations, we developed a modified auto-encoder (MAE) (eFigure 1),<sup>11</sup> an AI based method for real time forecasting of the new and cumulative confirmed cases of Covid-19 under various interventions around the world.<sup>12-13</sup> Transfer learning was used to train the MAE.<sup>13</sup> Weights between 0 and 1 were assigned to an intervention variable for the different degrees of interventions – zero being no intervention and one being complete. Complete intervention included keeping social distance, washing hands, strict travel restriction, no large group gatherings, mandatory quarantine, restricted public transportation, keep social distance, school closing and closure of all non-essential business, including manufacturing. Using real data

to evaluate the consequences of specific intervention is infeasible. We considered four intervention scenarios. The first intervention scenario started intervention on March 24, 2020 with a weight of 0.5, one week later transitioned to a complete intervention with a weight of 1. The second intervention started with a weight of 0 for the first week, a weight of 0.5 for the second week, and two weeks later transitioned to a complete intervention with a weight of 1. The third intervention was delaying two weeks, limited in the third week and complete in the fourth week. The fourth intervention delayed actions for three weeks, limited in the fourth weeks and complete in the fifth week. For each scenario, we investigated how the degree and timing of the intervention determined the peak time and case ending time, the peak number and maximum number of cases and forecasting the peak and maximum number of new and cumulative cases in more than 150 countries across the world.

## **Data Collection**

The analysis is based on the surveillance data of confirmed cumulative and new Covid-19 cases in the world up to March 16, 2020. Data on the number of cumulative, new and death cases of Covid-19 from January 20, 2020 to March 16 were obtained from WHO (<https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>). Data included the total numbers of cumulated, newly confirmed and attributed deaths across 152 countries.

## **Results**

### **Later intervention makes it difficult to stop the spread of Covid-19**

To demonstrate that the MAE is an accurate forecasting method, the MAE was applied to confirmed accumulated cases of Covid-19 across 152 countries. The intervention indicator for

China and other countries was set to 1 and 0, respectively. eTable 1 presented the one-step to five-step errors for forecasting the cumulative cases, starting from March 12, 2020. In all cases, the average forecasting accuracies of the MAE were less than 2.5% (eTable 1).

Table 1 shows the forecasting results of Covid-19 in 30 countries and worldwide under later intervention (Scenario 4). The total peak number of cumulative cases and new cases in the world with later intervention could reach 75,249,909 and 10,086,085, respectively. If every country in the world undertook later intervention, the total number of cases in the world could reach as high as 255,392,154 and community transmission of Covid-19 continuing until January 10, 2021. The top 10 countries with a high average increasing number of cases were Italy, Spain, Iran, Germany, USA, France, Switzerland, Belgium, UK and Austria. To show the dynamics of COVID-19 development, eFigure 2 (G) and (H) plot curves of the number of cumulative cases and new cases in seven major infected countries: Iran, Spain, Italy, Germany, USA, France and China under scenario 4.

### **New strategies are needed to curb the spread of COVID-19**

There is urgent need to develop new strategies to curb the spread of COVID-19.<sup>1</sup> We investigate whether complete interventions can control the spread of COVID-19 and how early complete interventions will reduce the peak time and the number, and the final total number of cases across the world. Table 2 shows the forecasted results of COVID-19 in 30 countries and worldwide under early complete intervention (Scenario 1). We can observe dramatic reduction of the cases of COVID-19. The forecasted total number of cases in the world was reduced by early complete intervention to 1,530,276 from nearly 255 million based on 4 weeks delay intervention. In other words, 99.4% of the potential cases would be eliminated by early complete intervention. The duration time was reduced from 356 days to 232 days, and the end time would

change from January 10, 2021 to September 8, 2020. eFigure 2 (A) and (B) plot curves of the number of cumulative cases and new cases in six major infected countries: Iran, Spain, Italy, Germany, USA and France under scenario 1.

To investigate intervention measures between early complete and 4 week delay intervention, eTables 2 and 3 show the results under scenarios 2 and 3, respectively. eFigure 2(C), (D), (E) and (F) plot transmission dynamics of Covid-19 with plotted curves of the cumulative cases and new cases in the six major infected countries under scenarios 2 and 3, respectively.

### **Comparisons among intervention strategies**

To further illustrate the impact of interventions on the spread of COVID-19, we compared the effects of the four intervention scenarios on the transmission dynamics of Covid-19 across the world. Figure 1 plots the world reported and forecasted time curves of the cumulative and newly confirmed cases of Covid-19 under the four intervention scenarios. The ratios of the world number of final cases across the four scenarios were 1:4.26:19.16:166.9 and ratios of case duration under the four intervention scenarios were 1:1.01:1.38:1.53. These results demonstrate that intervention time delays have serious consequences. 4 week delay intervention may increase the number of cases by 166.9 fold and will delay the ending time by almost four months. Delaying intervention will substantially speed the spread of coronavirus.

e Figure 3 plots the time-case curves for the top six infected countries: Iran, Spain, Italy, Germany, USA, France and China. The time-case curve under the 4 week delay intervention was shifted more than one month to the right and was much steeper than that of under the early intervention. Delaying intervention will substantially increase the number of cumulative cases of Covid-19.



eFigure 4 shows the case-fatality rate curve as a function of the Date where the case-fatality rate was defined as the ratio of the number of deaths over the number of cumulative cases in the world. The average case-fatality rate was 3.5%.

## **Discussion**

Real-time forecasting are more accurate than epidemiologic transmission model where the model parameters may not applicable in practice. We estimated the duration, peak time and ending time, peak number and maximum number of cumulative cases of Covid-19 under four intervention scenarios for 152 countries in the world. This provided critical information for government and health authorities to consider urgent public health response planning to slow the spread of Covid-19. We demonstrated that aggressive interventions are urgently needed. Otherwise, we will face disastrous consequences.

## References

1. Callaway E, Time to use the p-word? Coronavirus enters dangerous new phase, Nature (2020) doi: 10.1038/d41586-020-00551-1
2. Li Q, Guan X, Wu P, et al. Early transmission dynamics in Wuhan, China, of novel Coronavirus-infected pneumonia. N Engl J Med. Jan 29. doi: 10.1056/NEJMoa2001316. [Epub ahead of print].
3. Wu JT, Leung K, Leung GM. Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. Lancet. 2020; 395(10225):689-697.
4. Zhao S, Musa SS, Lin Q et al. Estimating the unreported number of novel Coronavirus (2019-nCoV) cases in China in the first half of January 2020: A data-driven modelling analysis of the early outbreak. J Clin Med. 2020 Feb 1;9(2). pii: E388. doi: 10.3390/jcm9020388.
5. Kucharski A, Russell T, Diamond C, Liu Y, CMMID nCoV working group, Edmunds J, Funk S, Eggo R. Analysis and projections of transmission dynamics of nCoV in Wuhan (2020) [https://cmmid.github.io/ncov/wuhan\\_early\\_dynamics/index.html](https://cmmid.github.io/ncov/wuhan_early_dynamics/index.html).
6. Tuite AR, Fisman DN. Reporting, epidemic growth, and reproduction numbers for the 2019 novel coronavirus (2019-nCoV) epidemic. Ann Intern Med. 2020 Feb 5. doi: 10.7326/M20-0358. [Epub ahead of print].
7. Hellewell J, Abbott S, Gimma A, et al. Centre for the Mathematical Modelling of Infectious Diseases COVID-19 Working Group, Funk S1, Eggo RM2. Feasibility of controlling COVID-19 outbreaks by isolation of cases and contacts. Lancet Glob Health. 2020 Feb 28. pii: S2214-109X(20)30074-7. doi: 10.1016/S2214-109X(20)30074-7. [Epub ahead of print].
8. Li R, Pei S, Chen B, Song Y, Zhang T, Yang W, Shaman J. Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV2). Science. 2020 Mar 16. pii: eabb3221. doi: 10.1126/science.abb3221. [Epub ahead of print].
9. Funk S, Camacho A, Kucharski AJ, Eggo RM, Edmunds WJ. Real-time forecasting of infectious disease dynamics with a stochastic semi-mechanistic model. Epidemics. 2018; 22:56-61.

10. Johansson MA, Apfeldorf KM, Dobson S et al. An open challenge to advance probabilistic forecasting for dengue epidemics. *Proc Natl Acad Sci U S A*. 2019;116(48):24268-24274.
11. Charte D, Charte F, García S, Jesus MJD, Herrera F. A practical tutorial on autoencoders for nonlinear feature fusion: Taxonomy, models, software and guidelines. *Information Fusion*. 2019; 44:78–96.
12. Yuan X, Huang B, Wang Y, Yang C, Gui W. Deep Learning-based feature representation and its application for soft sensor modeling with variable-wise weighted SAE. *IEEE Trans on Industrial informatics*. 2018; 14(7): 3235-3243.
13. F. Zhuang Z. Qi K. Duan D et al. He. A Comprehensive Survey on Transfer Learning. *arXiv:1911.02685*. (2019)

## Legend

**Figure 1.** The reported and forecasted curves of the cumulative and new confirmed cases of Covid-19 in the world as a function of days from January 20, 2020 to July 28, 2020.




Table 1. Spread of Covid-19 in 30 countries and world wide under 4 weeks delay intervention.

State	Peak Time	End Time	Duration	Peak (Cum)	Peak (New)	Current Case	End Case
Total	4/17/2020	1/10/2021	356	75249909	10086085	170568	255392154
ITALY	4/17/2020	1/10/2021	346	14945480	1999429	24747	53281848
SPAIN	4/17/2020	1/10/2021	345	10080564	1351788	7753	33196999
IRAN	4/17/2020	1/6/2021	322	8556153	1146663	14991	27343905
GERMANY	4/17/2020	1/10/2021	349	6532219	875856	4838	21864400
USA	4/17/2020	1/10/2021	356	4532725	607493	4740	16644849
FRANCE	4/17/2020	1/10/2021	352	4263429	572051	5380	14555999
SWITZERLAND	4/17/2020	1/10/2021	320	3092785	414952	2200	9772913
BELGIUM	4/17/2020	1/5/2021	336	2835657	380783	1085	8727195
UK	4/17/2020	1/10/2021	345	1624266	218542	1395	6349494
AUSTRIA	4/17/2020	1/10/2021	320	1156505	156173	959	4206694
NORWAY	4/17/2020	1/10/2021	319	1214800	163068	1077	3894919
MALAYSIA	4/17/2020	1/10/2021	351	1081414	144904	553	3750555
GREECE	4/17/2020	11/4/2020	252	1047665	141301	331	3595859
NETHERLANDS	4/17/2020	1/10/2021	318	881147	118402	1135	3080802
PORTUGAL	4/17/2020	1/10/2021	314	675964	91093	245	2104149
FINLAND	4/17/2020	1/10/2021	347	578886	77668	267	1923049
ESTONIA	4/17/2020	1/10/2021	319	607872	81796	205	1902652
SLOVENIA	4/17/2020	1/10/2021	312	598294	80475	219	1891314
ISRAEL	4/17/2020	1/10/2021	324	526864	71296	200	1867519
CANADA	4/17/2020	1/10/2021	350	480352	64450	304	1792760
CZECHIA	4/17/2020	1/8/2021	313	500323	67284	298	1708210
ICELAND	4/17/2020	1/10/2021	315	438161	59381	138	1570527
ROMANIA	4/17/2020	1/10/2021	319	383176	51910	158	1389549
QATAR	4/17/2020	1/10/2021	316	428531	57690	401	1245999
BRAZIL	4/17/2020	1/6/2021	315	374378	50246	200	1218993
AUSTRALIA	4/17/2020	1/10/2021	352	353747	47491	298	1190874

KOREA	4/17/2020	10/30/2020	284	296036	38849	8236	1019408
POLAND	4/17/2020	1/5/2021	308	287008	38725	150	985182

Table 2. Spread of Covid-19 in 30 countries and world wide under early complete intervention (one week from March 16 intervention).

State	Peak Time	End Time	Duration	Peak (Cum)	Peak (New)	Current Case	End Case
Total	2020/3/28	2020/9/8	232	951799	108853	170568	1530276
ITALY	2020/3/27	2020/9/8	222	161276	19998	24747	261790
SPAIN	2020/3/28	2020/8/20	202	117400	15268	7753	187157
IRAN	2020/3/27	2020/6/14	116	95104	12039	14991	157269
GERMANY	2020/3/28	2020/7/22	177	73998	9933	4838	129654
USA	2020/3/27	2020/6/6	138	47058	6454	4740	83921
CHINA	2020/2/5	2020/4/29	100	31432	5236	81077	83103
FRANCE	2020/3/27	2020/8/2	191	45186	5933	5380	81593
SWITZERLAND	2020/3/28	2020/9/6	194	34665	5031	2200	61734
BELGIUM	2020/3/28	2020/6/4	121	29479	4487	1085	52925
UK	2020/3/28	2020/6/2	123	19348	2467	1395	31006
NORWAY	2020/3/28	2020/6/22	117	13631	1985	1077	26386
AUSTRIA	2020/3/29	2020/6/5	101	14394	1825	959	24550
GREECE	2020/3/29	2020/6/10	105	13525	1922	331	22467
MALAYSIA	2020/3/28	2020/7/4	161	11271	1705	553	20985
NETHERLANDS	2020/3/26	2020/5/13	76	8097	1232	1135	16080
KOREA	2020/2/29	2020/5/22	123	3150	813	8236	15649
PORTUGAL	2020/3/30	2020/6/2	92	8578	1157	245	14841
FINLAND	2020/3/30	2020/7/22	175	7707	1037	267	13817
ESTONIA	2020/3/30	2020/6/21	116	7928	1036	205	13382
SLOVENIA	2020/3/28	2020/6/28	116	5856	958	219	12717
ISRAEL	2020/3/29	2020/6/30	130	5865	878	200	10838
ICELAND	2020/3/29	2020/6/23	114	4854	800	138	10679
CZECHIA	2020/3/28	2020/6/20	111	5653	772	298	9586
CANADA	2020/3/28	2020/7/8	164	5330	739	304	9282

QATAR	2020/3/29	2020/6/11	103	4794	652	401	8206
ROMANIA	2020/3/29	2020/5/21	85	4158	627	158	7754
AUSTRALIA	2020/3/28	2020/6/13	141	4117	585	298	7430
BRAZIL	2020/3/28	2020/6/11	106	4017	584	200	7162
DENMARK	2020/3/12	2020/6/19	114	615	353	898	6083



## Materials and Methods

### Data Sources

Data on the confirmed, new and death cases of Covid-19 from January 20, 2020 to March 16 were from WHO (<https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>). Data included the total numbers of accumulated, new confirmed and death cases in the world and across 152 countries. The data were organized in a matrix with the rows representing the whole world and countries and columns representing the number of the new confirmed cases each day. The confirmed cases of each country were a time series. Let  $t_{ij}$  be the number of the confirmed cases of the  $j^{th}$  day within the  $i^{th}$  country. Let  $Z$  be a  $I \times m$  dimensional matrix. The element  $Z_{ij}$  is the number of the confirmed new cases of Covid-19 on the  $j^{th}$  day, starting with January 20, 2020 in the  $i^{th}$  country.

### Modified Auto-encoder for Modeling Time Series

Modified auto-encoders (MAE) (1,2) were used to forecast the number of the accumulative and new confirmed cases of Covid-19. Unlike the classical auto-encoder where the number of nodes in the layers usually decreases from the input layer to the latent layers, the numbers of the nodes in the input, the first latent layer, the second latent layer and output layers in the MSAE were 8, 32, 4 and 1, respectively (eFigure 1). We view a segment of time series with 8 days as a sample of data and take 128 segments of time series as the training samples. One element from the data matrix  $Z$  is randomly selected as a start day of the segment and select its 7 successive days as the other days to form a segment of time series. Let  $i$  be the index of the segment and  $j_i$  be the column index of the matrix  $Z$  that was selected as the starting day. The  $i^{th}$  segment time series

can be represented as  $\{Z_{j_i}, Z_{j_i+1}, \dots, Z_{j_i+7}\}$ . Data were normalized to  $X_{j_i+k} = \frac{Z_{j_i+k}}{S}$ ,  $k = 0, 1, \dots, 7$ , where  $S = \frac{1}{8} \sum_{k=0}^7 Z_{j_i+k}$ . Let  $Y_i = \frac{Z_{j_i+8}}{S}$  be the normalized number of cases to forecast. If  $S = 0$ , then set  $Y_i = 0$ . The loss function was defined as

$$L = \sum_{i=1}^{128} W_i (Y_i - \hat{Y}_i)^2,$$


where  $Y_i$  was the observed number of the cases in the forecasting day of the  $i^{th}$  segment time series and  $\hat{Y}_i$  was its forecasted number of cases by the MAE, and  $W_i$  were weights. If  $j_i$  was in the interval  $[1, 12]$ , then  $W_i = 1$ . If  $j_i$  was in the interval  $[13, 24]$ , then  $W_i = 2$ , etc. The back propagation algorithm was used to estimate the weights and bias in the MSAE. Repeat training processed 5 times. The average forecasting  $\hat{Y}_i, i = 1, \dots, 152$  will be taken as a final forecasted number of the accumulated confirmed cases for each country.

### Forecasting Procedures


The trained MAE was used for forecasting the future number of cumulative cases of Covid-19 for each country. Consider the  $i^{th}$  country. Assume that the number of new confirmed cases of Covid-19 on the  $j^{th}$  day that needs to be forecasted is  $x_{ij}$ . Let  $H$  be a  $152 \times 8$  dimensional matrix and  $h_{il} = x_{ij-9+l}, i = 1, \dots, 152$ , and  $l = 1, \dots, 8$ . Let  $g_i = \frac{1}{8} \sum_{l=1}^8 h_{il}, i = 1, \dots, 152$  be the average of the  $i^{th}$  row of the matrix  $H$ . Let  $U$  be the normalized matrix of  $H$  where  $u_{il} = \frac{h_{il}}{g_i}, i = 1, \dots, 152$ , and  $l = 1, \dots, 8$ . The output of the MAE is the forecasted number of new confirmed cases and is denoted as  $\hat{v}_i = f(u_{i1}, \dots, u_{i8}, \theta), i = 1, \dots, 152$ , where  $\theta$  represented the estimated parameters in the trained MAE. The one-step forecasting of the number of new confirmed cases of Covid-19 for each country is given by  $\hat{Y}_i = \hat{v}_i g_i, i = 1, \dots, 152$ .

The recursive multiple-step forecasting involved using a one-step model multiple times where the prediction for the preceding time step was used as an input for making a prediction on the following time step. For example, for forecasting the number of new confirmed cases for the one more next day, the predicted number of new cases in one-step forecasting would be used as an observational input in order to predict day 2. Repeat the above process to obtain the two-step forecasting. The summation of the final forecasted number of new confirmed cases for each country was taken as the prediction of the total number of new confirmed cases of Covid-19 worldwide.


**eFigure 1** Architecture of modified autoencoder.




**eFigure 2** Trajectory of COVID-19 in the seven most infected countries: Iran, Spain, Italy, Germany, USA, France and China as a function of days from January 21 to June 19, 2020. (A), (C), (E) and (G) Forecasted curves of the newly confirmed cases of COVID-19 under scenarios 1, 2, 3 and 4, respectively. (B), (D), (F) and (H) Forecasted curves of the cumulative confirmed cases of COVID-19 under scenarios 1, 2, 3 and 4, respectively.



**eFigure 3** Time-case plot of the top seven infected countries: Iran, Spain, Italy, Germany, USA, France and China. (A) Time-case plot under intervention scenario 1; (B) Time-case plot under intervention scenario 2; (C) Time-case plot under intervention scenario 3 and (D) Time-case plot under intervention scenario 4.



**eFigure 4** Case-fatality curve for the world.



eTable 1. One- to five-step forecasting errors.

	<b>Reported</b>	<b>1-step predicted</b>	<b>1-step errors</b>	<b>2-step error</b>	<b>3-step error</b>	<b>4-step error</b>	<b>5-step error</b>
<b>3/12/2020</b>	125774	126272	0.40%				
<b>3/13/2020</b>	133774	130278	-2.61%	0.32%			
<b>3/14/2020</b>	143864	144715	0.59%	-3.58%	-0.10%		
<b>3/15/2020</b>	155618	153628	-1.28%	2.51%	-4.08%	0.03%	
<b>3/16/2020</b>	170568	163932	-3.89%	-2.02%	2.26%	-4.80%	0.34%
	<b>Average Absolute Error</b>		1.75%	2.11%	2.15%	2.42%	0.34%



eTable 2. Spread of Covid-19 in 30 countries and world wide under two weeks delay intervention.

State	Peak Time	End Time	Duration	Peak (Cum)	Peak (New)	Current Case	End Case
Total	4/3/2020	9/11/2020	235	3657852	493023	170568	6522982
ITALY	4/3/2020	9/8/2020	222	727996	96948	24747	1307179
SPAIN	4/3/2020	8/20/2020	202	477245	64939	7753	852807
IRAN	4/3/2020	7/23/2020	155	413873	54653	14991	710755
GERMANY	4/3/2020	7/23/2020	178	309434	43000	4838	549478
USA	4/3/2020	9/2/2020	226	216943	29222	4740	381178
FRANCE	4/3/2020	8/18/2020	207	204820	27442	5380	363355
SWITZERLAND	4/3/2020	9/6/2020	194	145504	20435	2200	257680
BELGIUM	4/3/2020	7/3/2020	150	132216	19337	1085	237907
UK	4/3/2020	9/11/2020	224	77356	10561	1395	138340
NORWAY	4/3/2020	6/22/2020	117	57561	8410	1077	105766
AUSTRIA	4/4/2020	7/23/2020	149	62095	7971	959	103793
MALAYSIA	4/4/2020	7/4/2020	161	57868	7303	553	93940
CHINA	2/5/2020	6/6/2020	138	31432	5236	81077	91305
GREECE	4/3/2020	7/20/2020	145	48448	6958	331	88863
NETHERLANDS	4/3/2020	8/2/2020	157	42387	5724	1135	75051
PORTUGAL	4/3/2020	6/22/2020	112	30116	4706	245	59791
SLOVENIA	4/3/2020	7/15/2020	133	27465	4190	219	56030
ESTONIA	4/3/2020	7/20/2020	145	27602	4221	205	55039
FINLAND	4/4/2020	7/22/2020	175	31047	4302	267	53472
ISRAEL	4/4/2020	6/30/2020	130	27678	3705	200	45801
CZECHIA	4/3/2020	6/20/2020	111	23470	3259	298	41366
CANADA	4/3/2020	7/8/2020	164	22704	3198	304	40782
ICELAND	4/4/2020	6/23/2020	114	22483	3087	138	37996
BRAZIL	4/3/2020	7/9/2020	134	17542	2509	200	34112
ROMANIA	4/4/2020	6/15/2020	110	19969	2759	158	33605
QATAR	4/3/2020	6/11/2020	103	18725	2701	401	33116

KOREA	4/5/2020	5/31/2020	132	24100	1873	8236	31670
AUSTRALIA	4/3/2020	7/3/2020	161	16648	2310	298	29334
POLAND	4/3/2020	6/13/2020	102	13287	1908	150	24239
INDONESIA	4/4/2020	7/28/2020	149	12137	1811	117	23177

---

eTable 3. Spread of Covid-19 in top 30 countries and world wide under three weeks delay intervention.

State	Peak Time	End Time	Duration	Peak (Cum)	Peak (New)	Current Case	End Case
Total	4/10/2020	12/4/2020	319	16528763	2221889	170568	29313739
ITALY	4/10/2020	9/8/2020	222	3278431	439028	24747	5693059
SPAIN	4/10/2020	8/20/2020	202	2206610	297488	7753	3919623
IRAN	4/10/2020	8/21/2020	184	1882888	252756	14991	3360378
GERMANY	4/10/2020	8/31/2020	217	1426977	192760	4838	2521231
USA	4/10/2020	10/10/2020	264	992158	133015	4740	1801181
FRANCE	4/10/2020	9/23/2020	243	933029	124787	5380	1674855
SWITZERLAND	4/10/2020	9/6/2020	194	677240	92147	2200	1210668
BELGIUM	4/10/2020	12/4/2020	304	620500	85031	1085	1114935
UK	4/10/2020	9/11/2020	224	354043	47385	1395	639280
NORWAY	4/10/2020	8/2/2020	158	266353	36481	1077	477043
AUSTRIA	4/10/2020	7/23/2020	149	251230	34028	959	446229
MALAYSIA	4/10/2020	8/7/2020	195	235384	32064	553	417237
GREECE	4/10/2020	8/23/2020	179	227442	30714	331	404882
NETHERLANDS	4/10/2020	11/8/2020	255	192602	25705	1135	353763
PORTUGAL	4/10/2020	7/5/2020	125	147451	20653	245	263904
ESTONIA	4/10/2020	8/4/2020	160	132648	18392	205	239417
SLOVENIA	4/10/2020	9/1/2020	181	130582	18060	219	236385
FINLAND	4/10/2020	8/30/2020	214	125404	17387	267	221710
ISRAEL	4/10/2020	8/13/2020	174	113848	15554	200	203392
CZECHIA	4/10/2020	7/7/2020	128	109112	14697	298	194123
CANADA	4/10/2020	10/17/2020	265	104602	14127	304	184096
QATAR	4/10/2020	8/19/2020	172	94166	13163	401	172432
ICELAND	4/10/2020	7/20/2020	141	94695	13174	138	167504
ROMANIA	4/10/2020	9/17/2020	204	82758	11451	158	147628
BRAZIL	4/10/2020	7/10/2020	135	81666	11128	200	145770
AUSTRALIA	4/10/2020	7/6/2020	164	77308	10457	298	138540

CHINA	2/5/2020	6/18/2020	150	31432	5236	81077	127241
KOREA	4/10/2020	7/29/2020	191	70196	8899	8236	123488
POLAND	4/10/2020	8/24/2020	174	62393	8447	150	111894
EGYPT	4/10/2020	8/10/2020	178	58876	8110	126	106174

