

ROCnReg: An R Package for Receiver Operating Characteristic Curve Inference with and without Covariate Information

María Xosé Rodríguez-Álvarez

BCAM - Basque Center for Applied Mathematics
and IKERBASQUE, Basque Foundation for Science

Vanda Inácio

University of Edinburgh

Abstract

The receiver operating characteristic (ROC) curve is the most popular tool used to evaluate the discriminatory capability of diagnostic tests/biomarkers measured on a continuous scale when distinguishing between two alternative disease states (e.g, diseased and nondiseased). In some circumstances, the test's performance and its discriminatory ability may vary according to subject-specific characteristics or different test settings. In such cases, information-specific accuracy measures, such as the covariate-specific and the covariate-adjusted ROC curve are needed, as ignoring covariate information may lead to biased or erroneous results. This paper introduces the R package **ROCnReg** that allows estimating the pooled (unadjusted) ROC curve, the covariate-specific ROC curve, and the covariate-adjusted ROC curve by different methods, both from (semi) parametric and nonparametric perspectives and within Bayesian and frequentist paradigms. From the estimated ROC curve (pooled, covariate-specific or covariate-adjusted), several summary measures of discriminatory accuracy, such as the (partial) area under the ROC curve and the Youden index, can be obtained. The package also provides functions to obtain ROC-based optimal threshold values using several criteria, namely, the Youden index criterion and the criterion that sets a target value for the false positive fraction. For the Bayesian methods, we provide tools for assessing model fit via posterior predictive checks, while model choice can be carried out via several information criteria. Numerical and graphical outputs are provided for all methods. The package is illustrated through the analyses of data from an endocrine study where the aim is to assess the capability of the body mass index to detect the presence or absence of cardiovascular disease risk factors. The package is available from the Comprehensive R Archive Network (CRAN) at <https://CRAN.R-project.org/package=ROCnReg>.

Keywords: accuracy measures, Bayesian, diagnostic tests, ROC curve, covariate-specific ROC curve, covariate-adjusted ROC curve, optimal thresholds, R.

1. Introduction

Before a diagnostic test is approved for being routinely used in practice, its ability to distinguish say, diseased from nondiseased individuals, must be narrowly evaluated. Throughout we assume that the true disease status of the individuals is known and the task is, compared to the truth, to quantify how accurate the test being investigated is. Before proceeding, it is worth noting that although our focus is on medical diagnosis, the problem of binary classifi-

cation is a much wider one, finding applications in fields as diverse as biology and finance, to name only two.

The receiver operating characteristic (ROC) curve (Metz 1978) is, unarguably, the most popular tool used for evaluating the discriminatory ability of continuous-outcome diagnostic tests. The ROC curve displays the false positive fraction (FPF) against the true positive fraction (TPF) for all possible threshold values used to dichotomise the test result. The ROC curve thus provides a global description of the trade-off between the FPF and the TPF of the test as the threshold changes. Plenty of parametric and semi/nonparametric methods are available for estimating ROC curves, either from frequentist or Bayesian viewpoints and we refer the interested reader to Pepe (1998, Chapter 5), Zhou, McClish, and Obuchowski (2011, Chapter 4), Gonçalves, Subtil, Oliveira, and Bermudez (2014) and Inácio, Rodríguez-Álvarez, and Gayoso-Diz (2020), and references therein.

However, it is known that in many situations, a test's outcome and, possibly, its discriminatory capacity, can be affected by additional information (covariates); Pepe (2003, pp 48–49) provides several examples of covariates that can affect the result of a diagnostic test. For instance, patient characteristics, such as age and gender, are important covariates to be considered as diagnostic accuracy is likely to vary according to them. In these cases pooling the test outcomes regardless of their covariate values may lead to erroneous or, at least, oversimplified conclusions and decisions. Interest should therefore be focused on assessing the accuracy of the test, but taking into account covariate information. Two different ROC-based measures that incorporate covariate information have been proposed: the covariate-specific or conditional ROC curve (see, e.g., Pepe 2003, Chapter 6) and the covariate-adjusted ROC curve (Janes and Pepe 2009). The formal definition of both curves is given in Section 2. In brief, a covariate-specific ROC curve is an ROC curve that conditions on a specific covariate value, thus describing the accuracy of the test in the 'subpopulation' defined by that covariate value. On the other hand, the covariate-adjusted ROC curve is a weighted average of covariate-specific ROC curves. Regarding estimation, since the seminal paper of Pepe (1998), a plethora of methods have been proposed in the literature for the estimation of the covariate-specific ROC curve and associated summary measures. Without being exhaustive, we mention the work of Faraggi (2003), Rodríguez-Álvarez, Roca-Pardiñas, and Cadarso-Suárez (2011b,a), Inácio de Carvalho, Jara, Hanson, and de Carvalho (2013), and Inácio de Carvalho, de Carvalho, and Branscum (2017). A detailed review can be found in Rodríguez-Álvarez, Tahoces, Cadarso-Suárez, and Lado (2011c), Pardo-Fernández, Rodríguez-Álvarez, and van Keilegom (2014) and Inácio *et al.* (2020). With respect to the covariate-adjusted ROC curve, estimation has been discussed in Janes and Pepe (2009), Rodríguez-Álvarez *et al.* (2011b), Guan, Qin, and Zhang (2012), and Inácio de Carvalho and Rodríguez-Álvarez (2018).

In a slightly different context, in the machine learning community, the topic of covariate-dependent classification has received much attention recently, being related to the concept of *fairness* (see, e.g., Hutchinson and Mitchell 2019). As an example, Buolamwini (2017) reports that the evaluation of four gender classifiers revealed that a significant gap exists when comparing gender classification accuracies of females versus males.

There are a few R (R Core Team 2020) packages for ROC curve analysis available on the Comprehensive R Archive Network (CRAN) and, as far as we are aware, all of them implementing frequentist approaches. The package **sROC** (Wang 2012) contains functions to perform nonparametric, kernel-based, estimation of ROC curves, while **pROC** (Robin,

Turck, Hainard, Tiberti, Lisacek, Sanchez, and Müller 2011) offers a set of tools to visualise, smooth, and compare ROC curves, and **nsROC** (Pérez Fernández, Martínez Cambior, Filzmoser, and Corral Blanco 2018) also allows estimating ROC curves, building confidence bands, as well as comparing several curves both for dependent and independent data (i.e., data arising from paired and unpaired designs, respectively). However, covariate information cannot be explicitly taken into account in any of these packages. Packages **ROCRegression** (available at <https://bitbucket.org/mxrodriguez/rocgression>) and **npROCRegression** (Rodríguez-Alvarez and Roca-Pardinas 2017) provide routines to estimate semi-parametrically and nonparametrically, under a frequentist framework, the covariate-specific ROC curve. We also mention **OptimalCutpoints** (López-Ratón, Rodríguez-Álvarez, Cadarso-Suárez, and Gude-Sampedro 2014) and **ThresholdROC** (Perez Jaume, Skaltsa, Pallarès, and Carrasco Jordan 2017) that provide a collection of functions for point and interval estimation of optimal thresholds for continuous diagnostic tests. To the best of our knowledge, there is no statistical software package implementing Bayesian inference for ROC curves and associated summary indices and optimal thresholds.

To close this gap, in this paper we introduce the **ROCnReg** package that allows conducting Bayesian inference for the (pooled or marginal) ROC curve, the covariate-specific ROC curve, and the covariate-adjusted ROC curve. For the sake of generality, frequentist approaches are also implemented. Specifically, in what concerns estimation of the pooled ROC curve, **ROCnReg** implements the frequentist empirical estimator described in Hsieh and Turnbull (1996), the kernel-based approach proposed of Zou, Hall, and Shapiro (1997), the Bayesian Bootstrap method of Gu, Ghosal, and Roy (2008), and the Bayesian nonparametric method based on a Dirichlet process mixture of normal distributions model proposed by Erkanli, Sung, Jane Costello, and Angold (2006). Regarding the covariate-specific ROC curve, **ROCnReg** implements the frequentist normal method of Faraggi (2003) and its semiparametric counterpart as described in Pepe (1998), the kernel-based approach of Rodríguez-Álvarez *et al.* (2011b), and the Bayesian nonparametric model, based on a single-weights dependent Dirichlet process mixture of normal distributions, proposed by Inácio de Carvalho *et al.* (2013). As for the covariate-adjusted ROC curve, the **ROCnReg** package allows estimation using the frequentist semiparametric approach of Janes and Pepe (2009), the frequentist nonparametric method discussed in Rodríguez-Álvarez *et al.* (2011b), and the recently proposed Bayesian nonparametric estimator of Inácio de Carvalho and Rodríguez-Álvarez (2018). Table 1 shows a summary of all methods implemented in the package. In addition, **ROCnReg** also provides functions to obtain ROC-based optimal thresholds to perform the classification/diagnosis using two different criteria, namely, the Youden index and the criterion that sets a target value for the false positive fraction. These are implemented for both the ROC curve, the covariate-specific and the covariate-adjusted ROC curve. A detailed description of the methods is presented in Section 3.

The remainder of the paper is organised as follows. In Section 2 we formally introduce the (pooled or marginal) ROC curve, the covariate-specific ROC curve, and the covariate-adjusted ROC curve. The description of the estimation methods implemented in the **ROCnReg** package is given in Section 3. In Section 4 the usage of the main functions and capabilities of **ROCnReg** are described and illustrated using a real example. The paper concludes with a discussion in Section 5.

Method	Description
Pooled ROC curve	
emp	(Frequentist) empirical estimator (Hsieh and Turnbull 1996).
kernel	(Frequentist) kernel-based approach (Zou <i>et al.</i> 1997).
BB	Bayesian bootstrap method (Gu <i>et al.</i> 2008).
dpm	Nonparametric Bayesian approach based on a Dirichlet process mixture of normal distributions (Erkanli <i>et al.</i> 2006).
Covariate-specific ROC curve	
sp	(Frequentist) parametric and semiparametric induced ROC regression approach (Pepe 1998; Faraggi 2003)
kernel	Nonparametric (kernel-based) induced ROC regression approach (Rodríguez-Álvarez <i>et al.</i> 2011b).
bnp	Nonparametric Bayesian model based on a single-weights dependent Dirichlet process mixture of normal distributions (Inácio de Carvalho <i>et al.</i> 2013).
Covariate-adjusted ROC curve	
sp	(Frequentist) semiparametric method (Janes and Pepe 2009).
kernel	Nonparametric (kernel-based) induced ROC regression approach (Rodríguez-Álvarez <i>et al.</i> 2011b).
bnp	Nonparametric Bayesian model based on a single-weights dependent Dirichlet process mixture of normal distributions and the Bayesian bootstrap (Inácio de Carvalho and Rodríguez-Álvarez 2018).

Table 1: Overview of ROC estimation methods included in the **ROCnReg** package.

2. Notation and definitions

This section sets out the formal definition of the pooled or marginal ROC curve, the covariate-specific ROC curve, and the covariate-adjusted ROC curve. Also, it describes the most commonly used summary measures of discriminatory accuracy, namely, the area under the ROC curve, the partial area under the ROC curve, and the Youden index. For conciseness, we intentionally avoid giving too many details and refer the interested reader to Pepe (2003) (and references therein) for an extensive account of many aspects of ROC curves with and without covariates.

In what follows, we denote as Y the outcome of the diagnostic test and as D the binary variable indicating the presence ($D = 1$) or absence ($D = 0$) of disease. We also assume that along with Y and the true disease status D , a covariate vector \mathbf{X} is also available, and that it may encompass both continuous and categorical covariates. For ease of notation, the covariate vector \mathbf{X} is assumed to be the same in both the diseased ($D = 1$) and nondiseased ($D = 0$) populations, although this is not always necessarily the case in practice (e.g., disease stage is, obviously, a disease-specific covariate). By a slight abuse of notation, we use the subscripts D and \bar{D} to denote (random) quantities conditional on, respectively, $D = 1$ and $D = 0$. For example, Y_D and $Y_{\bar{D}}$ denote the test outcomes in the diseased and nondiseased populations.

2.1. Pooled ROC curve

In the case of a continuous-outcome diagnostic test, the classification is usually made by comparing the test result Y against a threshold c . If the outcome is equal or above the threshold, $Y \geq c$, the subject will be considered as diseased. On the other hand, if the test result is below the threshold, $Y < c$, he/she will be classified as nondiseased. The ROC curve is then defined as the set of all possible false positive fractions, $\text{FPF}(c) = \text{P}(Y \geq c \mid D = 0) = \text{P}(Y_{\bar{D}} \geq c)$, and true positive fractions, $\text{TPF}(c) = \text{P}(Y \geq c \mid D = 1) = \text{P}(Y_D \geq c)$, that can be obtained by varying the threshold value c , i.e.,

$$\{(\text{FPF}(c), \text{TPF}(c)) : c \in \mathbb{R}\}.$$

It is common to represent the ROC curve as $\{(p, \text{ROC}(p)) : p \in [0, 1]\}$, where

$$p = \text{FPF}(c) = 1 - F_{\bar{D}}(c), \quad \text{ROC}(p) = 1 - F_D\{F_{\bar{D}}^{-1}(1 - p)\}, \quad (1)$$

with $F_D(y) = \text{P}(Y_D \leq y)$ and $F_{\bar{D}}(y) = \text{P}(Y_{\bar{D}} \leq y)$ denoting the cumulative distribution function (CDF) of Y in the nondiseased and diseased groups, respectively. Several indices can be used as global summary measures of the accuracy of a test. The most widely used is the area under the ROC curve (AUC), defined as

$$\text{AUC} = \int_0^1 \text{ROC}(p) dp. \quad (2)$$

In addition to its geometric definition, the AUC has also a probabilistic interpretation (see, e.g., [Pepe 2003](#), p. 78)

$$\text{AUC} = \text{P}(Y_D \geq Y_{\bar{D}}), \quad (3)$$

that is, the AUC is the probability that a randomly selected diseased subject has a higher test outcome than that of a randomly selected nondiseased subject. The AUC takes values between 0.5, in the case of an uninformative test that classifies individuals no better than chance, and 1.0 for a perfect test. We note that an AUC below 0.5 simply means that the classification rule should be reversed. As it is clear from its definition, the AUC integrates the ROC curve over the whole range of FPFs. Depending on the clinical circumstances, however, interest might lie only on a relevant interval of FPFs or TPFs, which leads to the notion of partial area under the ROC curve (pAUC). The pAUC over a range of FPFs $(0, u_1)$, where u_1 is typically low and represents the largest acceptable FPF, is defined as

$$\text{pAUC}(u_1) = \int_0^{u_1} \text{ROC}(p) dp. \quad (4)$$

On the other hand, the pAUC over a range of TPFs $(v_1, 1)$, where v_1 is typically large and represents the lowest acceptable TPF, is defined as

$$\text{pAUC}_{\text{TPF}}(v_1) = \int_{v_1}^1 \text{ROC}_{\text{TNF}}(p) dp, \quad (5)$$

where ROC_{TNF} is a 270° rotation of the ROC curve, which can be expressed as

$$\text{ROC}_{\text{TNF}}(p) = F_{\bar{D}}\{F_D^{-1}(1 - p)\}. \quad (6)$$




Figure 1: (a) Shaded area in blue represents the partial area under the ROC curve over the interval $(0, u_1)$ of FPFs. (b) Shaded area in blue represents the partial area under the ROC curve over the interval $(v_1, 1)$ of TPFs. (c) The same as in (b) but now represented as an area over the true negative fraction ROC curve.

The curve (6) is referred to as the true negative fraction (TNF) ROC curve, since $\text{TNF} (= 1 - \text{FPF})$ is plotted on the y -axis. We shall highlight that the argument p in the ROC curve stands for a false positive fraction, whereas in the ROC_{TNF} curve it stands for a true positive fraction. In Figure 1 we graphically illustrate the two partial areas.

We shall highlight that the argument p in the ROC curve stands for a false positive fraction, whereas in the ROC_{TNF} curve it stands for a true positive fraction.

Another summary index of diagnostic accuracy is the Youden index (Shapiro 1999; Youden 1950)

$$\text{YI} = \max_c \{ \text{TPF}(c) - \text{FPF}(c) \} \quad (7)$$

$$= \max_c \{ F_{\bar{D}}(c) - F_D(c) \} \quad (8)$$

$$= \max_p \{ \text{ROC}(p) - p \}. \quad (9)$$

The YI ranges from 0 to 1, taking the value of 0 in the case of an uninformative test and 1 for a perfect test. As for the AUC, a YI below 0 means that the classification rule should be reversed. The value c^* which maximises Equation (7) (or, equivalently, Equation (8)) is frequently used in practice to classify subjects as diseased or nondiseased. It should be noted that this index is equivalent to the Kolmogorov–Smirnov measure of distance between the distributions of Y_D and $Y_{\bar{D}}$ (Pepe 2003, p. 80).

2.2. Covariate-specific ROC curve

The conditional or covariate-specific ROC curve, given a covariate value \mathbf{x} , is defined as

$$\text{ROC}(p | \mathbf{x}) = 1 - F_D\{F_{\bar{D}}^{-1}(1 - p | \mathbf{x}) | \mathbf{x}\}, \quad (10)$$

where $F_{\bar{D}}(y | \mathbf{x}) = \Pr(Y_{\bar{D}} \leq y | \mathbf{X}_{\bar{D}} = \mathbf{x})$ and $F_D(y | \mathbf{x}) = \Pr(Y_D \leq y | \mathbf{X}_D = \mathbf{x})$ are the conditional CDFs of the test in the nondiseased and diseased groups, respectively. In this case,

a number of possibly different ROC curves (and therefore accuracies) may be obtained for different values of \mathbf{x} . Thus, the covariate-specific ROC curve is an important tool that helps to understand and determine the optimal and suboptimal populations where to apply the tests on (i.e., it allows determining the populations, defined by or homogeneous with respect to \mathbf{x} , where the diagnostic test has a ‘good’ or ‘poor’ discriminatory capacity). Similarly to the unconditional case, the covariate-specific TNF-ROC curve is given by

$$\text{ROC}_{\text{TNF}}(p | \mathbf{x}) = F_{\bar{D}}\{F_D^{-1}(1 - p | \mathbf{x}) | \mathbf{x}\}, \quad (11)$$

and the covariate-specific AUC, pAUC, and Youden index are

$$\text{AUC}(\mathbf{x}) = \int_0^1 \text{ROC}(p | \mathbf{x}) dp, \quad (12)$$

$$\text{pAUC}(u_1 | \mathbf{x}) = \int_0^{u_1} \text{ROC}(p | \mathbf{x}) dp, \quad (13)$$

$$\text{pAUC}_{\text{TPF}}(v_1 | \mathbf{x}) = \int_{v_1}^1 \text{ROC}_{\text{TNF}}(p | \mathbf{x}) dp, \quad (14)$$

$$\text{YI}(\mathbf{x}) = \max_c |\text{TPF}(c | \mathbf{x}) - \text{FPF}(c | \mathbf{x})| \quad (15)$$

$$= \max_c |F_{\bar{D}}(c | \mathbf{x}) - F_D(c | \mathbf{x})| \quad (16)$$

$$= \max_p |\text{ROC}(p | \mathbf{x}) - p|. \quad (17)$$

The value $c_{\mathbf{x}}^*$ that achieves the maximum in (15) (or (16)) is called the optimal covariate-specific YI threshold and can be used to classify a subject, with covariate value \mathbf{x} , as diseased or nondiseased.

2.3. Covariate-adjusted ROC curve

The covariate-specific ROC curve and associated AUC, pAUCs, and YI described in Section 2.2 depict the accuracy of the test for specific covariate values. However, it would be undoubtedly useful to have a global summary measure that also takes covariate information into account. Such summary measure was developed by [Janes and Pepe \(2009\)](#), who proposed the covariate-adjusted ROC (AROC) curve, defined as

$$\text{AROC}(p) = \int \text{ROC}(p | \mathbf{x}) dH_D(\mathbf{x}), \quad (18)$$

where $H_D(\mathbf{x}) = \text{P}(\mathbf{X}_D \leq \mathbf{x})$ is the CDF of \mathbf{X}_D . That is, the AROC curve is a weighted average of covariate-specific ROC curves, weighted according to the distribution of the covariates in the diseased group. Equivalently, as shown by [Janes and Pepe \(2009\)](#), the AROC curve can also be expressed as

$$\begin{aligned} \text{AROC}(p) &= \text{P}\{Y_D > F_{\bar{D}}^{-1}(1 - p | \mathbf{X}_D)\} \\ &= \text{P}\{1 - F_{\bar{D}}(Y_D | \mathbf{X}_D) \leq p\}. \end{aligned} \quad (19)$$

As will be seen in Section 3, Expression (19) is very convenient when it comes to estimating the AROC curve. Also, it emphasises that the AROC curve at a FPF of p is the overall TPF when the thresholds used for defining a positive test result are covariate-specific and chosen to ensure that the FPF is p in each subpopulation defined by the covariate values.

In contrast to the pooled ROC curve (see Expressions (1) and (6)) and the covariate-specific ROC curve (see Expressions (10) and (11)), the AROC curve (and its 270° rotation) cannot be expressed in terms of the (conditional) CDFs of the test in each group. This does not, however, preclude the possibility of defining AROC-based summary accuracy measures, yet more care is needed. Thus, for the AROC curve, the area under the AROC, as well as the partial areas and YI are expressed as follows

$$\text{AAUC} = \int_0^1 \text{AROC}(p) dp, \quad (20)$$

$$\text{pAAUC}(u_1) = \int_0^{u_1} \text{AROC}(p) dp, \quad (21)$$

$$\text{pAAUC}_{\text{TPF}}(v_1) = \int_{\text{AROC}^{-1}(v_1)}^1 \text{AROC}(p) dp - \{1 - \text{AROC}^{-1}(v_1)\} v_1, \quad (22)$$

$$\text{YI}_{\text{AROC}} = \max_p \{ \text{AROC}(p) - p \}. \quad (23)$$

Note, in particular, that the expressions for both the partial area under the AROC curve over a range of TPFs (see also Figure 1) and for the YI are defined in terms of the AROC curve. For the YI, once the value that achieves the maximum in (23) is calculated, say p^* , covariate-specific threshold values can be obtained as follows

$$c_{\mathbf{x}}^* = F_{\bar{D}}^{-1}(1 - p^* | \mathbf{X}_D = \mathbf{x}).$$

Note that, by construction, these threshold values will ensure that the FPF is p^* in each subpopulation defined by the covariate values; however, the TPF may vary with the covariate values, i.e.,

$$\text{TPF}(c_{\mathbf{x}}^*) = 1 - F_D(c_{\mathbf{x}}^* | \mathbf{X}_D = \mathbf{x}).$$

To finish this part, we mention that when the accuracy of a test is not affected by covariates, this does not necessarily mean that the covariate-specific ROC curve (which in this case is the same for all covariate values) coincides with the pooled ROC curve. It does coincide, however, with the AROC curve (see [Janes and Pepe 2009](#); [Pardo-Fernández et al. 2014](#); [Inácio de Carvalho and Rodríguez-Álvarez 2018](#), for more details). As such, in all cases where covariates affect the results of the test, even though they might not affect its discriminatory capacity, inferences based on the pooled ROC curve might be misleading. In such cases, the AROC curve should be used instead. This also applies to the selection of (optimal) threshold values, which might be covariate-specific (i.e., possibly different for different covariate values).

3. Methods

In this section we describe the different methods for ROC curve inference (with and without covariate information) implemented in the **ROCnReg** package.

3.1. Pooled ROC curve

Let $\{y_{\bar{D}i}\}_{i=1}^{n_{\bar{D}}}$ and $\{y_{Dj}\}_{j=1}^{n_D}$ be two independent random samples of test outcomes from the nondiseased and diseased groups of size $n_{\bar{D}}$ and n_D , respectively.

Empirical estimator

The function `pooledROC.emp` estimates the pooled ROC curve using the empirical estimator proposed by [Hsieh and Turnbull \(1996\)](#), which consists in estimating the CDFs of the test in each group by its empirical counterpart, that is,

$$\hat{F}_{\bar{D}}(y) = \frac{1}{n_{\bar{D}}} \sum_{i=1}^{n_{\bar{D}}} I(y_{\bar{D}i} \leq y), \quad \hat{F}_D(y) = \frac{1}{n_D} \sum_{j=1}^{n_D} I(y_{Dj} \leq y).$$

These empirical estimates are then plugged into Equations (1) and (6) to obtain, respectively, an estimate of the ROC and ROC_{TNF} curves.

In what concerns estimation of the AUC (Expression (3)), pAUC , and pAUC_{TNF} (Expressions (4) and (5), respectively) these are computed empirically by means of the Mann–Whitney U statistic. With respect to the Youden index (and associated threshold value), it is obtained by maximising, over a grid of possible threshold values, the expression in (8), with F_D and $F_{\bar{D}}$ being replaced by their empirical estimators.

Kernel estimator

The function `pooledROC.kernel` estimates the pooled ROC curve using the kernel-based estimator proposed by [Zou et al. \(1997\)](#) and [Zou, Tempany, Fielding, and Silverman \(1998\)](#), which is based on estimating the CDFs of the test as follows

$$\hat{F}_{\bar{D}}(y) = \frac{1}{n_{\bar{D}}} \sum_{i=1}^{n_{\bar{D}}} \Phi\left(\frac{y - y_{\bar{D}i}}{h_{\bar{D}}}\right), \quad \hat{F}_D(y) = \frac{1}{n_D} \sum_{j=1}^{n_D} \Phi\left(\frac{y - y_{Dj}}{h_D}\right),$$

where $\Phi(y)$ stands for the standard normal distribution evaluated at y . For the bandwidths, $h_{\bar{D}}$ and h_D , which control the amount of smoothing, two options are popular. Silverman's rule of thumb ([Silverman 1986](#), p. 48), which sets the bandwidth as

$$h_d = 0.9 \min\{\text{SD}(\mathbf{y}_d), \text{IQR}(\mathbf{y}_d)/1.34\} n_d^{-0.2}, \quad d \in \{\bar{D}, D\},$$

where $\text{SD}(\mathbf{y}_d)$ and $\text{IQR}(\mathbf{y}_d)$ are the standard deviation and interquartile range, respectively, of $\mathbf{y}_d = (y_{d1}, \dots, y_{dn_d})$. Another alternative criterion is to select the bandwidth by using least squares cross-validation ([Wand and Jones 1994](#), Chapter 3).

Here, both the AUC, pAUC , and pAUC_{TNF} (Expressions (2), (4), and (5)) are computed numerically using Simpson's rule. Regarding the Youden index (and associated threshold value), it is obtained by maximising, over a grid of possible threshold values, Expression (8), with F_D and $F_{\bar{D}}$ being replaced by their kernel estimators.

Uncertainty estimation for both the empirical and kernel estimators is conducted through bootstrap resampling.

Bayesian bootstrap estimator

The function `pooledROC.bb` implements the Bayesian bootstrap (BB) approach proposed by [Gu et al. \(2008\)](#). Their estimator relies on the notion of placement value ([Pepe 2003](#), Chapter 5), which is simply a standardisation of the test outcomes with respect to a reference group. Specifically, $U_D = 1 - F_{\bar{D}}(Y_D)$ is to be interpreted as a standardisation of a diseased test

outcome with respect to the distribution of test results in the nondiseased population. The ROC curve can be regarded as the CDF of U_D

$$\mathbb{P}(U_D \leq p) = \mathbb{P}\{1 - F_{\bar{D}}(Y_D) \leq p\} = 1 - F_D\{F_{\bar{D}}^{-1}(1 - p)\} = \text{ROC}(p), \quad 0 \leq p \leq 1. \quad (24)$$

This representation in (24) of the ROC curve provided the rationale for the two-step algorithm of Gu *et al.* (2008), which can be described as follows. Let S be the number of iterations.

Step 1: Computation of the placement value based on the BB.

For $s = 1, \dots, S$, let

$$U_{Dj}^{(s)} = \sum_{i=1}^{n_{\bar{D}}} q_{1i}^{(s)} I(y_{\bar{D}i} \geq y_{Dj}), \quad j = 1, \dots, n_D,$$

where $(q_{11}^{(s)}, \dots, q_{1n_{\bar{D}}}^{(s)}) \sim \text{Dirichlet}(n_{\bar{D}}; 1, \dots, 1)$.

Step 2: Generate a realisation of the ROC curve. Based on (24), generate a realisation of $\text{ROC}^{(s)}(p)$, the cumulative distribution function of $(U_{D1}^{(s)}, \dots, U_{Dn_D}^{(s)})$, where

$$\text{ROC}^{(s)}(p) = \sum_{j=1}^{n_D} q_{2j}^{(s)} I(U_{Dj}^{(s)} \leq p), \quad (q_{21}^{(s)}, \dots, q_{2n_D}^{(s)}) \sim \text{Dirichlet}(n_D; 1, \dots, 1).$$

The BB estimate of the ROC curve is obtained by averaging over the ensemble of ROC curves $\{\text{ROC}^{(1)}(p), \dots, \text{ROC}^{(S)}(p)\}$, that is,

$$\widehat{\text{ROC}}^{\text{BB}}(p) = \frac{1}{S} \sum_{s=1}^S \text{ROC}^{(s)}(p),$$

and a $(1 - \alpha) \times 100\%$ pointwise credible band can be obtained from the $\alpha/2 \times 100\%$ and $(1 - \alpha/2) \times 100\%$ percentiles of the same ensemble ($\alpha \in (0, 1)$). Note that these pointwise credible bands for the ROC curve are to be interpreted as credible intervals for the TPFs.

The Bayesian bootstrap estimator leads to closed-form expressions for the AUC and pAUC, which are, respectively, given by

$$\begin{aligned} \text{AUC}^{(s)} &= \int_0^1 \text{ROC}^{(s)}(p) dp = 1 - \sum_{j=1}^{n_D} q_{2j}^{(s)} U_{Dj}^{(s)}, \\ \text{pAUC}^{(s)}(u_1) &= \int_0^{u_1} \text{ROC}^{(s)}(p) dp = u_1 - \sum_{j=1}^{n_D} q_{2j}^{(s)} \min\{u_1, U_{Dj}^{(s)}\}. \end{aligned}$$

It is easy to show that

$$\text{pAUC}_{\text{TPF}}^{(s)}(v_1) = \int_{v_1}^1 \text{ROC}_{\text{TNF}}^{(s)}(p) dp = \sum_{i=1}^{n_{\bar{D}}} q_{1i}^{(s)} \max\{v_1, U_{\bar{D}i}^{(s)}\} - v_1,$$

where

$$U_{\bar{D}i}^{(s)} = \sum_{j=1}^{n_D} q_{2j}^{(s)} I(y_{Dj} \geq y_{\bar{D}i}), \quad i = 1, \dots, n_{\bar{D}},$$

and it is also easy to demonstrate that the ROC_{TNF} curve is the survival function of the placement value $U_{\bar{D}} = 1 - F_D(Y_{\bar{D}})$. With respect to the Youden index, it is obtained by maximising, over a grid of possible threshold values, the following expression

$$\text{YI}^{(s)} = \max_c \left\{ F_{\bar{D}}^{(s)}(c) - F_D^{(s)}(c) \right\},$$

where

$$F_{\bar{D}}^{(s)}(c) = \sum_{i=1}^{n_{\bar{D}}} q_{1i}^{(s)} I(y_{\bar{D}i} \leq c) \quad \text{and} \quad F_D^{(s)}(c) = \sum_{j=1}^{n_D} q_{2j}^{(s)} I(y_{Dj} \leq c).$$

As for the ROC curve, point estimates for the AUC, pAUC, pAUC_{TPF}, YI, and c^* can be obtained by averaging over the respective ensembles of S realisations, with credible bands derived from the percentiles of such ensembles.

Dirichlet process mixture of normal distributions estimator

The Bayesian nonparametric approach, based on a Dirichlet process mixture (DPM) of normal distributions, for estimating the pooled ROC curve (Erkanli *et al.* 2006) is implemented in the `pooledROC.dpm` function. In this case, as implicit by the name, the CDFs of the test outcomes in each group are estimated via a Dirichlet process mixture of normal distributions, that is, it is assumed that the CDF, say in the diseased group (the one in the nondiseased group, \bar{D} , follows analogously), is of the form

$$F_D(y) = \int \Phi(y \mid \mu, \sigma^2) dG_D(\mu, \sigma^2), \quad G_D \sim \text{DP}(\alpha_D, G_D^*(\mu, \sigma^2)), \quad (25)$$

where $\Phi(y \mid \mu, \sigma^2)$ denotes the CDF of the normal distribution with mean μ and variance σ^2 evaluated at y . Here $G_D \sim \text{DP}(\alpha_D, G_D^*)$ is used to denote that the mixing distribution G_D follows a Dirichlet process (DP) (Ferguson 1973) with centring distribution G_D^* , for which $E(G_D) = G_D^*$, and precision parameter α_D . Usually, due to conjugacy reasons, $G_D^*(\mu, \sigma^2) \equiv N(\mu \mid m_{D0}, S_{D0})\Gamma(\sigma^{-2} \mid a_D, b_D)$ and this is the centring distribution used by the `pooledROC.dpm` function. Note that here S_{D0} denotes the variance of the normal distribution and a_D and b_D are, respectively, the shape and rate parameters of the gamma distribution. All hyperparameter values are fixed.

For ease of posterior simulation and because it provides a highly accurate approximation, we make use of the truncated stick-breaking representation of the DP (Ishwaran and James 2001), according to which G_D can be written as

$$G_D(\cdot) = \sum_{l=1}^{L_D} \omega_{Dl} \delta_{(\mu_{Dl}, \sigma_{Dl}^2)}(\cdot),$$

where $(\mu_{Dl}, \sigma_{Dl}^2) \stackrel{\text{iid}}{\sim} G_D^*(\mu, \sigma^2)$, for $l = 1, \dots, L_D$, and the weights follow the so-called (truncated) stick-breaking construction: $\omega_{D1} = v_{D1}$, $\omega_{Dl} = v_{Dl} \prod_{r < l} (1 - v_{Dr})$, $l = 2, \dots, L_D$, and $v_{D1}, \dots, v_{D, L_D-1} \stackrel{\text{iid}}{\sim} \text{Beta}(1, \alpha_D)$. Further, one must set $v_{DL_D} = 1$ in order to ensure that the weights add up to one. Further, one must set $v_{DL_D} = 1$ in order to ensure that the weights add up to one. With regard to the parameter α_D , in **ROCnReg** a prior distribution is placed on it. In particular, and due to conjugacy reasons, a gamma distribution is considered, i.e.,

$\alpha_D \sim \Gamma(a_{\alpha_D}, b_{\alpha_D})$. The CDF in (25) can therefore be written as

$$F_D(y) = \sum_{l=1}^{L_D} \omega_{Dl} \Phi(y \mid \mu_{Dl}, \sigma_{Dl}^2),$$

where we shall note that L_D is not the exact number of components expected to be observed, but rather an upper bound on it, as some of the components may be unoccupied. Some comments are in order regarding the specification of the hyperparameters' values. In what concerns the centring distribution, m_{D0} represents the prior belief about the components' means and S_{D0} represents the confidence in such prior belief. Similarly, the values of a_D and b_D can be chosen to represent prior belief about the components' variance. Of course, when setting these parameters, it is crucial to consider the measurement scale of the data. By default, test outcomes are standardised (so that the resulting mean is zero and the variance is one) in the `pooledROC.dpm` function and the default values are as follows

$$m_{D0} = 0, \quad S_{D0} = 10, \quad a_D = 2, \quad b_D = 0.5.$$

Because test outcomes are standardised we expect the means of the components to be near zero and hence why $m_{D0} = 0$. The parameter S_{D0} then controls where the drawn μ_{Dl} can lie and the value of 10 implies that approximately 95% of the values roughly lie within -6 and 6 . Further, note that $a_D = 2$ leads to a prior with infinite variance that is centred around a finite mean ($b_D = 0.5$) and therefore favours variances less than one. Considering that the standardised data have a variance of one, it is reasonable to expect the within component variance to be smaller than the overall variance. The option of not standardising the test outcomes is also available in `pooledROC.dpm` and in such a case the defaults for the centring distribution hyperparameters' values are as following

$$m_{D0} = \bar{y}_D, \quad S_{D0} = 100s_D^2/n_D, \quad a_D = 2, \quad b_D = s_D^2/2,$$

with $\bar{y}_D = \frac{1}{n_D} \sum_{j=1}^{n_D} y_{Dj}$ and $s_D^2 = \frac{1}{n_D-1} \sum_{j=1}^{n_D} (y_{Dj} - \bar{y}_D)^2$. Regarding the precision parameter of the DP, α_D , it has a direct relationship with the number of occupied mixture components. One possible strategy for specifying α_D is to fix it to a small value to favour a small number of occupied components relative to the sample size. In the `pooledROC.dpm` function we set $\alpha_D = 1$, a commonly used default value (Gelman, Carlin, Stern, Dunson, Vehtari, and Rubin 2013, p. 553). Lastly, by default $L_D = 10$. Before proceeding we shall emphasise that these two configurations of hyperparameters values (for standardised and not standardised test outcomes) have proved to work well for a different range of test outcomes distributions but it is certainly not our goal to encourage users to use it blindly and thought should be dedicated to this important task. Nevertheless, output from the function `pooledROC.dpm` may be post processed and (informal) model fit diagnostics obtained; see more in Section and in the Appendices.

Because the full conditional distributions for all model parameters are available in closed-form, posterior simulation can be easily conducted through Gibbs sampler (see the details, for instance, in Ishwaran and James 2002). At iteration s of the Gibbs sampler procedure, the ROC curve is computed as

$$\text{ROC}^{(s)}(p) = 1 - F_D^{(s)} \left\{ F_D^{-1(s)}(1-p) \right\}, \quad s = 1, \dots, S,$$

with

$$F_D^{(s)}(y) = \sum_{l=1}^{L_D} \omega_{Dl}^{(s)} \Phi\left(y \mid \mu_{Dl}^{(s)}, \sigma_{Dl}^{2(s)}\right), \quad F_{\bar{D}}^{(s)}(y) = \sum_{k=1}^{L_{\bar{D}}} \omega_{\bar{D}k}^{(s)} \Phi\left(y \mid \mu_{\bar{D}k}^{(s)}, \sigma_{\bar{D}k}^{2(s)}\right), \quad (26)$$

and where the inversion is performed numerically. There is a closed-form expression for the AUC (Erkanli *et al.* 2006) given by

$$\text{AUC}^{(s)} = \sum_{k=1}^{L_{\bar{D}}} \sum_{l=1}^{L_D} \omega_{\bar{D}k}^{(s)} \omega_{Dl}^{(s)} \Phi\left(\frac{b_{kl}^{(s)}}{\sqrt{1 + a_{kl}^{2(s)}}}\right), \quad b_{kl}^{(s)} = \frac{\mu_{Dl}^{(s)} - \mu_{\bar{D}k}^{(s)}}{\sigma_{Dl}^{(s)}}, \quad a_{kl}^{(s)} = \frac{\sigma_{\bar{D}k}^{(s)}}{\sigma_{Dl}^{(s)}}.$$

Also, when $L_D = L_{\bar{D}} = 1$, there are closed-form expressions for the pAUC and pAUC_{TPF} which are used in the package (see Hillis and Metz 2012). For the pAUC/pAUC_{TPF}, when $L_D > 1$ or $L_{\bar{D}} > 1$, the integrals are approximated numerically using Simpson's rule. The Youden index/optimal threshold is computed as in the Bayesian bootstrap method, with the obvious difference that here the CDFs are expressed as in (26). At the end of the sampling procedure, we have an ensemble of S ROC curves and AUCs/pAUCs/pAUC_{TPFS}/YIs/optimal thresholds which, as before, allows obtaining point and interval estimates.

3.2. Covariate-specific ROC curve

We now let $\{(\mathbf{x}_{\bar{D}i}, y_{\bar{D}i})\}_{i=1}^{n_{\bar{D}}}$ and $\{(\mathbf{x}_{Dj}, y_{Dj})\}_{j=1}^{n_D}$ be two independent random samples of test outcomes and covariates from the nondiseased and diseased groups of size $n_{\bar{D}}$ and n_D , respectively. Further, for all $i = 1, \dots, n_{\bar{D}}$ and $j = 1, \dots, n_D$, let $\mathbf{x}_{\bar{D}i} = (x_{\bar{D}i,1}, \dots, x_{\bar{D}i,q})^\top$ and $\mathbf{x}_{Dj} = (x_{Dj,1}, \dots, x_{Dj,q})^\top$ be q -dimensional vectors of covariates, which can be either continuous or categorical.

Induced semiparametric linear model

The function `cROC.sp` implements the induced ROC approaches proposed by Faraggi (2003) and Pepe (1998). Both authors assume a location-scale regression model of the following form for the test outcomes in each group

$$Y_{\bar{D}} = \tilde{\mathbf{X}}_{\bar{D}}^\top \boldsymbol{\beta}_{\bar{D}} + \sigma_{\bar{D}} \varepsilon_{\bar{D}}, \quad Y_D = \tilde{\mathbf{X}}_D^\top \boldsymbol{\beta}_D + \sigma_D \varepsilon_D, \quad (27)$$

where $\tilde{\mathbf{X}}_{\bar{D}}^\top = (1, \mathbf{X}_{\bar{D}}^\top)$ and $\boldsymbol{\beta}_{\bar{D}} = (\beta_{\bar{D}0}, \dots, \beta_{\bar{D}q})^\top$ is a $(q+1)$ -dimensional vector of (unknown) regression coefficients; $\tilde{\mathbf{X}}_D$ and $\boldsymbol{\beta}_D$ are analogously defined. The error terms $\varepsilon_{\bar{D}}$ and ε_D have mean zero, variance one, are independent of each other and of the covariate, and have distribution functions given by $F_{\varepsilon_{\bar{D}}}$ and F_{ε_D} , respectively. Under these assumptions, we have

$$F_{\bar{D}}(y \mid \mathbf{x}) = F_{\varepsilon_{\bar{D}}}\left(\frac{y - \tilde{\mathbf{x}}^\top \boldsymbol{\beta}_{\bar{D}}}{\sigma_{\bar{D}}}\right) \quad \text{and} \quad F_D(y \mid \mathbf{x}) = F_{\varepsilon_D}\left(\frac{y - \tilde{\mathbf{x}}^\top \boldsymbol{\beta}_D}{\sigma_D}\right), \quad (28)$$

with $\tilde{\mathbf{x}}^\top = (1, \mathbf{x}^\top)$.

The approaches of Faraggi (2003) and Pepe (1998) differ in the assumptions made about the error terms. More concretely, Faraggi (2003)'s method assumes that the error term in both groups follows a standard normal distribution, i.e., $F_{\varepsilon_{\bar{D}}}(y) = F_{\varepsilon_D}(y) = \Phi(y)$, and can be summarised by the following three steps:

1. Estimate the regression coefficients $\beta_{\bar{D}}$ and β_D by ordinary least squares, on the basis of the samples $\{(\mathbf{x}_{\bar{D}i}, y_{\bar{D}i})\}_{i=1}^{n_{\bar{D}}}$ and $\{(\mathbf{x}_{Dj}, y_{Dj})\}_{j=1}^{n_D}$, respectively.
2. Estimate $\hat{\sigma}_{\bar{D}}^2$ as

$$\hat{\sigma}_D^2 = \frac{\sum_{j=1}^{n_D} (y_{Dj} - \tilde{\mathbf{x}}_{Dj}^\top \hat{\beta}_D)^2}{n_D - q - 1},$$

with $\hat{\sigma}_{\bar{D}}^2$ similarly estimated.

3. For a given covariate vector \mathbf{x} , compute the covariate-specific ROC curve as follows

$$\widehat{\text{ROC}}(p | \mathbf{x}) = 1 - \Phi \left\{ a(\mathbf{x}) + b\Phi^{-1}(1 - p) \right\}, \quad (29)$$

where

$$a(\mathbf{x}) = \tilde{\mathbf{x}}^\top \frac{(\hat{\beta}_{\bar{D}} - \hat{\beta}_D)}{\hat{\sigma}_D}, \quad \text{and} \quad b = \frac{\hat{\sigma}_{\bar{D}}}{\hat{\sigma}_D}. \quad (30)$$

Regarding the covariate-specific AUC, pAUC, and pAUC_{TPF} (Expressions (12), (13), and (14)) they admit closed-form expressions (see Hillis and Metz 2012).

As an alternative, Pepe (1998) suggests to estimate the CDF of the errors in each group by the corresponding empirical CDF of the estimated standardised residuals. Therefore, the first two steps of the estimation procedure remain the same, but now we have the following extra step

$$\hat{F}_{\varepsilon_D}(y) = \frac{1}{n_D} \sum_{j=1}^{n_D} I(\hat{\varepsilon}_{Dj} \leq y), \quad \hat{\varepsilon}_{Dj} = \frac{y_{Dj} - \tilde{\mathbf{x}}_{Dj}^\top \hat{\beta}_D}{\hat{\sigma}_D}.$$

The empirical CDF of the standardised residuals in the nondiseased group is estimated in a similar fashion. The covariate-specific ROC curve is finally computed in an analogous way as for the method of Faraggi (2003) as

$$\widehat{\text{ROC}}(p | \mathbf{x}) = 1 - \hat{F}_{\varepsilon_D} \left\{ a(\mathbf{x}) + b\hat{F}_{\varepsilon_D}^{-1}(1 - p) \right\}.$$

Here, the covariate-specific AUC and pAUC (Expressions (12) and (13)) also admit closed forms. However, especially for large datasets, their calculation can be very time-consuming. As a consequence, in **ROCnReg** they are computed numerically using Simpson's rule; in our experience Simpson's rule provides almost identical results to the ones obtained using the closed-form expressions. In what concerns the covariate-specific pAUC_{TNF}, it is interesting to note that

$$\widehat{\text{ROC}}_{\text{TNF}}(p | \mathbf{x}) = 1 - \hat{F}_{\varepsilon_D} \left\{ a^*(\mathbf{x}) + b^*\hat{F}_{\varepsilon_D}^{-1}(1 - p) \right\},$$

with

$$a^*(\mathbf{x}) = \tilde{\mathbf{x}}^\top \frac{(\hat{\beta}_D - \hat{\beta}_{\bar{D}})}{\hat{\sigma}_{\bar{D}}} \quad \text{and} \quad b^* = \frac{\hat{\sigma}_D}{\hat{\sigma}_{\bar{D}}}.$$

The covariate-specific pAUC_{TNF} (Expression (14)) is then computed numerically using Simpson's rule based on the previous expressions.

Finally, in pretty much the same way as for the pooled ROC curve, the covariate-specific Youden index (and associated threshold value) is obtained by maximising, over a grid of possible threshold values, the expression in (16), making use of result (28).

Induced kernel-based approach

The kernel-based approach of González-Manteiga, Pardo-Fernández, and van Keilegom (2011) and Rodríguez-Álvarez *et al.* (2011b) is available in the `cROC.kernel`. Differently to all the other estimating approaches for the covariate-specific ROC curve presented in this section, it can only deal with one continuous covariate. Similarly to the approaches of Pepe (1998) and Faraggi (2003), it also assumes a location-scale regression model for the test outcomes in each group, but the effect of the covariate is not assumed to be linear and the variance is allowed to depend on the covariate. Specifically, the models postulated in each group are as follows

$$Y_{\bar{D}} = \mu_{\bar{D}}(X_{\bar{D}}) + \sigma_{\bar{D}}(X_{\bar{D}})\varepsilon_{\bar{D}}, \quad Y_D = \mu_D(X_D) + \sigma_D(X_D)\varepsilon_D, \quad (31)$$

where $\mu_D(x) = \mathbf{E}(Y_D | X_D = x)$ and $\sigma_D^2(x) = \mathbf{VAR}(Y_D | X_D = x)$ are the regression and variance functions, respectively, with $\mu_{\bar{D}}(x)$ and $\sigma_{\bar{D}}^2(x)$ being analogously defined. The error terms $\varepsilon_{\bar{D}}$ and ε_D have mean zero, variance one, are independent of each other and of the covariate, and have distribution functions given by $F_{\varepsilon_{\bar{D}}}$ and F_{ε_D} , respectively.

Both the regression and variance functions are estimated using local polynomial kernel smoothers (Fan and Gijbels 1996). In particular, local constant (Nadaraya–Watson) or local linear estimators are employed for the regression function, whereas for the variance function only local constant estimators are used. Estimation in **ROCnReg** makes use of the R package **np** by Hayfield and Racine (2008). We note that estimation proceeds in a sequential manner: 1) the regression function, say in the diseased group and denoted by $\hat{\mu}_D$, is estimated first on the basis of $\{(x_{Dj}, y_{Dj})\}_{j=1}^{n_D}$, and 2) the variance function is estimated next on the basis of the sample $\{(x_{Dj}, [y_{Dj} - \hat{\mu}_D(x_{Dj})]^2)\}$. Both steps involve the selection of a bandwidth parameter which is done via cross-validation. As in the model of Pepe (1998), the CDFs F_{ε_D} and $F_{\varepsilon_{\bar{D}}}$ are estimated via the empirical CDF of the standardised residuals, that is,

$$\hat{F}_{\varepsilon_D}(y) = \frac{1}{n_D} \sum_{j=1}^{n_D} I(\hat{\varepsilon}_{Dj} \leq y), \quad \hat{\varepsilon}_{Dj} = \frac{y_{Dj} - \hat{\mu}_D(x_{Dj})}{\hat{\sigma}_D(x_{Dj})}$$

with the empirical CDF of the standardised residuals in the nondiseased group estimated analogously. Finally, the covariate-specific ROC curve is computed in an analogous way as before as

$$\widehat{\text{ROC}}(p | x) = 1 - \hat{F}_{\varepsilon_D} \left\{ a(x) + b(x) \hat{F}_{\varepsilon_{\bar{D}}}^{-1}(1 - p) \right\},$$

where

$$a(x) = \frac{\hat{\mu}_{\bar{D}}(x) - \hat{\mu}_D(x)}{\hat{\sigma}_D(x)} \quad \text{and} \quad b(x) = \frac{\hat{\sigma}_{\bar{D}}(x)}{\hat{\sigma}_D(x)}.$$

Estimation of the covariate-specific AUC, pAAUC, pAUC_{TNF}, and YI follows a similar reasoning as the one described previously for the induced semiparametric linear model when no assumptions are made regarding the distribution of the error terms.

For both the induced semiparametric linear model and the induced kernel approach, uncertainty quantification is done through a bootstrap of the residuals. For further details see, for instance, Rodríguez-Álvarez *et al.* (2011b).

Bayesian nonparametric approach based on a dependent Dirichlet process mixture of normal distributions

The Bayesian nonparametric approach for conducting inference about the covariate-specific ROC curve of Inácio de Carvalho *et al.* (2013), which is based on a single-weights dependent

Dirichlet process mixture of normal distributions De Iorio, Johnson, Müller, and Rosner (2009), is implemented in the function `cROC.bnp`. By opposition to the previously described approaches to ROC regression, this method rests on directly modelling the CDF of test outcomes separately in the diseased and nondiseased groups. In a single-weights dependent Dirichlet process mixture of normals model (De Iorio *et al.* 2009), the conditional CDF in the diseased group is modelled as follows

$$F_D(y_{Dj} | \mathbf{X}_D = \mathbf{x}_{Dj}) = \int \Phi(y_{Dj} | \mu_D(\mathbf{x}_{Dj}, \boldsymbol{\beta}), \sigma^2) dG_D(\boldsymbol{\beta}, \sigma^2), \quad G_D \sim \text{DP}(\alpha_D, G_D^*(\boldsymbol{\beta}, \sigma^2)),$$

with the conditional CDF in the nondiseased group \bar{D} following in an analogous manner. As in the no-covariate case, by making use of Sethuraman's truncated representation of the DP, we can write the conditional CDF as

$$\begin{aligned} F_D(y_{Dj} | \mathbf{x}_{Dj}) &= \sum_{l=1}^{L_D} \omega_{Dl} \Phi(y_{Dj} | \mu_D(\mathbf{x}_{Dj}, \boldsymbol{\beta}_{Dl}), \sigma_{Dl}^2), \\ \omega_{D1} &= v_{D1}, \quad \omega_{Dl} = v_{Dl} \prod_{r<l} (1 - v_{Dr}), \quad l = 2, \dots, L_D, \\ v_{Dl} &\stackrel{\text{iid}}{\sim} \text{Beta}(1, \alpha_D), \quad l = 1, \dots, L_D - 1, \quad v_{DL_D} = 1. \end{aligned}$$

$$\text{var}(y_{Dj} | \mathbf{x}_{Dj}) = \sum_{l=1}^{L_D} \omega_{Dl} \sigma_{Dl}^2 + \sum_{l=1}^{L_D} \omega_{Dl} \left\{ \mu_D(\mathbf{x}_{Dj}, \boldsymbol{\beta}_{Dl}) - \left(\sum_{l=1}^{L_D} \omega_{Dl} \mu_D(\mathbf{x}_{Dj}, \boldsymbol{\beta}_{Dl}) \right)^2 \right\}.$$

Note that by assuming that the weights, w_{Dl} , do not vary with covariates, the model might have limited flexibility in practice (MacEachern 2000). This issue can, however, be largely mitigated by using a flexible formulation for $\mu_D(\mathbf{x}_{Dj}, \boldsymbol{\beta}_{Dl})$, which is needed not only for the model to be able to recover nonlinear trends, but also to recover flexible shapes that might arise due to a dependence of the weights on the covariates. As such, the function `cROC.bnp` in **ROCnReg** allows to model the mean function of each component using an additive smooth structure

$$\mu_D(\mathbf{x}_{Dj}, \boldsymbol{\beta}_{Dl}) = \beta_{Dl0} + f_{Dl1}(x_{Dj,1}) + \dots + f_{Dlq}(x_{Dj,q}), \quad l = 1, \dots, L_D, \quad (32)$$

where the smooth functions, f_{Dlm} ($m = 1, \dots, q$), are approximated using a linear combination of B-splines basis functions. To avoid notational burden we have assumed that all q covariates are continuous and modelled in a flexible way. However, the function `cROC.bnp` can also deal with categorical covariates, linear effects of continuous covariates, as well as interactions. For the reasons mentioned before, we recommend that all continuous covariates are modelled as in (32). Nonetheless, posterior predictive checks, as illustrated in Section , can also be used to informally validate the fitted model. We write

$$\mu_D(\mathbf{x}_{Dj}, \boldsymbol{\beta}_{Dl}) = \mathbf{z}_{Dj}^\top \boldsymbol{\beta}_{Dl}, \quad l = 1, \dots, L_D, \quad j = 1, \dots, n_D, \quad (33)$$

where \mathbf{z}_{Dj}^\top is the j th row of the design matrix that contains the intercept, the continuous covariates that are modelled in a linear way (if any), the cubic B-splines basis representation for those modelled in a flexible way, the categorical covariates (if any), and their interaction(s) (if believed to exist). Also, $\boldsymbol{\beta}_{Dl}$ collects, for the l th component, the regression coefficients associated with the aforementioned covariates. For the covariate effects modelled using cubic

B-splines, an important issue is the selection of the number and location of the knots at which to anchor the basis functions, as this has the potential to impact inferences, more so for the former than the latter. The selection of the number of knots can be assisted by a model selection criterion, for example, (the adaptation to the case of mixture models of) the deviance information criterion (DIC) (Celeux, Forbes, Robert, and Titterton 2006), the log pseudo marginal likelihood (LPML) (Geisser and Eddy 1979), and the widely applicable information criterion (WAIC) (Gelman, Hwang, and Vehtari 2014). In turn, for the location of the interior knots themselves we follow Rosenberg (1995) and use the quantiles of the covariate values.

The regression coefficients and variances associated with each of the L_D components are sampled from the conjugate centring distribution $(\boldsymbol{\beta}_{Dl}, \sigma_{Dl}^{-2}) \stackrel{\text{iid}}{\sim} N_{Q_D}(\mathbf{m}_D, \mathbf{S}_D) \Gamma(a_D, b_D)$, with conjugate hyperpriors $\mathbf{m}_D \sim N(\mathbf{m}_{D0}, \mathbf{S}_{D0})$ and $\mathbf{S}_D^{-1} \sim \text{Wishart}(\nu_D, (\nu_D \Psi_D)^{-1})$ (a Wishart distribution with degrees of freedom ν_D and expectation Ψ_D^{-1}) and where Q_D is the dimension of the vector \mathbf{z}_{Dj} . Hyperparameters \mathbf{m}_{D0} and Ψ_D must be chosen to represent the prior belief about the regression coefficients associated to each mixture component and about their covariance matrix, respectively, whereas \mathbf{S}_{D0} and ν_D are chosen to represent the confidence in the prior belief of \mathbf{m}_{D0} and Ψ_D , respectively. As in the no-covariate case, by default in `cROC.bnp` test outcomes and covariates are standardised, which not only facilitates specification of the hyperparameter values but also improves the mixing of the Markov chain Monte Carlo (MCMC) chains. The default values are as follows:

$$\mathbf{m}_{D0} = \mathbf{0}_{Q_D}, \quad \mathbf{S}_{D0} = 10I_{Q_D}, \quad \nu_D = Q_D + 2, \quad \Psi_D = I_{Q_D}, \quad a_D = 2, \quad b_D = 0.5.$$

When test outcomes and covariates are not standardised, the defaults are the following:

$$\mathbf{m}_{D0} = \hat{\boldsymbol{\beta}}_D, \quad \mathbf{S}_{D0} = \hat{\boldsymbol{\Sigma}}_D, \quad \nu_D = Q_D + 2, \quad \Psi_D = 30\hat{\boldsymbol{\Sigma}}_D, \quad a_D = 2, \quad b_D = \hat{\sigma}_D^2/2,$$

where $\hat{\boldsymbol{\beta}}_D$ and $\hat{\sigma}_D$ are the least squares estimates from fitting the linear model $y_{Dj} = \mathbf{z}_{Dj}^\top \boldsymbol{\beta}_D + \sigma_D \varepsilon_{Dj}$, where $E(\varepsilon_{Dj}) = 0$, $\text{var}(\varepsilon_{Dj}) = 1$, and $\hat{\boldsymbol{\Sigma}}_D$ is the estimated covariance matrix of $\hat{\boldsymbol{\beta}}_D$. With regard to the specification of α_D and L_D , as in the DPM model (no-covariate case) we set them, respectively, to 1 and 10. The blocked Gibbs sampler is used to simulate draws from the posterior distribution and details about it can be found, for instance, in the Supplementary Materials of Inácio de Carvalho *et al.* (2017).

Similarly to the analogous model for the no covariate case, at iteration s of the Gibbs sampler procedure, the covariate-specific ROC curve is computed as

$$\text{ROC}^{(s)}(p | \mathbf{x}) = 1 - F_D^{(s)} \left\{ F_D^{-1(s)}(1 - p | \mathbf{x}) | \mathbf{x} \right\}, \quad s = 1, \dots, S,$$

with

$$F_D^{(s)}(y | \mathbf{x}) = \sum_{l=1}^{L_D} \omega_{Dl}^{(s)} \Phi \left(y | \mathbf{z}^\top \boldsymbol{\beta}_{Dl}^{(s)}, \sigma_{Dl}^{2(s)} \right), \quad F_D^{-1(s)}(y | \mathbf{x}) = \sum_{k=1}^{L_D} \omega_{Dk}^{(s)} \Phi \left(y | \mathbf{z}^\top \boldsymbol{\beta}_{Dk}^{(s)}, \sigma_{Dk}^{2(s)} \right), \quad (34)$$

and where the inversion is performed numerically. A point estimate for $\text{ROC}(p | \mathbf{x})$ can be obtained by computing the mean (or the median) of the ensemble $\{\text{ROC}^{(1)}(p | \mathbf{x}), \dots, \text{ROC}^{(S)}(p | \mathbf{x})\}$, with pointwise credible bands derived from the percentiles of the ensemble. Although the results presented in Erkanli *et al.* (2006) can be extended to derive a closed-form expression

for the covariate-specific AUC, for computational reasons, in **ROCnReg** the integral in (12) is approximated using Simpson's rule, and the same applies for the partial areas. Conditionally on a specific covariate value, the computation of the Youden index and of the optimal threshold proceeds in a similar way as in the DPM model (see [Inácio de Carvalho *et al.* 2017](#), for details). As for the covariate-specific ROC curve, point and interval estimates can be obtained from the corresponding covariate-specific ensemble of each summary measure.

3.3. Covariate-adjusted ROC curve

All estimators for the covariate-adjusted ROC curve make use of Equation (19) and rely on the following three steps

1. Estimation of the conditional distribution of test outcomes in the nondiseased group, $F_{\bar{D}}(y_{\bar{D}i} | \mathbf{x}_{\bar{D}i})$.
2. Computation of the placement value $U_D = 1 - F_{\bar{D}}(Y_D | \mathbf{X}_D)$ where, by a slight abuse of notation, we are designating it by the same letter used for the unconditional case.
3. Estimation of the cumulative distribution function of U_D .

Frequentist approaches

The approaches used for estimation of the AROC curve proposed by [Janes and Pepe \(2009\)](#) and [Rodríguez-Álvarez *et al.* \(2011b\)](#) only differ in Step 1. Specifically, once one has an estimate of the conditional CDF in the nondiseased group, say $\hat{F}_{\bar{D}}(\cdot | \mathbf{x})$, Step 2 in the two approaches consists of trivially computing the diseased placement values as

$$\hat{U}_{Dj} = 1 - \hat{F}_{\bar{D}}(y_{Dj} | \mathbf{x}_{Dj}), \quad j = 1, \dots, n_D.$$

Next, in Step 3, the AROC curve at a false positive fraction of p is estimated via the empirical distribution function of the placement values calculated in the previous step, $\{\hat{U}_{Dj}\}_{j=1}^{n_D}$, that is,

$$\widehat{\text{AROC}}(p) = \frac{1}{n_D} \sum_{j=1}^{n_D} I(\hat{U}_{Dj} \leq p).$$

With regard to Step 1, both authors assume a location-scale regression model for the test outcomes in the nondiseased group and, as such and as explained in the previous section, the conditional CDF of the test results can be written using the CDF of the regression errors, i.e.,

$$F_{\bar{D}}(y | \mathbf{x}) = F_{\varepsilon_{\bar{D}}} \left(\frac{y - \mu_{\bar{D}}(\mathbf{x})}{\sigma_{\bar{D}}(\mathbf{x})} \right).$$

While [Janes and Pepe \(2009\)](#) assume a location-scale model of the form of (27), [Rodríguez-Álvarez *et al.* \(2011b\)](#) rely on specification (31). The estimation of the mean and variance functions follow exactly the same procedures as those described in the induced semiparametric linear model (for [Janes and Pepe 2009](#)) and induced kernel-based approach (for [Rodríguez-Álvarez *et al.* 2011b](#)) for the covariate specific ROC curve (the only difference being that here we only need to perform the estimation for the nondiseased group). At last, and also as in the estimators for the covariate-specific ROC curve, $F_{\varepsilon_{\bar{D}}}$ can be either assumed to be the

standard normal distribution or left unspecified and estimated empirically on the basis of the standardised residuals. In both cases, the AAUC and pAAUC can be computed as follows

$$\widehat{\text{AAUC}} = \int_0^1 \text{AROC}(p) dp = 1 - \frac{1}{n_D} \sum_{j=1}^{n_D} \widehat{U}_{Dj},$$

$$\text{pAAUC}(u_1) = \int_0^{u_1} \text{AROC}(p) dp = u_1 - \frac{1}{n_D} \sum_{j=1}^{n_D} \min \{u_1, \widehat{U}_{Dj}\},$$

whereas the $\text{pAAUC}_{\text{TNF}}$ is computed as in Equation (22) using numerical integration methods (function `integrate` in R package `stats`).

Bayesian nonparametric approach

Recently, Inácio de Carvalho and Rodríguez-Álvarez (2018) proposed a Bayesian nonparametric estimator for the AROC curve that combines a dependent Dirichlet process mixture model and the Bayesian bootstrap. As in the Bayesian nonparametric approach for estimating the covariate-specific ROC curve, in Step 1, rather than assuming a location-scale regression model for the test outcomes in the nondiseased group, the entire conditional distribution is modelled using a single-weights dependent Dirichlet process mixture of normal distributions. Again, here, we also recommend to use cubic B-splines transformations of all continuous covariates. Using the same notation as before, we model the conditional density as

$$F_{\bar{D}}(y_{\bar{D}i} | \mathbf{x}_{\bar{D}i}) = \sum_{l=1}^{L_{\bar{D}}} \omega_{\bar{D}l} \Phi(y_{\bar{D}i} | \mathbf{z}_{\bar{D}i}^{\top} \boldsymbol{\beta}_{\bar{D}l}, \sigma_{\bar{D}l}^2).$$

The same prior distributions and default values as in the `cROC.bnp` function are adopted for $\boldsymbol{\beta}_{\bar{D}l}$ and $\sigma_{\bar{D}l}^2$. Once Step 1 has been completed and given a posterior sample from the parameters of interest, the corresponding realisation of the placement value of a diseased subject in the nondiseased population is easily computed as

$$U_{Dj}^{(s)} = 1 - F_{\bar{D}}^{(s)}(y_{Dj} | \mathbf{x}_{Dj}) = \sum_{l=1}^{L_{\bar{D}}} \omega_{\bar{D}l}^{(s)} \Phi(y_{Dj} | \mathbf{z}_{Dj}^{\top} \boldsymbol{\beta}_{\bar{D}l}^{(s)}, \sigma_{\bar{D}l}^{2(s)}), \quad j = 1, \dots, n_D, \quad s = 1, \dots, S.$$

Finally, in Step 3, the cumulative distribution function of $\{U_{Dj}^{(s)}\}_{j=1}^{n_D}$ is estimated through the Bayesian bootstrap

$$\text{AROC}^{(s)}(p) = \sum_{j=1}^{n_D} q_j^{(s)} I(U_{Dj}^{(s)} \leq p), \quad (q_1^{(s)}, \dots, q_{n_D}^{(s)}) \sim \text{Dirichlet}(n_D; 1, \dots, 1).$$

As before, closed-form expressions do exist for the AAUC and pAAUC

$$\text{AAUC}^{(s)} = \int_0^1 \text{AROC}^{(s)}(p) dp = 1 - \sum_{j=1}^{n_D} q_j^{(s)} U_{Dj}^{(s)},$$

$$\text{pAAUC}^{(s)}(u_1) = \int_0^{u_1} \text{AROC}^{(s)}(p) dp = u_1 - \sum_{j=1}^{n_D} q_j^{(s)} \min \{u_1, U_{Dj}^{(s)}\},$$

and the $\text{pAAUC}_{\text{TNF}}$ (Equation (22)) is computed using numerical integration methods. With regard to the YI, it is obtained by directly plugging in $\text{AROC}^{(s)}(p)$ in Expression (23).

A point estimate for $\text{AROC}(p)$ can be obtained by computing the mean (or the median) of the ensemble $\{\text{AROC}^{(1)}(p), \dots, \text{AROC}^{(S)}(p)\}$, that is,

$$\widehat{\text{AROC}}(p) = \frac{1}{S} \sum_{s=1}^S \text{AROC}^{(s)}(p),$$

and the percentiles of the ensemble can be used to provide pointwise credible bands/credible intervals. The same applies for the AAUC and pAAUC.

4. Package presentation and illustration

This section describes the main functions in the **ROCnReg** package and illustrates their usage using endocrine data from a cross-sectional study carried out by the Galician Endocrinology and Nutrition Foundation (FENGA). A detailed description of this dataset can be found in [Tomé Martínez de Rituerto, Botana, Cadarso-Suárez, Rego-Iraeta, Fernández-Mariño, Mato, Solache, and Perez-Fernandez \(2009\)](#). It has also been previously analysed in [Rodríguez-Álvarez *et al.* \(2011b,a\)](#) and [Inácio de Carvalho and Rodríguez-Álvarez \(2018\)](#). For confidentiality reasons, the data used in this paper correspond to a synthetic dataset that was obtained by mimicking the original one and can be found in the **ROCnReg** package under the name `endosyn`. A summary of the data follows.

```
R> library("ROCnReg")
R> data("endosyn")
R> summary(endosyn)
```

cvd_idf	age	gender	bmi
Min. :0.0000	Min. :18.25	Men :1317	Min. :12.60
1st Qu.:0.0000	1st Qu.:29.57	Women:1523	1st Qu.:23.19
Median :0.0000	Median :39.28		Median :26.24
Mean :0.2433	Mean :41.43		Mean :26.69
3rd Qu.:0.0000	3rd Qu.:50.84		3rd Qu.:29.74
Max. :1.0000	Max. :84.66		Max. :46.20

The dataset is comprised of 2840 individuals (1317 men and 1523 women, variable `gender`), with an `age` range between 18 and 85 years old. Variable `bmi` contains the body mass index (BMI) values, and `cvd_idf` is the variable that indicates the presence (1) or absence (0) of two or more cardiovascular disease (CVD) risk factors. Following previous studies, the CVD risk factors considered include raised triglycerides, reduced HDL-cholesterol, raised blood pressure, and raised fasting plasma glucose. Note that from the 2840 individuals, about 24% present two or more CVD risk factors.

Using the **ROCnReg** package, in the subsequent sections we will illustrate how to ascertain, through the pooled ROC curve, the discriminatory capacity of the BMI (which acts as our diagnostic test in this example) in differentiating individuals with two or more CVD risk factors (those belonging to the diseased class D) from those having none or just one CVD

risk factor (and that therefore belong to the nondiseased group \bar{D}). Also, in Section 4.2 we will show how to evaluate, through the covariate-specific ROC curve, the possible modifying effect of `age` and `gender` on the discriminatory capacity of the BMI. Finally, Section 4.3 focuses on the covariate-adjusted ROC curve as a global summary measure of the BMI discriminatory ability, when taking the `age` and `gender` effects into account. In the Appendices we show the usage of the package for those methods not described in the main text.

4.1. Pooled ROC curve

The **ROCnReg** package allows estimating the pooled ROC curve by means of the four methods described in Section 3. Recall that function `pooledROC.emp` implements the empirical estimator, `pooledROC.kernel` the kernel-based approach, and `pooledROC.BB` and `pooledROC.dpm` correspond, respectively, to the Bayesian bootstrap estimator and the approach based on a Dirichlet process mixture (of normal distributions). The input arguments in the functions are method-specific (details can be found in the manual accompanying the package), but in all cases numerical and graphical summaries can be obtained by calling the functions `print.pooledROC`, `summary.pooledROC`, and `plot.pooledROC`, which can be abbreviated by `print`, `summary`, and `plot`.

By way of example, we present here the syntax using the `pooledROC.dpm` function. Recall that our aim is to ascertain, using the `endosyn` dataset, the discriminatory capacity of the BMI in differentiating individuals with two or more CVD risk factors from those having none or just one CVD risk factor.

```
R> set.seed(123, "L'Ecuyer-CMRG") # for reproducibility
R> pROC_dpm <- pooledROC.dpm(marker = "bmi", group = "cvd_idf",
+   tag.h = 0, data = endosyn, standardise = TRUE, p = seq(0, 1, l = 101),
+   ci.level = 0.95, compute.lpm1 = TRUE, compute.WAIC = TRUE, compute.DIC = TRUE,
+   pauc = pauccontrol(compute = TRUE, focus = "FPF", value = 0.1),
+   density = densitycontrol(compute = TRUE), prior.h = priorcontrol.dpm(),
+   prior.d = priorcontrol.dpm(),
+   mcmc = mcmccontrol(nsave = 8000, nburn = 2000, nskip = 1),
+   parallel = "snow", ncpus = 2, cl = NULL)
```

Before describing in detail the previous call, we first present the control functions that are used. In particular

```
pauccontrol(compute = FALSE, focus = c("FPF", "TPF"), value = 1)
```

can be used to indicate whether the pAUC should be computed (by default it is not computed), and in case it is computed (i.e., `compute = TRUE`), whether the `focus` should be placed on restricted FPFs (pAUC, see (4)) or on restricted TPFs (pAUC_{TPF}, see (5)). In both cases, the upper bound u_1 (if focus is the FPF) or the lower bound v_1 (if focus is the TPF) should be indicated in the argument `value`. In addition to the pooled ROC curve, AUC, and pAUC (if required), the function `pooledROC.dpm` also allows computing the probability density function (PDF) of the test outcomes in both the diseased and nondiseased groups. In order to do so, we use

```
densitycontrol(compute = FALSE, grid.h = NA, grid.d = NA)
```

By default, PDFs are not returned by the function `pooledROC.dpm`, but this can be changed by setting `compute = TRUE`, and through `grid.h` and `grid.d` the user can specify a grid of test results where the PDFs are to be evaluated in, respectively, the nondiseased and diseased groups. Value `NA` signals auto initialisation, with default a vector of length 200 in the range of the test results. Regarding the hyperparameters for the Dirichlet process mixture of normals model (used for the estimation of the PDFs/CDFs of the test outcomes in each group), they can be controlled using

```
priorcontrol.dpm(m0 = NA, S0 = NA, a = 2, b = NA, alpha = 1, L = 10)
```

A detailed description of these hyperparameters is found in Section . Finally, to set the various parameters controlling the MCMC procedure (which in our case is simply a Gibbs sampler) we use

```
mcmccontrol(nsave = 8000, nburn = 2000, nskip = 1)
```

Here `nsave` is an integer value with the total number of scans to be saved, `nburn` is the number of burn-in scans, and `nskip` is the thinning interval. Unless due to memory usage reasons, we recommend not thinning and instead monitoring the effective sample size of the MCMC chain.

Coming back to the `pooledROC.dpm` function, through `marker` the user specifies the name of the variable containing the test results; in our case, these are the values of the BMI. The name of the variable that distinguishes diseased (two or more CVD risk factors, D) from nondiseased individuals (none or one CVD risk factor, \bar{D}) is represented by the argument `group`, and the value codifying nondiseased individuals in `group` is specified by `tag.h`. The `data` argument is a data frame containing the data and all needed variables. Setting `standardise = TRUE` (the default) will standardise (i.e., subtract the mean and divide by the standard deviation) the test outcomes. The set of FPFs at which to estimate the pooled ROC curve is specified in the argument `p`, and argument `ci.level` allows specifying the level for the credible intervals (by default: 0.95). The LPML, WAIC, and DIC are computed by setting, respectively, the arguments `compute.lpml`, `compute.WAIC`, and `compute.DIC` to `TRUE`. Argument `pauc` is an (optional) list of values to replace the default values returned by the function `pauccontrol`. Here, we ask for the pAUC to be computed, with focus on restricted FPFs and upper bound $u_1 = 0.1$. Similarly, the argument `density` is an (optional) list of values to replace the default values returned by the function `densitycontrol`, as it is the argument `mcmc`. Through `prior.h` and `prior.d` arguments we specify the hyperparameters in the nondiseased and diseased classes, respectively. Again, both arguments are (optional) lists of values to replace the default values returned by the function `priorcontrol.dpm`. We shall remember that different hyperparameters' default values are set depending on whether test outcomes are standardised or not. Finally, arguments `parallel`, `ncpus` and `cl` allow to perform parallel computations (based on the R-package `parallel`). In particular, through `parallel` the user specifies the type of parallel operation: either "no" (default), "multicore" (not available on Microsoft Windows operating systems) or "snow". Argument `ncpus` is used to indicate the number of processes to be used in parallel operation (when `parallel = "multicore"` or `parallel = "snow"`), and `cl` is an optional parallel or snow cluster to be used when `parallel = "snow"`. If `cl` is not supplied (as in our example), a cluster on the local machine is created for the duration of the call.

A numerical summary of the fitted model can be obtained by calling the function `summary`, that provides, among other information, the estimated AUC (posterior mean) and 95% credible interval (recall that, in the call to the function, we set `ci.level = 0.95`) and, if required, the LPML, WAIC, and DIC, separately, in the nondiseased (denoted here as Group H) and diseased (Group D) classes.

```
R> summary(pROC_dpm)
```

Call:

```
pooledROC.dpm(marker = "bmi", group = "cvd_idf", tag.h = 0, data = endosyn,
  standardise = TRUE, p = seq(0, 1, l = 101), ci.level = 0.95,
  compute.lpml = TRUE, compute.WAIC = TRUE, compute.DIC = TRUE,
  pauc = pauccontrol(compute = TRUE, focus = "FPF", value = 0.1),
  density = densitycontrol(compute = TRUE), prior.h = priorcontrol.dpm(L = 10),
  prior.d = priorcontrol.dpm(L = 10), mcmc = mcmccontrol(nsave = 8000,
  nburn = 2000, nskip = 1), parallel = "snow", ncpus = 2)
```

Approach: Pooled ROC curve - Bayesian DPM

```
-----
Area under the pooled ROC curve: 0.759 (0.74, 0.777)*
Partial area under the pooled ROC curve (FPF = 0.1): 0.168 (0.139, 0.199)*
* Credible level: 0.95
```

Model selection criteria:

	Group H	Group D
WAIC	12490.485	4017.063
WAIC (Penalty)	8.431	5.468
LPML	-6245.247	-2008.541
DIC	12490.276	4016.920
DIC (Penalty)	8.326	5.396

Sample sizes:

	Group H	Group D
Number of observations	2149	691
Number of missing data	0	0

To complement these numerical results, the **ROCnReg** package also provides graphical results that can be used to further explore the fitted model. Specifically, the function `plot` depicts the estimated pooled ROC curve and AUC (posterior means), jointly with `ci.level`×100% (pointwise) credible intervals (here 95%)

```
R> plot(pROC_dpm, cex.main = 1.5, cex.lab = 1.5, cex.axis = 1.5, cex = 1.5)
```

The result of the above code is shown in Figure 2.

By means of `density = densitycontrol(compute = TRUE)` in the call to the function, the estimates of the PDFs of the BMI in both classes are to be returned. This information can






Figure 2: Graphical results as provided by the `plot.pooledROC` function for an object of class `pooledROC.dpm`. Posterior mean and 95% pointwise credible band for the pooled ROC curve and corresponding posterior mean and 95% credible interval for the AUC.

be accessed through component `dens` in object `pROC_dpm` (i.e., `pROC_dpm$dens`), which is a list with elements `h` and `d` associated with the nondiseased and diseased groups, respectively. Each of the two elements is itself another list of two components: (1) `grid`, a vector that contains the grid of test results at which the PDFs have been evaluated (estimated); and (2) `dens`, a matrix with the PDFs at each iteration of the MCMC procedure. We can use these results to plot, e.g. the posterior mean (and 95% pointwise credible bands) of the PDF of the BMI in the nondiseased and diseased populations (see Figure 3(a), obtained using the R package `ggplot2` by Wickham 2016). As can be observed, the estimated densities obtained under the DPM method follow very closely the histograms of the data. Further, the estimated densities available in `dens` can be used, as advised by Gelman *et al.* (2013) (p. 553), to monitor convergence of the MCMC chains. The well-known label switching problem often leads to poor mixing of the chains of the component-specific parameters, but this may not impact convergence and mixing of the induced density/distribution of interest. For instance, Figure 4 shows trace plots of the MCMC iterations (after burn-in) of the PDFs of the BMI in the two groups, for different (and randomly selected) values of the BMI, and Figure 5 depicts the corresponding effective sample sizes and Geweke statistics (obtained using the R package `coda` by Plummer, Best, Cowles, and Vines 2006). Note that all plots give evidence of a good mixing and do not suggest lack of convergence. For conciseness, the R-code for producing Figures 3(a), 4 and 5 is not provided here, but in the replication code that accompanies the paper (<https://bitbucket.org/mxrodriguez/rocnreg>).

It is worth noting that the function `pooledROC.dpm` also allows fitting a normal distribution in each group; this is just a particular case (for which $L_D = L_{\bar{D}} = 1$) of the more general DPM model. In order to fit such model, one simply needs to set $L = 1$ in the `prior.d` and



(a) DPM model with 10 mixture components in each group



(b) Normal model in each group

Figure 3: Histogram of the (observed) BMI and posterior mean jointly along with 95% point-wise credible bands (red lines) of the PDF of the BMI obtained using (a) a Dirichlet process mixture of normals model (object `pROC_dpm`); and (b) a normal model (object `pROC_normal`). Left: Nondiseased individuals (none or one CVD risk factor). Right: Diseased individuals (two or more CVD risk factors).

`prior.h` arguments. The code follows.

```
R> pROC_normal <- pooledROC.dpm(marker = "bmi", group = "cvd_idf",
+   tag.h = 0, data = endosyn,
+   standardise = TRUE, p = seq(0, 1, l = 101), ci.level = 0.95,
+   compute.lpml = TRUE, compute.WAIC = TRUE, compute.DIC = TRUE,
+   pauc = pauccontrol(compute = TRUE, focus = "FPF", value = 0.1),
+   density = densitycontrol(compute = TRUE),
+   prior.h = priorcontrol.dpm(L = 1), prior.d = priorcontrol.dpm(L = 1),
+   mcmc = mcmccontrol(nsave = 8000, nburn = 2000, nskip = 1),
+   parallel = "snow", ncpus = 2, cl = NULL)
```

For the sake of space we omit from the summary the call to the function




Figure 4: Trace plots of the MCMC draws (after burn-in) of the PDFs of the BMI based on the model `pROC_dpm`. Results are shown separately for the nondiseased and diseased populations and for different values of the BMI.

```
R> summary(pROC_normal)
```

```
Call: [...]
```

```
Approach: Pooled ROC curve - Bayesian DPM
```

```
-----
```

```
Area under the pooled ROC curve: 0.748 (0.728, 0.768)*
```

```
Partial area under the pooled ROC curve (FPF = 0.1): 0.224 (0.194, 0.253)*
```

```
* Credible level: 0.95
```

```
Model selection criteria:
```

	Group H	Group D
WAIC	12639.952	4049.004
WAIC (Penalty)	2.431	2.267
LPML	-6319.976	-2024.502
DIC	12639.505	4048.714
DIC (Penalty)	1.986	1.987

```
Sample sizes:
```

```
Group H    Group D
```




Figure 5: Effective sample size and Geweke statistic of the MCMC chains (after burn-in) and of the PDFs of the BMI based on model `pROC_dpm` in the nondiseased and diseased populations. In both cases, results are shown along BMI values.

Number of observations	2149	691
Number of missing data	0	0

The fit of the DPM and normal models, in each group, can be compared on the basis of the WAIC, DIC, and/or the LPML. Remember that for the LPML, the higher its value, the better the model fit, while for the WAIC and DIC it is the other way around. By comparing these values, provided in the summary of each fitted model, we can conclude that the three criteria favour, in both the diseased and (especially in the) nondiseased groups, the more general DPM model. This is also corroborated by the plot of the fitted densities in each group shown in Figure 3(b).

We now estimate the pooled ROC curve using the empirical estimator (function `pooledROC.emp`), and comparisons with the results obtained using the DPM approach are provided.

```
R> pROC_emp <- pooledROC.emp(marker = "bmi", group = "cvd_idf",
+   tag.h = 0, data = endosyn, p = seq(0, 1, l = 101),
```

```
+ pauc = pauccontrol(compute = TRUE, focus = "FPF", value = 0.1), B = 500,
+ ci.level = 0.95, parallel = "snow", ncpus = 2)
```

```
R> summary(pROC_emp)
```

```
Call: [...]
```

```
Approach: Pooled ROC curve - Empirical
```

```
-----
Area under the pooled ROC curve: 0.76 (0.743, 0.778)*
```

```
Partial area under the pooled ROC curve (FPF = 0.1): 0.169 (0.14, 0.201)*
```

```
* Confidence level: 0.95
```

```
Sample sizes:
```

	Group H	Group D
Number of observations	2149	691
Number of missing data	0	0

Note that the posterior means for the AUC and pAUC obtained using the DPM method (0.759 and 0.168, respectively) are very similar to the point estimates using the empirical approach (0.760 and 0.169). This can also be observed when plotting the estimated ROC curves under the two methods (Figure 6).




Figure 6: Estimated ROC curve using the empirical approach (black line) and using the DPM method (red line—posterior mean).

We finish this section by showing how to use **ROCnReg** to obtain an (optimal) threshold value which could be further used to ‘diagnose’ an individual as diseased (two or more CVD risk

factors) or nondiseased (none or only one CVD risk factor). To that aim, and for `pooledROC` objects (i.e., those obtained using functions `pooledROC.dpm`, `pooledROC.BB`, `pooledROC.emp`, and `pooledROC.kernel`), we use the function `compute.threshold.pooledROC`, which allows obtaining (optimal) threshold values using two criteria: the YI and the one that sets a target value for the FPF. For illustration, we show here the results using the YI criterion.

```
R> th_pROC_dmp <- compute.threshold.pooledROC(pROC_dpm, criterion = "YI",
+   ci.level = 0.95, parallel = "snow", ncpus = 2)
R> th_pROC_dmp
```

```
$call
```

```
compute.threshold.pooledROC(object = pROC_dpm, criterion = "YI",
  ci.level = 0.95, parallel = "snow", ncpus = 2)
```

```
$thresholds
```

```
      est      ql      qh
26.46877 26.07129 26.85029
```

```
$YI
```

```
      est      ql      qh
0.4045776 0.3721684 0.4366298
```

```
$FPF
```

```
      est      ql      qh
0.3808336 0.3469580 0.4159478
```

```
$TPF
```

```
      est      ql      qh
0.7854112 0.7528575 0.8161865
```

The function returns the posterior mean (`est`) and `ci.level` \times 100% (here 95% since `ci.level` = 0.95) credible interval (lower bound: `ql`, upper bound: `qh`) for the YI and associated threshold value, as well as for the FPF and TPF associated with this cutoff value. For our example, the (posterior mean of the) YI is 0.40 and the YI-based threshold value is a BMI value of 26.5, which falls in the nutritional status defined as pre-obesity by the World Health Organization. By using this YI-based threshold value, we would have a FPF of 0.38 and a TPF of 0.79.

4.2. Covariate-specific ROC curve

We now turn our attention to the inclusion of covariates in ROC analysis. As shown in Table 1 and Section 3, with **ROCnReg** the user can estimate the covariate-specific ROC curve by means of three approaches: function `cROC.sp` implements the frequentist parametric and semiparametric induced ROC regression estimator, `cROC.kernel` corresponds to the nonparametric, kernel-based, counterpart of `cROC.sp`, and `cROC.bnp` stands for the Bayesian approach based on a single-weights dependent Dirichlet process mixture of normal distributions. As for the functions in **ROCnReg** for estimating the pooled ROC curve, the input arguments are method-specific, and we refer the reader to the manual for details. For

all methods, numerical and graphical summaries are obtained using functions `print.cROC`, `summary.cROC` and `plot.cROC`. Also, for objects of class `cROC.bnp`, **ROCnReg** provides the function `predictive.checks`, which implements tools for assessing model fit via posterior predictive checks.

Recall that, when including covariate information in ROC analysis, interest resides in evaluating if and how the discriminatory capacity of the test varies with such covariates. In particular, in our endocrine study we aim at evaluating the possible effect of both `age` and `gender` in the discriminatory capacity of the BMI. In what follows, we do that using the `cROC.bnp` function, and two different models are fitted. One which considers a normal distribution in each group and that incorporates the `age` effect in a linear way, and a second one which caps the maximum number of mixture components in each group at 10 (i.e., $L_D = L_{\bar{D}} = 10$) and that models the `age` effect using cubic B-splines (and thus allows for a nonlinear effect of age). Following Rodríguez-Álvarez *et al.* (2011b,a), both models consider the interaction between `age` and `gender`. For clarity, we first focus on the code that models `age` effect in a linear way, and use it to describe in detail the different arguments of the `cROC.bnp` function.

```
R> # Dataframe for predictions
R> agep <- seq(22, 80, l = 30)
R> endopred <- data.frame(age = rep(agep,2),
+   gender = factor(rep(c("Women", "Men"), each = length(agep))))

R> set.seed(123, "L'Ecuyer-CMRG") # for reproducibility
R> cROC_bp <- cROC.bnp(formula.h = bmi ~ gender*age,
+   formula.d = bmi ~ gender*age, group = "cvd_idf", tag.h = 0,
+   data = endosyn, newdata = endopred,
+   standardise = TRUE, p = seq(0, 1, l = 101), ci.level = 0.95,
+   compute.lpml = TRUE, compute.WAIC = TRUE, compute.DIC = TRUE,
+   pauc = pauccontrol(compute = FALSE),
+   prior.h = priorcontrol.bnp(L = 1), prior.d = priorcontrol.bnp(L = 1),
+   density = densitycontrol(compute = TRUE),
+   mcmc = mcmccontrol(nsave = 8000, nburn = 2000, nskip = 1),
+   parallel = "snow", ncpus = 2)
```

As can be seen, many arguments coincide with those of the function `pooledROC.dpm` (described in Section 4.1). We thus focus here on those that are specific to `cROC.bnp`. The arguments `formula.h` and `formula.d` are `formula` objects specifying the model for the regression functions (see Equation (33)) in, respectively, the nondiseased and diseased classes. They are similar to the formula used with the `glm` function, except that nonlinear functions (modelled by means of cubic B-splines) can be added using function `f` (an example will follow later in this section). Note that in both cases, the left-hand side of the formulas should include the name of the test/marker (in our case `bmi`). In our application, and for both groups, the model for the component's means includes, in addition to the linear effect of `age` and `gender`, the (linear) interaction between these two covariates (i.e., `gender*age` \equiv `gender + age + gender:age`). Through the `newdata` argument the user can specify a new data frame containing the values of the covariates at which the covariate-specific ROC curve and AUC (and also pAUC and PDFs, if required) are to be computed. Finally, `prior.h` (the same holds for `prior.d`) is an (optional) list of values to replace the defaults returned by `priorcontrol.bnp`

```
priorcontrol.bnp(m0 = NA, S0 = NA, nu = NA, Psi = NA, a = 2, b = NA,
  alpha = 1, L = 10)
```

which allows setting the hyperparameters for the single-weights dependent Dirichlet process mixture of normals model (see Section 3 and the manual accompanying the package for more details). In our example, we only modified the upper bound for the number of components in the mixture model (by default $L = 10$) and set it to 1. With this configuration, the model for the covariate-specific ROC curve can be regarded as a Bayesian counterpart of the induced ROC approach proposed by Faraggi (2003) and we denote it as the Bayesian normal linear model (for the test outcomes).

In this case, the `summary` of the fitted model provides the following information.

```
R> summary(cROC_bp)
```

```
Call: [...]
```

```
Approach: Conditional ROC curve - Bayesian nonparametric
```

```
-----
```

Parametric coefficients

Group H:

	Post. mean	Post. quantile 2.5%	Post. quantile 97.5%
(Intercept)	26.1459	25.8765	26.4096
genderWomen	-0.9160	-1.2726	-0.5680
age	1.1949	0.9180	1.4690
genderWomen:age	1.1948	0.8455	1.5394

Group D:

	Post. mean	Post. quantile 2.5%	Post. quantile 97.5%
(Intercept)	29.1865	28.7625	29.6115
genderWomen	2.0826	1.3705	2.7665
age	0.6578	0.2162	1.0904
genderWomen:age	-0.7711	-1.4655	-0.0956

ROC curve:

	Post. mean	Post. quantile 2.5%	Post. quantile 97.5%
(Intercept)	-0.6959	-0.8177	-0.5776
genderWomen	-0.6863	-0.8695	-0.5046
age	0.1229	0.0045	0.2415
genderWomen:age	0.4499	0.2745	0.6245
b	0.9391	0.8824	0.9975

Model selection criteria:

	Group H	Group D
--	---------	---------

WAIC	12174.986	4007.980
WAIC (Penalty)	6.283	5.646
LPML	-6087.492	-2003.990
DIC	12173.664	4007.329
DIC (Penalty)	4.994	5.053

Sample sizes:

	Group H	Group D
Number of observations	2149	691
Number of missing data	0	0

The first aspect to note is that, in this case, the `summary` function does not provide the estimated AUC as there is one (possibly different) AUC for each combination of covariate values. Also, given that: (1) only one component has been considered for modelling the CDFs of test results in the diseased and nondiseased groups, and (2) covariate effects have been modelled in a linear way, the `summary` function provides the posterior mean (and quantiles) of the (parametric) coefficients associated with the regression functions (see Equation (27)) and with the covariate-specific ROC curve (see Equation (30)). We note that since in the call to the function we have specified `standardise = TRUE` (and consequently both the test outcomes and covariates are standardised), the regression coefficients are on the scale of the standardised covariates. If we focus on the coefficients for the covariate-specific ROC curve, it seems that the discriminatory capacity of the BMI decreases with age, with the decrease being more pronounced in women (note that the expression of the covariate-specific ROC curve in (29) implies that positive coefficients correspond to a decrease in discriminatory capacity). These results are possibly better judged by plotting the estimated covariate-specific ROC curves and associated AUCs. This can be done using the `plot` function. For the covariate-specific ROC curve, the depicted graphics will depend on the number and nature of the covariates included in the analyses. In particular, for our application, we obtain, separately for men and women, the covariate-specific ROC curves (and AUCs) along age. These are shown in Figure 7, obtained using the code

```
R> op <- par(mfrow = c(2,2))
R> plot(cROC_sp, ask = FALSE)
R> par(op)
```

Although in this example we have modelled the `age` effect linearly and only one mixture component was considered, **ROCnReg** also allows for modelling the effect of continuous covariates in a nonlinear way, either using cubic B-spline basis expansions (through the function `cROC.bnp`) or kernel-based smoothers (via the function `cROC.kernel`). Also, as noted before, using only one mixture component for the single-weights dependent Dirichlet process mixture of normals model (function `cROC.bnp`) is equivalent to consider a (Bayesian) normal model, which might be too restrictive for most data applications. In what follows, we provide more flexibility to the model for the covariate-specific ROC curve by means of (1) increasing the number of mixture components, and (2) modelling the `age` effect in a nonlinear way (recall our considerations in Section ?? about the lack of flexibility of the single-weights dependent Dirichlet process mixture of normals model when covariates effects on the components’




Figure 7: Graphical results as provided by the `plot.cROC` function for an object of class `cROC.bnp`. Results for the model that includes the linear interaction between `age` and `gender` and one mixture component. Top row: Posterior mean of the covariate-specific ROC curve along age, separately for men and women. Bottom row: Posterior mean and 95% pointwise credible band for the covariate-specific AUC along age, separately for men and women.

means are modelled linearly). The former is done by modifying the value of `L` in the arguments `prior.h` and `prior.d`, with 10 being the default value. Regarding the latter, this is done by making use of the function `f` when specifying the component's mean functions through `formula.h` and `formula.d`. In particular, in our application we are interested in modelling the factor-by-curve interaction between `age` and `gender` (i.e., we model the `age` effect 'separately' for men and women). This is done using, e.g., `bmi ~ gender + f(age, by = gender, K = c(3,5))`. Through the argument `K` we indicate the number of internal knots that are used for constructing the cubic B-spline basis that is used to approximate the nonlinear effect of `age` (with the quantiles of `age` used to anchor the knots). Note that we

can specify a different number of internal knots for men and women ($K = c(3,5)$), where the order of vector K should match the ordering of the levels of the factor `gender`. We also note that to assist in the selection of the number of interior knots (in **ROCnReg** the location is always based on the quantiles of the corresponding covariates), the user can make use of the WAIC, DIC, and/or LPML. For instance, for this application, we fitted different models with different number of internal knots and we have chosen the model that provided the lowest WAIC (this was done in both the nondiseased and diseased populations and we remark that the number of knots does not need to be the same in the two populations). The final model is shown below.

```
R> # Levels of gender, and its ordering.
R> # Needed if we want to specify different
R> # number of knots for men and women
R> levels(endosyn$gender)

[1] "Men"    "Women"

R> set.seed(123, "L'Ecuyer-CMRG") # for reproducibility
R> cROC_bnp <- cROC.bnp(
+   formula.h = bmi ~ gender + f(age, by = gender, K = c(0,0))
+   formula.d = bmi ~ gender + f(age, by = gender, K = c(4,4)),
+   group = "cvd_idf", tag.h = 0, data = endosyn, newdata = endopred,
+   standardise = TRUE, p = seq(0, 1, l = 101), ci.level = 0.95,
+   compute.lpml = TRUE, compute.WAIC = TRUE, compute.DIC = TRUE,
+   pauc = pauccontrol(compute = FALSE),
+   prior.h = priorcontrol.bnp(L = 10), prior.d = priorcontrol.bnp(L = 10),
+   density = densitycontrol(compute = TRUE),
+   mcmc = mcmccontrol(nsave = 8000, nburn = 2000, nskip = 1),
+   parallel = "snow", ncpus = 2)

R> summary(cROC_bnp)
```

Call: [...]

Approach: Conditional ROC curve - Bayesian nonparametric

Model selection criteria:

	Group H	Group D
WAIC	11833.000	3909.828
WAIC (Penalty)	31.236	38.583
LPML	-5916.766	-1955.449
DIC	11829.750	3904.532
DIC (Penalty)	29.611	35.934

Sample sizes:

	Group H	Group D
Number of observations	2149	691
Number of missing data	0	0

```
R> op <- par(mfrow = c(2,2))
R> plot(cROC_sp, ask = FALSE)
R> par(op)
```




Figure 8: Graphical results as provided by the `plot.cROC` function for an object of class `cROC.bnp`. Results for the model that includes the factor-by-curve interaction between age and gender and 10 mixture components. Top row: Posterior mean of the covariate-specific ROC curve along age, separately for men and women. Bottom row: Posterior mean and 95% pointwise credible band for the covariate-specific AUC along age, separately for men and women.

The graphical results are shown in Figure 8. Note that, especially for women, age displays a

marked nonlinear effect. Recall that for objects of class `cROC.bnp`, and if required in the call to the function, the `summary` function provides, separately for the diseased and nondiseased groups, the WAIC, LPML, and DIC. Note that, in both cases, the three criteria support the use of the more flexible model that uses cubic B-splines and 10 mixture components for modelling the distribution of the BMI (model `cROC$_bnp`) over the more restrictive Bayesian normal linear model (model `cROC$_bp`). Because the WAIC, LPML, and DIC are relative criteria, posterior predictive checks are also available in **ROCnReg** through the function `predictive.checks`. Specifically, the function generates replicated datasets from the posterior predictive distribution in both class D and \bar{D} and compares them to the test values (BMI values in our application) using specific statistics. For the choice of such statistics we follow [Gabry, Simpson, Vehtari, Betancourt, and Gelman \(2019\)](#), who suggest choosing statistics that are ‘orthogonal’ to the model parameters. Since we are using a location-scale mixture of normals model for the test outcomes, we use here the skewness and kurtosis and check how well the posterior predictive distribution captures these two quantities.

```
R> op <- par(mfrow = c(2,3))
R> pc_cROC_bp <- predictive.checks(cROC_bp,
+   statistics = c("kurtosis", "skewness"), devnew = FALSE)
R> par(op)
```

```
R> op <- par(mfrow = c(2,3))
R> pc_cROC_bnp <- predictive.checks(cROC_bnp,
+   statistics = c("kurtosis", "skewness"), devnew = FALSE)
R> par(op)
```


Results are shown in Figure 9. As can be seen, the model that includes the factor-by-curve interaction between `age` and `gender` and 10 mixture components performs quite well in capturing both quantities, while the Bayesian normal linear model fails to do so. Also shown in Figure 9 (and provided by function `predictive.checks`) are the kernel density estimates of 500 randomly selected datasets drawn from the posterior predictive distribution, in each group, compared to the kernel density estimate of the observed BMI (in each group). Again, the more flexible model, as opposed to the Bayesian normal linear model, is able to simulate data that are very much similar to the observed BMI values.

As for the pooled ROC curve (Section 4.1), **ROCnReg** also provides a function that allows obtaining (optimal) threshold values for the covariate-specific ROC curve. For illustration, instead of the threshold values based on the Youden index, we now use the criterion that sets a target value for the FPF. The code for model `cROC_bnp`, when setting the $FPF = 0.3$, is as follows.


```
R> th_fpf_cROC_bnp <- compute.threshold.cROC(cROC_bnp,
+   criterion = "FPF", FPF = 0.3, newdata = endopred,
+   ci.level = 0.95, parallel = "snow", ncpus = 2)
R> names(th_fpf_cROC_bnp)
```

```
[1] "newdata"      "thresholds" "TPF"        "FPF"        "call"
```

In addition to the data frame `newdata` containing the covariate values at which the thresholds are computed, the function `compute.threshold.cROC` also returns the covariate-specific



(a) Model including the linear interaction between **age** and **gender** and one component



(b) Model including the factor-by-curve interaction between **age** and **gender** and 10 mixture components

Figure 9: Graphical results as provided by the `predictive.checks` function for an object of class `cROC.bnp`. Histograms of the statistics *skewness* and *kurtosis* computed from 8000 draws from the posterior predictive distribution in the diseased and nondiseased group. The red line is the estimated statistic from the observed BMI values. The right-hand side plots show the kernel density estimate of the observed BMI (solid black line), jointly with the kernel density estimates for 500 simulated datasets drawn from the posterior predictive distributions.

`thresholds` corresponding to the specified FPF as well as the covariate-specific TPF attached to these thresholds. In both cases, the function returns the posterior mean and the `ci.level` $\times 100\%$ (here 95%) pointwise credible intervals. Although **ROCnReg** does not pro-

vide a function for plotting the results obtained using `compute.threshold.cROC`, graphical results can be easily obtained. For simplicity, we only show here the code for the covariate-specific threshold values (`thresholds`), but a similar code can be used to plot the covariate-specific TPFs. Both plots are shown in Figure 10. As can be observed, for a FPF of 0.3, the BMI age-specific thresholds tend to increase with age both for men and women, although for the latter there is a slight decrease after an age of about 70 years old. The age-specific TPFs corresponding to the thresholds for which the FPF is 0.3 show a nonlinear behaviour and these are in general higher for women than for men (of the same age).

```
df <- data.frame(age = th_fpf_cROC_bnp$newdata$age,
+   gender = th_fpf_cROC_bnp$newdata$gender,
+   y = th_fpf_cROC_bnp$thresholds[[1]][,"est"],
+   ql = th_fpf_cROC_bnp$thresholds[[1]][,"ql"],
+   qh = th_fpf_cROC_bnp$thresholds[[1]][,"qh"])

R> g0 <- ggplot(df, aes(x = age, y = y, ymin = ql, ymax = qh)) +
+   geom_line() +
+   geom_ribbon(alpha = 0.2) +
+   labs(title = "Covariate-specific thresholds for a FPF = 0.3",
+   x = "Age (years)", y = "BMI") +
+   theme(strip.text.x = element_text(size = 20),
+   plot.title = element_text(hjust = 0.5, size = 20),
+   axis.text = element_text(size = 20),
+   axis.title = element_text(size = 20)) +
+   facet_wrap(~gender)
R> print(g0)
```

For conciseness we have not shown here how to perform convergence diagnostics of the MCMC chains for models fitted using the function `cROC.bnp`. In very much the same way as shown in the previous section for the object `pROC.dpm`, using the information contained in component `dens` in the list of returned values (if required), one can produce trace plots of the conditional densities at some sampled values, as well as obtain the corresponding effective sample sizes and Geweke statistics. Some results are provided in the Appendices, and the associated code can be found in the replication code that accompanies this paper.

4.3. Covariate-adjusted ROC curve

In this section we illustrate how to conduct inference about the covariate-adjusted ROC curve using **ROCnReg**. Similarly to the covariate-specific ROC curve, three approaches are available, namely, function `AROC.sp` implements the frequentist approaches that postulate that test outcomes in the nondiseased group follow a linear model with the CDF of the error term being either a standard normal distribution or estimated via the empirical CDF of the standardised residuals, `AROC.kernel` corresponds to the kernel-based counterpart, and `AROC.bnp` implements the Bayesian (nonparametric) approach based on a single-weights dependent Dirichlet process mixture of normal distributions and the Bayesian bootstrap.

Recall that the AROC curve is a global summary measure of diagnostic accuracy that takes covariate information into account. In the context of our endocrine application we seek to




Figure 10: Top row: Posterior mean (solid black line) and 95% pointwise credible band for the BMI threshold values, along age, corresponding to a FPF of 0.3. Bottom row: Posterior mean (solid black line) and 95% pointwise credible band of the TPFs, along age, corresponding to the BMI threshold values for which FPF = 0.3.

study the overall discriminatory capacity of the BMI for detecting the presence of CVD risk factors when adjusting for age and gender. Here we focus on how to estimate the AROC curve using the `AROC.bnp` function. The function syntax is exactly similar to the one of `cROC.bnp`, with the only difference being that we only need to specify the arguments related to the nondiseased population. The code and respective summary follow.

```
R> set.seed(123, "L'Ecuyer-CMRG") # for reproducibility
R> AROC_bnp <- AROC.bnp(
+   formula.h = bmi ~ gender + f(age, by = gender, K = c(0,0)),
+   group = "cvd_idf", tag.h = 0, data = endosyn, standardise = TRUE,
+   p = seq(0, 1, l = 101), ci.level = 0.05,
+   compute.lpml = TRUE, compute.WAIC = TRUE, compute.DIC = TRUE,
```

```
+ pauc = pauccontrol(compute = FALSE), prior = priorcontrol.bnp(L = 10),
+ density = densitycontrol.aroc(compute = FALSE),
+ mcmc = mcmccontrol(nsave = 8000, nburn = 2000, nskip = 1),
+ parallel = "snow", ncpus = 2)
```

```
R> summary(AROC_bnp)
```

```
Call: [...]
```

```
Approach: AROC Bayesian nonparametric
```

```
-----
Area under the covariate-adjusted ROC curve: 0.656 (0.629, 0.684)*
* Credible level: 0.95
```

```
Model selection criteria:
```

	Group H
WAIC	11833.000
WAIC (Penalty)	31.236
LPML	-5916.766
DIC	11829.750
DIC (Penalty)	29.611

```
Sample sizes:
```

	Group H	Group D
Number of observations	2149	691
Number of missing data	0	0

The area under the AROC curve is 0.656 (95% credible interval: (0.629, 0.684)) thus revealing a reasonable good ability of the BMI to detect the presence of CVD risk factors when teasing out the age and gender effects. As for the pooled ROC curve and the covariate-specific ROC curve, a plot function is also available (result in Figure 11(a)).

```
R> plot(AROC_bnp, cex.main = 1.5, cex.lab = 1.5, cex.axis = 1.5, cex = 1.3)
```

We finish with a comparison of the AROC curve with the pooled ROC curve that was obtained earlier by using a DPM model with 10 components in each group. In Figure 11(b) we show the plots of the two curves and, as can be noticed, the pooled ROC curve lies well above the AROC curve, thus evidencing the need for incorporating covariate information into the analysis.

```
R> plot(AROC_bnp$p, AROC_bnp$ROC[,1],
+ type = "l", xlim = c(0,1), ylim = c(0,1),
+ xlab = "FPF", ylab = "TPF",
+ main = "Pooled ROC curve vs AROC curve",
+ cex.main = 1.5, cex.lab = 1.5, cex.axis = 1.5, cex = 1.5)
R> lines(AROC_bnp$p, AROC_bnp$ROC[,2], col = 1, lty = 2)
```

```

R> lines(AROC_bnp$p, AROC_bnp$ROC[,3], col = 1, lty = 2)
R> lines(pROC_dpm$p, pROC_dpm$ROC[,1], col = 2)
R> lines(pROC_dpm$p, pROC_dpm$ROC[,2], col = 2, lty = 2)
R> lines(pROC_dpm$p, pROC_dpm$ROC[,3], col = 2, lty = 2)
R> abline(0, 1, col = "grey", lty = 2)

```




Figure 11: (a) Age and gender-adjusted ROC curve: posterior mean and 95% pointwise credible band. (b) Age and gender-adjusted ROC curve (in black) and pooled ROC curve (estimated using a DPM of normals model) (in red). Solid lines represent the posterior means and dashed lines the 95% pointwise credible bands.

4.4. Computational aspects

We finish this section with some comments on computational aspects. In our experience, the methods with the largest computing times are those implemented in `cROC.bnp` when $L_{\bar{D}} > 1$ and in `cROC.kernel` when confidence bands are to be constructed. In the first case, the main reason behind the computational burden is the need to invert $F_{\bar{D}}(\cdot | \mathbf{x})$ in order to obtain the covariate-specific ROC curve (see equation (10)). Note that when $L_{\bar{D}} > 1$, the conditional distribution function in the nondiseased group is given by a mixture of normal distributions and the corresponding quantile function needs to be computed for each covariate(s) value we might be interested in and for each iteration of the Gibbs sampler procedure. Regarding the `cROC.kernel` function, the computing time is mainly driven by the number of bootstrap samples used for constructing the confidence bands. In Table 2 we show the time, in seconds, needed for fitting the pooled, the covariate-specific, and the covariate-adjusted ROC curve using the Bayesian nonparametric and the kernel approaches for the synthetic endocrine data and when both parallel (with 2 and 4 processes) and no parallel options are used. We note that for the Bayesian approaches we also computed the densities/conditional densities, as

well as the WAIC, LPML, and DIC, which further increase the computing time (in the case of the AROC curve these were only computed in the nondiseased population). With respect to the kernel-based approach (in this case the fit is done separately for men and women and the corresponding results are presented in the Supplementary Material) we have used 500 bootstrap samples to construct the confidence bands. As it can be appreciated, for these two intense tasks, using 4 processes drastically improves the computation time. All computations were performed in a iMac with 3.6GHz quad core Intel i7 processor and 32GB RAM running under a macOS Catalina 10.15.5 operating system.

	No parallel	Snow (2 cores)	Snow (4 cores)
<code>pooledROC.dpm</code>	138	118	111
<code>pooledROC.kernel</code>	376	196	105
<code>cROC.bnp</code>	2052	1117	680
<code>cROC.kernel</code>	Men: 1159 Women: 1885	Men: 528 Women: 916	Men: 279 Women: 466
<code>AROC.bnp</code>	126	115	112
<code>AROC.kernel</code>	Men: 847 Women: 1707	Men: 404 Women: 833	Men: 214 Women: 438

Table 2: Time in seconds (rounded to the nearest second) needed to fit the pooled, the covariate-specific, and the covariate-adjusted ROC curve for the Bayesian nonparametric and kernel approaches.

5. Summary and future plans

In this paper we have introduced the capabilities of the R package **ROCnReg** for conducting inference about the pooled ROC curve, the covariate-specific ROC curve, and the covariate-adjusted ROC curve and their associated summary indices. As we have illustrated, the current version of the package provides several options for estimating ROC curves, both under frequentist and Bayesian paradigms, either parametrically, semiparametrically, or nonparametrically. To the best of our knowledge, this is the first software package implementing Bayesian inference for ROC curves. Several additions/extensions are planned in the future and these, among others, include:

- Implement the most time-consuming parts in C or C++.
- Incorporate methods for non-binary disease status (e.g., no disease, mild disease, severe disease). That is, implement ROC surface models.
- Implement new (optimal) threshold criteria (e.g., YI including costs).

Computational details

The results in this paper were obtained using R 4.0.3 with the **ROCnReg** 1.0-5 package. The **ROCnReg** package has multiple dependencies: **graphics**, **grDevices**, **parallel**, **splines**, **stats**, **moments** (Komsta and Novomestky 2015), **nor1mix** (Maechler 2019), **Matrix** (Bates

and Maechler 2019), **spatstat** (Baddeley and Turner 2005), **np** (Hayfield and Racine 2008), **lattice** (Sarkar 2008), **MASS** (Venables and Ripley 2002) and **pbivnorm** (Genz and Kenkel 2015). R itself and all packages used are available from the Comprehensive R Archive Network (CRAN) at <https://CRAN.R-project.org/>.

Acknowledgments

MX Rodríguez-Álvarez was funded by project MTM2017-82379-R (AEI/FEDER, UE), by the Basque Government through the BERC 2018-2021 program and Elkartek project 3KIA (KK-2020/00049), and by the Spanish Ministry of Science, Innovation, and Universities (BCAM Severo Ochoa accreditation SEV-2017-0718).

References

- Baddeley A, Turner R (2005). “**spatstat**: An R Package for Analyzing Spatial Point Patterns.” *Journal of Statistical Software*, **12**(6), 1–42. URL <http://www.jstatsoft.org/v12/i06/>.
- Bates D, Maechler M (2019). **Matrix**: *Sparse and Dense Matrix Classes and Methods*. R package version 1.2-17, URL <https://CRAN.R-project.org/package=Matrix>.
- Buolamwini JA (2017). *Gender Shades: Intersectional Phenotypic and Demographic Evaluation of Face Datasets and Gender Classifiers*. Ph.D. thesis, Massachusetts Institute of Technology, School of Architecture and Planning, Program in Media Arts and Sciences.
- Celeux G, Forbes F, Robert CP, Titterton DM (2006). “Deviance Information Criteria for Missing Data Models.” *Bayesian Analysis*, **1**(4), 651–673.
- De Iorio M, Johnson WO, Müller P, Rosner GL (2009). “Bayesian Nonparametric Nonproportional Hazards Survival Modeling.” *Biometrics*, **65**(3), 762–771.
- Dunn PK, Smyth GK (1996). “Randomized Quantile Residuals.” *Journal of Computational and Graphical Statistics*, **5**(3), 236–244.
- Erkanli A, Sung M, Jane Costello E, Angold A (2006). “Bayesian Semi-Parametric ROC Analysis.” *Statistics in Medicine*, **25**(22), 3905–3928.
- Fan J, Gijbels I (1996). *Local Polynomial Modelling and Its Applications*. Monographs on Statistics and Applied Probability. Chapman & Hall/CRC.
- Faraggi D (2003). “Adjusting Receiver Operating Characteristic Curves and Related Indices for Covariates.” *Journal of the Royal Statistical Society D*, **52**(2), 179–192.
- Ferguson TS (1973). “A Bayesian Analysis of Some Nonparametric Problems.” *The Annals of Statistics*, **1**(2), 209–230.
- Gabry J, Simpson D, Vehtari A, Betancourt M, Gelman A (2019). “Visualization in Bayesian Workflow.” *Journal of the Royal Statistical Society A*, **182**(2), 389–402.

- Geisser S, Eddy WF (1979). “A Predictive Approach to Model Selection.” *Journal of the American Statistical Association*, **74**(365), 153–160.
- Gelman A, Carlin JB, Stern HS, Dunson DB, Vehtari A, Rubin DB (2013). *Bayesian Data Analysis*. Chapman and Hall/CRC.
- Gelman A, Hwang J, Vehtari A (2014). “Understanding Predictive Information Criteria for Bayesian Models.” *Statistics and Computing*, **24**(6), 997–1016.
- Genz A, Kenkel B (2015). **pbivnorm**: *Vectorized Bivariate Normal CDF*. R package version 0.6.0, URL <https://CRAN.R-project.org/package=pbivnorm>.
- Gonçalves L, Subtil A, Oliveira MR, Bermudez P (2014). “ROC Curve Estimation: An Overview.” *REVSTAT–Statistical Journal*, **12**(1), 1–20.
- González-Manteiga W, Pardo-Fernández JC, van Keilegom I (2011). “ROC Curves in Non-Parametric Location-Scale Regression Models.” *Scandinavian Journal of Statistics*, **38**(1), 169–184.
- Gu J, Ghosal S, Roy A (2008). “Bayesian Bootstrap Estimation of ROC Curve.” *Statistics in Medicine*, **27**, 5407–5420.
- Guan Z, Qin J, Zhang B (2012). “Information Borrowing Methods for Covariate-Adjusted ROC Curve.” *Canadian Journal of Statistics*, **40**(3), 569–587.
- Hayfield T, Racine JS (2008). “Nonparametric Econometrics: The **np** Package.” *Journal of Statistical Software*, **27**(5). URL <http://www.jstatsoft.org/v27/i05/>.
- Hillis SL, Metz CE (2012). “An Analytic Expression for the Binormal Partial Area under the ROC Curve.” *Academic Radiology*, **19**(12), 1491 – 1498.
- Hsieh F, Turnbull BW (1996). “Nonparametric and Semiparametric Estimation of the Receiver Operating Characteristic Curve.” *The Annals of Statistics*, **24**(1), 25–40.
- Hutchinson B, Mitchell M (2019). “50 Years of Test (Un)Fairness: Lessons for Machine Learning.” In *Proceedings of the Conference on Fairness, Accountability, and Transparency, FAT* ’19*, pp. 49–58. ACM, New York, NY, USA. ISBN 978-1-4503-6125-5.
- Inácio V, Rodríguez-Álvarez MX, Gayoso-Diz P (2020). “Statistical evaluation of medical tests.” *Annual Review of Statistics and Its Application*, **8**.
- Inácio de Carvalho V, de Carvalho M, Branscum AJ (2017). “Nonparametric Bayesian Covariate-Adjusted Estimation of the Youden Index.” *Biometrics*, **73**(4), 1279–1288.
- Inácio de Carvalho V, Jara A, Hanson TE, de Carvalho M (2013). “Bayesian Nonparametric ROC Regression Modeling.” *Bayesian Analysis*, **8**(3), 623–646.
- Inácio de Carvalho V, Rodríguez-Álvarez MX (2018). “Bayesian Nonparametric Inference for the Covariate-Adjusted ROC Curve.” *arXiv:1806.00473 [stat.ME]*.
- Ishwaran H, James LF (2001). “Gibbs Sampling Methods for Stick-Breaking Priors.” *Journal of the American Statistical Association*, **96**(453), 161–173.

- Ishwaran H, James LF (2002). “Approximate Dirichlet Process Computing in Finite Normal Mixtures: Smoothing and Prior Information.” *Journal of Computational and Graphical Statistics*, **11**(3), 508–532.
- Janes H, Pepe MS (2009). “Adjusting for Covariate Effects on Classification Accuracy using the Covariate-Adjusted Receiver Operating Characteristic Curve.” *Biometrika*, **96**(2), 371–382.
- Komsta L, Novomestky F (2015). **moments**: Moments, cumulants, skewness, kurtosis and related tests. R package version 0.14, URL <https://CRAN.R-project.org/package=moments>.
- López-Ratón M, Rodríguez-Álvarez MX, Cadarso-Suárez C, Gude-Sampedro F (2014). “**OptimalCutpoints**: an R Package for Selecting Optimal Cutpoints in Diagnostic Tests.” *Journal of Statistical Software*, **61**(8), 1–36.
- MacEachern SN (2000). “Dependent Dirichlet Processes.” *Unpublished manuscript, Department of Statistics, The Ohio State University*, pp. 1–40.
- Maechler M (2019). **nor1mix**: Normal aka Gaussian (1-d) Mixture Models (S3 Classes and Methods). R package version 1.3-0, URL <https://CRAN.R-project.org/package=nor1mix>.
- Metz CE (1978). “Basic Principles of ROC Analysis.” *Seminars in Nuclear Medicine*, **8**(4), 283 – 298.
- Pardo-Fernández JC, Rodríguez-Álvarez MX, van Keilegom I (2014). “A Review on ROC Curves in the Presence of Covariates.” *REVSTAT–Statistical Journal*, **12**(1), 21–41.
- Pepe MS (1998). “Three Approaches to Regression Analysis of Receiver Operating Characteristic Curves for Continuous Test Results.” *Biometrics*, **54**(1), 124–135.
- Pepe MS (2003). *The Statistical Evaluation of Medical Tests for Classification and Prediction*. Oxford Statistical Sciences Series. Oxford University Press, New York.
- Pérez Fernández S, Martínez Camblor P, Filzmoser P, Corral Blanco N (2018). “nsROC: An R package for non-standard ROC curve analysis.” *The R Journal*, **10**(2).
- Perez Jaume S, Skaltsa K, Pallarès N, Carrasco Jordan JL (2017). “**ThresholdROC**: Optimum Threshold Estimation Tools for Continuous Diagnostic Tests in R.” *Journal of Statistical Software*, **82**(4), 1–21.
- Plummer M, Best N, Cowles K, Vines K (2006). “CODA: Convergence Diagnosis and Output Analysis for MCMC.” *R News*, **6**(1), 7–11. URL <https://journal.r-project.org/archive/>.
- R Core Team (2020). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez JC, Müller M (2011). “**pROC**: An Open-Source Package for R and S+ to Analyze and Compare ROC Curves.” *BMC Bioinformatics*, **12**, 77.

- Rodríguez-Álvarez MX, Roca-Pardinas J (2017). **npROCRegression**: *Kernel-Based Non-parametric ROC Regression Modelling*. R package version 1.0-5, URL <https://CRAN.R-project.org/package=npROCRegression>.
- Rodríguez-Álvarez MX, Roca-Pardiñas J, Cadarso-Suárez C (2011a). “A New Flexible Direct ROC Regression Model: Application to the Detection of Cardiovascular Risk Factors by Anthropometric Measures.” *Computational Statistics & Data Analysis*, **55**(12), 3257–3270.
- Rodríguez-Álvarez MX, Roca-Pardiñas J, Cadarso-Suárez C (2011b). “ROC curve and Co-variates: Extending Induced Methodology to the Non-Parametric Framework.” *Statistics and Computing*, **21**(4), 483–499.
- Rodríguez-Álvarez MX, Tahoces PG, Cadarso-Suárez C, Lado MJ (2011c). “Comparative Study of ROC Regression Techniques – Applications for the Computer-Aided Diagnostic System in Breast Cancer Detection.” *Computational Statistics & Data Analysis*, **55**(1), 888–902.
- Rosenberg PS (1995). “Hazard Function Estimation Using B-Splines.” *Biometrics*, **51**(3), 874–887.
- Sarkar D (2008). **Lattice**: *Multivariate Data Visualization with R*. Springer-Verlag, New York. ISBN 978-0-387-75968-5, URL <http://lmdvr.r-forge.r-project.org>.
- Shapiro DE (1999). “The Interpretation of Diagnostic Tests.” *Statistical Methods in Medical Research*, **8**(2), 113–134.
- Silverman BW (1986). *Density Estimation for Statistics and Data Analysis*. Chapman & Hall/CRC.
- Tomé Martínez de Rituerto MA, Botana MA, Cadarso-Suárez C, Rego-Iraeta A, Fernández-Mariño A, Mato JA, Solache I, Perez-Fernandez R (2009). “Prevalence of Metabolic Syndrome in Galicia (NW Spain) on Four Alternative Definitions and Association with Insulin Resistance.” *Journal of Endocrinological Investigation*, **32**(6), 505–511.
- Venables WN, Ripley BD (2002). *Modern Applied Statistics with S*. Fourth edition. Springer, New York. ISBN 0-387-95457-0, URL <http://www.stats.ox.ac.uk/pub/MASS4>.
- Wand MP, Jones MC (1994). *Kernel Smoothing*. Chapman & Hall/CRC.
- Wang XF (2012). **sROC**: *Nonparametric Smooth ROC Curves for Continuous Data*. R package version 0.1-2, URL <https://CRAN.R-project.org/package=sROC>.
- Wickham H (2016). **ggplot2**: *Elegant Graphics for Data Analysis*. Springer-Verlag New York. ISBN 978-3-319-24277-4. URL <https://ggplot2.tidyverse.org>.
- Youden WJ (1950). “Index for Rating Diagnostic Tests.” *Cancer*, **3**(1), 32–35.
- Zhou XH, McClish DK, Obuchowski NA (2011). *Statistical Methods in Diagnostic Medicine*. John Wiley & Sons, New York.
- Zou KH, Hall WJ, Shapiro DE (1997). “Smooth Non-Parametric Receiver Operating Characteristic (ROC) Curves for Continuous Diagnostic Tests.” *Statistics in Medicine*, **16**(19), 2143–2156.

Zou KH, Tempany CM, Fielding JR, Silverman SG (1998). “Original Smooth Receiver Operating Characteristic Curve Estimation from Continuous Data: Statistical Methods for Analyzing the Predictive Value of Spiral CT of Ureteral Stones.” *Academic Radiology*, **5**(10), 680–687.

A. Further computational tools for pooled ROC curve

A.1. Informal model diagnostics for the Bayesian methods: quantile residuals

We illustrate the use of quantile residuals (Dunn and Smyth 1996) which can be helpful for models fitted using the function `pooledROC.dpm`, for instance, in deciding if $L_D = 1$ or $L_D > 1$ (the same obviously applies to the nondiseased group). Quantile residuals are based on the well-known fact that for a continuous random variable, say Y_D , with CDF given by F_D , one has that $F_D(Y_D) \sim U(0, 1)$. As a consequence, quantile residuals defined by $\hat{r}_{Dj} = \Phi^{-1}\{\hat{F}_D(y_{Dj})\}$, for $j = 1, \dots, n_D$, should follow, approximately, a standard normal distribution if a correct model has been specified. A quantile-quantile (QQ) plot can then be used to determine deviations of the quantile residuals from the standard normal distribution. Below we provide the code to construct, for the diseased population, such QQ plot using output from the object `pROC_dpm` (that assumed $L_D = L_{\bar{D}} = 10$) obtained using the function `pooledROC.dpm`. The code for the nondiseased population follows in a similar manner, and is provided in the R replication code that accompanies this paper.

```
R> library("nor1mix")
R> traj <- matrix(0, nrow = pROC_dpm$mcmc$nsave,
+   ncol = length(pROC_dpm$marker$d))
R> lgrid <- length(pROC_dpm$marker$d)
R> grid <- qnorm(ppoints(lgrid))
R> for (l in 1:pROC_dpm$mcmc$nsave) {
+   aux <- norMix(mu = pROC_dpm$fit$d$mu[l,],
+   sigma = pROC_dpm$fit$d$sd[l,],
+   w = pROC_dpm$fit$d$probs[l,])
+   traj[l, ] <- quantile(qnorm(pnorMix(pROC_dpm$marker$d, aux)),
+   ppoints(lgrid), type = 2)
+ }
R> l.band <- apply(traj, 2, quantile, prob = 0.025)
R> trajhat <- apply(traj, 2, mean)
R> u.band <- apply(traj, 2, quantile, prob = 0.975)

R> op <- par(pty = "s")
R> plot(grid, trajhat, xlab = "Theoretical Quantiles",
+   ylab = "Sample Quantiles", main = "Nondiseased population",
+   cex.main = 2, cex.lab = 1.5, cex.axis = 1.5)
R> lines(grid, l.band, lty = 2, lwd = 2)
R> lines(grid, u.band, lty = 2, lwd = 2)
R> abline(a = 0, b = 1, col = "red", lwd = 2)
R> par(op)
```

The resulting QQ plots are shown in Figure 12(a) and show virtually no deviations from the standard normal distribution quantiles, thus revealing a good fit of the DPM model that assumes 10 mixture components in both the diseased and nondiseased groups. In contrast, those QQ plots obtained when fitting a normal model in each group (i.e., $L_D = L_{\bar{D}} = 1$;

model `pROC_normal` in the main manuscript), clearly show some deviations from the assumed normal distribution quantiles (see Figure 12(b)). The code used follows.

```
R> traj_normal <- matrix(0, nrow = pROC_normal$mcmc$nsave,
+   ncol = length(pROC_normal$marker$d))
R> lgrid_normal <- length(pROC_normal$marker$d)
R> grid_normal <- qnorm(ppoints(lgrid_normal))
R> for (l in 1:pROC_normal$mcmc$nsave) {
+   traj_normal[l, ] <- quantile(qnorm(pnorm(pROC_normal$marker$d,
+   pROC_normal$fit$d$mu[l], pROC_normal$fit$d$sd[l])),
+   ppoints(lgrid_normal), type = 2)
+ }
R> l.band_normal <- apply(traj_normal, 2, quantile, prob = 0.025)
R> trajhat_normal <- apply(traj_normal, 2, mean)
R> u.band_normal <- apply(traj_normal, 2, quantile, prob = 0.975)

R> op <- par(pty = "s")
R> plot(grid, trajhat, xlab = "Theoretical Quantiles",
+   ylab = "Sample Quantiles", main = "Nondiseased population",
+   cex.main = 2, cex.lab = 1.5, cex.axis = 1.5)
R> lines(grid, l.band, lty = 2, lwd = 2)
R> lines(grid, u.band, lty = 2, lwd = 2)
R> abline(a = 0, b = 1, col = "red", lwd = 2)
R> par(op)
```

A.2. Bayesian bootstrap and kernel estimators of the pooled ROC curve

The following code is used to fit the Bayesian bootstrap ROC curve estimator for the pooled ROC curve (function `pooledROC.BB`). The number of iterations considered is $B = 5000$ and, for the sake of illustration, we also compute the partial area under the curve corresponding to true positive fractions (TPF) or sensitivities between 0.8 and 1.

```
R> set.seed(123, "L'Ecuyer-CMRG") # for reproducibility
R> pROC_BB <- pooledROC.BB(marker = "bmi", group = "cvd_idf",
+   tag.h = 0, data = endosyn, p = seq(0, 1, l = 101), ci.level = 0.95,
+   pauc = pauccontrol(compute = TRUE, focus = "TPF", value = 0.8),
+   B = 5000, parallel = "snow", ncpus = 2)

R> summary(pROC_BB)
```

Call: [...]

Approach: Pooled ROC curve - Bayesian bootstrap

Area under the pooled ROC curve: 0.76 (0.74, 0.779)*

Partial area under the pooled ROC curve (FPF = 0.1): 0.17 (0.14, 0.201)*




Figure 12: Quantile residuals of the BMI data versus the theoretical quantiles of the standard normal distribution. The circles represent the posterior mean quantiles over all posterior samples, while the dashed lines represent the corresponding 95% credible bands. Top row: DPM model with 10 components in each group (model `pROC_dpm` in the main manuscript). Bottom row: normal model in each group (model `pROC_normal` in the main manuscript).

* Credible level: 0.95

Sample sizes:

	Group H	Group D
Number of observations	2149	691
Number of missing data	0	0

Note that all partial areas' values returned are normalised and, as such, what is being reported, in this case, is

$$\text{pAUC}_{\text{TPF}(0.8)}/(1 - 0.8).$$

The estimated pooled ROC curve and AUC (posterior means), jointly with 95% credible intervals, are obtained as follows

```
R> plot(pROC_BB, cex.main = 1.5, cex.lab = 1.5, cex.axis = 1.5, cex = 1.5)
```

The result is shown in Figure 13.




Figure 13: Graphical results as provided by the `plot.pooledROC` function for an object of class `pooledROC.BB`. Posterior mean and 95% pointwise credible band of the pooled ROC curve and corresponding AUC (posterior mean and 95% credible interval).

We shall present now the syntax associated to the kernel estimator of the pooled ROC curve (function `pooledROC.kernel`). In terms of arguments, `bw` specifies how the bandwidth should be computed, with `SRT` standing for Silverman's rule of thumb and `UCV` for least squares cross-validation. Additionally, here `B` stands for the number of bootstrap replications used to compute the confidence intervals/bands. The syntax follows.

```
R> set.seed(123, "L'Ecuyer-CMRG") # for reproducibility
R> pROC_kernel <- pooledROC.kernel(marker = "bmi", group = "cvd_idf",
+   tag.h = 0, data = endosyn, p = seq(0, 1, l = 101), bw = "SRT",
+   B = 500, ci.level = 0.95, method = "coutcome",
+   pauc = pauccontrol(compute = TRUE, focus = "TPF", value = 0.8),
+   parallel = "snow", ncpus = 2)

R> summary(pROC_kernel)
```

Call: [...]

Approach: Pooled ROC curve - Kernel-based

 Area under the pooled ROC curve: 0.755 (0.737, 0.774)*

Partial area under the specificity pooled ROC curve (Se = 0.8): 0.408 (0.373, 0.446)*

* Confidence level: 0.95

	Group H	Group D
Bandwidths:	0.867	1.019

Bandwidth Selection Method: Silverman's rule-of-thumb

Sample sizes:

	Group H	Group D
Number of observations	2149	691
Number of missing data	0	0

Graphical results are obtained with the following code, and present in Figure 14

```
R> plot(pROC_kernel, cex.main = 1.5, cex.lab = 1.5, cex.axis = 1.5,
+      cex = 1.5)
```

We finish this section with a comparison of the estimated pooled ROC curves obtained using all methods incorporated in **ROCnReg** (Figure 15).

```
R> plot(pROC_emp$p, pROC_emp$ROC[,1], type = "s", xlim = c(0,1),
+      ylim = c(0,1), xlab = "FPF", ylab = "TPF",
+      main = "Pooled ROC curve - Different approaches",
+      cex.main = 1.5, cex.lab = 1.5, cex.axis = 1.5)
+ lines(pROC_dpm$p, pROC_dpm$ROC[,1], col = 2)
+ lines(pROC_BB$p, pROC_BB$ROC[,1], col = 3)
+ lines(pROC_kernel$p, pROC_kernel$ROC[,1], col = 4)
+ abline(0, 1, col = "grey", lty = 2)
+ legend("topleft", legend = c("Empirical", "DPM", "BB", "Kernel"),
+      lty = 1, col = 1:4, bty = "n", lwd = 2, cex = 1.5)
```

B. Further computational tools for the covariate-specific ROC curve

B.1. Convergence assessments for the Bayesian methods: trace plots, effective sample sizes, and Geweke statistics

We start this section by including, for model `cROC_bnp` in Section 4.2, some trace plots of the MCMC draws (after burn-in) of the conditional PDFs of BMI (Figure 16) and corresponding




Figure 14: Graphical results as provided by the `plot.pooledROC` function for an object of class `pooledROC.kernel`. Estimate and 95% pointwise bootstrap confidence interval of the pooled ROC curve and corresponding AUC.




Figure 15: ROC curve estimated using the different approaches implemented in **ROCnReg**. ‘Empirical’ stands for the empirical estimator, ‘DPM’ for the Dirichlet process mixture of normal distributions estimator, ‘BB’ for the Bayesian bootstrap approach, and ‘Kernel’ for the kernel estimator.

effective sample sizes and Geweke statistics (Figure 17 and 18, respectively). All plots show good mixing of the MCMC chains and do not suggest lack of convergence. For conciseness, the R-code for producing Figures 16, 17, and 18 is not provided here, but in the R replication code that accompanies this paper.




Figure 16: Trace plots of the MCMC draws (after burn-in) of the conditional PDFs of BMI based on model `cROC_bnp`. Results are shown separately for the nondiseased and diseased population, for different combinations of age and gender (covariates) and for different values of the BMI.




Figure 17: Effective sample size of the MCMC chains (after burn-in) of the conditional PDFs of BMI based on model `pROC_dpm`. Results are shown separately for the nondiseased and diseased population and for different combinations of `age` and `gender`. In all cases, results are shown along BMI values.

B.2. Informal model diagnostics for the Bayesian methods: quantile residuals and conditional densities

As in the no covariates case, we also present the quantile residuals for both the Bayesian normal linear model (designated as model `cROC_bp` in Section “Package presentation and illustration” of the main manuscript) and for the Bayesian nonparametric model with 10 mixture components and a factor by curve interaction (model `cROC_bnp` in the main text). The results are shown in Figure 19 and, as for the unconditional case, they show virtually no deviation from the quantiles of the standard normal distribution, in both the diseased and nondiseased groups, for the Bayesian nonparametric method. Further, in Figures 20 and 21 we also show the histograms of the BMI for selected age intervals and for both men and women, along with the estimated (by model `cROC_bnp`) conditional distribution of BMI given age and gender, in the nondiseased and diseased groups. As it can be appreciated, the estimated conditional densities follow quite nicely histograms of the BMI for the different age intervals and genders. For conciseness, the R-code for producing these figures is not provided here, but in the R replication code that accompanies the paper.





Figure 18: Geweke statistic of the MCMC chains (after burn-in) of the conditional PDFs of BMI based on model `cROC_bnp`. Results are shown separately for the nondiseased and diseased population and for different combinations of `age` and `gender`. In all cases, results are shown along BMI values.


B.3. Frequentist estimators of the covariate-specific ROC curve

We now turn our attention on how to estimate the covariate-specific ROC curve using the induced (semiparametric) linear model (function `cROC.sp`). As for the Bayesian linear model described in the main manuscript, for both nondiseased and diseased groups, the model for the regression functions includes, in addition to the linear effect of `age` and `gender`, the (linear) interaction between the two (i.e., `gender*age` \equiv `gender + age + gender:age`). Also, by specifying `est.cdf = "normal"`, we assume that the error term in both groups follows a standard normal distribution. Finally, uncertainty estimation for this method is based on the bootstrap, and through argument `B = 500`, we indicate the number of resamples. As usual, numeric and graphical summaries are obtained using, respectively, functions `summary` and `plot` (see Figure 22).

```
R> set.seed(123, "L'Ecuyer-CMRG") # for reproducibility
R> cROC_sp <- cROC.sp(formula.h = bmi ~ gender*age,
+   formula.d = bmi ~ gender*age, group = "cvd_idf", tag.h = 0,
+   data = endosyn, newdata = endopred, est.cdf = "normal",
+   p = seq(0, 1, l = 101), B = 500, ci.level = 0.95,
+   parallel = "snow", ncpus = 2)
```



(a) BNP model with 10 mixture components and a factor by curve interaction in each group



(b) Bayesian normal linear model in each group

Figure 19: Quantile residuals of the BMI data versus the theoretical quantiles of the standard normal distribution. The circles represent the posterior mean quantiles over all posterior samples, while the dashed lines represent the corresponding 95% credible bands. Top row: BNP model with 10 mixture components and a factor by curve interaction in each group (model `cROC_bnp` in the main manuscript). Bottom row: Bayesian normal linear model in each group (model `cROC_bp` in the main manuscript).

```
R> summary(cROC_sp)
```

```
Call: [...]
```




Figure 20: Histograms of the BMI for selected age intervals and for men and women, along with the pointwise mean (continuous blue lines) and 95% credible bands (shaded blue areas) for the conditional distribution of BMI given age and gender, in the nondiseased group, under the BNP model with 10 mixture components and a factor by curve interaction (model `cROC_bnp` in the main manuscript). The ages of 30, 40, and 50 correspond, respectively and approximately, to the 25th, 50th, and 75th percentiles of the age variable in the whole dataset.

Approach: Conditional ROC curve - semiparametric

Parametric coefficients

Group H:

	Estimate	Quantile 2.5%	Quantile 97.5%
(Intercept)	22.7670	21.9331	23.5509
genderWomen	-4.2942	-5.2247	-3.3091
age	0.0885	0.0680	0.1091
genderWomen:age	0.0885	0.0645	0.1124

Group D:

	Estimate	Quantile 2.5%	Quantile 97.5%
(Intercept)	26.9171	25.5389	28.2935
genderWomen	4.7363	2.2900	7.1651
age	0.0440	0.0161	0.0719
genderWomen:age	-0.0515	-0.0954	-0.0075

ROC curve:




Figure 21: Histograms of the BMI for selected age intervals and for men and women, along with the pointwise mean (continuous blue lines) and 95% credible bands (shaded blue areas) for the conditional distribution of BMI given age and gender, in the diseased group, under the BNP model with 10 mixture components and a factor by curve interaction (model `cROC_bnp` in the main manuscript). The ages of 30, 40, and 50 correspond, respectively and approximately, to the 25th, 50th, and 75th percentiles of the age variable in the whole dataset.

	Estimate	Quantile 2.5%	Quantile 97.5%
(Intercept)	-0.9482	-1.3376	-0.6040
genderWomen	-2.0633	-2.6498	-1.4612
age	0.0102	0.0026	0.0183
genderWomen:age	0.0320	0.0191	0.0442
b	0.9378	0.8728	1.0044

Model selection criteria:

	Group H	Group D
AIC	12173.673	4007.215
BIC	12202.037	4029.905

Sample sizes:

	Group H	Group D
Number of observations	2149	691
Number of missing data	0	0

```
R> op <- par(mfrow = c(2,2))
R> plot(cROC_sp, ask = FALSE)
R> par(op)
```




Figure 22: Graphical results as provided by the `plot.cROC` function for an object of class `cROC.sp`. Results for a model that includes the linear interaction between `age` and `gender`. Top row: Estimate of the covariate-specific ROC curve along age, separately for men and women. Bottom row: Estimate and 95% pointwise bootstrap confidence interval of the covariate-specific AUC along age, separately for men and women.

When it comes to estimating the covariate-specific ROC curve using the induced kernel-based approach (function `cROC.kernel`), we should keep in mind that it can only deal with one continuous covariate. As a consequence, and for our endocrine study, we evaluate `age` effect separately for men and women, i.e., we fit two different models. The code follows. Note that in contrast with the other functions for estimating the covariate-specific ROC curve, function `cROC.kernel` expects as arguments `marker` and `covariate`, where the user specifies, respectively, the name of the variables that contain the test results (in our example `bmi`) and the covariate (`age`). Uncertainty estimation for this method is also based on the bootstrap, and through argument `B = 500`, we indicate the number of resamples. Numeric and graphical summaries are obtained using, respectively, functions `summary` and `plot`. The

graphical results, for both men and women, are shown in Figure 23.

```
R> # For prediction
R> agep <- seq(22, 80, l = 30)
R> endopred_ker <- data.frame(age = agep)

R> set.seed(123, "L'Ecuyer-CMRG") # for reproducibility
R> cROC_kernel_men <- cROC.kernel(marker = "bmi", covariate = "age",
+   group = "cvd_idf", tag.h = 0, data = subset(endosyn, gender == "Men"),
+   newdata = endopred_ker, p = seq(0, 1, l = 101), B = 500, ci.level = 0.95,
+   parallel = "snow", ncpus = 2)

R> summary(cROC_kernel_men)

Call: [...]

Approach: Conditional ROC curve - Kernel-based
-----

Regression functions:

           Group H      Group D
Bandwidth:  5.767820    6.477821

Kernel Estimator: Local-Constant
Bandwidth Selection Method: Least Squares Cross-Validation
Continuous Kernel Type: Second-Order Gaussian

Variance functions:

           Group H      Group D
Bandwidth:  6.489771    18.567326

Kernel Estimator: Local-Constant
Bandwidth Selection Method: Least Squares Cross-Validation
Continuous Kernel Type: Second-Order Gaussian

Sample sizes:

           Group H      Group D
Number of observations      899      418
Number of missing data       0       0

R> cROC_kernel_women <- cROC.kernel(marker = "bmi", covariate = "age",
+   group = "cvd_idf", tag.h = 0, data = subset(endosyn, gender == "Women"),
+   newdata = endopred_ker, p = seq(0, 1, l = 101), B = 500, ci.level = 0.95,
+   parallel = "snow", ncpus = 2)

R> summary(cROC_kernel_women)
```

Call: [...]

Approach: Conditional ROC curve - Kernel-based

Regression functions:

	Group H	Group D
Bandwidth:	3.993242	4.308757

Kernel Estimator: Local-Constant

Bandwidth Selection Method: Least Squares Cross-Validation

Continuous Kernel Type: Second-Order Gaussian

Variance functions:

	Group H	Group D
Bandwidth:	18.250813	10.490069

Kernel Estimator: Local-Constant

Bandwidth Selection Method: Least Squares Cross-Validation

Continuous Kernel Type: Second-Order Gaussian

Sample sizes:

	Group H	Group D
Number of observations	1250	273
Number of missing data	0	0

```
R> op <- par(mfcol = c(2,2))
R> plot(cROC_kernel_women, ask = FALSE)
R> plot(cROC_kernel_men, ask = FALSE)
R> par(op)
```

C. Frequentist estimators of the AROC curve

We finish this document by presenting the code for estimating the covariate-adjusted ROC curve (AROC curve) using the induced semiparametric linear model (function `AROC.sp`) and the kernel-based approach (function `AROC.kernel`). We avoid giving many details, and simply present the code for fitting the models and obtaining the numerical and graphical summaries. It is important to note that, since the kernel-based approach only deals with one continuous covariate, the AROC curve in this case is estimated separately in men and women. This is to be differentiated from the AROC curve obtained by including both `age` and `gender`, which reflects the discriminatory capacity solely due to the `bmi` while teasing out both `age` and `gender` effects.

```
R> # Induced semiparametric linear model
```




Figure 23: Graphical results as provided by the `plot.cROC` function for an object of class `cROC.kernel`. Top row: Estimate of the covariate-specific ROC curve along age, separately for men (left) and women (right). Bottom row: Estimate and 95% pointwise bootstrap confidence interval of the covariate-specific AUC along age, separately for men (left) and women (right). Results in this case were obtained separately for men and women.

```
R> set.seed(123, "L'Ecuyer-CMRG") # for reproducibility
R> AROC_sp <- AROC.sp(formula.h = bmi ~ gender*age, group = "cvd_idf",
+   tag.h = 0, data = endosyn, est.cdf = "normal", p = seq(0, 1, l = 101),
+   B = 500, ci.level = 0.95, parallel = "snow", ncpus = 2)
```

```
R> summary(AROC_sp)
```

Call: [...]

Approach: AROC semiparametric

 Area under the covariate-adjusted ROC curve: 0.638 (0.612, 0.664)*
 * Confidence level: 0.95

Parametric coefficients (Group H):

	Estimate	Quantile 2.5%	Quantile 97.5%
(Intercept)	22.7670	22.0029	23.6441
genderWomen	-4.2942	-5.4177	-3.3017
age	0.0885	0.0658	0.1080
genderWomen:age	0.0885	0.0637	0.1147

Model selection criteria:

	Group H
AIC	12173.673
BIC	12202.037

Sample sizes:

	Group H	Group D
Number of observations	2149	691
Number of missing data	0	0

R> plot(AROC_sp, cex.main = 1.5, cex.lab = 1.5, cex.axis = 1.5, cex = 1.3)

R> # Kernel-based approach

R> set.seed(123, "L'Ecuyer-CMRG") # for reproducibility

R> AROC_kernel_men <- AROC.kernel(marker = "bmi", covariate = "age",
 + group = "cvd_idf", tag.h = 0, data = subset(endosyn, gender == "Men"),
 + p = seq(0, 1, l = 101), B = 500, ci.level = 0.95, parallel = "snow", ncpus = 2)

R> summary(AROC_kernel_men)

Call: [...]

Approach: AROC Kernel-based

 Area under the covariate-adjusted ROC curve: 0.668 (0.636, 0.708)*
 * Confidence level: 0.95

Regression function:

	Group H
Bandwidth:	5.767820




Figure 24: Graphical results as provided by the `plot.AROC` function for an object of class `AROC.sp`. Estimate and 95% pointwise bootstrap confidence interval of the age/gender adjusted ROC curve (AROC) and corresponding AUC.

Kernel Estimator: Local-Constant
 Bandwidth Selection Method: Least Squares Cross-Validation
 Continuous Kernel Type: Second-Order Gaussian

Variance function:

Group H
 Bandwidth: 6.489771

Kernel Estimator: Local-Constant
 Bandwidth Selection Method: Least Squares Cross-Validation
 Continuous Kernel Type: Second-Order Gaussian

Sample sizes:

	Group H	Group D
Number of observations	899	418
Number of missing data	0	0

```
R> set.seed(123, "L'Ecuyer-CMRG") # for reproducibility
R> AROC_kernel_women <- AROC.kernel(marker = "bmi", covariate = "age",
+   group = "cvd_idf", tag.h = 0, data = subset(endosyn, gender == "Women"),
+   p = seq(0, 1, l = 101), B = 500, ci.level = 0.95, parallel = "snow", ncpus = 2)

R> summary(AROC_kernel_women)
```

Call: [...]

Approach: AROC Kernel-based

 Area under the covariate-adjusted ROC curve: 0.673 (0.636, 0.716)*
 * Confidence level: 0.95

Regression function:

 Group H
 Bandwidth: 3.993242

Kernel Estimator: Local-Constant
 Bandwidth Selection Method: Least Squares Cross-Validation
 Continuous Kernel Type: Second-Order Gaussian

Variance function:

 Group H
 Bandwidth: 18.250813

Kernel Estimator: Local-Constant
 Bandwidth Selection Method: Least Squares Cross-Validation
 Continuous Kernel Type: Second-Order Gaussian

Sample sizes:

	Group H	Group D
Number of observations	1250	273
Number of missing data	0	0

```
R> op <- par(mfcol = c(1,2))
R> plot(AROC_kernel_women, main = "AROC kernel-based \n Women",
+      cex.main = 1.5, cex.lab = 1.5, cex.axis = 1.5, cex = 1.7)
R> plot(AROC_kernel_men, main = "AROC kernel-based \n Men",
+      cex.main = 1.5, cex.lab = 1.5, cex.axis = 1.5, cex = 1.7)
R> par(op)
```




Figure 25: Graphical results as provided by the `plot.AROC` function for an object of class `AROC.kernel`. Estimate and 95% pointwise bootstrap confidence interval of the age adjusted ROC curve (AROC) and corresponding AUC. Analyses were done separately for women (left plot) and men (right plot).

Affiliation:

María Xosé Rodríguez-Álvarez
 BCAM - Basque Center for Applied Mathematics
and
 IKERBASQUE, Basque Foundation for Science
 Alameda de Mazarredo, 14
 E-48009 Bilbao, Basque Country, Spain
 E-mail: mxrodriguez@bcamath.org

Vanda Inácio
 School of Mathematics
 University of Edinburgh
 James Clerk Maxwell Building, The King's Buildings, Peter Guthrie Tait Road
 EH9 3FD Edinburgh, Scotland, United Kingdom
 E-mail: vanda.inacio@ed.ac.uk