

Flat phase of polymerized membranes at two-loop order

O. Coquand,^{1,2,*} D. Mouhanna,^{1,†} and S. Teber^{3,‡}

¹*Sorbonne Université, CNRS, Laboratoire de Physique Théorique de la Matière Condensée, 75005 Paris, France*

²*Institut für Materialphysik im Weltraum, Deutsches Zentrum für Luft- und Raumfahrt, Linder Höhe, 51147 Köln, Germany*

³*Sorbonne Université, CNRS, Laboratoire de Physique Théorique et Hautes Energies, LPTHE, 75005 Paris, France*

We investigate two complementary field-theoretical models describing the flat phase of polymerized – phantom – membranes by means of a two-loop, weak-coupling, perturbative approach performed near the upper critical dimension $D_{uc} = 4$, extending the one-loop computation of Aronovitz and Lubensky [Phys. Rev. Lett. **60**, 2634 (1988)]. We derive the renormalization group equations within the modified minimal subtraction scheme, then analyze the corrections coming from two-loop with a particular attention paid to the anomalous dimension and the asymptotic infrared properties of the renormalization group flow. We finally compare our results to those provided by nonperturbative techniques used to investigate these two models.

I. INTRODUCTION

Fluctuating surfaces are ubiquitous in physics (see, e.g., Refs.[1, 2]). One meets them within the context of high-energy physics [3–6], initially through high-temperature expansions of lattice gauge theories, then in the large- N limit of gauge theories, in two-dimensional quantum gravity, in string theory as world-sheet of string and, finally, in brane theory. They also occur as a fundamental object of biophysics where surfaces – called in this context *membranes* – constitute the building blocks of living cells such as erythrocyte [2, 7]. Last but not least, fluctuating surfaces – or membranes – have provided, in condensed matter physics, an extremely suitable model to describe both qualitatively and quantitatively sheets of graphene [8, 9] or graphenelike materials (see, e.g., [10] and references therein).

Two types of membranes should be distinguished regarding their critical or, more generally, long-distance properties: fluid membranes and polymerized membranes [11, 12]. The specificity of fluid membranes is that the molecules are essentially free to diffuse inside the structure. The consequence of this lack of fixed connectivity is the absence of elastic properties. As a result, the free energy of the membrane depends only on its shape – its curvature – and not on a specific coordinate system. Early studies [13–15] have shown that strong height – out-of-plane – fluctuations occur in such systems in such a way that the normal-normal correlation functions exponentially decay with the distance over a typical persistence length $\xi \sim e^{4\pi\kappa/T}$ in a way similar to what happens in the two-dimensional $O(N)$ model. As a consequence, there is no long-range orientational order in fluid membranes – that are always *crumpled* – in agreement with the Mermin-Wagner theorem [16].

Polymerized – or tethered – membranes are more remarkable. Indeed due to the fact that molecules are

tied together through a potential, they display a fixed internal connectivity giving rise to elastic – shearing and stretching – contributions to the free energy. It has been substantiated that, in these conditions, the coupling between the out-of-plane and in-plane fluctuations leads to a drastic reduction of the former [17]. This makes possible the existence of a phase transition between a disordered *crumpled* phase at high temperatures and an ordered *flat* phase with long-range order between the normals at low-temperatures [1, 18–22], analogous to that occurring in ferro/antiferro magnets – see, however, below. Although the nature – first or second order – of this crumpled-to-flat transition is still under debate [23–30] and the mere existence of a crumpled phase for realistic, i.e., self-avoiding [31], membranes seems to be compromised, there is no doubt about the existence of a stable flat phase.

Let us consider a D -dimensional membrane embedded in the d -dimensional Euclidean space. The location of a point on the membrane is realized by means of a D -dimensional vector \mathbf{x} whereas a configuration of the membrane in the Euclidean space is described through the embedding $\mathbf{x} \rightarrow \mathbf{R}(\mathbf{x})$ with $\mathbf{R} \in \mathbb{R}^d$. One assumes the existence of a low temperature, flat phase, defined by $\mathbf{R}^0(\mathbf{x}) = (\mathbf{x}, \mathbf{0}_{d_c})$ where $\mathbf{0}_{d_c}$ is the null vector of co-dimension $d_c = d - D$ and one decomposes the field \mathbf{R} into $\mathbf{R}(\mathbf{x}) = [\mathbf{x} + \mathbf{u}(\mathbf{x}), \mathbf{h}(\mathbf{x})]$ where \mathbf{u} and \mathbf{h} represent D longitudinal – phonon – and $d - D$ transverse – flexural – modes, respectively. The action of a flat phase configuration \mathbf{R} is given by [17–20, 22, 32]

$$S[\mathbf{R}] = \int d^D x \left\{ \frac{\kappa}{2} (\Delta \mathbf{R})^2 + \frac{\lambda}{2} u_{ii}^2 + \mu u_{ij}^2 \right\} \quad (1)$$

where u_{ij} is the strain tensor that parametrizes the fluctuations around the flat phase configuration $\mathbf{R}^0(\mathbf{x})$: $u_{ij} = \frac{1}{2}(\partial_i \mathbf{R} \cdot \partial_j \mathbf{R} - \partial_i \mathbf{R}^0 \cdot \partial_j \mathbf{R}^0) = \frac{1}{2}(\partial_i \mathbf{R} \cdot \partial_j \mathbf{R} - \delta_{ij})$. In Eq. (1), κ is the bending rigidity constant whereas λ and μ are the Lamé coefficients; stability considerations require that κ , μ , and the bulk modulus $B = \lambda + 2\mu/D$ be all positive.

The most remarkable fact arising from the analysis of (1) is that, in the flat phase, the normal-normal corre-

*Electronic address: coquand@lptmc.jussieu.fr

†Electronic address: mouhanna@lptmc.jussieu.fr

‡Electronic address: teber@lpthe.jussieu.fr

lation functions display long-range order from the upper critical (uc) dimension $D_{uc} = 4$ down to the lower critical (lc) dimension $D_{lc} < 2$ [18, 22]. While in apparent contradiction with the Mermin-Wagner theorem [16], this result can be explained in the following way. At long distances, or low momenta, typically given by $q \ll \sqrt{\mu/\kappa}, \sqrt{\lambda/\kappa}$, the term $(\Delta \mathbf{u})^2$ in (1) can be neglected with respect to the terms of the type $(\partial_i u_j)^2$ entering in the strain tensor u_{ij} , which are, thus, promoted to the rank of kinetic terms of the field u [33]. It follows immediately from power counting considerations that the nonlinear term in the phonon field \mathbf{u} appearing in u_{ij} , i.e. $\partial_i u_k \partial_j u_k$, is irrelevant; it can thus also be discarded. Under these assumptions the strain tensor u_{ij} is given by:

$$u_{ij} \simeq \frac{1}{2} [\partial_i u_j + \partial_j u_i + \partial_i \mathbf{h} \cdot \partial_j \mathbf{h}] . \quad (2)$$

It follows that action (1) is now quadratic in the phonon field \mathbf{u} and one can integrate over it exactly. This leads to an *effective* action depending only on the flexural field \mathbf{h} . In Fourier space, this effective action reads [34, 35]

$$S_{\text{eff}}[\mathbf{h}] = \frac{\kappa}{2} \int_{\mathbf{k}} k^4 |\mathbf{h}(\mathbf{k})|^2 + \frac{1}{4} \int_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} \mathbf{h}(\mathbf{k}_1) \cdot \mathbf{h}(\mathbf{k}_2) R_{ab,cd}(\mathbf{q}) k_1^a k_2^b k_3^c k_4^d \mathbf{h}(\mathbf{k}_3) \cdot \mathbf{h}(\mathbf{k}_4) , \quad (3)$$

where $\int_{\mathbf{k}} = \int d^D k / (2\pi)^D$ and $\mathbf{q} = \mathbf{k}_1 + \mathbf{k}_2 = -\mathbf{k}_3 - \mathbf{k}_4$. The fourth order, \mathbf{q} -transverse tensor, $R_{ab,cd}(\mathbf{q})$ is given by [34, 35]

$$R_{ab,cd}(\mathbf{q}) = \frac{\mu(D\lambda + 2\mu)}{\lambda + 2\mu} N_{ab,cd}(\mathbf{q}) + \mu M_{ab,cd}(\mathbf{q}) \quad (4)$$

where one has defined the two mutually orthogonal tensors:

$$N_{ab,cd}(\mathbf{q}) = \frac{1}{D-1} P_{ab}^T(\mathbf{q}) P_{cd}^T(\mathbf{q})$$

$$M_{ab,cd}(\mathbf{q}) = \frac{1}{2} [P_{ac}^T(\mathbf{q}) P_{bd}^T(\mathbf{q}) + P_{ad}^T(\mathbf{q}) P_{bc}^T(\mathbf{q})] - N_{ab,cd}(\mathbf{q})$$

where $P_{ab}^T(\mathbf{q}) = \delta_{ab} - q_a q_b / \mathbf{q}^2$ is the transverse projector. Note that, in $D = 2$, the tensor $M_{ab,cd}$ vanishes identically and the effective action (3) is parametrized by only one coupling constant which turns out to be proportional to Young's modulus [17, 34, 35]: $K_0 = 4\mu(\lambda + \mu)/(\lambda + 2\mu)$. The key point is that the momentum-dependent interaction (4) is nonlocal and gives rise to a phonon-mediated interaction between flexural modes which is of the long-range kind. More precisely, this interaction contains terms such that the product $R(|\mathbf{x} - \mathbf{y}|)|\mathbf{x} - \mathbf{y}|^2$ is not an integrable function in $D = 2$ as required by the Mermin-Wagner theorem [16] (see [36] for a detailed discussion). This flat phase is characterized by power-law behaviour

for the phonon-phonon and flexural-flexural modes correlation functions [18, 20, 22, 32]:

$$G_{uu}(q) \sim q^{-(2+\eta_u)} \quad \text{and} \quad G_{hh}(q) \sim q^{-(4-\eta)} \quad (5)$$

where η and η_u are nontrivial anomalous dimensions. In fact, it follows from Ward identities associated with the remaining partial rotation invariance of (1) – see below – that η and η_u are not independent quantities and one has $\eta_u = 4 - D - 2\eta$ [18, 20, 22, 32]. Interestingly, Eq. (5) provide also an implicit equation for the lower critical dimension D_{lc} defined as the dimension below which there is no more distinction between phonon and flexural modes. One gets from Eq. (5): $D_{lc} - 2 + \eta(D_{lc}) = 0$ [14, 18, 22]. It results from this expression that the lower critical dimension D_{lc} , as well as the associated anomalous dimension $\eta(D_{lc})$, are no longer given by a power-counting analysis around a Gaussian fixed point, as it occurs for the $O(N)$ model, but by a nontrivial computation of fluctuations. This implies, in particular, that there is no well-defined perturbative expansion of the flat phase theory near the lower critical dimension D_{lc} based on the study of a nonlinear σ – hard-constraints – model [19].

On the other hand, the soft-mode, Landau-Ginzburg-Wilson, model (1) does not suffer from the same kind of pathology; a standard, ϵ -expansion about the upper critical dimension D_{uc} is feasible and has been performed at leading order, a long-time ago, in the seminal works of Aronovitz et al. [18, 22] and Gutter et al. [20, 32] who have determined the renormalization group (RG) equations and the properties of the flat phase near $D = 4$. This perturbative approach faces, however, several drawbacks that explain why it has not been pushed forward until now: (i) It involves an intricate momentum and tensorial structure of the propagators and vertices that render the diagrammatic extremely rapidly growing in complexity with the order of perturbation [37]. (ii) The dimension of physical membranes, $D = 2$, is “far away” from D_{uc} . Clearly, high orders of the perturbative series, followed by suitable resummation techniques, are needed to get quantitatively trustable results. The difficulty of carrying such a task is, however, increased by the first drawback. (iii) The massless theory is manageable with current modern techniques whereas, with the $1/q^4$ -form of the flexural mode propagator G_{hh} in (5), one apparently faces the problem of dealing with infrared divergences. (iv) The use of the dimensional regularization and, more precisely, the modified minimal subtraction ($\overline{\text{MS}}$) scheme, which is by far the most convenient one, can enter in conflict with the D -dependence of physical quantities or properties – see below.

In this context, several nonperturbative methods – with respect to the parameter $\epsilon = 4 - D$ – have been employed in order to tackle the physics directly in $D = 2$. Among them, the $1/d_c$ expansion have been early performed at leading order [19, 20, 22, 32, 33] and, very recently, at next-to-leading order [38]. An improvement of the $1/d_c$ approximation that consists in replacing, within this last approach, the bare propagator and vertices by

their dressed and screened counterparts leads to the so-called self-consistent screening approximation (SCSA) that has also been used at leading [34, 35, 39, 40] and next-to-leading order [41]. Finally, a technique working in all dimensions D and d , called nonperturbative renormalization group (NPRG) – see below – has been employed to investigate various kinds of membranes at leading order of the so-called derivative expansion [26, 28, 42–45] and within an approach taking into account the full derivative dependence of the action [46, 47]. Therefore, within the whole spectrum of approaches used to investigate the properties of the flat phase of membranes, it is only for the weak-coupling perturbative approach that the next-to-leading order is still missing (see however [48]). This is clearly a flaw as the subleading corrections of any approach generally provide valuable insights on the structure of the whole theory. They also convey useful information about the accuracy of complementary approaches.

We propose here to fill this gap and to investigate the properties of the flat phase of polymerized membranes at two-loop order in the coupling constants, near $D_{uc} = 4$, considering successively the flexural-phonon, *two-field*, model (1) and then the flexural-flexural, *effective* model (3). We compute the RG functions of these two models, analyze their fixed points and compute the corresponding anomalous dimensions. Finally, we compare these results together and, then, with those obtained from nonperturbative methods. Note that, due to the length of the computations and expressions involved, we restrict here ourselves to the main results; details will be given in a forthcoming publication [37].

II. THE TWO-FIELD MODEL

A. The perturbative approach

We first consider the two-field model (1) truncated by means of the long distance approximations Eq. (2) and $(\Delta \mathbf{u})^2 \simeq 0$. The perturbative approach proceeds as usual: one expresses the action in terms of the phonon and flexural fields \mathbf{u} and \mathbf{h} then get the propagators and 3 and 4-point vertices, see [20, 37]. A crucial issue is that, although the truncations of action (1) above break its original $O(d)$ symmetry, a partial rotation invariance remains [20, 32]:

$$\begin{aligned} \mathbf{h} &\mapsto \mathbf{h} + \mathbf{A}_i x_i \\ u_i &\mapsto u_i - \mathbf{A}_i \cdot \mathbf{h} - \frac{1}{2} \mathbf{A}_i \cdot \mathbf{A}_j x_j \end{aligned}$$

where \mathbf{A}_i is any set of D vectors $\in \mathbb{R}^{d_c}$. From this property follow Ward identities for the effective action Γ [20, 32]:

$$\int d^D x \left(\mathbf{h} \frac{\delta \Gamma}{\delta u_i} - x_i \frac{\delta \Gamma}{\delta \mathbf{h}} \right) = 0. \quad (6)$$

One easily shows that this equation is solved by – the truncated form of – (1), thereby ensuring the renormalizability of the theory. Moreover, from (6), one can derive successive identities relating various n -points to $(n-1)$ -point functions in such a way that only the renormalizations of phonon and flexural modes propagators are required. This is a tremendous simplification of the computation which, nevertheless, preserves a nontrivial algebra. Also, as previously mentioned, an apparent difficulty comes from the structure of the – bare – flexural mode propagator $G_{hh}(q) \sim 1/q^4$ and the masslessness of the theory that suggests that the perturbative expansion could be plagued by severe infrared divergencies. In this respect, one has first to note that the masslessness of the theory and the form of the propagators (5) are somewhat contrived as they originate from the derivative character of (1) relying itself from the lack of translational invariance of the embedding $\mathbf{x} \rightarrow \mathbf{R}(\mathbf{x})$. It appears that the *natural* objects that should be ideally considered are the tangent-tangent correlation functions $\tilde{G} \sim \langle \partial_i \mathbf{R} \cdot \partial_i \mathbf{R} \rangle$ whose Fourier transforms are, for fixed-connectivity membranes, proportional to the position-position ones $G \sim \langle \mathbf{R} \cdot \mathbf{R} \rangle$ with a factor of \mathbf{q}^2 [22] and are, consequently, infrared safe. In practice, however, employing the latter correlation functions is both preferable and innocuous as its use only implies the appearance of tadpoles that cancel order by order in perturbation theory. One can, thus, proceed using dimensional regularization in the conventional way ignoring the occurrence of possible infrared poles [49].

B. The renormalization group equations

One introduces the renormalized fields \mathbf{h}_R and \mathbf{u}_R through $\mathbf{h} = Z^{1/2} \kappa^{-1/2} \mathbf{h}_R$ and $\mathbf{u} = Z \kappa^{-1} \mathbf{u}_R$ and the renormalized coupling constants λ_R and μ_R through :

$$\begin{aligned} \lambda &= k^\epsilon Z^{-2} \kappa^2 Z_\lambda \lambda_R \\ \mu &= k^\epsilon Z^{-2} \kappa^2 Z_\mu \mu_R \end{aligned} \quad (7)$$

where k is the renormalization momentum scale and $\epsilon = 4 - D$. Within the $\overline{\text{MS}}$ scheme, one introduces the scale $\bar{k}^2 = 4\pi e^{-\gamma_E} k^2$ where γ_E is the Euler constant. One then defines the β -functions $\beta_{\lambda_R} = \partial_t \lambda_R$ and $\beta_{\mu_R} = \partial_t \mu_R$, with $t = \ln \bar{k}$. As usual, in order to write these quantities in terms of the field and coupling constant renormalizations $Z = Z(\lambda_R, \mu_R, \epsilon)$, $Z_\lambda = Z_\lambda(\lambda_R, \mu_R, \epsilon)$ and $Z_\mu = Z_\mu(\lambda_R, \mu_R, \epsilon)$ one expresses the independence of the bare coupling constants λ and μ with respect to t : $d\lambda/dt = d\mu/dt = 0$. Using (7) and defining the anomalous dimension

$$\eta = \beta_{\lambda_R} \frac{\partial \ln Z}{\partial \lambda_R} + \beta_{\mu_R} \frac{\partial \ln Z}{\partial \mu_R}$$

one gets from these conditions:

$$\beta_{\lambda_R} \partial_{\lambda_R} \ln(\lambda_R Z_\lambda) + \beta_{\mu_R} \partial_{\mu_R} \ln(\lambda_R Z_\lambda) = -\epsilon + 2\eta$$

$$\beta_{\lambda_R} \partial_{\lambda_R} \ln(\mu_R Z_\mu) + \beta_{\mu_R} \partial_{\mu_R} \ln(\mu_R Z_\mu) = -\epsilon + 2\eta$$

where only simple poles in ϵ of Z , Z_λ and Z_μ have to be considered, see [37].

Computations have been performed independently by means of (i) conventional renormalization – counterterms – method (ii) BPHZ [50–52] renormalization scheme with the help of the LITERED *mathematica* package for the reduction two-loop integrals [53]. Both computations have required techniques for computing massless Feynman diagram calculations that are reviewed in, e.g., Ref.[54].

Omitting the R indices on the renormalized coupling constants one gets, after involved computations [37]

$$\begin{aligned} \beta_\mu &= -\epsilon\mu + 2\mu\eta + \frac{d_c\mu^2}{6(16\pi^2)} \left(1 + \frac{227}{180}\eta^{(0)}\right) \\ \beta_\lambda &= -\epsilon\lambda + 2\lambda\eta + \frac{d_c(6\lambda^2 + 6\lambda\mu + \mu^2)}{6(16\pi^2)} \\ &\quad - \frac{d_c(378\lambda^2 - 162\lambda\mu - 17\mu^2)}{1080(16\pi^2)}\eta^{(0)} - \frac{d_c^2\mu(3\lambda + \mu)^2}{36(16\pi^2)^2} \end{aligned} \quad (8)$$

where

$$\begin{aligned} \eta &= \eta^{(0)} + \eta^{(1)} = \frac{5\mu(\lambda + \mu)}{16\pi^2(\lambda + 2\mu)} \\ &\quad - \frac{\mu^2((340 + 39d_c)\lambda^2 + 4(35 + 39d_c)\lambda\mu + (81d_c - 20)\mu^2)}{72(16\pi^2)^2(\lambda + 2\mu)^2} \end{aligned} \quad (9)$$

C. Fixed points analysis

Equations (8) and (9) constitute the first set of our main results. These equations extend to two-loop order those of Aronovitz and Lubensky [18]. One first recalls the properties of the one-loop RG flow [18, 20, 22, 32], then, considers the full two-loop equations (8) and (9).

1. One-loop order

At one-loop order there are four fixed points, see Fig. 1:

(i) the Gaussian one P_1 for which $\mu_1^* = 0$, $\lambda_1^* = 0$ and $\eta_1 = 0$; it is twice unstable.




FIG. 1: The schematic RG flow diagram (not to scale) on the plane (μ, λ) . The stability region of action (1) is delimited by the line $2\lambda + \mu = 0$ on which lies the fixed point P_3 at one-loop order and the line $\mu = 0$ on which lies the fixed point P_2 . The dashed line corresponds to the one-loop attractive subspace $3\lambda + \mu = 0$ where the stable fixed point P_4 stands. At two-loop order, P_4 does not stand exactly on the line $3\lambda + \mu = 0$ anymore whereas P_3 is ejected out the stability region.

(ii) The – shearless – fixed point P_2 with $\mu_2^* = 0$, $\lambda_2^* = 16\pi^2\epsilon/d_c$ and $\eta_2 = 0$ which lies on the stability line $\mu = 0$; it is once unstable.

(iii) The infinitely compressible fixed point P_3 with $\mu_3^* = 96\pi^2\epsilon/(20 + d_c)$, $\lambda_3^* = -48\pi^2\epsilon/(20 + d_c)$ and $\eta_3 = 10\epsilon/(20 + d_c)$, for which the bulk modulus B vanishes, i.e. $2\lambda_3^* + \mu_3^* = 0$. It is thus located on the corresponding stability line; it is once unstable.

(iv) The flat phase fixed point P_4 for which $\mu_4^* = 96\pi^2\epsilon/(24 + d_c)$, $\lambda_4^* = -32\pi^2\epsilon/(24 + d_c)$ and $\eta_4 = 12\epsilon/(24 + d_c)$. It is fully stable and, thus, controls the flat phase at long distance. At one-loop order, this fixed point is located on the stable line $3\lambda + \mu = 0$ – that, in D dimensions, generalizes to the line $(D + 2)\lambda + 2\mu = 0$.

2. Two-loop order

At two-loop order there are still four fixed points. For the two first ones, nothing changes whereas, for the two last ones, the situation changes only marginally:

(i) the Gaussian fixed point P_1 remains twice unstable.

(ii) The once unstable fixed point P_2 keeps the same coordinates as at one-loop order – with in particular $\mu_2^* = 0$ – thus the associated anomalous dimension, which is proportional to μ , see (9), still vanishes: $\eta_2 = O(\epsilon^3)$.

(iii) At the other once unstable fixed point P_3 , whose

coordinates and associated exponent are given in Table I, the bulk modulus B becomes now slightly negative – and of order ϵ^2 – see Fig. 1. It follows that, at this order, P_3 is ejected out of the stability region. However, we emphasize that this fact fully depends on the technique or – two-field or effective – formulation of the theory – see below. It is, thus, likely that this is an artifact of the present computation. So one can still consider P_3 as potentially present in the genuine flow diagram of membranes.

μ_3^*	$\frac{96\pi^2 \epsilon}{20 + d_c} + \frac{80\pi^2(-d_c + 232)}{3(20 + d_c)^3} \epsilon^2$
λ_3^*	$-\frac{48\pi^2 \epsilon}{20 + d_c} - \frac{8\pi^2(9d_c^2 + 265d_c + 2960)}{3(20 + d_c)^3} \epsilon^2$
η_3	$\frac{10 \epsilon}{20 + d_c} - \frac{d_c(37d_c + 950)}{6(20 + d_c)^3} \epsilon^2$

TABLE I: Coordinates μ_3^* and λ_3^* of the fixed point P_3 and the corresponding anomalous dimension η_3 at order ϵ^2 obtained from the two-field model.

(iv) P_4 remains fully stable and, thus, still controls the flat phase. Its coordinates and associated anomalous dimension are given in Table II. As a noticeable point one indicates that this fixed point no longer lies on the line $(D + 2)\lambda + 2\mu = (6 - \epsilon)\lambda + 2\mu = 0$ – with a distance of order ϵ^2 as expected – which is, thus, no longer an attractive line in the infrared.

As can be seen in Table II the anomalous dimension at P_4 is only very slightly modified with respect to its one-loop order value. The extrapolation of our result for η_4 to $D = 2$, i.e. $\epsilon = 2$ and $d_c = 1$ leads, at one and two-loop orders, to $\eta_4^{1l} = 24/25 = 0.96$ and $\eta_4^{2l} = 2856/3125 \simeq 0.914$. These values are obviously only indicative and are in no way supposed to provide a quantitatively accurate prediction in $D = 2$. However, one can note that the two-loop correction moves the value of η_4 towards the right direction if one refers to the generally accepted numerical data that lie in the range of $[0.72, 0.88]$ [55–65].

μ_4^*	$\frac{96\pi^2 \epsilon}{24 + d_c} - \frac{32\pi^2(47d_c + 228)}{5(24 + d_c)^3} \epsilon^2$
λ_4^*	$-\frac{32\pi^2 \epsilon}{24 + d_c} + \frac{32\pi^2(19d_c + 156)}{5(24 + d_c)^3} \epsilon^2$
η_4	$\frac{12 \epsilon}{24 + d_c} - \frac{6d_c(d_c + 29)}{(24 + d_c)^3} \epsilon^2$

TABLE II: Coordinates μ_4^* and λ_4^* of the flat phase fixed point P_4 and the corresponding anomalous dimension η_4 at order ϵ^2 obtained from the two-field model.

III. THE FLEXURAL MODE EFFECTIVE MODEL

A. The perturbative approach

We have also considered an alternative approach to the flat phase theory of membranes which is given by the flexural mode effective model (3). There are three main reasons to tackle directly this model. The first one is formal and consists in showing that one can treat, at two-loop order, a model with a nonlocal interaction. The second reason is that this provides a nontrivial check of the previous computations. Indeed the field-content, the (unique) four-point nonlocal vertex as well as the whole structure of the perturbative expansion of the effective model (3) are considerably different from those of the two-field model so that the agreement between the two approaches is a very substantial fact. The last reason to investigate this model is that it involves a new coupling constant $b = \mu(D\lambda + 2\mu)/(\lambda + 2\mu)$, see (4), which: (i) is directly proportional to the bulk modulus B associated with a stability line of the model and (ii) incorporates a D -dependence which, as b is considered as a coupling constant in itself, will be kept from the influence of the dimensional regularization.

B. The renormalization group equations

As in the two-field model, one introduces the renormalized field \mathbf{h}_R through $\mathbf{h} = Z^{1/2}\kappa^{-1/2}\mathbf{h}_R$, the renormalized coupling constants b_R and μ_R through

$$\begin{aligned} b &= k^\epsilon Z^{-2} \kappa^2 Z_b b_R \\ \mu &= k^\epsilon Z^{-2} \kappa^2 Z_\mu \mu_R \end{aligned} \quad (10)$$

and the β -functions $\beta_{b_R} = \partial_t b_R$ and $\beta_{\mu_R} = \partial_t \mu_R$. Using (10) to express the independence of the bare coupling constants b and μ with respect to t and defining the anomalous dimension

$$\eta = \beta_{b_R} \frac{\partial \ln Z}{\partial b_R} + \beta_{\mu_R} \frac{\partial \ln Z}{\partial \mu_R}$$

the β -functions β_{b_R} and β_{μ_R} read:

$$\beta_{b_R} \partial_{b_R} \ln(b_R Z_b) + \beta_{\mu_R} \partial_{\mu_R} \ln(b_R Z_b) = -\epsilon + 2\eta$$

$$\beta_{b_R} \partial_{b_R} \ln(\mu_R Z_\mu) + \beta_{\mu_R} \partial_{\mu_R} \ln(\mu_R Z_\mu) = -\epsilon + 2\eta.$$

After a rather heavy algebra and using the same techniques as for the two-field model one gets:

$$\begin{aligned} \beta_\mu &= -\epsilon\mu + 2\mu\eta + \frac{d_c \mu^2}{6(16\pi^2)} \left(1 + \frac{107b + 574\mu}{216(16\pi^2)}\right) \\ \beta_b &= -\epsilon b + 2b\eta + \frac{5d_c b^2}{12(16\pi^2)} \left(1 + \frac{178\mu - 91b}{216(16\pi^2)}\right) \end{aligned} \quad (11)$$

and:

$$\eta = \frac{5(b+2\mu)}{6(16\pi^2)} + \frac{5(15d_c - 212)b^2 + 1160b\mu - 4(111d_c - 20)\mu^2}{2592(16\pi^2)^2}. \quad (12)$$

C. Fixed point analysis

Equations (11) and (12) constitute our second set of results. We now analyze their content.

1. One-loop order

At one-loop one finds four fixed points:

(i) the Gaussian one P_1 with $\mu_1^* = 0$, $b_1^* = 0$ and $\eta_1 = 0$, which is twice unstable.

(ii) A fixed point, P'_2 with $\mu_2'^* = 0$, $b_2'^* = 192\pi^2\epsilon/5(d_c + 4)$ and $\eta_2' = 2\epsilon/(d_c + 4)$. This fixed point has no counterpart within the two-field model where b is a function of λ and μ and, in particular, proportional to μ ; it is once unstable.

(iii) The infinitely compressible fixed point P_3 with $\mu_3^* = 96\pi^2\epsilon/(d_c + 20)$, $b_3^* = 0$ and $\eta_3 = 10\epsilon/(20 + d_c)$, for which the bulk modulus B vanishes. It thus identifies with the fixed point P_3 of the two-field model; it is once unstable.

(iv) The fixed point P_4 with $\mu_4^* = 96\pi^2\epsilon/(24 + d_c)$, $b_4^* = 192\pi^2\epsilon/5(d_c + 24)$ and $\eta_4 = 12\epsilon/(24 + d_c)$ which is fully stable and controls the flat phase. It is located on the stable line $5b - 2\mu = 0$ – corresponding to $(D+1)b - 2\mu = 0$ in D dimensions – equivalent to the line $3\lambda + \mu = 0$ in the two-field model. It fully identifies with the fixed point P_4 of that model.

Note finally that, as said above, in $D = 2$, the tensor $M_{ab,cd}$ vanishes, which is equivalent to the condition $\mu = 0$. This implies that the coordinates of the fixed points all obey this condition. As a consequence, in $D = 2$, only one nontrivial fixed point, P'_2 , remains.

2. Two-loop order

At two-loop order, as in the two-field model, the one-loop picture is not radically changed.

(i) The Gaussian fixed point P_1 remains twice unstable.

(ii) At P'_2 , $\mu_2'^*$ still strictly vanishes whereas $b_2'^*$ is only slightly modified, see Table III. This fixed point, as well as its anomalous dimension η_2' has been first obtained at two-loop order by Mauri and Katsnelson [48] in a very recent study of the Gaussian curvature interaction (CGI) model – see below.

$\mu_2'^*$	0
$b_2'^*$	$\frac{192\pi^2\epsilon}{5(4+d_c)} + \frac{32\pi^2(61d_c+424)}{75(4+d_c)^3}\epsilon^2$
η_2'	$\frac{2\epsilon}{4+d_c} + \frac{d_c(d_c-2)}{6(4+d_c)^3}\epsilon^2$

TABLE III: Coordinates $\mu_2'^*$ and $b_2'^*$ and the corresponding anomalous dimension η_2' of the fixed point P'_2 at order ϵ^2 obtained from the effective model; P'_2 has been first obtained in [48].

(iii) The fixed point P_3 is interesting as it has a direct counterpart in the two-field model, which allows to study the modifications induced by the change in model. Its coordinates, see Table IV, differ from those of the two-field model, see Table I, in particular as they still obey the condition $b_3^* = 0$ – or $B = 0$ – that puts P_3 just on the boundary of the stability region of the theory. This fact is an indication that, within the two-loop approach of the two-field model, the location of the fixed point P_3 out of the stability region is very likely an artifact of the model or of its perturbative approach. This could also be a drawback of the dimensional regularization that seems to mismanage D -dependent quantities such as the hypersurface $B = 0$. *Nevertheless* the anomalous dimension η_3 , see Table IV, coincides exactly with the two-field result, see Table I, which is a strong check of our computations.

μ_3^*	$\frac{96\pi^2\epsilon}{20+d_c} - \frac{80\pi^2(13d_c+8)}{3(20+d_c)^3}\epsilon^2$
b_3^*	0
η_3	$\frac{10\epsilon}{20+d_c} - \frac{d_c(37d_c+950)}{6(20+d_c)^3}\epsilon^2$

TABLE IV: Coordinates μ_3^* and b_3^* of the fixed point P_3 and the corresponding anomalous dimension η_3 at order ϵ^2 obtained from the effective model.

(iv) Finally the fixed point P_4 remains stable and controls the flat phase. Its coordinates and associated exponent η_4 are given in Table V. In the same way as for the fixed point P_3 , the coordinates of P_4 at two-loop order differ from those obtained from the two-field model, see Table II. *Also*, these coordinates *do not* obey the condition $(D+1)b_4^* - 2\mu_4^* = (5-\epsilon)b_4^* - 2\mu_4^* = 0$ corresponding to the one-loop stability line. *Nevertheless*, again the anomalous dimension η_4 coincides exactly with the two-field model result, see Table II.

μ_4^*	$\frac{96\pi^2 \epsilon}{24 + d_c} - \frac{32\pi^2(77d_c + 948)}{5(24 + d_c)^3} \epsilon^2$
b_4^*	$\frac{192\pi^2 \epsilon}{5(24 + d_c)} + \frac{64\pi^2(121d_c + 3804)}{25(24 + d_c)^3} \epsilon^2$
η_4	$\frac{12 \epsilon}{24 + d_c} - \frac{6d_c(d_c + 29)}{(24 + d_c)^3} \epsilon^2$

TABLE V: Coordinates μ_4^* and b_4^* of the flat phase fixed point P_4 and the corresponding anomalous dimension η_4 at order ϵ^2 obtained from the effective model.

IV. COMPARISON WITH PREVIOUS APPROACHES

We now discuss our results compared to the other techniques – or other models – that have been used to investigate the flat phase of membranes.

SCSA. The SCSA has been studied early [34] to investigate the properties of membranes in any dimension D . It is generally employed using the effective action (3) which is more suitable than (1) to establish self-consistent equations. By construction, this approach is one-loop exact. It is also exact at first order in $1/d_c$ and, finally, at $d_c = 0$. Even more remarkably, comparing the anomalous dimensions η'_2 , η_3 and η_4 obtained in this context to the two-loop results, see Table VI, one observes that

the first one is *exact* at order ϵ^2 whereas the latter ones are almost exact at this order as only the coefficients in ϵ^2/d_c^2 differ slightly from those of our exact results.

There are two important features of the SCSA approach that should be underlined. First, the solution with a vanishing bare modulus $b = 0$, thus corresponding to the fixed point P_3 , leads to a vanishing long-distance effective modulus $b(\mathbf{q}) = 0$ [35], in agreement with our results $b_3^* = 0$. Second, under the conditions fulfilled to reach the scaling behaviour associated with the fixed point P_4 , one observes the asymptotic infrared behaviour [34, 35]:

$$\frac{\lambda(\mathbf{q})}{\mu(\mathbf{q})} \underset{\mathbf{q} \rightarrow \mathbf{0}}{\sim} -\frac{2}{D+2} \quad (13)$$

in any dimension D – which is equivalent to the condition $(D+2)\lambda + 2\mu = 0$ or, equivalently, $(D+1)b - 2\mu = 0$ discussed above. This property has been proposed to work at all orders of the SCSA and even to be exact [41] which leads us to wonder about the genuine location of the fixed point P_4 found perturbatively at two-loop order that violates condition (13).

We finally recall that, in $D = 2$, one gets, at leading order, $\eta_{SCSA}^{D=2,l} = 0.821$ [34, 35] and, at next-to-leading order, $\eta_{SCSA}^{D=2,nl} = 0.789$ [41] which is inside the range of values given above and close to some of the most recent results obtained by means of numerical computations (see, e.g., [62] that provides $\eta \simeq 0.79$).

	Two-loop expansion	SCSA	NPRG
η'_2	$\frac{2\epsilon}{4 + d_c} + \frac{d_c(d_c - 2)}{6(4 + d_c)^3} \epsilon^2$	$\frac{2\epsilon}{4 + d_c} + \frac{d_c(d_c - 2)}{6(4 + d_c)^3} \epsilon^2$	$\frac{2\epsilon}{4 + d_c} + \frac{d_c(10 + 3d_c)}{12(4 + d_c)^3} \epsilon^2$
η_3	$\frac{10\epsilon}{20 + d_c} - \frac{d_c(37d_c + 950)}{6(20 + d_c)^3} \epsilon^2$	$\frac{10\epsilon}{20 + d_c} - \frac{d_c(37d_c + 890)}{6(20 + d_c)^3} \epsilon^2$	$\frac{10\epsilon}{20 + d_c} - \frac{d_c(69d_c + 1430)}{12(20 + d_c)^3} \epsilon^2$
η_4	$\frac{12\epsilon}{24 + d_c} - \frac{6d_c(d_c + 29)}{(24 + d_c)^3} \epsilon^2$	$\frac{12\epsilon}{24 + d_c} - \frac{6d_c(d_c + 30)}{(24 + d_c)^3} \epsilon^2$	$\frac{12\epsilon}{24 + d_c} - \frac{d_c(11d_c + 276)}{2(24 + d_c)^3} \epsilon^2$

TABLE VI: Anomalous dimensions η'_2 , η_3 and η_4 obtained from the two-loop expansion of either the two-field or the effective model (this paper) – column 1 – from the SCSA [34, 35] – column 2 – and from the NPRG [26] – column 3. The two-loop value of η'_2 has been first obtained by [48].

NPRG. This approach is, as the SCSA, nonperturbative in the dimensional parameter $\epsilon = 4 - D$. It is based on the use of an exact RG equation that controls the evolution of a modified, running effective action with the running scale [66] (see [67–72] for reviews). Approximations of this equation are needed and consist in truncating the running effective action in powers of the field-derivatives (and, if necessary, of the field itself). They however lead to RG equations that remain nonpertur-

bative both in ϵ and in $1/d_c$. Such a procedure, called derivative expansion, has been validated empirically at order 4 in the derivative of the field [73, 74] and, more recently, up to order 6 [75], since one observes a rapid convergence of the physical quantities with the order in derivative. More formal argument for the *convergence* of the series – in contrast to the asymptotic nature of the usual, perturbative, series – have also been given in [75]. One should have in mind that this approach, although

nonperturbative and, as the SCSA, exact in a whole domain of parameters – at leading order in ϵ , in $1/d_c$, in the coupling constant controlling the interaction near the lower-critical dimension, at $d_c = 0$ – is nevertheless not exact and generally misses the next-to-leading order of the perturbative approaches. For instance, reproducing *exactly* the weak-coupling expansion at two-loop order requires the knowledge of the infinite series in derivatives [76, 77]. Yet, for a given field theory, the ability of the NPRG to reproduce satisfactorily this subleading contribution is a very good indication of its efficiency. The NPRG equations for the flat phase of membranes have been derived at the first order in derivative expansion in [26] and then with help of ansatz involving the full derivative content in [46, 47]. We give in Table VI, column 3, the anomalous dimensions obtained within this approach [26] and re-expanded here at second order in ϵ . First, one notes that, as in the SCSA case, the leading order result is exactly reproduced. Then one can observe that the next-to-leading order is also numerically close or very close to those obtained within the two-loop computation.

It is also interesting to mention that, for the SCSA, the coordinates of the fixed point P_3 obey the condition of vanishing bulk modulus

$$B = O(\epsilon^3) \quad (14)$$

whereas those of the fixed point P_4 obey the identity:

$$(6 - \epsilon)\lambda_4^* + 2\mu_4^* = O(\epsilon^3). \quad (15)$$

The properties (14) and (15) are, in fact, true nonperturbatively in ϵ at least within the first order in the derivative expansion performed in [26] and, again, in agreement with the SCSA result (13).

Finally, one should recall that the result obtained in $D = 2$ by means of the NPRG approach [26, 45] $\eta_{NPRG}^{D=2} = 0.849(3)$ is also very close to that provided by several numerical approaches (see, e.g., [59, 61, 65] that lead to $\eta \simeq 0.85$).

GCI model. We conclude by quoting a very recent – and first – two-loop, weak-coupling perturbative approach to membranes that has been performed by Mauri and Katsnelson [48] on a variant of the effective model (3) named Gaussian curvature interaction (GCI) model. It is obtained by generalizing to any dimension D the

simplified form of the usual effective model (3), i.e. with $M_{ab,cd} = 0$, valid in the particular case $D = 2$. As a consequence the authors of [48] get a – unique – nontrivial fixed point which, in our context, is nothing but the fixed point P'_2 . One of the main results of their analysis is that the two-loop anomalous dimension η'_2 coincides exactly with the corresponding SCSA result, a fact which is also observed in Table VI. Our analysis of the complete theory shows that, for the stable fixed point P_4 , a small discrepancy between the two-loop and the SCSA results occurs.

V. CONCLUSION

We have performed the two-loop, weak coupling analysis of the two models describing the flat phase of polymerized membranes. We have determined the RG equations and the anomalous dimensions at this order. We have identified the fixed points, analyzed their properties and computed the corresponding anomalous dimensions. First, one notes that although the coordinates of the fixed points, as well as several D -dependent quantities, vary from one model to the other, the anomalous dimensions at the fixed points are very robust as we get the same values from the two models. This provides a very strong check of our computations. It remains nevertheless to understand more profoundly the interplay between the dimensional regularization used here and these D -dependent quantities that are inherent in theories with space-time symmetries, such as the present one. Second, the very good agreement between the anomalous dimensions computed in our paper with those obtained from the SCSA and NPRG approaches is a confirmation of the extreme efficiency of these last methods in the context of the theory of the flat phase of polymerized membranes. As said, these two approaches have in common that they both reproduce exactly – by construction – the leading order of all usual perturbative approaches. This, however, does not explain their singular achievements here which more likely rely on the very nature of the flat phase of membranes itself. This is under investigation.

Acknowledgments

We wish to thank warmly J. Gracey, M. Kompaniets and K. J. Wiese for very fruitful discussions.

-
- [1] D. R. Nelson, T. Piran, and S. Weinberg, eds., *Proceedings of the Fifth Jerusalem Winter School for Theoretical Physics* (World Scientific, Singapore, 2004), 2nd ed.
 - [2] M. J. Bowick and A. Travesset, *Phys. Rep.* **344**, 255 (2001).
 - [3] A. M. Polyakov, *Gauge Field and Strings* (Gordon and Breach, 1987).
 - [4] F. David, *Phys. Rep.* **184**, 221 (1989).
 - [5] J. Wheeler, *J. Phys. A* **27**, 3323 (1994).
 - [6] See the contribution of F. David in [1].
 - [7] C. F. Schmidt, K. Svoboda, N. Lei, I. B. Petsche, L. E. Berman, C. R. Safinya, and G. S. Grest, *Science* **259**, 952 (1993).
 - [8] K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, Y. Zhang, S. V. Dubonos, I. V. Gregorieva, and A. A. Firsov, *Science* **306**, 666 (2004).

- [9] K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, M. I. Katsnelson, I. V. Gregorieva, S. V. Dubonos, and A. A. Firsov, *Nature* **438**, 197 (2005).
- [10] M. I. Katsnelson, *Graphene: Carbon in Two Dimensions* (Cambridge University Press, Cambridge, U.K., 2012).
- [11] See the contribution of D. R. Nelson in Ref.[1].
- [12] D. R. Nelson, *Defects and Geometry in Condensed Matter Physics* (Cambridge University Press, Cambridge, UK, 2002).
- [13] P.-G. de Gennes and C. Taupin, *J. Phys. Chem* **86**, 2294 (1982).
- [14] L. Peliti and S. Leibler, *Phys. Rev. Lett.* **54**, 1690 (1985).
- [15] W. Helfrich, *J. Phys. France* **46**, 1263 (1985).
- [16] N. D. Mermin and H. Wagner, *Phys. Rev. Lett.* **17**, 1133 (1966).
- [17] D. R. Nelson and L. Peliti, *J. Phys. (Paris)* **48**, 1085 (1987).
- [18] J. A. Aronovitz and T. C. Lubensky, *Phys. Rev. Lett.* **60**, 2634 (1988).
- [19] F. David and E. Guitter, *Europhys. Lett.* **5**, 709 (1988).
- [20] E. Guitter, F. David, S. Leibler, and L. Peliti, *J. Phys. (Paris)* **50**, 1787 (1989).
- [21] M. Paczuski and M. Kardar and D. R. Nelson, *Phys. Rev. Lett.* **60**, 2638 (1988).
- [22] J. A. Aronovitz, L. Golubovic, and T. C. Lubensky, *J. Phys. (Paris)* **50**, 609 (1989).
- [23] J.-P. Kownacki and H. Diep, *Phys. Rev. E* **66**, 066105 (2002).
- [24] H. Koibuchi and N. Kusano and A. Nidaira and K. Suzuki and M. Yamada, *Phys. Rev. E* **69**, 066139 (2004).
- [25] H. Koibuchi and T. Kuwahata, *Phys. Rev. E* **72**, 026124 (2005).
- [26] J.-P. Kownacki and D. Mouhanna, *Phys. Rev. E* **79**, 040101(R) (2009).
- [27] H. Koibuchi and A. Shobukhov, *Int. J. Mod. Phys. C* **25**, 1450033 (2014).
- [28] K. Essafi, J.-P. Kownacki, and D. Mouhanna, *Phys. Rev. E* **89**, 042101 (2014).
- [29] U. Satoshi and H. Koibuchi, *J. Stat. Phys.* **162**, 701 (2016).
- [30] R. Cuerno, R. Gallardo Caballero, A. Gordillo-Guerrero, P. Monroy and J. J. Ruiz-Lorenzo, *Phys. Rev. E* **93**, 022111 (2016).
- [31] M. J. Bowick, A. Cacciuto, G. Thoroleifsson, and A. Travesset, *Eur. Phys. J. E* **5**, 149 (2001).
- [32] E. Guitter, F. David, S. Leibler, and L. Peliti, *Phys. Rev. Lett.* **61**, 2949 (1988).
- [33] I. V. Gornyi, V. Yu. Kachorovskii and A. D. Mirlin, *Phys. Rev. B* **92**, 155428 (2015).
- [34] P. Le Doussal and L. Radzihovsky, *Phys. Rev. Lett.* **69**, 1209 (1992).
- [35] P. Le Doussal and L. Radzihovsky, *Ann. Phys. (N.Y.)* **392**, 340 (2018).
- [36] O. Coquand, *Phys. Rev B* **100**, 125406 (2019).
- [37] O. Coquand, D. Mouhanna, and S. Teber, unpublished (2020).
- [38] D. R. Saykin, I. V. Gornyi, V. Yu. Kachorovskii and I. S. Burmistrov, *Ann. Phys. (N.Y.)* **414**, 168108 (2020).
- [39] K. V. Zakharchenko, R. Roldán, A. Fasolino and M. I. Katsnelson, *Phys. Rev. B* **82**, 125435 (2010).
- [40] R. Roldán, A. Fasolino, K. V. Zakharchenko, and M. I. Katsnelson, *Phys. Rev. B* **83**, 174104 (2011).
- [41] D. Gazit, *Phys. Rev. E* **80**, 041117 (2009).
- [42] K. Essafi, J.-P. Kownacki, and D. Mouhanna, *Phys. Rev. Lett.* **106**, 128102 (2011).
- [43] O. Coquand and D. Mouhanna, *Phys. Rev. E* **94**, 032125 (2016).
- [44] O. Coquand, K. Essafi, J.-P. Kownacki, and D. Mouhanna, *Phys. Rev E* **97**, 030102(R) (2018).
- [45] O. Coquand, K. Essafi, J.-P. Kownacki, and D. Mouhanna, *Phys. Rev. E* **101**, 042602 (2020).
- [46] F. L. Braghin and N. Hasselmann, *Phys. Rev. B* **82**, 035407 (2010).
- [47] N. Hasselmann and F. L. Braghin, *Phys. Rev. E* **83**, 031137 (2011).
- [48] A. Mauri and M.I. Katsnelson, *Nucl. Phys. B* **956**, 115040 (2020).
- [49] V. A. Smirnov and K. G. Chetyrkin, *Theor. Math. Phys.* **56**, 770 (1984).
- [50] N. N. Bogoliubov and O. S. Parasiuk, *Acta Math.* **97**, 227 (1957).
- [51] K. Hepp, *Commun. Math. Phys.* **2**, 301 (1966).
- [52] W. Zimmermann, *Commun. Math. Phys.* **15**, 208 (1969).
- [53] R. N. Lee, *J. Phys.: Conf. Ser.* **523**, 012059 (2014).
- [54] A. V. Kotikov and S. Teber, *Phys. Part. Nucl.* **50**, 1 (2019).
- [55] E. Guitter, S. Leibler, A. Maggs, and F. David, *J. Phys. (Paris)* **51**, 1055 (1990).
- [56] Z. Zhang, H. T. Davis, and D. M. Kroll, *Phys. Rev. E* **48**, R651 (1993).
- [57] M. J. Bowick et. al., *J. Phys. (France) I* **6**, 1321 (1996).
- [58] G. Gompper and D. M. Kroll, *J. Phys: Condens. Matter* **9**, 8795 (1997).
- [59] J. H. Los, M. I. Katsnelson, O. V. Yazyev, K. V. Zakharchenko, and A. Fasolino, *Phys. Rev. B* **80**, 121405(R) (2009).
- [60] A. Tröster, *Phys. Rev. B* **87**, 104112 (2013).
- [61] D. Wein and F. Wang, *J. Chem. Phys.* **141**, 144701 (2014).
- [62] A. Tröster, *Phys. Rev. B* **91**, 022132 (2015).
- [63] J. H. Los, A. Fasolino and M. I. Katsnelson, *Phys. Rev. Lett.* **116**, 015901 (2016).
- [64] A. Kosmrlj and D.R. Nelson, *Phys. Rev. X* **7**, 011002 (2017).
- [65] J. Hasik, E. Tosatti and R. Martonak, *Phys. Rev. B* **97**, 140301(R) (2018).
- [66] C. Wetterich, *Phys. Lett. B* **301**, 90 (1993).
- [67] C. Bagnuls and C. Bervillier, *Phys. Rep.* **348**, 91 (2001).
- [68] J. Berges, N. Tetradis, and C. Wetterich, *Phys. Rep.* **363**, 223 (2002).
- [69] B. Delamotte, D. Mouhanna, and M. Tissier, *Phys. Rev. B* **69**, 134413 (2004).
- [70] J. Pawłowski, *Ann. Phys. (N.Y.)* **322**, 2831 (2007).
- [71] O. Rosten, *Phys. Rep.* **511**, 177 (2012).
- [72] B. Delamotte, *Lect. Notes Phys.* **852**, 49 (2012).
- [73] L. Canet, B. Delamotte, D. Mouhanna, and J. Vidal, *Phys. Rev. B* **68**, 064421 (2003).
- [74] G. De Polsi, I. Balog, M. Tissier and N. Wschebor, *Phys. Rev. E* **101**, 042113 (2020).
- [75] I. Balog, H. Chaté, B. Delamotte, M. Marohni and N. Wschebor, *Phys. Rev. Lett.* **123**, 240604 (2019).
- [76] T. Papenbrock and C. Wetterich, *Z. Phys. C* **65**, 519 (1995).
- [77] T. R. Morris and J. F. Tighe, *J. High Energy Phys.* **08**, 007 (1999).