ON A CLASS OF STOCHASTIC PARTIAL DIFFERENTIAL EQUATIONS WITH MULTIPLE INVARIANT MEASURES

BÁLINT FARKAS, MARTIN FRIESEN, BARBARA RÜDIGER, AND DENNIS SCHROERS

Abstract. In this work we investigate the long-time behavior, that is the existence and characterization of invariant measures as well as convergence of transition probabilities, for Markov processes obtained as the unique mild solution to stochastic partial differential equations in a Hilbert space. Contrary to the existing literature where typically uniqueness of invariant measures is studied, we focus on the case where uniqueness of invariant measures fails to hold. Namely, using a generalized dissipativity condition combined with a decomposition of the Hilbert space, we prove the existence of multiple limiting distributions in dependence of the initial state of the process, and study convergence of transition probabilities in the Wasserstein 2-distance. Finally, we show that these results contain Lévy driven Ornstein-Uhlenbeck processes, the Heath-Jarrow-Morton-Musiela equation as well as stochastic partial differential equations with delay as a particular case.

1. INTRODUCTION

Stochastic partial differential equations arise in the modelling of applications in mathematical physics (e.g. Navier-Stokes equations [\[21,](#page-25-0) [17,](#page-25-1) [9,](#page-25-2) [35\]](#page-26-0) or stochastic non-linear Schrödinger equations $[4, 13]$ $[4, 13]$, biology (e.g. catalytic branching processes $[12, 28]$ $[12, 28]$), and finance (e.g. forward prices [\[23,](#page-25-6) [36,](#page-26-1) [15\]](#page-25-7)). While the construction of solutions to the underlying stochastic equations is an important mathematical issue, having applications in mind it is indispensable to also study their specific properties. Among them, an investigation of the long-time behavior of solutions, that is existence and uniqueness of invariant measures and convergence of transition probabilities, are often important and at the same time also challenging mathematical topics. In this work we investigate the long-time behavior of mild solutions to the stochastic partial differential equation of the form

$$
dX_t = (AX_t + F(X_t))dt + \sigma(X_t)dW_t + \int_E \gamma(X_t,\nu)\widetilde{N}(dt,d\nu), \qquad t \ge 0 \tag{1.1}
$$

on a separable Hilbert space H, where $(A, D(A))$ is the generator of a strongly continuous semigroup $(S(t))_{t\geq0}$ on H, $(W_t)_{t\geq0}$ is a Q-Wiener process and $\widetilde{N}(dt, d\nu)$ denotes a compensated Poisson random measure. The precise conditions need to be imposed on these objects will be formulated in the subsequent sections.

In the literature the study on the existence and uniqueness of invariant measures often relies on different variants of a *dissipativity condition*. The simplest form of such a dissipativity condition is: There exists $\alpha > 0$ such that

$$
\langle Ax - Ay, x - y \rangle_H + \langle F(x) - F(y), x - y \rangle_H \le -\alpha \|x - y\|_H^2, \qquad x, y \in D(A). \tag{1.2}
$$

Indeed, if [\(1.2\)](#page-0-0) is satisfied, σ and γ are globally Lipschitz-continuous, and α is large enough, then there exists a unique invariant measure for the Markov process obtained

Date: May 5, 2020.

²⁰¹⁰ Mathematics Subject Classification. Primary 37L40, 60J25; Secondary 37A30, 37L55.

from [\(1.1\)](#page-0-1), see, e.g., [\[30,](#page-25-8) Section 16], [\[10,](#page-25-9) Chapter 11, Section 6], and [\[34\]](#page-26-2). Note that (1.2) is satisfied, if F is globally Lipschitz continuous and $(A, D(A))$ satisfies for some $\beta > 0$ large enough the inequality $\langle Ax, x \rangle_H \leq -\beta ||x||_H^2$, $x \in D(A)$, i.e. $(A, D(A))$ is the generator of a strongly continuous semigroup satisfying $||S(t)||_{L(H)} \le e^{-\beta t}$. Here and below we denote by $L(H)$ the space of bounded linear operators from H to H and by $\|\cdot\|_{L(H)}$ its operator norm. For weaker variants of the dissipativity condition (e.g. cases where [\(1.2\)](#page-0-0) only holds for $||x||_H, ||y||_H \geq R$ for some $R > 0$), in general one can neither guarantee the existence nor uniqueness of an invariant measure. Hence, to treat such cases, additional arguments, e.g. coupling methods, are required. Such arguments have been applied to different stochastic partial differential equations on Hilbert spaces in [\[31,](#page-25-10) [32,](#page-25-11) [33\]](#page-26-3) where existence and, in particular, uniqueness of invariant measures was studied. We also mention [\[8,](#page-25-12) [22\]](#page-25-13) for an extension of Harris-type theorems for Wasserstein distances, and [\[24,](#page-25-14) [20\]](#page-25-15) for extensions of coupling methods.

In contrast to the aforementioned methods and applications, several stochastic models exhibit phase transition phenomena where uniqueness of invariant measures fails to hold. For instance, the generator $(A, D(A))$ and drift F appearing in the Heath-Jarrow-Morton-Musiela equation do not satisfy (1.2) , but instead F is globally Lipschitz continuous and the semigroup generated by $(A, D(A))$ satisfies $||S(t)x - Px||_H \leq$ $e^{-\alpha t} \|x - Px\|_H$ for some projection operator P. Based on this property it was shown in [\[36,](#page-26-1) [34\]](#page-26-2) that the Heath-Jarrow-Morton-Musiela equation has infinitely many invariant measures parametrized by the initial state of the process, see also Section 6. Another example is related to stochastic Volterra equations as studied, e.g., in [\[6\]](#page-24-1). There, using a representation of stochastic Volterra equations via SPDEs and combined with some arguments originated from the study of the Heath-Jarrow-Morton-Musiela equation, the authors studied existence of limiting distributions allowing, in particular, that these distributions depend on the initial state of the process.

In this work we provide a general and unified approach for the study of multiple invariant measures and, moreover, we show that with dependence on the initial distri-bution the law of the mild solution of [\(1.1\)](#page-0-1) is governed in the limit $t \to \infty$ by one of the invariant measures. In particular, we show that the methods developed in [\[36,](#page-26-1) [34,](#page-26-2) [6\]](#page-24-1) can be embedded as a special case of a general framework where one replaces [\(1.2\)](#page-0-0) by a weaker dissipativity condition, which we call *generalized dissipativity condition*:

(GDC) There exists a projection operator P_1 on the Hilbert space H and there exist constants $\alpha > 0, \beta \geq 0$ such that, for $x, y \in D(A)$, one has:

$$
\langle Ax - Ay, x - y \rangle_H + \langle F(x) - F(y), x - y \rangle_H
$$

$$
\leq -\alpha ||x - y||_H^2 + (\alpha + \beta) ||P_1x - P_1y||_H^2.
$$

Restricting $x, y \in D(A)$ to the subspace $x, y \in \text{ker}(P_1) \cap D(A)$ shows that (GDC) contains the classical dissipativity condition on the subspace ker(P_1) $\subset H$. Contrary, restricting x, y to ran(P_1) ∩ $D(A)$ does not yield the classical dissipativity condition but instead contains the additional term $||P_1x - P_1y||_H^2$ which describes the influence of the non-dissipative part of the drift. Sufficient conditions and additional remarks, e.g., on this condition are collected in the end of Section 2 while particular examples are discussed in Sections $5 - 7$. Let us mention that (GDC) is satisfied if F is globally Lipschitz continuousmthe semigroup $(S(t))_{t\geq 0}$ is symmetric (i.e. $S(t)^* = S(t)$ for $t \geq 0$), and uniformly convergent to P_1 (i.e. $||S(t) - P_1||_{L(H)} \longrightarrow 0$ as $t \to \infty$).

Roughly speaking, we will show that under condition (GDC) and some restrictions on the projected coefficients P_1F , $P_1\sigma$, and $P_1\gamma$, the Markov process obtained from [\(1.1\)](#page-0-1) has for each initial data $X_0 = x$ a limiting distribution π_x depending only on P_1x .

Moreover, the transition probabilities converge exponentially fast in the Wasserstein 2 distance to this limiting distributions. In order to prove this result, we first decompose the state space H according to

$$
H = H_0 \oplus H_1, \qquad x = P_0 x + P_1 x, \quad P_0 := I - P_1,
$$

where I denotes the identity operator on H, and then investigate the components P_0X_t and P_1X_t separately. Based on an idea taken from [\[37\]](#page-26-4), we construct, for each $\tau \geq 0$, a coupling of X_t and $X_{t+\tau}$. This coupling will be then used to efficiently estimate the Wasserstein 2-distance for the solution started at two different points.

This work is organized as follows. In Section 2 we state the precise conditions imposed on the coefficients of the SPDE [\(1.1\)](#page-0-1), discuss some properties of the solution and then provide sufficient conditions for the generalized dissipativity condition (GDC). Based on condition (GDC) we derive in Section 3 an estimate on the trajectories of the process when started at two different initial points, i.e. we estimate the L^2 -norm of $X_t^x - X_t^y$ t when $x \neq y$. Based on this estimate, we state and prove our main results in Section 4. Examples are then discussed in the subsequent Sections $5 - 7$. In Section 5 we explicitly characterize the limiting distributions of the Lévy driven Ornstein-Uhlenbeck process with an operator $(A, D(A))$ generating an uniformly convergent semigroup. The Heath-Jarrow-Morton-Musiela equation is then considered in Section 6 for which we first show that the main results of Section 4 contain [\[36,](#page-26-1) [34\]](#page-26-2), and then extend these results by characterizing its limiting distributions more explicitly. Finally, we apply our results in Section 7 to an SPDE with delay.

2. Preliminaries

2.1. **Framework and notation.** Here and throughout this work, $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P})$ is a filtered probability space satisfying the usual conditions. Let U be a separable Hilbert space and $W = (W_t)_{t \in \mathbb{R}_+}$ be a Q-Wiener process with respect to $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ on $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P})$, where $Q: U \to U$ is a non-negative, symmetric, trace class operator. Let E be a Polish space, $\mathcal E$ the Borel- σ -field on E, and μ a σ -finite measure on $(E, \mathcal E)$. Let $N(dt, d\nu)$ be a $(\mathcal{F}_t)_{t>0}$ -Poisson random measure with compensator $dt\mu(d\nu)$ and denote by $\tilde{N}(dt, d\nu) = N(dt, d\nu) - dt\mu(d\nu)$ the corresponding compensated Poisson random measure. Suppose that the random objects $(W_t)_{t>0}$ and $N(dt, d\nu)$ are mutually independent.

In this work we investigate the long-time behavior of mild solutions to the stochastic partial differential equation

$$
\begin{cases} dX_t^x = (AX_t^x + F(X_t^x))dt + \sigma(X_t^x)dW_t + \int_E \gamma(X_t^x, \nu)\widetilde{N}(dt, d\nu), & t \ge 0, \\ X_0^x = x \in L^2(\Omega, \mathcal{F}_0, \mathbb{P}; H), \end{cases}
$$
(2.1)

where $(A, D(A))$ is the generator of a strongly continuous semigroup $(S(t))_{t>0}$ on H, $H \ni x \mapsto F(x) \in H$ and $H \ni x \mapsto \sigma(x) \in L_2^0$ are Borel measurable mappings, and $(x, \nu) \mapsto \gamma(x, \nu)$ is measurable from $(H \times E, \mathcal{B}(H) \otimes \mathcal{E})$ to $(H, \mathcal{B}(H))$. Here $\mathcal{B}(H)$ denotes the Borel- σ -algebra on H, and $L_2^0 := L_2^0(H)$ is the Hilbert space of all Hilbert-Schmidt operators from U_0 to H, where $U_0 := Q^{1/2}U$ is a separable Hilbert space endowed with the scalar product

$$
\langle x, y \rangle_0 := \langle Q^{-1/2} x, Q^{-1/2} y \rangle_U = \sum_{k \in \mathbb{N}} \frac{1}{\lambda_k} \langle x, e_k \rangle_U \langle e_k, y \rangle_U, \qquad \forall x, y \in U_0,
$$

and $Q^{-1/2}$ denotes the pseudoinverse of $Q^{1/2}$. Here $(e_j)_{j\in\mathbb{N}}$ denotes an orthogonal basis of eigenvectors of Q in U with corresponding eigenvalues $(\lambda_j)_{j\in\mathbb{N}}$. For comprehensive introductions to integration concepts in infinite dimensional settings we refer to [\[10\]](#page-25-9) for the case of Q-Wiener processes and to [\[26\]](#page-25-16) for compensated Poisson random measures as integrators. Throughout this work we suppose that the coefficients F, σ, γ are Lipschitz continuous. More precisely:

(A1) There exist constants $L_F, L_{\sigma}, L_{\gamma} \geq 0$ such that for all $x, y \in H$

$$
||F(x) - F(y)||_H^2 \le L_F ||x - y||_H^2,
$$

\n
$$
||\sigma(x) - \sigma(y)||_{L_2^0(H)}^2 \le L_\sigma ||x - y||_H^2,
$$

\n
$$
\int_E ||\gamma(x, \nu) - \gamma(y, \nu)||_H^2 \mu(d\nu) \le L_\gamma ||x - y||_H^2.
$$
\n(2.2)

Moreover we suppose that

$$
\int_{E} \|\gamma(0,\nu)\|_{H}^{2}\mu(d\nu) < \infty.
$$
\n(2.3)

Note that condition [\(2.3\)](#page-3-0) implies that the jumps satisfy the usual growth conditions, i.e.

$$
\int_{E} \|\gamma(x,\nu)\|_{H}^{2}\mu(d\nu) \le 2 \int_{E} \|\gamma(x,\nu) - \gamma(0,\nu)\|_{H}^{2}\mu(d\nu) + 2 \int_{E} \|\gamma(0,\nu)\|_{H}^{2}\mu(d\nu) \n\le 2 \max \left\{ L_{\gamma}, \int_{E} \|\gamma(0,\nu)\|_{H}^{2}\mu(d\nu) \right\} (1 + \|x\|_{H}^{2}).
$$

Moreover, it follows from (GDC) and (A1) that

$$
\langle Ax, x \rangle_H \le \left(\beta + \sqrt{L_F}\right) ||x||_H^2, \qquad x \in D(A).
$$

Hence $A - (\beta + \sqrt{L_F})$ is dissipative and thus by the Lumer-Phillips theorem the semigroup $(S(t))_{t\geq0}$ generated by $(A, D(A))$ is quasi-contractive, i.e.

$$
||S(t)x||_H \le e^{\left(\beta + \sqrt{L_F}\right)t} ||x||_H, \quad x \in H. \tag{2.4}
$$

Then, under conditions (GDC) and (A1), for each initial condition $x \in L^2(\Omega, \mathcal{F}_0, \mathbb{P}; H)$ there exists a unique cádlág, $(F_t)_{t>0}$ -adapted, mean square continuous, mild solution $(X_t^x)_{t\geq 0}$ to [\(2.1\)](#page-2-0) such that, for each $T > 0$, there exists a constant $C(T) > 0$ satisfying

$$
\mathbb{E}\left[\sup_{t\in[0,T]}\|X_t^x\|_H^2\right] \le C(T)\left(1+\mathbb{E}\left[\|x\|_H^2\right]\right) \tag{2.5}
$$

and, for all $x, y \in L^2(\Omega, \mathcal{F}_0, \mathbb{P}; H)$,

$$
\mathbb{E}\left[\|X_t^x - X_t^y\|_H^2\right] \le C(T)\mathbb{E}\left[\|x - y\|_H^2\right], \qquad t \in [0, T].\tag{2.6}
$$

This means that $(X_t^x)_{t\geq 0}$ satisfies $\mathbb{P}\text{-a.s.}$

$$
X_t^x = S(t)x + \int_0^t S(t-s)F(X_s^x)ds + \int_0^t S(t-s)\sigma(X_s^x)dW_s
$$

+
$$
\int_0^t \int_E S(t-s)\gamma(X_s^x, \nu)\widetilde{N}(ds, d\nu), \qquad t \ge 0,
$$
 (2.7)

where all (stochastic) integrals are well-defined, see, e.g., [\[1\]](#page-24-2), [\[26\]](#page-25-16), and [\[16\]](#page-25-17). The obtained solution is a Markov process whose transition probabilities $p_t(x, dy) = \mathbb{P}[X_t^x \in dy]$ are measurable with respect to x. Denote by $(p_t)_{t\geq0}$ its transition semigroup, i.e., for each bounded measurable function $f : H \longrightarrow \mathbb{R}$, $p_t f$ is given by

$$
p_t f(x) = \mathbb{E}\left[f(X_t^x)\right] = \int_H f(y)p_t(x, dy), \qquad t \ge 0, \quad x \in H.
$$

Using the continuous dependence on the initial condition, see (2.6) , it can be shown that $p_t f \in C_b(H)$ for each $f \in C_b(H)$, i.e. the transition semigroup is C_b -Feller.

In this work we investigate the the existence of invariant measures and convergence of the transition probabilities towards these measures for the Markov process $(X_t^x)_{t\geq 0}$ with particular focus on the cases where uniqueness of invariant measures fails to hold. By slight abuse of notation, we denote by p_t^* the adjoint operator to p_t defined by

$$
p_t^* \rho(dx) = \int_H p_t(y, dx) \rho(dy), \qquad t \ge 0.
$$

Recall that a probability measure π on $(H, \mathcal{B}(H))$ is called *invariant measure* for the semigroup $(p_t)_{t\geq 0}$ if and only if $p_t^*\pi = \pi$ holds for each $t \geq 0$. Let $\mathcal{P}_2(H)$ be the space of Borel probability measures ρ on $(H,\mathcal{B}(H))$ with finite second moments. Recall that $\mathcal{P}_2(H)$ is separable and complete when equipped with the *Wasserstein-2-distance*

$$
\mathcal{W}_2(\rho, \widetilde{\rho}) = \inf_{G \in \mathcal{H}(\rho, \widetilde{\rho})} \left(\int_{H^2} ||x - y||_H^2 G(dx, dy) \right)^{\frac{1}{2}}, \qquad \rho, \widetilde{\rho} \in \mathcal{P}_2(H), \tag{2.8}
$$

where $\mathcal{H}(\rho, \tilde{\rho})$ denotes the set of all couplings of $(\rho, \tilde{\rho})$, i.e. Borel probability measures on $H \times H$ whose marginals are given by ρ and $\tilde{\rho}$, respectively, see [\[38,](#page-26-5) Section 6] for a general introduction to couplings and Wasserstein distances.

2.2. Discussion of generalized dissipativity condition. In this section we briefly discuss the condition

$$
\langle Ax, x \rangle_H \le -\lambda_0 \|x\|_H^2 + (\lambda_0 + \lambda_1) \|P_1 x\|_H^2, \qquad x \in D(A), \tag{2.9}
$$

where $\lambda_0 > 0$ and $\lambda_1 \geq 0$. Note that, if [\(2.9\)](#page-4-0) and condition [\(2.1\)](#page-2-0) are satisfied, then

$$
\langle Ax - Ay, x - y \rangle_H + \langle F(x) - F(y), x - y \rangle_H
$$

\n
$$
\leq \langle Ax - Ay, x - y \rangle_H + \sqrt{L_F} ||x - y||_H^2
$$

\n
$$
\leq - \left(\lambda_0 - \sqrt{L_F} \right) ||x - y||_H^2 + \left(\lambda_0 + \lambda_1 \right) ||P_1x - P_1y||_H^2,
$$
\n(2.10)

i.e. the generalized dissipativity condition (GDC) is satisfied for $\alpha = \lambda_0 - \sqrt{L_F}$ and $\beta = \lambda_1 + \sqrt{L_F}$, provided that $\lambda_0 > \sqrt{L_F}$.

Proposition 2.1. *Suppose that there exists an orthogonal decomposition* $H = H_0 \oplus H_1$ *of* H into closed linear subspaces $H_0, H_1 \subset H$ such that $(S(t))_{t>0}$ leaves H_0 and H_1 *invariant and there exist constants* $\lambda_0 > 0$ *and* $\lambda_1 \geq 0$ *satisfying*

$$
||S(t)x_0||_H \le e^{-\lambda_0 t} ||x_0||_H
$$
, $||S(t)x_1||_H \le e^{\lambda_1 t} ||x_1||_H$, $\forall t \ge 0$.

for all $x_0 \in H_0$ *and* $x_1 \in H_1$ *. Then* [\(2.9\)](#page-4-0) *holds for* P_1 *being the orthogonal projection operator onto* H_1 .

Proof. Let P_0 be the orthogonal projection operator onto H_0 . Since $(S(t))_{t>0}$ leaves the closed subspace H_0 invariant, its restriction $(S(t)|_{H_0})_{t\geq0}$ onto H_0 is a strongly continuous semigroup of contractions on H_0 with generator $(A_0, D(A_0))$ being the H_0 part of A, that is

$$
A_0x = Ax, \qquad x \in D(A_0) = \{ y \in D(A) \cap H_0 \mid Ay \in H_0 \}.
$$

Since H_0 is closed and $S(t)$ leaves H_0 invariant, it follows that $Ay = \lim_{t\to 0} \frac{S(t)y-y}{t} \in H_0$ for $y \in D(A) \cap H_0$, i.e. $D(A_0) = D(A) \cap H_0$ and $P_0 : D(A) \to D(A_0)$. Let P_1 the orthogonal projection operator onto H_1 . Arguing exactly in the same way shows that the restriction $(S(t)|_{H_1})_{t\geq 0}$ is a strongly continuous semigroup of contractions on H_1 with generator $(A_1, D(A_1))$ given by $A_1x = Ax$ and $x \in D(A_1) = D(A) \cap H_1$ so that $P_1: D(A) \to D(A_1)$. Since $S(t)$ leaves H_0 and H_1 invariant, we obtain $P_0S(t) = S(t)P_0$, $P_1S(t) = S(t)P_1$ from which we conclude that $AP_1x = P_1Ax$ and $AP_0x = P_0Ax$ for $x \in D(A).$

Since $(e^{\lambda_0 t}S(t)|_{H_0})_{t\geq 0}$ is a strongly continuous semigroup of contractions on H_0 with generator $A_0 + \lambda_0 I$, and $(e^{-\lambda_1 t}S(t)|_{H_1})_{t \geq 0}$ is a strongly continuous semigroup of contractions on H_1 with generator $A_1 - \lambda_1 I$, we have by the Lumer-Phillips theorem (see [\[29,](#page-25-18) Theorem 4.3])

 $\langle A_0 x_0, x_0 \rangle_H \leq -\lambda_0 \|x_0\|_H^2$ and $\langle A_1 x_1, x_1 \rangle_H \leq \lambda_1 \|x_1\|_H^2$, $x_0 \in H_0$, $x_1 \in H_1$. Hence we find that

$$
\langle Ax, x \rangle_H = \langle Ax, P_0x \rangle_H + \langle Ax, P_1x \rangle_H
$$

= $\langle P_0Ax, P_0x \rangle_H + \langle P_1Ax, P_1x \rangle_H$
= $\langle A_0P_0x, P_0x \rangle_H + \langle A_1P_1x, P_1x \rangle_H$
 $\leq -\lambda_0 \|P_0x\|_H^2 + \lambda_1 \|P_1x\|_H^2$
= $-\lambda_0 \|x\|_H^2 + (\lambda_0 + \lambda_1) \|P_1x\|_H^2$,

where the last equality follows from $H_0 \perp H_1$. This proves the assertion.

At this point it is worthwhile to mention that Onno van Gaans has investigated in [\[37\]](#page-26-4) ergodicity for a class of Lévy driven stochastic partial differential equations where the semigroup $(S(t))_{t\geq0}$ was supposed to be hyperbolic. Proposition [2.1](#page-4-1) also covers this case, provided that the hyperbolic decomposition is orthogonal. The conditions of previous proposition are satisfied whenever $(S(t))_{t>0}$ is a symmetric, uniformly convergent semigroup.

Remark 2.2. *Suppose that* $(S(t))_{t>0}$ *is a strongly continuous semigroup on* H *and there exists an orthogonal projection operator* P *on* H *and* $\lambda_0 > 0$ *such that*

$$
||S(t)x - Px||_H \le e^{-\lambda_0 t} ||x - Px||_H, \qquad t \ge 0, \quad x \in H. \tag{2.11}
$$

Then the conditions of Proposition [2.1](#page-4-1) *are satisfied for* $H_0 = \text{ker}(P)$ *and* $H_1 = \text{ran}(P)$ *with* $\lambda_0 > 0$ *and* $\lambda_1 = 0$ *. In particular,* $(S(t))_{t \geq 0}$ *is a semigroup of contractions.*

The following example shows that [\(2.9\)](#page-4-0) can also be satisfied for non-symmetric and non-convergent semigroups.

Example 2.3. Let $H = \mathbb{R}^2$, $H_0 = \mathbb{R} \times \{0\}$, $H_1 = \{0\} \times \mathbb{R}$, and denote by P_0, P_1 the *projection operators onto* H_0 *and* H_1 *, respectively. Let* A *be given by* $A =$ $\begin{pmatrix} -1 & 1 \\ 0 & 1 \end{pmatrix}$. *Then*

$$
\left\langle \begin{pmatrix} x \\ y \end{pmatrix}, A \begin{pmatrix} x \\ y \end{pmatrix} \right\rangle = -x^2 + xy + y^2
$$

\n
$$
\leq -\frac{1}{2}(x^2 + y^2) + 2y^2
$$

\n
$$
= -\frac{1}{2} ||(x, y)||_H^2 + 2||P_1(x, y)||_H^2,
$$

i.e. [\(2.9\)](#page-4-0) *holds for* $\lambda_0 = \frac{1}{2}$ $rac{1}{2}$ and $\lambda_1 = \frac{3}{2}$ $\frac{3}{2}$ *. Since* $e^{tA} = \begin{pmatrix} e^{-t} & \frac{e^{t} - e^{-t}}{2} \\ 0 & e^{t} \end{pmatrix}$ \setminus *, it is clear that neither the conditions of Proposition [2.1](#page-4-1) nor of Remark [2.2](#page-5-0) are satisfied.*

3. Key stability estimate

Define, for $x, y \in D(A)$, the function

$$
\mathcal{L}(\|\cdot\|_H^2)(x, y) := 2\langle A(x - y) + F(x) - F(y), x - y \rangle_H + \|\sigma(x) - \sigma(y)\|_{L_2^0(H)}^2 + \int_E \|\gamma(x, \nu) - \gamma(y, \nu)\|_H^2 \mu(d\nu).
$$

Remark that under the additional assumption that [\(1.1\)](#page-0-1) has a strong solution, the function

$$
\mathcal{L}(\|\cdot\|_H^2)(z) := 2\langle A(z)+F(z),z\rangle_H + \|\sigma(z)\|_{L_2^0(H)}^2 + \int_E \|\gamma(z,\nu)\|_H^2 \mu(d\nu).
$$

is simply the generator $\mathcal L$ applied to the unbounded function $||z||_H^2$, see, e.g.. [\[2,](#page-24-3) equation (3.4)]). Since we work with mild solutions instead, all computations given below require to use additionally Yosida approximations for the mild solution of [\(1.1\)](#page-0-1).

Below we first prove a Lyapunov-type estimate for $\mathcal{L}(\|\cdot\|_H^2)$ and then deduce from that by an application of the Itô-formula for mild solutions to (2.1) an estimate for the L^2 -norm of $\hat{X}_t^x - X_t^y$ $_t^y$.

Lemma 3.1. *Assume that condition (GDC) and (A1) are satisfied. Then*

$$
\mathcal{L}(\|\cdot\|_H^2)(x,y) \leq -(2\alpha - L_{\sigma} - L_{\gamma})\|x - y\|_H^2 + 2(\alpha + \beta)\|P_1x - P_1y\|_H^2 \tag{3.1}
$$

holds for $x, y \in D(A)$ *.*

Proof. Using first (A1) and then (GDC) we find that

$$
\mathcal{L}(\|\cdot\|_{H}^{2})(x,y) \le (L_{\sigma} + L_{\gamma}) \|x - y\|_{H}^{2} \n+ 2\langle Ax - Ay, x - y \rangle_{H} + 2\langle F(x) - F(y), x - y \rangle_{H} \n\le -(2\alpha - L_{\sigma} - L_{\gamma}) \|x - y\|_{H}^{2} + 2(\alpha + \beta) \|P_{1}x - P_{1}y\|_{H}^{2}.
$$

This proves the asserted inequality.

The following is our key stability estimate.

Proposition 3.2. *Suppose that (GDC) and (A1) are satisfied, that*

$$
\varepsilon := 2\alpha - L_{\sigma} - L_{\gamma} > 0,\tag{3.2}
$$

and suppose that

$$
\sup_{x \in H} \int_{E} \|\gamma(x,\nu)\|^4 \mu(d\nu) < \infty. \tag{3.3}
$$

Then, for each $x, y \in L^2(\Omega, \mathcal{F}_0, \mathbb{P}; H)$ *and all* $t \geq 0$ *,*

$$
\mathbb{E}\left[\|X_t^x - X_t^y\|_H^2\right]
$$

$$
\leq e^{-\varepsilon t} \mathbb{E}\left[\|x - y\|_H^2\right] + 2(\alpha + \beta) \int_0^t e^{-\varepsilon(t-s)} \mathbb{E}\left[\|P_1 X_s^x - P_1 X_s^y\|_H^2\right] ds.
$$
 (3.4)

Proof. To simplify the notation, denote by $(X_t)_{t>0}$ the mild solution to [\(1.1\)](#page-0-1) with initial condition x and by $(Y_t)_{t>0}$ the mild solution to [\(1.1\)](#page-0-1) with initial condition y. Moreover, we write $(X_t^n)_{t\geq 0}$ and $(Y_t^n)_{t\geq 0}$ for the strong solutions to the corresponding Yosida -approximation systems

$$
\begin{cases} dX_t^n = AX_t^n + R_n F(X_t^n)dt + R_n \sigma(X_t^n) dW_t + \int_E R_n \gamma(X_t^n, \nu) \widetilde{N}(dt, d\nu), \\ X_0^n = R_n x, \quad t \ge 0 \end{cases}
$$

and

$$
\begin{cases} dY_t^n = AY_t^n + R_n F(Y_t^n)dt + R_n \sigma(Y_t^n)dW_t + \int_E R_n \gamma(Y_t^n, \nu) \widetilde{N}(dt, d\nu), \\ Y_0^n = R_n y, \quad t \ge 0 \end{cases}
$$

where $R_n = n(n - A)^{-1}$ for $n \in \mathbb{N}$ with $n > \alpha + \beta + \sqrt{L_F} =: \lambda$. By [\(2.4\)](#page-3-2) we find for each $n \geq 1 + \lambda$ the inequality

$$
||R_nz||_H \leq \frac{n}{n-\lambda}||z||_H \leq (1+\lambda)||z||_H.
$$

By classical properties of the resolvent (see [\[29,](#page-25-18) Lemma 3.2]), one clearly has $R_n z \to z$ as $n \to \infty$ in H. Moreover, by properties of the Yosida approximation of mild solutions of SPDEs (compare e.g. with Appendix A2 in [\[26\]](#page-25-16) or Section 2 in [\[2\]](#page-24-3)) we have

$$
\lim_{n \to \infty} \mathbb{E} \left[\sup_{t \in [0,T]} \|X_t^n - X_t\|_H^2 + \sup_{t \in [0,T]} \|Y_t^n - Y_t\|_H^2 \right] = 0, \qquad \forall T > 0
$$

and hence there exists a subsequence (which is again denoted by n) such that $X_t^n \longrightarrow X_t$ and $Y_t^n \longrightarrow Y_t$ hold a.s. for each $t \geq 0$. Following a method proposed in [\[2\]](#page-24-3) we verify that sufficient conditions are satisfied to apply the generalized Itô-formula from Theorem [A.2](#page-23-0) to the function $F(t, z) := e^{\varepsilon t} ||z||_H^2$, where $\varepsilon = 2\alpha - L_{\sigma} - L_{\gamma}$ is given by [\(3.2\)](#page-6-0):

$$
X_t^n - Y_t^n = R_n(x - y) + \int_0^t \{A(X_s^n - Y_s^n) + R_n(F(X_s^n) - F(Y_s^n))\} ds
$$

+
$$
\int_0^t R_n(\sigma(X_s^n) - \sigma(Y_s^n)) dW_s + \int_0^t \int_E R_n(\gamma(X_s^n, \nu) - \gamma(Y_s^n, \nu)) \widetilde{N}(ds, d\nu).
$$

Observe that, by condition (A1) and [\(3.3\)](#page-6-1), one has

$$
\int_{0}^{t} \int_{E} ||R_{n}(\gamma(X_{s}^{n}, \nu) - \gamma(Y_{s}^{n}, \nu))||_{H}^{2} \mu(d\nu) ds
$$

+
$$
\int_{0}^{t} \int_{E} ||R_{n}(\gamma(X_{s}^{n}, \nu) - \gamma(Y_{s}^{n}, \nu))||_{H}^{4} \mu(d\nu) ds
$$

$$
\leq (1 + \lambda)^{2} \int_{0}^{t} \int_{E} ||\gamma(X_{s}^{n}, \nu) - \gamma(Y_{s}^{n}, \nu)||_{H}^{2} \mu(d\nu) ds
$$

+
$$
8(1 + \lambda)^{4} \int_{0}^{t} \int_{E} (||\gamma(X_{s}^{n}, \nu)||_{H}^{4} + ||\gamma(Y_{s}^{n}, \nu)||_{H}^{4}) \mu(d\nu) ds
$$

$$
\leq L_{\gamma}(1 + \lambda)^{2} \int_{0}^{t} ||X_{s}^{n} - Y_{s}^{n}||_{H}^{2} ds
$$

+
$$
16(1 + \lambda)^{4} t \sup_{z \in H} \int_{E} ||\gamma(z, \nu)||_{H}^{4} \mu(d\nu) < \infty.
$$

Thus we can apply the generalized Itô-formula from Theorem $A.2$ and obtain (similar) to (3.5) in [\[2\]](#page-24-3))

$$
e^{\varepsilon t} \|X_t^n - Y_t^n\|_H^2 - \|R_n(x - y)\|_H^2
$$

= $\int_0^t \langle 2e^{\varepsilon s}(X_s^n - Y_t^n), R_n(\sigma(X_s^n) - \sigma(Y_s^n))dW_s \rangle_H$
+ $\int_0^t e^{\varepsilon s} [\varepsilon \|X_s^n - Y_s^n\|_H^2 + \mathcal{L}_n(\|\cdot\|_H^2)(X_s^n, Y_s^n)] ds$
+ $\int_0^t \int_E e^{\varepsilon s} [\|X_s^n - Y_s^n + R_n(\gamma(X_s^n, \nu) - \gamma(Y_s^n, \nu))\|_H^2 - \|X_s^n - Y_s^n\|_H^2] \widetilde{N}(ds, d\nu),$
(3.5)

where we used, for $z, w \in D(A)$, the notation

$$
\mathcal{L}_n(\|\cdot\|_H^2)(z,w) := 2\langle z - w, A(z - w) + R_n(F(z) - F(w)) \rangle_H + \|R_n(\sigma(z) - \sigma(w))\|_{L_2^0(H)}^2 + \int_E \|R_n(\gamma(z, \nu) - \gamma(w, \nu))\|_H^2 \mu(d\nu).
$$

Taking expectations in [\(3.5\)](#page-8-0) yields

$$
e^{\varepsilon t} \mathbb{E} \left[\|X_t^n - Y_t^n\|_H^2 \right] - \mathbb{E} \left[\|R_n(x - y)\|_H^2 \right]
$$

=
$$
\mathbb{E} \left[\int_0^t e^{\varepsilon s} \left(\varepsilon \|X_s^n - Y_s^n\|_H^2 + \mathcal{L}_n (\|\cdot\|_H^2) (X_s^n, Y_s^n) \right) ds \right].
$$
 (3.6)

Lemma [3.1](#page-6-2) yields

$$
e^{\varepsilon t} \mathbb{E} \left[\|X_t^n - Y_t^n\|_H^2 \right] - \mathbb{E} \left[\|R_n(x - y)\|_H^2 \right] - 2(\alpha + \beta) \int_0^t e^{\varepsilon s} \mathbb{E} \left[\|P_1 X_s^n - P_1 Y_s^n\|_H^2 \right] ds
$$

$$
\leq \mathbb{E} \left[\int_0^t e^{\varepsilon s} (-\mathcal{L}(\|\cdot\|_H^2)(X_s^n, Y_s^n) + \mathcal{L}_n(\|\cdot\|_H^2)(X_s^n, Y_s^n)) ds \right].
$$

Below we prove that the right-hand-side tends to zero as $n \to \infty$, which would imply the assertion of this theorem. To prove the desired convergence to zero we apply the generalized Lebesgue Theorem (see [\[26,](#page-25-16) Theorem 7.1.8]). For this reason we have to prove that

$$
\mathcal{L}(\|\cdot\|_{H}^{2})(X_{s}^{n}, Y_{s}^{n}) - \mathcal{L}_{n}(\|\cdot\|_{H}^{2})(X_{s}^{n}, Y_{s}^{n}) \to 0
$$
\n(3.7)

holds a.s. for each $s > 0$ as $n \to \infty$ and, moreover, there exists a constant $C > 0$ such that

$$
|\mathcal{L}(\|\cdot\|_H^2)(X_s^n, Y_s^n) - \mathcal{L}_n(\|\cdot\|_H^2)(X_s^n, Y_s^n)| \le C \|X_s^n - Y_s^n\|_H^2.
$$
 (3.8)

We start with the proof of [\(3.7\)](#page-8-1). Denote $F_s^n := F(X_s^n) - F(Y_s^n)$, $\sigma_s^n := \sigma(X_s^n) - \sigma(Y_s^n)$ and $\gamma_s^n(\nu) := \gamma(X_s^n, \nu) - \gamma(Y_s^n, \nu)$ and analogously $F_s := F(X_s) - F(Y_s)$, $\sigma_s := \sigma(X_s) - \sigma(X_s)$ $\sigma(Y_s)$ and $\gamma_s(\nu) := \gamma(X_s, \nu) - \gamma(Y_s, \nu)$ for each $n \in \mathbb{N}, s \geq 0$ and $\nu \in E$. Then

$$
\begin{aligned} &\left| \left(\mathcal{L}(\|\cdot\|_H^2)(X_s^n, Y_s^n) - \mathcal{L}_n(\|\cdot\|_H^2)(X_s^n, Y_s^n)) \right| \right. \\ &\left. \leq 2|\langle X_s^n - Y_s^n, F_s^n - R_n F_s^n \rangle_H| + ||\sigma_s^n||_{L_2^0}^2 - ||R_n \sigma_s^n||_{L_2^0}^2 \right| \\ &\left. + \left| \int_E ||\gamma_s^n(\nu)||_H^2 - ||R_n \gamma_s^n(\nu)||_H^2 \mu(d\nu) \right| \right| \\ &\qquad=: I_1 + I_2 + I_3. \end{aligned}
$$

For the first term I_1 we estimate

 $I_1 \leq 2||X_s^n - Y_s^n||_H ||F_s^n - R_n F_s^n||_H$

10 BÁLINT FARKAS, MARTIN FRIESEN, BARBARA RÜDIGER, AND DENNIS SCHROERS

$$
\leq 2\|X_s^n - Y_s^n\|_H \left(\|F_s^n - F_s\|_H + \|F_s - R_n F_s\|_H + \|R_n F_s - R_n F_s^n\|_H \right) \leq 2\|X_s^n - Y_s^n\|_H \left(\|F_s^n - F_s\|_H + \|F_s - R_n F_s\|_H + (1 + \lambda) \|F_s - F_s^n\|_H \right).
$$

Using that $X_s^n \to X_s$ and $Y_s^n \to Y_s$ as a.s. for some subsequence (also denoted by n), we easily find that the right-hand side tends to zero. The convergence of the second term follows from

$$
I_2 = \left| \|\sigma_s^n\|_{L_2^0} - \|R_n \sigma_s^n\|_{L_2^0} \right| \left(\|\sigma_s^n\|_{L_2^0} + \|R_n \sigma_s^n\|_{L_2^0} \right)
$$

\n
$$
\leq (2 + \lambda) \sqrt{L_\sigma} \|\sigma_s^n - R_n \sigma_s^n\|_{L_2^0} \|X_s^n - Y_s^n\|_H
$$

\n
$$
\leq (2 + \lambda)^2 \sqrt{L_\sigma} \|X_s^n - Y_s^n\|_H \left(\|\sigma_s^n - \sigma_s\|_{L_2^0} + \|\sigma_s - R_n \sigma_s\|_{L_2^0} + \|\sigma_s - \sigma_s^n\|_{L_2^0} \right).
$$

It remains to show the convergence of the third term. First, observe

$$
I_3 \le (2 + \lambda) \int_E ||\gamma_s^n(\nu) - R_n \gamma_s^n(\nu) ||_H ||\gamma_s^n(\nu) ||_H \mu(d\nu)
$$

\n
$$
\le (2 + \lambda) \int_E \left(||\gamma_s^n(\nu) - \gamma_s(\nu) ||_H + ||\gamma_s(\nu) - R_n \gamma_s(\nu) ||_H \right) ||\gamma_s^n(\nu) ||_H
$$

\n
$$
+ ||R_n \gamma_s(\nu) - R_n \gamma_s^n(\nu) ||_H \right) ||\gamma_s^n(\nu) ||_H \mu(d\nu)
$$

\n
$$
\le (2 + \lambda) \left(\int_E ||\gamma_s^n(\nu) ||_H^2 \mu(d\nu) \right)^{\frac{1}{2}} \left[\left(\int_E ||\gamma_s^n(\nu) - \gamma_s(\nu) ||_H^2 \mu(d\nu) \right)^{\frac{1}{2}} \right]
$$

\n
$$
+ \left(\int_E ||\gamma_s(\nu) - R_n \gamma_s(\nu) ||_H^2 \mu(d\nu) \right)^{\frac{1}{2}} + \left(\int_E ||R_n \gamma_s(\nu) - R_n \gamma_s^n(\nu) ||_H^2 \mu(d\nu) \right)^{\frac{1}{2}} \right]
$$

\n
$$
\le \sqrt{2} (2 + \lambda)^2 L_\gamma ||X_s^n - Y_s^n||_H (||X_s^n - X_s||_H + ||Y_s^n - Y_s||_H)
$$

\n
$$
+ (2 + \lambda) \sqrt{L_\gamma} ||X_s^n - Y_s^n||_H \left(\int_E ||\gamma_s(\nu) - R_n \gamma_s(\nu) ||_H^2 \mu(d\nu) \right)^{\frac{1}{2}}
$$

\n
$$
= I_3^1 + I_3^2
$$

where the last inequality follows from condition $(A1)$ combined with the inequality

$$
||R_n \gamma_s(\nu) - R_n \gamma_s^n(\nu)||_H^2
$$

\n
$$
\leq (1 + \lambda)^2 ||\gamma_s(\nu) - \gamma_s^n(\nu)||_H^2
$$

\n
$$
\leq 2(1 + \lambda)^2 (||\gamma(X_s, \nu) - \gamma(Y_s, \nu)||_H^2 + ||\gamma(X_s^n, \nu) - \gamma(Y_s^n, \nu)||_H^2).
$$

The first expression I_1^1 clearly tends to zero as $n \to \infty$. For the second expression I_3^2 we use the inequality $\|\gamma_s(\nu) - R_n \gamma_s(\nu)\|_H^2 \leq 2(2+\lambda)^2 \|\gamma_s(\nu)\|_H^2$ so that dominated convergence theorem is applicable, which shows that $I_3^2 \to 0$ as $n \to \infty$ a.s.. This proves [\(3.7\)](#page-8-1). Concerning [\(3.8\)](#page-8-2), we find that

$$
\begin{split}\n&\left| \left(\mathcal{L}(\|\cdot\|_{H}^{2}) (X_{s}^{n}, Y_{s}^{n}) - \mathcal{L}_{n}(\|\cdot\|_{H}^{2}) (X_{s}^{n}, Y_{s}^{n}) \right) \right| \\
&\leq 2|\langle X_{s}^{n} - Y_{s}^{n}, F_{s}^{n} - R_{n}F_{s}^{n}\rangle_{H}| + |||\sigma_{s}^{n}||_{L_{2}^{0}(H)}^{2} - ||R_{n}\sigma_{s}^{n}||_{L_{2}^{0}(H)}^{2} \\
&+ \left| \int_{E} ||\gamma_{s}^{n}(\nu)||_{H}^{2} - ||R_{n}\gamma_{s}^{n}(\nu)||_{H}^{2} \mu(d\nu) \right| \\
&\leq 2(2 + \lambda) \|X_{s}^{n} - Y_{s}^{n}||_{H} \|F_{s}^{n}||_{H} + (1 + (1 + \lambda)^{2}) \left[\|\sigma_{s}^{n}||_{L_{2}^{0}(H)}^{2} + \int_{E} ||\gamma_{s}^{n}(\nu)||_{H}^{2} \mu(d\nu) \right] \\
&\leq 2(2 + \lambda)L_{F} \|X_{s}^{n} - Y_{s}^{n}||_{H}^{2} + (1 + (1 + \lambda)^{2}) \left(L_{\sigma} + L_{\gamma} \right) \|X_{s}^{n} - Y_{s}^{n}||_{H}^{2}.\n\end{split}
$$

Hence the generalized Lebesgue Theorem is applicable, and thus the assertion of this theorem is proved. \Box

Note that condition (3.3) is used to guarantee that the Itô-formula [A.2](#page-23-0) for Hilbert space valued jump diffusions can be applied for $(x,t) \to e^{t\varepsilon} ||x||_H^2$. The assertion of Proposition [3.2](#page-6-3) is also true when $\varepsilon \leq 0$, but will be only applied for the case when $\varepsilon > 0$.

4. Convergence to limiting distribution

4.1. The Case of Vanishing Coefficients. It follows from Proposition [3.2](#page-6-3) that, for $\varepsilon > 0$, one has an estimate on the L²-norm of the difference $X_t^x - X_t^y$ t^y . Such an estimate alone does neither imply the existence nor uniqueness of an invariant distribution. However, if the coefficients F, σ, γ vanish at H_1 , then we may characterize the limiting distributions in L^2 .

Theorem 4.1. *Suppose that (GDC) holds with a projection operator* P_1 , $(A1)$, (3.3) *,* [\(3.2\)](#page-6-0) are satisfied, that $(S(t))_{t>0}$ leaves $H_0 := \text{ran}(I - P_1)$ invariant, and that $\text{ran}(P_1) \subset$ ker(A)*. Moreover, assume that*

$$
P_1 F \equiv 0, \ P_1 \sigma \equiv 0, \ P_1 \gamma \equiv 0. \tag{4.1}
$$

Fix $x \in L^2(\Omega, \mathcal{F}_0, \mathbb{P}; H)$ *and suppose that*

$$
F(P_1x) = 0, \ \sigma(P_1x) = 0, \ \gamma(P_1x, \cdot) = 0 \tag{4.2}
$$

holds for this fixed choice of x*. Then*

$$
\mathbb{E}\left[\|X_t^x - P_1x\|_H^2\right] \le e^{-\varepsilon t} \mathbb{E}\left[\|(1 - P_1)x\|_H^2\right].
$$

In particular, let ρ *be the law of* $x \in L^2(\Omega, \mathcal{F}_0, \mathbb{P}; H)$ *and* ρ_1 *be the law of* P_1x *, respectively. Then* ρ_1 *is an invariant measure.*

Proof. Fix $x \in L^2(\Omega, \mathcal{F}_0, \mathbb{P}; H)$ with property (4.2) and set $P_0 = I - P_1$. Since $(S(t))_{t \geq 0}$ leaves H_0 invariant we find that $S(t)P_0 = P_0S(t)$ and hence we obtain $P_1S(t)P_0 =$ $P_1P_0S(t) = 0$. Moreover, using [\(4.1\)](#page-10-1) we find that

$$
P_1 X_t^x = P_1 S(t)x = P_1 S(t) P_0 x + P_1 S(t) P_1 x = P_1 x
$$

where we have that $S(t)P_1 = P_1$ due to $ran(P_1) \subset ker(A)$. From this we conclude that $(P_0 X_t^x)_{t \geq 0}$ satisfies

$$
P_0 X_t^x = P_0 S(t)x + \int_0^t P_0 S(t-s) F(X_s^x) ds + \int_0^t P_0 S(t-s) \sigma(X_s^x) dW_s
$$

+
$$
\int_0^t \int_E P_0 S(t-s) \gamma(X_s^x) \tilde{N}(ds, d\nu)
$$

=
$$
S(t) P_0 x + \int_0^t S(t-s) P_0 F(P_1 x + P_0 X_t^x) ds + \int_0^t S(t-s) P_0 \sigma(P_1 x + P_0 X_s^x) dW_x
$$

+
$$
\int_0^t \int_E S(t-s) P_0 \gamma(P_1 x + P_0 X_s^x) \tilde{N}(ds, d\nu)
$$

=
$$
S(t) P_0 x + \int_0^t S(t-s) \tilde{F}(P_0 X_t^x) ds + \int_0^t S(t-s) \tilde{\sigma}(P_0 X_s^x) dW_s
$$

$$
\int_0^t \int_E S(t-s) \tilde{\gamma}(P_0 X_s^x) \tilde{N}(ds, d\nu),
$$

where we have set $\widetilde{F}(y) := P_0 F(P_1 x + y), \ \widetilde{\sigma}(y) := P_0 \sigma(P_1 x + y)$ and $\widetilde{\gamma}(y, \nu) := P_0 \gamma(P_1 x + y)$ y, ν for all $y \in H_0$ and $\nu \in E$. Since these coefficients share the same Lipschitz estimates as F, σ and γ , we can apply Proposition [3.2](#page-6-3) to the process $(P_0 X_t^x)_{t\geq0}$ obtained from the above auxiliary SPDE, we obtain

$$
\mathbb{E}[\|X_t^x - P_1x\|_H^2] = \mathbb{E}[\|P_0X_t^x\|_H^2] = \mathbb{E}[\|P_0X_t^x - P_0X_t^0\|_H^2] \le e^{-\varepsilon t}\mathbb{E}[\|P_0x\|_H^2],
$$

where we have used that $P_0X_t^0 = 0$ due to [\(4.2\)](#page-10-0).

This theorem can be applied, for instance, to the Heath-Jarrow-Morton-Musiela equation, see Section 6. Below we discuss two simple examples showing that, in general, conditions $ran(P_1) \subset ker(A)$ and (4.1) cannot be omitted.

Example 4.2. Let $H = \mathbb{R}^2$ and $(W_t)_{t \geq 0}$ be a 2-dimensional standard Brownian motion. (a) Let $X_t = (X_t^1, X_t^2) \in H = \mathbb{R}^2$ be given by

$$
dX_t = \begin{pmatrix} -1 & 0 \\ 0 & 0 \end{pmatrix} X_t dt + \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} dW_t.
$$

Then (GDC) holds for $P_1 =$ $\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ *, and* $H_1 = \text{ran}(P_1) = \{0\} \times \mathbb{R} = \text{ker}(A)$ *. The semigroup is given by* $S(t) = \begin{pmatrix} e^{-t} & 0 \\ 0 & 1 \end{pmatrix}$ and leaves $H_0 = \text{ran}(I - P_1) =$ $\mathbb{R} \times \{0\}$ *invariant. Finally we have* $\varepsilon = 2 > 0$ *and* [\(4.1\)](#page-10-1)*, while while condition* (4.1) *is not satisfied. By* $X_t^2 = W_t^2$, *it is clear that* X_t^2 *does not have a limiting* $distribution. Hence also X_t cannot have a limiting distribution.$

(b) Let $Y_t = (Y_t^1, Y_t^2) \in H = \mathbb{R}^2$ be the solution of

$$
dY_t = \begin{pmatrix} -1 & 1 \\ 0 & 1 \end{pmatrix} Y_t dt + \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} dW_t.
$$

Then (GDC) holds with P¹ *being the projection onto the second coordinate. Hence* [\(4.1\)](#page-10-1) *holds, while* $\text{ran}(P_1) \subset \text{ker}(A)$ *is not satisfied. Since* $Y_t^2 = e^t Y_0^2 +$ $\int_0^t e^{t-s} dW_s^2$ it is clear that Y_t^2 does not have a limiting distribution. Hence also Y^t *cannot have a limiting distribution.*

4.2. The General Case. In Theorem [4.1](#page-10-2) we have assumed (4.1) , (4.2) , and that $(S(t))_{t>0}$ leaves H_0 invariant. Below we continue with the more general case. Namely, for the projection operator P_1 given by condition (GDC) we decompose the process X_t^x according to $X_t^x = P_0 X_t^x + P_1 X_t^x$, with $P_0 = I - P_1$ and suppose that:

(A2) The process $P_1 X_t^x$ is deterministic of the form

$$
P_1 X_t^x = P_1 S(t) P_1 x + \int_0^t P_1 S(t - s) P_1 F(P_1 X_t^x) ds.
$$
\n(4.3)

Our next condition imposes a control on this solution:

(A3) For each $x \in H_1 = \text{ran}(P_1)$ there exists $\tilde{X}^x_\infty \in H_1$ and constants $C(x) > 0$, $\delta(x) \in (0, |\varepsilon|)$ such that

$$
||P_1 X_t^x - \widetilde{X}_{\infty}^x||_H^2 \le C(x) e^{-\delta(x)t}, \qquad t \ge 0.
$$

Note that, if $P_1F(P_1) = 0$ then condition (A3) reduces to a condition on the limiting behavior of the semigroup $(S(t))_{t>0}$ when restricted to $H_1 = \text{ran}(P_1)$. In such a case condition (A3) is, for instance, satisfied if $H_1 \subset \text{ker}(A)$. The following is our main result for this case.

Theorem 4.3. *Suppose that condition (GDC) holds for some projection operator* P1*, that conditions (A1) – (A3),* [\(3.3\)](#page-6-1) *and* [\(3.2\)](#page-6-0) *are satisfied. Then the following assertions hold:*

(a) For each $x \in H$ there exists an invariant measure $\pi_{\delta_x} \in \mathcal{P}_2(H)$ for the Markov *semigroup* $(p_t)_{t>0}$ *and a constant* $K(\alpha, \beta, \varepsilon, h) > 0$ *such that*

$$
\mathrm{W}_2(p_t(x,\cdot), \pi_{\delta_x}) \le K(\alpha, \beta, \varepsilon, x) e^{-\frac{\delta(x)}{2}t}, \qquad t \ge 0.
$$

(b) Suppose, in addition to the conditions of $(A3)$, that there are constants δ *and* C*, such that*

$$
\delta(x) \ge \delta > 0 \quad and \quad C(x) \le C(1 + ||x||_H)^4, \qquad x \in H. \tag{4.4}
$$

Then, for each $\rho \in \mathcal{P}_2(H)$ *, there exists an invariant measure* $\pi_{\rho} \in \mathcal{P}_2(H)$ *for the Markov semigroup* $(p_t)_{t>0}$ *and a constant* $K(\alpha, \beta, \varepsilon) > 0$ *such that*

$$
\mathrm{W}_2(p_t^*\rho, \pi_\rho) \le K(\alpha, \beta, \varepsilon) \int_H (1 + \|x\|_H)^2 \rho(dx) e^{-\frac{\delta}{2}t}, \qquad t \ge 0.
$$

The proof of this theorem relies on the key stability estimate formulated in Proposition [3.2](#page-6-3) and is given at the end of this section. So far we have only shown the existence of invariant measures parametrized by the initial state of the process. However, under the given conditions it can also be shown that π_{δ_x} as well as π_ρ depend only on the H_1 part of x or ρ , respectively.

Corollary 4.4. *Suppose that condition (GDC) holds for some projection operator* P_1 , *that conditions (A1) – (A3),* [\(3.3\)](#page-6-1) *and* [\(3.2\)](#page-6-0) *are satisfied. Then the following assertions hold:*

- (a) Let $x, y \in H$ be such that $P_1x = P_1y$. Then $\pi_{\delta_x} = \pi_{\delta_y}$.
- *(b) Suppose, in addition, that* [\(4.4\)](#page-12-0) *holds.* Let $\rho, \tilde{\rho} \in \mathcal{P}_2(H)$ *be such that* $\rho \circ P_1^{-1} = \tilde{\rho} \circ \tilde{\rho} \circ \tilde{\rho}^{-1}$ $\widetilde{\rho} \circ P_1^{-1}$. Then $\pi_{\rho} = \pi_{\widetilde{\rho}}$.

Next we turn to a proof of Theorem [4.3](#page-12-1) and Corollary [4.4.](#page-12-2)

4.3. Construction of a coupling. Let $x \in H$ and let $(X_t^x)_{t \geq 0}$ be the unique mild so-lution to [\(2.7\)](#page-3-3). Below we construct for given $\tau \geq 0$ a coupling for the law of $(X_t^x, X_{t+\tau}^x)$. Let $(Y_t^{x,\tau})$ $(t^{x,\tau}_{t})_{t\geq 0}$ be the unique mild solution to the SPDE

$$
Y_t^{x,\tau} = S(t)x + \int_0^t S(t-s)F(Y_s^{x,\tau})ds + \int_0^t S(t-s)\sigma(Y_s^{x,\tau})dW_s^{\tau} + \int_0^t \int_E S(t-s)\gamma(Y_s^{x,\tau}, \nu)\tilde{N}^{\tau}(ds, d\nu), \qquad t \ge 0,
$$
\n(4.5)

where $W_s^{\tau} = W_{\tau+s} - W_{\tau}$, $\widetilde{N}^{\tau}(ds, d\nu) = \widetilde{N}(ds + \tau, d\nu) - \widetilde{N}(ds, d\nu)$ is a Q-Wiener process and a Poisson random measure with respect to the filtration $(\mathcal{F}_s^{\tau})_{s\geq 0}$ defined by \mathcal{F}_s^{τ} = $\mathcal{F}_{s+\tau}$.

Lemma 4.5. *Suppose that (GDC), (A1),* [\(3.3\)](#page-6-1) *and* [\(3.2\)](#page-6-0) *are satisfied. Then for each* $x \in H$ *and* $t, \tau \geq 0$ *the following assertions hold:*

 (a) $Y_t^{x,\tau}$ $t^{x,\tau}$ has the same law as X_t^x .

(b) It holds that

$$
\mathbb{E} [||Y_t^{x,\tau} - X_{t+\tau}^x||_H^2] \le e^{-\varepsilon t} \mathbb{E} [||x - X_\tau^x||_H^2] + 2(\alpha + \beta) \int_0^t e^{-\varepsilon(t-s)} \mathbb{E} [||P_1 Y_s^{x,\tau} - P_1 X_{s+\tau}^x||_H^2] ds.
$$

Proof. (a) Since [\(2.7\)](#page-3-3) has a unique solution it follows from the Yamada-Watanabe Theorem (see [\[25\]](#page-25-19)) that also uniqueness in law holds for this equation. Since the driving noises N^{τ} and W^{τ} in [\(4.5\)](#page-12-3) have the same law as N and W from [\(2.7\)](#page-3-3), it follows that the unique solution to (4.5) has the same law as the solution to (2.7) . This proves the assertion.

(b) Set $X_t^{x,\tau}$ $t^{x,\tau} := X^x_{t+\tau}$, then by direct computation we find that

$$
X_{t}^{x,\tau} = S(t)S(\tau)x + \int_{0}^{t+\tau} S(t+\tau-s)F(X_{s}^{x})ds + \int_{0}^{t+\tau} S(t+\tau-s)\sigma(X_{s}^{x})dW_{s}
$$

+
$$
\int_{0}^{t+\tau} \int_{E} S(t+\tau-s)\gamma(X_{s}^{x}, \nu)\tilde{N}(ds, d\nu)
$$

=
$$
S(t)S(\tau)x + S(t)\int_{0}^{\tau} S(\tau-s)F(X_{s}^{x})ds + S(t)\int_{0}^{\tau} S(\tau-s)\sigma(X_{s}^{x})dW_{s}
$$

+
$$
S(t)\int_{0}^{\tau} \int_{E} S(\tau-s)\gamma(X_{s}^{x}, \nu)\tilde{N}(ds, d\nu)
$$

+
$$
\int_{\tau}^{t+\tau} S(t+\tau-s)F(X_{s}^{x})ds + \int_{\tau}^{t+\tau} S(t+\tau-s)\sigma(X_{s}^{x})dW_{s}
$$

+
$$
\int_{\tau}^{t+\tau} \int_{E} S(t+\tau-s)\gamma(X_{s}^{x}, \nu)\tilde{N}(ds, d\nu)
$$

=
$$
S(t)X_{0}^{x,\tau} + \int_{0}^{t} S(t-s)F(X_{s}^{x,\tau})ds + \int_{0}^{t} S(t-s)\sigma(X_{s}^{x,\tau})dW_{s}^{\tau}
$$

+
$$
\int_{0}^{t} \int_{E} S(t-s)\gamma(X_{s}^{x,\tau}, \nu)\tilde{N}^{\tau}(ds, d\nu),
$$

where in the last equality we have used, for appropriate integrands $\Phi(s, \nu)$ and $\Psi(s)$, that

$$
\int_{\tau}^{\tau+t} \Psi(s)dW_s = \int_0^t \Psi(s+\tau)dW_s^{\tau},
$$

$$
\int_{\tau}^{\tau+t} \int_E \Phi(s,\nu)\widetilde{N}(ds,d\nu) = \int_0^t \int_E \Phi(s+\tau,\nu)\widetilde{N}^{\tau}(ds,d\nu).
$$

Hence $(X_t^{x,\tau})$ $\mathcal{F}_t^{\pi,\tau}$)_t also solves [\(4.5\)](#page-12-3) with $\mathcal{F}_0^{\tau} = \mathcal{F}_{\tau}$ and initial condition $X_0^{\pi,\tau} = X_{\tau}^{\pi}$. Consequently, the assertion follows from Proposition [3.2](#page-6-3) applied to $X_t^{x,\tau}$ $t^{x,\tau}$ and $Y_t^{x,\tau}$ $\overline{t}^{x,\tau}$. \Box

4.4. Proof of Theorem [4.3.](#page-12-1)

Proof of Theorem [4.3.](#page-12-1) Fix $x \in H$ and recall that $p_t(x, \cdot)$ denotes the transition proba-bilities of the Markov process obtained from [\(2.7\)](#page-3-3). Below we prove that $(p_t(x, \cdot))_{t\geq 0}$ $\mathcal{P}_2(H)$ is a Cauchy sequence with respect to the Wasserstein distance W₂. Fix $t, \tau \geq 0$. We treat the cases $\tau \in (0, 1]$ and $\tau > 1$ separately.

Case $0 < \tau \leq 1$: Then using the coupling lemma [4.5.](#page-12-4)(b) yields

$$
W_2(p_{t+\tau}(x,\cdot), p_t(x,\cdot)) \leq \left(\mathbb{E}\left[\|Y_t^{x,\tau} - X_{t+\tau}^x\|_H^2\right]\right)^{1/2} \n\leq e^{-\frac{\varepsilon}{2}t} \left(\mathbb{E}\left[\|X_{\tau}^x - x\|_H^2\right]\right)^{1/2} \n+ \sqrt{2(\alpha + \beta)} \left(\int_0^t e^{-\varepsilon(t-s)} \mathbb{E}\left[\|P_1 Y_s^{x,\tau} - P_1 X_{s+\tau}^x\|_H^2\right] ds\right)^{1/2} \n=: I_1 + I_2.
$$

The first term I_1 can be estimated by

$$
I_1 \le e^{-\frac{\varepsilon}{2}t} \sup_{s \in [0,1]} \left(\mathbb{E} \left[\|X_s^x - x\|_H^2 \right] \right)^{1/2}.
$$

To estimate the second term I_2 we first observe that by condition (A2) we have $P_1 Y_s^{x,\tau} =$ $P_1 X_s^x$ and hence by condition (A3) one has for each $s \geq 0$ that

$$
\mathbb{E} \left[\|P_1 Y_s^{x,\tau} - P_1 X_{s+\tau}^x\|_H^2 \right] \le 2 \|P_1 Y_s^{x,\tau} - \tilde{X}_{\infty}^x\|_H^2 + 2 \|P_1 X_{s+\tau}^x - \tilde{X}_{\infty}^x\|_H^2
$$

$$
\le 4C(x) e^{-\delta(x)s}.
$$

This readily yields

$$
\int_0^t e^{-\varepsilon(t-s)} \mathbb{E} \left[\|P_1 Y_s^{x,\tau} - P_1 X_{s+\tau}^x\|_H^2 \right] ds
$$

\n
$$
\leq 4C(x) \int_0^t e^{-\varepsilon(t-s)} e^{-\delta(x)s} ds
$$

\n
$$
= 4C(x) e^{-\varepsilon t} \frac{e^{(\varepsilon-\delta(x))t} - 1}{\varepsilon - \delta(x)}
$$

\n
$$
\leq 4C(x) \frac{e^{-\delta(x)t}}{\varepsilon - \delta(x)}.
$$

Inserting this into the definition of I_2 gives

$$
I_2 \le 2\sqrt{\frac{(\alpha+\beta)C(x)}{\varepsilon-\delta(x)}}e^{-\frac{\delta(x)}{2}t}.
$$

Case $\tau > 1$: Fix some $N \in \mathbb{N}$ with $\tau < N < 2\tau$ and define a sequence of numbers $(a_n)_{n=0,...,N}$ by

$$
a_n := \frac{\tau}{N} n, \qquad n = 0, \dots, N.
$$

Then $a_0 = 0$, $a_N = \tau$ and $a_n - a_{n-1} = \frac{\tau}{N} =: \varkappa \in (\frac{1}{2})$ $(\frac{1}{2}, 1)$ for $n = 1, \ldots, N$. Hence we obtain from the coupling Lemma [4.5.](#page-12-4)(b)

$$
W_{2}(p_{t+\tau}(x,\cdot), p_{t}(x,\cdot))
$$

\n
$$
\leq \sum_{n=1}^{N} W_{2}(p_{t+a_{n}}(x,\cdot), p_{t+a_{n-1}}(x,\cdot))
$$

\n
$$
\leq \sum_{n=1}^{N} \left(\mathbb{E} \left[\left\| Y_{t+a_{n-1}}^{x,x} - X_{t+a_{n-1}+\varkappa}^{x} \right\|_{H}^{2} \right] \right)^{1/2}
$$

\n
$$
\leq \sum_{n=1}^{N} e^{-\frac{\varepsilon}{2}(t+a_{n-1})} \left(\mathbb{E} \left[\left\| X_{\varkappa}^{x} - x \right\|_{H}^{2} \right] \right)^{1/2}
$$

\n
$$
+ \sqrt{2(\alpha + \beta)} \sum_{n=1}^{N} \left(\int_{0}^{t+a_{n-1}} e^{-\varepsilon(t+a_{n-1}-s)} \mathbb{E} \left[\left\| P_{1} Y_{s}^{x,x} - P_{1} X_{s+\varkappa}^{x} \right\|_{H}^{2} \right] ds \right)^{1/2}
$$

\n
$$
=: I_{1} + I_{2}.
$$

For the first term I_1 we use $\varkappa > \frac{1}{2}$ $\frac{1}{2}$ so that

$$
\sum_{n=1}^{N} e^{-\frac{\varepsilon}{2}\varkappa(n-1)} \le \sum_{n=0}^{\infty} e^{-\frac{\varepsilon}{4}n} = \left(1 - e^{-\frac{\varepsilon}{4}}\right)^{-1},
$$

from which we obtain

$$
I_1 = e^{-\frac{\varepsilon}{2}t} \sup_{s \in [0,1]} \left(\mathbb{E}[\|X_s^x - x\|_H^2] \right)^{\frac{1}{2}} \sum_{n=1}^N e^{-\frac{\varepsilon}{2}t(n-1)}
$$

$$
\leq \sup_{s \in [0,1]} \left(\mathbb{E}[\|X_s^x - x\|_H^2] \right)^{\frac{1}{2}} \left(1 - e^{-\frac{\varepsilon}{4}} \right)^{-1} e^{-\frac{\varepsilon}{2}t}.
$$

To estimate the second term I_2 we first observe that by condition (A2) we have $P_1 Y_s^{x,\tau} =$ $P_1 X_s^x$ and hence by condition (A3), one has for $s \geq 0$

$$
\mathbb{E} [||P_1 Y_s^{x,\varkappa} - P_1 X_{s+\varkappa}^x||_H^2] \le 2||P_1 Y_s^{x,\varkappa} - \tilde{X}_{\infty}^x||_H^2 + 2||P_1 X_{s+\varkappa}^x - \tilde{X}_{\infty}^x||_H^2
$$

$$
\le 4C(x)e^{-\delta(x)s}.
$$

Hence we find that

$$
\int_0^{t+a_{n-1}} e^{-\varepsilon(t+a_{n-1}-s)} \mathbb{E} \left[\|P_1 Y_s^{x,\varkappa} - P_1 X_{s+\varkappa}^x\|_H^2 \right] ds
$$

\n
$$
\leq 4C(x) \int_0^{t+a_{n-1}} e^{-\varepsilon(t+a_{n-1}-s)} e^{-\delta(x)s} ds
$$

\n
$$
= 4C(x) e^{-\varepsilon(t+a_{n-1})} \frac{e^{(\varepsilon-\delta(x))(t+a_{n-1})} - 1}{\varepsilon - \delta(x)}
$$

\n
$$
\leq 4C(x) \frac{e^{-\delta(x)t}}{\varepsilon - \delta(x)} e^{-\delta(x)a_{n-1}}
$$

\n
$$
\leq 4C(x) \frac{e^{-\delta(x)t}}{\varepsilon - \delta(x)} e^{-\frac{\delta(x)}{2}(n-1)}
$$

where the last inequality follows from $a_{n-1} = \varkappa(n-1) \geq \frac{1}{2}(n-1)$. From this we readily derive the estimate

$$
I_2 \leq 2\sqrt{\frac{(\alpha+\beta)C(x)}{\varepsilon-\delta(x)}} \left(1 - e^{-\frac{\delta(x)}{4}}\right)^{-1} e^{-\frac{\delta(x)}{2}t}.
$$

Hence we obtain

$$
\mathrm{W}_2(p_{t+\tau}(x,\cdot), p_t(x,\cdot)) \le K(\alpha,\beta,\varepsilon,x) e^{-\frac{\delta(x)}{2}t}, \qquad t,\tau \ge 0,
$$
\n(4.6)

where the constant $K(\alpha, \beta, \varepsilon, x) > 0$ is given by

$$
K(\alpha, \beta, \varepsilon, x) = K(\varepsilon)(1 + \|x\|_H) + 2\sqrt{\frac{(\alpha + \beta)C(x)}{\varepsilon - \delta(x)}} \left(1 - e^{-\frac{\delta(x)}{4}}\right)^{-1}
$$

with another constant $K(\varepsilon) > 0$. This implies that, for each $x \in H$, $(p_t(x, \cdot))_{t \geq 0}$ has a limit in $\mathcal{P}_2(H)$. Denote this limit by π_{δ_x} . Assertion (a) now follows by taking the limit $\tau \to \infty$ in [\(4.6\)](#page-15-0) and using the fact that $K(\alpha, \beta, \varepsilon, x)$ is independent of τ .

It remains to prove assertion (b). First observe that, using $\delta(x) \ge \delta > 0$ and $C(x) \le$ $C(1 + ||x||_H)^4$, we have

$$
K(\alpha, \beta, \varepsilon, x) \le (1 + ||x||_H)^2 \widetilde{K}(\alpha, \beta, \varepsilon)
$$

for some constant $\widetilde{K}(\alpha, \beta, \varepsilon)$. Note that

$$
p_t^*\rho(dy) = \int_H p_t(z, dy)\rho(dz)
$$
 and $p_{t+\tau}^*\rho(dy) = \int_H p_{t+\tau}(z, dy)\rho(dz).$

Hence using first the convexity of the Wasserstein distance and then [\(4.6\)](#page-15-0) we find that

$$
W_2(p_{t+\tau}^* \rho, p_t^* \rho) \le \int_H W_2(p_{t+\tau}(x, \cdot), p_t(x, \cdot)) \rho(dx)
$$

$$
\le \widetilde{K}(\alpha, \beta, \varepsilon) \int_H (1 + ||x||_H)^2 \rho(dx) \cdot e^{-\frac{\delta}{2}t}.
$$

Since $\rho \in \mathcal{P}_2(H)$, the assertion is proved.

4.5. Proof of Corollary [4.4.](#page-12-2)

Proof of Corollary [4.4.](#page-12-2) Recall that, by condition (A2) the process $P_1 X_t^x$ is deterministic of the form

$$
P_1 X_t^x = P_1 S(t) P_1 x + \int_0^t P_1 S(t - s) F(P_1 X_s^x) ds.
$$

Since F is globally Lipschitz continuous by condition $(A1)$, it follows that this equation has for each $x \in H$ a unique solution. From this we readily conclude that $P_1 X_t^x = P_1 X_t^y$ t holds for all $t \geq 0$, provided that $P_1x = P_1y$. Hence Proposition [3.2](#page-6-3) yields for such x, y

$$
\mathbb{E}\left[\|X_t^x - X_t^y\|_H^2\right] \le e^{-\varepsilon t} \|x - y\|_H^2, \qquad \forall t \ge 0.
$$
 (4.7)

Then for each $x, y \in H$ with $P_1x = P_1y$ and each $t \geq 0$ we obtain

$$
W_2(\pi_{\delta_x}, \pi_{\delta_y}) \le W_2(\pi_{\delta_x}, p_t(x, \cdot)) + W_2(p_t(x, \cdot), p_t(y, \cdot)) + W_2(p_t(y, \cdot), \pi_{\delta_y})
$$

$$
\le W_2(\pi_{\delta_x}, p_t(x, \cdot)) + e^{-\frac{\varepsilon}{2}t} \|x - y\|_H + W_2(p_t(y, \cdot), \pi_{\delta_y}).
$$

Letting $t \to \infty$ yields $\pi_{\delta_x} = \pi_{\delta_y}$ and hence assertion (a) is proved.

To prove assertion (b), let $\rho, \tilde{\rho} \in \mathcal{P}_2(H)$ be such that $\rho \circ P_1^{-1} = \tilde{\rho} \circ P_1^{-1}$. Then

$$
W_2(\pi_\rho, \pi_{\widetilde{\rho}}) \le W_2(\pi_\rho, p_t^*\rho) + W_2(p_t^*\rho, p_t^*\widetilde{\rho}) + W_2(p_t^*\widetilde{\rho}, \pi_{\widetilde{\rho}})
$$

Again, by letting $t \to \infty$, it suffices to prove that

$$
\limsup_{t \to \infty} \mathcal{W}_2(p_t^* \rho, p_t^* \widetilde{\rho}) = 0.
$$
\n(4.8)

Let G be a coupling of $(\rho, \tilde{\rho})$. Using the convexity of the Wasserstein distance and Proposition [3.2](#page-6-3) gives

$$
W_2(\mathcal{P}_t^*\rho, \mathcal{P}_t^*\tilde{\rho})
$$

\n
$$
\leq \int_{H \times H} W_2(p_t(x, \cdot), p_t(y, \cdot)) G(dx, dy)
$$

\n
$$
\leq \int_{H \times H} \left(\mathbb{E} \left[\|X_t^x - X_t^y\|_H^2 \right] \right)^{1/2} G(dx, dy)
$$

\n
$$
\leq \int_{H \times H} e^{-\frac{\varepsilon}{2}t} \|x - y\|_H G(dx, dy)
$$

\n
$$
+ \sqrt{2(\alpha + \beta)} \int_{H^2} \left(\int_0^t e^{-\varepsilon(t-s)} \mathbb{E} \left[\|P_1 X_s^x - P_1 X_s^y\|_H^2 \right] ds \right)^{1/2} G(dx, dy)
$$

\n
$$
=: I_1 + I_2.
$$

The first term I_1 satisfies

$$
I_1 \le \left(2 + \int_H \|x\|_H^2 \rho(dx) + \int_H \|y\|_H^2 \widetilde{\rho}(dy)\right) e^{-\frac{\varepsilon}{2}t}.
$$

18 BÁLINT FARKAS, MARTIN FRIESEN, BARBARA RÜDIGER, AND DENNIS SCHROERS

For the second term we first use (A2) so that $P_1 X_s^x = P_1 X_s^{P_1x}, P_1 X_s^y = P_1 X_s^{P_1y}$ and hence we find for each $T > 0$ a constant $C(T) > 0$ such that for $t \in [0, T]$

$$
I_2 = \sqrt{2(\alpha + \beta)} \int_{H_1^2} \left(\int_0^t e^{-\varepsilon(t-s)} \|P_1 X_s^x - P_1 X_s^y\|_H^2 ds \right)^{1/2} G(dx, dy)
$$

= $C(T) \left(\int_{H^2} \|P_1 x - P_1 y\|_H^2 G(dx, dy) \right)^{1/2}.$

Let us choose a particular coupling G as follows: By disintegration we write $\rho(dx)$ $\rho(x_1, dx_0) (\rho \circ P_1^{-1})(dx_1), \widetilde{\rho}(dx) = \widetilde{\rho}(x_1, dx_0) (\widetilde{\rho} \circ P_1^{-1})(dx_1) = \widetilde{\rho}(x_1, dx_0) (\rho \circ P_1^{-1})(dx_1)$ where $\rho(x_1, dx_0)$, $\tilde{\rho}(x_1, dx_0)$ are transition kernels defined on $H_1 \times \mathcal{B}(H_0)$ and we have used that $(\rho \circ P_1^{-1})(dx_1) = (\tilde{\rho} \circ P_1^{-1})(dx_1)$. Then G is, for $A, B \in \mathcal{B}(H)$, given by

$$
G(A \times B) := \int_{H^2} \mathbb{1}_A(x_0, x_1) \mathbb{1}_B(y_0, y_1) \rho(x_1, dx_0) \widetilde{\rho}(y_1, dy_0) \widetilde{G}(dx_1, dy_1),
$$

where \tilde{G} is a probability measure on H_1^2 given, for $A_1, B_1 \in \mathcal{B}(H_1)$, by

$$
\widetilde{G}(A_1 \times B_1) = (\rho \circ P_1^{-1})(A_1 \cap B_1) = \rho \left(\{ x \in H \mid P_1 x \in A_1 \cap B_1 \} \right).
$$

For this particular choice of G we find that

$$
\int_{H^2} ||P_1x - P_1y||_H^2 G(dx, dy) = \int_{H_1^2} \int_{H_0^2} ||x_1 - y_1||_H^2 \rho(x_1, dx_0) \widetilde{\rho}(y_1, dy_0) \widetilde{G}(dx_1, dy_1)
$$

$$
= \int_{H_1^2} ||x_1 - y_1||_H^2 \widetilde{G}(dx_1, dy_1) = 0
$$

and hence $I_2 = 0$. This proves [\(4.8\)](#page-16-0) and completes the proof.

5. Ornstein-Uhlenbeck process on Hilbert space

Let H be a separable Hilbert space and let $(Z_t)_{t>0}$ be a H-valued Lévy process on a stochastic basis $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t>0}, \mathbb{P})$ with the usual conditions. Then, following [\[3\]](#page-24-4), it has characteristic exponent Ψ of Lévy-Khinchine form, i.e.

$$
\mathbb{E}\left[e^{i\langle u,Z_t\rangle_H}\right] = e^{t\Psi(u)}, \qquad u \in H, \quad t > 0,
$$

with Ψ given by

$$
\Psi(u) = \mathrm{i} \langle b, u \rangle_H - \frac{1}{2} \langle Qu, u \rangle_H + \int_H \left(e^{\mathrm{i} \langle u, z \rangle_H} - 1 - \mathrm{i} \langle u, z \rangle_H \mathbb{1}_{\{|z\|_H \le 1\}} \right) \mu(dz),
$$

where $b \in H$ denotes the drift, Q denotes the covariance operator being a positive, symmetric, trace-class operator on H, and μ is a Lévy measure on H. Let $(S(t))_{t\geq 0}$ be a strongly continuous semigroup on H . The Ornstein-Uhlenbeck process driven by $(Z_t)_{t\geq0}$ is the unique mild solution to

$$
dX_t^x = AX_t^x dt + dZ_t, \qquad X_0^x = x \in H, \quad t \ge 0,
$$

where $(A, D(A))$ denotes the generator of $(S(t))_{t\geq0}$, i.e. $(X_t^x)_{t\geq0}$ satisfies

$$
X_t^x = S(t)x + \int_0^t S(t-s)dZ_s, \qquad t \ge 0.
$$

The characteristic function of $(X_t^x)_{t\geq 0}$ is given by

$$
\mathbb{E}\left[e^{i\langle u,X_t^x\rangle_H}\right] = \exp\left(i\langle S(t)x, u\rangle_H + \int_0^t \Psi(S(r)^*u)dr\right), \qquad u \in H, \quad t \ge 0.
$$

For additional properties, references and related results we refer to the review article [\[3\]](#page-24-4) where also the existence, uniqueness and properties of invariant measures are discussed. Following these results, the Ornstein-Uhlenbeck process has a unique invariant measure provided that $(S(t))_{t\geq0}$ is uniformly exponentially stable, that is

$$
\exists \alpha > 0, \ M \ge 1: \qquad \|S(t)\|_{L(H)} \le M e^{-\alpha t}, \qquad t \ge 0,
$$

and the Lévy measure μ satisfies a log-integrability condition for its big jumps

$$
\int_{\{|z\|_H > 1\}} \log(1 + \|z\|_H) \mu(dz) < \infty. \tag{5.1}
$$

Below we show that for a uniformly convergent semigroup $(S(t))_{t\geq0}$ the corresponding Ornstein-Uhlenbeck process may admit multiple invariant measures parameterized by the range of the limiting projection operator of the semigroup.

Theorem 5.1. *Suppose that* $(S(t))_{t>0}$ *is uniformly exponentially convergent, i.e. there exists a projection operator* P *on* H *and constants* $M \geq 1$, $\alpha > 0$ *such that*

$$
||S(t)x - Px||_H \le M||x||_H e^{-\alpha t}, \qquad t \ge 0, x \in H.
$$
 (5.2)

Suppose that the Lévy process satisfies the following conditions:

- *(i)* The drift b satisfies $Pb = 0$.
- *(ii)* The covariance operator Q satisfies $PQu = 0$ for all $u \in H$.
- *(iii)* The Lévy measure μ *is supported on* ker(P) and satisfies [\(5.1\)](#page-18-0).

Then

$$
X_t^x \longrightarrow Px + X_\infty^0, \qquad x \in H,
$$

in law, where X^0_∞ *is an H*-valued random variable determined by

$$
\mathbb{E}\left[e^{\mathrm{i}\langle u, X_{\infty}^0 \rangle_H}\right] = \exp\left(\int_0^{\infty} \Psi(S(r)^*u) dr\right).
$$

In particular, the set of all limiting distributions for the Ornstein-Uhlenbeck process $(X_t^x)_{t\geq 0}$ *is given by* $\{\delta_x * \mu_\infty \mid x \in \text{ker}(P)\}\$ *, where* μ_∞ *denotes the law of* X_∞^0 *.*

Proof. We first prove the existence of a constant $C > 0$ such that

$$
\int_0^\infty |\Psi(S(r)^*u)| dr \le C(||u||_H + ||u||_H^2), \qquad u \in H. \tag{5.3}
$$

To do so we estimate

$$
|\Psi(S(r)^*u)| \le |\langle b, S(r)^*u \rangle| + |\langle QS(r)^*u, S(r)^*u \rangle|
$$

+
$$
\int_{\{|z||_H \le 1\}} |e^{i\langle S(r)^*u, z \rangle} - 1 - i\langle S(r)^*u, z \rangle| \mu(dz)
$$

+
$$
\int_{\{|z||_H > 1\}} |e^{i\langle S(r)^*u, z \rangle} - 1| \mu(dz)
$$

= $I_1 + I_2 + I_3 + I_4.$

We find by [\(5.2\)](#page-18-1) that $||S(r)x||_H \leq Me^{-\alpha r}||x||_H$ for all $x \in \text{ker}(P)$ and hence

$$
I_1 = |\langle S(r)b, u \rangle| \le ||u||_H ||S(r)b||_H \le ||u||_H Me^{-\alpha r} ||b||_H.
$$

For the second term I_2 we use $\text{ran}(Q) \subset \text{ker}(P)$ so that

$$
||S(r)Qu||_H \le Me^{-\alpha r}||Qu||_H \le e^{-\alpha r}||Q||_{L(H)}||u||_H.
$$

This yields $||QS(r)^*||_{L(H)} = ||S(r)Q||_{L(H)} \leq Me^{-\alpha r} ||Q||_{L(H)}$ and hence $I_2 = |\langle QS(r)^*u, S(r)^*u \rangle|$

$$
\leq \|QS(r)^*u\|_H \|S(r)^*u\|_H
$$

\n
$$
\leq M \|u\|_H \|QS(r)^*u\|_H
$$

\n
$$
\leq M \|u\|_H^2 \|Q\|_{L(H)} M e^{-\alpha r}.
$$

For the third term I_3 we obtain

$$
I_3 \leq C \int_{\{||z||_H \leq 1\}} |\langle S(r)^* u, z \rangle|^2 \mu(dz)
$$

= $C \int_{\{||z||_H \leq 1\} \cap \ker(P)} |\langle u, S(r)z \rangle|^2 \mu(dz)$
 $\leq C ||u||_H^2 e^{-\alpha r} \int_{\{||z||_H \leq 1\}} ||z||_H^2 \mu(dz),$

where $C > 0$ is a generic constant. Proceeding similarly for the last term, we obtain

$$
I_3 \leq C \int_{\{||z||_H > 1\}} \min\{1, |\langle S(r)^*u, z \rangle| \} \mu(dz)
$$

\n
$$
\leq C \int_{\{||z||_H > 1\} \cap \ker(P)} \min\{1, ||u||_H e^{-\alpha r} ||z||_H \} \mu(dz)
$$

\n
$$
\leq C ||u||_H e^{-\alpha r} \left(\mu(\{||z||_H > 1\}) + \int_{\{||z||_H > 1\}} \log(1 + ||z||_H) \mu(dz) \right),
$$

where we have used, for $a = ||u||_H e^{-\alpha r}$, $b = ||z||_H$, the elementary inequalities $\min\{1, ab\} < C \log(1 + ab)$

$$
n{1, ab} \le C \log(1 + ab)
$$

\n
$$
\le C \min\{\log(1 + a), \log(1 + b)\} + C \log(1 + a) \log(1 + b)
$$

\n
$$
\le C a (1 + \log(1 + b)),
$$

see [\[18,](#page-25-20) appendix]. Combining the estimates for I_1, I_2, I_3, I_4 we conclude that [\(5.3\)](#page-18-2) is satisfied. Hence, using

$$
\lim_{t \to \infty} \langle S(t)x, u \rangle = \langle Px, u \rangle
$$

we find that

$$
\lim_{t \to \infty} \mathbb{E}\left[e^{i\langle u, X_t^x \rangle}\right] = \exp\left(i\langle Px, u \rangle + \int_0^\infty \Psi(S(r)^*u) dr\right). \tag{5.4}
$$

Since, in view of [\(5.3\)](#page-18-2), $u \mapsto \int_0^\infty \Psi(S(r)^*u) dr$ is continuous at $u = 0$, the assertion follows from Lévy's continuity theorem combined with the particular form of (5.4) . \Box

Below we briefly discuss an application of this result to a stochastic perturbation of the Kolmogorov equation associated with a symmetric Markov semigroup. Let X be a Polish space and η a Borel probability measure on X. Let $(A, D(A))$ be the generator of a symmetric Markov semigroup $(S(t))_{t\geq 0}$ on $H := L^2(X, \eta)$. Then there exists, for each $f \in D(A)$, a unique solution to the Kolmogorov equation (see, e.g., [\[29\]](#page-25-18))

$$
\frac{dv(t)}{dt} = Av(t), \qquad v(0) = f.
$$

Below we consider an additive stochastic perturbation of this equation in the sense of Itô, i.e. the stochastic partial differential equation

$$
dv(t) = Av(t)dt + dZ_t, \qquad v(0) = f,
$$
\n
$$
(5.5)
$$

where $(Z_t)_{t\geq 0}$ is an $L^2(X,\eta)$ -valued Lévy process with characteristic function Ψ . Let $(v(t); f)$ _t ≥ 0 be the unique mild solution to this equation.

Corollary 5.2. Suppose that the semigroup generated by $(A, D(A))$ on $L^2(X, \eta)$ satisfies [\(5.2\)](#page-18-1) *with the projection operator*

$$
P_1v = \int_X v(x)\eta(dx),
$$

and assume that the Lévy process $(Z_t)_{t>0}$ *satisfies the conditions (i) – (iii) of Theorem [5.1.](#page-18-3) Then*

$$
v(t; f) \longrightarrow \int_X f(x)\eta(dx) + v(\infty), \qquad t \to \infty
$$

in law, where $v(\infty)$ *is a random variable whose characteristic function is given by*

$$
\mathbb{E}\left[e^{\mathrm{i}\langle u,v(\infty)\rangle_{L^2}}\right] = \exp\left(\int_0^\infty \Psi(S(r)^*u)dr\right).
$$

6. The Heath-Jarrow-Mortion-Musiela equation

The Heath-Jarrow-Morton-Musiela equation (HJMM-equation) describes the term structure of interest rates in terms of its forward rate dynamics modelled, for $\beta > 0$ fixed, on the space of forward curves

$$
H_{\beta} = \{h : \mathbb{R}_{+} \to \mathbb{R} : h \text{ is absolutely continuous and } ||h||_{\beta} < \infty\}, \qquad (6.1)
$$

$$
||h||_{\beta}^{2} = |h(\infty)|^{2} + \int_{0}^{\infty} (h'(x))^{2} e^{\beta x} dx < \infty
$$

Such space was first motivated and introduced by Filipovic [\[14\]](#page-25-21). Note that $h(\infty) :=$ $\lim_{x\to\infty} h(x)$ exists, whenever $h \in H_\beta$. It is called the *long rate* of the forward curve h. The HJMM-equation on H_β is given by

$$
\begin{cases} dX_t = (AX_t + F_{HJMM}(\sigma, \gamma)(X_t)) dt + \sigma(X_t) dW_t + \int_E \gamma(X_t, \nu) \widetilde{N}(dt, d\nu), \\ X_0 = h_0 \in L^2(\Omega, \mathcal{F}_0, \mathbb{P}; H_\beta) \end{cases} \tag{6.2}
$$

where $(W_t)_{t\geq0}$ is a Q-Wiener process and $\widetilde{N}(dt, d\nu)$ is a compensated Poisson random measure on E with compensator $dt\mu(d\nu)$ as defined in Section 2 for $H := H_\beta$, and

- (i) A is the infinitesimal generator of the shift semigroup $(S(t))_{t\in\mathbb{R}_+}$ on H_β , that is $S(t)h(x) := h(x+t)$ for all $t, x \geq 0$.
- (ii) $h \mapsto \sigma(h)$ is a $\mathcal{B}(H_\beta)/\mathcal{B}(L_2^0)$ -measurable mapping from H_β into $L_2^0(H_\beta)$ and $(h, \nu) \mapsto \gamma(h, \nu)$ is $\mathcal{B}(H_\beta) \otimes \mathcal{E}/\mathcal{B}(H_\beta)$ -measurable mapping from $H_\beta \times E$ into H_{β} .
- (iii) The drift is of the form

$$
F_{HJMM}(\sigma, \gamma)(h) = \sum_{j \in \mathbb{N}} \sigma^j(h) \Sigma^j(h) - \int_E \gamma(h, \nu)(e^{\Gamma(h, \nu)} - 1)\mu(d\nu),
$$

with $\sigma^j(h) = \sqrt{\lambda_j} \sigma(h)e_j,$
 $\Sigma^j(h)(t) = \int_0^t \sigma^j(h)(s)ds$ and $\Gamma(h, \nu)(t) = -\int_0^t \gamma(h, \nu)(s)ds.$

The special form of the drift stems from mathematical finance and is sufficient for the absence of arbitrage opportunities. We denote the space of all forward rates with long rate equal to zero by

$$
H_{\beta}^{0} = \{ h \in H_{\beta} : h(\infty) = 0 \}.
$$

For the construction of a unique mild solution to [\(6.2\)](#page-20-0) the following conditions have been introduced in [\[11\]](#page-25-22):

22 BÁLINT FARKAS, MARTIN FRIESEN, BARBARA RÜDIGER, AND DENNIS SCHROERS

- (B1) $\sigma: H_{\beta} \to L_2^0(H_{\beta}^0), \gamma: H_{\beta} \times E \to H_{\beta'}^0$ are Borel measurable for some $\beta' > \beta$.
- (B2) There exists a function $\Phi: E \to \mathbb{R}_+$ such that $\Phi(\nu) \geq |\Gamma(h,\nu)(t)|$ for all $h \in H_\beta$, $\nu \in E$ and $t \geq 0$.
- (B3) There is an $M \geq 0$ such that, for all $h \in H_\beta$, and some $\beta' > \beta$

$$
\|\sigma(h)\|_{L_2^0(H_\beta)} \le M, \quad \int_E e^{\Phi(\nu)} \max\{\|\gamma(h,\nu)\|_{\beta'}^2, \|\gamma(h,\nu)\|_{\beta'}^4\} \mu(d\nu) \le M.
$$

(B4) The function $F_2: H_\beta \to H_\beta^0$ defined by

$$
F_2(h) = -\int_E \gamma(h,\nu) \left(e^{\Gamma(h,\nu)} - 1 \right) \mu(d\nu)
$$

has the weak derivative given by

$$
\frac{d}{dx}F_2(h) = \int_E \gamma(h,\nu)^2 \left(e^{\Gamma(h,\nu)}\right) \mu(d\nu) - \int_E \left(\frac{d}{dx}\gamma(h,\nu)\right) \left(e^{\Gamma(h,\nu)} - 1\right) \mu(d\nu).
$$

(B5) There are constants L_{σ} , $L_{\gamma} > 0$ such that, for all $h_1, h_2 \in H_{\beta}$, we have

$$
\|\sigma(h_1) - \sigma(h_2)\|_{L_2^0(H_\beta)}^2 \le L_\sigma \|h_1 - h_2\|_{\beta}^2,
$$

$$
\int_E e^{\Phi(\nu)} \|\gamma(h_1, \nu) - \gamma(h_2, \nu)\|_{\beta'}^2 \mu(d\nu) \le L_\gamma \|h_1 - h_2\|_{\beta}^2.
$$

The following is the basic existence and uniqueness result for the Heath-Jarrow-Morton-Musiela equation [\(6.2\)](#page-20-0).

Theorem 6.1. [\[11\]](#page-25-22) *Suppose that conditions (B1) – (B5) are satisfied. Then* F_{HJMM} : $H_{\beta} \longrightarrow H_{\beta}^0$ and there exists a constant $L_F > 0$ such that, for each $h_1, h_2 \in H_{\beta}$,

$$
||F_{HJMM}(h_1) - F_{HJMM}(h_2)||_{\beta}^2 \le L_F ||h_1 - h_2||_{\beta}^2.
$$
 (6.3)

This constant can be choosen as

$$
L_F = \frac{\max(L_{\sigma}, L_{\gamma})\sqrt{M}}{\beta} \left(\sqrt{6M\sqrt{2}} + \sqrt{\frac{8}{\beta^3} + \frac{16}{\beta}} + \sqrt{\frac{16(1 + \frac{1}{\sqrt{\beta}})^2 + 48}{(\beta' - \beta)}} \right). \tag{6.4}
$$

Moreover, for each initial condition $h \in L^2(\Omega, \mathcal{F}_0, \mathbb{P}; H_\beta)$ *there is a unique adapted, cádlág mild solution* $(r_t)_{t>0}$ *to* [\(6.2\)](#page-20-0).

Proof. This result can be found essentially in [\[11\]](#page-25-22), where the bound on L_F is an immediate result from its derivation.

Using the space of all functions with zero long rate we obtain the decomposition

$$
H_{\beta} = H_{\beta}^0 \oplus \mathbb{R}, \qquad h = (h - h(\infty)) + h(\infty),
$$

where $h(\infty) \in \mathbb{R}$ is identified with a constant function. Denote by

$$
P_0 h = h - h(\infty) \text{ and } P_1 h = h(\infty)
$$

the corresponding projections onto H_{β}^0 and \mathbb{R} , respectively. Such a decomposition of H_β was first used in [\[36\]](#page-26-1) to study invariant measures for the HJMM-equation driven by a Q -Wiener process. An extension to the Lévy driven HJMM-equation was then obtained in [\[34\]](#page-26-2). The next theorem shows that the results of Section 4 contain the HJMM-equation as a particular case.

Theorem 6.2. Suppose that conditions $(B1) - (B5)$ are satisfied. If

$$
\beta > 2\sqrt{L_F} + L_{\sigma} + L_{\gamma},\tag{6.5}
$$

then for each initial distribution ρ *on* H_β *with finite second moments there exists an invariant measure* π_{ρ} *and it holds that*

$$
\mathrm{W}_{2}(\mathcal{P}_{t}^{*}\rho,\pi_{\rho}) \leq K \left(1 + \int_{H_{\beta}} \|h\|_{H_{\beta}}^{2} \rho(dh) + \int_{H_{\beta}} \|h\|_{H_{\beta}}^{2} \pi_{\rho}(dh)\right) e^{-\frac{\beta - 2\sqrt{L_{F}} - L_{\sigma} - L_{\gamma}}{2}t} \tag{6.6}
$$

for some constant $K = K(\beta, \sigma, \gamma) > 0$ *. Moreover, given* $\rho, \tilde{\rho}$ such that $\rho \circ P_1^{-1} = \tilde{\rho} \circ P_1^{-1}$, *then* $\pi_{\rho} = \pi_{\widetilde{\rho}}$ *.*

Proof. Observe that the assertion is an immediate consequence of Theorem [4.3](#page-12-1) and Corollary [4.4.](#page-12-2) Below we briefly verify the assumptions given in these statements. Condition $(A1)$ follows from $(B1)$, $(B5)$, and (6.3) while the growth condition (3.3) is satisfied by (B3) and the fact that $\|\cdot\|_{\beta} \leq \|\cdot\|_{\beta'}$ for $\beta < \beta'$. It is not difficult to see that

$$
||S(t)h - P_1h||_{\beta} \le e^{-\frac{\beta}{2}t}||h - P_1h||_{\beta}, \qquad t \ge 0
$$

and that $(S(t))_{t\geq0}$ leaves H^0_β as well as $\mathbb{R} \subset H_\beta$ invariant. Hence Remark [2.2](#page-5-0) yields that

$$
\langle Ah, h \rangle \le -\frac{\beta}{2} ||h||_{\beta}^2 + \frac{\beta}{2} ||P_1h||_{\beta}^2, \qquad h \in D(A).
$$

It follows from the considerations in Section 2 (see [\(2.10\)](#page-4-2)) that (GDC) is satisfied for $\alpha = \frac{\beta}{2}$ – $\sqrt{L_F}$. Consequently, $\varepsilon = \beta - 2\sqrt{L_F} - L_{\sigma} - L_{\gamma}$ and [\(3.2\)](#page-6-0) holds due to [\(6.5\)](#page-22-0). Since the coefficients map into H^0_β and $S(t)P_1h = h(\infty) = P_1h$, conditions (A2), (A3) and [\(4.4\)](#page-12-0) are trivially satisfied. The particular form of the estimate [\(6.6\)](#page-22-1) follows from the proof of Theorem [4.3.](#page-12-1)

Comparing our result with [\[36,](#page-26-1) [34\]](#page-26-2), we allow for a more general jump noise and prove convergence in the stronger Wasserstein distance with an exponential rate. Moreover, assuming that the volatilities map constant functions onto zero, i.e.

$$
\sigma(c) \equiv 0, \quad \gamma(c, \nu) \equiv 0, \qquad \forall c \in \mathbb{R} \subset H_{\beta}, \ \nu \in E \tag{6.7}
$$

shows that $F(c) \equiv 0$ and hence also [\(4.2\)](#page-10-0) is satisfied. Hence we may apply Theorem [4.1](#page-10-2) to characterize these invariant measures more explicitly.

Corollary 6.3. Suppose that conditions $(B1) - (B5)$ are satisfied, that (6.5) and (6.7) *hold. Then*

$$
\mathbb{E}\left[\|X_t^h - h(\infty)\|_{\beta}^2\right] \le \mathbb{E}\left[\|h - h(\infty)\|_{\beta}^2\right] e^{-\left(\beta - 2\sqrt{L_F} - L_{\sigma} - L_{\gamma}\right)t}
$$

for each $h \in L^2(\Omega, \mathbb{F}_0, \mathbb{P}, H)$ *.*

We close this section by applying our results for the particular example of coefficients as introduced in [\[34\]](#page-26-2).

Example 6.4. *Take*

$$
\sigma^{1}(h)(x) := \int_{x}^{\infty} \min\left(e^{-\beta y}, \ |h'(y)|\right) dy
$$

 \int *and* $\sigma^j \equiv 0$ *for* $j \geq 2$ *. Then*

$$
\|\sigma(h)\|_{L_2^0}^2 = \|\sigma^1(h)\|_{\beta}^2 \le \int_0^\infty (e^{-2\beta x}) e^{\beta x} dx = \frac{1}{\beta} =: M
$$

24 BÁLINT FARKAS, MARTIN FRIESEN, BARBARA RÜDIGER, AND DENNIS SCHROERS

and since min(a, b₁) − min(a, b₂) ≤ |b₁ − b₂| *for* a, b₁, b₂ ∈ ℝ₊*,* we also have

$$
\begin{aligned} \|\sigma(h_1) - \sigma(h_2)\|_{L_2^0}^2 &= \|\sigma^1(h) - \sigma^1(h_2)\|_{\beta}^2 \\ &= \int_0^\infty (\min(\mathrm{e}^{-\beta x}, |h_1'(x)|) - \min(\mathrm{e}^{-\beta x}, |h_2'(x)|))^2 \mathrm{e}^{\beta x} dx \\ &\le \int_0^\infty (h_1'(x) - h_2'(x))^2 \mathrm{e}^{\beta x} dx \\ &\le \|h_1 - h_2\|_{\beta}^2. \end{aligned}
$$

This shows that the Lipschitz condition for σ *holds, in particular, for* $L_{\sigma} = 1$ *. Consequently, by taking* $\gamma \equiv 0$ *, the conditions (B1) – (B5) are satisfied with* $L_{\sigma} = 1$ *and* $L_{\gamma} = 0$ and $M = \frac{1}{\beta}$ for the Lipschitz and growth constants. By [\(6.4\)](#page-21-1) we get

$$
L_F = \frac{1}{\sqrt{\beta^3}} \left(\sqrt{\frac{6\sqrt{2}}{\beta}} + \sqrt{\frac{8}{\beta^3} + \frac{16}{\beta}} + \sqrt{\frac{16(1 + \frac{1}{\sqrt{\beta}})^2 + 48}{(\beta' - \beta)}} \right),
$$

for all $\beta' > \beta$. Choosing $\beta \geq 3$ and $\beta' > \beta$ large enough such that $L_F < 1$, we find that $2\sqrt{L_F} + L_{\sigma} + L_{\gamma} < 3 = \beta,$

i.e. [\(6.5\)](#page-22-0) *is satisfied. It is clear that* $\sigma(c) \equiv 0$ *for each constant function c. Hence Corollary [6.3](#page-22-3) is applicable.*

APPENDIX A. ITÔ FORMULA

Below we recall an Itô formula for Hilbert space valued semimartingales of the form

$$
X(t) = X(0) + \int_0^t a(s)ds + \int_0^t \sigma(s)dW_s + \int_0^t \int_E \gamma(s,\nu)\widetilde{N}(ds,d\nu),
$$

where a and σ are as before and $(\gamma(t, \nu))_{t>0}$ is a predictable, H-valued stochastic process for each $\nu \in E$ such that

$$
\mathbb{E}\left[\int_0^t \int_E \|\gamma(s,\nu)\|_H^2 \mu(d\nu)ds\right] < \infty
$$

and

$$
\mathbb{E}\left[\int_0^t \|\sigma(s)\|_{L_2^0}^2 ds\right] < \infty.
$$

For this purpose we first introduce the class of quasi-sublinear functions.

Definition A.1 (Sublinear Functions). *A continuous, non-decreasing function* h : $\mathbb{R}_+ \to \mathbb{R}_+$ *is called quasi-sublinear, if there exists a constant* $C > 0$ *such that*

$$
h(x + y) \le C(h(x) + h(y))
$$

$$
h(xy) \le C(h(x)h(y))
$$

for all $x, y \geq 0$ *.*

The following Itô-Formula is a combination of [\[19\]](#page-25-23) and [\[27\]](#page-25-24).

Theorem A.2 (Generalized Itô-Formula). Let $F \in C^2(\mathbb{R}_+ \times H, \mathbb{R})$ and suppose there *exist quasi-sublinear functions* $h_1, h_2 : \mathbb{R}_+ \to \mathbb{R}_+$ *such that for all* $t \geq 0$ *and* $x \in H$

$$
||F_x(t,x)||_H \le h_1(||x||_H), \qquad ||F_{xx}(t,x)||_{L(H,L(H,\mathbb{R}))} \le h_2(||x||_H)
$$

and

$$
\int_0^t \int_E ||\gamma(s,\nu)||_H^2 \mu(d\nu)ds + \int_0^t \int_E h_1(||\gamma(s,\nu)||_H)^2 ||\gamma(s,\nu)||_H^2 \mu(d\nu)ds
$$

+
$$
\int_0^t \int_E h_2(||\gamma(s,\nu)||_H) ||\gamma(s,\nu)||_H^2 \mu(d\nu)ds < \infty
$$

Then $\mathbb{P}\text{-}almost surely for each $t > 0$:$

$$
\int_0^t \|F_t(s, X(s))\|_H ds + \int_0^t \int_E |F(s, X(s) + \gamma(s, \nu)) - F(s, X(s))|^2 \mu(d\nu) ds
$$

+
$$
\int_0^t \int_E |F(s, X(s) + \gamma(s, \nu)) - F(s, X(s)) - \langle F_x(s, X(s)), \gamma(s, \nu) \rangle_H) |\mu(d\nu) ds < \infty.
$$

Moreover, the generalized Itô-formula holds $\mathbb{P}\text{-almost surely for each $t \geq 0$ and$

$$
F(t, X(t)) = F(0, X(0)) + \int_0^t \mathcal{L}F(s, X(s))ds
$$

+
$$
\int_0^t \langle F_x(s, X(s)), \sigma(s)dW_s \rangle_H
$$

+
$$
\int_0^{t+} \int_E \{ F(s, X(s-) + \gamma(s, \nu)) - F(s, X(s-)) \} \widetilde{N}(ds, d\nu)
$$

where $LF(x, X(s))$ *is given by*

$$
\mathcal{L}F(s, X(s))
$$
\n
$$
= \int_0^t \{F_t(s, X(s)) + \langle F_x(s, X(s)), a(s) \rangle_H\} ds
$$
\n
$$
+ \frac{1}{2} \int_0^t \text{tr}\left[F_{xx}(s, X(s))\sigma(s)Q\sigma(s)^*\right] ds
$$
\n
$$
+ \int_0^t \int_E \{F(s, X(s) + \gamma(s, \nu)) - F(s, X(s)) - \langle F_x(s, X(s)), \gamma(s, \nu) \rangle_H\} \mu(d\nu) ds
$$

Acknowledgements. Dennis Schroers was funded within the project STORM: Stochastics for Time-Space Risk Models, from the Research Council of Norway (RCN). Project number: 274410.

REFERENCES

- 1. S. Albeverio, V Mandrekar, and B. Rüdiger, Existence of mild solutions for stochastic differential equations and semilinear equations with non-gaussian lévy noise, Stochastic Processes and their Applications 119 (2009), no. 3, 835 – 863.
- 2. Sergio Albeverio, Leszek Gawarecki, Vidyadhar Mandrekar, Barbara Rüdiger, and Barun Sarkar, Itô Formula for Mild Solutions of SPDEs with Gaussian and non-Gaussian noise and Applications to Stability Properties, Random Operators and Stochastic Equations 25 (2017), no. 2.
- 3. David Applebaum, Infinite dimensional Ornstein-Uhlenbeck processes driven by Lévy processes, Probab. Surv. 12 (2015), 33–54. MR 3385977
- 4. Viorel Barbu, Michael Röckner, and Deng Zhang, Stochastic nonlinear Schrödinger equations, Nonlinear Anal. 136 (2016), 168–194. MR 3474409
- 5. András Bátkai and Susanna Piazzera, Semigroups for delay equations, Research Notes in Mathematics, vol. 10, A K Peters, Ltd., Wellesley, MA, 2005. MR 2181405
- 6. Fred Espen Benth, Nils Detering, and Paul Kruehner, Stochastic Volterra integral equations and a class of first order stochastic partial differential equations, arXiv:1903.05045 (2019).
- 7. Joris Bierkens and Onno van Gaans, Dissipativity of the delay semigroup, J. Differential Equations 257 (2014), no. 7, 2418–2429. MR 3228971

26 BÁLINT FARKAS, MARTIN FRIESEN, BARBARA RÜDIGER, AND DENNIS SCHROERS

- 8. Oleg Butkovsky, Subgeometric rates of convergence of Markov processes in the Wasserstein metric, Ann. Appl. Probab. 24 (2014), no. 2, 526–552. MR 3178490
- 9. Giuseppe Da Prato and Arnaud Debussche, Ergodicity for the 3D stochastic Navier-Stokes equations, J. Math. Pures Appl. (9) 82 (2003), no. 8, 877–947. MR 2005200
- 10. Giuseppe Da Prato and Jerzy Zabczyk, Stochastic Equations in Infinite Dimensions, Encyclopedia of Mathematics and its Applications, Cambridge University Press, 2014.
- 11. Stefan Tappe Damir Filipović and Josef Teichmann, Term Structure Models Driven by Wiener Processes and Poisson Measures: Existence and Positivity, SIAM Journal on Financial Mathematics 1 (2010), no. 1, 523–554.
- 12. Donald A. Dawson, Alison M. Etheridge, Klaus Fleischmann, Leonid Mytnik, Edwin A. Perkins, and Jie Xiong, Mutually catalytic branching in the plane: infinite measure states, Electron. J. Probab. 7 (2002), No. 15, 61. MR 1921744
- 13. A. de Bouard and A. Debussche, The stochastic nonlinear Schrödinger equation in H^1 , Stochastic Anal. Appl. 21 (2003), no. 1, 97–126. MR 1954077
- 14. Damir Filipović, Consistency Problems for HJM Interest Rate Models, Springer, Berlin; Heidelberg ; New York ; Barcelona ; Hong Kong ; London ; Milan ; Paris ; Singapore ; Tokyo, 2001.
- 15. Damir Filipović, Term-Structure Models: A Graduate Course, Springer Finance, Springer Berlin Heidelberg, 2009.
- 16. Damir Filipović, Stefan Tappe, and Josef Teichmann, Jump-Diffusions in Hilbert Spaces: Existence, Stability and Numerics, Stochastics 82 (2010), no. 5, 475–520.
- 17. Franco Flandoli, Dissipativity and invariant measures for stochastic Navier-Stokes equations, NoDEA Nonlinear Differential Equations Appl. 1 (1994), no. 4, 403–423. MR 1300150
- 18. Martin Friesen, Peng Jin, and Barbara Rüdiger, Stochastic equation and exponential ergodicity in Wasserstein distances for affine processes, Ann. Appl. Prob. (2020), (to appear).
- 19. Leszek Gawarecki and Vidyadhar Mandrekar, Stochastic Differential Equations in Infinite Dimensions: with Applications to Stochastic Partial Differential Equations, Probability and Its Applications, Springer Berlin Heidelberg, 2010.
- 20. M. Hairer, J. C. Mattingly, and M. Scheutzow, Asymptotic coupling and a general form of Harris' theorem with applications to stochastic delay equations, Probab. Theory Related Fields 149 (2011), no. 1-2, 223–259. MR 2773030
- 21. Martin Hairer and Jonathan C. Mattingly, Ergodicity of the 2D Navier-Stokes equations with degenerate stochastic forcing, Ann. of Math. (2) 164 (2006), no. 3, 993–1032. MR 2259251
- 22. $____________\.\$ Yet another look at Harris' ergodic theorem for Markov chains, Seminar on Stochastic Analysis, Random Fields and Applications VI, Progr. Probab., vol. 63, Birkhäuser/Springer Basel AG, Basel, 2011, pp. 109–117. MR 2857021
- 23. David Heath, Robert Jarrow, and Andrew Morton, Bond Pricing and the Term Structure of Interest Rates: A New Methodology for Contingent Claims Valuation, Econometrica 60 (1992), 77–105.
- 24. Alexei Kulik and Michael Scheutzow, Generalized couplings and convergence of transition probabilities, Probab. Theory Related Fields 171 (2018), no. 1-2, 333–376. MR 3800835
- 25. Thomas G. Kurtz, The Yamada-Watanabe-Engelbert theorem for general stochastic equations and inequalities, Electron. J. Probab. 12 (2007), 951–965. MR 2336594
- 26. Vidyadhar Mandrekar and Barbara Rüdiger, Stochastic Integration in Banach Spaces: Theory and Applications (Probability Theory and Stochastic Modelling Book 73), Springer, 2014.
- 27. Vidyadhar Mandrekar, Barbara Rüdiger, and Stefan Tappe, Itô's Formula for Banach-space-valued Jump Processes Driven by Poisson Random Measures, Seminar on Stochastic Analysis, Random Fields and Applications VII, Springer Basel, 2013, pp. 171–186.
- 28. Leonid Mytnik, Uniqueness for a mutually catalytic branching model, Probab. Theory Related Fields 112 (1998), no. 2, 245–253. MR 1653845
- 29. A. Pazy, Semigroups of linear operators and applications to partial differential equations, Applied Mathematical Sciences, vol. 44, Springer-Verlag, New York, 1983. MR 710486
- 30. Szymon Peszat and Jerzy Zabczyk, Stochastic Partial Differential Equations with Lévy Noise: An Evolution Equation Approach, Encyclopedia of Mathematics and its Applications, Cambridge University Press, 2007.
- 31. Enrico Priola, Armen Shirikyan, Lihu Xu, and Jerzy Zabczyk, Exponential ergodicity and regularity for equations with Lévy noise, Stochastic Process. Appl. 122 (2012), no. 1, 106–133. MR 2860444
- 32. Enrico Priola, Lihu Xu, and Jerzy Zabczyk, Exponential mixing for some SPDEs with Lévy noise, Stoch. Dyn. 11 (2011), no. 2-3, 521–534. MR 2836539
- 33. Enrico Priola and Jerzy Zabczyk, On linear evolution equations for a class of cylindrical Lévy noises, Stochastic partial differential equations and applications, Quad. Mat., vol. 25, Dept. Math., Seconda Univ. Napoli, Caserta, 2010, pp. 223–242. MR 2985090
- 34. Anna Rusinek, Mean Reversion for HJMM Forward Rate Models, Advances in Applied Probability 42 (2010), no. 2, 371391.
- 35. S. S. Sritharan and P. Sundar, Large deviations for the two-dimensional Navier-Stokes equations with multiplicative noise, Stochastic Process. Appl. **116** (2006), no. 11, 1636–1659. MR 2269220
- 36. Michael Tehranchi, A Note on Invariant Measures for HJM Models, Finance and Stochastics 9 (2005), no. 3, 389–398.
- 37. Onno van Gaans, Invariant measures for stochastic evolution equations with Hilbert space valued $Lévy$ noise, 2005.
- 38. Cédric Villani, Optimal transport, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 338, Springer-Verlag, Berlin, 2009, Old and new. MR 2459454

(Bálint Farkas) SCHOOL OF MATHEMATICS AND NATURAL SCIENCES, UNIVERSITY OF WUPPERTAL, **GERMANY**

E-mail address, Bálint Farkas: farkas@math.uni-wuppertal.de

(Martin Friesen) School of Mathematics and Natural Sciences, University of Wuppertal, **GERMANY**

E-mail address, Martin Friesen: friesen@math.uni-wuppertal.de

(Barbara Rüdiger) SCHOOL OF MATHEMATICS AND NATURAL SCIENCES, UNIVERSITY OF WUPPERtal, Germany

E-mail address, Barbara Rüdiger: ruediger@uni-wuppertal.de

(Dennis Schroers) Department of Mathematics, University of Oslo E-mail address, Dennis Schroers: dennissc@math.uio.no