

New Approximation Algorithms for Maximum Asymmetric Traveling Salesman and Shortest Superstring

Katarzyna Paluch
Institute of Computer Science, University of Wrocław
abraka@cs.uni.wroc.pl

Abstract

In the maximum asymmetric traveling salesman problem (Max ATSP) we are given a complete directed graph with nonnegative weights on the edges and we wish to compute a traveling salesman tour of maximum weight. In this paper we give a fast combinatorial $\frac{7}{10}$ -approximation algorithm for Max ATSP. It is based on techniques of *eliminating* and *diluting* problematic subgraphs with the aid of *half-edges* and a method of edge coloring. (A *half-edge* of edge (u, v) is informally speaking “either a head or a tail of (u, v) ”). A novel technique of *diluting* a problematic subgraph S consists in a seeming reduction of its weight, which allows its better handling.

The current best approximation algorithms for Max ATSP, achieving the approximation guarantee of $\frac{2}{3}$, are due to Kaplan, Lewenstein, Shafrir, Sviridenko (2003) and Elbassioni, Paluch, van Zuylen (2012). Using a result by Mucha, which states that an α -approximation algorithm for Max ATSP implies a $(2 + \frac{11(1-\alpha)}{9-2\alpha})$ -approximation algorithm for the shortest superstring problem (SSP), we obtain also a $(2\frac{33}{76} \approx 2,434)$ -approximation algorithm for SSP, beating the previously best known (having an approximation factor equal to $2\frac{11}{23} \approx 2,4782$.)

1 Introduction

In the maximum asymmetric traveling salesman problem (Max ATSP) we are given a complete directed graph $G = (V, E)$ with nonnegative weights on the edges and we wish to compute a traveling salesman tour of maximum weight. The problem is known to be APX-hard [26] and the current best approximation algorithms for it are due to Kaplan, Lewenstein, Shafir, Sviridenko [13] obtained in 2003 and Elbassioni, Paluch, van Zuylen [25] published in 2012. Both of them achieve the approximation ratio of $\frac{2}{3}$, the former is based on linear programming and the other is combinatorial and simpler. Besides being an interesting problem in itself, Max ATSP is also of particular interest because of its applications to a number of related problems. For example, an α -approximation algorithm for Max ATSP implies a $(2 + \frac{11(1-\alpha)}{9-2\alpha})$ -approximation algorithm for SSP, which was shown by Mucha [20]. The shortest superstring problem is defined as follows. We are given n strings s_1, s_2, \dots, s_n over a given alphabet Σ and we want to find a shortest string s such that each s_i for $i, 1 \leq i \leq n$ is a substring of s . SSP arises in DNA sequencing and data compression. Currently the best approximation algorithm for SSP is due to Mucha [20] and achieves an approximation factor of $2\frac{11}{23}$. For a long time the best approximation algorithm for SSP was the one given by Sweedyk [27] in 1999 with an approximation factor of $2\frac{1}{2}$. Any α -approximation algorithm for Max ATSP implies also an algorithm with the same guarantee for the maximal compression problem defined by Tarhio and Ukkonen [28].

We devise a combinatorial $\frac{7}{10}$ -approximation algorithm for Max ATSP, thus proving

Theorem 1 *There exists a $\frac{7}{10}$ -approximation algorithm for the maximum asymmetric traveling salesman problem.*

Using the result of Mucha [20], we obtain

Corollary 1 *There exists a $2\frac{33}{76}$ -approximation algorithm for the shortest superstring problem.*

The presented results are a simpler and weaker version of [22].

The approach we have adopted is as follows. We start by computing a maximum weight cycle cover C_{max} of G , where a cycle cover C of graph G is defined as a set of directed cycles of G such that each vertex of G belongs to exactly one cycle of C . A maximum weight cycle cover of G can be found in polynomial time by a reduction to maximum weight matching. Let opt denote the weight of a traveling salesman tour of G of maximum weight. The weight of an edge e will be denoted as $w(e)$ and for any subset E' of edges E by $w(E')$ we will mean $\sum_{e \in E'} w(e)$. Since a traveling salesman tour is a cycle cover of G (consisting of just one cycle), we know that $w(C_{max}) \geq opt$. By removing the lightest edge from each cycle of C_{max} , we obtain a collection of vertex-disjoint paths, which can be arbitrarily patched to form a tour. Removing the lightest edge from cycle c of length k results in a path of weight at least $\frac{k-1}{k}w(c)$. Since C_{max} may contain cycles of length two (2-cycles), in the worst case the obtained tour may have weight equal to $\frac{1}{2}w(C_{max})$. If we could find a maximum weight cycle cover of G without cycles of length two (2-cycles) or three (3-cycles or *triangles*), then we would achieve a $\frac{3}{4}$ -approximation, but, unfortunately finding a maximum weight cycle cover without 2-cycles is APX-hard [6].

Eliminating and diluting problematic subgraphs with the aid of half-edges Since 2- and 3-cycles in a maximum weight cycle cover are an obstacle to getting a $\frac{7}{10}$ -approximation, we would like to somehow get rid of them. To this end we use a technique of eliminating problematic subgraphs with the aid of *half-edges* - a half-edge of edge (u, v) is informally speaking “either a head or a tail of (u, v) ”. Half-edges have already been introduced in [25].

They have also been employed in [1],[10], [24]. Here we further develop this approach and show how to eliminate even more complex subgraphs. We already know that computing a maximum weight cycle cover without 2- and 3-cycles is hard. What we propose instead is to find a cycle cover C' *improving on* C_{max} in the sense it does not contain certain 2- and 3-cycles from C_{max} as well as some other difficult subgraphs but possibly contains half-edges and has weight at least opt . Let us note that it is the requirement that the weight of C' is an upper bound on opt that makes the task difficult. Without it finding new cycle covers avoiding prescribed configurations is easy and we would not even have to resort to using half-edges. We believe that the method utilizing half-edges provides a handy and relatively easy way of obtaining new cycle covers (or sometimes matchings) improving on previous ones in a certain manner and having weight upper or lower bounding opt , respectively. Additionally, half-edges in such cycle covers can be either completely discarded or extended to full edges, yielding regular cycle covers. Such an approach is often substantially easier than extracting a good cycle cover from the fractional solution of an appropriate linear program. For example, note that the method of obtaining two cycle covers of weight at least $2opt$ and without any common 2-cycle in [13] is very complicated.

We deal with problematic subgraphs by either *eliminating* or *diluting* them. If C_{max} contains at least one 2-cycle or triangle, we compute a cycle cover of G that does not contain any 2-cycle or triangle that already belongs to C_{max} but may contain 2-cycles or triangles that are not in C_{max} or half-edges. Such a cycle cover C_1 is going to be called a *relaxed cycle cover* C_1 *improving* C_{max} . Also we will ensure that a computed C_1 has weight at least opt . In some cases C_1 would suffice to build a traveling salesman tour of weight at least $\frac{7}{10}opt$. To (try to) extract such a tour from C_1 and C_{max} we build a multigraph G_1 consisting of 4 copies of C_{max} and 10 copies of C_1 . Each occurrence of an edge e in C_{max} contributes 4 copies of e to G_1 and each occurrence of e in C_1 contributes 10 copies of e to G_1 . If C_1 contains only one half-edge of a certain edge e , then C_1 contributes 5 copies of e to G_1 . The number of copies of edge e in G_1 may be equal to up to 14. The total weight of edges of G_1 is at least $14opt$. We would like to divide edges of G_1 into 20 sets Z_1, \dots, Z_{20} in such a way that each Z_i ($1 \leq i \leq 20$) is a collection of vertex-disjoint paths. One of the sets Z_1, \dots, Z_{20} would then have to have weight at least $\frac{7}{10}opt$ and by patching it to a tour, we would obtain the desired solution. Dividing edges of G_1 into 20 sets can be viewed as coloring them with 20 colors so that each color class contains vertex-disjoint paths. Such coloring will also be called a *path-20-coloring* of G_1 . We can see that we are not able to path-20-color G_1 if C_1 contains a *tricky* triangle t , which is a triangle that shares an edge with a 2-cycle of C_{max} . This is because a subgraph of G_1 induced on the vertices of t contains 38 edges, 4 of which belong to an edge oppositely oriented to an edge of t . Therefore we would need 21 colors to path-color it. In the paper we show that if C_1 does not contain a tricky triangle, then we are able to color G_1 as required.

To safeguard against tricky triangles in C_1 , we introduce a technique of *diluting*, one of the main new techniques of the paper. It consists in allowing a tricky triangle t to occur in C_1 , but in a *diluted* form, by which we mean that although it contains all its edges, its weight is seemingly appropriately decreased, which enables its path-coloring. In other words, this technique succeeds (in a way) in altering the weights of edges in an unalterable (fixed) graph!

Methods of edge coloring For coloring G_1 we present a method, which we think is interesting in its own right. One of the surprisingly simple ideas on which this method is based is as follows: let S be a subset of V and $e = (u, v)$ an edge going into S (i.e. $u \notin S$ and $v \in S$), which is colored with a color k . Then if there exists no edge $e' = (u', v')$ outgoing from S (i.e. such that $u' \in S$ and $v' \notin S$) which is colored k , then e does not belong to any cycle, whose

all edges are colored k . Using this idea in an inductive way is very helpful in the process of coloring.

Coloring of multigraphs considered in this paper is also related to *the linear arboricity conjecture*, which asserts that every k -regular digraph can be path- $(k + 1)$ -colored ([21], [2]). This relationship is more visible while path-3-coloring a 2-regular 2-digraph or path-4-coloring a 3-regular digraph. Path-3-coloring a 2-digraph is a special very short case of our method of path-coloring and we obtain a method of path-4-coloring 3-regular digraphs obtained from 2 copies of one cycle cover C_o and 1 copy of another one C'_o .

We are convinced that the presented techniques will find many other applications, not only in the context of traveling salesman problems.

Previous and related results The history of approximating the problems of maximum asymmetric traveling salesman and shortest superstring is quite long as is shown by the following lists of papers [18], [7], [29], [9], [14], [3], [3], [8], [27], [13], [25], [20] and [11], [14] [5], [17], [13], [25].

Other variants of the maximum traveling salesman problem that have been considered are among others: the maximum symmetric traveling salesman problem (MAX TSP), in which the underlying graph is undirected - currently the best known approximation ratio is $\frac{4}{5}$ [10], the maximum symmetric traveling salesman problem, in which the edge weights satisfy the triangle inequality - the best approximation factor is $\frac{7}{8}$ [15], the maximum asymmetric traveling salesman problem with a triangle inequality - the best approximation ratio is $\frac{35}{44}$ [16].

2 Outline of algorithm

Suppose we have computed a maximum weight cycle cover C_{max} of a given complete directed graph $G = (V, E)$. We will say that a cycle c is **hard** if it belongs to C_{max} and each edge e of c satisfies $w(e) > \frac{3}{10}w(c)$. We are going to call cycles of length i , i.e. consisting of i edges, **i -cycles**. Also, 3-cycles will be called **triangles**. Let us notice that only 2-cycles and triangles can be hard. By $c = (v_1, v_2, \dots, v_i)$ we denote an i -cycle consisting of edges $(v_1, v_2), \dots, (v_{i-1}, v_i), (v_i, v_1)$. If C_{max} does not contain a hard cycle, then we can easily build a traveling salesman tour of weight at least $\frac{7}{10}w(C_{max}) \geq \frac{7}{10}opt$. If C_{max} contains at least one hard cycle, we would like to obtain another cycle cover C_1 , which does not contain any hard cycle from C_{max} (i.e. for each hard cycle c of C_{max} , not all edges of c are contained in C_1), has weight at least opt and enables us to build a tour of weight at least $\frac{7}{10}opt$. Let us remark here that computing a cycle cover of weight at least opt and not containing any hard cycle is hard. For comparison, note that computing a maximum weight cycle cover without any 2-cycles is NP-hard [6]. For this reason, we are going to relax the notion of a cycle cover and allow it to contain **half-edges** - a half-edge of edge (u, v) is informally speaking "half of the edge (u, v) that contains either a head or a tail of (u, v) ". We formally define half-edges and cycle covers allowing half-edges later. For now one may think of C_1 as a standard cycle cover.

To extract a tour of weight at least $\frac{7}{10}opt$ from C_{max} and C_1 , we are going to build a multigraph G_1 consisting of 4 copies of C_{max} and 10 copies of C_1 . More precisely, G_1 contains 4 copies of each edge $e \in C_{max} \setminus C_1$, 10 copies of each $e \in C_1 \setminus C_{max}$ and 14 copies of each $e \in C_1 \cap C_{max}$. We would like to color each edge of G_1 with one of 20 colors so that edges of the same color form a collection of disjoint paths or, in other words, we would like to **path-20-color** G_1 or **path-color it with 20 colors**. We may notice that it is not possible, if C_1 contains one of the following:

1. a 2-cycle or triangle of C_{max} .

2. a triangle oppositely oriented to a triangle of C_{max} .
3. a 2-cycle $c = (u, v)$ such that one of its edges belongs to C_{max} . This is because G_1 contains in this case 24 edges connecting u and v (14 in one direction and 10 in the other) and thus we would need 24 colors to path-color G_1 .
4. a triangle $t = (p, q, r)$ such that a 2-cycle (q, r) belongs to C_{max} . In this case G_1 contains a subgraph consisting of 14 copies of (q, r) , 10 copies of each of (p, q) , (r, p) and 4 copies of (r, q) , which is clearly non-path-20-colorable.

We later show that if C_1 does not contain any of the above cycles, then G_1 built from C_{max} and C_1 in the manner described above is always path-20-colorable. Ideally, we would like the enumerated cycles not to occur in C_1 at all. However, not all of them are bad for our purposes, because sometimes it is easy to replace some edges of these cycles with other ones, so that we obtain a path-20-colorable multigraph. For example, if C_1 contains a triangle $t = (p, q, r)$ such that a 2-cycle (q, r) belongs to C_{max} and $w(r, q) \leq \frac{3}{2}w(q, r)$, we can replace 4 copies of (r, q) with 6 copies of (q, r) and make the subgraph on p, q, r path-20-colorable. Also, this way we do not diminish the overall weight of the subgraph.

Below we define a set of cycles that are **tricky**. The occurrence of any such cycle c in C_1 means that G_1 is non-path-20-colorable and we cannot remedy this by local replacements of edges. A cycle of G oppositely oriented to c is denoted as $opp(c)$. A cycle c' is said to be a **subcycle** of c if every vertex of c' belongs to c . For any multisubgraph G' of G , by $mult_{G'}(e)$ we denote the number of copies of e occurring in G' and for any subset V' of vertices of G' by $E_{G'}(V')$ we denote the set of edges of G' connecting any two vertices of V' .

Let $S = (V_S, E_S)$ be a multisubgraph of G . For any $v \in V_S$, by $indeg_S(v)$, $outdeg_S(v)$ we denote, respectively, the indegree and outdegree of v in S . Let G_1/S denote a multigraph $(G_1 \setminus E_{G_1}(S)) \cup E_S$. We say that a multisubgraph S of G is **amenable** if (i) any path-20-coloring of $G_1 \setminus S$ can be extended to path-20-coloring of G_1/S and (ii) every vertex $v \in V_S$ satisfies $indeg_{G_1/S}(v) \leq 17$ or $outdeg_{G_1/S}(v) \leq 17$. (The degrees are required to satisfy this condition, because we want to leave the possibility of adding 3 copies of some edge $e \in G$ incident to any vertex v to the multigraph G_1/S .)

We define a vertex/edge surrounding of c (with respect to C_{max}) denoted $sur^v(c)$ and $sur^e(c)$, respectively. For a triangle $t = (p, q, r) \in C_{max}$ we have $sur^v(t) = \{p, q, r\}$ and $sur^e(t) = \{(p, q), (q, r), (r, p)\}$. For a triangle $t = (p, q, r)$ such that the 2-cycle (q, r) belongs to C_{max} , we have $sur^v(t) = \{p, q, r\}$ and $sur^e(t) = \{(q, r), (r, q)\}$. Let c be a 2-cycle (u, v) and $(u', u), (v, v')$ two edges of C_{max} . Then $sur^v(c) = \{u', u, v, v'\}$ and $sur^e(c) = \{(u', u), (u, v), (v, v')\}$.

A cycle c is **tricky** if it belongs to type 1, 3 or 4 enumerated above and no amenable subgraph on $sur^v(c)$ has weight at least $10w(c) + 4w(sur^e(c))$. A triangle t that belongs to C_{max} is called a **3-triangle** and a triangle of type 4 - a **2-triangle**.

We later prove that any two tricky 2-cycles are vertex-disjoint. We observe also that for any tricky 3-triangle t holds that if t is not vertex-disjoint with some other tricky cycle c , then c is a sub-2-cycle of t . On the other hand tricky 2-triangles do not even have to be edge-disjoint with other tricky 2-triangles. We require that C_1 does not contain any tricky 2-cycle or tricky 3-triangle or a triangle oppositely oriented to a tricky 3-triangle. As for tricky 2-triangles, we are going to forbid only their subset in C_1 .

Let t be a tricky 2-triangle $t = (p, q, r)$ such that $c = (q, r)$ is a 2-cycle of C_{max} . We call p its **t-point** and the 2-cycle (q, r) its **t-cycle**. We have already observed that $w(r, q) > \frac{3}{2}w(q, r)$, because otherwise we could take 4 copies of c , 10 copies of t and replace in it 4 copies of $w(r, q)$

with 6 copies of (q, r) , obtaining thus an amenable subgraph. Let $\Delta(c) = w(r, q) - \frac{3}{2}w(q, r)$. To the 2-cycle c we assign weight $w'(c) = w(q, r) + \Delta(c)$. Also, by $\kappa(t)$ we denote $\frac{w(r, q)}{10}$. Among the set of all tricky 2-triangles we are going to distinguish a set R of its representatives and require that C_1 does not contain any tricky triangle from R or if it does, then every such triangle t is *diluted*, by which we mean that in C_1 it has weight equal to $w(t) - \kappa(t)$. We explain below how this is possible.

To identify the set R , we construct a bipartite graph $H = (C \cup P, E_t)$, where C contains all t-cycles and P t-points. An edge (c, p) belongs to E_t iff there exists a tricky triangle t such that c is its t-cycle and p its t-point. Let $T_1 \cup T_2 \cup \dots \cup T_k$ be a partition of the set of vertices of C such that t-cycles belonging to the same T_i have equal weight w' and for each $i < j$ and any $c_i \in T_i, c_j \in T_j$, it holds $w'(c_i) > w'(c_j)$. We assign *ranks* to edges of H in the following manner. Any edge of E_t incident to a vertex of T_i has rank i . We are going to compute a *rank-maximal* matching N of H , which is a matching of H containing a maximum number of rank one edges and subject to this condition a maximum number of rank two edges and so on. A rank-maximal matching can be computed in polynomial time [12].

One can observe that

Fact 1 *Any rank-maximal matching of H is a maximum matching of H .*

As the set R representing tricky 2-triangles we set tricky triangles corresponding to the edges of N .

Let $t = (p, q, r)$ be a tricky 2-triangle with a t-cycle $c = (q, r)$. Observe that if C_1 contained a diluted t , i.e. with weight in C_1 decreased by $\kappa(t)$, then $10(w(t) - \kappa(t)) + 4w(c)$ has the same weight as 14 copies of (q, r) , 10 copies of each of (p, q) , (r, p) and 3 copies of (r, q) , which forms a path-20-colorable subgraph on p, q, r . Hence, a diluted triangle of R can be allowed in C_1 .

To be able to compute a cycle cover C_1 of weight at least *opt* and which does not contain any problematic cycle, we are going to allow it to contain *half-edges*, defined as follows. Let $\tilde{G} = (\tilde{V}, \tilde{E})$ be a graph obtained from G by splitting each edge $(u, v) \in E$ with a vertex $x_{(u, v)}$ into two edges $(u, x_{(u, v)})$ and $(x_{(u, v)}, v)$ having weights such that $w(u, x_{(u, v)}) + w(x_{(u, v)}, v) = w(u, v)$. Each of the edges $(u, x_{(u, v)}), (x_{(u, v)}, v)$ is called *a half-edge (of (u, v))*. By saying that an edge (u, v) of G belongs to a subset $\tilde{C} \subseteq \tilde{E}$, we will mean that both half-edges of (u, v) belong to \tilde{C} .

We say that $\tilde{C} \subseteq \tilde{E}$ does not contain a cycle c of G , if \tilde{C} does not contain all edges of c , i.e., there exists at least one edge e of c such that at least one half-edge of e does not belong to \tilde{C} .

To deal with tricky triangles from the set R , we need to further extend the graph \tilde{G} . For each tricky triangle $t \in R$, we add two new vertices v_t, v'_t and two loops: e_t incident to v_t and e'_t incident to v'_t with weights $w(e_t) = -\kappa(t), w(e'_t) = \kappa(t)$. We call this graph $\hat{G} = (\hat{V}, \hat{E})$. (Note that this is a supergraph of \tilde{G} . The idea behind these new loops is as follows. For each tricky triangle $t = (p, q, r) \in R$, C_1 either does not contain t or it does contain t and also a loop e_t . This implies that the weight of such t in C_1 can be viewed as though it were equal to $w(t) - \kappa(t)$, i.e., it means that t is diluted, which enables the coloring of the subgraph on p, q, r . By saying that C_1 contains a diluted t , we mean that C_1 contains t and also a loop e_t .

Definition 1 *A relaxed cycle cover improving C_{max} is a subset $\hat{C} \subseteq \hat{E}$ such that*

- (i) *each vertex in V has exactly one outgoing and one incoming half-edge in \hat{C} ;*
- (ii) *for any tricky 2-cycle or 3-triangle c , \hat{C} does not contain c or $opp(c)$.*

- (iii) for any tricky 2-triangle $t \in R$, \hat{C} either does not contain t or contains a diluted t .
- (iv) if \hat{C} contains only one half-edge of edge (u, v) , then (u, v) belongs to a tricky 2-cycle, 3-triangle or 2-triangle of R or to a triangle oppositely oriented to a tricky 3-triangle.

A relaxed cycle cover C improving C_{max} , or a relaxed cycle cover C for short, consists of directed cycles and/or directed paths. A directed cycle of C corresponds to a directed cycle of the original graph G and a directed path ends and begins with a vertex in $\tilde{V} \setminus V$.

The outline of a $\frac{7}{10}$ -approximation algorithm for Max ATSP is as follows.

1. Compute a maximum weight cycle cover C_{max} of G .
2. If C_{max} does not contain a hard cycle, extract from C_{max} a set \mathcal{P} of vertex-disjoint paths of weight at least $\frac{7}{10}w(C_{max})$ and go to Step 6.
3. Compute a relaxed cycle cover C_1 improving C_{max} with weight $w(C_1) \geq opt$.
4. Compute a multigraph G_1 with weight $w(G_1) \geq 4w(C_{max}) + 10w(C_1)$ and path-20-color G_1 omitting non-path-20-colorable subgraphs. If the whole G_1 is path-colored, go to Step 6.
5. Compute exchange sets E_1, F_1 such that $G_2 = G_1 \setminus E_1 \cup F_1$ is path-20-colorable. Extend the existing coloring of G_1 to that of G_2 .
6. Extend a set \mathcal{P} of vertex-disjoint paths of weight at least $\frac{7}{10}opt$ to a tour of G .

3 Computation of a relaxed cycle cover C_1

To compute a relaxed cycle cover C_1 improving C_{max} we construct the following undirected graph $G' = (V', E')$. For each vertex v of G we add two vertices v_{in}, v_{out} to V' . For each edge (u, v) that belongs to a tricky 2-cycle, 3-triangle or 2-triangle of R we add vertices e_{uv}^1, e_{uv}^2 , called **subdivision vertices** of (u, v) , an edge (e_{uv}^1, e_{uv}^2) of weight 0 and edges $(u_{out}, e_{uv}^1), (v_{in}, e_{uv}^2)$ having weights such that $w(u_{out}, e_{uv}^1) + w(v_{in}, e_{uv}^2) = w(u, v)$. Edges $(u_{out}, e_{uv}^1), (v_{in}, e_{uv}^2)$ are also called **half-edges** of (u, v) . If (u, v) does not belong to t or $opp(t)$ such that t is a tricky 3-triangle, then each of the half-edges of (u, v) gets weight $\frac{1}{2}w(u, v)$. For every other edge $(u, v) \in E$ we add an edge (u_{out}, v_{in}) of weight $w(u, v)$.

Next we build so-called gadgets. For each tricky 2-cycle c on vertices u and v , we add vertices γ_u^c and γ_v^c and edges $(\gamma_u^c, e_{uv}^1), (\gamma_u^c, e_{vu}^2), (\gamma_v^c, e_{vu}^1), (\gamma_v^c, e_{uv}^2)$ with weight 0. The gadget is shown in Figure 1. If c is not a subcycle of a tricky 3-triangle or 2-triangle of R , each of the half-edges of (u, v) gets weight $\frac{1}{2}w(u, v)$ and each of the half-edges of (v, u) gets weight $\frac{1}{2}w(v, u)$.




Figure 1: A gadget for a 2-cycle (q, r) .

Let t be any tricky 3-triangle $t = (p, q, r)$. Among edges of $opp(t)$ we choose one with maximum weight. Suppose that the chosen edge is (r, q) . For each such t , we build the following gadget. We add vertices $\gamma_p^{t-}, \gamma_p^{t+}$ and connect them to vertices e_{qp}^2, e_{rp}^2 and e_{qp}^1, e_{rp}^1 , respectively, via edges of weight 0. Each of the edges $(e_{qp}^2, p_{in}), (e_{rp}^2, p_{in})$ gets weight $\frac{1}{2} \max\{w(q, p), w(r, p)\}$. Thus, $w(q_{out}, e_{qp}^1) = w(q, p) - \frac{1}{2} \max\{w(q, p), w(r, p)\}$, $w(r_{out}, e_{rp}^1) = w(r, p) - \frac{1}{2} \max\{w(q, p), w(r, p)\}$. We proceed analogously for pairs of edges $(r, q), (p, q)$ and $(p, r), (q, r)$. Thus, we add vertices $\gamma_q^{t-}, \gamma_q^{t+}$ and connect them to vertices e_{pq}^2, e_{rq}^2 and e_{pq}^1, e_{rq}^1 , respectively, via edges of weight 0, and we add vertices $\gamma_r^{t-}, \gamma_r^{t+}$ and connect them to vertices e_{pr}^2, e_{qr}^2 and e_{pr}^1, e_{qr}^1 , respectively, via edges of weight 0. Each of the edges $(e_{pq}^2, q_{in}), (e_{rq}^2, q_{in})$ gets weight $\frac{1}{2} \max\{w(p, q), w(r, q)\}$ and each of the edges $(e_{pr}^2, r_{in}), (e_{qr}^2, r_{in})$ gets weight $\frac{1}{2} \max\{w(p, r), w(q, r)\}$. Additionally, if the 2-cycle $c = (r, q)$ is not tricky, we add a gadget for c , which is the same as the gadget for a tricky 2-cycle.

The gadget is depicted in Figure 2.

We say that a half-edge is **incoming** if it is a half-edge of some edge (u, v) incident to v . A half-edge of (u, v) incident to u is called **outgoing**.

Let e_1, e_2 denote two different edges of G incident with the same vertex v . Assume that a relaxed cycle cover \tilde{C} contains exactly one half-edges of each of e_1, e_2 . We say that these two half-edges are **crossing** if exactly one of them is incident to v and **non-crossing** otherwise.

A *quasi relaxed cycle cover* denotes a relaxed cycle cover that does not satisfy point (iii) of Definition 1. We say that a (quasi) relaxed cycle cover \tilde{C} is **non-integral** on a set S of edges if there exists some edge $e \in S$ such that \tilde{C} contains only one half-edge of e .

We say that a half-edge e_h of \tilde{C} is within a set of edges $F \subseteq E$ if e_h is a half-edge of some edge in F . Let $w(\tilde{C})_t$ denote the total weight of half-edges of \tilde{C} within $t \cup opp(t)$.

Definition 2 A tricky triangle $t = (p, q, r)$ is said to be **harmonious** in a relaxed cycle cover \tilde{C} if \tilde{C} satisfies the following:

1. The difference between the numbers of edges of \tilde{C} incoming to t and outgoing of t , denoted $dif(t)$, is either zero or two.
2. If $dif(t) = 0$, then \tilde{C} is integral on t .
3. If $dif(t) = 2$, then depending on the configuration of edges incoming to and outgoing of t , $10w(\tilde{C})_t + 4w(t)$ is upper bounded by:
 - $\max\{10w(t) - 5w(r, p), 10(w(p, q) + w(q, p))\}$, if \tilde{C} contains one edge outgoing of t , incident to r and three edges incoming to t ,
 - $\max\{10w(t) - 5w(q, r), 10(w(p, q) + w(q, p))\}$, if \tilde{C} contains one incoming to t incident to r and three edges outgoing of t ,
 - $\max\{10w(t) + 5w(r, p), 10(w(p, r) + w(p, q) + w(q, p)), 10(w(p, q) + w(p, r) + w(r, p))\}$, if \tilde{C} contains two edges outgoing of t , incident to p and q and no edge incoming to t ,
 - $\max\{10w(t) + 5w(q, r), 10(w(q, r) + w(r, p) + w(p, r)), 10(w(p, r) + w(r, q) + w(q, r))\}$, if \tilde{C} contains two edges incoming to t incident to p and q and no edge outgoing of t .

Lemma 1 Any perfect matching of G' yields a quasi relaxed cycle cover \tilde{C} with the following properties:




Figure 2: A gadget for a triangle $t = (p, q, r)$.

- (i) for each problematic 2-cycle (u, v) , if \tilde{C} contains two half-edges from $\{(u, x_{(u,v)}), (x_{(u,v)}, v), (v, x_{(v,u)}), (x_{(v,u)}, u)\}$, then they either belong to the same edge or are crossing - thus one of them is incident with u and the other with v and are either both incoming or both outgoing.
- (ii) for each tricky 3-triangle $t = (p, q, r)$, t is harmonious in \tilde{C} .

The proof is in Section 5.

We construct an undirected graph G'' by extending and modifying G' as follows. For each tricky triangle $t = (p, q, r)$ such that (q, r) is a 2-cycle of C_{max} , we add the following gadget. We add vertices $a_{\{p,q,r\}}, b_{\{p,q,r\}}$ and connect them to vertices e_{pq}^1, e_{rp}^2 and e_{pq}^2, e_{rp}^1 , respectively, via edges of weight $\frac{\kappa(t)}{2}$ and connect $a_{\{p,q,r\}}$ and $b_{\{p,q,r\}}$ via an edge of weight 0. We also decrease the weight of each of the edges $(r_{out}, e_{rq}^1), (e_{rq}^2, q_{in}), (q_{out}, e_{qr}^1), (e_{qr}^2, r_{in})$ by $\frac{\kappa(t)}{2}$.

Lemma 2 Any perfect matching of G'' yields a relaxed cycle cover \hat{C} with the following properties. Let $t = (p, q, r)$ be a tricky triangle of R , where $c = (q, r)$ is its t -cycle.

- (i) If \hat{C} is non-integral on $t \cup c$, then it contains (i) crossing half-edges within c or (ii) crossing half-edges within $\{(r, p), (p, q)\}$.
- (ii) \hat{C} contains all edges of t if and only if it also contains a loop incident to v_t .
- (iii) If \hat{C} contains a loop incident to v_t' , then it contains no half-edges within c .

Proof. Let M be a perfect matching of G'' . If M does not contain the edge $(a_{\{p,q,r\}}, b_{\{p,q,r\}})$ (which means that these vertices are matched via edges of weight $\frac{\kappa(t)}{2}$) and does not contain any edge corresponding to a half-edge of an edge of c , then \hat{C} contains the loop e_t' with weight $\kappa(t)$. If M contains edges corresponding to all half-edges within t , then \hat{C} contains all edges of t as well as the loop e_t with weight $-\kappa(t)$. In other respects the proof is similar to that of Lemma 1. \square

Theorem 2 Any perfect matching of G'' yields a relaxed cycle cover C_1 improving C_{max} . A maximum weight perfect matching of G'' yields a relaxed cycle cover C_1 improving C_{max} such that $w(C_1) \geq opt$.

Proof. The first statement follows from the preceding two lemmas. The second statement follows from the fact that a traveling salesman tour is also a cycle cover that does not contain any 2-cycles or triangles unless the whole graph has two or three vertices. \square

4 Path-coloring

Once we have computed a maximum weight cycle cover C_{max} and a relaxed cycle cover C_1 our next task is to construct and color a multigraph G_1 . The constructed multigraph is required to have weight at least $4w(C_{max})+10w(C_1)$ and be path-20-colorable. On the high level, to satisfy the first requirement, we build G_1 by taking 4 copies of C_{max} and 10 copies of C_1 , obtaining possibly a multigraph with non-path-20-colorable subgraphs. To remedy the multigraph, we replace certain edges of such non-path-20-colorable subgraphs with other ones in a way that preserves the required weight and makes the multigraph G_1 path-20-colorable or facilitates the coloring. The precise construction is described below.

4.1 Preprocessing via alternating cycles

The preprocessing described below is needed for coloring 2-cycles of C_1 (the proof of Lemma 11) and it can be skipped during the first reading.

Before building the multigraph G_1 we modify C_1 so that it differs from C_{max} in a minimal way. An **alternating cycle** in $C_{max} \oplus C_1$ is a sequence of edges of the form $(v_1, v_2), (v_3, v_2), (v_3, v_4), (v_5, v_4), \dots, (v_{k-1}, v_k), (v_1, v_k)$, in which edges belong alternately to $C_{max} \setminus C_1$ and $C_1 \setminus C_{max}$. By applying an alternating cycle C_{alt} to C_1 we mean the operation, whose result is $C_1 \oplus C_{alt}$, in which we treat C_{alt} as a set of edges. An alternating cycle C_{alt} is **good** if $C'_1 = C_1 \oplus C_{alt}$ does not contain a problematic cycle or a tricky triangle of R , i.e., C'_1 is a relaxed cycle cover improving C_{max} .

Fact 2 *Let C_{alt} be a good alternating cycle and $C'_1 = C_1 \oplus C_{alt}$. Then $w(C'_1) \geq w(C_1)$.*

Proof. Since C_{max} is a maximum weight cycle cover of G , $w(C_{max} \oplus C_{alt}) \leq w(C_{max})$. Therefore, $w(C'_1) \geq w(C_1)$. \square

We apply good alternating cycles to C_1 until it is no longer possible. We still call a new relaxed cycle cover C_1 .

Lemma 3 *After preprocessing, it holds that no alternating cycle is good.*

4.2 Construction of G_1

In this section we assume that G does not contain any tricky triangles and that C_1 does not contain any **strange 2-cycles** - a 2-cycle is said to be strange if exactly one of its edges belongs to C_{max} . This in particular means that each half-edge of C_1 belongs to a tricky 2-cycle. If C_1 contains exactly one half-edge of each edge of a 2-cycle c of G , then c is called a **halfy 2-cycle** of C_1 . In this section we also assume that each halfy 2-cycle c of C_1 belongs to C_{max} .

We start the construction of G_1 by taking 4 copies of C_{max} and 10 copies of C_1 , by which we mean the following. Let $mult(e)$ denote the number of copies of edge $e \in G$ contained in G_1 . At the beginning for each edge $e \in G$, we set $mult(e) = 0$. Next, for each $e \in C_{max}$, we increase $mult(e)$ by 4 and further, for each $e \in C_1$ (note that $e \in C_1$ means that the whole edge

e belongs to C_1), we increase $\text{mult}(e)$ by 10. Subsequently, for each e such that C_1 contains only a half-edge of e , we increase $\text{mult}(e)$ by 5. Clearly, the thus obtained G_1 has weight equal to exactly $4w(C_{max}) + 10w(C_1)$.

4.3 Path-coloring

Let \mathcal{K} denote $\{i \in N : 1 \leq i \leq 20\}$. To **path-20-color** a multigraph G_1 , or to **path-color** it, means to assign a color of \mathcal{K} to each edge of G_1 in such a way that each color class consists of vertex-disjoint paths. Equivalently, we will be interested in path-coloring the underlying simple graph G_1 , in which to each edge e of G_1 we will assign a subset $\text{col}(e)$ of colors of \mathcal{K} such that the size of $\text{col}(e)$ equals the number of copies of e in the multigraph G_1 , i.e., $|\text{col}(e)| = \text{mult}(e)$ (and each color class consists of vertex-disjoint paths).

A path-coloring of G_1 will be carried out gradually. In the process each edge e of G_1 can be either **colored** - when it has $\text{mult}(e)$ colors assigned to it, or **uncolored** - when it is assigned no color. A cycle c is called **monochromatic** if there exists a color i of \mathcal{K} such that each edge of c is colored with i - c is then a monochromatic cycle of color i . Of course, a (partially) path-colored G_1 cannot contain a monochromatic cycle. We will say that an edge e is **safe** if no matter how we color the so far uncolored edges, it is guaranteed not to belong to any monochromatic cycle. For example suppose that u has three incident edges in G_1 - $e_1 = (u, v), e_2 = (z, u), e_3 = (z', u)$ such that $\text{mult}(e_1) = \text{mult}(e_2) = 4, \text{mult}(e_3) = 10$ and $\text{col}(e_1) = \{1, 2, 3, 4\}, \text{col}(e_2) = \{5, 6, 7, 8\}$ and $\text{col}(e_3) = \{11, \dots, 20\}$. Also, u has no other outgoing edge in G_1 . Then, clearly, e_1 is safe. By saying that an edge e is k -safe we will mean that e is guaranteed not to belong to a monochromatic cycle of color k .

If S denotes any subset of vertices of G_1 , then S^+ denotes a set of edges $\{(u, v) \in G_1 : u \in S, v \notin S\}$ and analogously, $S^- = \{(u, v) \in G_1 : u \notin S, v \in S\}$.

In path-coloring G_1 we are going to heavily use the following very helpful observation:

Observation 1 *Suppose that edge $e \in S^-$ is colored with k and no edge of S^+ is uncolored or colored with k . Then e is k -safe. Analogously, if edge $e \in S^+$ is colored with k and no edge of S^- is uncolored or colored with k , then e is k -safe.*

Recall that C_1 consists of cycles and paths. Any path p of C_1 ends with a half-edge of some edge e . Such edge e is called a **border** of p . All paths of C_1 occurring in this section end with borders belonging to tricky 2-cycles of C_{max} . Notice that each halfy 2-cycle of C_1 either has two incoming paths of C_1 or two outgoing paths of C_1 . Apart from borders, we distinguish two other types of edges of C_{max} . An edge $e = (u, v) \in C_{max}$ that is not a border is called a **ray** if u and v belong to two different cycles of C_1 or two different paths of C_1 or one of them belongs to a path of C_1 and the other to a cycle of C_1 . Otherwise, it is called a **chord**. Note that a chord e may also belong to C_1 . A ray $r = (u, v)$ incident to a vertex on a cycle c or path p of C_1 is said to be a ray of c or correspondingly p . If vertex v belongs to e , then r is said to be an **inray** of e (and c or p). Otherwise, it is called its **outray**.

Using Observation 1 we can apply the following simple method of coloring rays of C_1 .

Lemma 4 *Let c be a cycle of C_1 such that each of its incident rays is uncolored or safe. Then we are able to color all uncolored rays of c in such a way that each one of them is safe.*

Proof. It is easy to guarantee that each newly colored ray is safe - it suffices if we color inrays and outrays of c with disjoint sets of colors, i.e., we partition \mathcal{K} into $Z^-(c)$ and $Z^+(c)$ and each uncolored inray of c is colored with colors of $Z^-(c)$ and each uncolored outray of c with colors

of $Z^+(c)$. Then by Observation 1 and the fact that each previously colored ray is already safe, each ray of c is safe. \square

For paths of C_1 we can in fact apply the same method:

Lemma 5 *Let p be a cycle of C_1 such that each of its incident rays is uncolored or safe. Then we are able to color all uncolored rays of p in such a way that each one of them is safe.*

Proof. We use the same method as in the lemma above. Rays are not the only outgoing/incoming edges of p . There are also borders. However, the only cycle any border belongs to is a halfy 2-cycle and any 2-cycle consists of two borders (of two different paths of C_1). Thus by this observation, Observation 1 and the fact that each previously colored ray is already safe, each ray of p is safe. \square

Coloring rays so that they are safe does not mean, however, that there always exists a possibility of coloring the remaining edges of G_1 so that we do not create a monochromatic cycle. Let us consider a few examples.

If $c = (p, q, r, s)$ is a 4-cycle of C_1 with 4 inrays, each colored with $\{1, 2, 3, 4\}$ and 4 outrays, each colored with $\{5, 6, 7, 8\}$, then the only colors we can use on any edge of c are those belonging to $Z = \mathcal{K} \setminus \{1, 2, \dots, 8\}$. Any color of Z can be used on at most three edges of c and each edge of c has to be assigned 10 different colors. Thus we would need at least $40/3 > 13$ different colors, but have only 12. Therefore, it is not possible to path-color c .

We can notice that, if instead of a 4-cycle we had a 6-cycle c with 6 inrays, each colored with $\{1, 2, 3, 4\}$ and 6 outrays, each colored with $\{5, 6, 7, 8\}$, then we would be able to path-color c . Suppose now that we have a 4-cycle $c = (p, q, r, s)$ the same as above except for the fact that one of its outrays is colored with $\{1, 2, 3, 4\}$. We of course assume that all rays are safe. It turns out that in this case we can path-color c , because one edge of c can be colored with colors of $\{5, 6, 7, 8\}$ and then we need $(40 - 4)/3 = 12$ colors for the rest.

Below we define *blocked* cycles of C_1 and prove that any cycle of C_1 that is not blocked can be path-20-colored.

Two edges e_1, e_2 are said to be *coincident* if there exists vertex v such that either $e_1 = (v, v_1), e_2 = (v, v_2)$ or $e_1 = (v_1, v), e_2 = (v_2, v)$. We say that two edges e_1, e_2 are *diverse* if $col(e_1) \cap col(e_2) = \emptyset$. Let us note that coincident edges must be diverse.

We define the flexibility of $e = (u, v) \in C_1$, denoted $flex(e)$, as follows. Let $e_1 = (u, v'), e_2 = (u', v)$ be edges of C_{max} coincident with e (it is possible that $e_1 = e_2$ as well as one or both of e_1, e_2 do not exist because they have been removed during the modification of G_1). Then $flex(e) = 10 - |col(e_1) \cup col(e_2)|$. Thus, if e_1 and e_2 are colored with two non-empty disjoint sets of colors, then $flex(e) = 2$. For each cycle c of C_1 we define its flexibility $flex(c)$ and colorfulness $kol(c)$. The flexibility of c is defined as $flex(c) = \sum_{e \in c} flex(e)$. Colorfulness $kol(c)$ denotes the number of colors of \mathcal{K} used so far for coloring the edges of G_1 incident to c . For a subset E' of edges of E by $mult(E')$ we denote $\sum_{e \in E'} mult(e)$. By $\lambda(c)$ we denote the length of a cycle c . For a cycle c with at least one chord, $chor(c) = 4$; a cycle c with no chords has $chor(c) = 0$. Using the above notions we define the characteristic $\chi(c)$ of a cycle c of C_1 as follows. If c has (i) at least two chords or (ii) one chord and $\lambda(c) > 2$, then $\chi(c) = 20$. Otherwise, $\chi(c) = flex(c) + kol(c) - chor(c)$. A cycle c of C_1 is said to be *blocked* if $\chi(c) < 20$ and *unblocked* otherwise.

Lemma 6 *Let c be a cycle of C_1 that is not blocked and such that each of its incident rays is colored and safe. Then we are able to color all edges and chords of c in such a way that each one of them is safe.*

Similarly, as cycles of C_1 may be blocked, paths of C_1 can become non-path-20-colorable too. Or, more precisely, halfy 2-cycles of C_1 can become blocked. Let $p_1 = (u_1, \dots, u_k)$ denote a path of C_1 . Then both u_1 and u_k belong to two different halfy 2-cycles $c_1 = (u_1, v_1)$ and $c_2 = (u_k, w_l)$ of C_1 . Thus C_1 contains also paths $p_2 = (v_1, \dots, v_{k'})$ and $p_3 = (w_1, \dots, w_l)$, though it may happen that $p_2 = p_3$. If p_1 consists of more than one edge, then C_{max} contains edges $a_1 = (u'_2, u_2), a' = (u_{k-1}, u'')$, none of which is a border. Each of these edges is called an **antenna** (of p_1). a_1 is also said to be an antenna of c_1 and a' of c_2 .




Figure 3: Antennas a_1, a_2 of a halfy 2-cycle (u_1, v_1) .

Fact 3 *Let $c = (u_1, v_1)$ be a halfy 2-cycle of C_1 with two antennas a_1, a_2 . Then, in any path-coloring of G_1 the antennas a_1 and a_2 have to be diverse.*

Proof. Suppose that C_1 contains paths $p_1 = (u_1, u_2, \dots, u_k)$ and $p_2 = (v_1, v_2, \dots, v_l)$. Then the antennas a_1 and a_2 have the form $a_1 = (u'_2, u_2)$ and $a_2 = (v'_2, v_2)$. We know that $mult(a_1) = mult(a_2) = 4$, $mult(u_1, u_2) = mult(v_1, v_2) = mult(u_1, v_1) = mult(v_1, u_1) = 10$. The situation is depicted in Figure 3. Since c is a 2-cycle, its edges have to be diverse. Also, we may notice that (u_1, u_2) has to be colored in the same way as (v_1, u_1) and (v_1, v_2) in the same way as (u_1, v_1) . Therefore (u_1, u_2) and (v_1, v_2) have to be diverse. Also $col(u_1, u_2) \cup col(v_1, v_2) = \mathcal{K}$. Since a_1 and (u_1, u_2) have to be diverse and so do a_2 and (v_1, v_2) , a_1 and a_2 have to be diverse as well. \square

If a_1, a_2 are two antennas of a 2-cycle c and a_1 is already colored but a_2 not, then we say that colors of $col(a_1)$ are **forbidden** on a_2 . A halfy 2-cycle c of C_1 is said to be **blocked**, if it has two antennas and they are not diverse. The multigraph G_1 is **blocked** if at least one cycle or halfy 2-cycle of C_1 is blocked. The multigraph G_1 is **safe** if each of its colored edges is safe.

We say that a cycle or path of C_1 is **unprocessed** if at least one of its rays is uncolored. To **process** a cycle/path of C_1 means to color its rays so that each of them is safe and G_1 is not blocked, assuming that before starting to process this cycle or path, G_1 is safe and unblocked.

We are now ready to state the algorithm for path-20-coloring G_1 .

Algorithm Color7

while there exists an unprocessed cycle of C_1
 $c \leftarrow$ an unprocessed cycle of C_1 with a minimal number of uncolored rays;
 process c ;

while there exists an unprocessed path of C_1
 $p \leftarrow$ any unprocessed path of C_1 ;
 process p ;

color the remaining uncolored edges in such a way that each of them is safe;

In what follows we prove the correctness of Algorithm Color7.

Let B denote the set of uncolored edges of G_1 . We divide the flexibility of each edge e having coincident edges $e_1, e_2 \in C_{max}$ using three components $flex^0(e) = 10 - mult(e_1) - mult(e_2)$, $flex^+(e) = |col(e_1) \cap col(e_2)|$ and $blank(e) = mult(e_1 \cap B) + mult(e_2 \cap B)$. Thus $flex(e) = flex^0(e) + flex^+(e) + blank(e)$. As a result the flexibility of each cycle c of C_1 consists of three components as well - $flex(c) = flex^0(c) + flex^+(c) + blank(c)$.

We say that two rays $r_1 = (u, u'), r_2 = (v', v)$ of a 2-cycle c are **complementary (on c)** if either (u, v) or (v', u') is an edge of c .

Lemma 7 *Let c be a cycle of C_1 with u incident uncolored edges of C_{max} . Assume that c has (i) no chords or (ii) one chord and $\lambda(c) = 3$. Then $\chi(c) = kol(c) + flex^0(c) + flex^+(c) + blank(c) - chor(c) \geq kol(c) + 2\lambda(c) + 4u + flex^+(c)$.*

As a consequence:

1. A cycle c of length $\lambda(c) > 2$ can be blocked only if it has at most one uncolored ray.
2. A 2-cycle is blocked only if some two of its non-complementary rays r_1, r_2 are not diverse.

Proof. Any chord e of c contributes $2mult(e)$ to $blank(c)$. Therefore, $blank(c) - chor(c) = 4u$. For any edge e of c , it holds $flex^0(e) = 2$. The claim follows. \square

Lemma 8 *Let r_1, r_2 be two edges of C_{max} coincident with an edge e belonging to a cycle c of C_1 . Suppose also that r_1 is uncolored. There exists a set $Z \subseteq \mathcal{K}$ of colors, the application of any color of which on r_1 increases $kol(c) + flex^+(c)$ by one, i.e., $kol(c) + flex^+(c)$ increases by $|col(r_1) \cap Z|$ after coloring r_1 . If r_2 is uncolored, then Z has $20 - kol(c)$ elements. Otherwise, Z is of size $20 - kol(c) + mult(r_2)$.*

Proof. By coloring r_1 with a color not occurring yet on the rays of c , we increase $kol(c)$ by 1. There are $20 - kol(c)$ such colors. Additionally, if r_2 is already colored, then by coloring r_1 with any color assigned to r_2 , we increase $flex^+(e)$ and thus also $flex^+(c)$. \square

For a cycle c of C_1 , $\rho(c)$ indicates the maximum multiplicity of r_2 from Lemma 8, i.e., the maximum multiplicity of a colored ray r_2 of c incident to $e \in c$ such that the other edge r_1 of C_{max} incident to e is an uncolored ray of c .

Lemma 9 *Suppose that at step S we want to color a set U of uncolored edges, where U consists of either (i) a subset of uncolored rays of a cycle c of C_1 or (ii) an antenna of a halfy 2-cycle c of C_1 . Then, assuming that G_1 is unblocked, there always exists a number $\Delta'(c)$ and a set $Z \subseteq \mathcal{K}$ such that by using $\Delta'(c)$ different colors of Z on U , we guarantee that c does not become blocked. Depending on additional conditions, $\Delta'(c)$ and $|Z|$ can be expressed as the following functions of a certain $\Delta(c) \leq \Delta'(c)$:*

0 . If c has at least two chords or one chord and $\lambda(c) > 3$, then $\Delta'(c) = 0$. In the remaining points we assume that c has no chords or one chord and $\lambda(c) = 3$.

1. If c is a 2-cycle with r colored rays, then $\Delta'(c) = \text{mult}(U)$ and $|Z| = 20 - 4r + \rho(c)$.
2. If c has one uncolored ray, no chords and $\lambda(c) > 2$, then $\Delta'(c) = 4 - \Delta(c) \geq 0$, where $\Delta(c) = \text{flex}^+(c) + \text{kol}(c) - 10$ and $|Z| \geq 14 - \Delta(c)$.
3. Assume that c has exactly two uncolored incident edges of C_{\max} and $\lambda(c) > 2$. Then $|Z| \geq 12 - \Delta(c) + \rho(c)$, where $\Delta(c) = \text{flex}^+(c) + \text{kol}(c) - 8$. If we color only one ray of c , then $\Delta'(c) = 2 - \Delta(c)$, otherwise $\Delta'(c) = 6 - \Delta(c)$.
4. Assume that c has at least $u \geq 3$ uncolored rays and $\lambda(c) > 2$. Then $|Z| \geq 20 - \text{flex}^+(c) - \text{kol}(c) + \rho(c)$. If we color $u - 2$ rays of c , then $\Delta'(c) = 0$; if $u - 1$, then $\Delta'(c) = \min\{10 - \text{flex}^+(c) - \text{kol}(c), 0\}$; if we color all u rays of c , then $\Delta'(c) = \min\{14 - \text{flex}^+(c) - \text{kol}(c), 0\}$.
5. If U consists of an antenna of c , then $\Delta'(c) = 4$ and $|Z| \geq 16$.

The proof of this lemma is contained in the proof of Lemma 20.

Fact 4 1. Each edge of C_{\max} is an antenna of at most two different halfy 2-cycles of C_1 .

2. If an edge $e \in C_{\max}$ is an antenna of two halfy 2-cycles, then it is not incident to a cycle of C_1 .

Lemma 10 Let c such that $\lambda(c) > 2$ be an unprocessed cycle of C_1 that at some step of Algorithm Color7 has a minimal number of uncolored rays. Then it is always possible to process c .

Proof. We divide the set of colors \mathcal{K} into two sets $Z^+(c)$ and $Z^-(c)$. Next, we color each uncolored inray of c with one of the colors of $Z^-(c)$ and each uncolored outray of c with one of the colors of $Z^+(c)$. This way each newly colored ray is safe - by Observation 1 and the assumption that all previously colored rays are safe.

Now, we prove that we can carry out the above in such a way that no cycle or halfy 2-cycle of C_1 becomes blocked.

Suppose first that c has exactly one uncolored ray r . By Lemma 9 there exists $\Delta(c) \leq 4$ and a $(12 - \Delta(c))$ -element set $Z \subseteq K$ such that by coloring r with $4 - \Delta(c)$ colors of Z we guarantee that c does not become blocked. If r is incident to another cycle of C_1 or is an antenna of a halfy 2-cycle of C_1 , then we may also have to ensure that this (halfy) cycle denoted as c' does not become blocked. Regardless of whether c' is a cycle or a halfy 2-cycle of C_1 , by Lemma 9 we know that there exists an analogous number $\Delta(c') \leq 4$ and an at least $(12 - \Delta(c'))$ -element set $Z' \subseteq K$ such that coloring r with $4 - \Delta(c')$ colors of Z' guarantees that c' does not become blocked. Because $20 \geq |Z \cup Z'| = |Z| + |Z'| - |Z \cap Z'| \geq 24 - \Delta(c) - \Delta(c') - |Z \cap Z'|$, we obtain that $|Z \cap Z'| \geq 4 - \Delta(c) - \Delta(c')$. If $\Delta(c) + \Delta(c') \geq 4$, then $(4 - \Delta(c)) + (4 - \Delta(c')) \leq 4$ and we can simply use $4 - \Delta(c)$ colors of Z and $4 - \Delta(c')$ colors of Z' to color r . Otherwise, we use $4 - \Delta(c) - \Delta(c')$ colors of $Z \cap Z'$, $\Delta(c')$ colors of $Z \setminus Z'$ and $\Delta(c)$ colors of $Z' \setminus Z$. This way neither c nor c' will become blocked.

Suppose now that c has exactly two uncolored rays r_1, r_2 . By Lemma 9 it is enough to color r_1, r_2 with $6 - \Delta(c)$ colors of an at least $(12 - \Delta(c))$ -element set $Z \subseteq K$. If r_1 is a ray or an antenna of a (halfy) cycle c' of C_1 , then by the argument above, we can color r_1 so that

at least $4 - \Delta(c)$ colors belong to Z and at least $4 - \Delta(c')$ to Z' . This means that we have already used at least $4 - \Delta(c)$ colors of Z . To guarantee that c is not blocked, it suffices to use at most 2 additional (not already used on r_1) colors of Z . If r_1 and r_2 are also the last two uncolored rays of c' , then we have to use $6 - \Delta(c')$ colors of Z' , which means that it suffices to color r_2 with two additional colors of Z' . If r_2 is a ray of a different cycle c'' of C_1 , then by Lemma 9 it is enough to color r_2 with $2 - \Delta(c'') \geq 2$ colors of an at least $(12 - \Delta(c''))$ -element set Z'' . Thus both these cases are easy to handle - since r_2 has to be colored with 4 colors, we use 2 colors of Z and two colors of either Z' or Z'' , depending on whether r_2 is incident to the same cycle c' or not.

If c has exactly 3 uncolored rays, then on the one hand $col(c) + flex^+(c) \geq 8$ and on the other it suffices to color the uncolored rays of c with $14 - kol(c) - flex^+(c) \leq 6$ colors that increase $kol(c)$. If the uncolored rays of c are also the last 3 uncolored rays of some different cycle c' , then we may also need to use 6 colors of $\mathcal{K} \setminus col(c')$. This can be easily achieved as we may use up to 12 different colors for coloring 3 rays. □

Lemma 11 *Let c such that $\lambda(c) = 2$ be an unprocessed cycle of C_1 that at some step of Algorithm Color7 has a minimal number of uncolored rays. Then it is always possible to process c .*

Proof. If c has only one uncolored ray, then the proof is the same as in Lemma 10.

Let O_2 denote a set of cycles of C_1 such that each cycle c of O_2 has no chords and has exactly two uncolored rays coincident with the same edge e of c .

Claim 1 *It never happens that each of the uncolored rays of O_2 is incident to another cycle of O_2 .*

Proof. Suppose to the contrary that each of the uncolored rays of O_2 is incident to another cycle of O_2 . Then the rays of these cycles together with edges of these cycles, with which the uncolored rays are coincident form a good alternating cycle contradicting Lemma 3. □

By this claim, we can always choose for processing a 2-cycle that either does not belong to O_2 or belongs to O_2 but one of its rays is not incident to a cycle of O_2 .

Suppose that c belongs to O_2 and has two uncolored rays r_1, r_2 . Rays r_1, r_2 have to be diverse with the already colored rays of c . Thus, there exists an at least 12-element set Z such that we have to color r_1, r_2 with 8 colors of Z . If r_1, r_2 are also rays of another cycle c' , then by the above claim, $c' \notin O_2$. If r_1, r_2 are also the last uncolored rays of c' , then there exists a number $\Delta(c') \leq 4$ and an at least $(12 - \Delta(c') + \rho(c'))$ -element set $Z' \subseteq K$ such that coloring r_1, r_2 with $8 - \Delta(c')$ colors of Z' guarantees that c' does not become blocked. Since $\rho(c') = 4$, we get that $|Z \cap Z'| \geq 8 - \Delta(c')$, which means that we have enough colors at our disposal. If r_1, r_2 are not the last uncolored rays of c' , then the task is even easier, because either Z' has more colors or we have to use fewer than $8 - \Delta(c')$ colors of Z' .

Consider now the case when $r_i, i \in \{1, 2\}$ is a ray or antenna of c_i and $c_1 \neq c_2$. Then the coloring of r_1, r_2 is the most difficult when both r_1 and r_2 are antennas. In such a case by Lemma 9 there exist sets Z_1, Z_2 , each of size at least 16 such that coloring r_i with 4 colors of Z_i guarantees that c_i is not blocked. Since we have that $|Z \cap (Z_1 \cup Z_2)| \geq 8$, it is also possible to color r_1, r_2 , so than none of the cycles c, c_1, c_2 is blocked.

If c has more than two uncolored rays, then processing c is easy, because we can use all colors of \mathcal{K} so as not to block c and even if each of the uncolored rays r_i is an antenna, then

there exists an at least 16-element set Z_i , which can be used for coloring r_i . \square

If a is an antenna of two different halfy 2-cycles, then it is said to be a **bilateral antenna**.

Lemma 12 *Let p be an unprocessed path of C_1 . Then it is possible to process it.*

Proof. The proof is similar to the one above. Since paths are processed after cycles of C_1 , the only thing we have to take care of is that antennas of the same halfy 2-cycle are diverse. The path p has at most two incident bilateral antennas a_1, a_2 and if it does, then at most one of them is an inray and at most one an outray of p . Assume that a_1 is an inray and a_2 an outray. (They may also be chords.) Each a_i may have to be diverse with two different edges. Thus for each a_i it may happen that up to 8 colors are forbidden on it. Let Z_i denote the set of colors forbidden on a_i . We partition \mathcal{K} into two 10-element sets $Z^-(p)$ and $Z^+(p)$ so that $|Z^-(p) \setminus Z_1| \geq 5$ and $|Z^+(p) \setminus Z_2| \geq 5$. To achieve this we divide $\mathcal{K} \setminus (Z_1 \cap Z_2)$ (almost) equally between $Z^-(p)$ and $Z^+(p)$. Since $|Z_1 \cap Z_2| \leq 10$, it is always possible. Then we are able to color each ray of p so as to ensure that each antenna is diverse with required antennas. \square

5 Missing proofs

We start with two auxiliary lemmas.

Lemma 13 *Let t be a tricky 3-triangle and a any of its edges. Then $\frac{9}{31}w(t) < w(a) < \frac{2}{5}w(t)$. Let p be any two-edge path contained in $opp(t)$. Then $w(p) < \frac{28}{37}w(t)$.*

Proof. We show that if at least one of these statements does not hold, then there exists an amenable subgraph on t of weight at least $14w(t)$, which contradicts the fact that t is tricky.

Suppose that $w(a) \geq \frac{2}{5}w(t)$ for some edge a of t . Then the other two edges have weights $w(b) = \frac{3}{10}w(t) + \delta, w(c) = \frac{3}{10}w(t) - \delta$ for some $\delta \geq 0$. By taking 20 copies of a , 17 of b and 3 of c , we obtain an amenable subgraph on t of weight at least $14w(t)$.

Suppose now that $\frac{9}{31}w(t) \geq w(a)$ for some edge a of t . Then the other two edges have weights $w(b) = \frac{11}{31}w(t) + \delta, w(c) = \frac{11}{31}w(t) - \delta$ for some $\delta \geq 0$. By taking 20 copies of a , 17 of b and 3 of c , we obtain an amenable subgraph on t of weight at least $14w(t)$.

Assume now that there exists a two-edge path p contained in $opp(t)$ having weight $w(p) \geq \frac{28}{37}w(t)$. Therefore its edges a and b have weights $w(a) = \frac{14}{37}w(t) + \delta, w(b) = \frac{14}{37}w(t) - \delta$ for some $\delta \geq 0$. By taking 20 copies of a and 17 of b , we obtain an amenable subgraph on t of weight at least $14w(t)$. \square

The corollary of this lemma is the following:

Lemma 14 *Let t be a tricky 3-triangle and a any of its edges. Let a', b', c' denote edges of $opp(t)$ such that $w(a') \geq w(b') \geq w(c')$. Then $w(b') < \frac{14}{37}w(t)$.*

Proof of Lemma 1

First we show that any perfect matching M of G' yields a relaxed cycle cover. For any edge $(u_{out}, e_{uv}^1) \in M$, we add a half-edge $(u, x_{(uv)})$ to \tilde{C} and for any edge $(u_{in}, e_{vu}^2) \in M$, we add a half-edge $(x_{(vu)}, u)$ to \tilde{C} .

Since M is a perfect matching of G' each vertex u_{in} and each vertex u_{out} has an incident edge in M . Hence, each vertex in V has exactly one outgoing and one incoming half-edge in \tilde{C} .

If an edge $(u, v) \in E$ does not belong to any tricky 2-cycle, 3-triangle or 2-triangle of R , then it is replaced with an edge (u_{out}, v_{in}) in E' . Thus, if \tilde{C} contains only one half-edge of some edge e , then e must belong to one of the mentioned tricky cycles.

For every tricky 3-triangle t , G' contains eight additional vertices $\gamma_p^{t+}, \gamma_p^{t-}, \dots, \gamma_r^c, \gamma_q^c$ each of which excludes exactly one half-edge within $t \cup opp(t)$ from \tilde{C} . This exclusion follows from the fact that each of these vertices is matched to some vertex $e_{(uv)}^1$ or $e_{(uv)}^2$, such that $(u, v) \in t \cup opp(t)$. If one of these additional vertices is matched to a vertex $e_{(uv)}^1$, then a half-edge $(u, x_{(uv)})$ does not belong to \tilde{C} . Similarly, if it is matched to a vertex $e_{(uv)}^2$, then a half-edge $(x_{(uv)}, v)$ does not belong to \tilde{C} . Therefore, for each tricky triangle t , at least eight half-edges within $t \cup opp(t)$ do not belong to \tilde{C} , which means that \tilde{C} contains at most four half-edges within $t \cup opp(t)$, hence contains neither t nor $opp(t)$.

Next, we deal with the properties stated in the current lemma. The proof of the first property is very similar to the proof of Lemma 2 in [25].

Let $t = (p, q, r)$ be a tricky 3-triangle such that (r, q) is the chosen edge with maximum weight among edges of $opp(t)$. Let us notice that we may assume that \tilde{C} is integral on $\{(q, p), (r, p)\}$ because if M contains one half-edge of (q, p) , then it also contains one half-edge of (r, p) and these half-edges are crossing. Also, the half-edges incoming to p have the same weight. Therefore such two half-edges in M can be replaced in \tilde{C} by that one of the edges $\{(q, p), (r, p)\}$, whose half-edge incident to q or r is contained in M .

Applying the same kind of reasoning we can prove that:

Claim 2 *Let c denote the 2-cycle (q, r) . If vertices γ_q^c and γ_r^c are matched in M to the subdivision vertices of the same edge of c , then M yields a relaxed cycle cover \tilde{C} , which is integral on t .*

We denote the half-edges $(u, x_{(u,v)})$ and $(x_{(u,v)}, v)$ as u_v^{\rightarrow} and v_u^{\leftarrow} .

Assume first that γ_q^c and γ_r^c are matched in M to $e_{q,r}^2$ and $e_{r,q}^2$. This means that γ_q^{t-} must be matched to $e_{p,q}^2$ and γ_r^{t-} to $e_{p,r}^2$. Vertices $\gamma_q^{t+}, \gamma_r^{t+}$ may be matched in M in one of the following ways:

1. $e_{p,q}^1$ and $e_{p,r}^1$. Thus, $e_{q,r}^2$ and $e_{r,q}^2$ must be matched to q_{out} and r_{out} , which in turn means that neither can $e_{q,p}^1$ be matched to q_{out} nor $e_{r,p}^1$ to r_{out} . Therefore, \tilde{C} contains two outgoing half-edges within the 2-cycle (q, r) and no half-edges within $\{(p, q), (p, r), (q, p), (r, p)\}$.
2. $e_{r,q}^1$ and $e_{q,r}^1$. This would mean that both $e_{p,q}^1$ and $e_{p,r}^1$ must be matched to p_{out} , which is impossible. Therefore this case cannot occur.
3. $e_{r,q}^1$ and $e_{p,r}^1$. \tilde{C} contains half-edges $q_r^{\rightarrow}, p_q^{\rightarrow}$, and either the edge (q, p) or no half-edge within $\{(r, p), (q, p)\}$.
4. $e_{p,q}^1$ and $e_{p,r}^1$. \tilde{C} contains half-edges $p_r^{\rightarrow}, r_q^{\rightarrow}$ and either the edge (r, p) or no half-edge within $\{(r, p), (q, p)\}$.

Assume next that γ_q^c and γ_r^c are matched in M to $e_{q,r}^1$ and $e_{r,q}^1$. This means that γ_q^{t+} must be matched to $e_{p,q}^1$ and γ_r^{t+} to $e_{p,r}^1$. Vertices $\gamma_q^{t-}, \gamma_r^{t-}$ may be matched in M in one of the following ways:

1. $e_{p,q}^2$ and $e_{p,r}^2$. \tilde{C} contains two incoming half-edges within the 2-cycle (q, r) .
2. $e_{r,q}^2$ and $e_{q,r}^2$. This means that $e_{p,q}^2$ must be matched to q_{in} and $e_{p,r}^2$ to r_{in} . Therefore, \tilde{C} contains $q_p^{\leftarrow}, r_p^{\leftarrow}$ and no half-edge within (q, r) .
3. $e_{r,q}^2$ and $e_{p,r}^2$. \tilde{C} contains half-edges $q_p^{\leftarrow}, r_q^{\leftarrow}$.
4. $e_{p,q}^2$ and $e_{q,r}^2$. \tilde{C} contains half-edges $q_r^{\leftarrow}, r_p^{\leftarrow}$.

In all the above four cases \tilde{C} may contain additionally one of the edges $(q, p), (r, p)$.

We now want to show that $w(\tilde{C})_t$ satisfies the conditions described in the definition of a harmonious triangle t .

Suppose first that \tilde{C} is not integral and contains two half-edges within $t \cup opp(t)$. By the observation above we know that none of these half-edges belongs to (q, p) or (r, p) . If \tilde{C} does not contain a half-edge of (r, q) (the heaviest edge of $opp(t)$), then to prove that t is harmonious in this case it suffices to show that $5w(a') + 5w(c') + 4w(t) \leq 10w(t) - 5w(a)$ holds for any two different edges a', c' of $\{(p, q), (p, r), (q, r)\}$ and for any edge a of t . This is equivalent to showing $5w(a') + 5w(c') + 5w(a) \leq 6w(t)$. Suppose to the contrary that $5w(a') + 5w(c') + 5w(a) > 6w(t)$. But then if $a', c' \in opp(t)$, by Lemma 13 $5(w(a') + w(c')) < 5 \cdot \frac{28}{37}w(t)$ and $w(a) < \frac{2}{5}w(t)$, which means that $5w(a') + 5w(c') + 5w(a) < 6w(t)$ - a contradiction. If only one of a', c' belongs to $opp(t)$, we have that $5w(a') + 5w(c') + 5w(a) < 10 \cdot \frac{2}{5}w(t) + 5 \cdot \frac{14}{37}w(t) < 6w(t)$ - again a contradiction. If both a' and c' belong to t , then $w(a') + w(c') < \frac{22}{31}w(t)$ and hence $5w(a') + 5w(c') + 5w(a) < 6w(t)$. The proof holds also when \tilde{C} contains two half-edges within $opp(t)$, one of which may be a half-edge (r, q) , because then $5(w(a') + w(c')) < 5 \cdot \frac{28}{37}w(t)$.

Assume now that \tilde{C} contains two half-edges, one of which is a half-edge of (r, q) and the other is within t . We notice that then \tilde{C} must contain one half-edge of (q, r) , because if M contains one half-edge of (r, q) and one half-edge of (p, q) , then these half-edges are crossing and can be replaced by one edge - either (r, q) or (p, q) . Now we prove that $5w(r, q) + 5w(q, r) + 4w(t) \leq \max\{10w(t) - 5w(a), 10(w(r, q) + w(q, r))\}$ holds for any edge a of t different from (q, r) . Let $b = (q, r)$ and $b' = (r, q)$. Suppose that $5w(b) + 5w(b') + 4w(t) > 10w(t) - 5w(a)$. This means that $w(b') + w(b) > \frac{6}{5}w(t) - w(a)$. If $5w(b) + 5w(b') + 4w(t)$ is also greater than $10(w(b) + w(b'))$, then $w(b) + w(b') < \frac{4}{5}w(t)$. However, by Lemma 13 $\frac{6}{5}w(t) - w(a) > \frac{4}{5}w(t)$ - a contradiction.

Let us now consider the cases when \tilde{C} is not integral and contains four half-edges within $t \cup opp(t)$. Hence, \tilde{C} contains the whole edge (r, p) or the whole edge (q, p) , because \tilde{C} can contain at most two half-edges within the remaining edges within $t \cup opp(t)$. We can notice that if \tilde{C} contains (r, p) , then we have an identical proof as above for the case of two half-edges within $t \cup opp(t)$ because we then add $10w(a)$ to both sides of the inequality. We analyze now the cases when \tilde{C} contains (q, p) :

1. all four half-edges are within $opp(t)$. We show that $5w(a') + 5w(b') + 10w(c') \leq 6w(t) + 5w(a)$ if $a', b', c' \in opp(t), a \in t$ and $c' \neq (r, q)$. Since $5w(a') + 5w(b') + 10w(c') = 5w(opp(t)) + 5w(c') \leq 5w(t) + \frac{70}{37}w(t) < 7w(t)$ and $6w(t) + 5w(a) > 7\frac{14}{31}w(t)$, the inequality indeed holds.
2. \tilde{C} contains also two incoming half-edges within (q, r) . We notice that it cannot happen that $5w(b) + 5w(b') + 4w(t) + 10w(c') > 10w(t) + 5w(a)$, because it would mean that $w(b') + 2w(c') > \frac{6}{5}w(t)$. However, by Lemma 13 $w(c') + w(b') < \frac{28}{37}w(t)$ and $w(c') < \frac{14}{37}w(t)$, which means that $w(b') + 2w(c') < \frac{42}{37}w(t) < \frac{6}{5}w(t)$.

3. \tilde{C} contains $q_p^{\leftarrow}, r_p^{\leftarrow}$. Then we show that $5w(c) + 5w(a') + 10w(c') + 4w(t) \leq 10w(t) + 5w(b)$, which is equivalent to $w(c) - w(b) + w(a') + 2w(c') \leq \frac{6}{5}w(t)$. a', c' are two edges of $opp(t)$ which do not contain a maximum weight edge b' . Hence $w(a') + 2w(c') \leq w(opp(t)) \leq w(t)$. Also, $w(c) - w(b) < \frac{2}{5}w(t) - \frac{9}{31}w(t) < \frac{1}{5}w(t)$.
4. \tilde{C} contains half-edges $q_p^{\leftarrow}, r_q^{\leftarrow}$. We show that $5w(b) + 5w(c) + 10w(c') + 4w(t) \leq 10w(t) + 5w(b)$, which is equivalent to $w(c) + 2w(c') \leq \frac{6}{5}w(t)$. We know that $2w(c') < \frac{28}{37}w(t) < \frac{4}{5}w(t)$. On the other hand, $w(c) < \frac{2}{5}w(t)$.

Proof of Lemma 6

Let $e = (u, v)$ be any edge of c and $e_1 = (u, v'), e_2 = (u', v)$ edges of C'_{max} coincident with it. Then e has to be colored with colors of $\mathcal{K} \setminus (col(e_1) \cup col(e_2))$, i.e., it cannot be colored with any color assigned to the edge coincident with it.

Suppose first that c does not contain any chords. If each color $k \in \mathcal{K}$ is assigned to some ray of c , then we are already done. By coloring each edge e of c with any $mult(e)$ colors of \mathcal{K} that are not assigned to any edges of C_{max} coincident with e , we achieve that each color $k \in \mathcal{K}$ is *not* assigned to some edge of c , thus c is not monochromatic with respect to any color of \mathcal{K} . Otherwise, if $kol(c) < 20$ and thus not every color $k \in \mathcal{K}$ is assigned to some ray of c , we still have that $\chi(c) = kol(c) + flex(c) \geq 20$. Therefore, $flex(c) = \sum_{e \in c} flex(e) \geq 20 - kol(c)$. This means that for each edge e of c with $flex(e) > 0$ we can choose $flex(e)$ colors of \mathcal{K} , which will *not* appear on e . Recall that e is colored with $mult(e)$ colors and $flex(e) = 10 - |col(e_1) \cup col(e_2)|$, where e_1, e_2 are edges coincident with e . This way we can distribute all colors of \mathcal{K} not assigned to any rays of c among edges of c and ensure that each edge of \mathcal{K} , which is not assigned to any ray of c , does not appear on some edge of c . Hence, c will again not be monochromatic w.r.t. any color of \mathcal{K} .

Let us also remark that during processing of cycles we keep an invariant that if c has some rays, then some two of them are diverse.

Assume now that c contains some chords. Observe that if c has no incident bows (edges of multiplicity 5), then the flexibility of each edge e of c satisfies $flex^0(e) \geq 2 + \min\{0, 8 - kol(c)\}$. Thus for the flexibility of the whole cycle c it holds: $flex^0(c) \geq 2\lambda(c) + \min\{0, 8 - kol(c)\}$. This means that if c has length at least 6, we can color the chords of c however we like and c will not become blocked - $\chi(c) \geq 20$. Using the same argument as above we can then always color the edges of c in such a way that each color of \mathcal{K} does not appear on some edge of c .

There remains the question of how to ensure that each chord is safe. If a chord e of c is contained in a directed path P consisting of edges of C_{max} that contains some ray r of c , then to guarantee that e is safe, we can simply color all edges of P between e and the closest ray r' of c (together with e) with the same set of colors as r' . Notice that if c has length smaller than 6 and has no uncolored chords after this procedure, then $flex^+(c) \geq 4$, thus $kol(c) + flex(c) \geq 20$. Assume then now that we have already colored all such chords of c .

Assume that c has the form $(v_1, v_2, \dots, v_{\lambda(c)})$, then we say that a sequence $R(i, j) = (v_i, v_{i+1}, \dots, v_j)$ is **a row of subcycles** if (i) for each $k, i \leq k \leq j$ there exists an uncolored subcycle c_k of c going through v_k and for any two i_1, i_2 such that $i \leq i_1 < i_2 \leq j$ subcycles c_{i_1} and c_{i_2} are different and (ii) it is maximal - i.e. it cannot be extended.

A row $R(i, j)$ begins (corr. ends) in one of three ways:

1. a marker - when edges of C_{max} incident to v_{i-1} (corr. v_{j+1}) are already colored,
2. a twist - if a subcycle c' of c going through v_{i-1} goes also through one of the vertices v_{i+1}, \dots, v_j (corr. c' going through v_{j+1} goes also through one of the vertices v_i, \dots, v_{j-1}),

3. a broad subcycle - if a subcycle c_i goes also through v_{i-1} (corr. c_j goes also through v_{j+1}).

To color a row $R(i, j)$ of subcycles in a **zebra manner** means to (i) choose two disjoint four-element sets of colors B and W and (ii) color all edges of C_{max} outgoing of v_i, \dots, v_j using all colors of B and the remaining edges of subcycles c_i, c_{i+1}, \dots, c_j with 4 colors of W .

Claim 3 *Suppose that a row $R(i, j)$ of subcycles is colored in a zebra manner and $e = (v', v_{i-1})$ is an edge of C_{max} . Then*

1. *If $R(i, j)$ begins with a marker and $B = col(e)$, then all edges of subcycles c_i, c_{i+1}, \dots, c_j are safe except for possibly the edge outgoing of v_j - some edge (v_j, v'') of c_j .*
2. *If $R(i, j)$ begins with a twist, then all edges of subcycles c_i, c_{i+1}, \dots, c_j are safe except for possibly the edge (v_j, v'') of c_j and the edge $e' = (v_{i-1}, v_i)$ of c has $flex(e') \geq 4$.*
3. *If $R(i, j)$ begins with a broad cycle, then all edges of subcycles c_i, c_{i+1}, \dots, c_j are safe except for possibly the edges: (v_j, v'') of c_j and (v''', v_i) of c_i ; the edge $e' = (v_{i-1}, v_i)$ of c has $flex^+(e') \geq 4$ or (v''', v_i) is safe.*

Suppose first that $kol(c) + flex^0(c) \geq 20$ and no chord is a bow (an edge of multiplicity 5). We then only have to color the chords and the edges of the cycle c so that each one of them is safe. We consider each row $R(i, j)$ of subcycles separately. Suppose first that it begins with a marker. We then choose for B some 4 colors contained in $col(e_1) \cup col(e_2)$ and color $R(i, j)$ in a zebra manner. If $R(i, j)$ ends with a twist, then all edges contained in the subcycles of $R(i, j)$ are safe as a result. Otherwise, the only edge, which may be not safe is (v_j, v'') . If $R(i, j)$ ends with a broad cycle, then $flex^+(v_j, v_{j+1}) \geq 4$ and we can forbid 4 colors of W on (v_j, v_{j+1}) thus ensuring that (v_j, v'') is safe. If $R(i, j)$ ends with a marker, then let e_3, e_4 denote the edges of C_{max} incident to v_{j+1} . To W' we add $\min\{4, |(col(e_3) \cup col(e_4)) \setminus B|\}$ colors of $(col(e_3) \cup col(e_4)) \setminus B$. Note that (v_j, v'') is safe w.r.t. each such color. If W' has fewer than 4 such colors, we notice that $flex^+(v_j, v_{j+1}) \geq 4 - |W'|$, because $4 - |W'|$ colors of B must be also present on (v''', v_{j+1}) . Therefore we can forbid all the remaining colors of $W \setminus W'$ on (v_j, v_{j+1}) and thus ensure that (v_j, v'') is safe. If $R(i, j)$ begins and ends with a twist, then by coloring it in a zebra manner, we guarantee that all edges of its subcycles are safe. Similarly, if $R(i, j)$ begins and ends with a broad cycle, then $flex^+(v_{i-1}, v_i) \geq 4, flex^+(v_j, v_{j+1}) \geq 4$ and we can forbid colors of B on (v_{i-1}, v_i) and colors of W on (v_j, v_{j+1}) .

If any of the subcycles c' in a row $R(i, j)$ contains a bow, then c' cannot be a broad cycle (because a broad cycle contains an edge of multiplicity at least 14 and a 2-cycle containing a bow has edges with multiplicities 4 and 5). If $kol(c) \geq 20$ or $kol(c) + flex^0(c) \geq 20+$ the number of chords that are bows, then we color any bow occurring in a row $R(i, j)$ with a color $k' \notin W \cup B$ and forbid k' on one edge of c , using thus one unit of $flex^0(c)$.

We deal now with cycles with chords such that $flex^0(c) + kol(c) < 20$ and without incident bows. Note that such c has length between 4 and 5. In this case we additionally need to get an extra $flex^+(c) \geq 4$. Except for the case when $R(i, j)$ begins and ends with a broad cycle, we either get it for free by the above claim or can ensure it by choosing B in such a way that $flex^+(v_{i-1}, v_i) \geq 4$. If a row $R(i, j)$ begins and ends with a broad cycle, then we do not color it in a zebra manner but in an **alternate zebra manner**: we use the same 4 colors of B on edges $(v_i, v'_i), (v'_{i+1}, v_{i+1}), (v_{i+2}, v'_{i+2}), \dots$ and the same 4 colors of W on $(v'_i, v_i), (v_{i+1}, v'_{i+1}), (v'_{i+2}, v_{i+2}), \dots$ thus increasing $flex^+(c)$ sufficiently.

Finally, we deal with cycles with chords such that $flex^0(c) + kol(c) < 20$ and having incident bows. The outline of the proof in this case is the following. Let c' be an uncolored subcycle. It is contained in two rows $R(i, j)$ and $R(i', j')$. Sometimes we need to get an extra $flex^+(c) \geq 7$ (for example, if c is a square). We are then able to get an extra $flex^+(c)$ of 4 from each of the rows. \square

6 Construction of G_1 in the presence of tricky triangles and tricky 2-cycles

We show how to modify the multigraph G_1 built in the previous section, when G contains tricky triangles or C_1 contains strange 2-cycles. We say that a tricky triangle $t = (p, q, r)$ is **halfy** if C_1 contains exactly one half-edge of some edge of t or of $opp(t)$.

The main new features are going to be the following 3 types of subgraphs, shown in Figures 4 and 5, arising on tricky triangles:

1. a subgraph on p, q, r such that C_{max} contains a halfy 3-triangle $t = (p, q, r)$ and C_1 contains exactly four edges incident to t : either three incoming edges and one outgoing of t or three outgoing and one incoming. W.l.o.g. assume that C_1 contains edges $(p', p), (r', r), (q, q_1), (q_2, q)$. We proceed as indicated in the definition of a harmonious triangle. G_1 then contains either (i) 10 copies of each of $(r, p), (q, r)$ and 5 copies of (p, q) or (ii) 10 copies of each of $(p, r), (r, p)$ and 0 copies of $(p, q), (q, r)$. We choose the option of maximum weight. If option (ii) is maximum, then we treat C_{max} as though it contained the 2-cycle (p, r) and not the triangle (p, q, r) and in coloring G_1 we do not treat (p, q, r) as a tricky triangle. Any edge $e \in C_{max}$ such that $mult(e) = 10$ is called a **b-edge**. A subgraph of G_1 on p, q, r contains thus two b-edges.
2. a subgraph on p, q, r such that C_{max} contains a halfy 2-triangle $t = (p, q, r)$ with a t-cycle $c = (q, r)$ and C_1 contains exactly four edges incident to t , three of which are incident to c . W.l.o.g. assume that two edges of C_1 are incident to q and that C_1 contains an edge (r_1, r) . Then $mult(r, q) = 10$ and $mult(q, r) = 5$, hence (r, q) is a b-edge.
3. a subgraph on q, r such that C_{max} contains a 2-cycle $c = (q, r)$, which is a t-cycle of a tricky 2-triangle t and C_1 contains a loop e'_t . Then C_1 contains four edges incident to c and $mult(q, r) = 5, mult(r, q) = 4$. We call the edge (q, r) a **bow**.

The following two types of subgraphs can be treated in a very similar way as a subgraph surrounding a halfy 2-cycle (Figure 6):

1. a subgraph on p, q, r such that C_{max} contains a halfy 3-triangle $t = (p, q, r)$ and C_1 contains exactly two edges incident to t : either two incoming or two outgoing. W.l.o.g. assume that C_1 contains edges (p', p) and (q', q) . We again proceed as indicated in the definition of a harmonious triangle. G_1 contains then either (i) 10 copies of each of $(p, q), (r, p)$ and 15 copies of (q, r) or (ii) 10 copies of each of $(p, r), (r, q), (q, r)$ or (iii) 10 copies of each of $(q, r), (r, p), (p, r)$. We choose the option with maximum weight. If option (ii) or (iii) is maximum, then we treat C_{max} as though it contained the 2-cycle (r, q) or (r, p) and not the triangle (p, q, r) and in coloring G_1 we do not treat (p, q, r) as a tricky triangle. We call the edges $(p', p''), (q', q'')$ of C_{max} **antennas** of t and require that they are diverse.




Figure 4: Halfy triangles with 3 incoming and 1 outgoing edges of C_1 : a tricky 3- and 2-triangle $t = (p, q, r)$.




Figure 5: A bow.

- a subgraph on p, q, r such that G contains a halfy 2-triangle $t = (p, q, r)$, where $c = (q, r)$ is its t -cycle and C_1 contains exactly two edges incident to t : either two incoming or two outgoing. W.l.o.g. assume that C_1 contains edges (p', p) and (q', q) . G_1 contains then 5 copies of each of (p, q) , (r, q) , 4 copies of (r, p) and 14 copies of (q, r) . We call the edges (p', p'') , (q', q'') of C_{max} **antennas** of t and require that they are diverse. We call the edge (p, p''') of C_{max} a **weak antenna** of t and require that it is **weakly diverse** with (p', p'') , by which we mean that $|col(p, p''') \setminus col(p', p'')| \geq 2$.




Figure 6: Halfy triangles with two incoming edges of C_1 : a tricky 3- and 2-triangle $t = (p, q, r)$.

If C_1 contains some strange 2-cycle or tricky triangles, then the multigraph G_1 contains non-path-20-colorable subgraphs. We deal with such non-colorable subgraphs at the end by finding exchange sets F_1, F_2 and extending the partial path-20-coloring.

If C_1 contains a 2-cycle or triangle of C_{max} , then such a cycle is not tricky and then we can replace it with other edges so as to receive an amenable subgraph.

In all other aspects the obtained multigraph G_1 has almost the same properties as in previous sections, i.e., each vertex has at most two incoming and two outgoing edges: two in C_{max} and two in C_1 and thus indegree and outdegree at most 4.

Below we give a detailed description of the construction of G_1 .

6.1 Tricky 2-cycles

A 2-cycle $c = (u, v)$ of G is **strange** if exactly one of the edges of c belongs to C_{max} . Let $c = (u, v)$ be a strange 2-cycle such that $(u, v) \in C_{max}$ and $(u', u), (v, v')$ are its incident edges of C_{max} . If c is not a subcycle of a triangle of C_{max} and both (i) $w(u, v) > \frac{3}{4}(w(u', u) + w(v, v'))$ and (ii) $w(v, u) > \frac{3}{4}(w(u', u) + w(v, v'))$, then it is said to be **in corrigible**.

Lemma 15 *Let $c = (u, v)$ be an incorrigible 2-cycle. Then at most one of the vertices u, v is part of an incorrigible 2-cycle $c' \neq c$. If $c = (u, v)$ and $c' = (u, v')$ are two incorrigible 2-cycles such that $(u, v), (u, v'), (v', v'') \in C_{max}$ and $w(u, v) \geq w(u, v')$, then $w(v', v'') < \frac{w(u, v')}{3}$.*

Proof. Suppose to the contrary that both u and v are part of incorrigible 2-cycles $c_1 = (u', u)$ and $c_2 = (v, v')$ different from c . Assume that edges $(u, v), (u', u), (v, v')$ belong to C_{max} and let $(u'', u'), (v', v'')$ be edges of C_{max} adjacent to u' and v' . Since the 2-cycles c_1 and c_2 are incorrigible, we have that $w(u', u) > \frac{3}{4}(w(u, v) + w(u'', u'))$ and $w(v, v') > \frac{3}{4}(w(u, v) + w(v', v''))$, which implies that $w(u', u) > \frac{3}{4}w(u, v)$ and $w(v, v') > \frac{3}{4}w(u, v)$. It means that $\frac{3}{4}(w(u', u) + w(v, v')) > \frac{9}{8}w(u, v) \geq w(u, v)$. Therefore $c = (u, v)$ is not incorrigible - a contradiction.

Let $c = (u, v)$ and $c' = (u, v')$ be two incorrigible 2-cycles such that $(u, v), (u, v'), (v', v'') \in C_{max}$ and $w(u, v) \geq w(u, v')$. Then, because c' is incorrigible, $w(u, v') > \frac{3}{4}(w(u, v) + w(v', v''))$. Hence, $w(v', v'') < \frac{4}{3}w(u, v') - w(u, v)$. Since $w(u, v) \geq w(u, v')$, we get that $w(v', v'') < \frac{1}{3}w(u, v')$. \square

For 2-cycles, which are not subcycles of triangles of C_{max} we in fact use a different definition of a tricky cycle than the one presented earlier.

A 2-cycle $c = (u, v)$ of G , which is not a subcycle of any triangle of C_{max} is tricky if it satisfies one of the following:

1. c is hard (by definition it then belongs to C_{max}).
2. $c = (u, v)$ is an incorrigible 2-cycle that is either vertex-disjoint with any other incorrigible 2-cycle or there exists an incorrigible 2-cycle $c' = (u, v')$ such that $(u, v), (u, v') \in C_{max}$ and $w(u, v) > w(u, v')$.

It easily follows that any two tricky 2-cycles are vertex-disjoint.

6.2 Strange halfy 2-cycles

If C_1 contains exactly one half-edge of each edge of a 2-cycle c of G , then c is called a **halfy** 2-cycle of C_1 . Additionally, if a halfy 2-cycle c of C_1 is such that exactly one of the edges of c belongs to C_{max} , then it is said to be a **strange halfy 2-cycle of C_1** .

In the way shown below we deal with each strange halfy 2-cycle that is not a subcycle of a tricky 3-triangle. To facilitate the subsequent coloring of G_1 , we modify it as follows. We are also going to modify C_{max} . To avoid confusion, we denote the modified C_{max} as C'_{max} . Let $c = (u, v)$ be any strange halfy 2-cycle of C_1 . Suppose that $(u, v) \in C_{max}$. Let $(u', u), (v, v') \in C_{max}$ be the two edges of C_{max} incident to (u, v) . We remove all copies of $(u', u), (v, v')$ (8

in total) from G_1 and replace them with 1 additional copy of (u, v) and 4 additional copies of (v, u) - as a result $\text{mult}(u, v) = \text{mult}(v, u) = 10$ and $\text{mult}(u', u) = \text{mult}(v, v') = 0$. Thus, (u, v) and (v, u) are b-edges. Also, in C'_{max} , we replace edges $(u', u), (v, v')$ with one edge (v, u) . As a consequence of this modification, we obtain a multigraph G_1 , in which each halfy 2-cycle $c = (u, v)$ of C_1 (not only strange but also one with both edges in C_{max}) has no incident edges of C_{max} apart from those already belonging to c and both edges of c are b-edges.

Lemma 16 *The weight of the thus modified G_1 is at least $4w(C_{max}) + 10w(C_1)$.*

Proof. This follows from the fact that each halfy and strange 2-cycle (u, v) of C_1 is incorrigible. Assuming that it is (u, v) that belongs to C_{max} and (u, v) is incorrigible, by the definition we get that $w(u, v) > \frac{1}{4}(w(u', u) + w(v, v'))$ and $w(v, u) > \frac{3}{4}(w(u', u) + w(v, v'))$, which means that $w(u, v) + 5w(v, u) \geq 4(w(u', u) + w(v, v'))$. □

6.3 Tricky 3-triangles

For any tricky 3-triangle, which is halfy in C_1 , we proceed as described at the beginning of this section. This is justified by Lemma 1, which says that any such triangle is harmonious in C_1 .

6.4 Tricky 2-triangles

Lemma 17 *Let $t = (p, q, r)$ be a tricky 2-triangle of C_1 with a t -cycle $c = (q, r)$. Then*

1. $w(r, q) > \max\{\frac{w(t)}{2}, \frac{3}{2}w(q, r)\}$.
2. Let $\Delta = w(r, q) - \frac{3}{2}w(q, r)$. Then $\min\{w(p, q), w(r, p)\} > \frac{3}{5}\Delta + \frac{w(q, r)}{2}$.
3. Let $\epsilon = w(r, q) - \frac{w(t)}{2}$. Then $w(q, r) \geq \frac{w(t)}{3} - \epsilon$.

Proof.

If point 1 did not hold, we could replace 4 copies of (r, q) with two copies of t and obtain a path-20-colorable subgraph or replace 4-copies of (r, q) with 6 copies of (q, r) .

Let $a = (q, r), d = (r, q), b_1 = (p, q), b_2 = (r, p)$.

We now prove point 2. We have that $w(d) = \frac{3}{2}w(a) + \Delta$. We notice that in order for t to be tricky it has to hold that $4w(c) + 10w(t) > 10w(a) + 10w(b_2) + 10w(d)$, because the subgraph consisting of 10 copies of each of a, b_2, d is path-20-colorable. This means that $20w(a) + 10w(b_1) + 10w(b_2) + 4\Delta > 25w(a) + 10w(b_2) + 10\Delta$, which implies that $w(b_1) > \frac{3}{5}\Delta + \frac{w(a)}{2}$. We obtain the same estimation for $w(b_2)$ if we consider the subgraph consisting of 10 copies of each of a, b_1, d . Then it must hold that $4w(c) + 10w(t) > 10w(a) + 10w(b_1) + 10w(d)$.

To prove point 3, suppose to the contrary that $a < \frac{w(t)}{3} - \epsilon$. We will show that in such a case t is not tricky. Notice that $10w(t) + 4w(c) < 10w(t) + \frac{10}{3}w(t) = \frac{40}{3}w(t)$.

Since $a < \frac{w(t)}{3} - \epsilon$, $b + w(b_2) > \frac{w(t)}{3} - \epsilon$. Suppose that $b \geq w(b_2)$. Let $b = \frac{w(t)}{3} + \epsilon/2 + \delta$. Then $w(b_2) > \frac{w(t)}{3} + \epsilon/2 - \delta$.

If b_1 is coincident with an edge of C_{max} with multiplicity 5 (a bow), then we take 15 copies of b_1 , 13 copies of b_2 and 12 copies of a . Hence, the whole subgraph has weight greater than $5w(t) + 7.5\epsilon + 15\delta + \frac{13}{3}w(t) + 6.5\epsilon - 13\delta + 4w(t) - 12\epsilon > \frac{40}{3}w(t)$.

If b_1 is not coincident with an edge of C_{max} with multiplicity 5, then we take 16 copies of b , 12 copies of b_2 and 12 copies of a . Hence, the whole subgraph has also weight greater than $\frac{40}{3}w(t)$. \square

Let $t = (p, q, r)$ be a tricky triangle such that $c = (r, q)$ is a 2-cycle of C_{max} . Let $a = (q, r)$, $d = (r, q)$, $b_1 = (p, q)$, $b_2 = (r, p)$.

1. C_1 contains two crossing half-edges within b_1, b_2 and no half-edges within c .

Then C_1 contains either (r_1, r) and two edges incident to q or (q_1, q) and two edges incident to r . W.l.o.g. assume that the first case holds. G_1 contains then 10 copies of d and 5 copies of a .

To prove that $10w(d) + 5w(a) \geq 5w(b_1) + 5w(b_2) + 4w(c)$, we use Lemma 17 point 3. Let $\epsilon = w(d) - \frac{w(t)}{2}$ and suppose that $10w(d) + 5w(a) < 5w(b_1) + 5w(b_2) + 4w(c)$. Then $10w(d) + 5w(a) < 5w(b_1) + 5w(b_2) + 4w(a) + 4w(d)$, which implies that $6w(d) + w(a) < 5w(b_1) + 5w(b_2)$. Hence $3w(t) + 6\epsilon + w(a) < 5(w(t) - w(a))$. Thus $w(a) < \frac{w(t)}{3} - \epsilon$, which contradicts Lemma 17 point 3.

2. C_1 contains two crossing half-edges within b_1, b_2 and two crossing half-edges within c .

Then C_1 contains either (i) (p_1, p) and (q, q_1) or (ii) (r_1, r) and (p, p_1) . W.l.o.g. assume that case (i) holds. Then G_1 contains 15 copies of d and depending on which is more convenient either (i) 5 copies of a and 5 copies of that edge from b_1, b_2 which has greater weight or (ii) 5 copies of each of b_1, b_2 . To facilitate the coloring of G_1 , we modify G'_1 in such a way that we replace edges $(p, q), (q, q_1)$ with one edge (p, q_1) . The edge (p, q_1) has multiplicity 10 in G'_1 . G'_1 does not contain any of the remaining edges of t or c . The restriction regarding this modification is such that if G'_1 contains (q_1, p) and $mult(q_1, p) = 14$, then we do not perform it and instead remove from G'_1 all edges of t and c as well as (q, q_1) .

Note that the required weight is equal to $W = 4w(c) + 5w(c) + 5w(b_1) + 5w(b_2) = 9w(a) + 9w(d) + 5w(b_1) + 5w(b_2)$. We need to prove that $15w(d) + 5w(a) + 5 \max\{w(b_1), w(b_2)\} \geq W$ and $15w(d) + 5w(b_1) + 5w(b_2) \geq W$.

Let us prove the first one. Notice that $\max\{w(b_1), w(b_2)\} \geq \frac{w(t) - w(a)}{2}$. Suppose that $15w(d) + 5w(a) + 5 \max\{w(b_1), w(b_2)\} < W$. This means that $4w(a) + 5 \max\{w(b_1), w(b_2)\} > 6w(d)$. Using $w(d) = \frac{w(t)}{2} + \epsilon$, we get that $1.5w(a) + 2.5w(t) > 3w(t) + 6\epsilon$. Hence $w(a) > \frac{w(t)}{3} + 4\epsilon$. But then $w(d) < \frac{3}{2}w(a)$, which contradicts Lemma 17 point 1.

3. C_1 contains two crossing half-edges within $(p, q), (r, p)$ and one whole edge within c .

Then C_1 contains either (i) (p_1, p) and (q_1, q) or (ii) (p, p_1) and (q, q_1) . The case is similar to the case for the tricky 3-triangle. We do not modify anything in G_1 . W.l.o.g. assume that case (i) holds. Edges $(p_1, p_2), (q_1, q_2)$ of C'_{max} are called the **antennas** of t and required to be diverse.

4. C_1 does not contain any half-edges within c but contains a loop e'_t .

Since $10w(e'_t) = w(r, q)$, G_1 contains 5 copies of (r, q) (and not 4 as usual). Such an edge (r, q) is called a **bow**.

5. C_1 contains all edges of t and a loop e_t .

Then $10w(t) + 4w(c) + 10w(e_t) = 14w(q, r) + 10(w(p, q) + w(r, p)) + 3w(r, q)$ and G_1 indeed contains 14 copies of (q, r) , 10 copies of each of (p, q) , (r, p) and 3 copies of (r, q) , which is a path-20-colorable subgraph.

6.5 Strange 2-cycles

Finally, we show what we do about strange 2-cycles of C_1 . Let $c = (u, v)$ be a strange 2-cycle of C_1 and suppose C_{max} contains (u, v) , (u', u) , (v, v') .

Since c belongs to C_1 , it means that $4w(v, u) \leq 6w(v', v)$ or $4w(v, u) \leq 6w(u, u')$. Suppose that the first case holds. Then G_1 will contain 6 copies of (v, u) (instead of 10) and additionally, 6 copies of (v', v) . To facilitate the path-coloring of G_1 , G'_1 is modified as follows. If $v' = u'$, G'_1 contains a 2-cycle (u', u'') , where u'' is a new vertex and the number of copies of (u', u'') and (u'', u) is equal to respectively 10 and 5 and G'_1 does not contain vertices u, v or their adjacent edges. If $v' \neq u'$, G'_1 does not contain u or v or their adjacent edges and contains instead 4 copies of the edge (u', v') , which is treated as though it belonged to C'_{max} .

Theorem 3 $w(G_1) \geq 10w(C_1) + 4w(C_{max})$

The proof follows from the above discussion.

7 Path-coloring in the presence of tricky triangles

As previously, i.e., in Section 4, we would like to take advantage of Observation 1 and color rays in portions by coloring all rays of one cycle or path in one step. Here, however, the situation is somewhat more complicated because of b-edges. Consider an edge $e = (u, v)$ belonging to some path or cycle s of C_1 . Suppose that it is coincident with two rays $r = (u, u')$, $r' = (v', v)$ of this path or cycle, thus r is an inray and r' an outray of s . If r is a b-edge, meaning that $mult(r) = 10$, then it is impossible to color r and r' with disjoint sets of colors of \mathcal{K} . This is because, both r and r' have to be diverse with e . However, $mult(e) = 10$, which implies that afterwards the coloring it must hold that $col(e) \cup col(r) = \mathcal{K}$, hence $col(r')$ must be a subset of $col(r)$. To deal efficiently with coloring of b-edges, we divide the set of colors $col(r)$ assigned to any b-edge r into r 's **own colors**, denoted $col'(r)$ and **colors inherited** by r , denoted $col''(r)$. If in the above example, r' is not a b-edge, then colors inherited by r are such that $col''(r) = col(r')$ and r' is called an **ally** of r . Below we define **allies** for all b-edges. They help to control, which colors are inherited by which edges. For every b-edge, its inherited colors come from its ally.

The division is such that for any $r, r' \in C_{max}$ coincident with an edge e of C_1 such that r is a b-edge, it holds that (i) $col''(r) \supset col(r')$, if r' is not a b-edge, (ii) $col''(r) \supset col'(r')$, otherwise and (iii) $|col'(r)| = |col''(r)| = 5$ (which holds after r is fully colored). For example, if e is coincident with two b-edges r and r' , then it holds that $col(r) = col(r')$ and $col'(r) = col''(r')$, $col'(r') = col''(r)$, thus half of the set $col(r)$ are r 's own colors and half are inherited from r' .

Now, we define allies. Let $r = (u, v)$ be a b-edge. It is coincident with one edge of C_1 . Suppose that r is coincident with an edge $e_1 = (v_1, v)$ of C_1 . Then there exists an edge $r'_1 = (v_1, v'_1)$ belonging to C'_{max} . We call r'_1 an **ally** of r and denote as $al(r) = r'_1$. (If such an edge r'_1 does not exist, that we can add an artificial edge of this form. In reality it means that we have more flexibility in coloring r .) The situation is symmetric if r is coincident with an edge $e_2 = (u, u_1)$ of C_1 . Then the edge $r'_2 = (u_1, u'_1) \in C'_{max}$ is an **ally** of r and denoted as $al(r) = r'_2$.

Let us now examine what methods we can use to ensure that any b-edge is safe with respect to inherited colors. To this end, we appropriately define what it means for a halfy triangle to be **blocked** and **cooperative**. As for colors owned by b-edges we can apply Observation 1 to ensure their safety.

We say that two antennas a_1, a_2 of a halfy cycle c of C_1 are **diverse** if the sets of its own colors are disjoint, i.e., if $col'(a_1) \cap col'(a_2) = \emptyset$.

7.1 Halfy triangles

Let $t = (p, q, r)$ be a halfy triangle consisting of edges $a = (p, q), b = (q, r), c = (r, p)$ and $(p_1, p), (q_1, q), (q, q_2), (r_1, r)$ edges of C_1 and $(r_1, r'_1), (q''_2, q_2), (q_1, q'_1), (p_1, p'_1)$ edges of C'_{max} .

Suppose that G'_1 contains 10 copies of $d = (r, q)$ and 5 copies of b and C_1 contains edges $(q_1, q), (q, q_2), (r_1, r)$. We call b an **s-edge** of t . If $b' = (r_1, r'_1) \in C'_{max}$ does not belong to a halfy 2-cycle of C_1 , then b' is said to be an **outer antenna** of t and d an **inner antenna** of t . Since d is a b-edge, it has an ally d' and 5 colors assigned to it are inherited, i.e., $col''(d) \supset col'(d')$. To be able to guarantee that d is safe with respect to each inherited color $k \in col''(d)$, we require that the antennas of t are diverse (i.e. that $col'(d)$ and $col'(b')$ are disjoint) and we say that t is **blocked** if this condition is not satisfied. (The situation is symmetric if t contains $b = (r, q)$, G'_1 contains 10 copies of $d = (q, r)$ and 5 copies of b and C_1 contains edges $(q_1, q), (q, q_2), (r, r_1)$.)

Let $Z(t)$ denote a subset of $col''(d) \setminus col'(b')$ such that $k \in Z(t)$ if d is not already safe w.r.t. k at the moment of coloring d . $Z(t)$ may of course contain all colors of $col''(d) \setminus col'(b')$.

Notice that if we color b in such a way that we assign $|Z(t)|$ colors of $col'(b') \setminus col''(d)$ to b (i.e. $|(col'(b') \setminus col''(d)) \cap col(b)| \geq |Z(t)|$), then $flex^+(r_1, r) \geq |Z(t)|$, which means that we can forbid any color of $Z(t)$ on (r_1, r) and hence are able to color (r_1, r) so that no color of $Z(t)$ occurs on it (any color of $col'(b')$ does not occur on (r_1, r) anyway). This is how we are going to ensure that d is safe with respect to inherited colors. Observe that since the outer antenna b' is required to be diverse with the inner antenna d (i.e. $col'(d) \cap col'(b') = \emptyset$), there is no risk of assigning to b any color already assigned to d and hence creating a monochromatic 2-cycle.

To summarise, we say that edge b is **shadowed** or that we **shadow** b if (i) it is colored in such a way that $|(col'(b') \setminus col''(d)) \cap col(b)| \geq |Z(t)|$ and if (ii) no color of $Z(t)$ occurs on (r_1, r) . Also, if b' is a b-edge, then it views the edge b as though it were colored with colors of $Z(t)$ and treats them as $col'(b)$. Thus, if b' is a b-edge, then $col''(b') \supseteq Z(t)$.

For example, suppose that d' and b' are already colored and $col(d') = \{1, 2, 3, 4, 5\}$ and $col(b') = \{6, 7, 8, 9, 10\}$ and we want to color d and b (because we are processing a path or cycle of C_1 containing edge $(q_1, q), (q, q_2)$). Because d is a b-edge and d' is its ally, the colors $col''(d)$ inherited by d are $\{1, 2, 3, 4, 5\}$. We assign own colors of d to d so that they are disjoint with $col(b')$, for example, $col'(d) = \{11, 12, 13, 14, 15\}$. Therefore, $col(d) = col'(d) \cup col''(d)$. Next, we shadow b . Since, $Z(t) = \{1, 2, 3, 4, 5\}$, we assign all colors of b' to b , thus $col(b) = \{6, 7, 8, 9, 10\}$. Also, we assign colors of $\mathcal{K} \setminus (Z(t) \cup col(b))$ to (r_1, r) . Hence, $col(r_1, r) = \{11, 12, \dots, 20\}$. This means that d is safe w.r.t. each color of $col''(d)$.

We say that such a halfy triangle t is **cooperative** if

- t is not blocked,
- its s-edge is shadowed,
- b and d are diverse.

Suppose next that $t = (p, q, r)$ is a halfy triangle such that G'_1 contains 10 copies of each of $a = (p, q)$, $c = (r, p)$, 5 copies of $b = (q, r)$ and C_1 contains edges (p_1, p) , (q_1, q) , (q, q_2) , (r_1, r) . (The situation is symmetric when C_1 contains (p, p_1) , (q_1, q) , (q, q_2) , (r, r_1) .) We call b an **s-edge** of t , a the **main b-edge** of t and c the **secondary b-edge** of t . Both a and c are b-edges and their allies are denoted as a' and c' , respectively. An edge $b' = (r_1, r_2) \in C'_{max}$ is said to be an **ally** of the s-edge b .

If c' does not belong to a halfy 2-cycle of C_1 , then c' is said to be an **outer antenna** of t and a an **inner antenna** of t . A triangle t is said to be **blocked**, if its antennas are not diverse. Notice that since c' is an ally of c , the fact that antennas of t are diverse implies that $col''(c) \cap col'(a) = \emptyset$.

Whenever possible, we color c in such a way that each color k inherited by a ($k \in col''(a)$) is assigned to c . More precisely, for any inherited color k assigned to a , if k is not already assigned to c ($k \in col''(a) \setminus col''(c)$), we assign it to $col'(c)$ unless it is forbidden on c , because c is an antenna of some halfy cycle of C_1 . (It can be proved that if a color k is forbidden on c , then a is safe w.r.t. k .) Additionally, if $col'(c) \neq col''(a)$, then we ensure that $|col(a) \cap col(c)| \leq 5$. (If $col'(c) = col''(a)$, then $|col(a) \cap col(c)| \leq 5$ always holds.) For each color $k \in col(a) \cap col(c)$, we are going to guarantee that both a and c are safe w.r.t. k by *shadowing* b similarly as in the case above and as explained below.

Let $Z(t) = \{k \in (col(a) \cap col(c)) \setminus col'(b') : k \text{ is such that } a \text{ is not safe w.r.t. } k \text{ at the moment of coloring } a\}$. We want to ensure that a and c are safe with respect to each color $k \in Z(t)$. To this end, we color b in such a way that b is assigned at least $|Z(t)|$ colors of $col'(b')$, i.e., $|col'(b') \cap col(b)| \geq |Z(t)|$. Then we can color (r, r_1) so that no color of $Z(t)$ occurs on it, i.e., b is **shadowed**.

To sum up, we say that a halfy triangle t is **cooperative** if

- t is not blocked and $|col(a) \cap col(c)| \leq 5$,
- its s-edge is shadowed,
- no color k occurs on every edge of t .

Let $e(t)$ be a b-edge of a halfy triangle t , then an s-edge contained in t is said to be **associated** with $e(t)$.

7.2 Algorithm

When coloring rays of a cycle or path s of C_1 , we may not be able to color b-edges and s-edges incident to s fully, because their allies have not been colored yet. For this reason, we introduce the notion of precoloring. To **precolor** an edge r means to:

- color r , if r is neither a b-edge nor an s-edge,
- color r with 5 colors denoted as $col'(r)$, if r is a b-edge but not a secondary b-edge,
- leave r uncolored, if r is a secondary b-edge or an s-edge.

Below we show that we can guarantee that each ray of a given cycle c is safe by using a similar approach as previously, where we colored inrays and outrays with disjoint sets of colors. The modification consists in the fact that for b-edges, we only require that colors owned by them, i.e., sets $col'(r)$ obey this partition.

Lemma 18 *Let c be a cycle of C_1 such that each of its incident rays is uncolored or safe. Then we are able to precolor uncolored rays of c in such a way that under the condition that each halfy triangle incident to c is cooperative, each ray of c is safe.*

Proof. We partition \mathcal{K} into $Z^-(c)$ and $Z^+(c)$. We would like to color each uncolored inray of c with colors of $Z^-(c)$ and each uncolored outray of c with colors of $Z^+(c)$. For every uncolored ray r such that $\text{mult}(r) \leq 5$ and which is not an s-edge, this is indeed how we proceed. For every ray r of c , which is a b-edge, we assign 5 colors of either $Z^+(c)$ or $Z^-(c)$ to $\text{col}'(r)$, depending on whether r is an inray or an outray. Note that a ray of a cycle can never be a secondary b-edge, because both endpoints of a secondary b-edge belong to paths of C_1 . The other colors assigned to r are inherited from the ally of r .

As for any s-edge e , we have already observed that it can only belong to a monochromatic cycle c , which is a (sub)cycle of a halfy triangle t which contains e . If t is cooperative, then it is guaranteed not to happen.

Let k be a color assigned to some ray r of c , also possibly at some later point after the precoloring of c . If r is an s-edge, then we have already shown above that under the condition the halfy triangle t containing r is cooperative, r is safe. Assume next that r is not an s-edge. If r is an inray and $k \in \text{col}'(r)$, then if r was precolored before processing c , it is safe by the assumption. If r was precolored during processing c , then $k \in Z^-(c)$. Any potential monochromatic cycle c' containing r must contain some outray r' of c . If any outray r' of c is colored k , then r' was either colored k before we started precoloring rays of c or r' is an s-edge or k belongs to colors inherited by r' , i.e., $k \in \text{col}''(r')$. In all these three cases, however, r' is guaranteed to be safe w.r.t. k under the condition that a halfy triangle containing r' is cooperative. This means that r is safe under the condition that each halfy triangle incident to c is cooperative. \square

We say that an edge is **conditionally safe** if it is guaranteed not to belong to a monochromatic cycle under the condition that all halfy triangles are cooperative.

We say that G'_1 is blocked if there exists a cycle, a halfy 2-cycle or tricky triangle of C_1 that is blocked. Otherwise, G'_1 is unblocked.

To **process** a cycle or path s of C_1 means to precolor all its rays in such a way that all of them are conditionally safe and G'_1 is unblocked, assuming that before starting to process this cycle or path, G'_1 is safe and unblocked.

Algorithm Color7

```

while there exists an unprocessed cycle of  $C_1$  without any incident b-edge
     $c \leftarrow$  an unprocessed cycle of  $C_1$  without any incident b-edge with a minimal number of uncolored rays;
    process  $c$ ;

process all cycles of  $C_1$  with an incident b-edge;

while there exists an unprocessed path of  $C_1$ 
     $p \leftarrow$  any unprocessed path of  $C_1$ ;
    process  $p$ ;

color fully all b-edges and s-edges;
color the remaining uncolored edges in such a way that each of them is safe and  $G'_1$  does not become blocked.

```

By $\beta(c)$ we denote the number of bows incident to c .

Lemma 19 *Let c be a cycle of C_1 . Then $\chi(c) = \text{kol}(c) + \text{flex}^0(c) + \text{flex}^+(c) + \text{blank}(c) \geq \text{kol}(c) + 2\lambda(c) - \beta(c) + \text{flex}^+(c)$. If the number of uncolored rays and chords of c is equal to r , then $\chi(c) \geq \text{kol}(c) + 2\lambda(c) - \beta(c) + 4r + \text{flex}^+(c)$.*

As a consequence:

1. A cycle of length greater than 2 and one uncolored chord or two uncolored rays cannot be blocked.
2. A 2-cycle is blocked only if some two of its non-complementary rays r_1, r_2 are not diverse.

Proof. Let e be an edge of c . If none of its coincident edges of C'_{max} is colored, then $flex(e) = 10$. If exactly one of its coincident edges of C'_{max} is colored, then $flex(e) = 6$. Otherwise, $flex(e) \geq \max\{10 - kol(c), 2\}$. \square

We now present an extended version of Lemma 9. The main difference comes from the fact that G'_1 may contain bows, which occurs in point 5. Recall that a bow is an edge of multiplicity 5 contained in a 2-cycle of C'_{max} .

Lemma 20 *Suppose that at step S we want to color a set U of uncolored edges, where U consists of either (i) a subset of uncolored rays of a cycle c of C_1 or (ii) an antenna of a halfy cycle c of C_1 . Then, assuming that G'_1 is unblocked, there always exists a number $\Delta'(c)$ and a set $Z \subseteq \mathcal{K}$ such that by using $\Delta'(c)$ different colors of Z on U , we guarantee that c does not become blocked. Depending on additional conditions, $\Delta'(c)$ and $|Z|$ can be expressed as the following functions of a certain $\Delta(c) \leq \Delta'(c)$:*

0. *If c has at least two chords or one chord and $\lambda(c) > 3$, then $\Delta'(c) = 0$. In the remaining points we assume that c has no chords or one chord and $\lambda(c) = 3$.*
1. *If c is a 2-cycle with r colored rays, then $\Delta'(c) = mult(U)$ and $|Z| = 20 - 4r + \rho(c)$.*
2. *If c has one uncolored ray, no chords and $\lambda(c) > 2$, then $\Delta'(c) = 4 - \Delta(c) \geq 0$, where $\Delta(c) = flex^+(c) + kol(c) - 10$ and $|Z| \geq 12 - \Delta(c)$.*
3. *Assume that c has exactly two uncolored incident edges of C_{max} and $\lambda(c) > 2$. Then $|Z| \geq 12 - \Delta(c) + \rho(c)$, where $\Delta(c) = flex^+(c) + kol(c) - 8$. If we color only one ray of c , then $\Delta'(c) = 2 - \Delta(c)$, otherwise $\Delta'(c) = 6 - \Delta(c)$.*
4. *Assume that c has at least $u \geq 3$ uncolored rays and $\lambda(c) > 2$. Then $|Z| \geq 20 - flex^+(c) - kol(c) + \rho(c)$. If we color $u - 2$ rays of c , then $\Delta'(c) = 0$; if $u - 1$, then $\Delta'(c) = \min\{10 - flex^+(c) - kol(c), 0\}$; if we color all u rays of c , then $\Delta'(c) = \min\{14 - flex^+(c) - kol(c), 0\}$.*
5. *If U consists of an antenna of c , then $\Delta'(c) = 4$ and $|Z| \geq 15$.*

Proof. If c has no chords and r uncolored rays, then $\chi(c) = kol(c) + 2\lambda(c) - \beta(c) + 4r + flex^+(c)$.

Case: c has exactly one uncolored ray and no chords.

We can notice that if c has exactly one uncolored ray e incident to vertex $v \in c$, then it cannot belong to a 2-cycle c' of C'_{max} , because both edges of any 2-cycle of C'_{max} are colored during the same step. Thus at any point of the execution of Algorithm Color 7 either both edges of such 2-cycle are uncolored or both are colored. This means that e is not a bow and also that v has no incident bow. Therefore, $\beta(c) \leq \lambda(c) - 1$ and $mult(e) = 4$. Thus $2\lambda(c) - \beta(c)$ is minimum when $\beta(c) = \lambda(c) - 1$ and $\lambda(c) = 3$ and amounts to 4. By Lemma 8 there exists a set Z of colors the application of any color of which increases $flex^+(c) + kol(c)$. To guarantee that c does not become blocked it suffices to use $mult(e) - \min\{0, \chi(c) - 20\}$ colors of Z . Let

us estimate $\chi(c) - 20$. We have $\chi(c) - 20 = kol(c) + flex^+(c) + 2\lambda(c) - \beta(c) + 4 - 20 \geq kol(c) + flex^+(c) - 12$. Let $\Delta(c) = \min\{0, kol(c) + flex^+(c) - 12\}$. By Lemma 8 the size of Z is at least $24 - kol(c) \geq 12 + 12 - kol(c) - flex^+(c) = 12 - \Delta(c)$.

Case: c has exactly two uncolored incident edges e, f of C'_{max} .

Suppose first that the currently colored rays of c do not contain a bow. Then $kol(c) + flex^+(c) \geq 8$ and $flex^0(c) + blank(c) \geq 14$, because either e, f do not contain a bow and then $blank(c) = 8$ and $flex^0(c) \geq 6$ or e, f contain a bow and then $blank(c) = 9$ and $flex^0(c) \geq 5$. Hence $\chi(c) \geq 22$, because $\chi(c) = kol(c) + flex^+(c) + flex^0(c) + blank(c)$. Thus, to guarantee that c does not become blocked, it suffices to use $mult(U) - 2 - \min\{0, \chi(c) - 22\}$ colors of Z . We define $\Delta(c)$ as follows: $\chi(c) - 22 \geq kol(c) + flex^+(c) + flex^0(c) + blank(c) - 22 \geq kol(c) + flex^+(c) - 8 = \Delta(c)$.

Suppose next that the currently colored rays of c contain a bow. Then $kol(c) + flex^+(c) \geq 13$ (because we keep an invariant that if there exists an edge e of c with both rays colored and an incident bow, then $kol(c) + flex^+(c) \geq 13$ only among these three rays of c incident to e) and $flex^0(c) + blank(c) \geq 12 + x$, where $x = 1$ if e, f are coincident with the same edge of c and $x = 0$, otherwise. Hence, $\chi(c) \geq 25 + x$. For the case when $|U| = 1$, we can color the ray of U in any way, since currently $\chi(c) \geq 25$, hence $\Delta'(c) = 0$. For the case when $|U| = 2$, it suffices to use $mult(U) - 5 - x - \min\{0, \chi(c) - 25\}$ colors of Z . We define $\Delta(c)$ as follows: $\chi(c) - 25 = kol(c) + flex^+(c) + flex^0(c) + blank(c) - 25 \geq kol(c) + flex^+(c) - 12 - x = \Delta(c)$.

By Lemma 8 the size of Z is at least $20 - kol(c)$, when e and f are coincident with the same edge of c . If the currently colored rays of c do not contain a bow, then $20 - kol(c) \geq 12 + 8 - kol(c) - flex^+(c) = 12 - \Delta(c)$. Otherwise, $20 - kol(c) \geq 7 + 13 - kol(c) - flex^+(c) = 7 - \Delta(c)$.

When e and f are not coincident with the same edge of c , then the size of Z is at least $24 - kol(c)$, which is greater or equal $16 - \Delta(c)$, if c has no incident colored bow and greater or equal $12 - \Delta(c)$, otherwise. \square

Lemma 21 *Let c be an unprocessed cycle of C_1 that at some step of Algorithm Color7 has a minimal number of uncolored rays, no incident b-edges and $\lambda(c) > 2$. Then it is always possible to process c .*

Proof.

If c has exactly one uncolored ray r or its already colored rays do not contain a bow, then the proof is the same as in Lemma 10. Assume then now that c already has a colored ray, which is a bow. If c has two uncolored incident edges e, f of C'_{max} forming a set U , then we only have to use $mult(U) - 5 - x - \Delta(c)$ colors of the set Z , which has size at least $12 - \Delta(c)$ if e, f are not coincident with the same edge or $7 - \Delta(c)$, otherwise. The first case is analogous to those considered in Lemma 10. In the second one, e, f cannot contain a bow and thus we only have to use $8 - 5 - 1 = 2$ colors of Z , which is easily achieved.

If c has at least 3 uncolored rays, then the proof is almost the same as the proof of Lemma 10. \square

Lemma 22 *Let c be an unprocessed 2-cycle of C_1 that at some step of Algorithm Color7 has a minimal number of uncolored rays and no incident b-edges. Then it is always possible to process c .*

Proof. If $c = (v, u)$ has no incident bow, then the proof is almost the same as that of Lemma 11. Suppose now that c has an bow r incident to vertex u of c . It means that r belongs to a 2-cycle $c' = (u, u')$ of C_{max} and u' does not lie on c . If u' also lies on a 2-cycle $c_1 = (u', v')$ of C_1 , we also process c_1 , i.e., we process c and c_1 (and possibly some other 2-cycles) during the same step. W.l.o.g. we may assume that rays r_1, r_2 of c incident to v and rays r_3, r_4 of c_1 incident to v' are not bows. We then treat these two 2-cycles c, c_1 as though they were one 2-cycle of C_1 with four rays r_1, r_2, r_3, r_4 . We then complete the coloring on bows accordingly, i.e., if r_1 is an inray of c and r_3 an outray of c_1 , we assign any colors of $col(r_1) \cup col(r_3)$ to the edge (u, u') . Similarly, we assign any colors of $col(r_2) \cup col(r_4)$ to the edge (u', u) . \square

Lemma 23 *Let c be an unprocessed cycle of C_1 that has an incident b-ray. Then it is always possible to process c .*

Proof. We show that we can always ensure that c is not blocked by coloring the rays of c in such a way that $kol(c) + flex^+(c) + flex^0(c) \geq 20$.

To process c , we need to color all rays of c , which are not s-edges. We partition \mathcal{K} into two disjoint sets $Z^+(c)$ and $Z^-(c)$. If we precolor each uncolored outray with colors of Z^+ and each uncolored inray with colors of $Z^-(c)$, then each such newly colored ray becomes safe. As for s-edges, we have more freedom in coloring them and do not have to observe this partition. Recall that each s-edge e is guaranteed to be safe (as long as it is not assigned the same color k as the other b-edge(s) of the same halfy triangle).

While coloring the rays of c , we also have to ensure that no other cycle or halfy cycle of C_1 becomes blocked. We do not need to concern ourselves with blocking cycles of C_1 different from c , because cycles of C_1 with no incident b-rays are already processed and by the current lemma we are always capable of processing a cycle of C_1 with an incident b-ray. Thus we only have to take care of halfy cycles. Since each ray of c is an antenna of at most one halfy cycle, every ray of c has to be diverse with at most one edge. We make the following useful observation.

Claim 4 *Let r be a b-ray of c , r' an s-edge associated with r and k a color not occurring on any ray of c . Then we can always assign k to at least one of r, r' and be able to color the halfy triangle containing r in a cooperative manner.*

Proof. Let t be a halfy triangle containing r . If k is not forbidden on r , then we are done. Suppose then that k is forbidden on r , which means that k is assigned to the outer antenna a of t - we have a requirement that r has to be diverse with a . If t is a tricky 2-triangle, then it means that while shadowing r' , we will assign colors of $col'(a)$ to $col'(r')$ and hence we will assign k to r' . More precisely, we will assign k to r' as long as $k \notin col''(r)$. However, if $k \in col''(r)$, then k is assigned to r . Either way k can appear on r or r' . If t is a tricky 3-triangle and k is assigned to a , then we do not want to assign k to r , because we then would have to shadow r' w.r.t. k and possibly additional 5 inherited colors of r . But we can notice that if k does not occur on any ray of c , we do not have to shadow r' w.r.t. k , because r is safe w.r.t. it. Even if k is assigned to inherited colors of some b-ray r_1 of c , then we make sure that r_1 is safe w.r.t. all inherited colors. This completes the proof. \square

In view of the above claim, we notice that if the number of uncolored rays of c , which are not s-edges is at least 4, then we can easily guarantee that $kol(c) + flex^0(c) \geq 20$ by assigning different colors to $col'(r)$ of each uncolored ray or by assigning them to an appropriate s-edge.

(For example, suppose that c has one b-ray r , one s-edge r' and 3 uncolored rays and for each of the rays the same 5 colors of Z' are forbidden. Then by Claim 4 we can assign Z' to r' and use colors of $\mathcal{K} \setminus Z'$ on the uncolored rays.) The situation is analogous if c has already one colored ray and the number of uncolored rays of c , which are not s-edges is at least 3 or generally if $kol(c) + flex^+(c) + flex^0(c) + blank(c) - 5|\{\text{s-edges incident to } c\}| \geq 20$. In all these cases we are simply able to use $20 - kol(c) - flex^+(c) - flex^0(c)$ new colors (not already included in $col(c)$) on the uncolored rays or s-edges.

Let us next observe that in the situation when all uncolored rays of c , which are not s-edges, are either all outrays or all inrays, all rays of c are safe under the condition that each b-ray of c is diverse with an associated s-edge. Thus we do not have to shadow s-edges incident to c and can use new colors on them. This means that in such situations we are always able to use $20 - kol(c) - flex^+(c) - flex^0(c)$ new colors on uncolored rays of c , because $\chi(c) = kol(c) + flex^+(c) + flex^0(c) + blank(c) \geq 20$. We can use the same argument for any 2-cycle with exactly one incident b-ray, even if it has two uncolored inrays or two uncolored outrays.

To illustrate the above reasoning consider the following example. Suppose that c has only two uncolored rays: an incoming b-ray r and an s-edge r' associated with it. In this case c already has at least 4 colored rays (or it has some number of colored rays and chords). Thus it already holds that $kol(c) + flex^+(c) + flex^0(c) \geq 10$. Also, the ally al of r is already colored. Let $Z' = col'(al)$. We of course have to assign Z' to r . Since c does not have an uncolored outray, which is not an s-edge, r is safe w.r.t. every color of Z' . Hence we do not have to shadow r' . Next we assign 5 new colors to r and 5 different new colors to r' . This way we increase $kol(c)$ by 10. As a result $kol(c) + flex^+(c) + flex^0(c) \geq 20$.

We are thus left with the following three cases: c has one b-inray and one b-outray and either (i) one more b-ray and no other chords or rays or (ii) two rays incident to the same vertex and $kol(c) + flex^+(c) + flex^0(c) < 10$ or (iii) c is a 2-cycle.

In the first two cases it means that c is a triangle. Therefore, in all these three cases one of the edges of c is coincident both with a b-outray r_1 and a b-inray r_2 . Let us note that r_1, r_2 will be colored with the same 10 colors belonging to the set Z and none of the s-edges s_1, s_2 associated with, respectively, r_1 and r_2 will be colored with any element of Z . Hence, r_1, r_2 are going to contribute 10 new colors to $col(c)$ and s_1, s_2 are going to either contribute at least another 5 to $kol(c)$ or increase $flex^+(c)$. Either way, s_1, s_2 will increase $kol(c) + flex^+(c)$ by 5. If c already has some colored rays or is 2-cycle, then it means that we have already guaranteed that $kol(c) + flex^+(c) + flex^0(c)$ reaches at least 20. If we have the first case (i), then let r_3 be the third b-ray of c . Let us observe that r_3 's ally is either s_1 or s_2 . It suffices, if we precolor r_3 with colors disjoint with Z . In this way the contribution of r_1, r_2, r_3 to $col(c)$ amounts to 15 colors and edges s_1, s_2 will increase $kol(c) + flex^+(c)$ by another 5. \square

Fact 5 1. Each edge of C'_{max} is an antenna of at most two halfy cycles of C_1 .

2. If an edge $e \in C'_{max}$ is an antenna of two halfy cycles, then it is not incident to a cycle of C_1 .

Lemma 24 It is always possible to process a path of C_1 .

Proof. Since paths are processed after cycles of C_1 , only halfy cycles of C_1 can become blocked. To prevent this, we have to ensure that antennas of the same halfy cycle are diverse or weakly diverse. The path p has two outer antennas a_1, a_2 and at most one of them is an inray and

at most one the outray of p . (Each one of them may also be a chord of p .) Assume that a_1 is an inray and a_2 the outray. Each a_i may have to be diverse with two different antennas. Additionally, each a_i may be accompanied by either an inner antenna of the same halfy cycle or a weak antenna. In each case we call it a'_i . Observe that any inner antenna a'_i may also have to be diverse with two other antennas, one of which is always a_i . A weak antenna a'_i only needs to be weakly diverse with a_i , but may have to be diverse with some other antenna b_j . The case is most difficult when all four antennas exist and all are bilateral. Note that no other ray of p is a bilateral antenna. Let Z_i and Z'_i denote the set of colors forbidden on a_i and a'_i , respectively. Let us note that if a_i is uncolored, then $|Z'_i| \leq 5$. If a'_i is an uncolored inner antenna, then $|Z_i| \leq 5$. If a'_i is a weak antenna (or if p has no antenna a'_i), then $|Z_i|$ may be equal to 10. Assume that none of the four antennas is already colored, as the other cases are contained in this one.

Suppose first that a'_1 and a'_2 are weak antennas. We partition \mathcal{K} into two 10-element sets $Z^-(p)$ and $Z^+(p)$ so that $|Z^-(p) \setminus Z_1| \geq 5$, $|Z^-(p) \setminus (Z_1 \cup Z'_1)| \geq 7$ and $|Z^+(p) \setminus Z_2| \geq 5$, $|Z^+(p) \setminus (Z_2 \cup Z'_2)| \geq 7$. To this end, we divide $Z = \mathcal{K} \setminus (Z_1 \cap Z_2)$ (almost) equally between $Z^-(p)$ and $Z^+(p)$ in such a way that $Z \setminus Z_1$ goes to $Z^-(p)$ and $Z \setminus Z_2$ to $Z^+(p)$. Since $|Z_1 \cap Z_2| \leq 10$ and hence $|Z| \geq 10$ each of $Z^-(p) \setminus Z_1, Z^+(p) \setminus Z_2$ contains at least 5 colors. Additionally, we divide $Z' = ((Z_1 \cap Z_2) \setminus (Z_1 \cap Z_2))$ (almost) equally between $Z^-(p)$ and $Z^+(p)$ in such a way that $Z' \setminus Z'_1$ goes to $Z^-(p)$ and $Z' \setminus Z'_2$ to $Z^+(p)$. Since $|Z'_1 \cap Z'_2| \leq 5$, we get that $\frac{1}{2}(|Z| + |Z'|) \geq \frac{1}{2}(20 - |Z_1 \cap Z_2| + |Z_1 \cap Z_2| - |Z'_1 \cap Z'_2|) \geq 7$, which means that each of the sets $Z^-(p) \setminus (Z_1 \cup Z'_1), Z^+(p) \setminus (Z_2 \cup Z'_2)$ contains at least 7 elements. To finish the partition of \mathcal{K} , we divide the set $Z'_1 \cap Z'_2$ in such a way that each of the sets $Z^-(p)$ and $Z^+(p)$ has exactly 10 elements.

We can check that having such sets $Z^-(p)$ and $Z^+(p)$ enables us to color each ray of p so that each antenna is (weakly) diverse with required antenna. For any ray r , which is not any of the four antennas a_1, a_2, a'_1, a'_2 , r has at most 5 colors forbidden on it and since each of the sets $Z^-(p), Z^+(p)$ contains 10 elements, we are able to color r in a required manner. To color a_1 and a'_1 , we assign to them colors of $Z^-(p) \setminus (Z_1 \cup Z'_1)$ (of which there are at least 7) so that $|col'(a_1) \cap col'(a'_1)| \leq 3$.

Suppose next that a'_1, a'_2 are inner antennas. Then we want to partition \mathcal{K} into two 10-element sets $Z^-(p)$ and $Z^+(p)$ so that $|Z^-(p) \setminus (Z_1 \cup Z'_1)| \geq 10$ and $|Z^+(p) \setminus (Z_2 \cup Z'_2)| \geq 10$. Observe that it is not possible if $Z_1 \cap Z_2 \cap Z'_1 \cap Z'_2 \neq \emptyset$. However, this can be avoided, by requiring, for example that antennas b_1, b'_1 are diverse (thus b_1 has to be diverse with c_1 and b'_1 instead of c_1 and a_1 at the moment of coloring b_1), where b_1, b'_1 are antennas, with which a_1, a'_1 have to be diverse. The case when one of a'_1, a'_2 is a weak antenna and the other an inner one is very similar.

After computing such a partition, we are able to color each ray of p so as to ensure that each antenna is (weakly) diverse with required antennas. \square

7.3 Full coloring of b-edges and s-edges

During the processing of paths and cycles of C_1 we do not assign colors $col'(e)$ for any edge e , which is either a secondary b-edge or an s-edge. As a result some b-edges cannot be assigned their inherited colors $col''(e)$. This happens for any b-edge, whose ally is an s-edge or a secondary b-edge.

To be able to complete the process of coloring all b-edges and s-edges, we introduce a directed graph $D = (V_D, E_D)$, which shows the dependencies between halfy triangles containing

such edges. The vertex set of D consists of all halfy triangles. For a halfy triangle t , we denote by v_t a vertex representing it in D . The edge set E_D contains an edge $(v_t, v_{t'})$ iff the ally of the main b-edge of t' is either an s-edge of t or a secondary b-edge of t . The direction of an edge $(v_t, v_{t'})$ reflects the fact that t needs to be fully colored to be able to complete the coloring of t' . Note that each vertex of V_D has at most one incoming edge and at most two outgoing edges. To ***D-process*** a directed path or cycle s of D means to complete the coloring of each halfy triangle t corresponding to any vertex on s in such a way that G_1 remains unblocked.

Notice that any two cycles of D are vertex-disjoint.

Lemma 25 *It is possible to color an s-edge of a tricky 2-triangle t in such a way that it does not prevent any halfy triangle from being colored in a cooperative manner.*

Proof. If the b-edge $e(t)$ of t is already fully colored, then we color the s-edge $s(t)$ in a standard manner described earlier. Assume now that $e(t)$ is not fully colored. It means that the ally $al(e(t))$ of $e(t)$ is not precolored, i.e., its own colors are not yet assigned. Therefore, $al(e(t))$ is an s-edge or a secondary b-edge of some tricky triangle t_0 . Let $a(t)$ denote the antenna of t and e_{01} an edge of C_1 coincident with both $e(t)$ and $al(e(t))$.

If $a(t)$ is already precolored and no color $k \in col'(a(t))$ is forbidden on $s(t)$, then we claim that we can assign colors of $col'(a(t))$ to $s(t)$. To see this, suppose first that $al(e(t))$ is an s-edge of t_0 . Then $e(t_0)$ and e_{01} form a directed path P of length 2. If $e(t)$ is an antenna of t_0 (when t_0 is a 2-triangle) or an ally of $al(e(t))$ (when t_0 is a 3-triangle), then the standard procedure of shadowing $al(e(t))$ would involve assigning colors of $col'(e(t))$ to $al(e(t))$ and forbidding colors of $col''(e(t_0))$ on e_{01} by assigning colors of $col''(e(t_0))$ to $col''(e(t))$. We still do this if $col''(e(t_0)) \cap col(s(t)) = \emptyset$. On the other hand, if some color $k \in col''(e(t_0)) \cap col(s(t))$, then we can notice that $e(t_0)$ is already safe w.r.t. k , because $e(t_0)$ cannot belong to a monochromatic cycle of color k , since such a cycle would have to contain $s(t)$, which is impossible. This means that we do not have to do shadowing for any such color k . If $e(t)$ is neither an antenna of t_0 (when t_0 is a 2-triangle) nor an ally of $al(e(t))$ (when t_0 is a 3-triangle), then the reasoning is similar and the following. Let $al(al(e(t)))$ denote the ally of $al(e(t))$. Instead of doing the shadowing with the aid of $al(al(e(t)))$, we can instead use $e(t)$ and then the proof of the claim for the case when $al(e(t))$ is an s-edge of t_0 goes through. Suppose next that $al(e(t))$ is a secondary b-edge of t_0 , then again $e(t_0)$ and e_{01} form a directed path P of length 2 and for any $k \in col''(e(t_0)) \cap s(t)$, the edge $e(t_0)$ is safe w.r.t. k . Thus we do not assign it to $al(e(t))$ or hence also not assign it to $col''e(t)$. This finishes the proof of the claim.

Suppose next that $a(t)$ is already precolored but some colors of $col'(a(t))$ are forbidden on $s(t)$. It means that $s(t)$ is an antenna of some tricky triangle t_2 . Let e_{12} denote an edge of C_1 coincident with $s(t)$ and incident to e_{01} - the edges e_{01}, e_{12} form a directed path P_1 . We can notice that then $e(t_0)$ is safe w.r.t. any color $k \notin col'(e(t_2))$, because P_1 together with $e(t_2)$ form a directed path of length 3 and no color $k' \in col''(e(t_0))$ can occur on e_{12} . Therefore, we do not shadow $s(t_0)$ - instead we assign $col'(e(t_2))$ to $e(t)$, which ensures that $e(t_0)$ is safe w.r.t. any color. Also we assign $s(t)$ any 5 colors not forbidden on it, i.e. disjoint with $col'(e(t)) \cup col'(e(t_2))$.

Finally, we consider the case when $a(t)$ is not yet precolored. If $a(t)$ is a secondary b-edge of a 3-triangle t_3 , we can assign to $s(t)$ and $a(t)$ the same 5-element set of colors disjoint with $col'(e(t))$. Note that we do not have to do the shadowing for inherited colors of $e(t)$, because $col'(e(t)) = col'(e(t_3))$ and hence $e(t)$ is safe w.r.t. any color $k \in col''(e(t))$ (because such color either does not occur on $e(t_3)$ or on an edge of C_1 connecting t and t_3). If $a(t)$ is an s-edge, then for similar reasons we can assign to $s(t)$ and $a(t)$ the same 5-element set of colors disjoint with $col'(e(t))$. \square

Lemma 26 *It is possible to color an s-edge $s(t)$ of a tricky 3-triangle t in such a way that it does not prevent any halfy triangle from being colored in a cooperative manner. Moreover, if $s(t)$ is an ally of a main b-edge $e(t')$ of some halfy triangle t' , then it is possible fully color $e(t')$.*

Proof. We proceed in the manner described in the proof of the lemma above, where the ally of the s-edge of t plays the same role as the outer antenna of a tricky 2-triangle in Lemma 25. The only difference is that the ally a of the s-edge of t may be colored in the same way as the main b-edge of t . Then we can still assign colors of $col'(a)$ to $s(t)$ because the secondary b-edge of t is diverse with $e(t)$ i.e., $col''(e'(t)) \cap col'(e(t)) = \emptyset$.

If $s(t) = (u, v)$, then C_1 contains edges $(u, u_1), (v_1, v)$ coincident with $s(t)$ and C_{max} contains edges $a = (u_2, u_1), a' = (v_1, v_2)$. If the ally a of the s-edge of t is the main b-edge of a halfy triangle t' , then instead of assigning colors of $col'(a)$ to $s(t)$, we may assign colors of $col'(a')$ to it and then a' plays the role as the outer antenna of a tricky 2-triangle in Lemma 25.

If $col'(a) = col'(a')$ (more precisely, if $col'(a) \cap col'(a') \neq \emptyset$), then it still means that $e(t')$ is not colored fully. In such a case, we assign to $col''(e(t'))$ any 5 color set Z disjoint with $col(e(t))$. If later on, $col''(e(t))$ turns out to be different from Z , we can shadow $s(t)$ using a' , because $col(s(t)) = col'(a')$. \square

At this point the only not fully colored edges are either the main or the secondary b-edges of halfy triangles. Moreover, the main b-edge $e(t)$ of a halfy triangle t is not fully colored only if (i) its ally is the secondary b-edge of some halfy triangle t' or (ii) its ally is an s-edge $s(t')$ of some halfy triangle t' and $e(t)$ is the ally of $s(t')$, which also means that if t' is a tricky 2-triangle, then $e(t)$ is the antenna of t' .

Lemma 27 *It is possible to D-process each cycle of D .*

Proof. Let c be any cycle of D . By saying that a triangle t is on c , we mean that $v_t \in c$. Similarly, by $(t', t) \in c$ we mean $(v_{t'}, v_t) \in c$.

Let (t', t) be any edge of c . Let Z_1 denote $col'(e(t'))$. If t' is a 3-triangle, then Z_2 denotes $col''(e'(t'))$. Otherwise, if t' is a 2-triangle, then $Z_2 = col'(e(t'))$. Notice that Z_1 and Z_2 are disjoint (because the outer antenna of t' is equal to $e(t)$ if t' is a 2-triangle.) We assign any 5-color set Z disjoint with both Z_1 and Z_2 to $col''(e(t'))$. Next we color fully t' and all succeeding tricky triangles using only colors of $Z_3 = Z \cup Z_1 \cup Z_2$ on main b-edges.

We now argue that this is possible for each triangle t_1 on c . Suppose first that t_1 is a 2-triangle and we have just fully colored its main b-edge so that $col''(e(t_1)) \subset Z_3$. Let t_2 denote the tricky triangle succeeding t_1 on c . The ally of $e(t_2)$ is $s(t_1)$ as well as the antenna of t_1 is $e(t_2)$. We want to shadow $s(t_1)$ w.r.t. colors of $col''(e(t_1))$. We recall that if some color $k \in col''(e(t_1)) \cap col(s(t_2))$, then we do not shadow $s(t_1)$ w.r.t. k because $e(t_1)$ is already safe w.r.t. k . We assign all colors of $col''(e(t_1)) \setminus col(s(t_2))$ to $col''(e(t_2))$. If there are fewer than 5 of them, we add some colors of Z_3 .

Suppose next that t_1 is a 3-triangle and we have just fully colored its main b-edge $e(t_1)$ by adding 5 colors of Z_3 , i.e. $col''(e(t_1)) \subset Z_3$. We assign $col''(e(t_1)) \setminus (col''(e'(t_1)) \cup col(s(t_1)))$ to $col'(e'(t_1))$. Recall that we shadow $s(t)$ only w.r.t. colors of $col''(e(t_1)) \cap col(e'(t_1))$. If the successor t_2 of t_1 on c is such that the ally of $e(t_2)$ is $s(t_1)$, then we assign also $col''(e(t_1)) \setminus (col''(e'(t_1)) \cup col(s(t_1)))$ to $col''(e(t_2))$ and if there are fewer than 5 such colors, we complete the

set with an appropriate number of colors of Z_3 . Note that we also independently add some colors to $col'(e'(t_1))$, if there are already fewer than 5. These colors do not have to belong to Z_3 . If the ally of $e(t_2)$ is $e'(t_1)$, we also complete the set $col'(e'(t_1))$ with colors of Z_3 . Since we want to avoid at most $|col'(e(t_1)) \cup col''(e'(t_1))| \leq 10$ colors and $|Z_3| = 15$ we can always do that.

This way when we return to the triangle t' , we will not have to recolor it much. Anyway, we will be able to do it in such a way that the coloring of none of the remaining triangles on c will have to be changed. More precisely, consider the edge (t'', t') . Since t'' is now fully colored, $e(t')$ inherits some colors from one of the edges of t'' . The inherited colors are a subset of $Z \cup Z_1 \cup Z_2$. If the inherited colors are Z_1 or Z , then we assign Z to $e(t)$ and Z_2 to $s(t)$. If the inherited colors are Z_2 , then we leave the coloring as it is. In the case that the inherited colors I contain i colors of $Z \cup Z_1$ and $5 - i$ colors of Z_2 , we proceed analogously, i.e. we assign to $s(t)$: i colors of $Z_2 \setminus I$ and $5 - i$ colors of $Z \setminus I$. □

Lemma 28 *Let t be a tricky triangle with exactly two incident edges e_1, e_2 of C_1 and such that either both these edges are incoming or both are outgoing. Then it is possible to color the edges of t and e_1, e_2 in such a way that G'_1 is unblocked.*

Proof. Suppose that $t = (p, q, r)$ and $e_1 = (p', p), e_2 = (q', q)$. Assume also that a 2-cycle (q, r) belongs to C_{max} and that C'_{max} contains edges $(p', p''), (p, p'''), (p_4, p), (q', q'')$. W.l.o.g. suppose that $col(p', p'') = \{6, 7, 8, 9, 10\}, col(q', q'') = \{16, 17, 18, 19, 20\}$. Since (p, p''') is weakly diverse with (p', p'') at least 2 colors of (p, p''') do not occur on (p', p'') . This means that there exist 3 colors of $col(p', p'')$ which occur neither on e_1 nor on (p, p''') nor on (p_4, p) . Suppose that these are colors 8, 9, 10. We assign them to (r, p) . □

8 Return from G'_1 to G_1

While building the multigraph G_1 , we have modified it in two types of places (tricky 2-triangles and strange 2-cycles) creating the multigraph G'_1 .

Lemma 29 *Given a path-20-coloring of G'_1 , we can obtain a path-20-coloring of G_1 .*

Proof. Let $t = (p, q, r)$ be a tricky triangle of C_1 such that in G'_1 we have replaced the edges $(p, q), (q, q_1)$ with one edge.

We color the edges as follows. Edge (q, q_1) is colored in the same way as (p, q_1) in G'_1 . Suppose first that $w(r, p) \geq w(p, q)$. Edges $(p_2, p) \in C'_{max}$ and (p_1, p) are colored with at most 15 colors of \mathcal{K} . We color (r, p) with 5 colors of $\mathcal{K} \setminus (col(p_2, p) \cup col(p_1, p))$. For each color $k \in col(r, p)$, if $k \in col(p, p_3)$, then we assign k to (q, r) ; otherwise we assign k to (p, q) . Note that k cannot be assigned to (q, q_1) , because (p, q_1) is coincident with (p, p_3) in G'_1 . Hence (p, q_1) and (p, p_3) have to be diverse in G'_1 . Next we assign all 15 colors of $\mathcal{K} \setminus col(r, p)$ to (r, q) . We easily notice that each of the edges of t as well as the edge (r, q) are safe.

Suppose next that $w(r, p) < w(p, q)$. This case is, in fact, easier than the one above. We choose 5 colors from the set $\mathcal{K} \setminus (col(p, p_3) \cup col(q, q_1))$ and assign them to both (p, q) and (q, r) . Next, we assign all 15 colors of $\mathcal{K} \setminus col(p, r)$ to (r, q) . □

9 Completing the path-coloring

Let S denote a set of strange 2-cycles of C_1 .

We show that there exist so called *exchange sets* E_1 and F_1 with the following properties.

Lemma 30 *There exist sets $E_1, F_1 \subset E$ and an assignment $f : E_1 \rightarrow F_1$ satisfying the following conditions.*

1. Let c be any strange 2-cycle of C_1 . Then E_1 contains one edge e_1 of c and $f(e_1)$ is an edge of C_{max} incident to c (but not contained in c).
2. $F_1 = f(E_1)$ is a matching, i.e., no two edges of F_1 share a vertex.
3. $4w(E_1) \leq 6w(F_1)$.

Proof. For each strange 2-cycle c , the set E_1 contains an edge of c with minimum weight.

Let d be a cycle of C_{max} with length greater than 3 such that at least one strange 2-cycle of C_1 shares an edge with d .

Consider a 2-cycle $c = (u, v)$ belonging to S such that the edge (u, v) belongs to d . Let $(u', u), (v, v')$ be the edges of d adjacent to c . We call (u', u) an *incoming neighbour* of c and (v, v') an *outgoing neighbour* of c . If c is not incorrigible, then $\min\{w(u, v), w(v, u)\} \leq \frac{3}{4}(w(u', u) + w(v, v'))$. If d shares an edge only with strange 2-cycles which are not incorrigible, we set F_1 as either the set of all incoming neighbours or the set of all outgoing neighbours, choosing the one with maximum weight. Since strange 2-cycles are vertex-disjoint, the obtained set F_1 is a matching.

If c is incorrigible, then $w(u, v) \leq \max\{w(u', u), w(v, v')\}$ and F_1 contains the neighbour of c with maximum weight. If that neighbouring edge e is also adjacent to another cycle c' of S , then 6 copies of e are sufficient for removing 4 copies of (u, v) and 4 copies of an edge of c' . \square

Let R' denote the set of all tricky triangles of C_1 . They correspond to a matching N' of H . Notice that $N \cap N' = \emptyset$, because no tricky triangle of R (corresponding to N) can occur in C_1 . Thus $N \cup N'$ forms a set of alternating paths and cycles. Since N is a maximum matching of H , each alternating path P that contains at least one edge of N' has even length - thus the number of edges of N' on P equals the number of edges of N . For each alternating cycle and each alternating path of even length we replace some edges of triangles of R' with edges belonging to triangles represented by edges of N belonging to the same path or cycle. More precisely, suppose that an alternating path P or cycle C consists of a sequence of edges $e_1, f_1, \dots, e_i, f_i, \dots, e_k, f_k$ such that for $1 \leq i \leq k$ it holds that $e_i \in N', f_i \in N$ and edges e_i, f_i have a common vertex in $V(H)$. Then we replace some edges of each tricky triangle t_i of C_1 corresponding to edge e_i with some edges of a tricky triangle (not occurring in C_1) corresponding to edge f_i .

We now describe the exact procedure of replacement.

Let $t_i = (p, q, r)$ be a tricky triangle of C_1 with a t-cycle $c_i = (q, r)$. Recall that $\Delta(c) = w(r, q) - 1.5w(q, r)$. In G_1 we take 14 copies of (q, r) , 10 copies of each of $(p, q), (r, p)$ and 3 copies of (r, q) . This means that we are lacking only one copy of (r, q) , i.e.,:

Fact 6 *The weight of the induced subgraph $G_1(t_i)$ of G_1 on vertices p, q, r satisfies:*

$$w(G_1(t_i)) = 4w(c_i) + 10w(t_i) - w(q, r).$$

Consider alternating paths and cycles of $N \cup N'$. Each one of them consists of some sequence of edges $e_1, f_1, \dots, e_i, f_i, \dots, e_k, f_k$. For any alternating cycle C , we can additionally arrange the edges on C so that that a common vertex of any two edges e_i and f_i on C in H corresponds to a 2-cycle c_i . Let (e_i, f_i) be any pair of edges from such alternating cycle or path and suppose that a tricky triangle t_i of C_1 corresponding to e_i has the form $t_i = (p, q, r)$. If the common vertex of e_i and f_i in H corresponds to a 2-cycle c_i , then a tricky triangle represented by f_i has the form $t'_i = (p', q, r)$. We add either (p', q) or (r, p') to F_2 (and also 3 copies of the edge added to F_2 to G_1). If F_1 contains an edge incoming to p' , we choose (p', q) , otherwise (r, p') . If, on the other hand, the common vertex of e_i and f_i in H corresponds to the vertex p , then a tricky triangle represented by f_i has the form $t'_i = (p, q', r')$. In this case we add either (p, q') or (r', p) to F_2 (and also 3 copies of the chosen edge to G_1). If F_1 contains an edge incoming to p , we choose (p, q') , otherwise (r', p) . We call a tricky triangle t'_i of R corresponding to the edge f_i a *rescuer* of t_i .

In the next lemma we are going to prove that the total weight of edges added to G_1 makes up for the deficiencies in the weights of the subgraphs induced by vertices of tricky triangles of C_1 .

Lemma 31 *Let N'_2 denote the set of all t-cycles of tricky triangles of C_1 . If c is a t-cycle of a tricky triangle, then $\alpha(c)$ denotes the weight of the lighter edge of c . We have:*

$$\sum_{c \in N'_2} 1.5\alpha(c) + \Delta(c) \leq 3w(F_2)$$

To prove it we show the following lemma.

Lemma 32 *Let c_1, c_2 be two 2-cycles such that $w'(c_1) = w'(c_2)$ and let $\mu(c_i)$ denote the minimum weight edge of a tricky triangle incident to c_i . Then $3\mu(c_1) \geq 1.5\alpha(c_2) + \Delta(c_2)$.*

Proof. Suppose that $\alpha(c_2) = \alpha(c_1) + \epsilon$, $\Delta(c_2) = \Delta(c_1) - \epsilon$.

We know that $\mu(c_1) > 0.6\Delta(c_1) + \frac{\alpha(c_1)}{2}$. We show that $3(0.6\Delta(c_1) + \frac{\alpha(c_1)}{2}) \geq 1.5\alpha(c_2) + \Delta(c_2)$. This is equivalent to $0.8\Delta(c_1) \geq \frac{\epsilon}{2}$, which is true because $\Delta(c_1) \geq \epsilon$.

Next we show that $3(\frac{3}{5}(\Delta(c_1) - \epsilon) + \frac{1}{2}(\alpha(c_1) + \epsilon)) \geq \frac{3}{2}\alpha(c_1) + \Delta(c_1)$. This is equivalent to $\frac{4}{5}\Delta(c_1) \geq \frac{3}{10}\epsilon$, which holds because $\Delta(c_1) \geq \epsilon$. \square

Next we show that we are able to extend the current path-coloring of G_1 to the subgraphs containing strange 2-cycles and tricky 2-triangles of C_1 .

We start with subgraphs containing tricky triangles of C_1 . We proceed in the order dictated by directed paths and cycles of a graph H^{dir} , which is a compressed and directed version of the graph H . H^{dir} is obtained from H as follows. For each tricky triangle t of C_1 we identify as one vertex v_t four vertices in total: all vertices of t as well as the t-cycle c of t . Let $e = (u, v)$ be any edge of N . It then corresponds to a tricky triangle t' of R . If t' is a rescuer of a tricky triangle t of C_1 , we direct the counterpart of e in H^{dir} from v_t .

We first deal with directed cycles of H^{dir} .

Lemma 33 *Let c_H be any directed cycle of H^{dir} . We are able to extend the partial coloring of G_1 to the edges of tricky triangles covered by c_H and the edges of F_2 of their rescuers.*

Proof. Let t_1, \dots, t_k be the order of tricky triangles of C_1 , in which they (or more precisely, the vertices representing them) occur on c_H . Assume that each t_i has the form $t_i = (q_i, r_i, p_i)$, where (q_i, r_i) is a t-cycle of t_i . This means that for each $1 \leq i \leq k$ a rescuer t'_i of t_i has

the form $t'_i = (q_i, r_i, p'_i)$, where p'_i lies on t_{i+1} (indices are taken modulo k). The vertex p_i is incident to two edges $e_i = (s_i, p_i)$, $e'_i = (p_i, s'_i)$ belonging to C_{max} , which are already colored. We can assume that e_i and e'_i are diverse. Thus we use either 8 or 9 edges of \mathcal{K} on e_i and e'_i . We consider each t_i in turn. We assign 3 colors either to $f_i = (r_i, p'_i)$ or to $f'_i = (p'_i, q_i)$. We do it in such a way that:

- A color assigned to f_i or f'_i does not occur on any of $e_{i+1}, e'_{i+1}, f_{i-1}, f'_{i-1}$.
- A color occurring on e_i may be assigned to f_i but not f'_i . Similarly, a color occurring on e'_i may be assigned to f'_i but not f_i .

We now show that we are able to assign colors to each f_i or f'_i to satisfy the above. Suppose that we consider t_i . At most 9 colors of \mathcal{K} are used on e_i, e'_i . Possibly, t_{i-1} was considered before and thus one of f_{i-1}, f'_{i-1} is already colored with 3 colors. Let $Z_1 = col(e_i) \cup col(e'_i) \cup col(f_{i-1}) \cup col(f'_{i-1})$, $Z_2 = col(e_{i+1}) \cup col(e'_{i+1}) \cup col(f_{i+1}) \cup col(f'_{i+1})$ and $d = \max\{|col(e_i) \setminus Z_2|, |col(e'_i) \setminus Z_2|\}$. We use $\min\{3, d\}$ colors of either $col(e_i)$ or $col(e'_i)$ on correspondingly either f_i or f'_i . If we have applied 3 colors, we are done. Note that $|Z_1| \leq 12$ and $|Z_2| \leq 12$. We have $|Z_1 \cup Z_2| = |Z_1| + |Z_2| - |Z_1 \cap Z_2| \leq mult(e_i) + mult(e'_i) + 15 - |Z_1 \cap Z_2|$. Also, $|Z_1 \cap Z_2| \geq mult(e_i) + mult(e'_i) - 2d$. Therefore, $|Z_1 \cup Z_2| \leq 15 + 2d$. This means that, if $d < 3$, there are at least $3 - d$ colors of \mathcal{K} , none of which belongs to either Z_1 or Z_2 and then we use $3 - d$ such colors. □

Lemma 34 *Let p_H be any directed cycle of H^{dir} . We are able to extend the partial coloring of G_1 to the edges of tricky triangles covered by p_H and the edges of F_2 of their rescuers.*

Proof. Let t_1, \dots, t_k be the order of tricky triangles of C_1 , in which they (or more precisely, the vertices representing them) occur on p_H . Assume that each t_i has the form $t_i = (q_i, r_i, p_i)$, where (q_i, r_i) is a t-cycle of t_i . This means that for each $1 \leq i \leq k$ a rescuer t'_i of t_i has the form $t'_i = (q_i, r_i, p'_i)$, where for each $i > 1$, p'_i lies on t_{i-1} . The vertex p_i is incident to two edges $e_i = (s_i, p_i)$, $e'_i = (p_i, s'_i)$ belonging to C_{max} , which are already colored.

Before coloring G_1 , whenever possible, we replace the edges e_i and e'_i with one edge $e''_i = (s_i, s'_i)$. The only cases when we do not perform such a replacement is when (i) $s_i = s'_i$ and then (s_i, p_i) is a 2-cycle of C_{max} , (ii) (s_i, s'_i) is a 2-cycle of C_1 , (iii) there is a triangle of C_1 containing s_i and s'_i . Thus, apart from the three cases described above, edges e_i and e'_i are colored with the same 4 colors of \mathcal{K} .

Suppose that $f_i = (r_i, p'_i) \in F_2$. There are 6 colors forming set Z_i available for coloring it and we have to choose 3. There exists such set Z_i , because G_1 contains one edge of C_{max} incoming to p'_i , colored with 4 colors and one edge of C_1 incoming to p'_i , colored with 10 colors and f_i has to be diverse with both of them. Let $d_i = (s_i, x_i)$, $d'_i = (x'_i, s'_i)$ be two edges of C_1 . Each of them is colored with 10 colors. It is possible that there exists one or two edges of F_2 of the form $\tilde{f}_i = (s_i, y_i)$, $\tilde{f}'_i = (y'_i, s'_i)$, which are already colored. Any edge of F_2 is colored with 3 colors. If we want to recolor e_i , we have to ensure that e_i is diverse with both d_i and \tilde{f}_i . It means that we have at least 7 colors (set Z_i^1) at our disposal for coloring e_i . By the same token, we have at least 7 colors (set Z_i^2) available for coloring e'_i .

Let us first consider the case when e_i and e'_i are colored with the same 4 colors. If $|col(e_i) \cap Z_i| \leq 3$ we color f_i with 3 colors of $Z_i \setminus col(e_i)$. In the other case, we recolor e_i and e'_i by replacing one fixed color $k \in Z_i \cap col(e_i)$ with k_1 on e_i and with k_2 on e'_i . Colors k_1, k_2 are

such that $k_1 \neq k_2$ and $k_j \in Z_i^j \setminus \text{col}(e_i)$ for $j \in \{1, 2\}$. We then use k and two other colors of $Z_i \setminus \text{col}(e_i)$ on f_i .

Let us now deal with the three cases when e_i and e'_i are not replaced with one edge. If $s_i = s'_i$, then F_2 contains at most one of the edges $\tilde{f}_i = (s_i, y_i), \tilde{f}'_i = (y'_i, s_i)$. It means that at least one of the sets Z_i^1, Z_i^2 contains 10. We can notice that if some color k belongs to $Z_i^1 \cap Z_i^2$, then if we use k on exactly one of e_i, e'_i then that edge will be safe with respect to k , because neither d_i nor d'_i is colored with k . We need to assign $\text{mult}(e_i) + \text{mult}(e'_i) \leq 9$ different colors to e_i and e'_i . Note that color z assigned to f_i can be assigned only to e_i and not to e'_i but we need to ensure that it will not belong to a monochromatic cycle. We recolor e_i, e'_i as follows:

- If $|(Z_i^1 \cup Z_i^2) \setminus Z_i| \geq 6$, then we color e_i, e'_i using at most 3 colors of Z_i in total and assign the remaining colors (unassigned to either e_i or e'_i) to f_i .
- If $|(Z_i^1 \cup Z_i^2) \setminus Z_i| = 4 + x$, where $x \in \{0, 1\}$, then $|Z_i^1 \cap Z_i^2| \geq 7 - x$, because $|Z_i^1 \cup Z_i^2| = |Z_i^1| + |Z_i^2| - |Z_i^1 \cap Z_i^2| \geq 17 - |Z_i^1 \cap Z_i^2|$. This means that $Z_i^1 \cap Z_i^2$ contains at least $2 - x$ elements of Z_i . We assign $2 - x$ colors of $Z_i^1 \cap Z_i^2 \cap Z_i$ to e_i and also f_i , $1 + x$ other colors of Z_i to f_i and the remaining 3 colors of Z can be assigned to either e_i or e'_i . We also use $9 - (5 - x) = 4 + x$ colors of $(Z_i^1 \cup Z_i^2) \setminus Z_i$ to complete the coloring of e_i and e'_i .

□

References

- [1] Anna Adamaszek, Matthias Mnich, Katarzyna Paluch: New Approximation Algorithms for (1, 2)-TSP. ICALP 2018: 9:1-9:14
- [2] Noga Alon, Joel H. Spencer. The probabilistic method. 2000.
- [3] Chris Armen and Clifford Stein. Improved Length Bounds for the Shortest Superstring Problem. In *Proceedings of the 4th International Workshop on Algorithms and Data Structures, WADS '95*, volume 955 of *Lecture Notes in Computer Science*, pages 494–505. Springer 1995.
- [4] Chris Armen and Clifford Stein. A $2\frac{2}{3}$ -Approximation Algorithm for the Shortest Superstring Problem. In *Proceedings of the 7th Annual Symposium on Combinatorial Pattern Matching, CPM '96*, volume 1075 of *Lecture Notes in Computer Science*, pages 87–101. Springer 1996.
- [5] Markus Bläser. An $8/13$ -approximation algorithm for the asymmetric maximum TSP. *J. Algorithms*, 50(1):23–48, 2004.
- [6] Markus Bläser and Bodo Manthey. Two approximation algorithms for 3-cycle covers. In *Proceedings of the 5th International Workshop on Approximation Algorithms for Combinatorial Optimization*, volume 2462 of *Lecture Notes in Computer Science*, pages 40–50. Springer, 2002.
- [7] Avrim Blum and Tao Jiang and Ming Li and John Tromp and Mihalis Yannakakis. Linear Approximation of Shortest Superstrings. *J. ACM*, 41(4):630–647, 1994.

- [8] Dany Breslauer, Tao Jiang, and Zhigen Jiang. Rotations of periodic strings and short superstrings. *J. Algorithms*, 24(2):340–353, 1997.
- [9] Artur Czumaj and Leszek Gasieniec and Marek Piotrów and Wojciech Rytter. Sequential and Parallel Approximation of Shortest Superstrings. *J. Algorithms*, 23(1):74–100, 1997.
- [10] Szymon Dudycz, Jan Marcinkowski, Katarzyna E. Paluch, Bartosz Rybicki: A $4/5$ - Approximation Algorithm for the Maximum Traveling Salesman Problem. IPCO 2017: 173-185
- [11] M. L. Fisher, G. L. Nemhauser, and L. A. Wolsey. An analysis of approximations for finding a maximum weight Hamiltonian circuit. *Oper. Res.*, 27(4):799–809, 1979.
- [12] Robert W. Irving, Telikepalli Kavitha, Kurt Mehlhorn, Dimitrios Michail, Katarzyna E. Paluch: Rank-maximal matchings. *ACM Trans. Algorithms* 2(4): 602-610 (2006)
- [13] Haim Kaplan, Moshe Lewenstein, Nira Shafir, and Maxim Sviridenko. Approximation algorithms for asymmetric tsp by decomposing directed regular multigraphs. *J. ACM*, 52(4):602–626, 2005. Preliminary version appeared in FOCS’03.
- [14] S. Rao Kosaraju, James K. Park, and Clifford Stein. Long tours and short superstrings (preliminary version). In *Proceedings of the 35th Annual Symposium on Foundations of Computer Science*, pages 166–177, 1994.
- [15] Lukasz Kowalik and Marcin Mucha. Deterministic $7/8$ -approximation for the metric maximum tsp. *Theor. Comput. Sci.*, 410(47-49):5000–5009, 2009.
- [16] Lukasz Kowalik and Marcin Mucha. $35/44$ -approximation for asymmetric maximum tsp with triangle inequality. *Algorithmica*, 59(2):240–255, 2011.
- [17] Moshe Lewenstein and Maxim Sviridenko. A $5/8$ approximation algorithm for the maximum asymmetric tsp. *SIAM J. Discrete Math.*, 17(2):237–248, 2003.
- [18] M. Li. Towards a DNA sequencing theory. In *Proceedings of the 31st Annual Symposium on Foundations of Computer Science*, pages 125–134, 1990.
- [19] L.Lovasz, M. D.Plummer. Matching Theory. 1986.
- [20] Marcin Mucha. Lyndon words and short superstrings. *SODA 2013*.
- [21] A. Nakayama, B. Peroche. Linear arboricity of digraphs. *Networks*, 17:39-53, 1987.
- [22] Katarzyna E. Paluch: Better Approximation Algorithms for Maximum Asymmetric Traveling Salesman and Shortest Superstring. CoRR abs/1401.3670 (2014)
- [23] Katarzyna E. Paluch, Marcin Mucha, and Aleksander Madry. A $7/9$ - approximation algorithm for the maximum traveling salesman problem. In *Proceedings of the 12th International Workshop on Approximation Algorithms for Combinatorial Optimization*, volume 5687 of *Lecture Notes in Computer Science*, pages 298–311. Springer, 2009.
- [24] Katarzyna E. Paluch: Maximum ATSP with Weights Zero and One via Half-Edges. *Theory Comput. Syst.* 62(2): 319-336 (2018)

- [25] Katarzyna E. Paluch and Khaled M. Elbassioni and Anke van Zuylen. Simpler Approximation of the Maximum Asymmetric Traveling Salesman Problem. In *Proceedings of the 29th Symposium on Theoretical Aspects of Computer Science, STACS'2012*, Leibniz International Proceedings of Informatics 14, pages 501–506, 2012.
- [26] Christos H. Papadimitriou and Mihalis Yannakakis. The traveling salesman problem with distances one and two. *Mathematics of Operations Research*, 18:1–11, 1993.
- [27] Z. Sweedyk. A $2\frac{1}{2}$ -Approximation Algorithm for Shortest Superstring. *SIAM J. Comput.*, 29(3):954–986, 1999.
- [28] Jorma Tarhio and Esko Ukkonen. A greedy approximation algorithm for constructing shortest common superstrings. *Theor. Comput. Sci.*, 57:131–145, 1988.
- [29] Shang-Hua Teng and F. Frances Yao. Approximating Shortest Superstrings. *SIAM J. Comput.*, 26(2): 410–417, 1997.