FEEDBACK STABILIZATION OF LINEAR AND BILINEAR UNBOUNDED SYSTEMS IN BANACH SPACE

K. AMMARI, S. EL ALAOUI, AND M. OUZAHRA

ABSTRACT. We consider linear control systems of the form $\dot{y}(t) = Ay(t) - \mu BCy(t)$ where μ is a positive real parameter, A is the state operator and generates a linear C_0 -semigroup of contractions S(t) on a Banach space X, B and C are respectively the operators of control and observability, which are defined in appropriate spaces in which they are unbounded in some sense. We aim to show the exponential stability of the above system under sufficient conditions which are expressed in term of admissibility and observability properties. The uniform exponential stabilization using bilinear control is considered as well. Applications to transport and heat equations are also provided.

Contents

1.	Introduction	1
2.	The main results	3
2.1.	Preliminary on linear semigroups	3
2.2.	The stabilization results	5
3.	Applications	11
3.1.	Stabilization of unbounded bilinear systems	11
3.2.	Examples	12
References		15

1. Introduction

Let us consider the following linear control system

(1)
$$\dot{y}(t) = Ay(t) + Bu(t), \ t \ge 0, \ y(0) = y_0 \in X$$

augmented with the output z(t) = Cy(t), $t \ge 0$, where X and U are two Banach spaces representing respectively, the state and observation/control space, $A: D(A) \subset X \to X$ is the system operator, which generates a C_0 -semigroup of contractions S(t) on X, the space X is endowed with norm $\|\cdot\|_X$, and let X_{-1} denote the completion of X w.r.t to the norm $\|x\|_{-1} := \|(A-\eta I)^{-1}x\|_X$, $x \in X$ for some (or equivalently all) η in resolvent

 $^{2010\} Mathematics\ Subject\ Classification.\ 93 D15.$

Key words and phrases. Exponential stabilization; unbounded control operator; linear system; bilinear system.

set $\rho(A)$ of $A, B \in \mathcal{L}(U, X_{-1})$ is the control operator and $C \in \mathcal{L}(W, U)$ is the observation operator, where W is a Banach space such that the injections $X_1 \hookrightarrow W \hookrightarrow X$ are continuous $(X_1 \text{ being the space } D(A) \text{ equipped with the graph norm})$. Then, closing the system (1) with the control $u(t) = -\mu Cy(t)$, $(\mu > 0 \text{ is the gain control})$ one obtains the following Cauchy problem

(2)
$$\dot{y}(t) = Ay(t) - \mu BCy(t), \ t > 0, \ y(0) = y_0 \in X,$$

which is well-posed in X whenever $A-\mu BC$ is a generator of a C_0 -semigroup on X (cf. [20], Section II.6).

We further consider the bilinear system

(3)
$$\dot{y}(t) = Ay(t) + v(t)\mathcal{B}y(t), y(0) = y_0 \in X$$

The well-posedeness of the systems like (1) and (3) has been studied in many works using different approaches (see e.g. [1, 13, 14, 23, 27, 35, 38]). In several practical situation, the modeling gives rise to unbounded control systems of form (1) or (3), where the closed loop operator is of type Weiss-Staffans, Miyadera-Voigt or Desch-Schappacher (see e.g. [5, 21, 26, 32, 34, 35, 36, 37]). This fact often occurs when the control is exercised through the boundary or a point for systems governed by partial differential equations.

The problem of feedback stabilization of some classes of linear and nonlinear systems has been investigated in case of bounded and unbounded control operators in [2, 3, 4, 5, 6, 7, 8, 9]. Feedback stabilization of the bilinear system (3) has been investigated in the case of a bounded control operator by numerous authors using various control approaches, such as quadratic control laws, sliding mode control, piecewise constant feedback and optimal control laws (see [10, 31] and the references therein). Recently, the question of stabilization of bilinear systems with unbounded control operator has been treated in [12, 18, 19, 30]. In [12], the author considered the case where A is self-adjoint and B is positive self-adjoint and bounded from some subspace V of H to its dual space V', then he established the weak and strong stabilizability of the system (3) for all $y_0 \in D(A)$ using nonlinear control. Moreover, in [18, 19] it has been supposed that the linear operator B is relatively bounded w.r.t A from H to an extension X of Hwith a continuous embedding $H \hookrightarrow X$. Then, under an exact observability condition, it has been shown that (3) is strongly stabilizable, and a polynomial decay estimate of the stabilized state has been provided in the case of positive self-adjoint control operator. In [30], the exponential stabilizability of bilinear systems has been considered for Miyadera's control operator, and the stabilizing control is a switching one which leads to a closed-loop system like (2) evolving in a reflexive state space. More recently, the case of nonreflexive state space was considered in the context of bounded control operator [31]. In this paper, we deal with a wide class of linear/bilinear systems evolving on a nonreflexive state space with unbounded control operators, including control operators of type Weiss-Staffans, Miyadera-Voigt or Desch-Schappacher. Then we will give sufficient conditions for exponential

stabilizability of infinite dimensional systems that can be described by the systems (1) or (3).

The paper is organized as follows: In the second section, we provide some tools that will be required for the stabilization problem, then state and show the main result in which we present sufficient conditions for exponential stabilization of the linear system (1) with a feedback control involving the output, which leads to closed-loop operator of Weiss-Staffans's type. Next, we provide applications to bilinear system (3) with control operator of Miyadera-Voigt or Desch-Schappacher type. Applications to transport and heat equations are presented as well.

2. The main results

- 2.1. Preliminary on linear semigroups. Let us recall some notions and properties related to linear C_0 -semigroups.
 - The duality pairing between the space X and its dual X* is denoted by ⟨·,·⟩, where X* is the set of all bounded linear forms on X and the pairing between y ∈ X and φ ∈ X* is denoted by ⟨y, φ⟩. The duality map J from X to X* is in general a multi-valued operator; i.e. for each y ∈ X, J(y) is by definition the (nonempty) set of all φ ∈ X* such that ⟨y, φ⟩ = ||y||²_X = ||φ||², where ||·|| denotes the norm of X* associated to ||·||_X.
 - A one parameter family S(t), $t \geq 0$, of bounded linear operators from a Banach space X into X is a semigroup on X if (i) S(0) = I, (the identity operator on X) and (ii) S(t+s) = S(t)S(s) for every $t, s \geq 0$. A semigroup S(t) of bounded linear operators on X is a C_0 semigroup if in addition $\lim_{t\to 0^+} S(t)x = x$ for every $x \in X$. This property guarantees the continuity of the semigroup on \mathbb{R}^+ . Moreover, one can show (see [33], p. 4) that for every C_0 semigroup S(t), there exist constants $\omega \geq 0$ and $M \geq 1$ such that

$$(4) ||S(t)|| \le Me^{\omega t}, \ \forall t \ge 0.$$

If $\omega=0$ and $M=1,\,S(t)$ is called a C_0 -semigroup of contractions. The linear operator A defined by $Ax=\lim_{t\to 0^+}\frac{S(t)x-x}{t}$ for $x\in X$ such that $\lim_{t\to 0^+}\frac{S(t)x-x}{t}$ exists in X, is the infinitesimal generator of the C_0 -semigroup S(t). The linear space $D(A):=\{x\in X:\lim_{t\to 0^+}\frac{S(t)x-x}{t}\in X\}$ is the domain of A.

The infinitesimal generator of a contraction C_0 —semigroup is dissipative, i.e., for every $y \in D(A)$ and for all $y^* \in J(y)$ we have $Re\langle Ay, y^* \rangle \leq 0$ (see [33], pp. 14-15).

• For $x \in D(A)$, we have $Ax = \frac{d^+S(t)x}{dt}|_{t=0}$ and $y(t) := S(t)y_0$ is differentiable and lies in D(A) for all t > 0, and is the unique solution

of the Cauchy problem: $\dot{y}(t) = Ay(t), t > 0$, $y(0) = y_0$. Moreover, for every $y_0 \in X$; $y(t) = S(t)y_0$ is called mild solution of this Cauchy problem.

- If A is the infinitesimal generator of a C_0 -semigroup S(t), then D(A) (the domain of A) is dense in X and A is a closed linear operator. Moreover, according to Hille-Yosida's Theorem (see for instance [33], p. 20), a linear operator A is the infinitesimal generator of a C_0 -semigroup S(t) satisfying (4) if and only if (i) A is closed and $\mathcal{D}(A)$ is dense in X, and (ii) the resolvent set $\rho(A)$ of A contains the ray $(\omega, +\infty)$ and $||R(\lambda, A)^n|| \leq \frac{M}{(\lambda \omega)^n}$ for $\lambda > \omega$, n = 1, 2, ... In particular, a closed operator A with densely domain D(A) in X is the infinitesimal generator of a C_0 -semigroup of contractions on X if and only if the resolvent set $\rho(A)$ of A contains \mathbb{R}^+ and for all $\lambda > 0$; $||\lambda R(\lambda, A)|| \leq 1$ (see [33], p. 8).
- The C_0 -semigroup S(t) may be extended to a C_0 -semigroup $S_{-1}(t)$ on X_{-1} , whose generator is the extension $A_{-1}:D(A_{-1}):=X\subset X_{-1}\to X_{-1}$ of $A:D(A)\subset X\to X$ to a m-dissipative operator from X to X_{-1} . In particular, we have $A_{-1}y=Ay$ for all $y\in D(A)$ (see [20], p. 126). Using the integral representation of the resolvent one obtains $R(\lambda,A_{-1})y=R(\lambda,A)y,\ \forall y\in X,$ for all $\lambda\in\rho(A_{-1}).$ Recall also that $R(\lambda,A_{-1})y\in X,\ \forall y\in X_{-1}$ and $R(\lambda,A_{-1})y=R(\lambda,A)y,\ \forall y\in X,\ \forall \lambda\in\rho(A_{-1}).$ Moreover, If A is dissipative, then so is A_{-1} : For $z\in X$, we have $\|S_{-1}(t)z\|_{-1}=\|S(t)z\|_{-1}=\|R(\eta:A)S(t)z\|_X\leq \|R(\eta:A)z\|_X=\|z\|_{-1}.$ Then by density of X in X_{-1} , we conclude that $S_{-1}(t)$ is a contraction on X_{-1} .
- We have $\sup_{\lambda \in \rho(A_{-1})} \|\lambda R(\lambda, A_{-1})B\|_{\mathcal{L}(U, X_{-1})} < \infty$, where $\rho(A_{-1})$ is the resolvent set of A_{-1} . Moreover, by the closed graph theorem we have that $\lambda R(\lambda, A_{-1})B \in \mathcal{L}(U, X)$. For instance for $W = U = X_1$ and $Range(B) \subset X$, i.e., $B \in \mathcal{L}(X_1, X)$ (which is the case of Miyadera's operators), we have for all (real) $\lambda \in \rho(A_{-1})$ large enough,

$$\|\lambda R(\lambda, A_{-1})B\|_{\mathcal{L}(X_1, X)} = \|\lambda R(\lambda, A)B\|_{\mathcal{L}(X_1, X)} \le$$
$$\|B\|_{\mathcal{L}(X_1, X)}, \ \forall \lambda > 0.$$

In fact this is also true for any admissible operator B in the sense of (h_2) below (see [35], p. 219), that is there exists K > 0 such that

(5)
$$\|\lambda R(\lambda, A_{-1})B\|_{\mathcal{L}(U,X)} \leq K$$
, for all λ large enough.

In general this property does not imply the admissibility of B. (see [14, 15, 16, 17, 22, 25, 27, 28, 39] for some discussions and partial results about this implication).

• Let $\mathfrak{X} \oplus \mathfrak{X}_{-1}$ be a direct (algebraic) decomposition in X_{-1} between two subspaces \mathfrak{X} and \mathfrak{X}_{-1} such that $\mathfrak{X} \subset X$ and $X \cap \mathfrak{X}_{-1} = \{0\}$, and let $P_{\mathfrak{X}}$ denote the projection on \mathfrak{X} according to the above decomposition. Moreover, if K is a linear operator such that $Range(K) \subset \mathbb{R}$

 $\mathfrak{X} \oplus \mathfrak{X}_{-1}$, then we set $_{X}K := P_{\mathfrak{X}}K$.

Note that ${}_XK$ depends on the choice of \mathfrak{X} and \mathfrak{X}_{-1} , so in the sequel, we suppose that such a choice is made. In this case, we have $Range(K) = Y \oplus Z$ with $Y := \mathfrak{X} \cap Range(K) \subset X$ and $Z := \mathfrak{X}_{-1} \cap Range(K)$ so $Z \cap X = (0)$. Moreover, we can also write $K = K_1 + K_2$ with $Range(K_1) \subset X$ and $Range(K_2) \cap X = \{0\}$, and we have $K_1 = {}_XK$.

2.2. The stabilization results. In this part we consider the stability of the system (2). The first task is to guarantee the existence and uniqueness of the solution. Note that if the operator $\mu BC \in \mathcal{L}(W, X_{-1})$ is a Weiss-Staffans perturbation for A, then (see e.g. [1, 37]) the closed loop system (2) is well-posed. More precisely, for small gain control $\mu > 0$, the operator $(A - \mu BC)|_X$ (i.e. the part of $A_{-1} - \mu BC$ on X) with domain $D_{\mu} := \{y \in W : (A_{-1} - \mu BC)y \in X\}$ generates a C_0 -semigroup T(t) on X satisfying the following variation of constants formula (V.C.F)

(6)
$$T(t)y_0 = S(t)y_0 - \mu \int_0^t S_{-1}(t-s)BCT(s)y_0, \ \forall y_0 \in D_\mu.$$

Note that in general, we have $D(A) \cap D((BC)|_X) \subset D_\mu$. Moreover, if $W \subset X_1 \cup D((BC)|_X)$, then we have $(A_{-1} - \mu BC)|_X = D(A) \cap D((BC)|_X)$.

This motivates the consideration of the assumptions $(h_1) - (h_4)$ below.

 (h_1) the well-posedness assumption: there exists $\alpha_1 > 0$ such that for every $\mu \in (0, \alpha_1)$, the operator $(A_{-1} - \mu BC)|_X$ with domain $D := D((A_{-1} \cap BC)|_X)$ generates a C_0 -semigroup T(t) on X.

Now let us consider the following assumptions for some T, M > 0.

 (h_2) The admissibility assumption of $B \in \mathcal{L}(U, X_{-1})$:

$$\int_0^T S_{-1}(T-s)Bu(s)ds \in X, \, \forall \, u \in L^1(0,T;U),$$

which implies that

$$\left\| \int_{0}^{T} S_{-1}(T-s)Bu(s)ds \right\|_{X} \le M\|u\|_{1},$$

for all $u \in L^1(0,T;U)$ (or equivalently for all $u \in W^{1,1}(0,t_1;U)$). This also implies that the operator defined by

$$\mathcal{B}_T: u(\cdot) \in L^1(0, +\infty; U) \mapsto \int_0^T S_{-1}(T-s)Bu(s)ds$$

is bounded (see [1]).

Note that if X is reflexive, then the admissibility assumption (h_2) is equivalent to the boundedness of B (see [37]).

 (h_3) The admissibility assumption of $C \in \mathcal{L}(W, U)$:

$$\int_0^T \|CS(t)y\|_U dt \le M \|y\|_X, \ \forall y \in D(A).$$

 (h_4) Joint-admissibility of B and C:

$$\int_0^T \left\| C \int_0^r S_{-1}(r-s) Bu(s) ds \right\|_U dr \le M \|u\|_1, \ \forall u \in L^1(0,T;U)$$

with $||u||_1 = \int_0^T ||u(\tau)||_X d\tau$.

In the sequel, if there is no confusion, we use $\langle z, J(y) \rangle$ for any $y^* \in J(y)$ instead of $\langle z, y^* \rangle$. Also, for any functions $t \mapsto \zeta(t)$, we will write $\phi(\cdot) \in J(\zeta(\cdot))$ if $\phi(t) \in J(\zeta(t))$, $\forall t \geq 0$.

Now, for the stabilization results we further consider the following observation assumption.

 (h_5) The observability condition: for some $\delta > 0$ we have

(7)
$$\int_0^T \mathcal{R}e\langle_X(BC)S(t)y, J(S(t)y)\rangle dt \ge \delta \|S(T)y\|_X^2, \ \forall y \in D(A).$$

The estimate (7) may be seen as a null-exact controllability inequality in the sense of linear systems.

 (h_6) The function $F_J: y \mapsto \{\langle (BC)y, y^* \rangle; y^* \in J(y) \}$ is such that for all $y, z \in X$, there exists $(y^*, z^*) \in J(y) \times J(z)$ such that

$$|\langle_X(BC)y, y^*\rangle - \langle_X(BC)z, z^*\rangle| \le k_1 \left(||y||_{D(C)} + ||z||_{D(C)} \right) ||y - z||_X + k_2 (||y||_X + ||z||_X) ||C(y - z)||_U, \quad \forall y, z \in D(A) \subset W$$

for some constants $k_i \ge 0$, i = 1, 2, where $||y||_{D(C)} := ||y||_X + ||Cy||_U$.

This assumption is motivated by the fact that here, the state space is a general Banach space, i.e. without any smoothness property that evolves the duality mapping. In particular, if the state space X is smooth, so that J is Lipschitz-continuous and X is reflexive (see e.g. [30]), then (h_6) is verified under the admissibility of B, as in that case the operator BC will be bounded from W to X (see [37]).

Let us now state our main result.

Theorem 1. Let assumptions $(h_1)-(h_6)$ hold. Then there exists $\alpha > 0$ such that for any $\mu \in (0, \alpha)$, the closed-loop system (2) is exponentially stable.

Proof. According to assumption (h_1) , the operator $A_{BC} := (A_{-1} - \mu BC)|_X$ with domain $D(A_{BC}) = D(A) \cap D((BC)|_X) = D$ generates a C_0 -semigroup T(t) on X for $\mu > 0$ small enough (says $\mu \in (0, \alpha_1)$). Moreover, $y(t) := T(t)y_0$ is the unique mild solution of (2) and satisfies the following V.C.F

(8)
$$y(t) = S(t)y_0 - \mu \int_0^t S_{-1}(t-s)BCy(s)ds, \, \forall \, y_0 \in D.$$

Let $y_0 \in D$ be fixed. Then for all $t \geq 0$, we have $(A_{-1} - \mu BC)|_{X}y(t) \in X$. Moreover, y(t) has a weak derivative $A_{BC}y(t) = T(t)A_{BC}y_0$, which is weakly continuous and hence bounded

$$\left|\frac{d}{dt}\langle y(t), f\rangle\right| \le L\|A_{BC}y_0\|_X\|f\|, \ \forall f \in X^* \ (L > 0)$$

in any bounded time-interval. Thus y(t) (and so is $||y(t)||_X$) is Lipschitz continuous (recall that $||y||_X^2 = \sup_{f \in X^*, ||f|| \le 1} |\langle y, f \rangle|$). It follows that $||y(t)||_X$

is differentiable almost everywhere and (see [24]) for a.e t > 0 we have

(9)
$$\frac{d}{dt}||y(t)||_X^2 = 2\mathcal{R}e\langle A_{BC}y(t), J(y(t))\rangle.$$

Here J is the duality mapping of X (recall that $A_{BC}y(t) \in X$), which by integrating and using the dissipativeness of A implies

(10)
$$2\mu \int_{s}^{t} \mathcal{R}e\langle BCy(\tau), J(y(\tau))\rangle d\tau \leq \|y(s)\|_{X}^{2} - \|y(t)\|_{X}^{2}, \ t \geq s \geq 0.$$

According to (h_6) , we have

$$\mathcal{R}e\langle_{X}(BC)S(t)y_{0}, J(S(t)y_{0})\rangle \leq \\ K\left(\|S(t)y_{0}\|_{D(C)} + \|y(t)\|_{D(C)}\right)\|S(t)y_{0} - y(t)\|_{X} + \\ K\left(\|S(t)y_{0}\|_{X} + \|y(t)\|_{X}\right)\|C(S(t)y_{0} - y(t))\|_{U} + \\ \mathcal{R}e\langle BCy(t), J(y(t))\rangle$$

where $K = max(k_1, k_2)$, where we have used that $_X(BC)y(t) = BCy(t)$, as $y(t) \in X, \ \forall t \geq 0$.

From the admissibility assumption (h_3) , we have

$$\int_{0}^{T} \|CS(t)y_{0}\|_{U} dt \le M \|y_{0}\|_{X}$$

and for all $t \in [0, T]$, we have

$$||S(t)y_0 - y(t)||_X = \mu \left\| \int_0^t S_{-1}(t-s)BCy(s)ds \right\|_X$$

 $\leq \mu ||\mathcal{B}_T(Cy(\cdot))||_1,$

where \mathcal{B}_T is the bounded operator defined by

$$\mathcal{B}_T: u(\cdot) \in L^1(0, +\infty; U) \mapsto \int_0^T S_{-1}(T-s)Bu(s)ds.$$

Hence

$$||S(t)y_0 - y(t)||_X \le \mu M ||Cy(\cdot)||_1$$

where $||Cy(\cdot)||_1 := \int_0^T ||Cy(\tau)||_U d\tau$. It follows from this and (h_4) that

$$\int_0^T \|C(S(t)y_0 - y(t))\|_U dt = \mu \int_0^T \|C\int_0^t S_{-1}(t - s)BCy(s)ds\|_U dt$$

$$\leq \mu M \|Cy(\cdot)\|_1.$$

Let us estimate $||Cy(\cdot)||_1$. For every $t \geq 0$, we have $y(t) \in D \subset W$ and $S(t)y_0 \in D(A) \subset W$. Then, from the V.C.F, we derive

$$\begin{array}{ll} \int_0^T \|Cy(\tau)\|_U d\tau & \leq \int_0^T \|CS(\tau)y_0\|_U d\tau + \mu \int_0^T \|C\int_0^t S_{-1}(t-s)BCy(s)ds\|_U dt \\ & \leq M\|y_0\|_X + M\mu\|Cy(.)\|_1 \\ & = M\|y_0\|_X + M\mu \int_0^T \|Cy(\tau)\|_U d\tau. \end{array}$$

Hence for $0 < \mu < \alpha_2 := \inf(\alpha_1, \frac{1}{M})$, we have

(11)
$$\int_0^T \|Cy(\tau)\|_U d\tau \le \frac{M}{1 - M\mu} \|y_0\|_X.$$

Then

$$||y(t)||_X \le \left(1 + \frac{\mu M^2}{1 - M\mu}\right) ||y_0||_X.$$

We conclude that

$$\int_0^T \mathcal{R}e\langle_X(BC)S(t)y_0,J(S(t)y_0)\rangle dt \leq c\mu\|y_0\|_X^2 + \int_0^T \mathcal{R}e\langle BCy(t),J(y(t))\rangle dt$$

for some constant c > 0 which is independent of y_0 .

This together with (7) gives

$$\delta ||S(T)y_0||_X^2 - c\mu ||y_0||_X^2 \le \int_0^T \mathcal{R}e\langle BCy(t), J(y(t))\rangle dt.$$

Then taking y(t) instead of y_0 in the last estimate, it comes

(12)
$$\delta \|S(T)y(t)\|_X^2 - c\mu \|y(t)\|_X^2 \le \int_t^{t+T} \mathcal{R}e\langle BCy(s), J(y(s))\rangle ds.$$

From the variation of constants formula (8) we deduce that for all $t \geq T$, we have

$$||y(t)||_X \leq ||S(T)y_0||_X + \mu|| \int_0^t S_{-1}(t-s)BCy(s)ds||_X \leq ||S(T)y_0||_X + \mu M||Cy(\cdot)||_1.$$

Then taking y(kT) instead of y_0 , it comes via (11)

$$\begin{split} \|y(t)\|_{X} & \leq & \|S(T)y(kT)\|_{X} + \mu M \int_{kT}^{(k+1)T} \|Cy(s)\|_{U} ds \\ & \leq & \|S(T)y(kT)\|_{X} + \frac{\mu M^{2}}{1-M\mu} \|y(kT)\|_{X}, \ \forall t \in [kT, (k+1)T]. \end{split}$$

Thus for all $k \geq 0$, we have

(13)
$$||y((k+1)T)||_X^2 \le 2||S(T)y(kT)||_X^2 + 2\left(\frac{\mu M^2}{1-M\mu}\right)^2 ||y(kT)||_X^2.$$

This together with (10) and (12) implies

$$\mu\delta\bigg(\|y((k+1)T)\|_X^2 - 2\Big(\frac{\mu M^2}{1-M\mu}\Big)^2\|y(kT)\|_X^2\bigg) - 2c\mu^2\|y(kT)\|_X^2 \le \|y(kT)\|_X^2 - \|y((k+1)T)\|_X^2.$$

Hence

$$(1+\mu\delta)\|y((k+1)T)\|_X^2 \le \left(2\delta\mu(\frac{\mu M^2}{1-M\mu})^2 + 2c\mu^2 + 1\right)\|y(kT)\|_X^2, \ k \ge 0,$$

from which we derive

(14)
$$||y(kT)||_X^2 \le q^k ||y_0||_X^2, \ \forall k \ge 0$$

with $q:=\frac{1+2\mu^2\left(\delta\mu\left(\frac{M^2}{1-M\mu}\right)^2+c\right)}{1+2\mu\delta}$, which lies in (0,1) for $\mu\to 0^+$. Moreover, using the following well known property of linear C_0 –semigroups:

$$||y(t)||_X \le Ne^{wt}||y_0||_X, \ t \ge 0, \ (\text{for some constants } N, w > 0),$$

we deduce, by taking k = E(t/T), that $||y(t)||_X \leq N e^{wT} ||y(kT)||_X$ and hence by (14)

(15)
$$||y(t)||_X \le M' e^{-\sigma t} ||y_0||_X, \ \forall t \ge 0,$$

where M', σ are independent of y_0 .

This estimate extends to all initial data in X by density of D in X. \square

In the previous theorem, we have considered the case where the domain of the generator A_{BC} is $D(A) \cap D((BC)|_X)$. In the next result, we will state an other stabilization result which only requires that the domain of the generator A_{BC} is independent of the gain control μ , provided some further conditions are fulfilled. Let us consider the following assumption.

 (h_7) Compatibility condition: Range $(B_{\lambda}) \subset W$, for some/all $\lambda \in \rho(A)$ holds with $B_{\lambda} := \lambda R(\lambda, A_{-1})B$.

Notice that under the compatibility assumption, it comes from the closed graph theorem and the resolvent property that $Range(B_{\lambda}C) \subset \mathcal{W}$. Let us define the following scalar valued function

$$f_{BC}(y) = \limsup_{\lambda \to +\infty} \mathcal{R}e\langle B_{\lambda}Cy, J(y) \rangle.$$

Note that by (5), we have $f_{BC}(y) \in \mathbb{R}$, $\forall y \in X$. Moreover, if \overline{BC} is the operator defined by

$$\overline{BC}y:=\lim_{\lambda\to +\infty}B_{\lambda}Cy,\ \forall y\in D(\overline{BC}):=\{y\in W:\ \lim_{\lambda\to +\infty}B_{\lambda}Cy\ \text{exists in}\ X\},$$

then we have

(16)
$$f_{BC}(y) = \mathcal{R}e\langle \overline{BC}y, J(y)\rangle, \ \forall y \in D(\overline{BC}).$$

If in addition $Range(B) \subset X$, then the relation (16) holds in X. Let us consider the following assumptions:

 $(h_5)'$ The observability condition: for some $\delta > 0$ we have

(17)
$$\int_0^T f_{BC}(S(t)y)dt \ge \delta ||S(T)y||_X^2, \ \forall y \in D(A),$$

 $(h_6)'$ for all $y, z \in X$, there exists $(y^*, z^*) \in J(y) \times J(z)$ such that

$$|f_{BC}(y) - f_{BC}(z)| \le k_1 (||y||_{D(C)} + ||z||_{D(C)}) ||y - z||_X +$$

$$k_2(||y||_X + ||z||_X) ||C(y-z)||_U, \quad \forall y, z \in D(A) \subset W$$

for some constants $k_i \geq 0$, i = 1, 2.

From the proof of Theorem 1 we deduce the following result.

Corollary 1. Assume that for some $\mu_1 > 0$, the domain of $A_{BC} := (A_{-1} - \mu BC)|_X$ is independent of $\mu \in (0, \mu_1)$, and let assumptions $(h_2) - (h_4)$, $(h_5)' - (h_6)'$ and (h_7) hold.

Then there exists $\alpha > 0$ such that for any $\mu \in (0, \alpha)$, the closed-loop system (2) is exponentially stable.

Proof. Under the assumptions $(h_2) - (h_4)$ and the compatibility condition (h_7) , the operator $\mu BC \in \mathcal{L}(W, X_{-1})$ is a Weiss-Staffans perturbation for A. Accordingly (see [1]), the closed loop operator $(A_{-1} - \mu BC)|_X$ with domain $D(A_{BC}) = \{y \in W : (A_{-1} - \mu BC)y \in X\}$ generates a C_0 -semigroup $T(t)y_0$ on X and $y(t) := T(t)y_0$ is the unique mild solution of (1) and satisfies the formula (6) for all $y_0 \in D(A_{BC})$.

Let $y_0 \in D(A_{BC})$ be fixed. Then we have $(A_{-1} - \mu BC)|_{X}y(t) \in X$ for all $t \geq 0$, and $||y(t)||_{X}$ is differentiable almost everywhere and (9) holds for $y_0 \in D(A_{BC})$. Moreover, for every $y \in D(A_{BC})$, we have

$$\mathcal{R}e\langle\lambda R(\lambda,A_{-1})A_{BC}y,J(y)\rangle = \mathcal{R}e\langle\lambda R(\lambda,A_{-1})A_{-1}y,J(y)\rangle - \mu \mathcal{R}e\langle B_{\lambda}Cy,J(y)\rangle$$

Because A is dissipative, we have for all $y \in X$

$$\mathcal{R}e\langle R(\lambda, A_{-1})A_{-1}y, J(y) = \mathcal{R}e\langle -y + \lambda R(\lambda, A)y, J(y) \rangle$$

< 0.

It follows that

$$\mathcal{R}e\langle \lambda R(\lambda, A_{-1})A_{BC}y, J(y)\rangle \leq -\mu \mathcal{R}e\langle B_{\lambda}Cy, J(y)\rangle.$$

Since $y \in D(A_{BC})$, it comes that

$$\lambda R(\lambda, A_{-1})A_{BC}y = \lambda R(\lambda, A)A_{BC}y \to A_{BC}y \text{ in } X, \text{ as } \lambda \to +\infty.$$

Hence

$$\mathcal{R}e\langle A_{BC}y, J(y)\rangle \leq -\mu \lim_{\lambda \to +\infty} \sup \mathcal{R}e\langle B_{\lambda}Cy, J(y)\rangle.$$
$$= -\mu f_{BC}(y).$$

The remainder of the proof is exactly the same as in the proof of Theorem 1, which leads to the estimate (15) with constants M', σ which are independent of y_0 . Then we conclude by density of $D(A_{BC})$ in X.

Remark 1. From the proof of Theorem 1 (resp. Corollary 1), we can see that the results remain true if we assume that (h_5) (resp. $(h_5)'$) holds for some element of the duality set J(S(t)y) provided that (h_6) (resp. $(h_6)'$) holds for every $(y^*, z^*) \in J(y) \times J(z)$.

3. Applications

In this section we will apply the result of the previous section to the bilinear system (3). More precisely, we will investigate the exponential stability of (3) under the bang-bang feedback control $v(t) = -\mu \mathbf{1}_{\{t \geq 0; \mathcal{B}y(t) \neq 0\}}$. As special cases, we will consider Miyadera-Voigt's and Desch-Schappacher's control operators.

3.1. Stabilization of unbounded bilinear systems. First let us note that if \mathcal{B} is decomposable according to $\mathcal{B} = BC$ with B and C satisfying the conditions of Theorem 1, then we can see that (3) is exponentially stabilizable by the control $v(t) = -\mu \mathbf{1}_{\{t>0\}}$.

In the following corollary, we provide a result that extends the one of [30] to the case of non reflexive state space.

Corollary 2. Let $\mathcal{B} \in \mathcal{L}(X_1, X)$ be such that

 (m_1) there exists M > 0 such that

$$\int_0^T \|\mathcal{B}S(t)y\|_X dt \le M\|y\|_X, \ \forall y \in D(A),$$

 (m_2) there exists $\delta > 0$ such that

$$\int_{0}^{T} Re\langle \mathcal{B}S(t)y, J(S(t)y)\rangle dt \geq \delta ||y||_{X}^{2}, \ \forall y \in D(A),$$

 (m_3) the function $F_J: y \mapsto \{\langle \mathcal{B}y, y^* \rangle; y^* \in J(y)\}$ is such that for all $y, z \in X$, there exists $(y^*, z^*) \in J(y) \times J(z)$ such that

$$|\langle \mathcal{B}y, y^* \rangle - \langle \mathcal{B}z, z^* \rangle| \leq k_1 (||y||_{D(\mathcal{B})} + ||z||_{D(\mathcal{B})}) ||y - z||_X + k_2 (||y||_X + ||z||_X) ||y - z||_{D(\mathcal{B})}, \quad \forall y, z \in D(A),$$

for some constants $k_i \geq 0$, i = 1, 2.

Then there exists $\alpha > 0$ such that for any $\mu \in (0, \alpha)$, the control $v(t) = -\mu \mathbf{1}_{\{t \geq 0; \ \mathcal{B}y(t) \neq 0\}}$ guarantees the exponential stabilization of (3).

Proof. Let us first observe that under the assumption of the corollary, the operator $\mu\mathcal{B}$ may be seen as a Miyadera's perturbation of A, for $\mu > 0$ small enough, and we have $D((A_{-1} - \mu\mathcal{B})|_X) = D(A)$.

Moreover, we have $_{X}\mathcal{B} = \mathcal{B}$ relatively to the choice $\mathfrak{X} = X$ and $\mathfrak{X}_{-1} = \{0\}$.

Let us take $B = i : X \hookrightarrow X_{-1}$ (the embedding $X \hookrightarrow X_{-1}$) with U = X and $W = X_1$, so that $\mathcal{B} = C \in \mathcal{L}(X_1, X)$. Then, since $Range(\mathcal{B}) = Range(C) \subset X$, the compatibility condition (h_7) follows from the fact that $R(\lambda, A_{-1})C = R(\lambda, A)C$, while the others conditions of Theorem 1 are clearly satisfied. Moreover, the well-posedeness follows from [29, 36] (see

also [20], p. 199) and ([1], Theorem 18 and its remark).

Finally, by observing that $v(y)\mathcal{B}y = -\mu\mathcal{B}y$ with $v(y) = -\mu\mathbf{1}_{(y\notin\ker\mathcal{B})}$, we can see that the bilinear system (3), closed with the feedback control v(t) = $-\mu \mathbf{1}_{\{t>0: \mathcal{B}_{y}(t)\neq 0\}}$, leads to the system in closed-loop (2). Hence according to Theorem 1, we have the exponential stability for small gain control $\mu > 0$.

We have the following result regarding the case of Desch-Schapacher's control operator.

Corollary 3. Let $\mathcal{B} \in \mathcal{L}(X, X_{-1})$ be such that for some $\mu_1, T, \beta > 0$ we have

$$(ds)_1$$
 the domain of $A_{BC} := (A_{-1} - \mu BC)|_X$ is independent of $\mu \in (0, \mu_1)$, $(ds)_2$ for all $u \in L^1(0, T; X)$, we have $\int_0^T S_{-1}(T - s)\mathcal{B}u(s)ds \in X$,

$$(ds)_2$$
 for all $u \in L^1(0,T;X)$, we have $\int_0^1 S_{-1}(T-s)\mathcal{B}u(s)ds \in X$

 $(ds)_3$ there exists $\delta > 0$ such that

$$\int_0^T \limsup_{\lambda \to +\infty} Re \langle \mathcal{B}_{\lambda} S(t) y, J(S(t) y) \rangle dt \ge \delta ||S(T) y||_X^2, \ \forall y \in X,$$

 $(ds)_4$ for all $y, z \in X$, there exists $(y^*, z^*) \in J(y) \times J(z)$ such that

$$|\limsup_{\lambda \to +\infty} \langle \mathcal{B}_{\lambda} y, y^* \rangle - \limsup_{\lambda \to +\infty} \langle \mathcal{B}_{\lambda} z, z^* \rangle| \le k (\|y\|_X + \|z\|_X) \|y - z\|_X, \ \forall y, z \in X,$$

for some constant k > 0.

Then there is $\alpha > 0$ such that for any $\mu \in (0, \alpha)$, the control v(t) = $-\mu \mathbf{1}_{\{t>0: \mathcal{B}_{y}(t)\neq 0\}}$ guarantees the exponential stabilization of (3).

Proof. This follows from Corollary 1 by taking U = W := X, $C := I_X$ and $B := \mathcal{B} \in \mathcal{L}(X, X_{-1}).$

3.2. Examples.

Example 1. Let us consider the following system

(18)
$$\begin{cases} y_t(\cdot,t) = y_x(\cdot,t) - \mu \mathbf{B} y(\cdot,t), & in (0,1) \times (0,+\infty), \\ y(1,t) = 0, & in (0,+\infty), \\ y(\cdot,0) = y_0 \in L^1(0,1), & in (0,1). \end{cases}$$

Here, $X = L^1(0,1)$ and the duality map is given for all $y \in X$ by

$$J(y) = \{\xi \in L^{\infty}(\Omega): \ \xi(x) \in \ \operatorname{sign}(y(x)) \cdot \|y\|\},$$

where the sign function is defined by

$$sign(s) = \begin{cases} 1, & s > 0, \\ I, & s = 0, \text{ with } I = [-1, 1]. \\ -1, & s < 0. \end{cases}$$

The state operator is defined by Ay = y' with domain $D(A) = W_0^{1,1}(0,1) =$

 $\{y \in W^{1,1}(0,1): y(1)=0\}$ and generates the semigroup S(t) defined for all $y \in L^1(\Omega)$ by

$$(S(t)y)(\xi) = \begin{cases} y(\xi+t), & if \xi+t \leq 1\\ 0, & else. \end{cases}$$

Let $\phi = \alpha \delta_1 \in (W^{1,1}(0,1))'$ where δ_1 is the Dirac point evaluation in 1 and $\alpha \in \mathbb{R}$, and let us define the control operator for $y \in W^{1,1}(0,1)$ by $\mathbf{B}y = y + \phi(y)A_{-1}a$, where a(x) = 1, a.e. $x \in (0,1)$.

Let us consider the unbounded part of \mathcal{B} , which is defined by $y \mapsto \phi(y)A_{-1}a$, and which may be written in the form $BC: W^{1,1}(0,1) \to X_{-1}$, where $B \in \mathcal{L}(\mathbb{C}, X_{-1})$ is defined by $Bq = qA_{-1}a$, $q \in U := \mathbb{C}$ and $C \in \mathcal{L}(W, \mathbb{C})$ is defined by $Cy = \phi(y) = \alpha y(1)$, $\forall y \in W := W^{1,1}(0,1)$.

The admissibility properties of B and C as well as the compatibility condition can be checked for T=1 (see [1]), which implies the well-posedness of the above system (for $\mu > 0$ small enough). Let $\mu \in (0, \mu_1) \subset (0, 1)$ with $0 < \mu_1 < \frac{1}{\alpha}$ for which the well-posedness is guaranteed. Then

$$y \in D((BC)|_X) \Leftrightarrow \phi(y)A_{-1}a \in X$$

 $\Leftrightarrow \phi(y) = 0 \text{ (because } a \notin D(A))$
 $\Leftrightarrow \mathcal{B}y = y.$

Here, one can take $\mathfrak{X} = W^{1,1}(0,1)$ and $\mathfrak{X}_{-1} = span(A_{-1}a)$. Then we have ${}_{x}\mathcal{B}y = y, \ \forall y \in W^{1,1}(0,1)$ and so $\langle {}_{x}\mathcal{B}y, J(y) \rangle = \|y\|^2, \ \forall y \in W$. Thus the assumptions (h_5) & (h_6) clearly hold. Moreover, we have

$$y \in D((A_{-1} - \mu \mathcal{B})|_X) \Rightarrow A_{-1}(y - \mu \phi(y)a) \in X$$

$$\Rightarrow y - \mu \phi(y)a \in D(A)$$

$$\Rightarrow y(1) = \mu \phi(y)$$

$$\Rightarrow \mathcal{B}y = y \in X \text{ (recall that } 0 < \alpha \mu < 1)$$

$$\Rightarrow y \in D(\mathcal{B}|_X) \cap D(A).$$

We conclude by Theorem 1 that the system (18) is exponentially stable for $\mu > 0$ small enough.

Example 2. Let $\Omega = (0, +\infty)$ and let us consider the following system

(19)
$$\begin{cases} y_t(\cdot,t) = -y_x(\cdot,t) + v(t)(1+k(x))y(\cdot,t), & in (0,+\infty)^2 \\ y(0,t) = 0, & in (0,+\infty) \\ y(\cdot,0) = y_0 \in L^1(0,+\infty), & in (0,+\infty) \end{cases}$$

Here, $X = L^1(\Omega)$ is the state space, the parameter u(t) is the bilinear control and the corresponding solution $z(t) := y(\cdot, t) \in X$ is the state. The function k is such that $k \in L^1(0, +\infty)$ and $||k||_X < 1$.

The unbounded operator $A = -\frac{\partial}{\partial x}$ with domain

$$D(A) = \{ y \in W^{1,1}(\Omega); \ y(0) = 0 \}$$

generates a group of isometries S(t) on X, which is defined for all $y \in L^1(\Omega)$ by

$$S(t)y(\xi) = \begin{cases} y(\xi - t), & if \xi - t \ge 0 \\ 0, & else. \end{cases}$$

Let us define the operators $\mathcal{B} = y + k(x)y$ and By = k(x)y.

Note that if $k \notin L^{\infty}(0,+\infty)$, then \mathcal{B} is not a bounded operator on X.

Let us show that B is A-bounded. It comes from

$$|y(x)| = \left| \int_0^x y'(s)ds \right| \le \int_0^\infty |y'(s)|ds, \forall y \in D(A)$$

that $D(A) \subset D(B)$) and that for all $y \in D(A)$, we have

$$||ky||_X = \int_0^\infty |k(x)y(x)|dx \le ||y||_{D(A)} ||k||_X.$$

Hence $B \in \mathcal{L}(X_1, X)$.

• Admissibility of B. Let T > 0,

$$\int_{0}^{T} \|BS(t)y\|_{X} dt = \int_{0}^{T} \|k(x)S(t)y\|_{X} dt
= \int_{0}^{T} \int_{0}^{\infty} |k(x)y(x-t)| \mathbf{1}_{(0 \le t \le x)} dx dt
= \int_{0}^{T} \int_{0}^{\infty} |k(x+t)y(x)| dx dt
= \int_{0}^{\infty} |y(x)| \left(\int_{0}^{T} |k(x+t)| dt\right) dx
\le \|k\|_{X} \|y\|_{X}.$$

This implies the admissibility etimate (m_1) .

• Assumption (h_6) . For T > 0, we have

$$\langle k(x)y, J(y)\rangle = ||y||_X \int_0^\infty k(x)|y(x)|dx, \ y \in D(A).$$

Then

$$|\langle k(x)y, J(y) \rangle - \langle k(x)z, J(z) \rangle| \leq ||y||_X \int_0^\infty |k(x)(y(x) - z(x))| dx + ||y - z||_X \int_0^\infty |k(x)y(x)| dx = ||y||_X ||B(y - z)||_X + ||y - z||_X ||By||_X$$

which gives (h_6) .

• Observation. For T > 0, we have

$$\langle \mathcal{B}S(t)y, J(S(t)y) \rangle = \|S(t)y\|_X^2 + \langle BS(t)y, J(S(t)y) \rangle$$

$$\geq \|y\|_X^2 - \|y\|_X \|BS(t)y\|_X$$

then

$$\int_{0}^{T} \langle \mathcal{B}S(t)y, J(S(t)y) \rangle dt \geq \|y\|_{X}^{2} - \|y\|_{X} \int_{0}^{T} \|BS(t)y\|_{X} dt \\
\geq (1 - \|k\|_{X}) \|y\|_{X}^{2}.$$

Hence (m_2) holds for $||k||_X < 1$.

From Corollary 2, we conclude that the control $v(t) = -\mu \mathbf{1}_{\{t \geq 0: \ y(\cdot,t) \neq 0\}}$ ensures the exponential stability of the system (19).

Example 3. Consider the following system:

$$\begin{cases} y_t(\cdot,t) = y_{xx}(\cdot,t) + v(t) (y(\cdot,t) + y_x(\cdot,t)), & (x,t) \in [0,1] \times (0,+\infty), \\ y'(0,t) = y'(1,t) = 0, & t \in (0,+\infty), \\ y(\cdot,0) = y_0 \in \mathcal{C}_0([0,1]), & x \in [0,1] \end{cases}$$

where $u(t) \in \mathbb{R}$ is the control and $y(t) = y(\cdot, t)$ is the state.

The state space $X = \mathcal{C}_0([0,1])$ is equipped with the supremum norm, the operator $A = \partial_{xx}$ with domain $\mathcal{D}(A) = \{y \in \mathcal{C}^2([0,1]) : y'(0) = y'(1) = 0\}$ generates a contraction C_0 -semigroup S(t) in $X := \mathcal{C}_0([0,1])$ (see [20], pp. 93-94). The control operator is $\mathcal{B} = I + \partial_x$, then the A-boundedness of \mathcal{B} follows from the following inequalities

$$|y'(x)| \le \int_0^1 |y''(s)| ds \le ||y||_{D(A)}, \ y \in D(A).$$

In other words, $\mathcal{B} \in \mathcal{L}(X_1, X)$.

We will show that the stabilization assumptions previously considered are not all required. In particular, here we only need some elements of the duality set. For $f \in C_0([0,1])$ we have ([20], p. 93):

$$\Lambda(f) := \{ \varphi = f(s_0)\delta_0 : \ s_0 \in [0,1] \ is \ s.t \ |f(s_0)| = ||f|| = \max_{s \in [0,1]} |f(s)| \} \subset J(f),$$

where δ_{s_0} is any point measure supported by a point s_0 where |f| reaches its maximum. For the expression of the full duality map (see e.g. [11], p. 5).

Let $y \in D(A)$ and let $y^* = |y(s_0)|\delta_{s_0} \in J(y)$, i.e. $|y(s_0)| = ||y||$. Thus for $s_0 \in (0,1)$ we have $y'(s_0) = 0$, so taking into account the Neumann boundary conditions, we deduce that $y'(s_0) = 0$ for every $s_0 \in [0,1]$ such that $|y(s_0)| = ||y||$. Thus we have

$$\langle y', J(y) \rangle = |y'(s_0)||y(s_0)| = 0, \ \forall y \in D(A).$$

It follows that

$$\langle \mathcal{B}y, J(y) \rangle = ||y||^2, \ \forall y \in D(A).$$

Hence (h_5) and (h_6) are verified for any element of $\Lambda(f)$.

Now, from ([33], p. 82) we deduce that $A - \mu \mathcal{B}$ with domain $D(A - \mu \mathcal{B}) = D(A)$ is a generator on X for $\mu > 0$ small enough. Hence using again that $\langle \mathcal{B}y, J(y) \rangle = ||y||^2$ for all $y \in D(A)$, we derive directly from (9) that the feedback control $v(t) = -\mu \mathbf{1}_{\{t \geq 0: \ y(t) + y_x(t) \neq 0\}}$ results in an exponentially stable closed-loop system for a small gain control $\mu > 0$.

REFERENCES

- M. Adler, M. Bombieri and K. J. Engel, On perturbations of generators of C₀-Semigroups, Abstr. Appl. Anal., 2014, Art. ID 213020, 13 pp.
- [2] F. Alabau-Boussouira and K. Ammari, Sharp energy estimates for nonlinearly locally damped PDEs via observability for the associated undamped system, J. Funct. Anal., 260 (2011), 2424–2450.

- [3] K. Ammari, A. Bchatnia and K. El Mufti, Non-uniform decay of the energy of some dissipative evolution systems, Z. Anal. Anwend., 36 (2017), 239–251.
- [4] K. Ammari, A. Bchatnia and K. El Mufti, Stabilization of the nonlinear damped wave equation via linear weak observability, *NoDEA Nonlinear Differential Equations Appl.*, **23** (2016), no. 2, Art. 6, 18 pp.
- [5] K. Ammari and S. Nicaise, Stabilization of elastic systems by collocated feedback, 2124, Springer, Cham, 2015.
- [6] K. Ammari and M. Ouzahra, Feedback stabilization for a bilinear control system under weak observability inequalities, Automatica J. IFAC., 113 (2020), 108821, 8 pp.
- [7] K. Ammari and F. Triki, On Weak Observability for Evolution Systems with Skew-Adjoint Generators, SIAM J. Math. Anal., 52 (2020), 1884–1902.
- [8] K. Ammari and M. Tucsnak, Stabilization of second order evolution equations by a class of unbounded feedbacks, *ESAIM Control Optim. Calc. Var.*, **6** (2001), 361–386.
- [9] K. Ammari and M. Tucsnak, Stabilization of Bernoulli-Euler beams by means of a pointwise feedback force, SIAM J. Control Optim., 39 (2000), 1160–1181.
- [10] J. M. Ball and M. Slemrod, Feedback stabilization of distributed semilinear control systems, Appl. Math. Opt., 5 (1979), 169–179.
- [11] V. Barbu and P. Korman, Analysis and control of nonlinear infinite dimensional systems, Vol. 190, Boston: Academic Press, 1993.
- [12] L. Berrahmoune, Stabilization of unbounded bilinear control systems in Hilbert space, Journal of Mathematical Analysis and Applications, 372 (2010), 645–655.
- [13] L. Berrahmoune, A note on admissibility for unbounded bilinear control systems, Bulletin of the Belgian Mathematical Society-Simon Stevin, 16 (2009), 193–204.
- [14] H. Bounit and A. Idrissi, Regular bilinear systems, IMA Journal of Mathematical Control and Information, 22 (2005), 26–57.
- [15] H. Bounit, A. Driouich and O. El-Mennaoui, A direct approach to the Weiss conjecture for bounded analytic semigroups, *Czechoslovak mathematical journal*, 60 (2010), 527–539.
- [16] H. Bounit, A. Driouich and O. El-Mennaoui, A direct approach to the weighted admissibility of observation operators for bounded analytic semigroups, *Semigroup Forum*, 83 (2011), 52–64.
- [17] H. Bounit and F. Maragh, On the regularity for some class of L^p-well-posed linear systems, Transnational Journal of Mathematical Analysis and Applications, 2 (2014), 37–65.
- [18] R. El Ayadi, M. Ouzahra and A. Boutoulout, Strong stabilization and decay estimate for unbounded bilinear systems, *International Journal of Control*, 85 (2012), 1497– 1505.
- [19] R. El Ayadi and M. Ouzahra, Feedback stabilization for unbounded bilinear systems using bounded control, IMA Journal of Mathematical Control and Information, 36 (2019), 1073–1087.
- [20] K. J. Engel and R. Nagel, One-parameter semigroups for linear evolution equations, With contributions by S. Brendle, M. Campiti, T. Hahn, G. Metafune, G. Nickel, D. Pallara, C. Perazzoli, A. Rhandi, S. Romanelli and R. Schnaubelt. Graduate Texts in Mathematics, 194. Springer-Verlag, New York, 2000.
- [21] G. Greiner, Perturbing the boundary-conditions of a generator, Houston Journal of mathematics, 13 (1987), 213–229.
- [22] B. H. Haak and P.C. Kunstmann, Weighted admissibility and wellposedness of linear systems in Banach spaces. SIAM journal on control and optimization, **45** (2007), 2094–2118.
- [23] S. Hadd, R. Manzo and A. Rhandi, Unbounded perturbations of the generator domain, Discrete Contin. Dyn. Syst, 35 (2015), 703–723.

- [24] T. Kato, Nonlinear semigroups and evolution equations, J. Math. Soc. Japan., 19 (1967), 508–520.
- [25] B. Jacob, H. Zwart and O. Staffans, Weak admissibility does not imply admissibility for analytic semigroups, *Systems & Control Letters*, **48** (2003), 341–350.
- [26] Z. H. Luo, B. Z. Guo and O. Morgul, Stability and Stabilisation of Infinitedimensional Systems with Applications, Communications and Control Engineering Series. Springer-Verlag London, Ltd., London, 1999.
- [27] F. Maragh, H. Bounit, A. Fadili and H. Hammouri, On the admissible control operators for linear and bilinear systems and the Favard spaces, Bulletin of the Belgian Mathematical Society-Simon Stevin, 21 (2014), 711–732.
- [28] C. Le Merdy. The weiss conjecture for bounded analytic semigroups, Journal of the London Mathematical Society, 67 (2003), 715–738.
- [29] I. Miyadera, On perturbation theory for semi-groups of operators. Tohoku Mathematical Journal, Second Series, 18 (1966), 299–310.
- [30] M. Ouzahra, Exponential stability of nondissipative linear system in Banach space and application to unbounded bilinear systems, Systems & Control Letters, 109 (2017), 53–62.
- [31] M. Ouzahra, Uniform exponential stabilization of nonlinear systems in Banach spaces, Mathematical Methods in the Applied Sciences, https://doi.org/10.1002/mma.6468 (2020).
- [32] A. Ö. Özer, Stabilization results for well-posed potential formulations of a current-controlled piezoelectric beam and their approximations, *Applied Mathematics & Optimization*, https://doi.org/10.1007/s00245-020-09665-4 (2020).
- [33] A. Pazy, Semi-groups of linear operators and applications to partial differential equations, Springer Verlag, New York, 1983.
- [34] R. Risken, The Fokker-Planck equation: Methods of solutions and applications, Springer Verlag, Berline, 1996.
- [35] O. Staffans, Well-posed linear systems, Vol. 103. Cambridge University Press, 2005.
- [36] J. Voigt, On the perturbation theory for strongly continuous semigroups, Mathematische Annalen, 229 (1977), 163–171.
- [37] G. Weiss, Admissibility of unbounded control operators, SIAM Journal on Control and Optimization, 27 (1977), 527–545.
- [38] G. Weiss, Regular linear systems with feedback, Math. Control Signals Syst., 7 (1994), 23–57.
- [39] G. Weiss, Two conjectures on the admissibility of control operators. Birkhäuser Verlag, Basel, In F. Kappel, W. Desch, editors, Estimation and Control of Distributed Parameter Systems: 367–378, 1989.

UR Analysis and Control of PDE's, UR 13E64, Department of Mathematics, Faculty of Sciences of Monastir, University of Monastir, 5019 Monastir, Tunisia

E-mail address: kais.ammari@fsm.rnu.tn

DEPARTMENT OF MATHEMATICS & INFORMATICS, ENS. UNIVESITY OF SIDI MOHAMED BEN ABDELLAH. FES, MOROCCO

E-mail address: elalaoui.sfe18@gmail.com

DEPARTMENT OF MATHEMATICS & INFORMATICS, ENS. UNIVESITY OF SIDI MOHAMED BEN ABDELLAH. FES, MOROCCO

 $E ext{-}mail\ address: mohamed.ouzahra@usmba.ac.ma}$