arXiv:2006.00760v2 [nlin.AO] 11 Feb 2023

Quantum asymptotic phases reveal signatures of quantum synchronization

Yuzuru Kato^{1,*} and Hiroya Nakao²

¹Department of Complex and Intelligent Systems, Future University Hakodate, Hokkaido 041-8655, Japan ²Department of Systems and Control Engineering, Tokyo Institute of Technology, Tokyo 152-8552, Japan (Dated: February 14, 2023)

Synchronization of quantum nonlinear oscillators has attracted much attention recently. To characterize the quantum oscillatory dynamics, we recently proposed a fully quantum-mechanical definition of the asymptotic phase, which is a key quantity in the synchronization analysis of classical nonlinear oscillators [1]. In this work, we further extend this theory and introduce multiple asymptotic phases using the eigenoperators of the adjoint Liouville superoperator of the quantum nonlinear oscillator associated with different fundamental frequencies. We analyze a quantum van der Pol oscillator with Kerr effect in the strong quantum regime and show that the system has several different fundamental frequencies. By introducing order parameters and power spectra in terms of the associated quantum asymptotic phases, we reveal that phase locking of the system with a harmonic drive at several different frequencies, an explicit quantum signature observed only in the strong quantum regime, can be interpreted as synchronization on a torus rather than a simple limit cycle.

^{*} Corresponding author: katoyuzu@fun.ac.jp

I. INTRODUCTION

Spontaneous rhythmic oscillations and synchronization are phenomena ubiquitously observed in various fields of science and technology [2–7]. Owing to the recent progress in nanotechnology, synchronization in micro- and nano-scale devices has been realized experimentally [8–12] and experimental demonstrations of quantum phase synchronization in spin-1 atoms [13], in nuclear spin systems [14], and on the IBM Q system [15] have been reported. A number of theoretical investigations have also been performed to reveal novel quantum signatures in synchronization [1, 16–46, 62].

In the strong quantum regime where only a small number of energy states participate in the system dynamics, the discrete nature of the energy spectrum can give rise to explicit quantum signatures, such as multiple phase locking at several different frequencies [16] and synchronization blockade [17, 18]. In characterizing quantum synchronization, several definitions of the system's oscillation phase have been proposed [16, 19–23, 43–46].

In classical mechanics, nonlinear dissipative systems exhibiting spontaneous rhythmic dynamics can be modeled as limit-cycle oscillators. The asymptotic phase [2–6] is a fundamental quantity for the analysis of synchronization, which is defined by the oscillator's vector field and increasing with a constant frequency in the basin of the limit-cycle attractor. It provides the basis for *phase reduction theory* [2–7], a standard dimensionality-reduction method to derive phase equations describing weakly-coupled oscillators.

We recently introduced the asymptotic phase for quantum oscillatory systems [1] by extending the definition of the asymptotic phase for classical stochastic oscillatory systems [47] based on the Koopman operator theory [48]. This quantum asymptotic phase is defined fully quantum-mechanically in terms of the eigenoperator of the system's adjoint Liouville superoperator associated with the slowest decaying mode and hence applicable even in the strong quantum regimes where we cannot rely on the limit-cycle trajectory in the classical limit as in the semiclassical regime [27, 28, 49].

In this study, we further extend this definition and introduce multiple quantum asymptotic phases in terms of the eigenoperators with several different fundamental frequencies, i.e., with the eigenvalues possessing the smallest absolute imaginary part in the individual branches of the eigenvalues near the imaginary axis, which play dominant roles in synchronization dynamics. As an example, we analyze a quantum van der Pol oscillator with Kerr effect and show that it possesses several dominant eigenvalues with different fundamental frequencies in the strong quantum regime. By introducing the order parameters and power spectra in terms of the associated quantum asymptotic phases with respective fundamental frequencies, we reveal a torus-like structure of multiple-frequency phase locking of the system with a harmonic drive [16], which is observed only in the strong quantum regime.

II. ASYMPTOTIC PHASES FOR QUANTUM OSCILLATORY SYSTEMS

In Ref. [1], we proposed a definition of the asymptotic phase for quantum oscillatory systems in terms of the eigenoperator of the adjoint Liouville superoperator associated with the slowest decaying mode, inspired by the definition of the asymptotic phase for classical stochastic oscillators [47, 48]. In this section, we briefly review this definition and extend it to introduce multiple quantum asymptotic phases by using the eigenoperators associated with several eigenvalues with different fundamental frequencies.

A. Quantum master equation

We consider a quantum nonlinear oscillatory system with a single degree of freedom coupled to reservoirs, and assume that the interactions of the system with the reservoirs are instantaneous and Markovian approximation can be employed. The time evolution of the system's density operator ρ is described by a quantum master equation [50, 51],

$$\dot{\rho} = \mathcal{L}\rho = -i[H,\rho] + \sum_{j=1}^{n} \mathcal{D}[C_j]\rho, \qquad (1)$$

where \mathcal{L} is a Liouville superoperator representing the evolution of ρ , H is a system Hamiltonian, C_j is a coupling operator between the system and *j*th reservoir (j = 1, ..., n), [A, B] = AB - BA is the commutator, $\mathcal{D}[C]\rho = C\rho C^{\dagger} - (\rho C^{\dagger}C + C^{\dagger}C\rho)/2$ is the Lindblad form (\dagger denotes Hermitian conjugate), and the reduced Planck's constant is set as $\hbar = 1$.

Introducing an inner product $\langle X, Y \rangle_{tr} = \text{Tr}(X^{\dagger}Y)$ of linear operators X and Y, we define the adjoint superoperator

 \mathcal{L}^* of \mathcal{L} satisfying $\langle \mathcal{L}^* X, Y \rangle_{tr} = \langle X, \mathcal{L}Y \rangle_{tr}$

$$\mathcal{L}^* X = i[H, X] + \sum_{j=1}^n \mathcal{D}^*[C_j] X,$$
(2)

where $\mathcal{D}^*[C]X = C^{\dagger}XC - (XC^{\dagger}C + C^{\dagger}CX)/2$ is the adjoint Lindblad form. This \mathcal{L}^* describes the evolution of an observable F as

$$\dot{F} = \mathcal{L}^* F. \tag{3}$$

In the Schrödinger picture, the density operator ρ evolves as in Eq. (1) while the observation operator F does not vary with time, and in the Heisenberg picture, F evolves as in Eq. (3) while ρ remains constant. The expectation value

$$\langle F \rangle = \operatorname{Tr}(\rho F) = \langle \rho, F \rangle_{tr}$$
 (4)

of F with respect to ρ is kept the same in both pictures (note that ρ is self-adjoint).

We assume that the Liouville superoperator \mathcal{L} has a set of eigensystem (an eigenvalue and right and left eigenoperators) { λ_k, U_k, V_k } satisfying

$$\mathcal{L}U_k = \lambda_k U_k, \quad \mathcal{L}^* V_k = \overline{\lambda_k} V_k, \quad \langle V_k, U_l \rangle_{tr} = \delta_{kl}, \tag{5}$$

for $k, l = 0, 1, 2, \ldots$, where the overline indicates complex conjugate [52]. We assume that among $\{\lambda_k\}_{k\geq 0}$, one eigenvalue λ_0 is always 0, which corresponds to the stationary state ρ_0 of the system satisfying $\mathcal{L}\rho_0 = 0$, and all other eigenvalues have negative real parts. This assumption also means that the system has no decoherence-free subspace [53]. Considering the system's oscillatory dynamics, we also assume that the eigenvalues of \mathcal{L} with the largest nonvanishing real part (i.e., with the slowest decay rate) are given by a complex-conjugate pair and denote them as Λ_1 and $\overline{\Lambda_1}$, where $\Omega_1 = \text{Im } \overline{\Lambda_1}$ gives the frequency of the slowest decay mode. One may also choose $\Omega_1 = \text{Im } \Lambda_1$, which reverses the direction of the asymptotic phase. The sign of Ω_1 can be chosen arbitrarily and will be fixed later.

Figure 1 shows typical examples of the eigenvalues of \mathcal{L} of the quantum vdP oscillator (see the next section for details). The eigenvalues form several branches. In the semiclassical regime (Fig. 1(b)), the rightmost branch is far apart from the other branches and thus it is the only dominant branch, while in the strong quantum regime (Fig. 1(a)), the branches are closer to each other and the system possess several comparably important branches. It can also be seen in the semiclassical regime (Fig. 1(b)) that the imaginary part of the eigenvalues are approximately integer multiples of the fundamental frequency (the smallest absolute imaginary part with the slowest decay rate) in each individual branch.

B. Phase-space representation

The density operator ρ can also be transformed into a quasiprobability distribution in the phase space [50, 51, 54]. We use the *P*-representation and describe ρ as

$$\rho = \int p(\boldsymbol{\alpha}) |\alpha\rangle \langle \alpha | d\boldsymbol{\alpha}, \tag{6}$$

where $|\alpha\rangle$ is a coherent state specified by a complex value α , or equivalently by a complex vector $\boldsymbol{\alpha} = (\alpha, \overline{\alpha})^T$, $p(\boldsymbol{\alpha})$ is a quasiprobability distribution of $\boldsymbol{\alpha}$, $d\boldsymbol{\alpha} = d\alpha d\overline{\alpha}$, and the integral is taken over the entire complex plane. The observable F is also transformed into a function in the phase space as

$$f(\boldsymbol{\alpha}) = \langle \boldsymbol{\alpha} | F | \boldsymbol{\alpha} \rangle,\tag{7}$$

where the operator F is arranged in the normal order [50, 51, 54]. By introducing the L^2 inner product $\langle g(\boldsymbol{\alpha}), h(\boldsymbol{\alpha}) \rangle_{\boldsymbol{\alpha}} = \int \overline{g(\boldsymbol{\alpha})} h(\boldsymbol{\alpha}) d\boldsymbol{\alpha}$ of two functions $g(\boldsymbol{\alpha})$ and $h(\boldsymbol{\alpha})$, the expectation value of F with respect to ρ is expressed as

$$\langle F \rangle = \operatorname{Tr}(\rho F) = \int d\boldsymbol{\alpha} p(\boldsymbol{\alpha}) f(\boldsymbol{\alpha}) = \langle p(\boldsymbol{\alpha}), f(\boldsymbol{\alpha}) \rangle_{\boldsymbol{\alpha}}.$$
 (8)

The time evolution of $p(\alpha)$ corresponding to Eq. (1) is described by a partial differential equation

$$\partial_t p(\boldsymbol{\alpha}) = L_{\boldsymbol{\alpha}} p(\boldsymbol{\alpha}),\tag{9}$$

where the differential operator L_{α} satisfies $\mathcal{L}\rho = \int L_{\alpha} p(\alpha) |\alpha\rangle \langle \alpha | d\alpha$. The explicit form of L_{α} can be calculated from Eq. (1) by using the standard calculus for the phase-space representation [50, 51, 54]. The corresponding evolution of $f(\alpha)$ in the Heisenberg picture is given by

$$\partial_t f(\boldsymbol{\alpha}) = L_{\boldsymbol{\alpha}}^+ f(\boldsymbol{\alpha}),\tag{10}$$

where the differential operator L^+_{α} is the adjoint of L_{α} with respect to the L^2 inner product, i.e., $\langle L^+_{\alpha}g(\alpha), h(\alpha)\rangle_{\alpha} = \langle g(\alpha), L_{\alpha}h(\alpha)\rangle_{\alpha}$, which satisfies $L^+_{\alpha}f(\alpha) = \langle \alpha | \mathcal{L}^*F | \alpha \rangle$.

The differential operator L_{α} also has a set of eigensystem (an eigenvalue and right and left eigenfunctions) $\{\lambda_k, u_k(\alpha), v_k(\alpha)\}$ which satisfies

$$L_{\alpha}u_k = \lambda_k u_k, \quad L_{\alpha}^+ v_k = \overline{\lambda_k} v_k, \quad \langle v_k, u_l \rangle_{\alpha} = \delta_{kl}.$$
⁽¹¹⁾

This eigensystem has a one-to-one correspondence with Eq. (5), where the eigenvalues $\{\lambda_k\}_{k\geq 0}$ are the same as those of \mathcal{L} ; the eigenfunctions u_k and v_k of L_{α} and L_{α}^+ are related to the eigenoperators U_k and V_k of \mathcal{L} and \mathcal{L}^* as

$$U_k = \int u_k(\boldsymbol{\alpha}) |\alpha\rangle \langle \alpha | d\boldsymbol{\alpha}, \quad v_k(\boldsymbol{\alpha}) = \langle \alpha | V_k | \alpha\rangle, \tag{12}$$

which follow from $\mathcal{L}U_k = \int u_k(\boldsymbol{\alpha}) \{\mathcal{L}|\alpha\rangle\langle\alpha|\} d\boldsymbol{\alpha} = \int \{L_{\boldsymbol{\alpha}}u_k(\boldsymbol{\alpha})\} |\alpha\rangle\langle\alpha|d\boldsymbol{\alpha} = \lambda_k U_k \text{ and } L_{\boldsymbol{\alpha}}^+ v_k = L_{\boldsymbol{\alpha}}^+ \langle\alpha|V_k|\alpha\rangle = \langle\alpha|\mathcal{L}^*V_k|\alpha\rangle = \overline{\lambda_k}\langle\alpha|V_k|\alpha\rangle = \overline{\lambda_k}v_k.$

C. Quantum asymptotic phase associated with the slowest decaying mode

In Ref. [1], we defined the quantum asymptotic phase function Φ_1 of the system state ρ as the argument (polar angle) of the expectation of the eigenoperator V_1 associated with the eigenvalue Λ_1 satisfying $\mathcal{L}^*V_1 = \overline{\Lambda_1}V_1$, namely,

$$\Phi_1(\rho) = \arg \langle V_1 \rangle = \arg \langle \rho, V_1 \rangle_{tr} = \arg \langle p(\boldsymbol{\alpha}), v_1(\boldsymbol{\alpha}) \rangle_{\boldsymbol{\alpha}}.$$
(13)

We showed that this quantum asymptotic phase yields appropriate phase values, namely, it always increases with a constant frequency Ω_1 as $\dot{\Phi}_1(\rho) = \Omega_1$ with the evolution of ρ even in the strong quantum regime and reproduces the conventional asymptotic phase in the semiclassical regime [1]. We note that $\Phi_1(\rho)$ is not defined when $\langle V_1 \rangle = 0$.

This quantum asymptotic phase is a natural extension of the asymptotic phase for stochastic limit-cycle oscillators defined in terms of the slowest decaying eigenfunction of the backward Kolmogorov (Fokker-Planck) operator [47] from the Koopman operator viewpoint [48].

D. Multiple quantum asymptotic phases associated with different fundamental frequencies

In this study, we further extend the definition in the previous subsection and introduce multiple quantum asymptotic phases associated with several different fundamental frequencies by using the principal eigenvalues on different eigenvalue branches near the imaginary axis, and use them to characterize quantum signatures of synchronization in the strong quantum regime.

As shown in Fig. 1(a), in the strong quantum regime, multiple branches of the eigenvalues with different fundamental frequencies can exist near the imaginary axis, suggesting that not only the eigenvalue Λ_1 on the rightmost branch but also the eigenvalues with the slowest decay rates on the other branches play important roles; for comparison, see Fig. 1(b) for a typical example of the eigenvalue Λ_1 in the semiclassical regime, where only a single dominant branch of eigenvalues exist. We denote these eigenvalues by $\Lambda_1, \Lambda_2, \Lambda_3, \ldots$ and their imaginary parts, i.e., fundamental frequencies, by $\Omega_j = \text{Im } \overline{\Lambda_j} \ (j \ge 1)$, and call $\{\Lambda_j\}_{j\ge 1}$ the *principal eigenvalues*. Here, the first principal eigenvalue Λ_1 and the fundamental frequency Ω_1 are those introduced in the previous subsection. These principal eigenvalues on the individual branches are shown by red dots in Fig. 1(a).

In a similar manner to the phase function Φ_1 , we introduce the *j*-th quantum asymptotic phase function Φ_j (j = 2, 3, ...) of ρ as the argument of the *P*-representation of the eigenoperator V_j associated with the principal eigenvalue Λ_j satisfying $\mathcal{L}^* V_j = \overline{\Lambda_j} V_j$, namely,

$$\Phi_j(\rho) = \arg \langle V_j \rangle = \arg \langle \rho, V_j \rangle_{tr} = \arg \langle p(\boldsymbol{\alpha}), v_j(\boldsymbol{\alpha}) \rangle_{\boldsymbol{\alpha}}.$$
(14)

Note that $\Phi_j(\rho)$ is not defined when $\langle V_j \rangle = 0$. Since

$$\frac{d}{dt}\langle V_j \rangle = \langle \dot{\rho}, V_j \rangle_{tr} = \langle \mathcal{L}\rho, V_j \rangle_{tr} = \langle \rho, \mathcal{L}^* V_j \rangle_{tr} = \overline{\Lambda_j} \langle \rho, V_j \rangle_{tr} = \overline{\Lambda_j} \langle V_j \rangle, \tag{15}$$

we obtain $\langle V_j \rangle(t) = \langle V_j \rangle(0) e^{\overline{\Lambda_j} t}$ and therefore

$$\frac{d}{dt}\Phi_j(\rho) = \frac{d}{dt}\arg\langle V_j\rangle = \operatorname{Im}\overline{\Lambda_j} = \Omega_j.$$
(16)

Thus, Φ_j always increases with a constant frequency Ω_j with the evolution of ρ and plays the role of the asymptotic phase for any $j = 1, 2, 3, \ldots$

It should be noted that $\Phi_j(\rho)$ cannot be defined for the stationary state ρ_0 , because ρ_0 is the eigenfunction of the Liouville superoperator \mathcal{L} with the eigenvalue $\lambda_0 = 0$ and hence $\langle V_j \rangle = \langle \rho_0, V_j \rangle_{tr} = 0$ from the biorthogonality in Eq. (5). As will be shown later, for the stationary state of a periodically driven system, $\langle V_j \rangle$ takes a non-zero value and the above definition of the phase functions can be used for the analysis of phase locking.

We stress that in the classical deterministic limit, Φ_j $(j \ge 2)$ is not independent from Φ_1 and does not provide additional information, because Ω_j $(j \ne 1)$ is equal to Ω_1 according to the Koopman operator theory [55–58]. In the semiclassical regime, Ω_j $(j \ne 1)$ differs only slightly from Ω_1 as shown in Fig. 1(b) due to the effect of weak quantum noise and the corresponding decay rate for $j \ge 2$ is much larger, so the corresponding mode does not play an important role. However, as we see in the next section, Φ_j $(j \ge 2)$ is distinctly different from Φ_1 and the corresponding decay rate Re $\overline{\Lambda_j}$ $(j \ge 2)$ is comparable to Re $\overline{\Lambda_1}$ in the strong quantum regime, hence the phase function Φ_j $(j \ge 2)$ yields independent information from Φ_1 .

III. QUANTUM VAN DER POL OSCILLATOR WITH KERR EFFECT

A. Eigenvalues of the Liouville superoperator

As an example, we consider a quantum van der Pol oscillator with Kerr effect. The master equation is given by [1, 16, 28]

$$\dot{\rho} = \mathcal{L}_0 \rho, \quad \mathcal{L}_0 \rho = -i \left[H, \rho \right] + \gamma_1 \mathcal{D}[a^{\dagger}] \rho + \gamma_2 \mathcal{D}[a^2] \rho, \tag{17}$$

where $H = \omega_0 a^{\dagger} a + K a^{\dagger 2} a^2$, ω_0 is the natural frequency of the oscillator, K is the Kerr parameter, and γ_1 and γ_2 are the decay rates for negative damping and nonlinear damping, respectively. We added a subscript 0 to the Liouville operator as we will introduce an additional external drive later.

We first consider a strong quantum regime with large γ_2 and K, where only a small number of energy states participate in the system dynamics and the discrete nature of the energy spectrum plays important roles in the dynamics. We set the parameters as $\gamma_1 = 0.1$ and $(\omega_0, \gamma_2, K)/\gamma_1 = (300, 4, 100)$, which are the same as in our previous study [1]. In the numerical calculation, we approximately truncated the density operator as a large-dimensional $N \times N$ matrix and mapped it into a N^2 -dimensional vector in the double-ket notation [59]. We can then represent the Liouville operator by a $N^2 \times N^2$ matrix and obtain the asymptotic phase in Eq.(13) from the eigensystem of this matrix [1].

Figure 1(a) shows the eigenvalues of \mathcal{L}_0 near the imaginary axis obtained numerically. We can identify several branches of the eigenvalues near the imaginary axis characterized by the principal eigenvalues $\Lambda_1, \Lambda_2, \Lambda_3, \ldots$ whose fundamental frequencies $\Omega_1 = \text{Im } \overline{\Lambda_1}, \Omega_2 = \text{Im } \overline{\Lambda_2}, \Omega_3 = \text{Im } \overline{\Lambda_3}, \ldots$ are different from each other. Moreover, their decay rates, which are characterized by Re $\overline{\Lambda_1}$, Re $\overline{\Lambda_2}$, Re $\overline{\Lambda_3}, \ldots$ are comparable to each other. This indicates that not only the quantum asymptotic phase Φ_1 associated with the principle eigenvalue Λ_1 of the slowest decaying mode [1] but also those associated with the other principal eigenvalues $\Lambda_2, \Lambda_3, \ldots$ can play important roles in the dynamics. We choose a negative value for each Ω_j so that the corresponding Φ_j increases from 0 to 2π in the counterclockwise direction.

In Ref. [16], it is shown that, in the strong quantum regime, $|m+1\rangle\langle m|$ is an approximate eigenoperator of \mathcal{L}_0 with the eigenvalue

$$\widetilde{\lambda}_m = i[-\omega_0 - 2mK] - \frac{1}{2} \{\gamma_1(2m+3) + 2\gamma_2 m^2\},\tag{18}$$

where m = 0, 1, 2, ..., namely, $\mathcal{L}_0^{-1} | m + 1 \rangle \langle m | \approx \tilde{\lambda}_m^{-1} | m + 1 \rangle \langle m |$. As shown in Fig. 1(a), these eigenvalues correspond to the principal eigenvalues of \mathcal{L}_0 , i.e., $\Lambda_j \approx \tilde{\lambda}_{j-1}$, and thus $V_j \approx |j\rangle \langle j-1|$ (j = 1, 2, 3, 4). This indicates that that periodic transitions between the adjacent discrete energy states play important roles in the oscillatory behavior in the strong quantum regime, where the difference in the transition frequencies arises due to the unequal spacing of the energy levels characterized by the Kerr parameter in Eq. (18).

As a comparison, we next consider the semiclassical regime where γ_2 and K are sufficiently small and the semiclassical approximation can be taken [27]. We set the parameters as $\gamma_1 = 1$ and $(\omega_0, \gamma_2, K)/\gamma_1 = (0.1, 0.05, 0.025)$, which are



FIG. 1. Eigenvalues of the Liouville superoperator \mathcal{L}_0 near the imaginary axis. (a) Strong quantum regime. (b) Semiclassical regime. The red dot on the light-blue branch represents the principal eigenvalue Λ_1 with the the slowest decay rate. (c) Classical limit (schematic). In (a, b, c), individual branches of eigenvalues are shown with different colors and the red dots represent the principal eigenvalues Λ_j (j = 1, 2, 3, 4 from the right) with the fundamental frequencies in individual branches. In (a), the dotted lines indicate $\tilde{\lambda}_m$ (m = 0, 1, 2, 3) in Eq. (18).

the same as those used in Ref. [1]. In this regime, we can approximate Eq. (9) by a quantum Fokker-Planck equation for $p(\alpha)$ and the system can be approximately described as a Stuart-Landau oscillator (Hopf normal form) subjected to small quantum noise [1, 16, 28]. (Therefore, the conventional quantum vdP oscillator is also called a quantum Stuart-Landau oscillator recently [37, 39] and a more appropriate model of the quantum van der Pol oscillator has also been proposed [39, 40].)

Figure 1(b) shows the eigenvalues of \mathcal{L}_0 near the imaginary axis, where the principal eigenvalues are shown by red dots on the individual branches; $\Lambda_1 = \mu_1 + i\Omega_1$ ($\mu_1 < 0$) is on the rightmost light-blue branch. In contrast to the strong quantum regime, the rightmost branch of the eigenvalues, approximately given by a parabola $\hat{\lambda}_n =$ $i\Omega_1 n + \mu_1 n^2$ ($n = 0, \pm 1, \pm 2, ...$) passing through $\overline{\Lambda_1}$, is isolated from other branches of eigenvalues with faster decay rates; the relative decay rate Re $\overline{\Lambda_2}$ /Re $\overline{\Lambda_1} \approx -0.996/(-0.0736) \approx 13.5$ is more than three times larger than Re $\overline{\Lambda_2}$ /Re $\overline{\Lambda_1} \approx -0.65/(-0.15) \approx 4.33$ in the strong quantum regime in Fig. 1(a), indicating that only the rightmost branch is dominant in the semiclassical regime. Also, the fundamental frequencies of the other branches, defined as the smallest absolute imaginary part of the eigenvalues, are approximately equal to Ω_1 ; the small differences in the fundamental frequencies arise from small quantum noise. Thus, it is sufficient to consider only Λ_1 and introduce a single phase function Φ_1 in this regime.

The system in the classical limit, i.e., in the limit of vanishing quantum noise, is described by the drift term of the approximate quantum Fokker-Planck equation for $p(\alpha)$, which represents the Stuart-Landau oscillator and possesses a stable limit-cycle solution [1, 16, 28]. Figure 1(c) shows a schematic diagram of the eigenvalues of the differential operator L^+_{α} , which are equivalent to those of the backward Liouville operator \mathcal{L}^*_0 , in the classical limit. They are given in the form $\lambda_c = m\kappa_1 + in\omega_1$ ($m = 0, 1, 2, \ldots, n = 0, \pm 1, \pm 2$), where κ_1 is the real part of the largest negative eigenvalue and ω_1 is the pure-imaginary eigenvalue with the smallest absolute imaginary part. Thus, Φ_j ($j \geq 2$) is identical to Φ_1 and does not provide additional information, because Ω_j ($j \neq 2$) is identical to Ω_1 .

The above results suggest that the phase function Φ_j yields independent information from Φ_1 only in the strong quantum regime. The existence of several dominant fundamental frequencies in the strong quantum regime suggests that the system behaves like a torus rather than a limit cycle with a single fundamental frequency, and that we need to consider the phase functions Φ_2, Φ_3, \ldots associated with $\Lambda_2, \Lambda_3, \ldots$ in addition to Φ_1 and Λ_1 . Figure 2 shows a schematic picture of the torus behavior of the system with two fundamental frequencies.

B. Asymptotic phase functions in the strong quantum regime

In this subsection, we examine the validity of the quantum asymptotic phase functions in the strong quantum regime.

Figure 3 shows the asymptotic phase functions $\Phi_j(\alpha)$ for j = 1, 2, 3, and 4 of the pure coherent state α on the



FIG. 2. Schematic diagram of the torus behavior of the system with two fundamental frequencies.



FIG. 3. Asymptotic phase functions of the quantum van der Pol oscillator with Kerr effect in the strong quantum regime. (a) Φ_1 , (b) Φ_2 , (c) Φ_3 , and (d) Φ_4 . The parameters are $\gamma_1 = 0.1$ and $(\omega_0, \gamma_2, K)/\gamma_1 = (300, 4, 100)$. Figures in the bottom row show enlargements of the regions near the origin in the corresponding figures in the top row. In all figures, (x, p) = (2.5, 0) is chosen as the phase origin where $\Phi_j = 0$ (j = 1, 2, 3, 4).

complex plane ($x = \text{Re } \alpha, p = \text{Im } \alpha$). The parameters are the same as in Fig. 1(a). These asymptotic phase functions look similar, but they are associated with different fundamental frequencies and slightly different from each other near the origin as shown in the enlarged figures. Thus, they capture different oscillatory dynamics of the system. Though not shown, we may also draw similar asymptotic phase functions Φ_j for $j \ge 5$, which are less dominant. Here, the asymptotic phase functions for different values of j look similar to each other since they are identical in the classical limit [55–58], but the differences in their eigenfrequencies brought by the strong quantum effect take an important role in analyzing strong quantum signatures in synchronization.

To demonstrate that these quantum asymptotic phase functions yield appropriate phase values, we consider free oscillatory relaxation of ρ from a pure coherent initial state $\rho = |\alpha_0\rangle\langle\alpha_0|$ with $\alpha_0 = 1$ at t = 0 and measured the evolution of the expectation values of V_j and their arguments, i.e., their asymptotic phases $\Phi_j(\rho) = \arg\langle V_j\rangle$ (j = 1, 2, 3, 4), as well as those of the annihilation operator a for comparison. It is noted that, though the initial condition is a pure coherent state, the system state quickly becomes mixed due to the interaction with the reservoirs.

We can confirm that each $\langle V_j \rangle$ exhibits exponentially damped harmonic oscillations as shown in Fig. 4(a, c, e, g), and correspondingly, each $\Phi_j(\rho)$ gives constantly varying phase values with frequency Ω_j as shown in Fig. 4(b, d, f, h),



FIG. 4. Evolution of the expectation values of V_j (j = 1, 2, 3, 4) and the annihilation operator a (a, c, e, g, i) and their arguments (b, d, f, h, j) from a pure coherent initial state in the strong quantum regime. The parameters are $\gamma_1 = 0.1$ and $(\omega_0, \gamma_2, K)/\gamma_1 = (300, 4, 100)$. (a, c, e, g) $\langle V_j \rangle$, (b, d, f, h) $\arg \langle V_j \rangle$, (i) $\langle a \rangle$, (j) $\arg \langle a \rangle$.

verifying the validity of the definition of the quantum asymptotic phases. It is noted that different asymptotic phases independently capture different oscillation modes in the evolution of ρ . In contrast, $\langle a \rangle$ shown in Fig. 4(i) exhibits more complex oscillatory dynamics and the simple argument $\arg\langle a \rangle$ does not vary constantly with time as shown in Fig. 4(j). Thus, $\arg\langle a \rangle$ cannot be considered the asymptotic phase, although it is often used to define power spectra in the analysis of quantum synchronization. It is noted that the asymptotic phase is quantitatively different from the geometric angle also in the classical limit when the Kerr effect is added. In the strong quantum regime, due to the strong Kerr effect, the difference between the asymptotic phase Ω_j and simple argument $\arg\langle a \rangle$ can be observed more clearly than in the semiclassical regime or in the classical limit [1].



FIG. 5. Dependence of the order parameters on the frequency detuning Δ (divided by K) and driving strength E (divided by γ_1). (a) $|S_a|$. (b-e) $|S_j|$ (j = 1, 2, 3, 4). The parameters are $\gamma_1 = 0.1$ and $(\gamma_2, K)/\gamma_1 = (4, 100)$.



FIG. 6. Power spectra. The parameters are $\gamma_1 = 0.1$ and $(\Delta, \gamma_2, K, E)/\gamma_1 = (0, 4, 100, 1)$. (a) P_a . (b-e) P_j (j = 1, 2, 3, 4).

C. Revealing multiple phase-locking structure

We now analyze quantum synchronization of the vdP oscillator with a harmonic drive in the strong quantum regime using the proposed asymptotic phases.

The master equation in the rotating frame of the frequency ω_d of the harmonic drive is

$$\dot{\rho} = (\mathcal{L}_0 + \mathcal{L}_1)\rho,\tag{19}$$

where \mathcal{L}_0 is given by Eq. (17) with $H = -\Delta a^{\dagger}a + Ka^{\dagger 2}a^2$ and the harmonic drive is represented by $\mathcal{L}_1\rho = -i[iE(a-a^{\dagger}),\rho]$, where $\Delta = \omega_d - \omega_0$ is the frequency detuning of the harmonic drive from the oscillator and E is the strength of the harmonic drive [16, 28]. We use the same parameters as in Fig. 1(a) and vary the detuning parameter Δ by controlling ω_d while keeping the natural frequency ω_0 fixed.

In Ref. [16], Lörch *et al.* showed that this system under strong Kerr effect exhibits multiple phase locking to the harmonic drive at several detuning frequencies $\Delta = 2mK$ (m = 0, 1, 2, ...), i.e., at $\omega_d = \omega_0 + 2mK$, observed as multiple sharp Arnold tongues, while the corresponding classical system exhibits only a single broad Arnold tongue. It is stressed that K is the Kerr parameter, hence this is not the ordinary higher-harmonic phase locking at the frequencies $\omega_d = (p/q)\omega_0$ with p, q = 1, 2, 3, ... $(p/q \neq 1)$; the multiple Arnold tongues are related to the periodic transitions between the adjacent discrete energy states. It is also noted that such multiple phase locking is robust to thermal noise under the finite temperature environment (See Supplemental Material in Ref. [16]).

In Ref. [16], the following order parameter S_a and power spectrum P_a defined using the annihilation operator a are used to analyze the system:

$$S_a = |S_a|e^{i\theta_a} = \frac{\langle a \rangle}{\sqrt{\langle a^{\dagger}a \rangle}},\tag{20}$$

$$P_{a}(\omega) = \int_{-\infty}^{\infty} d\tau e^{i\omega\tau} \left(\langle a^{\dagger}(\tau)a(0) \rangle - \langle a^{\dagger}(\tau) \rangle \langle a(0) \rangle \right), \qquad (21)$$

where the expectation is taken with respect to the steady-state density operator obtained from Eq. (19).

Here, in addition to these quantities, we introduce the order parameters and the power spectra in terms of the quantum asymptotic phases (or, more precisely, the corresponding principal Koopman eigenoperators) defined in this study. They are defined using the left eigenoperators V_i of \mathcal{L}_0 as

$$S_j = |S_j| e^{i\theta_j} = \frac{\langle V_j \rangle}{\sqrt{\langle V_j^{\dagger} V_j \rangle}},\tag{22}$$

$$P_{j}(\omega) = \int_{-\infty}^{\infty} d\tau e^{i\omega\tau} \left(\langle V_{j}^{\dagger}(\tau) V_{j}(0) \rangle - \langle V_{j}^{\dagger}(\tau) \rangle \langle V_{j}(0) \rangle \right),$$
(23)

where $V_j^{\dagger}(\tau) = e^{\mathcal{L}_0^* \tau} V_j^{\dagger}(0) = e^{\Lambda_j \tau} V_j^{\dagger}(0)$ and $V_j^{\dagger}(0) = V_j^{\dagger}$ (j = 1, 2, 3 and 4). Here, $|S_a|$ and $|S_j|$ quantify the phase coherence of the system, while θ_a and θ_j characterize the averaged phases of the system relative to the harmonic drive. We note here that we can use V_j , which is defined in the original frame, in the rotating frame with the external drive since the eigenoperators V_j changes only its phase factor by the coordinate rotation, i.e., $e^{i\omega_d a^{\dagger} a t} V_j e^{-i\omega_d a^{\dagger} a t} = V_j e^{i\Omega t}$ with a constant Ω (see Appendix A). The resulting power spectra in the rotating frame are simply shifted from the power spectra in the original frame by $-\Omega$. Using these two quantities in Eqs. (22) and (23), we can analyze both the phase coherence and frequency characteristics of the oscillator in quantum synchronization.

Figure 5 shows the dependence of the order parameters $|S_a|$ and $|S_j|$ on the detuning Δ and strength E of the harmonic drive. In Fig. 5(a) showing $|S_a|$, several Arnold tongues representing phase locking of the oscillator at different frequencies are observed [16]. Figures 5(b)-(e) show $|S_j|$ for j = 1, 2, 3 and 4, respectively. It is remarkable that the Arnold tongues in Fig. 5(a) are clearly decomposed into individual Arnold tongues around $\Delta = 2(j-1)K$ in Figs. 5(b)-5(e). This indicates that the order parameters in terms of the asymptotic phases can capture the phase locking dynamics at $\omega_d = \omega_0 + 2(j-1)K$ for different j individually.

Similarly, Fig. 6 shows the power spectra P_a and P_j for $\Delta = 0$. Multiple peaks of P_a in Fig. 6(a), which indicate multiple phase locking of the oscillator to the harmonic drive, are clearly decomposed into individual peaks around $\omega = \Delta - 2(j-1)K$ in Figs. 6(b)-(e) for P_j (j = 1, 2, 3, and 4). The Arnold tongue and power spectrum are sharper when the decay rate characterized by Re Λ_j is smaller. Though not shown, we can also detect even smaller tongues and peaks with $j \geq 5$.

The above results reveal that, in the strong quantum regime, the system behaves like a torus with several fundamental frequencies and each of the associated oscillating mode individually exhibits phase locking to the harmonic

IV. CONCLUSION

In this study, we defined multiple quantum asymptotic phases of quantum nonlinear oscillators in terms of the eigenoperators of the adjoint Liouville superoperator associated with several fundamental frequencies from the Koopman operator viewpoint, which extends our previous definition of the quantum asymptotic phase associated with the slowest decaying mode of the system. We introduced the order parameters and power spectra in terms of the proposed quantum asymptotic phases and applied them to the analysis of a quantum van der Pol oscillator with Kerr effect. We successfully revealed that the multiple phase locking of the system with a harmonic drive at several different frequencies [16], which is an explicit quantum signature observed only in the strong quantum regime, can be interpreted as synchronization on a torus rather than on a simple limit cycle.

Though not discussed in this paper, it will also be interesting to introduce the amplitude functions in addition to the quantum asymptotic phases by extending the definition for classical stochastic oscillators [48, 61], which can be defined using the eigenfunctions associated with the eigenvalues on the real axis. The phase-amplitude description that uses both the phases and amplitudes may be applied to the analysis of quantum complete synchronization [19]. Also, investigating the mutual synchronization between two quantum nonlinear oscillators [17, 29, 62] by using quantum asymptotic phase functions is also a future work. We may be able to introduce a new measure for quantum synchronization between two oscillators, e.g., the normalized correlator [20], by using quantum asymptotic phase functions. It may be also interesting to extend our definition of the asymptotic phase for quantum synchronization in non-Markov open quantum systems [63].

We expect that the proposed definition of the quantum asymptotic phases will serve as a fundamental tool for analyzing strong quantum effects in synchronization [16, 17] and will be useful for future applications of quantum synchronization in the growing fields of quantum technologies.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available upon reasonable request from the authors.

ACKNOWLEDGMENTS

Numerical simulations are performed by using QuTiP numerical toolbox [64, 65]. We acknowledge JSPS KAK-ENHI JPJSBP120202201, JP20J13778, JP22K14274, JP22K11919, JP22H00516, and JST CREST JP-MJCR1913 for financial support.

Appendix A: Proof of $e^{i\omega_d a^{\dagger} a t} V_j e^{-i\omega_d a^{\dagger} a t} = V_j e^{i\Omega t}$

The time evolution of $\tilde{V}_i(t) = e^{i\omega_d a^{\dagger} a t} V_i e^{-i\omega_d a^{\dagger} a t}$ is given by

resulting in the torus-like behavior.

$$\frac{dV_j(t)}{dt} = i\omega_d a^{\dagger} a e^{i\omega_d a^{\dagger} a t} V_j e^{-i\omega_d a^{\dagger} a t} + e^{i\omega_d a^{\dagger} a t} V_j (-i\omega_d a^{\dagger} a) e^{-i\omega_d a^{\dagger} a t} = i\omega_d e^{i\omega_d a^{\dagger} a t} [a^{\dagger} a, V_j] e^{-i\omega_d a^{\dagger} a t}.$$
 (A1)

Because the adjoint superoperator \mathcal{L}_0^* couples only the elements of the form $|n+k\rangle\langle m+k|$ to the elements of the form $|n+k'\rangle\langle m+k'|$ for an arbitrary pair of integers $\{n,m\}$ and integers k,k' [66, 67], the eigenoperators V_j can be explicitly written as

$$V_j = \sum_{n=0}^{\infty} v_j(n) |n\rangle \langle n+k|.$$
(A2)

Therefore, we obtain $[a^{\dagger}a, V_j] = -kV_j$ and

$$\frac{d\tilde{V}_j(t)}{dt} = i\omega_d e^{i\omega_d a^{\dagger} a t} [a^{\dagger} a, V_j] e^{-i\omega_d a^{\dagger} a t} = -i\omega_d k \tilde{V}_j(t).$$
(A3)

If we assume $\Omega = -i\omega_d k$, we obtain $\tilde{V}_i(t) = V_i e^{-i\omega_d kt} = V_i e^{i\Omega t}$.

- [1] Yuzuru Kato and Hiroya Nakao. A definition of the asymptotic phase for quantum nonlinear oscillators from the Koopman operator viewpoint. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 32(6):063133, 2022.
- [2] Arthur T Winfree. The geometry of biological time. Springer, New York, 2001.
- [3] Yoshiki Kuramoto. Chemical oscillations, waves, and turbulence. Springer, Berlin, 1984.
- [4] Arkady Pikovsky, Michael Rosenblum, and Jürgen Kurths. Synchronization: a universal concept in nonlinear sciences. Cambridge University Press, Cambridge, 2001.
- [5] Hiroya Nakao. Phase reduction approach to synchronisation of nonlinear oscillators. Contemporary Physics, 57(2):188–214, 2016.
- [6] G Bard Ermentrout and David H Terman. Mathematical foundations of neuroscience. Springer, New York, 2010.
- [7] SH Strogatz. Nonlinear dynamics and chaos. Westview Press, 1994.
- [8] Matthew H Matheny, Jeffrey Emenheiser, Warren Fon, Airlie Chapman, Anastasiya Salova, Martin Rohden, Jarvis Li, Mathias Hudoba de Badyn, Márton Pósfai, Leonardo Duenas-Osorio, et al. Exotic states in a simple network of nanoelectromechanical oscillators. *Science*, 363(6431):eaav7932, 2019.
- [9] Sören Kreinberg, Xavier Porte, David Schicke, Benjamin Lingnau, Christian Schneider, Sven Höfling, Ido Kanter, Kathy Lüdge, and Stephan Reitzenstein. Mutual coupling and synchronization of optically coupled quantum-dot micropillar lasers at ultra-low light levels. *Nature Communications*, 10(1):1539, 2019.
- [10] Hanuman Singh, S Bhuktare, A Bose, A Fukushima, K Yakushiji, S Yuasa, H Kubota, and Ashwin A Tulapurkar. Mutual synchronization of spin-torque nano-oscillators via oersted magnetic fields created by waveguides. *Physical Review Applied*, 11(5):054028, 2019.
- [11] MF Colombano, G Arregui, NE Capuj, A Pitanti, J Maire, A Griol, B Garrido, A Martinez, CM Sotomayor-Torres, and D Navarro-Urrios. Synchronization of optomechanical nanobeams by mechanical interaction. *Physical Review Letters*, 123(1):017402, 2019.
- [12] Jiteng Sheng, Xinrui Wei, Cheng Yang, and Haibin Wu. Self-organized synchronization of phonon lasers. *Physical Review Letters*, 124(5):053604, 2020.
- [13] Arif Warsi Laskar, Pratik Adhikary, Suprodip Mondal, Parag Katiyar, Sai Vinjanampathy, and Saikat Ghosh. Observation of quantum phase synchronization in spin-1 atoms. *Physical Review Letters*, 125(1):013601, 2020.
- [14] VR Krithika, Parvinder Solanki, Sai Vinjanampathy, and TS Mahesh. Observation of quantum phase synchronization in a nuclear-spin system. *Physical Review A*, 105(6):062206, 2022.
- [15] Martin Koppenhöfer, Christoph Bruder, and Alexandre Roulet. Quantum synchronization on the IBM Q system. Physical Review Research, 2(2):023026, 2020.
- [16] Niels Lörch, Ehud Amitai, Andreas Nunnenkamp, and Christoph Bruder. Genuine quantum signatures in synchronization of anharmonic self-oscillators. *Physical Review Letters*, 117(7):073601, 2016.
- [17] Niels Lörch, Simon E Nigg, Andreas Nunnenkamp, Rakesh P Tiwari, and Christoph Bruder. Quantum synchronization blockade: Energy quantization hinders synchronization of identical oscillators. *Physical Review Letters*, 118(24):243602, 2017.
- [18] Simon E Nigg. Observing quantum synchronization blockade in circuit quantum electrodynamics. Physical Review A, 97(1):013811, 2018.
- [19] A Mari, A Farace, N Didier, V Giovannetti, and R Fazio. Measures of quantum synchronization in continuous variable systems. *Physical Review Letters*, 111(10):103605, 2013.
- [20] Talitha Weiss, Andreas Kronwald, and Florian Marquardt. Noise-induced transitions in optomechanical synchronization. New Journal of Physics, 18(1):013043, 2016.
- [21] Michael R Hush, Weibin Li, Sam Genway, Igor Lesanovsky, and Andrew D Armour. Spin correlations as a probe of quantum synchronization in trapped-ion phonon lasers. *Physical Review A*, 91(6):061401, 2015.
- [22] Talitha Weiss, Stefan Walter, and Florian Marquardt. Quantum-coherent phase oscillations in synchronization. Physical Review A, 95(4):041802, 2017.
- [23] Noufal Jaseem, Michal Hajdušek, Parvinder Solanki, Leong-Chuan Kwek, Rosario Fazio, and Sai Vinjanampathy. Generalized measure of quantum synchronization. *Physical Review Research*, 2(4):043287, 2020.
- [24] Tony E Lee and HR Sadeghpour. Quantum synchronization of quantum van der Pol oscillators with trapped ions. *Physical Review Letters*, 111(23):234101, 2013.
- [25] Stefan Walter, Andreas Nunnenkamp, and Christoph Bruder. Quantum synchronization of a driven self-sustained oscillator. Physical Review Letters, 112(9):094102, 2014.
- [26] Sameer Sonar, Michal Hajdušek, Manas Mukherjee, Rosario Fazio, Vlatko Vedral, Sai Vinjanampathy, and Leong-Chuan Kwek. Squeezing enhances quantum synchronization. *Physical Review Letters*, 120(16):163601, 2018.

- [27] Yuzuru Kato, Naoki Yamamoto, and Hiroya Nakao. Semiclassical phase reduction theory for quantum synchronization. *Physical Review Research*, 1(3):033012, 2019.
- [28] Yuzuru Kato and Hiroya Nakao. Semiclassical optimization of entrainment stability and phase coherence in weakly forced quantum limit-cycle oscillators. *Physical Review E*, 101(1):012210, 2020.
- [29] Tony E Lee, Ching-Kit Chan, and Shenshen Wang. Entanglement tongue and quantum synchronization of disordered oscillators. *Physical Review E*, 89(2):022913, 2014.
- [30] Dirk Witthaut, Sandro Wimberger, Raffaella Burioni, and Marc Timme. Classical synchronization indicates persistent entanglement in isolated quantum systems. *Nature Communications*, 8:14829, 2017.
- [31] Alexandre Roulet and Christoph Bruder. Quantum synchronization and entanglement generation. *Physical Review Letters*, 121(6):063601, 2018.
- [32] Najmeh Es' haqi Sani, Gonzalo Manzano, Roberta Zambrini, and Rosario Fazio. Synchronization along quantum trajectories. *Physical Review Research*, 2(2):023101, 2020.
- [33] Yuzuru Kato and Hiroya Nakao. Enhancement of quantum synchronization via continuous measurement and feedback control. New Journal of Physics, 23(1):013007, 2021.
- [34] Yuzuru Kato and Hiroya Nakao. Instantaneous phase synchronization of two decoupled quantum limit-cycle oscillators induced by conditional photon detection. *Physical Review Research*, 3(1):013085, 2021.
- [35] Wenlin Li, Najmeh Es' haqi Sani, Wen-Zhao Zhang, and David Vitali. Quantum zeno effect in self-sustaining systems: Suppressing phase diffusion via repeated measurements. *Physical Review A*, 103(4):043715, 2021.
- [36] Minghui Xu, David A Tieri, EC Fine, James K Thompson, and Murray J Holland. Synchronization of two ensembles of atoms. *Physical Review Letters*, 113(15):154101, 2014.
- [37] W-K Mok, L-C Kwek, and H Heimonen. Synchronization boost with single-photon dissipation in the deep quantum regime. *Physical Review Research*, 2(3):033422, 2020.
- [38] Alexandre Roulet and Christoph Bruder. Synchronizing the smallest possible system. *Physical Review Letters*, 121(6):053601, 2018.
- [39] A Chia, LC Kwek, and C Noh. Relaxation oscillations and frequency entrainment in quantum mechanics. Physical Review E, 102(4):042213, 2020.
- [40] Lior Ben Arosh, MC Cross, and Ron Lifshitz. Quantum limit cycles and the rayleigh and van der Pol oscillators. Physical Review Research, 3(1):013130, 2021.
- [41] Albert Cabot, Gian Luca Giorgi, Fernando Galve, and Roberta Zambrini. Quantum synchronization in dimer atomic lattices. *Physical Review Letters*, 123(2):023604, 2019.
- [42] Albert Cabot, Gian Luca Giorgi, and Roberta Zambrini. Metastable quantum entrainment. New Journal of Physics, 23(10):103017, 2021.
- [43] Fernando Galve, Gian Luca Giorgi, and Roberta Zambrini. Quantum correlations and synchronization measures. In Lectures on General Quantum Correlations and their Applications, pages 393–420. Springer, 2017.
- [44] H Eneriz, DZ Rossatto, Francisco A Cárdenas-López, E Solano, and M Sanz. Degree of quantumness in quantum synchronization. Scientific Reports, 9(1):1–9, 2019.
- [45] Parvinder Solanki, Noufal Jaseem, Michal Hajdušek, and Sai Vinjanampathy. Role of coherence and degeneracies in quantum synchronization. *Physical Review A*, 105(2):L020401, 2022.
- [46] Wataru Setoyama and Yoshihiko Hasegawa. Lie algebraic quantum phase reduction. arXiv preprint arXiv:2208.12006, 2022.
- [47] Peter J Thomas and Benjamin Lindner. Asymptotic phase for stochastic oscillators. *Physical Review Letters*, 113(25):254101, 2014.
- [48] Yuzuru Kato, Jinjie Zhu, Wataru Kurebayashi, and Hiroya Nakao. Asymptotic phase and amplitude for classical and semiclassical stochastic oscillators via Koopman operator theory. *Mathematics*, 9(18):2188, 2021.
- [49] Ryan Hamerly and Hideo Mabuchi. Optical devices based on limit cycles and amplification in semiconductor optical cavities. *Physical Review Applied*, 4(2):024016, 2015.
- [50] Howard J Carmichael. Statistical Methods in Quantum Optics 1, 2. Springer, New York, 2007.
- [51] Crispin W Gardiner. Quantum Noise. Springer, New York, 1991.
- [52] Andy CY Li, F Petruccione, and Jens Koch. Perturbative approach to Markovian open quantum systems. Scientific Reports, 4:4887, 2014.
- [53] Daniel A Lidar, Isaac L Chuang, and K Birgitta Whaley. Decoherence-free subspaces for quantum computation. *Physical Review Letters*, 81(12):2594, 1998.
- [54] Kevin E Cahill and Roy J Glauber. Density operators and quasiprobability distributions. *Physical Review*, 177(5):1882, 1969.
- [55] Alexandre Mauroy, Igor Mezić, and Jeff Moehlis. Isostables, isochrons, and Koopman spectrum for the action-angle representation of stable fixed point dynamics. *Physica D: Nonlinear Phenomena*, 261:19–30, 2013.
- [56] Y Susuki Mauroy and I Mezic. The Koopman operator in systems and control, 2020.
- [57] Sho Shirasaka, Wataru Kurebayashi, and Hiroya Nakao. Phase-amplitude reduction of transient dynamics far from attractors for limit-cycling systems. Chaos, 27(2):023119, 2017.
- [58] Yoshiki Kuramoto and Hiroya Nakao. On the concept of dynamical reduction: the case of coupled oscillators. *Philosophical Transactions of the Royal Society A*, 377(2160):20190041, 2019.
- [59] Victor V Albert. Lindbladians with multiple steady states: theory and applications. arXiv preprint arXiv:1802.00010, 2018.

- [60] Peter Grindrod and Ebrahim L Patel. Phase locking to the n-torus. IMA Journal of Applied Mathematics, 81(1):152–164, 2016.
- [61] Alberto Pérez-Cervera, Benjamin Lindner, and Peter J Thomas. Isostables for stochastic oscillators. *Physical Review Letters*, 127(25):254101, 2021.
- [62] Stefan Walter, Andreas Nunnenkamp, and Christoph Bruder. Quantum synchronization of two van der Pol oscillators. Annalen der Physik, 527(1-2):131–138, 2015.
- [63] Göktuğ Karpat, İskender Yalçinkaya, Barış Çakmak, Gian Luca Giorgi, and Roberta Zambrini. Synchronization and non-markovianity in open quantum systems. *Physical Review A*, 103(6):062217, 2021.
- [64] JR Johansson, PD Nation, and Franco Nori. Qutip: An open-source python framework for the dynamics of open quantum systems. Computer Physics Communications, 183(8):1760–1772, 2012.
- [65] JR Johansson, PD Nation, and Franco Nori. Qutip 2: A python framework for the dynamics of open quantum systems. Computer Physics Communications, 184:1234–1240, 2013.
- [66] Stephen M Barnett and Stig Stenholm. Spectral decomposition of the Lindblad operator. Journal of Modern Optics, 47(14-15):2869–2882, 2000.
- [67] Hans-Jürgen Briegel and Berthold-Georg Englert. Quantum optical master equations: The use of damping bases. Physical Review A, 47(4):3311, 1993.