Coloring (P_5, gem) -free graphs with $\Delta - 1$ colors

Daniel W. Cranston^{*}

Hudson Lafayette[†]

Landon Rabern[‡]

June 4, 2020

Abstract

The Borodin–Kostochka Conjecture states that for a graph G, if $\Delta(G) \geq 9$ and $\omega(G) \leq \Delta(G) - 1$, then $\chi(G) \leq \Delta(G) - 1$. We prove the Borodin–Kostochka Conjecture for (P_5, gem) -free graphs, i.e., graphs with no induced P_5 and no induced $K_1 \vee P_4$.

1 Introduction

Every graph G satisifies $\chi(G) \leq \Delta(G) + 1$. (We denote by $\Delta(G)$, $\omega(G)$, and $\chi(G)$ the maximum degree, clique number, and chromatic number of G.) To see this, we greedily coloring the vertices of G in any order. In 1941, Brooks [2] strengthened this bound to $\chi(G) \leq \Delta(G)$.

Brooks' Theorem. Let G be a graph. If $\Delta(G) \geq 3$ and $\omega(G) \leq \Delta(G)$, then $\chi(G) \leq \Delta(G)$.

In 1977, Borodin and Kostochka [1] conjectured a further strengthening of Brooks' bound.

Borodin–Kostochka Conjecture. Let G be a graph. If $\Delta(G) \geq 9$ and $\omega(G) \leq \Delta(G) - 1$, then $\chi(G) \leq \Delta(G) - 1$.

If true, the Borodin–Kostochka Conjecture is best possible in the following two ways. First, if the hypothesis $\Delta(G) \geq 9$ is weakened to $\Delta(G) \geq 8$ then the conjecture is false, as witnessed by G_1 , on the left in Figure 1. Note that G_1 is a counterexample to this stronger version since $\Delta(G_1) = 8$ and $\omega(G_1) = 6 \leq 7 = \Delta(G_1) - 1$, but $\chi(G) = 8 > 7 = \Delta(G_1) - 1$. Second, if the hypothesis $\omega(G) \leq \Delta(G) - 1$ is strengthened to $\omega(G) \leq \Delta(G) - 2$, then the bound $\chi(G) \leq \Delta(G) - 1$ cannot be strengthened to $\chi(G) \leq \Delta(G) - 2$, as witnessed by G_2 , on the right in Figure 1. For each $t \geq 9$, note that G_2 is a counterexample to this stronger version, since $\Delta(G) = t \geq 9$ and $\omega(G_2) = \Delta(G_2) - 2$, but $\chi(G_2) = \Delta(G_2) - 1$.

By Brooks' Theorem, each graph G with $\chi(G) > \Delta(G) \ge 9$ contains $K_{\Delta(G)+1}$. So the Borodin– Kostochka Conjecture asserts that each G with $\chi(G) = \Delta(G) \ge 9$ contains $K_{\Delta(G)}$. This conjecture has been proved for many interesting classes of graphs, particularly those defined by forbidden subgraphs. In 2013, Cranston and Rabern [6] proved it for claw-free graphs.

^{*}Department of Mathematics and Applied Mathematics, Viriginia Commonwealth University, Richmond, VA; dcranston@vcu.edu

[†]Department of Mathematics and Applied Mathematics, Viriginia Commonwealth University, Richmond, VA; lafayettehl@vcu.edu

[‡]landon.rabern@gmail.com



Figure 1: Each bold edge denotes a complete bipartite graph. We have $\Delta(G_1) = 8$, $\omega(G_1) = 6$, but $\chi(G_1) = 8$. And we have $\Delta(G_2) = t$, $\omega(G_2) = t - 2$, but $\chi(G_2) = t - 1$.

Theorem 1.1 ([6]). Every claw-free graph with $\chi(G) \ge \Delta(G) \ge 9$ contains $K_{\Delta(G)}$.

The strongest partial result toward the Borodin–Kostochka Conjecture is due to Reed [11]. In 1999, he proved this conjecture for every graph G with $\Delta(G)$ sufficiently large.

Theorem 1.2 ([11]). Every graph with $\chi(G) = \Delta(G) \ge 10^{14}$ contains $K_{\Delta(G)}$.

Reed stated that a more careful analysis of his argument could reduce the lower bound to about 10^3 , but definitely not to 10^2 .

In 1998, Reed also conjectured [12] that $\chi(G) \leq \lceil (\omega(G) + \Delta(G) + 1)/2 \rceil$. This conjecture, now called Reed's Conjecture, is weaker than the Borodin–Kostochka Conjecture when $\omega(G) \in \{\Delta(G) - 1, \Delta(G) - 2\}$, equivalent to it when $\omega(G) \in \{\Delta(G) - 3, \Delta(G) - 4\}$, and stronger otherwise. Chudnovsky, Karthick, Maceli, and Maffray [4] proved Reed's Conjecture for all (P_5 , gem)-free graphs. In this note we use their structure theorem for (P_5 , gem)-free graphs, as well as one of their key lemmas [4, Theorem 3 and Lemma 2], to prove the Borodin–Kostochka Conjecture for the same class of graphs.

Main Theorem. Let G be a (P_5, gem) -free graph. If $\Delta(G) \ge 9$ and $\omega(G) \le \Delta - 1$, then $\chi(G) \le \Delta(G) - 1$.

2 Definitions and Notation

All of our graphs are finite and simple. Most of our definitions follow [13], but we highlight a few terms. By coloring we mean a proper vertex coloring. A graph is k-colorable (or has a k-coloring) if there exists a coloring $\varphi: V(G) \to \{1, \ldots, k\}$. Given two graphs S and T, the join, denoted $S \lor T$ is formed from their disjoint union by adding all edges with one endpoint in S and the other in T. For graphs H_1, \ldots, H_s , a graph G is (H_1, \ldots, H_s) -free if G does not contain any of H_1, \ldots, H_s as an induced subgraph. Thus, the class of (P_5, gem) -free graphs is all graphs that do not contain an induced P_5 or an induced gem. Here gem is $K_1 \lor P_4$; see Figure 2.

We use the following notation from [4]. Let G be a graph and $X, Y \subseteq V(G)$. Let [X, Y] denote the set of edges that have one end in X and other end in Y. If every vertex in X is adjacent to every vertex in Y, then X is *complete* to Y or [X, Y] is complete; if $[X, Y] = \emptyset$, then X is *anticomplete* to Y. A set X is a *homogeneous set* if every vertex with a neighbor in X is complete to X.

An expansion of a graph H is any graph G such that V(G) can be partitioned into |V(H)|non-empty sets Q_v , for each $v \in V(H)$, such that $[Q_u, Q_v]$ is complete if $uv \in E(H)$, and $[Q_u, Q_v]$



Figure 2: A P_5 and a gem.

is anticomplete if $uv \notin E(H)$. An expansion of a graph is a *clique expansion* if each Q_v is a clique, and is a P_4 -free expansion if each Q_v induces a P_4 -free graph.

For each $i \in \{1, \ldots, 10\}$, the class of graphs \mathcal{G}_i is all the P_4 -free expansions of the single graph shown in Case *i* below (in Figures 3–6), where each Q_i is a single vertex. The class \mathcal{G}_i^* is all clique expansions of the graph in Case *i*. The classes \mathcal{H} and \mathcal{H}^* are defined nearly analogously for the graph in Case 11. (There is a slight difference, which we explain when we handle Case 11.)

3 Lemmas and Reductions

In this section, we present lemmas that we will use to prove our Main Theorem. We will assume the Main Theorem is false and choose G to be a counterexample that is *vertex-critical*; that is $\chi(G-v) < \chi(G)$ for each $v \in V(G)$. Ultimately, we reach a contradiction, by constructing a $(\Delta(G)-1)$ -coloring of G. To construct this coloring, we repeatedly use the following easy lemma.

Lemma 3.1. Fix $k \in \mathbb{Z}^+$. Let G be a graph and let I_1, \ldots, I_t be pairwise disjoint independent sets of G. If $G - \bigcup_{i=1}^t I_i$ is (k-t-1)-degenerate, then $\chi(G) \leq k$.

Proof. Let G, k, and I_1, \ldots, I_t satisfy the hypotheses. If $G - \bigcup_{j=1}^t I_j$ is (k-t-1)-degenerate, then it has a (k-t)-coloring φ' . We extend φ' to a k-coloring of G by giving each I_j its own color. \Box

We will use Lemma 3.1 to 8-color G, typically with t = 2. Let $G' := G - \bigcup_{j=1}^2 I_j$. To prove that G' is 5-degenerate we give a vertex order $\sigma = (v_1, \ldots, v_n)$ such that each v_i has at most 5 neighbours earlier in σ . For a vertex partition $S_1 \uplus \cdots \uplus S_t$ of V(G) we write (S_1, \ldots, S_t) to denote a vertex order σ where all vertices in S_i come before all vertices in S_{i+1} , for each i, and vertices within each S_i are ordered arbitrarily.

Lemma 3.2. Every vertex-critical counterexample G to the Borodin–Kostochka Conjecture has $\delta(G) \ge \Delta(G) - 1$. In particular, $\delta(G) \ge 8$ and $|d(v) - d(w)| \le 1$ for all $v, w \in V(G)$.

Proof. Assume that G is a vertex-critical counterexample to the Borodin–Kostochka Conjecture. Suppose there exists $v \in V(G)$ such that $d(v) < \Delta(G) - 1$. Since G is vertex-critical, G - v has a $(\Delta(G) - 1)$ -coloring φ . To extend φ to v, we simply color v with a color unused on $N_G(v)$.

We want to reuse the idea in the proof of Lemma 3.2 to show that other induced subgraphs cannot appear in G. For this, we use notation and a lemma from [7]. For a graph H, a d_1 -assignment L gives to each $v \in V(H)$ a set (of allowable colors) L(v) such that $|L(v)| = d_H(v) - 1$. A graph H is d_1 -choosable if H has a proper coloring φ with $\varphi(v) \in L(v)$ for every d_1 -assignment L.

Lemma 3.3 ([7]). If G is vertex-critical and $\chi(G) = \Delta(G)$, then G cannot contain any nonempty, d_1 -choosable, induced subgraph H. So such a G contains neither of the following as induced subgraphs:

- $K_3 \vee 3K_2$ or
- $K_4 \vee H$, where V(H) contains two disjoint pairs of nonadjacent vertices.

Proof. We begin with the first statement. Suppose, to the contrary, that G contains an induced subgraph H, where G and H satisfy the hypotheses. Since G is vertex-critical, G - H has a $(\Delta(G) - 1)$ -coloring φ . For each $v \in V(H)$, let L(v) denote the colors in $\{1, \ldots, \Delta(G) - 1\}$ that are unused by φ on $N_G(v)$. Thus, $|L(v)| \geq \Delta(G) - 1 - (d_G(v) - d_H(v)) \geq d_H(v) - 1$. Now we can extend φ to H precisely because H is d_1 -choosable. This gives a $(\Delta(G) - 1)$ -coloring of G, which is a contradiction; this proves the first statement. The second statement follows from [7, Lemmas 3.3 & 3.10], where it is proved that the two subgraphs listed above are d_1 -choosable.

Copycat Lemma. Let G be a vertex-critical counterexample to the Borodin–Kostochka Conjecture. If G contains nonempty, disjoint homogeneous sets A and B such that A and B are cliques and $N(A) \subseteq N(B)$, then |A| > |B|.

Proof. Let G, A, and B satisfy the hypotheses. Suppose, for the sake of contradiction, that $|A| \leq |B|$. Since G is vertex-critical, G - A has a $(\Delta(G) - 1)$ -coloring φ . Since $|A| \leq |B|$, $N(A) \subseteq N(B)$, and B is a homogeneous set, we can extend φ to G by coloring A with colors used on B. Thus, G is not a counterexample to the Borodin–Kostochka Conjecture, a contradiction. \Box

A graph class \mathcal{G} is *hereditary* if for each $G \in \mathcal{G}$ we have $H \in \mathcal{G}$ for each induced subgraph H of G. Every class of graphs characterized by a list of forbidden induced subgraphs is a hereditary class; in particular, the class of (P_5, gem) -free graphs is hereditary.

Theorem 3.4 (Kostochka [10], Catlin [3]). Let \mathcal{G} be a hereditary class of graphs. If the Borodin– Kostochka Conjecture is false for some $G \in \mathcal{G}$, then it is false for some $G \in \mathcal{G}$ with $\Delta(G) = 9$.

Proof. We assume the Borodin–Kostochka Conjecture is true for all $G \in \mathcal{G}$ with $\Delta(G) = 9$ and show that it is true for all $G \in \mathcal{G}$. Suppose instead that some $G \in \mathcal{G}$ with $\Delta(G) > 9$ is a counterexample; in particular, $\omega(G) \leq \Delta(G) - 1$. Among all such G, choose one to minimize $\Delta(G)$. We want to find a maximum independent set I that intersects each clique of size $\Delta(G) - 1$. If $\omega(G) < \Delta(G) - 1$, then any maximum independent set I suffices. Otherwise, I is guaranteed by a result of King [9]. Let G' := G - I. Note that $\omega(G') \leq \Delta(G) - 2$ and $\Delta(G') \leq \Delta(G) - 1$, since I is maximum.

If $\Delta(G') \leq \Delta(G) - 3$, then we can greedily color G' with at most $\Delta(G) - 2$ colors. By using a new color on I, we get a $(\Delta(G) - 1)$ -coloring of G, a contradiction. If $\Delta(G') = \Delta(G) - 2$, then G' has a $(\Delta(G) - 2)$ -coloring by Brooks' Theorem, since $\omega(G') \leq \Delta(G) - 2$. By using a new color on I, we again get a $(\Delta(G) - 1)$ -coloring of G, a contradiction. So we must have $\Delta(G') = \Delta(G) - 1$ and $\omega(G') \leq \Delta(G) - 2$. Since $G' \in \mathcal{G}$ and $\Delta(G') < \Delta(G)$, we know that G' is not a counterexample to the Borodin–Kostochka Conjecture. In particular G' has a $(\Delta(G') - 1)$ -coloring. By using a new color on I, we extend this coloring of G' to a $(\Delta(G) - 1)$ -coloring of G, a contradiction.

We conclude this section with two results of Chudnovsky, Karthick, Maceli, and Maffray.

Theorem 3.5 ([4, Theorem 3]). If G is a connected (P_5 , gem)-free graph that contains an induced C_5 , then either $G \in \mathcal{H}$ or $G \in \mathcal{G}_i$, for some $i \in \{1, \ldots, 10\}$.

Theorem 3.6 ([4, 8]). Fix $i \in \{1, ..., 10\}$. For every $G \in \mathcal{G}_i$ (resp. $G \in \mathcal{H}$) there is $G^* \in \mathcal{G}_i^*$ (resp. $G^* \in \mathcal{H}^*$) such that $\omega(G) = \omega(G^*)$ and $\chi(G) = \chi(G^*)$. Further, G^* is an induced subgraph of G.

4 Proof of Main Result

Main Theorem. Let G be a (P_5, gem) -free graph. If $\Delta(G) \ge 9$ and $\omega(G) \le \Delta(G) - 1$, then $\chi(G) \le \Delta(G) - 1$.

Proof. Suppose the theorem is false. Let G be a counterexample that minimizes $\Delta(G)$. Further, we can choose G to be vertex-critical. By Theorem 3.4, we can assume that $\Delta(G) = 9$. Next we show that we can also assume that either $G \in \mathcal{G}_i^*$ for some $i \in \{1, \ldots, 10\}$ or $G \in \mathcal{H}^*$.

If G is perfect, then $\chi(G) = \omega(G) \leq \Delta(G) - 1$, a contradiction. Since G is not perfect, the Strong Perfect Graph Theorem [5] implies that G must contain an odd hole or odd anti-hole (that is, either G or its complement contains an induced odd cycle of length at least 5). Every odd hole of length at least 7 contains a P_5 as an induced subgraph. Similarly, every odd antihole of length at least 7 contains a gem as an induced subgraph. Since G is (P_5, gem) -free, G must contain a hole or antihole of length 5. In fact, these are both congruent to C_5 . So, by Theorem 3.5, either $G \in G_i$ for some $i \in \{1, \ldots, 10\}$ or $G \in \mathcal{H}$. Finally, by Theorem 3.6, we can assume that either $G \in \mathcal{G}_i^*$ or $G \in \mathcal{H}^*$. This is because G is vertex-critical. Since $\chi(G^*) = \chi(G)$, we conclude that $G^* \cong G$. For each Q_i , we write x_i, x'_i, x''_i , and x'''_i to denote arbitrary distinct vertices in Q_i (provided that such vertices exist). For independent sets I_1 and I_2 and $G - (I_1 \cup I_2)$, we write $(Q'_{j_1}, Q'_{j_2}, Q'_{j_3}, \ldots)$ to denote the vertex order $(Q_{j_1}, Q_{j_2}, Q_{j_3}, \ldots)$ restricted to $V(G) \setminus (I_1 \cup I_2)$. Now we consider these 11 cases in succession. Each case is independent of all others.

Case 1: $G \in \mathcal{G}_1^*$. Since $\Delta(G) = 9$, there exists $i \in \{1, \ldots, 5\}$ such that $|Q_i| \ge 4$. By symmetry, assume i = 1. If $|Q_5| \ge 2$ and $|Q_2| \ge 2$, then $G[\{x_1, x'_1, x''_1, x''_1, x_2, x'_2, x_5, x'_5\}] \cong K_4 \lor H$, where H contains two disjoint pairs of nonadjacent vertices; this subgraph is d_1 -choosable, which contradicts Lemma 3.3. Assume instead, by symmetry, that $|Q_5| = 1$. Lemma 3.2 implies that $1 \ge d(x_2) - d(x_1) = |N[x_2]| - |N[x_1]| = |Q_3| - |Q_5|$. Thus, $|Q_3| \le 2$. Since $8 \le d(x_4) =$ $|Q_3| + |Q_4| - 1 + |Q_5|$, we know $|Q_4| \ge 6$. But now $d(x_5) = |Q_4| + (|Q_5| - 1) + |Q_1| \ge 6 + (1 - 1) + 4 = 10$. This contradicts that $\Delta(G) = 9$.

Case 2: $G \in \mathcal{G}_2^*$. By the Copycat Lemma, $|Q_2| > |Q_6|$ and $|Q_5| > |Q_6|$; in particular, $|Q_2|, |Q_5| \ge 2$ since each Q_i is nonempty. If $|Q_6| \ge 2$, then let $I_1 = \{x_2, x_5, x_6\}$ and $I_2 = \{x'_2, x'_5, x'_6\}$. Now $G - (I_1 \cup I_2)$ is 5-degenerate with vertex order $(Q'_1, Q'_4, Q'_3, Q'_5, Q'_2, Q'_6)$; by Lemma 3.1, G is 8-colorable, a contradiction. Assume instead that $|Q_6| = 1$. If $|Q_1| \ge 2$, then let $I_1 = \{x_2, x_5, x_6\}$. Now $G - I_1$ is 6-degenerate with vertex order $(Q'_1, Q'_5, Q'_2, Q'_4, Q'_3, Q'_6)$; by Lemma 3.1, G is 8colorable, a contradiction. Assume instead that $|Q_1| = 1$. Now Lemma 3.2 implies that $1 \ge d(x_3) - d(x_2) = |N[x_3]| - |N[x_2]| = |Q_4| + |Q_6| - |Q_1| = |Q_4|$. By symmetry, $|Q_3| \le 1$. Thus, $d(x_6) = |Q_1| + |Q_3| + |Q_4| + |Q_6| - 1 = 3$, which contradicts that $\delta(G) \ge 8$.

Case 3: $G \in \mathcal{G}_3^*$. By the Copycat Lemma, $|Q_5| > |Q_6|$ and $|Q_3| > |Q_7|$; in particular,



Figure 3: Cases 1–3: $G \in \mathcal{G}_1^*, G \in G_2^*$, or $G \in G_3^*$.

 $|Q_5|, |Q_3| \ge 2$, since each Q_i is nonempty. If $|Q_4| \ge 2$, then let $I_1 = \{x_2, x_5, x_6\}$ and $I_2 = \{x_1, x_3, x_7\}$. Now $G - (I_1 \cup I_2)$ is 5-degenerate with vertex order $(Q'_4, Q'_3, Q'_5, Q'_2, Q'_1, Q'_6, Q'_7)$; by Lemma 3.1, G is 8-colorable, a contradiction. Assume instead that $|Q_4| = 1$. Lemma 3.2 implies that $1 \ge d(x_2) - d(x_3) = |N[x_2]| - |N[x_3]| = |Q_1| + |Q_7| - |Q_4| = |Q_1| + |Q_7| - 1$. Thus, $|Q_1| = |Q_7| = 1$. By symmetry, $|Q_2| = |Q_6| = 1$. So $d(x_6) = |Q_1| + |Q_7| + |Q_4| + |Q_6| - 1 = 3$, which contradicts that $\delta(G) \ge 8$.

Case 4: $G \in \mathcal{G}_4^*$. By the Copycat Lemma, $|Q_1| > |Q_5|$; so $|Q_1| \ge 2$ since each Q_i is nonempty. Let $I_1 := \{x_1, x_5, x_7\}$ and $I_2 := \{x'_1, x_3, x_6\}$. Now $G - (I_1 \cup I_2)$ is 5-degenerate with vertex order $(Q'_4, Q'_2, Q'_1, Q'_5, Q'_3, Q'_7, Q'_6)$; by Lemma 3.1, G is 8-colorable, a contradiction.



Figure 4: Cases 4–6: $G \in \mathcal{G}_4^*, G \in \mathcal{G}_5^*$, or $G \in \mathcal{G}_6^*$.

Case 5: $G \in \mathcal{G}_5^*$. By the Copycat Lemma, $|Q_5| > |Q_6|$ and $|Q_2| > |Q_7|$; so $|Q_5|, |Q_2| \ge 2$ since each Q_i is nonempty. If $|Q_1| \ge 3$, then $G[\{x_1, x'_1, x''_1, x_2, x'_2, x_5, x'_5, x_6, x_7\}] \cong K_3 \vee 3K_2$; this subgraph is d_1 -choosable, contradicting Lemma 3.3. Assume instead that $|Q_1| \le 2$. If $|Q_3| \ge 4$, then $G[\{x_3, x'_3, x''_3, x''_3, x_2, x_4, x_7, x_8\}] \cong K_4 \vee H$ where H contains two disjoint pairs of nonadjacent vertices; this subgraph is d_1 -choosable, contradicting Lemma 3.3. So $|Q_3| \le 3$. By symmetry, $|Q_4| \le 3$. Thus, $|Q_2|, |Q_5| \ge 4$, since $d(x_2), d(x_5) \ge 8$. However, now $d(x_1) = (|Q_1| - 1) + |Q_2| +$ $|Q_5| + |Q_6| + |Q_7| \ge (1 - 1) + 4 + 4 + 1 + 1 = 10$, which contradicts that $\Delta(G) = 9$.

Case 6: $G \in \mathcal{G}_{6}^{*}$. Let $I_1 := \{x_3, x_5, x_7\}$ and $I_2 := \{x_2, x_6, x_8\}$. Now $G - (I_1 \cup I_2)$ is 5-degenerate with vertex order $(Q'_4, Q'_1, Q'_8, Q'_5, Q'_7, Q'_6, Q'_3, Q'_2)$. By Lemma 3.1, G is 8-colorable, a contradiction.



Figure 5: Cases 7–8: $G \in \mathcal{G}_7^*$ or $G \in \mathcal{G}_8^*$.

Case 7: $G \in \mathcal{G}_7^*$. By the Copycat Lemma, $|Q_4| > |Q_7|$; so $|Q_4| \ge 2$ since each Q_i is nonempty. If $|Q_1| \ge 7$, then $d(x_2) = |Q_3| + (|Q_2| - 1) + |Q_6| + |Q_5| + |Q_1| \ge 10$, contradicting that $\Delta(G) = 9$. Thus, $|Q_1| \le 6$. Let $I_1 := \{x_4, x_6, x_7\}$ and $I_2 := \{x_2, x'_4, x_8\}$. Now $G - (I_1 \cup I_2)$ is 5-degenerate with vertex order $(Q'_5, Q'_3, Q'_4, Q'_7, Q'_2, Q'_8, Q'_6, Q'_1)$. By Lemma 3.1, G is 8-colorable, a contradiction.

Case 8: $G \in \mathcal{G}_8^*$. Let $I_1 := \{x_3, x_5, x_7\}$ and $I_2 := \{x_2, x_6, x_8\}$. Now $G - (I_1 \cup I_2)$ is 5-degenerate with order $(Q'_4, Q'_1, Q'_3, Q'_2, Q'_7, Q'_8, Q'_6, Q'_5)$; by Lemma 3.1, G is 8-colorable, a contradiction.

Case 9: $G \in \mathcal{G}_{9}^{\bullet}$. By the Copycat Lemma $|Q_{9}| > |Q_{5}|$; so $|Q_{9}| \ge 2$ since each Q_{i} is nonempty. Let $I_{1} := \{x_{3}, x_{5}, x_{7}, x_{9}\}$ and $I_{2} := \{x_{2}, x_{6}, x_{8}, x'_{9}\}$. Now $G - (I_{1} \cup I_{2})$ is 5-degenerate with order $(Q'_{4}, Q'_{1}, Q'_{3}, Q'_{2}, Q'_{7}, Q'_{8}, Q'_{9}, Q'_{6}, Q'_{5})$. By Lemma 3.1, G is 8-colorable, a contradiction.

Case 10: $G \in \mathcal{G}_{10}^*$. Let $I_1 := \{x_3, x_5, x_7\}$ and $I_2 := \{x_2, x_6, x_8\}$. Now $G - (I_1 \cup I_2)$ is 5-degenerate with order $(Q'_9, Q'_4, Q'_1, Q'_3, Q'_2, Q'_7, Q'_8, Q'_6, Q'_5)$. By Lemma 3.1, G is 8-colorable, a contradiction.



Figure 6: Cases 9–11: $G \in \mathcal{G}_9^*$, $G \in G_{10}^*$, or $G \in \mathcal{H}^*$.

Case 11: $G \in \mathcal{H}^*$. Let \mathcal{H} be the class of connected (P_5, gem) -free graphs G for which V(G) can be partitioned into non-empty sets A_1, \ldots, A_7 such that the following properties all hold: each A_i induces a P_4 -free graph, the vertex-set of each component of $G[A_7]$ is a homogeneous set, each edge with exactly one endpoint in $V(A_7)$ has the other endpoint in $V(A_6)$, and for all distinct $i, j \in \{1, \ldots, 6\}$, if a solid edge appears between A_i and A_j in the rightmost graph in Figure 6, then $[A_i, A_j]$ is complete, but if no edge appears then $[A_i, A_j] = \emptyset$. Let \mathcal{H}^* be the class of graphs that are in \mathcal{H} with A_1, \ldots, A_5 being cliques and also each component of A_7 being a clique. By Theorem 3.6, we know that if $G \in \mathcal{H}$, then in fact $G \in \mathcal{H}^*$.

If $|A_5| \leq \omega(A_6)$, then by criticality $G - A_5$ has an 8-coloring, call it φ . To extend φ to G, we color A_5 with colors used on a maximum clique in A_6 . This yields a proper 8-coloring of G, a contradiction. Thus, $|A_5| > \omega(A_6)$. By symmetry, $|A_2| > \omega(A_6)$. Since A_6 is nonempty by definition, $|A_2|, |A_5| \geq 2$. Since $d(a_6) \leq 9$ and each of A_1, A_3 , and A_4 is nonempty, each component of A_7 has size at most 6. If $|A_6| \geq 2$, then let $I_1 = \{a_2, a_5, a_6\}$ and $I_2 = \{a'_2, a'_5, a'_6\}$. Now $G - (I_1 \cup I_2)$ is 5-degenerate with order $A'_1, A'_5, A'_2, A'_4, A'_3, A'_6, A'_7$. By Lemma 3.1, G is 8-colorable, a contradiction. So assume instead that $|A_6| = 1$.

By criticality, there exists an 8-coloring φ of $G' := G - V(A_7)$. Each component of A_7 is a clique of size at most 6 that is adjacent to only a_6 in G'. Thus we can color each component of A_7 with 8 colors so that φ extends to G, which contradicts that $\chi(G) \ge \Delta(G) = 9$.

References

 O.V. Borodin and A.V. Kostochka. On an upper bound of a graph's chromatic number, depending on the graph's degree and density. *Journal of Combinatorial Theory Series B*, 23, no.2-3:247-250, 1977.

- [2] R.L. Brooks. On colouring the nodes of a network. Mathematical Proceedings of the Cambridge Philosophical Society, vol. 37, Cambridge Univ Press, pp. 194-197, 1941.
- [3] P.A. Catlin. Embedding subgraphs and coloring graphs under extremal degree conditions. ProQuest LLC, Ann Arbor, MI, Thesis (Ph.D.)-The Ohio State University, 1976.
- [4] M. Chudnovsky, T. Karthick, P. Maceli, and F. Maffray. Coloring graphs with no induced five-vertex path or gem. https://arxiv.org/abs/1810.06186, 2019.
- [5] M. Chudnovsky, N. Robertson, P. Seymour, R. Thomas. The strong perfect graph theorem. Annals of Mathematics Second Series, vol. 164, no. 1, 51-229, 2006.
- [6] D.W. Cranston and L. Rabern. Coloring claw-free graphs with Δ 1 colors. SIAM Journal of Discrete Math., vol. 27, no.1:534-549, 2013.
- [7] D.W. Cranston and L. Rabern. Coloring a graph with $\Delta 1$ colors: Conjectures equivalent to the Borodin–Kostochka Conjecture that appear weaker. *European Journal of Combinatorics*, 44 Part A:23-42, 2015.
- [8] T. Karthick and F. Maffray. Coloring (gem, co-gem)-free graphs. Journal of Graph Theory, vol. 89, no.3:288-303, 2018.
- [9] A.D. King, Hitting all maximum cliques with a stable set using lopsided independent transversals. *Journal of Graph Theory* vol. 67 no.4:300-305, 2011.
- [10] A.V. Kostochka. Degree, density, and chromatic number. Metody Diskret. Anal., no. 35:45-70, 1980, (in Russian).
- [11] B. Reed. A strengthening of Brooks' theorem. Journal of Combinatorial Theory Series B, 76, no.2:136-149, 1999.
- [12] B. Reed. ω , Δ , and χ . Journal of Graph Theory, 27, no. 4:177-212, 1998.
- [13] D.B. West. Introduction to graph theory. Prentice Hall, Inc., Upper Saddle River, NJ, 1996. xvi+512 pp. ISBN: 0-13-227828-6.