Curved fronts of bistable reaction-diffusion equations in spatially periodic media *

Hongjun Guo ^a, Wan-Tong Li ^b, Rongsong Liu ^a and Zhi-Cheng Wang ^b

Abstract

In this paper, curved fronts are constructed for spatially periodic bistable reaction-diffusion equations under the a priori assumption that there exist pulsating fronts in every direction. Some sufficient and some necessary conditions of the existence of curved fronts are given. Furthermore, the curved front is proved to be unique and stable. Finally, a curved front with varying interfaces is also constructed. Despite the effect of the spatial heterogeneity, the result shows the existence of curved fronts for spatially periodic bistable reaction-diffusion equations which is known for the homogeneous case.

Keywords. Pulsating front; Curved front; Spatially periodic reaction-diffusion equation; Uniqueness; Stability.

1 Introduction

In this paper, we consider spatially periodic reaction-diffusion equations of the type

$$u_t - \Delta u = f(x, y, u), \quad (t, x, y) \in \mathbb{R} \times \mathbb{R}^2,$$
 (1.1)

where $u_t = \frac{\partial u}{\partial t}$ and $\Delta = \partial_{xx} + \partial_{yy}$ denotes the Laplace operator with respect to the space variables $(x, y) \in \mathbb{R}^2$. The reaction term f(x, y, y, u) is assumed to be periodic in (x, y) and bistable in u. More precisely, we assume throughout this paper that

(F1) f(x, y, u) is continuous, of class C^{α} in (x, y) uniformly in $u \in [0, 1]$ with $\alpha \in (0, 1)$, of class C^{2} in u uniformly in $(x, y) \in \mathbb{R}^{2}$ with $f_{u}(x, y, u)$ and $f_{uu}(x, y, u)$ being Lipschitz continuous in $u \in \mathbb{R}$;

^a Department of Mathematics and Statistics, University of Wyoming, Laramie, USA

^b School of Mathematics and Statistics, Lanzhou University, Lanzhou, China

^{*}Guo and Liu are partially supported by by NSF grant 1826801. Li is supported by the National Natural Science Foundation of China under grants 11731005 and 11671180. Wang is supported by the National Natural Science Foundation of China under grant 11371179.

- (F2) f(x, y, u) is L-periodic with respect to (x, y) where $L = (L_1, L_2) \in \mathbb{R}^2$, that is, $f(x+k_1L_1, y+k_2L_2, u) = f(x, y, u)$ for any $k_1, k_2 \in \mathbb{Z}$;
- (F3) for every $(x, y) \in \mathbb{R}^2$, 0 and 1 are stable zeroes of $f(x, y, \cdot)$, that is,

$$f(x, y, 0) = f(x, y, 1) = 0,$$

and there exist $\lambda > 0$ and $\sigma \in (0, 1/2)$ such that

$$-f_u(x, y, u) \ge \lambda$$
 for all $(x, y, u) \in \mathbb{R}^2 \times [0, \sigma]$ and $(x, y, u) \in \mathbb{R}^2 \times [1 - \sigma, 1]$.

A typical example of f(x, y, u) is the cubic nonlinearity

$$f(x, y, u) = u(1 - u)(u - \theta_{x,y}),$$

where $\theta_{x,y} \in (0,1)$ is a *L*-periodic function. The equation (1.1) is a special generalization of the famous Allen-Cahn equation [1]. For mathematical convenience, we extend f(x,y,u) out of the interval $u \in [0,1]$ such that

$$-f_u(x, y, u) \ge \lambda \text{ for all } (x, y, u) \in \mathbb{R}^2 \times (-\infty, \sigma] \text{ and } (x, y, u) \in \mathbb{R}^2 \times [1 - \sigma, +\infty). \tag{1.2}$$

Then, f(x, y, u) is globally Lipschitz continuous in $u \in \mathbb{R}$.

Before going further on, we first recall some well-known results in the homogeneous case, that is,

$$u_t - \Delta u = f(u), \ (t, x) \in \mathbb{R} \times \mathbb{R}^N, \tag{1.3}$$

where f is of bistable type, that is, $f(0) = f(1) = f(\theta)$, f < 0 on $(0, \theta)$ and f > 0 on $(\theta, 1)$, for some $\theta \in (0, 1)$. For one-dimensional space, it follows from celebrated results due to Fife and McLeod [12] that (1.3) admits a unique (up to shifts) traveling front $\phi(x - c_f t)$ satisfying

$$0 < \phi < 1, \ \phi(-\infty) = 1 \text{ and } \phi(+\infty) = 0.$$

Moreover, the speed c_f has the sign of $\int_0^1 f(u)du$ and the front is globally and exponentially stable. A trivial extension of the traveling front to higher dimensional spaces is the planar front $\phi(x \cdot e - c_f t)$ where $e \in \mathbb{S}^{N-1}$ denotes the propagation direction. Notice that every level set of a planar front is a plane. Except planar fronts, more types of fronts are also known to exist in high dimensional spaces, such as V-shaped fronts, conical shaped fronts and pyramidal fronts, see Hamel et. al. [18], Ninomiya and Taniguchi [20] and Taniguchi [23, 24]. All these fronts are transition fronts connecting 0 and 1 defined by Hamel [16]. For above fronts, their interfaces between 0 and 1 can be given by their level sets and different shapes of interfaces actually show some structures of the solutions. One can roughly imagine a global appearance of such solutions in the framework of transition fronts by noticing that the solutions are approaching to 1 and 0 on one side and the other of the interfaces, respectively.

As far as a spatially periodic bistable reaction-diffusion equation considered, the situation is more complicated than the homogenous case. Because of the effect of hetereogeneities, there may even not exist transition fronts connecting states 0 and 1, see Zlatoš [32]. However, what we are concerned in this paper is the existence of curved fronts when there exist some fronts in every direction, that is, pulsating fronts. We now introduce the notion of pulsating front by referring to [2, 22, 27, 28, 29].

Definition 1.1 Denote a periodic cell by $\mathbb{T}^2 = [0, L_1] \times [0, L_2]$. A pair (U_e, c_e) with $U_e : \mathbb{R} \times \mathbb{T}^2 \to \mathbb{R}$ and $c_e \in \mathbb{R}$ is said to be a pulsating front of (1.1) with effective speed c_e in the direction $e \in \mathbb{S}$ connecting 0 and 1 if the two following conditions are satisfied:

(i) For every $\xi \in \mathbb{R}$, the profile $U_{\ell}(\xi, x, y)$ is L-periodic in (x, y) and satisfies

$$\lim_{\xi \to +\infty} U_e(\xi, x, y) = 0, \ \lim_{\xi \to -\infty} U_e(\xi, x, y) = 1, \ \ uniformly \ for \ (x, y) \in \mathbb{T}^2.$$

(ii) The map $u(t, x, y) := U_e((x, y) \cdot e - c_e t, x, y)$ is an entire (classical) solution of the parabolic equation (1.1).

We now recall some existence results of pulsating fronts for the general reaction-diffusion equation in spatially periodic media

$$u_{t} = \sum_{i} (a(x)u_{x_{i}})_{x_{i}} + \sum_{i} b_{i}(x)u_{x_{i}} + f(x, u), \ t \in \mathbb{R}, \ x \in \mathbb{R}^{N}.$$
(1.4)

For one dimensional case of (1.4) when f(x, u) = g(x)f(u), Nolen and Ryzhik [21] proved the existence of pulsating fronts with nonzero speed by provided with some restrictions for g and f. Moreover, Ducrot, Giletti and Matano [8] also got some existence results of pulsating fronts with a positive speed, if the solutions of (1.4) with some compactly supported initial conditions can converge locally uniformly to 1 as $t \to +\infty$. Still for one-dimensional case, Ding, Hamel and Zhao [6] applied the implicit function theorem and abstract results of Fang and Zhao [11] to get the existence of pulsating fronts for small period and large period. For higher dimensions, when the diffusivity matrix a is close to identity and f is independent of f, the existence of pulsating fronts is obtained by Xin [27, 28, 29] through refined perturbation arguments. Ducrot [7] also got some existence results of fronts connecting 0 and 1 in every direction for slowly varying medium and rapidly varying medium (that is, f << 1 and f >> 1 respectively when the reaction term is f (f > 1, f > 1, f > 1, f = 1, f = 1, f > 1, f = 1, f =

In this work, we aim to construct curved fronts by using some pulsating fronts with nonzero speeds. Therefore, we need to assume a priori that

(H1)
$$\int_{\mathbb{T}^2 \times [0,1]} f(x,y,u) dx dy du \neq 0,$$

(H2) for every unit vector $e \in \mathbb{R}^2$, the equation (1.1) admits a pulsating front $U_e((x, y) \cdot e - c_e t, x, y)$ with $c_e \neq 0$.

From the results of Ducrot [7] and Guo [14], one knows that if (H1), (H2) hold, the propagation speed c_e of the pulsating front in every direction has the sign of $\int_{\mathbb{T}^2 \times [0,1]} f(x,y,u) dx dy du$. We assume without loss of generality that

$$\int_{\mathbb{T}^2 \times [0,1]} f(x, y, u) dx dy du > 0, \tag{1.5}$$

which implies $c_e > 0$ for all $e \in \mathbb{S}$. Otherwise, one can replace u, f, $U_e(\xi, x, y)$ by $\tilde{u} = 1 - u$, g(x, y, u) = -f(x, y, 1 - u), $\tilde{U}_e(\xi, x, y) = 1 - U_e(-\xi, x, y)$ and consider the new pulsating front \tilde{U}_e with speed $-c_e$. From [3] and [14], the speed c_e and the profile U_e of the pulsating front are unique up to shifts in time for any direction e. We fix the pulsating front in every direction e by

$$U_e(0,0,0) = \frac{1}{2}.$$

From [14], we also know that $\partial_{\xi}U_e < 0$, the family $\{c_e\}_{e \in \mathbb{S}}$ is uniformly bounded with respect to e and the minimum and maximum of c_e can be reached with the following inequality

$$0 < \min_{e \in \mathbb{S}} c_e \le \max_{e \in \mathbb{S}} c_e < +\infty.$$

In the whole paper, we always assume that (F1)-(F3), (H1)-(H2) and (1.5) hold and we do not repeat it in the sequel. We now focus on construction of curved fronts by some pulsating fronts. To the best of our knowledge, few results of the existence of curved fronts are known for bistable reaction-diffusion in spatially periodic media. However, one can refer to [9, 10] for the existence of curved fronts of monostable and combustion reaction-diffusion equations with a periodic shear flow and refer to [4] for a space-time periodic monostable reaction-advection-diffusion equation. Although the pulsating front $U_e((x,y) \cdot e - c_e t, x, y)$ is not exactly planar, every level set is still bounded with a plane. Thus, the pulsating front is also called almost-planar in the framework of transition fronts, see [16]. We try to apply the ideas of Ninomiya and Taniguchi [20] which they used for homogeneous bistable case, to construct the curved fronts. But, since the profiles U_e and speeds c_e of pulsating fronts are different in general with respect to the direction e, we have to update their ideas.

We then claim our results. Let $\alpha \in (0, \pi)$. Then, by Assumption (H2), there exists a pulsating front in the direction $(\cos \alpha, \sin \alpha)$, that is,

$$U_{\alpha}(x\cos\alpha + y\sin\alpha - c_{\alpha}t, x, y).$$

For any $\alpha, \beta \in (0, \pi)$, define

$$U_{\alpha\beta}^{-}(t,x,y) := \max\{U_{\alpha}(x\cos\alpha + y\sin\alpha - c_{\alpha}t,x,y), U_{\beta}(x\cos\beta + y\sin\beta - c_{\beta}t,x,y)\}, \quad (1.6)$$

which is a subsolution of (1.1). Our first result shows the existence of a curved front which converges to pulsating fronts along its asymptotic lines under some conditions on angles α and β . The curved front is actually a transition front connecting 0 and 1 whose interfaces can be chosen as a V-shaped curve.

Theorem 1.2 For any $\theta \in (0, \pi)$, let $g(\theta) = c_{\theta}/\sin \theta$. For any $0 < \alpha < \beta < \pi$ such that

$$\frac{c_{\alpha}}{\sin \alpha} = \frac{c_{\beta}}{\sin \beta} := c_{\alpha\beta} > \frac{c_{\theta}}{\sin \theta} \text{ for any } \theta \in (\alpha, \beta), \ g'(\alpha) < 0 \text{ and } g'(\beta) > 0, \tag{1.7}$$

there exists an entire solution V(t, x, y) of (1.1) such that $V_t(t, x, y) > 0$ for all $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$ and

$$\lim_{R \to +\infty} \sup_{x^2 + (y - c_{\alpha\beta}t)^2 > R^2} \left| V(t, x, y) - U_{\alpha\beta}^{-}(t, x, y) \right| = 0.$$
 (1.8)

Remark 1.3 It seems that the conditions $g'(\alpha) < 0$ and $g'(\beta) > 0$ can not be removed by our methods. These conditions are actually true for homogeneous unbalanced bistable case with the reaction term having positive integration from 0 to 1 (α has to be smaller than $\pi/2$ in this case by symmetry and $\beta = \pi - \alpha$), but false for homogeneous balanced bistable case. Moreover, the V-shaped front exists in homogeneous unbalanced bistable case, but does not exist in homogeneous balanced bistable case, see [17]. Nevertheless, for the balanced case, there exist some fronts whose level sets have an exponential shape for 2-dimensional space and a paraboloidal shape for N-dimensional space with $N \geq 3$, see [5, 25, 26].

Remark 1.4 One can easily check that the curved front V(t, x, y) in Theorem 1.2 is a transition front connecting 0 and 1 (see [16] for the definition) with sets

$$\Gamma_t := \{x \leq 0, y \in \mathbb{R}; x \cos \alpha + y \sin \alpha - c_{\alpha} t\} \cup \{x > 0, y \in \mathbb{R}; x \cos \beta + y \sin \beta - c_{\beta} t\},\$$

 $\Omega_t^+ := \{x \le 0, y \in \mathbb{R}; x \cos \alpha + y \sin \alpha - c_\alpha t < 0\} \cup \{x > 0, y \in \mathbb{R}; x \cos \beta + y \sin \beta - c_\beta t < 0\},$ and

$$\Omega_t^- := \{x \le 0, y \in \mathbb{R}; x \cos \alpha + y \sin \alpha - c_\alpha t > 0\} \cup \{x > 0, y \in \mathbb{R}; x \cos \beta + y \sin \beta - c_\beta t > 0\}.$$

Notice that for any fixed t, Γ_t is a connected polyline since $c_{\alpha}/\sin \alpha = c_{\beta}/\sin \beta$ and the shape of Γ_t is invariant with respect to t. Moreover, by the definition of the global mean speed [16], the curved front V(t, x, y) has a global mean speed equal to $\min\{c_{\alpha}, c_{\beta}\}$, in the sense that

$$\frac{d(\Gamma_t, \Gamma_s)}{|t-s|} \to \min\{c_\alpha, c_\beta\}, \ as \ |t-s| \to +\infty.$$

Here, the distance d(A, B) between two subsets A and B of \mathbb{R}^2 , is defined by the smallest geodesic distance between pairs of points in A and B. Other definition of the distance \tilde{d} like

$$\tilde{d}(A, B) = \min \Big(\sup \{ d(x, B); x \in A \}, \sup \{ d(y, A); y \in B \} \Big),$$

could be used. Then, there holds that $d(A, B) \leq \tilde{d}(A, B)$ and the global mean speed is equal to $\max\{c_{\alpha}, c_{\beta}\}$, in the sense that

$$\frac{\tilde{d}(\Gamma_t, \Gamma_s)}{|t-s|} \to \max\{c_\alpha, c_\beta\}, \ as \ |t-s| \to +\infty.$$




Figure 1: An example of α and β satisfying (1.7).

This is different with the homogeneous case, in which the global mean speeds under these two definitions are the same, see [16] and see [15] for the underlying domains being exterior domains and domains with multiple branches.

We then show that the condition (1.7) is not empty, that is, it is satisfied when α close to 0 and β close to π , see Figure 1.

Corollary 1.5 There exist $0 < \alpha_1 < \beta_1 < \pi$ such that for any $\alpha \in (0, \alpha_1)$, there is $\beta \in (\beta_1, \pi)$ such that (1.7) holds for such α , β and there exists an entire solution V(t, x, y) of (1.1) satisfying (1.8).

Indeed, one can rotate the coordinate such that *y*-axis points to any direction. Therefore, Corollary 1.5 implies that for any two pulsating fronts whose propagation directions are close to reversed with each other, one can use them to construct a curved front.

Corollary 1.6 There exist $0 < \rho < 1$ such that for any directions e_1 , e_2 with $-1 < e_1 \cdot e_2 < -1 + \rho$, there exist a direction e_0 such that

$$\frac{c_{e_1}}{\sqrt{1 - (e_1 \cdot e_0)^2}} = \frac{c_{e_2}}{\sqrt{1 - (e_2 \cdot e_0)^2}} := c_{e_1 e_2}$$
 (1.9)

and there is an entire solution V(t, x, y) of (1.1) satisfying

$$\lim_{R \to +\infty} \sup_{((x,y)-c_{e_1e_2}te_0)^2 > R^2} \left| V(t,x,y) - U_{e_1e_2}^-(t,x,y) \right| = 0, \tag{1.10}$$

where

$$U^-_{e_1e_2}(t,x,y) := \max\{U_{e_1}((x,y)\cdot e_1 - c_{e_1}t,x,y), U_{e_2}((x,y)\cdot e_2 - c_{e_2}t,x,y)\}.$$

By Theorem 1.2, one knows that (1.7) is a sufficient condition for the existence of V(t, x, y) satisfying (1.8). However, we can not show that (1.7) is necessary, but can show that (1.7) without $g'(\alpha) < 0$ and $g'(\beta) > 0$ is necessary.

Theorem 1.7 If there are two angles α and β of $(0, \pi)$ and a constant $c_{\alpha\beta} > 0$ such that there exists an entire solution V(t, x, y) of (1.1) satisfying (1.8), then it holds

$$c_{\alpha\beta} = \frac{c_{\alpha}}{\sin \alpha} = \frac{c_{\beta}}{\sin \beta} > \frac{c_{\theta}}{\sin \theta} \text{ for any } \theta \in (\alpha, \beta).$$

Now, we show the uniqueness and the stability of the curved front V(t, x, y).

Theorem 1.8 For any fixed $0 < \alpha < \beta < \pi$ satisfying

$$\frac{c_{\alpha}}{\sin \alpha} = \frac{c_{\beta}}{\sin \beta} := c_{\alpha\beta},$$

the entire solution V(t, x, y) of (1.1) satisfying (1.8) is unique, that is, if there is an entire solution $V^*(t, x, y)$ satisfying (1.8), then $V^*(t, x, y) \equiv V(t, x, y)$.

Theorem 1.9 Let α and β be fixed angles satisfying (1.7) and V(t, x, y) be the entire solution of (1.1) satisfying (1.8). Let $0 \le u_0(x, y) \le 1$ be an initial value satisfying

$$\lim_{R \to +\infty} \sup_{x^2 + y^2 > R^2} \left| u_0(x, y) - U_{\alpha\beta}^{-}(0, x, y) \right| = 0.$$
 (1.11)

Then, the solution u(t, x, y) of (1.1) for t > 0 with $u(0, x, y) = u_0(x, y)$ satisfies

$$\lim_{t \to +\infty} ||u(t, x, y) - V(t, x, y)||_{L^{\infty}(\mathbb{R}^2)} = 0.$$

Next, we construct a transition front connecting 0 and 1 with varying interfaces. Such a kind of transition front is known in homogeneous case by [16], in which the solution is orthogonal symmetric with respect to y-axis and behaves as three planar fronts as $t \to -\infty$. However, in our case, this transition front can not be symmetric in general.

Theorem 1.10 Let α and β be fixed angles satisfying (1.7) and let $V_{\alpha\beta}(t, x, y)$ be the entire solution of (1.1) satisfying (1.8). Denote $e_{\alpha} = (\cos \alpha, \sin \alpha)$ and $e_{\beta} = (\cos \beta, \sin \beta)$. Assume that there exist another angle $\theta \in (\alpha, \beta)$ and a direction $e_{\theta} = (\cos \theta, \sin \theta)$ such that

- (i) for e_{α} and e_{θ} , there is a direction $e_{\alpha\theta}$ such that (1.9) holds for $e_1 = e_{\alpha}$, $e_2 = e_{\theta}$ and $e_0 = e_{\alpha\theta}$, it holds $h'(\alpha) < 0$ where $h(s) = c_s/(e_{\alpha\theta} \cdot (\cos s, \sin s))$ for $0 < s < \theta$ and there is an entire solution $V_{\alpha\theta}(t, x, y)$ satisfying (1.10).
- (ii) for e_{β} and e_{θ} , there is a direction $e_{\beta\theta}$ such that (1.9) holds for $e_1 = e_{\beta}$, $e_2 = e_{\theta}$ and $e_0 = e_{\beta\theta}$, it holds $h'(\beta) > 0$ where $h(s) = c_s/(e_{\beta\theta} \cdot (\cos s, \sin s))$ for $\theta < s < \pi$ and $e_0 = e_{\alpha\theta}$ and there is an entire solution $V_{\beta\theta}(t, x, y)$ satisfying (1.10).



Figure 2: Left: interface when t << -1; Right: interface when t >> 1.

Then, there exists an entire solution u(t, x, y) of (1.1) such that

$$u(t,x,y) \to \begin{cases} V_{\alpha\theta}(t,x,y), \text{ uniformly in the half plane } \{(x,y) \in \mathbb{R}^2; x < 0\}, \\ V_{\beta\theta}(t,x,y)\}, \text{ uniformly in the half plane } \{(x,y) \in \mathbb{R}^2; x > 0\}, \end{cases} \text{ as } t \to -\infty. (1.12)$$

and

$$u(t, x) \to V_{\alpha\beta}(t, x, y), \text{ as } t \to +\infty \text{ uniformly in } \mathbb{R}^2.$$
 (1.13)

The convergence in above theorem is in the sense of L^{∞} norm.

Remark 1.11 From the proof of Theorem 1.10, one can easily check that the entire solution u(t, x, y) is a transition front connecting 0 and 1 with the interfaces

$$\Gamma_{t} := \left\{ x \leq \frac{c_{\alpha} \sin \theta - c_{\theta} \sin \alpha}{\sin(\theta - \alpha)} t, y \in \mathbb{R}; x \cos \alpha + y \sin \alpha - c_{\alpha} t = 0 \right\} \cup \left\{ \frac{c_{\alpha} \sin \theta - c_{\theta} \sin \alpha}{\sin(\theta - \alpha)} t < x \right\}$$

$$\leq \frac{c_{\beta} \sin \theta - c_{\theta} \sin \beta}{\sin(\theta - \beta)} t, y \in \mathbb{R}; x \cos \theta + y \sin \theta - c_{\theta} t = 0 \right\} \cup \left\{ x > \frac{c_{\beta} \sin \theta - c_{\theta} \sin \beta}{\sin(\theta - \beta)} t, y \in \mathbb{R}; x \cos \beta + y \sin \beta - c_{\beta} t = 0 \right\}, \quad \text{for } t \leq 0,$$

and

$$\Gamma_t := \{x \le 0, y \in \mathbb{R}; x \cos \alpha + y \sin \alpha - c_{\alpha}t = 0\} \cup \{x > 0, y \in \mathbb{R}; x \cos \beta + y \sin \beta - c_{\beta}t = 0\}, \text{ for } t > 0,$$
see Figure 2.

Finally, we give an example showing that Theorem 1.10 is not empty.

Corollary 1.12 Assume that e_* is the direction such that the family of speeds $\{c_e\}_{e\in\mathbb{S}}$ reaches its minimal, that is, $c_{e_*} = \min_{e\in\mathbb{S}}\{c_e\}$. Then, there exist e_1 and e_2 close to e_* such that (1.9) holds for $e_0 = e_*$ and there is an entire solution $V_{e_1e_2}(t, x, y)$ of (1.1) satisfying (1.10). Moreover, there

exist a direction e_3 close to $-e_*$ and a direction e_{**} such that there is an entire solution u(t, x, y) of (1.1) such that

$$u(t,x,y) \rightarrow \begin{cases} V_{e_1e_2}(t,x,y), & \text{uniformly in the half plane } \{(x,y) \in \mathbb{R}^2; (x,y) \cdot e_{**} < 0\}, \\ V_{e_2e_3}(t,x,y)\}, & \text{uniformly in the half plane } \{(x,y) \in \mathbb{R}^2; (x,y) \cdot e_{**} > 0\}, \end{cases}$$

as
$$t \to -\infty$$
 and

$$u(t,x) \to V_{e_1e_3}(t,x,y)$$
, as $t \to +\infty$ uniformly in \mathbb{R}^2 .

We organize this paper as following. In Section 2, we first prove the existence of the curved front, that is, Theorem 1.2. Then, we give some examples showing that Theorem 1.2 is not empty. We also show a necessary condition for the existence of the curved front in this section. Section 3 is devoted to the proof of the uniqueness and stability of the curved front in Theorem 1.2. In Section 4, we construct a curved front with varying interfaces and give an example.

2 Existence of curved fronts

This section is devoted to the construction of a curved front satisfying Theorem 1.2. We will need some properties of the pulsating front, especially the differentiability of the profile U_e and the speed c_e with respect to the direction e.

2.1 Preliminaries

We will use the hyperbolic function sech(x) frequently in the sequel. Thus, we recall some known properties of it which can be checked easily.

Lemma 2.1 It holds that

$$|\operatorname{sech}'(x)|, |\operatorname{sech}''(x)| \leq \operatorname{sech}(x), \text{ for } x \in \mathbb{R},$$

and there is a positive constant p such that

$$\operatorname{sech}'(x) > 0$$
 for $x \le -p$, $\operatorname{sech}'(x) < 0$ for $x \ge p$ and $\operatorname{sech}''(x) > 0$ for $|x| \ge p$.

Then, we need a smooth V-shaped curve with $y = -x \cot \alpha$ and $y = -x \cot \beta$ being its asymptotic lines.

Lemma 2.2 For any $0 < \alpha < \beta < \pi$, there is a smooth function $\psi(x)$ for $x \in \mathbb{R}$ with $y = -x \cot \alpha$ and $y = -x \cot \beta$ being its asymptotic lines and there are positive constants k_1 , k_2 and K_1 such that

$$\begin{cases} \psi''(x) > 0, & for \ all \ x \in \mathbb{R} \\ -\cot \alpha < \psi'(x) < -\cot \beta, & for \ all \ x \in \mathbb{R} \\ k_1 \mathrm{sech}(x) \leq \psi'(x) + \cot \alpha \leq K_1 \mathrm{sech}(x), & for \ x < 0, \\ k_2 \mathrm{sech}(x) \leq -\cot \beta - \psi'(x) \leq K_1 \mathrm{sech}(x), & for \ x \geq 0, \\ \max(|\psi''(x)|, |\psi'''(x)|) \leq K_1 \mathrm{sech}(x), & for \ all \ x \in \mathbb{R}. \end{cases}$$

$$(2.1)$$




Figure 3: The function $\varphi(x)$.

Proof. Let $0 < \alpha < \beta < \pi$. Since $\alpha < \beta$, there are two positive constants a, b and a smooth function $\varphi(x)$ such that

$$\varphi(x) = \begin{cases} -x \cot \alpha, & x \le -a \\ -x \cot \beta, & x \ge b. \end{cases} \text{ and } \varphi''(x) > 0 \text{ for } -a < x < b.$$

An example of such a function is that one can take an incircle of the straight lines $y = -x \cot \alpha$ and $y = -x \cot \beta$ with tangent points $(-a, a \cot \alpha)$ and $(b, -b \cot \beta)$ and $\varphi(x)$ is made of the line $y = -x \cot \alpha$ for $x \le -a$, the arc of the incircle between -a and b, and the line $y = -x \cot \beta$ for $x \ge b$. One can mollify $\varphi(x)$ at $(-a, a \cot \alpha)$ and $(b, -b \cot \beta)$ such that $\varphi(x) \in C^{\infty}(\mathbb{R})$, see Figure 3. Define a smooth function $\psi(x)$ as following

$$\psi(x) := \varphi(x) + \rho \operatorname{sech}(x),$$

where $\rho > 0$ is a constant. Since sech"(x) is bounded and by Lemma 2.1, one can make ρ small enough and a, b sufficiently large such that

$$\psi''(x) > 0$$
 for all $x \in \mathbb{R}$.

Moreover, one can easily check that $\psi(x)$ satisfies all properties in (2.1). This completes the proof.

We now recall some properties of the pulsating front $U_e((x, y) \cdot e - c_e t, x, y)$. One can substitute the form $U_e((x, y) \cdot e - c_e t, x, y)$ into (1.1) and get that $(U_e(\xi, x, y), c_e)$ satisfies the semi-linear elliptic degenerate equation

$$c_e \partial_{\xi} U_e + \partial_{\xi\xi} U_e + 2\nabla_{x,y} \partial_{\xi} U_e \cdot e + \Delta_{x,y} U_e + f(x,y,U_e) = 0, \text{ for all } (\xi,x,y) \in \mathbb{R} \times \mathbb{T}^2.$$
 (2.2)

From [14, Lemma 2.1], we have the following lemma.

Lemma 2.3 For any pulsating front $(U_e(\xi, x, y), c_e)$ with $c_e > 0$, there exist $\mu_1 > 0$, $\mu_2 > 0$, $C_1 > 0$ and $C_2 > 0$ independent of e such that

$$0 < U_e(\xi, x, y) \le C_1 e^{-\mu_1 \xi} \text{ for } \xi > 0, \ (x, y) \in \mathbb{T}^2$$
$$0 < 1 - U_e(\xi, x, y) \le C_2 e^{\mu_2 \xi} \text{ for } \xi \le 0, \ (x, y) \in \mathbb{T}^2.$$

Then, by standard parabolic estimates applied to $u(t, x, y) = U_e((x, y) \cdot e - c_e t, x, y)$, one can get that $|\nabla_{x,y}u_t|$, $|u_{tt}|$, $|u_{tt}| \le Cu(t+1, x, y)$ for some constant C > 0 and $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$. Notice that $u_t(t, x, y) = -c_e \partial_{\xi} U_e((x, y) \cdot e - c_e t, x, y)$ with $c_e > 0$. Then, by Lemma 2.3, we have the following lemma.

Lemma 2.4 For any pulsating front $(U_e(\xi, x, y), c_e)$ with $c_e > 0$, there exist $\mu_3 > 0$ and $C_3 > 0$ independent of e such that

$$|\partial_{\xi}U_{e}(\xi,x,y)|, \ |\partial_{\xi\xi}U_{e}(\xi,x,y)|, \ |\nabla_{x,y}\partial_{\xi}U_{e}(\xi,x,y)| \leq C_{3}e^{-\mu_{3}|\xi|} \ \text{for } \xi \in \mathbb{R}, \ (x,y) \in \mathbb{T}^{2}.$$

We also need the following properties.

Lemma 2.5 For any C > 0, there is $0 < \delta < 1/2$ independent of e such that

$$\delta \le U_e(\xi, x, y) \le 1 - \delta, \text{ for } -C \le \xi \le C \text{ and } (x, y) \in \mathbb{T}^2, \tag{2.3}$$

and there is r > 0 independent of e such that

$$-\partial_{\xi} U_{e}(\xi, x, y) \ge r \text{ for for } -C \le \xi \le C \text{ and } (x, y) \in \mathbb{T}^{2}.$$
 (2.4)

Proof. Let $u(t, x, y) = U_e((x, y) \cdot e - c_e t, x, y)$. One can easily check that u(t, x, y) is a transition front connecting 0 and 1 with set $\{(t, x, y) \in \mathbb{R} \times \mathbb{R}^2; (x, y) \cdot e - c_e t = 0\}$ being its interfaces. Then, by [3, Theorem 1.2], one immediately has that there is $0 < \delta < 1/2$ such that

$$\delta < u(t, x, y) < 1 - \delta$$
, for $-C < (x, y) \cdot e - c_{\alpha}t < C$.

By continuity of U_e with respect to e, one has that δ can be independent of e.

The following proof for (2.4) can be simplified for the pulsating front U_e . However, we do it in a general way in purpose that such idea can be used to prove that the curved front which we construct later has similar properties. Notice that $u_t(t, x, y) > 0$ satisfies

$$(u_t)_t - \Delta u_t - f_u(x, y, u)u_t = 0$$
, for $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$.

Assume that there is a sequence $\{(t_n, x_n, y_n)\}_{n \in \mathbb{N}}$ of $\mathbb{R} \times \mathbb{R}^2$ such that $-C \leq (x_n, y_n) \cdot e - c_e t_n \leq C$ and $u_t(t_n, x_n, y_n) \to 0$ as $n \to +\infty$. Since f(x, y, u) is periodic in (x, y), there is $(x', y') \in \mathbb{R}^2$ such that $f(x + x_n, y + y_n, u) \to f(x + x', y + y', u)$ as $n \to +\infty$. Let $u_n(t, x, y) = u(t + t_n, x + x_n, y + y_n)$

and $v_n(t, x, y) = u_t(t + t_n, x + x_n, y + y_n)$. By standard parabolic estimates, $u_n(t, x, y)$ converges to a solution $u_{\infty}(t, x, y)$ of

$$u_t - \Delta u - f(x + x', y + y', u_\infty) = 0$$
, for $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$,

and $v_n(t, x, y)$ converges to a solution $v_{\infty}(t, x, y)$ of

$$v_t - \Delta v - f_u(x + x', y + y', u_\infty)v = 0$$
, for $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$.

Moreover, $v_{\infty}(t, x, y)$ satisfies $v_{\infty}(t, x, y) \ge 0$ and $v_{\infty}(0, 0, 0) = 0$. By the maximum principle, $v_{\infty}(t, x, y) \equiv 0$. Since $U_{e}(\xi, x, y) \to 1$ as $\xi \to -\infty$, there is R > 0 large enough such that

$$u(t, x, y) \ge 1 - \sigma$$
 for $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$ such that $(x, y) \cdot e - c_e t \le -R$

where σ is defined in (F3). Take $(x_*, y_*) \in \mathbb{R}^2$ such that $(x_*, y_*) \cdot e < -R - C$. Then, $v_\infty(t, x, y) \equiv 0$ implies that $u_t(t + t_n, x + x_* + x_n, y + y_* + y_n) \to 0$ as $n \to +\infty$ locally uniformly in $\mathbb{R} \times \mathbb{R}^2$. Notice that $(x_* + x_n, y_* + y_n) \cdot e - c_e t_n \leq -R$ and hence, $u(t_n, x_* + x_n, y_* + y_n) \geq 1 - \sigma$. Also notice that 1 is the only equilibrium of (1.1) over $1 - \sigma$ from (F3) and (1.2). It further implies that $u(t + t_n, x + x_* + x_n, y + y_* + y_n) \to 1$ locally uniformly in $\mathbb{R} \times \mathbb{R}^2$. Since (x_*, y_*) is fixed and $-C \leq (x_n, y_n) \cdot e - c_e t_n \leq C$, it reaches a contradiction with (2.3). This completes the proof. \Box

It follows from [14, Theorem 1.5] that U_e and c_e are differentiable with respect to e. Remember that U_e are normalized by $U_e(0,0,0)=1/2$ for all $e \in \mathbb{S}$. For any $b \in \mathbb{R}^2 \setminus \{0\}$, define

$$U_b = U_{\frac{b}{|b|}} \text{ and } c_b = c_{\frac{b}{|b|}}.$$
 (2.5)

Lemma 2.6 Let U_b and c_b be defined in (2.5). Then, U_b and c_b are doubly continuously Fréchet differentiable at any $b \in \mathbb{R}^N \setminus \{0\}$.

Let us denote the Fréchet derivatives up to second order of U_e and c_e with respect to e by U'_e , U''_e , c'_e and c''_e . From [14], one knows that U_e is continuous with respect to e in L^{∞} space. Since U_e is uniformly bounded for $e \in \mathbb{S}$, the Fréchet derivatives are all bounded in the sense that

$$||U_e'|| = \sup_{h \in \mathbb{R}^N} \frac{||U_e' \cdot h||_{L^{\infty}(\mathbb{R} \times \mathbb{T}^N)}}{|h|} < +\infty, \ ||U_e''|| = \sup_{(h,\rho) \in \mathbb{R}^N \times \mathbb{R}^N} \frac{||(U_e'' \cdot h) \cdot \rho||_{L^{\infty}(\mathbb{R} \times \mathbb{T}^N)}}{|h||\rho|} < +\infty,$$

and

$$||c'_e|| = \sup_{h \in \mathbb{R}^N} \frac{|c'_e \cdot h|}{|h|} < +\infty, \ ||c''_e|| = \sup_{(h,\rho) \in \mathbb{R}^N \times \mathbb{R}^N} \frac{|(c''_e \cdot h) \cdot \rho|}{|h||\rho|} < +\infty.$$

We also know from [14] that for any $h \in \mathbb{R}^2$, $\rho \in \mathbb{R}^2$, $U'_e \cdot h$ and $(U''_e \cdot h) \cdot \rho$ are differentiable with respect to ξ , x and y up to second order and these derivatives are bounded too. We then need the following properties of U'_e .

Lemma 2.7 For any $e \in \mathbb{S}$, there exist $\mu_4 > 0$ and $C_4 > 0$ independent of e such that

$$|(U'_e \cdot h)(\xi, x, y)|, |(\partial_{\xi} U'_e \cdot h)(\xi, x, y)| \le C_4 e^{-\mu_4 |\xi|} |h|, \text{ for any } h \in \mathbb{R}^2, \xi \in \mathbb{R} \text{ and } (x, y) \in \mathbb{T}^2.$$

Proof. Take a smooth nonincreasing function $p(\xi)$ such that

$$p(\xi) = 1$$
 for $\xi \le 0$ and $p(\xi) = e^{-r\xi}$ for $\xi \ge b$,

for some positive constants r and b. Here, one can make r and b to be small and large enough respectively such that

$$r \le \min\{\mu_1, \mu_2\},\tag{2.6}$$

and

$$c_e \left| \frac{p'(\xi)}{p(\xi)} \right| + \left| \frac{p''(\xi)}{p(\xi)} \right| \le \frac{\lambda}{2} \text{ for all } \xi \in \mathbb{R} \text{ and } e \in \mathbb{S},$$
 (2.7)

where $\lambda > 0$ is defined in (F3).

For every direction e, we define a function $V_e(\xi, x, y)$ by

$$V_e(\xi, x, y) := p^{-1}(\xi)U_e(\xi, x, y), \text{ for } \xi \in \mathbb{R} \text{ and } (x, y) \in \mathbb{T}^2.$$

By Lemma 2.3 and (2.6), one has

$$V_e(-\infty, x, y) = 1$$
 and $V_e(+\infty, x, y) = 0$, uniformly for $(x, y) \in \mathbb{T}^2$ and $e \in \mathbb{S}$,

and $V(\xi, x, y) \in L^2(\mathbb{R} \times \mathbb{T}^2)$. Since $U_e(\xi, x, y)$ satisfies (2.2), one can get that $V_e(\xi, x, y)$ satisfies

$$c_{e}\partial_{\xi}V + \partial_{\xi\xi}V_{e} + 2\nabla_{x,y}\partial_{\xi}V_{e} \cdot e + \Delta_{x,y}V_{e} + \frac{2p'}{p}\partial_{\xi}V_{e} + \frac{2p'}{p}\nabla_{x,y}V_{e} \cdot e + \frac{1}{p}f(x,y,pV_{e}) + \left(c_{e}\frac{p'}{p} + \frac{p''}{p}\right)V_{e} = 0, \text{ for } (\xi,x,y) \in \mathbb{R} \times \mathbb{T}^{2}.$$

From (F3) and (2.7), there is C > 0 such that

$$\begin{cases}
\frac{1}{p}f(x,y,pV_e) + \left(c_e \frac{p'}{p} + \frac{p''}{p}\right)V_e \le -\frac{\lambda}{2}V_e, & \text{for } (x,y) \in \mathbb{T}^2 \text{ and } \xi \ge C, \\
\frac{1}{p}f(x,y,pV_e) + \left(c_e \frac{p'}{p} + \frac{p''}{p}\right)V_e \ge \frac{\lambda}{2}(1 - V_e), & \text{for } (x,y) \in \mathbb{T}^2 \text{ and } \xi \le -C,
\end{cases} (2.8)$$

For any $e \in \mathbb{S}$, define a linear operator

$$M_e(v) := c_e \partial_{\xi} v + \partial_{\xi\xi} v + 2\nabla_{x,y} \partial_{\xi} v \cdot e + \Delta_{x,y} v + \frac{2p'}{p} \partial_{\xi} v + \frac{2p'}{p} \nabla_{x,y} v \cdot e - \beta v,$$

where $\beta > 0$ is a fixed real number and

$$v\in D:=\{v\in H^1(\mathbb{R}\times\mathbb{T}^N);\ \partial_{\xi\xi}v+2\nabla_y\partial_\xi v\cdot e+\Delta_yv\in L^2(\mathbb{R}\times\mathbb{T}^N)\},$$

see [14] for definitions of $L^2(\mathbb{R} \times \mathbb{T}^2)$, $H^1(\mathbb{R} \times \mathbb{T}^2)$ and their norms. The space D is endowed with the norm $\|v\|_D = \|v\|_{H^1(\mathbb{R} \times \mathbb{T}^N)} + \|\partial_{\xi\xi}v + 2\nabla_y\partial_{\xi}v \cdot e + \Delta_yv\|_{L^2(\mathbb{R} \times \mathbb{T}^N)}$. Then, by the similar proofs of Lemma 3.1, Lemma 3.2 and Lemma 3.3 in [6] and Lemma 2.7 in [14], one knows that M_e satisfies all the properties in Lemma 2.7 of [14], such as invertibility and boundedness. For any $e \in \mathbb{S}$, we then define

$$H_{e}(v) := c_{e}\partial_{\xi}v + \partial_{\xi\xi}v + 2\nabla_{x,y}\partial_{\xi}v \cdot e + \Delta_{x,y}v + \frac{2p'}{p}\partial_{\xi}v + \frac{2p'}{p}\nabla_{x,y}v \cdot e + f_{u}(y, pV_{e})v + \left(c_{e}\frac{p'}{p} + \frac{p''}{p}\right)v, \quad v \in D.$$

Notice that $H_e(v) = \widetilde{H}_e(pv)/p$ with $0 < p(\xi) \le 1$ where

$$\widetilde{H}_e(v) := c_e \partial_{\xi} v + \partial_{\xi \xi} v + 2 \nabla_v \partial_{\xi} v \cdot e + \Delta_v v + f_u(y, U_e) v, \quad v \in D.$$

By Lemma 2.9 in [14], one knows that the operator \widetilde{H}_e and its adjoint operator \widetilde{H}_e^* have algebraically simple eigenvalue 0 and the kernel of \widetilde{H}_e is generated by $\partial_{\xi}U_e$. Therefore, the operator H_e and its adjoint operator H_e^* also have algebraically simple eigenvalue 0 and the kernel of H_e is generated by $p^{-1}\partial_{\xi}U_e$. Moreover, the property that the range of H_e is closed in $L^2(\mathbb{R}) \times \mathbb{T}^2$ can be proved in the same line of the proof of [6, Lemma 4.1] by using (2.8).

Now, for any $e \in \mathbb{S}$, $v \in H^2(\mathbb{R} \times \mathbb{T}^2)$, $\vartheta \in \mathbb{R}$ and $\eta \in \mathbb{R}^2$, define

$$\begin{split} K_e(v,\vartheta,\eta) &= \vartheta \partial_\xi (V_e + v) + 2 \nabla_y \partial_\xi (V_e + v) \cdot \eta + \frac{2p'}{p} \nabla_{x,y} (V_e + v) \cdot \eta \\ &+ \frac{1}{p} f(y,p(V_e + v)) - \frac{1}{p} f(y,V_e) + \left(c_e \frac{p'}{p} + \frac{p''}{p} + \beta \right) v, \end{split}$$

and

$$G_e(v,\vartheta,\eta) := \left(v + M_e^{-1}(K_e(v,\vartheta,\eta)), \int_{\mathbb{R}^+ \times \mathbb{T}^N} \left[(V_e(\xi,y) + v(\xi,y))^2 - U_e^2(\xi,y) \right] dy d\xi \right).$$

By following the proof of [14, Lemma 2.10], one can get that for every $e \in \mathbb{S}$, the function $G_e: H^2(\mathbb{R} \times \mathbb{T}^N) \times \mathbb{R} \times \mathbb{R}^N \to D \times \mathbb{R}$ is continuous and it is continuously Fréchet differentiable with respect to (v, ϑ) and doubly continuously Fréchet differentiable with respect to η . For any $e \in \mathbb{S}^{N-1}$ and $(\tilde{v}, \tilde{\vartheta}) \in D \times \mathbb{R}$, define

$$Q_e(\tilde{v}, \tilde{\vartheta}) = \left(\tilde{v} + M_e^{-1}(\tilde{\vartheta}\partial_{\xi}V_e + f_u(y, U_e)\tilde{v} + \left(c_e\frac{p'}{p} + \frac{p''}{p} + \beta\right)\tilde{v}\right), 2\int_{\mathbb{R}^+ \times \mathbb{T}^N} V_e(\xi, y)\tilde{v}(\xi, y)dyd\xi\right),$$

which has the same form as $\partial_{(\nu,\vartheta)}G_e(0,0,0)$. By the properties of H_e and the same line of the proofs of [6, Lemma 3.3] and [14, Lemma 2.11], one can get that Q_e satisfies all properties in [14, Lemma 2.11], such as invertibility and boundedness.

As soon as we have all these properties of these operators, we can follow the same proof of [14, Theorem 1.5] to get that $V_b(\xi, x, y) = p^{-1}(\xi)U_b(\xi, x, y)$ is doubly Fréchet differentiable at any $b \in \mathbb{R}^2 \setminus \{0\}$. Moreover, $||V_e'||$ is bounded for any $e \in \mathbb{S}$.

Thus, by the definition of Fréchet differentiation, we have

$$(U'_e \cdot h)(\cdot, \cdot, \cdot) = p(\xi)(V'_e \cdot h)(\cdot, \cdot, \cdot), \text{ for any } e \in \mathbb{S} \text{ and } h \in \mathbb{R}^2.$$

Therefore, there exist a positive constant C_4 such that

$$|(U'_e \cdot h)(\xi, x, y)| \le p(\xi)||V'_e|||h| \le C_4 e^{-r\xi}|h| \text{ for } \xi \ge 0, (x, y) \in \mathbb{T}^2 \text{ and } h \in \mathbb{R}^2.$$
 (2.9)

By applying similar arguments to the other side, that is, ξ < 0, one can also get that there are positive constants C_5 and μ_5 such that

$$|(U'_{\alpha} \cdot h)(\xi, x, y)| \le C_5 e^{\mu_5 \xi} |h| \text{ for } \xi < 0, (x, y) \in \mathbb{T}^2 \text{ and } h \in \mathbb{R}^2.$$
 (2.10)

Lastly, we differentiate (2.2) at e on $h \in \mathbb{R}^2$ and get that

$$(c'_{e} \cdot h)\partial_{\xi}U_{e} + c_{e}\partial_{\xi}(U'_{e} \cdot h) + \partial_{\xi\xi}(U'_{e} \cdot h) + 2\nabla_{y}\partial_{\xi}U_{e} \cdot (h - (e \cdot h)e)$$

$$+ 2\nabla_{x,y}\partial_{\xi}(U'_{e} \cdot h) \cdot e + \Delta_{x,y}(U'_{e} \cdot h) + f_{u}(x, y, U_{e})(U'_{e} \cdot h) = 0.$$

By changing variables $\xi = (x, y) \cdot e - c_e t$, one has that $u(t, x) := (U'_e \cdot h)((x, y) \cdot e - c_e t, x, y)$ satisfies a parabolic equation

$$u_t - \Delta u = f_u(x, y, U_e)u + (c'_e \cdot h)\partial_{\varepsilon}U_e + 2\nabla_{x,y}\partial_{xi}U_e \cdot (h - (e \cdot h)e).$$

By parabolic estimates, Lemma 2.4 and (2.9)-(2.10), one can get that there are positive constants C_6 and μ_6 such that

$$|u_t(t, x, y)| \le C_6 e^{-\mu_6 |(x, y) \cdot e - c_e t|} |h|,$$

that is,

$$|(\partial_{\xi}U'_{e}\cdot h)(\xi,x,y)|\leq C_{6}e^{-\mu_{6}|\xi|}|h| \text{ for any } h\in\mathbb{R}^{2},\,\xi\in\mathbb{R} \text{ and } (x,y)\in\mathbb{T}^{2}.$$

This completes the proof.

2.2 Proof of Theorem 1.2

Take any two angles α , β of $(0, \pi)$ such that (1.7) holds. Let $\psi(x)$ be a smooth function satisfying Lemma 2.2 for α and β . Take a constant ϱ to be determined later. For every point (x, y) on the curve $y = \psi(\varrho x)/\varrho$, there is a unit normal

$$e(x) = (e_1(x), e_2(x)) = \left(-\frac{\psi'(\varrho x)}{\sqrt{\psi'^2(\varrho x) + 1}}, \frac{1}{\sqrt{\psi'^2(\varrho x) + 1}}\right). \tag{2.11}$$

By Lemma 2.2, every component of e(x) is differentiable with respect to x and

$$e(x) \to (\cos \alpha, \sin \alpha)$$
 as $x \to -\infty$ and $e(x) \to (\cos \beta, \sin \beta)$ as $x \to +\infty$.

Its derivatives can be denoted by

$$e'(x) = (e'_1(x), e'_2(x)) = \Big(-\frac{\varrho \psi''(\varrho x)}{(\psi'^2(\varrho x) + 1)^{\frac{3}{2}}}, -\frac{\varrho \psi'(\varrho x) \psi''(\varrho x)}{(\psi'^2(\varrho x) + 1)^{\frac{3}{2}}} \Big),$$

and

$$e''(x) = (e_1''(x), e_2''(x)) = \left(-\frac{\varrho^2 \psi'''(\varrho x)}{(\psi'^2(\varrho x) + 1)^{\frac{3}{2}}} + \frac{3\varrho^2 \psi'(\varrho x) \psi''^2(\varrho x)}{(\psi'^2(\varrho x) + 1)^{\frac{5}{2}}}, -\frac{\varrho^2 \psi''^2(\varrho x)}{(\psi'^2(\varrho x) + 1)^{\frac{3}{2}}} - \frac{\varrho^2 \psi'(\varrho x) \psi'''(\varrho x)}{(\psi'^2(\varrho x) + 1)^{\frac{3}{2}}} + \frac{3\varrho^2 \psi'^2(\varrho x) \psi''^2(\varrho x)}{(\psi'^2(\varrho x) + 1)^{\frac{5}{2}}}\right).$$

Therefore, by Lemma 2.2, there exist $K_2 > 0$ and $K_3 > 0$ such that

$$|e'(x)| \le \rho K_2 \operatorname{sech}(\gamma x) \text{ and } |e''(x)| \le \rho^2 K_3 \operatorname{sech}(\gamma x) \text{ for all } x \in \mathbb{R}.$$
 (2.12)

Remember that $U^-_{\alpha\beta}(t,x,y)$ defined by (1.6) is a subsolution of (1.1). Now, take a positive constant ε and we define

$$U^+(t, x, y) = U_{e(x)}(\xi(t, x, y), x, y) + \varepsilon \operatorname{sech}(\varrho x),$$

where

$$\xi(t, x, y) = \frac{y - c_{\alpha\beta}t - \psi(\varrho x)/\varrho}{\sqrt{\psi'^2(\varrho x) + 1}},$$
(2.13)

and $c_{\alpha\beta}$ is defined by (1.7). We prove that $U^+(t, x, y)$ is a supersolution of (1.1) for small ε and ϱ .

Lemma 2.8 There exist $\varepsilon_0 > 0$ and $\varrho(\varepsilon_0) > 0$ such that for any $0 < \varepsilon \le \varepsilon_0$ and $0 < \varrho \le \varrho(\varepsilon_0)$, the function $U^+(t, x, y)$ is a supersolution of (1.1) with $U^+_t > 0$. Moreover, it satisfies

$$\lim_{R \to +\infty} \sup_{x^2 + (y - c_{\alpha\beta}t)^2 > R^2} \left| U^+(t, x, y) - U^-_{\alpha\beta}(t, x, y) \right| \le 2\varepsilon, \tag{2.14}$$

and

$$U^{+}(t, x, y) \ge U_{\alpha\beta}^{-}(t, x, y), \text{ for all } t \in \mathbb{R} \text{ and } (x, y) \in \mathbb{R}^{2}.$$

$$(2.15)$$

Proof. We divide the proof into three steps.

Step 1: U^+ is a supersolution. We will pick $\varepsilon_0 > 0$ and $\varrho(\varepsilon)$ such that Lemma 2.8 holds. Assume that

$$\varepsilon_0 \leq \frac{\sigma}{2}$$
,

where $\sigma > 0$ is defined in (F3). More restrictions on ε_0 will be given later. One can compute that

$$\begin{split} LU^{+} &:= U_{t}^{+} - \Delta_{x,y}U^{+} - f(x,y,U^{+}) \\ &= \partial_{\xi}U_{e(x)}\xi_{t} - \partial_{\xi\xi}U_{e(x)}(\xi_{x}^{2} + \xi_{y}^{2}) - 2\nabla_{x,y}\partial_{\xi}U_{e(x)} \cdot (\xi_{x},\xi_{y}) - \Delta_{x,y}U_{e(x)} - \partial_{\xi}U_{e(x)}\xi_{xx} \\ &- U_{e(x)}'' \cdot e'(x) \cdot e'(x) - U_{e(x)}' \cdot e''(x) - 2\partial_{\xi}U_{e(x)}' \cdot e'(x)\xi_{x} - 2\partial_{x}U_{e(x)}' \cdot e'(x) \\ &- \varepsilon \rho^{2} \operatorname{sech}''(\rho x) - f(x,y,U^{+}), \end{split}$$

where $\partial_{\xi}U_{e(x)}$, $\partial_{\xi\xi}U_{e(x)}$, $\nabla_{x,y}\partial_{\xi}U_{e(x)}$, $\Delta_{x,y}U_{e(x)}$, $U''_{e(x)} \cdot e'(x) \cdot e'(x)$, $U''_{e(x)} \cdot e'(x)$, $\partial_{\xi}U'(e(x)) \cdot e'(x)$, $\partial_{\xi}U'_{e(x)} \cdot e'(x)$ are taking values at $(\xi(t,x,y),x,y)$ and U^+ , ξ_t , ξ_x , ξ_y are taking values at (t,x,y). By (2.13), it follows from a direct computation that

$$\xi_{t} = -\frac{c_{\alpha\beta}}{\sqrt{\psi'^{2}(\varrho x) + 1}},
\xi_{x} = -\frac{\varrho \psi'(\varrho x)\psi''(\varrho x)}{(\psi'^{2}(\varrho x) + 1)^{\frac{1}{2}}} \xi - \frac{\psi'(\varrho x)}{\sqrt{\psi'^{2}(\varrho x) + 1}},
\xi_{y} = \frac{1}{\sqrt{\psi'^{2}(\varrho x) + 1}},
\xi_{xx} = -\frac{\varrho^{2}\psi''^{2}(\varrho x) + \varrho^{2}\psi'(\varrho x)\psi''(\varrho x)}{\psi'^{2}(\varrho x) + 1} \xi + \frac{3\varrho^{2}\psi'^{2}(\varrho x)\psi''^{2}(\varrho x)}{(\psi'^{2}(\varrho x) + 1)^{2}} \xi + \frac{\varrho(\psi'^{2}(\varrho x) - 1)\psi''(\varrho x)}{(\psi'^{2}(\varrho x) + 1)^{\frac{3}{2}}},
\xi_{x}^{2} + \xi_{y}^{2} - 1 = \left(\frac{\varrho\psi'(\varrho x)\psi''(\varrho x)}{(\psi'^{2}(\varrho x) + 1)^{\frac{3}{2}}}\right)^{2} \xi^{2} + 2\left(\frac{\varrho\psi'(\varrho x)\psi''(\varrho x)}{(\psi'^{2}(\varrho x) + 1)^{\frac{3}{2}}}\right) \frac{\psi'(\varrho x)}{\sqrt{\psi'^{2}(\varrho x) + 1}} \xi\right).$$

By noticing that $\xi_v = e_2(x)$ and by (2.2), one has

$$LU^{+} = (c_{e(x)} - \xi_{t})\partial_{\xi}U_{e(x)} - \partial_{\xi\xi}U_{e(x)}(\xi_{x}^{2} + \xi_{y}^{2} - 1) - 2\partial_{x}\partial_{\xi}U_{e(x)}(\xi_{x} - e_{1}(x)) - \partial_{\xi}U_{e(x)}\xi_{xx}$$

$$- U''_{e(x)} \cdot e'(x) \cdot e'(x) - U'_{e(x)} \cdot e''(x) - 2\partial_{\xi}U'_{e(x)} \cdot e'(x)\xi_{x} - 2\partial_{x}U'_{e(x)} \cdot e'(x)$$

$$- \varepsilon\varrho^{2}\operatorname{sech}''(\varrho x) + f(x, y, U_{e(x)}) - f(x, y, U^{+}),$$
(2.17)

where $\partial_{\xi}U_{e(x)}$, $\partial_{\xi\xi}U_{e(x)}$, $\partial_{x}\partial_{\xi}U_{e(x)}$, $U''_{e(x)} \cdot e'(x) \cdot e'(x)$, $U'_{e(x)} \cdot e''(x)$, $\partial_{\xi}U'(e(x)) \cdot e'(x)$, $\partial_{x}U'_{e(x)} \cdot e'(x)$, $U'_{e(x)} \cdot e''(x)$, $U'_{e(x)$

$$|\partial_{\xi\xi} U_{e(x)}(\xi_x^2 + \xi_y^2 - 1)| + 2|\partial_x \partial_\xi U_{e(x)}(\xi_x - e_1(x))| + |\partial_\xi U_{e(x)}\xi_{xx}| \le C_5 \varrho \operatorname{sech}(\varrho x). \tag{2.18}$$

Since $||U'_e||$, $||U''_e||$, $||\partial_{\xi}U'_e||$, $||\partial_xU'_e||$ are bounded and by Lemma 2.7, (2.12), there is $C_6 > 0$ such that

$$|U_{e(x)}'' \cdot e'(x) \cdot e'(x)| + |U_{e(x)}' \cdot e''(x)| + 2|\partial_{\xi} U_{e(x)}' \cdot e'(x)\xi_x| + 2|\partial_{x} U_{e(x)}' \cdot e'(x)| \le C_6 \varrho \operatorname{sech}(\varrho x). \tag{2.19}$$

We claim that

Claim 2.9 There is $C_7 > 0$ such that

$$\xi_t - c_{e(x)} = \frac{c_{\alpha\beta}}{\sqrt{\psi'^2(\rho x) + 1}} - c_{e(x)} \ge C_7 sech(\rho x) > 0.$$
 (2.20)

We postpone the proof of this claim after the proof of this lemma.

Then, it follows from (2.17), (2.18), (2.19), (2.20), Lemma 2.1 and $\partial_{\varepsilon}U_{e} < 0$ that

$$LU^{+} \ge -\partial_{\xi} U_{e(x)} C_{7} \operatorname{sech}(\varrho x) - (C_{5} + C_{6}) \varrho \operatorname{sech}(\varrho x) - 2\varepsilon \varrho^{2} \operatorname{sech}(\varrho x) + f(x, y, U_{e(x)}) - f(x, y, U^{+}).$$
(2.21)

By Lemma 2.3, there is C > 0 such that

$$0 < U_e(\xi, x, y) \le \frac{\sigma}{2} \text{ for } \xi \ge C \text{ and } 0 < 1 - U_e(\xi, x, y) \le \frac{\sigma}{2} \text{ for } \xi \le -C.$$
 (2.22)

uniformly for $(x,y) \in \mathbb{T}^2$ and $e \in \mathbb{S}$. Then, for $(t,x,y) \in \mathbb{R} \times \mathbb{R}^2$ such that $\xi(t,x,y) \geq C$ and $\xi(t,x,y) \leq -C$ respectively, one has that $U^+(t,x,y) \leq \sigma/2 + \varepsilon \leq \sigma$ and $U^+(t,x,y) \geq 1 - \sigma/2$ respectively since $\varepsilon \leq \varepsilon_0 \leq \sigma/2$ and hence, it follows from (1.2) that

$$f(x, y, U_{e(x)}) - f(x, y, U^{+}) \ge \lambda \varepsilon \operatorname{sech}(\varrho x).$$
 (2.23)

Since $\partial_{\xi}U_e < 0$ and by (2.21), (2.23), one has that

$$LU^{+} \ge \left(-(C_5 + C_6)\varrho - 2\varepsilon\varrho^2 + \lambda\varepsilon\right)\operatorname{sech}(\varrho x) \ge 0,$$

by taking $0 < \varrho \le \varrho(\varepsilon)$ where $\varrho(\varepsilon) > 0$ is small enough such that

$$-(C_5 + C_6)\varrho - 2\varepsilon\varrho^2 + \lambda\varepsilon > 0, \text{ for all } 0 < \varrho \le \varrho(\varepsilon).$$
 (2.24)

Finally, for $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$ such that $-C \le \xi(t, x, y) \le C$, it follows from Lemma 2.5 that there is k > 0 such that

$$-\partial_{\xi}U_{e}(\xi, x, y) \ge k \text{ for all } e \in \mathbb{S}.$$
 (2.25)

Notice that

$$f(x, y, U_{e(x)}) - f(x, y, U^{+}) \ge -M\varepsilon \operatorname{sech}(\varrho x),$$
 (2.26)

where $M := \max_{(x,y,u) \in \mathbb{T}^2 \times \mathbb{R}} |f_u(x,y,u)|$. Thus, it follows from (2.21), (2.24), (2.25) and (2.26) that

$$LU^{+} \ge \left(kC_7 - (C_5 + C_6)\varrho - 2\varepsilon\varrho^2 - M\varepsilon\right) \operatorname{sech}(\varrho x) \ge \left(kC_7 - (\lambda + M)\varepsilon\right) \operatorname{sech}(\varrho x) \ge 0,$$

by taking $\varepsilon_0 = \min\{\sigma/2, kC_7/(\lambda + M)\}\$ and $0 < \varepsilon \le \varepsilon_0$.

Therefore, $LU^+ \ge 0$ for all $t \in \mathbb{R}$ and $(x, y) \in \mathbb{R}^2$. By the comparison principle, $U^+(t, x, y)$ is a supersolution of (1.1). The property $U_t^+ > 0$ comes from $\partial_{\xi} U_e < 0$ and $c_{\alpha\beta} > 0$.

Step 2: the proof of (2.14). Since $e(x) \to (\cos \alpha, \sin \alpha)$ as $x \to -\infty$ and by the definition of U'_e , there is $R_1 > 0$ such that

$$|U_{e(x)}(\xi(t, x, y), x, y) - U_{\alpha}(\xi(t, x, y), x, y)| \le ||U'_{\alpha}|||e(x) - (\cos \alpha, \sin \alpha)| + o(|e(x) - (\cos \alpha, \sin \alpha)|)$$

$$\le \frac{\varepsilon}{2}, \quad \text{for } x \le -R_1 \text{ and } t \in \mathbb{R}, y \in \mathbb{R}.$$

$$(2.27)$$

Notice that $1/\sqrt{\psi'^2(\varrho x)+1} \to \sin \alpha$ as $x \to -\infty$ and $c_{\alpha\beta}\sin \alpha = c_{\alpha}$. Then, by Lemma 2.2, one has that

$$\xi(t, x, y) \to x \cos \alpha + y \sin \alpha - c_{\alpha} t$$
, as $x \to -\infty$ for any $t \in \mathbb{R}$ and $y \in \mathbb{R}$.

Thus, there is $R_2 > 0$ such that

$$|U_{\alpha}(\xi(t,x,y),x,y) - U_{\alpha}(x\cos\alpha + y\sin\alpha - c_{\alpha}t,x,y)| \le \frac{\varepsilon}{2}, \text{ for } x \le -R_2 \text{ and } t \in \mathbb{R}, y \in \mathbb{R}.$$

Together with (2.27), it follows that

$$|U^{+}(t, x, y) - U_{\alpha}(x \cos \alpha + y \sin \alpha - c_{\alpha}t, x, y)| \le 2\varepsilon, \text{ for } x \le -\max\{R_{1}, R_{2}\} \text{ and } t \in \mathbb{R}, y \in \mathbb{R}.$$
(2.28)

Similarly, one can prove that there is $R_3 > 0$ such that

$$|U^{+}(t, x, y) - U_{\beta}(x \cos \beta + y \sin \beta - c_{\beta}t, x, y)| \le 2\varepsilon, \text{ for } x \ge R_{3} \text{ and } t \in \mathbb{R}, y \in \mathbb{R}.$$
 (2.29)

Now, for $-\max\{R_1, R_2\} \le x \le R_3$, we know that $\psi(\varrho x)$ and $\psi'(\varrho x)$ are bounded. Then, as $y - c_{\alpha\beta}t \to +\infty$, one has that

$$\xi(t, x, y) \to +\infty$$
 and $x \cos \alpha + y \sin \alpha - c_{\alpha}t \to +\infty$, for $-\max\{R_1, R_2\} \le x \le R_3$.

Thus, there is $R_4 > 0$ such that

$$0 < U_{e(x)}(\xi(t, x, y), x, y) \le \frac{\varepsilon}{2},$$

and

$$0 < U_{\alpha}(x\cos\alpha + y\sin\alpha - c_{\alpha}t, x, y), \ U_{\beta}(x\cos\beta + y\sin\beta - c_{\beta}t, x, y) \le \frac{\varepsilon}{2},$$

for $-\max\{R_1, R_2\} \le x \le R_3$ and $y - c_{\alpha\beta}t \ge R_4$. Hence,

$$|U^{+}(t, x, y) - U_{\alpha\beta}(t, x, y)| \le 2\varepsilon, \tag{2.30}$$

for $-\max\{R_1, R_2\} \le x \le R_3$ and $y - c_{\alpha\beta}t \ge R_4$. Similarly, since $U_{e(x)}(-\infty, x, y) = U_{\alpha}(-\infty, x, y) = 1$ uniformly for $(x, y) \in \mathbb{T}^2$, there is R_5 such that

$$|U^{+}(t, x, y) - U_{\alpha\beta}(t, x, y)| \le 2\varepsilon, \tag{2.31}$$

for $-\max\{R_1, R_2\} \le x \le R_3$ and $y - c_{\alpha\beta}t \le -R_5$.

On the other hand, for any fixed $r \in \mathbb{R}$ and any point $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$ such that $x \cos \alpha + y \sin \alpha - c_{\alpha}t = r$, one has that

$$x\cos\beta + y\sin\beta - c_{\beta}t = x\frac{\sin(\alpha - \beta)}{\sin\alpha} + \frac{\sin\beta}{\sin\alpha}r \to +\infty$$
, as $x \to -\infty$,

since $-\pi < \alpha - \beta < 0$ and $c_{\alpha}/\sin \alpha = c_{\beta} \sin \beta$. It implies that $U_{\beta}(x \cos \beta + y \sin \beta - c_{\beta}t, x, y) \to 0$ as $x \to -\infty$ for $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$ such that $x \cos \alpha + y \sin \alpha - c_{\alpha}t = r$. While, $U_{\alpha}(r, x, y) > 0$. Thus, there is $R_6 > 0$ such that

$$U^{-}_{\alpha\beta}(t,x,y) = U_{\alpha}(x\cos\alpha + y\sin\alpha - c_{\alpha}t), \text{ for } x \leq -R_6 \text{ and } t \in \mathbb{R}, y \in \mathbb{R}.$$

Similarly, there is $R_7 > 0$ such that

$$U_{\alpha\beta}^{-}(t, x, y) = U_{\beta}(x \cos \beta + y \sin \beta - c_{\beta}t), \text{ for } x \ge R_7 \text{ and } t \in \mathbb{R}, y \in \mathbb{R}.$$

Then, by (2.28)-(2.31), we have our conclusion (2.14).

Step 3: the proof of (2.15). We only have to prove that $U^+(t, x, y) \ge U_{\alpha}(x \cos \alpha + y \sin \alpha - c_{\alpha}t)$ and $U^+(t, x, y) \ge U_{\beta}(x \cos \beta + y \sin \beta - c_{\beta}t)$ for all $t \in \mathbb{R}$ and $(x, y) \in \mathbb{R}^2$.

Since $U_e(-\infty, x, y) = 1$ and $U_e(+\infty, x, y) = 0$ for any $(x, y) \in \mathbb{T}^2$ and $e \in \mathbb{S}$, there is C > 0 such that

$$0 < U_e(\xi, x, y) \le \sigma \text{ for } \xi \ge C \text{ and } (x, y) \in \mathbb{T}^2,$$

and

$$1 - \sigma \le U_e(\xi, x, y) < 1 \text{ for } \xi \le -C \text{ and } (x, y) \in \mathbb{T}^2,$$

where σ is defined in (F3). By (2.14) and letting $\varepsilon \le \sigma/4$, there is R > 0 such that

$$U^+(t,x,y) \le \sigma$$
, for $(t,x,y) \in \Omega_R^+$ and $U^+(t,x,y) \ge 1 - \sigma$, for $(t,x,y) \in \Omega_R^-$,

where

$$\Omega_R^+ := \{ (t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \le 0 \text{ and } x \cos \alpha + y \sin \alpha - c_\alpha t \ge c_\alpha R \} \cup \{ (t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x > 0 \text{ and } x \cos \beta + y \sin \beta - c_\beta t \ge c_\beta R \},$$

and

$$\Omega_R^- := \{ (t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \le 0 \text{ and } x \cos \alpha + y \sin \alpha - c_\alpha t \le -c_\alpha R \} \cup \{ (t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x > 0 \text{ and } x \cos \beta + y \sin \beta - c_\beta t \le -c_\beta R \}.$$

Notice that for any t, the boundaries of Ω_t^+ and Ω_t^- are connected polylines since $c_\alpha/\sin\alpha = c_\beta/\sin\beta$. By Lemma 2.5 and the definition of $U^+(t, x, y)$, there is $0 < \sigma' \le \sigma$ such that

$$\sigma' \leq U^+(t, x, y) \leq 1 - \sigma', \text{ for } (t, x, y) \in \mathbb{R} \times \mathbb{R}^2 \setminus (\Omega_R^+ \cup \Omega_R^-).$$

For any $\tau \in \mathbb{R}$, let $u_{\tau}(t, x, y) = U_{\alpha}(x \cos \alpha + y \sin \alpha - c_{\alpha}t + \tau)$. Let

$$\omega_{\tau}^+ := \{ (t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \cos \alpha + y \sin \alpha - c_{\alpha} t + \tau \ge C \},$$

and

$$\omega_{\tau}^{-} := \{ (t, x, y) \in \mathbb{R} \times \mathbb{R}^{2}; x \cos \alpha + y \sin \alpha - c_{\alpha} t + \tau \le -C \}.$$

Notice that since $\alpha < \beta$, one has that

$$\{(t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \cos \alpha + y \sin \alpha - c_{\alpha} t \le -c_{\alpha} R\} \subset \Omega_R^-$$

and

$$\Omega_R^+ \subset \{(t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \cos \alpha + y \sin \alpha - c_\alpha t \ge c_\alpha R\}.$$

Thus,

$$\mathbb{R} \times \mathbb{R}^2 \setminus (\omega_{\tau}^+ \cup \omega_{\tau}^-) \subset \Omega_{(C-\tau)/c_{\sigma}}^-$$
 and $\mathbb{R} \times \mathbb{R}^2 \setminus (\Omega_R^+ \cup \Omega_R^-) \subset \omega_{C+c_{\sigma}R}^+$.

Then, by (2.14), $U_e(-\infty, x, y) = 1$ and $U_e(+\infty, x, y) = 0$, there is $\tau_1 \ge c_\alpha R + C$ large enough such that for any $\tau \ge \tau_1$,

$$U^+(t, x, y) \ge 1 - \sigma' \ge u_\tau(t, x, y)$$
, for all $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2 \setminus (\omega_\tau^+ \cup \omega_\tau^-)$,

and

$$u_{\tau}(t, x, y) \leq \sigma' \leq U^{+}(t, x, y), \text{ for all } (t, x, y) \in \mathbb{R} \times \mathbb{R}^{2} \setminus (\Omega_{R}^{+} \cup \Omega_{R}^{-}).$$

Moreover, since $\tau \geq \tau_1 \geq R + C$, one has that

$$U^+(t,x,y) \geq 1 - \sigma \geq \sigma \geq u_\tau(t,x,y), \text{ for all } (t,x,y) \in \omega_\tau^+ \cap \Omega_R^-.$$

Thus, it follows that

$$u_{\tau}(t, x, y) \le U^{+}(t, x, y)$$
, for any $\tau \ge \tau_{1}$ and all $(t, x, y) \in \mathbb{R} \times \mathbb{R}^{2} \setminus (\omega_{\tau}^{-} \cup \Omega_{R}^{+})$. (2.32)

Also notice that

$$u_{\tau}(t,x,y), \ U^{+}(t,x,y) \geq 1 - \sigma \text{ in } \omega_{\tau}^{-} \text{ and } u_{\tau}(t,x,y), \ U^{+}(t,x,y) \leq \sigma \text{ in } \Omega_{R}^{+},$$

and f(x, y, u) is nonincreasing in $u \in (-\infty, \sigma]$ and $u \in [1 - \sigma, +\infty)$ for any $(x, y) \in \mathbb{T}^2$ by (1.2). By following similar proof as the proof of [3, Lemma 4.2] which mainly applied the sliding method and the linear parabolic estimates, one can get that

$$U^+(t, x, y) \ge u_{\tau}(t, x, y)$$
, in ω_{τ}^- and Ω_R^+ .

Combined with (2.32), one has that

$$U^+(t, x, y) \ge u_{\tau}(t, x, y)$$
, for any $\tau \ge \tau_1$ and all $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$.

Now, we decrease τ . Define

$$\tau_* = \inf\{\tau \in \mathbb{R}; U^+(t, x, y) \ge u_\tau(t, x, y) \text{ for all } (t, x, y) \in \mathbb{R} \times \mathbb{R}^2\}.$$

From above arguments, one knows that $\tau_* < +\infty$. Since $U^+(t, x, y) \to U_\alpha(x \cos \alpha + y \sin \alpha - c_\alpha t, x, y)$ as $x \to -\infty$, $U_\alpha(\xi, x, y)$ is decreasing in ξ and by the definition of $u_\tau(t, x, y)$, one also knows that $\tau_* \geq 0$. Assume that $\tau_* > 0$. If

$$\inf\{U^+(t,x,y)-u_{\tau_*}(t,x,y);(t,x,y)\in\mathbb{R}\times\mathbb{R}^2\setminus(\omega_{\tau_*}^-\cup\Omega_R^+)\}>0,$$

then there is $\eta > 0$ such that

$$U^+(t, x, y) \ge u_{\tau_* - \eta}(t, x, y)$$
, for $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2 \setminus (\omega_{\tau_* - \eta}^- \cup \Omega_R^+)$.

Then, one can apply the above arguments again and get that $U^+(t, x, y) \ge u_{\tau_* - \eta}(t, x, y)$ for all $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$ which contradicts the definition of τ_* . Thus,

$$\inf\{U^+(t,x,y)-u_{\tau_*}(t,x,y);(t,x,y)\in\mathbb{R}\times\mathbb{R}^2\setminus(\omega_{\tau_*}^-\cup\Omega_R^+)\}=0.$$

Since $\alpha < \beta$, there is a sequence $\{(t_n, x_n, y_n)\}_{n \in \mathbb{N}}$ in $\mathbb{R} \times \mathbb{R}^2 \setminus (\omega_{\tau_*}^- \cup \Omega_R^+)$ such that

$$-C - \tau_* \le x_n \cos \alpha + y_n \sin \alpha - c_\alpha t_n \le c_\alpha R,$$

and

$$U^{+}(t_{n}, x_{n}, y_{n}) - u_{\tau_{*}}(t_{n}, x_{n}, y_{n}) \to 0$$
, as $n \to +\infty$.

Then, there is $\xi_* \in \mathbb{R}$ such that $x_n \cos \alpha + y_n \sin \alpha - c_\alpha t_n \to \xi_*$ as $n \to +\infty$. Since $U^+(t, x, y) \to U_\alpha(x \cos \alpha + y \sin \alpha - c_\alpha t, x, y)$ as $x \to -\infty$, $U^+(t, x, y) \to U_\beta(x \cos \beta + y \sin \beta - c_\beta t, x, y)$ as $x \to +\infty$ with $\alpha < \beta$ and $\tau_* > 0$, one has that x_n is bounded and there is $x_* \in \mathbb{R}$ such that $x_n \to x_*$ as $n \to +\infty$. Again by $U^+(t, x, y) \to U_\alpha(x \cos \alpha + y \sin \alpha - c_\alpha t, x, y)$ as $x \to -\infty$ and by (2.28), there is R' > 0 such that

$$|U^{+}(t, x, y) - U_{\alpha}(x \cos \alpha + y \sin \alpha - c_{\alpha}t, x, y)| \le \varepsilon \text{ for } x \le -R' \text{ and } t \in \mathbb{R}, y \in \mathbb{R}.$$
 (2.33)

Let $v(t, x, y) = U^+(t, x, y) - u_{\tau_*}(t, x, y)$. Then, $v(t, x, y) \ge 0$ and

$$v(t, x, y) > 0$$
 for any $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$ such that $x \le -R$, $x \cos \alpha + y \sin \alpha - c_{\alpha} t = \xi_*$, (2.34)

by (2.33), $\tau_* > 0$ and taking ε sufficiently small. Since $U^+(t, x, y)$ is a supersolution and $u_{\tau_*}(t, x, y)$ is a solution of (1.1), we have that v(t, x, y) satisfies

$$v_t - \Delta v \ge -b(x, y)v$$
, for $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$,

where b(x, y) is bounded. Since $v(t_n, x_n, y_n) \to 0$ and by the linear parabolic estimates and x_n is bounded, one gets that

$$v(t_n - 1, x_n - R', y_n + \frac{R'\cos\alpha - c_\alpha}{\sin\alpha}) \to 0 \text{ as } n \to +\infty,$$

which contradicts (2.34). Thus, $\tau_* = 0$ and $U^+(t, x, y) \ge U_\alpha(x \cos \alpha + y \sin \alpha - c_\alpha t, x, y)$ for all $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$.

Similarly one can prove that $U^+(t, x, y) \ge U_{\beta}(x \cos \beta + y \sin \beta - c_{\beta}t, x, y)$ for all $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$. In conclusion, $U^+(t, x, y) \ge U^-_{\alpha\beta}(t, x, y)$ for all $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$.

Proof of Claim 2.9. Notice that

$$-\frac{\psi'(\rho x)}{\sqrt{\psi'^2(\rho x) + 1}} = e_1(x) \text{ and } \frac{1}{\sqrt{\psi'^2(\rho x) + 1}} = e_2(x).$$

Let $\hat{\theta}(x) = \arccos e_1(x)$. By Lemma 2.2, one can get that $\alpha < \hat{\theta}(x) < \beta$ for all $x \in \mathbb{R}$ and $\hat{\theta}(-\infty) = \alpha$, $\hat{\theta}(+\infty) = \beta$. Then, $e(x) = (\cos \hat{\theta}, \sin \hat{\theta})$ and

$$\frac{c_{e(x)}}{e_2(x)} = \frac{c_{\hat{\theta}}}{\sin \hat{\theta}}.$$

Thus,

$$\frac{c_{\alpha\beta}}{\sqrt{\psi'^2(\varrho x)+1}} - c_{e(x)} = \sin \hat{\theta} \left(c_{\alpha\beta} - \frac{c_{\hat{\theta}}}{\sin \hat{\theta}} \right).$$

Since $c_{\alpha\beta} > c_{\theta}/\sin\theta$ for any $\theta \in (\alpha,\beta)$ and $0 < \min\{\sin\alpha,\sin\beta\} \le \sin\hat{\theta} \le 1$, one only has to prove that

$$c_{\alpha\beta} - \frac{c_{\hat{\theta}}}{\sin \hat{\theta}} \ge C_7 \operatorname{sech}(\varrho x)$$
, for some positive constant C_7 and when $|x|$ is large. (2.35)

We only consider when x < 0 and similar arguments can be applied for x > 0. Define

$$g(\theta) = \frac{c_{\theta}}{\sin \theta}$$
, for $\theta \in (0, \pi)$.

Obviously, $g(\theta)$ is a C^2 function since c_e is doubly differentiable with respect to e. By (1.7), one has that $g'(\alpha) < 0$. Since $\hat{\theta}(x) \to \alpha$ as $x \to -\infty$, it then follows that

$$\frac{c_{\alpha}}{\sin \alpha} - \frac{c_{\hat{\theta}}}{\sin \hat{\theta}} = g'(\alpha)(\alpha - \hat{\theta}(x)) + o(|\alpha - \hat{\theta}(x)|), \text{ for } x \text{ negative enough.}$$
 (2.36)

Moreover, by (2.1), one has that

$$\hat{\theta}(x) - \alpha = \int_{-\infty}^{x} \hat{\theta}'(s)ds = -\int_{-\infty}^{x} \frac{e_1'(s)}{\sqrt{1 - e_1^2(s)}} ds = \int_{\infty}^{x} \frac{\varrho \psi''(\varrho s)}{\psi'^2(\varrho s) + 1} ds$$

$$\geq \frac{1}{|\psi'|_{L^{\infty}}^2 + 1} (\psi'(\varrho x) + \cot \alpha) \geq \frac{k_1}{|\psi'|_{L^{\infty}}^2 + 1} \operatorname{sech}(\varrho x).$$

One then can conclude (2.35) from (2.36) for x negative enough.

Now, we are ready to prove Theorem 1.2.

Proof of Theorem 1.2. Let $u_n(t, x)$ be the solution of (1.1) for $t \ge -n$ with initial data

$$u_n(-n, x, y) = U_{\alpha\beta}^-(-n, x, y).$$

By Lemma 2.8, one can get from the comparison principle that

$$U_{\alpha\beta}^{-}(t, x, y) \le u_n(t, x, y) \le U^{+}(t, x, y), \text{ for } t \ge -n \text{ and } (x, y) \in \mathbb{R}^2.$$
 (2.37)

Since $U_{\alpha\beta}^-(t, x, y)$ is a subsolution, the sequence $u_n(t, x, y)$ is increasing in n. Letting $n \to +\infty$ and by parabolic estimates, the sequence $u_n(t, x, y)$ converges to an entire solution V(t, x, y) of (1.1). By (2.37), one has that

$$U^-_{\alpha\beta}(t,x,y) \leq V(t,x,y) \leq U^+(t,x,y)$$
, for $t \in \mathbb{R}$ and $(x,y) \in \mathbb{R}^2$.

Then, it follows from Lemma 2.8 that (1.8) holds.

By $U_{\alpha\beta}^-(t, x, y)$ is increasing in t and the maximum principle, one has that $(u_n)_t(t, x, y) > 0$ for all $t \in (-n, +\infty)$ and $(x, y) \in \mathbb{R}^2$. By letting $n \to +\infty$ and the strong maximum principle, one concludes that $u_t(t, x, y) > 0$ for all $t \in \mathbb{R}$ and $(x, y) \in \mathbb{R}^2$. This completes the proof.

2.3 Proofs of Corollaries 1.5, 1.6 and Theorem 1.7

We then give some examples to show that Theorem 1.2 is not empty, that is, Corollaries 1.5, 1.6.

Proof of Corollary 1.5. Notice that c_{θ} and c'_{θ} are uniformly bounded for $\theta \in [0, \pi]$. Let $g(\theta) := c_{\theta} / \sin \theta$. Then,

$$g'(\theta) = \frac{c'_{\theta} \cdot (-\sin\theta, \cos\theta)}{\sin\theta} - \frac{c_{\theta}\cos\theta}{\sin^2\theta}.$$

Obviously, there are constants $0 < \alpha_1 < \beta_1 < \pi$ such that $g'(\theta) < 0$ for $\theta \in (0, \alpha_1)$ and $g'(\theta) > 0$ $\theta \in (\beta_1, \pi)$ since c'_e is bounded for any $e \in \mathbb{S}$ and $\sin \theta \to 0$ as $\theta \to 0$ or π . One can also notice that $g(\theta) \to +\infty$ as $\theta \to 0$ or $\theta \to \pi$. By continuity, one can take any $\alpha \in (0, \alpha_1)$ and there is $\beta \in (\beta_1, \pi)$ such that $g(\alpha) = g(\beta)$ and $g(\theta) < g(\alpha) = g(\beta)$ for all $\theta \in (\alpha, \beta)$.

Then, the conclusion of Corollary 1.5 follows from Theorem 1.2.

Proof of Corollary 1.6. Take two directions $e_1 = (\cos \theta_1, \sin \theta_1)$ and $e_2 = (\cos \theta_2, \sin \theta_2)$ where $\theta_1, \theta_2 \in (0, 2\pi)$. Assume without loss of generality that $\theta_2 > \theta_1$. Rotate the coordinate by changing variables as

$$\begin{cases} X = x \cos \theta + y \sin \theta, \\ Y = -x \sin \theta + y \cos \theta, \end{cases}$$

where θ varies from $\theta_2 - \pi/2$ to $\theta_1 + \pi/2$. Let u(t, x, y) be the solution of (1.1) and let $v(t, X, Y) = u(t, x, y) = u(t, X \cos \theta - Y \sin \theta, X \sin \theta + Y \cos \theta)$. Then, v(t, X, Y) satisfies

$$v_t - \Delta_{X,Y} v = g(X, Y, u), \quad (t, X, Y) \in \mathbb{R} \times \mathbb{R}^2, \tag{2.38}$$

where $g(X, Y, v) = f(X \cos \theta - Y \sin \theta, X \sin \theta + Y \cos \theta, v)$. Notice that g(X, Y, v) satisfies (F1)-(F3).

Assume without loss of generality that $\theta_2 - \theta_1 < \pi$. Otherwise, if $\theta_2 - \theta_1 > \pi$, we can take θ varying from θ_2 to $2\pi + \theta_1$. Then, under the new coordinate, directions e_1 and e_2 become $(\cos(\theta_1 + \pi/2 - \theta), \sin(\theta_1 + \pi/2 - \theta))$ and $(\cos(\theta_2 + \pi/2 - \theta), \sin(\theta_2 + \pi/2 - \theta))$ where $0 < \theta_1 + \pi/2 - \theta < \theta_2 + \pi/2 - \theta < \pi$. Since $\sin \theta$ is increasing in $[0, \pi/2]$ and decreasing in $[\pi/2, \pi]$, one has that

$$\frac{c_{e_1}}{\sin(\theta_1 + \pi/2 - \theta)}$$
 is increasing from $\frac{c_{e_1}}{\sin(\theta_1 - \theta_2 + \pi)}$ to $+\infty$ as θ varies from $\theta_2 - \pi/2$ to $\theta_1 + \pi/2$,

and

$$\frac{c_{e_2}}{\sin(\theta_2 + \pi/2 - \theta)}$$
 is decreasing from $+\infty$ to $\frac{c_{e_2}}{\sin(\theta_2 - \theta_1)}$ as θ varies from $\theta_2 - \pi/2$ to $\theta_1 + \pi/2$.

By continuity and for any $0 < \theta_2 - \theta_1 < \pi$, there is θ^* such that

$$\frac{c_{e_1}}{\sin(\theta_1 + \pi/2 - \theta^*)} = \frac{c_{e_2}}{\sin(\theta_2 + \pi/2 - \theta^*)}.$$

On the other hand, by the proof of Corollary 1.5, there is $0 < \alpha_1 < \pi$ small enough such that for $0 < \pi - (\theta_2 - \theta_1) < \alpha_1$, it holds

$$\frac{c_{e_1}}{\sin(\theta_1 + \pi/2 - \theta^*)} = \frac{c_{e_2}}{\sin(\theta_2 + \pi/2 - \theta^*)} > \frac{c_{\theta}}{\sin(\theta - \theta^*)} \text{ for } \theta_1 + \pi/2 < \theta < \theta_1 + \pi/2.$$

Then, it follows from Corollary 1.5 that there is an entire solution V(t, X, Y) of (2.38) satisfying (1.10). By taking $\rho = \cos(\pi - \alpha_1) - 1$ and $e_0 = (\cos \theta^*, \sin \theta^*)$, the conclusion of Corollary 1.6 immediately follows.

Now, we show that condition (1.7) without $g'(\alpha) < 0$ and $g'(\beta) > 0$ is necessary for the existence of the curved front in Theorem 1.2.

Proof of Theorem1.7. We first prove that

$$\frac{c_{\alpha}}{\sin \alpha} = \frac{c_{\beta}}{\sin \beta}.\tag{2.39}$$

Assume by contradiction that $c_{\alpha}/\sin \alpha \neq c_{\beta}/\sin \beta$. Take a sequence $\{t_n\}_{n\in\mathbb{N}}$ such that $t_n \to +\infty$. Then, for the sequence

$$(x_n, y_n) = \left(\frac{(c_\alpha \sin \beta - c_\beta \sin \alpha)t_n}{\sin(\beta - \alpha)}, \frac{(c_\alpha \cos \beta - c_\beta \cos \alpha)t_n}{\sin(\alpha - \beta)}\right),$$

one has that $x_n^2 + (y_n - c_{\alpha\beta}t_n)^2 \to +\infty$ as $n \to +\infty$ for any $c_{\alpha\beta} \in \mathbb{R}$ since $c_{\alpha}/\sin \alpha \neq c_{\beta}/\sin \beta$. Notice that for any n, there are k_n^1 , $k_n^2 \in \mathbb{Z}$ and x_n' , $y_n' \in [0, L_2)$ such that $x_n = k_n^1 L_1 + x_n'$ and $y_n = k_n^2 L_2 + y_n'$. Moreover, up to extract subsequences of x_n and y_n , there are $x_*' \in [0, L_1]$ and $y'_* \in [0, L_2]$ such that $x'_n \to x'_*$ and $y'_n \to y'_*$ as $n \to +\infty$. Since $f(x, y, \cdot)$ is L-periodic in (x, y), one has $f(x+x_n, y+y_n, \cdot) \to f(x+x'_*, y+y'_*, \cdot)$ as $n \to +\infty$. Let $v_n(t, x, y) = V(t+t_n, x+x_n, y+y_n)$. By standard parabolic estimates, $v_n(t, x, y)$, up to extract of a subsequence, converges to a solution $v_\infty(t, x, y)$ of

$$v_t - \Delta v = f(x + x'_*, y + y'_*, v), \text{ for } t \in \mathbb{R} \text{ and } (x, y) \in \mathbb{R}^2.$$
 (2.40)

By definitions of x_n and y_n , one can also have that

$$U_{\alpha\beta}^-(t+t_n,x+x_n,y+y_n)\to \hat{U}_{\alpha\beta}^-(t,x,y)$$
, as $n\to +\infty$ uniformly in $\mathbb{R}\times\mathbb{R}^2$,

where

$$\hat{U}_{\alpha\beta}^{-}(t, x, y) := \max\{U_{\alpha}(x\cos\alpha + y\sin\alpha - c_{\alpha}t, x + x'_{*}, y + y'_{*}), U_{\beta}(x\cos\beta + y\sin\beta - c_{\beta}t, x + x'_{*}, y + y'_{*})\}.$$

Moreover, by (1.8) and $x_n^2 + (y_n - c_{\alpha\beta}t_n)^2 \to +\infty$ as $n \to +\infty$, one gets that

$$v_n(t, x, y) \to \hat{U}_{\alpha\beta}^-(t, x, y)$$
 as $n \to +\infty$ locally uniformly in $\mathbb{R} \times \mathbb{R}^2$.

It implies that $v_{\infty}(t, x, y) = \hat{U}_{\alpha\beta}^{-}(t, x, y)$ which is impossible since $\hat{U}_{\alpha\beta}^{-}(t, x, y)$ is not a solution of (2.40). Therefore, (2.39) holds.

Then, we prove that

$$c_{\alpha\beta} = \frac{c_{\alpha}}{\sin \alpha} = \frac{c_{\beta}}{\sin \beta}.$$
 (2.41)

Assume by contradiction that $c_{\alpha\beta} \neq c_{\alpha}/\sin \alpha$. Take a sequence $(t_n)_{n\in\mathbb{N}}$ such that $t_n = L_2 n \sin \alpha/c_{\alpha} \rightarrow +\infty$ and consider the sequence

$$(x_n, y_n) = (0, \frac{c_\alpha}{\sin \alpha} t_n).$$

Notice that $x_n^2 + (y_n - c_{\alpha\beta}t_n)^2 \to +\infty$ as $n \to +\infty$ since $c_{\alpha\beta} \neq c_{\alpha}/\sin\alpha$, $t_nc_{\alpha}/\sin\alpha = nL_2$ and $U_{\alpha\beta}^-(t+t_n,x+x_n,y+y_n) = U_{\alpha\beta}^-(t,x,y)$. Then, one can make the similar arguments as above to get a contradiction. Thus, (2.41) holds.

At last, we prove that

$$\frac{c_{\theta}}{\sin \theta} < c_{\alpha\beta} = \frac{c_{\alpha}}{\sin \alpha} = \frac{c_{\beta}}{\sin \beta} \text{ for any } \theta \in (\alpha, \beta).$$

Assume by contradiction that there is $\theta \in (\alpha, \beta)$ such that $c_{\theta}/\sin \theta \ge c_{\alpha\beta}$. Then, two cases may occur: (i) $c_{\theta}/\sin \theta > c_{\alpha\beta}$; (ii) $c_{\theta}/\sin \theta = c_{\alpha\beta}$.

For case (i), take t = 0 and by (1.8), for any $\varepsilon > 0$, there is $R_{\varepsilon} > 0$ such that

$$\sup_{|(x,y)|>R_{\varepsilon}} \left| V(0,x,y) - U_{\alpha\beta}^{-}(0,x,y) \right| \le \varepsilon. \tag{2.42}$$

We claim that

Claim 2.10 *There exist constants* $\tau \in \mathbb{R}$ *and* $\delta > 0$ *such that*

$$V(t, x, y) \ge U_{\theta}(x \cos \theta + y \sin \theta - c_{\theta}t + \tau, x, y) - \delta e^{-\delta t}$$
, for $t \ge 0$ and $x \in \mathbb{R}^2$.

In order to not lengthen the proof, we postpone the proof of this claim after the proof of Theorem 1.7. Take a sequences $(t_n)_{n\in\mathbb{N}}$ such that $t_n \to +\infty$ as $n \to +\infty$ and $y_n = c_{\alpha\beta}t_n + R$ where R is a constant. Then, since $U_e(+\infty, x, y) = 0$ for all $e \in \mathbb{S}$ and $(x, y) \in \mathbb{T}^2$, one can take R large enough such that

$$\begin{split} U_{\alpha\beta}^{-}(t_{n},0,y_{n}) &= \max\{U_{\alpha}(y_{n}\sin\alpha - c_{\alpha}t_{n},0,y_{n}), U_{\beta}(y_{n}\sin\beta - c_{\beta}t_{n},0,y_{n})\}\\ &= \max\{U_{\alpha}(R\sin\alpha,0,y_{n}), U_{\beta}(R\sin\beta,0,y_{n})\} \leq \frac{1}{4}. \end{split}$$

By (1.8) and even if it means increasing R, one has that

$$V(t_n, 0, y_n) \le U_{\alpha\beta}^-(t_n, 0, y_n) + \frac{1}{4} \le \frac{1}{2} \text{ for all } n.$$
 (2.43)

However, since $c_{\theta}/\sin\theta > c_{\alpha\beta}$ and hence,

$$y_n \sin \theta - c_\theta t_n = (c_{\alpha\beta} \sin \theta - c_\theta)t_n + R \sin \theta \rightarrow -\infty$$
, as $n \rightarrow +\infty$,

it follows from Claim 2.10 that

$$V(t_n, 0, y_n) \ge U_{\theta}(y_n \sin \theta - c_{\theta}t_n + \tau, 0, y_n) \to 1 \text{ as } n \to +\infty,$$

which contradicts (2.43). Case (i) is ruled out.

Now we consider case (ii). Since $U_e(-\infty, x, y) = 1$ and $U_e(+\infty, x, y) = 0$ for any $(x, y) \in \mathbb{T}^2$ and $e \in \mathbb{S}$, there is C > 0 such that

$$0 < U_e(\xi, x, y) \le \sigma \text{ for } \xi \ge C \text{ and } (x, y) \in \mathbb{T}^2,$$

and

$$1 - \sigma \le U_e(\xi, x, y) < 1 \text{ for } \xi \le -C \text{ and } (x, y) \in \mathbb{T}^2,$$

where σ is defined in (F3). By (1.8), there is R > 0 such that

$$V(t, x, y) \leq \sigma$$
, for $(t, x, y) \in \Omega_R^+$ and $V(t, x, y) \geq 1 - \sigma$, for $(t, x, y) \in \Omega_R^-$,

where

$$\Omega_R^+ := \{ (t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \le 0 \text{ and } x \cos \alpha + y \sin \alpha - c_\alpha t \ge c_\alpha R \} \cup \{ (t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x > 0 \text{ and } x \cos \beta + y \sin \beta - c_\beta t \ge c_\beta R \},$$

and

$$\Omega_R^- := \{(t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \le 0 \text{ and } x \cos \alpha + y \sin \alpha - c_\alpha t \le -c_\alpha R\} \cup \{(t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \le 0 \text{ and } x \cos \alpha + y \sin \alpha - c_\alpha t \le -c_\alpha R\} \cup \{(t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \le 0 \text{ and } x \cos \alpha + y \sin \alpha - c_\alpha t \le -c_\alpha R\} \cup \{(t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \le 0 \text{ and } x \cos \alpha + y \sin \alpha - c_\alpha t \le -c_\alpha R\} \cup \{(t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \le 0 \text{ and } x \cos \alpha + y \sin \alpha - c_\alpha t \le -c_\alpha R\} \cup \{(t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \le 0 \text{ and } x \cos \alpha + y \sin \alpha - c_\alpha t \le -c_\alpha R\} \cup \{(t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \le 0 \text{ and } x \cos \alpha + y \sin \alpha - c_\alpha t \le -c_\alpha R\} \cup \{(t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \le 0 \text{ and } x \cos \alpha + y \sin \alpha - c_\alpha t \le -c_\alpha R\} \cup \{(t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \le 0 \text{ and } x \cos \alpha + y \sin \alpha - c_\alpha t \le -c_\alpha R\} \cup \{(t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \le 0 \text{ and } x \cos \alpha + y \sin \alpha - c_\alpha t \le -c_\alpha R\} \cup \{(t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \le 0 \text{ and } x \cos \alpha + y \sin \alpha - c_\alpha t \le -c_\alpha R\} \cup \{(t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \ge 0 \text{ and } x \cos \alpha + y \sin \alpha - c_\alpha t \le -c_\alpha R\} \cup \{(t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \ge 0 \text{ and } x \cos \alpha + y \cos$$

$$x > 0$$
 and $x \cos \beta + y \sin \beta - c_{\beta}t \le -c_{\beta}R$.

By a similar proof as of Lemma 2.5, there is $0 < \sigma' \le \sigma$ such that

$$\sigma' \leq V(t, x, y) \leq 1 - \sigma'$$
, for $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2 \setminus (\Omega_R^+ \cup \Omega_R^-)$.

For any $\tau \in \mathbb{R}$, let $u_{\tau}(t, x, y) = U_{\theta}(x \cos \theta + y \sin \theta - c_{\theta}t + \tau, x, y)$. Let

$$\omega_{\tau}^{+} := \{ (t, x, y) \in \mathbb{R} \times \mathbb{R}^{2}; x \cos \theta + y \sin \theta - c_{\theta}t + \tau \ge C \},$$

and

$$\omega_{\tau}^{-} := \{ (t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \cos \theta + y \sin \theta - c_{\theta} t + \tau \le -C \}.$$

Since $\alpha < \theta < \beta$ and $c_{\theta} / \sin \theta = c_{\alpha} / \sin \alpha = c_{\beta} / \sin \beta$, one can easily check that

$$\mathbb{R} \times \mathbb{R}^2 \setminus (\omega_{\tau}^+ \cup \omega_{\tau}^-) \subset \Omega_{(C-\tau)/c_{\alpha}}^-$$
 and $\mathbb{R} \times \mathbb{R}^2 \setminus (\Omega_R^+ \cup \Omega_R^-) \subset \omega_{C+c_{\alpha}R}^+$.

Then, by (1.8), $U_e(-\infty, x, y) = 1$ and $U_e(+\infty, x, y) = 0$, there is $\tau_1 \ge c_\alpha R + C$ large enough such that for any $\tau \ge \tau_1$,

$$V(t, x, y) \ge 1 - \sigma' \ge u_{\tau}(t, x, y)$$
, for all $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2 \setminus (\omega_{\tau}^+ \cup \omega_{\tau}^-)$,

and

$$u_{\tau}(t,x,y) \leq \sigma' \leq V(t,x,y), \text{ for all } (t,x,y) \in \mathbb{R} \times \mathbb{R}^2 \setminus (\Omega_R^+ \cup \Omega_R^-).$$

Moreover, since $\tau \geq \tau_1 \geq c_{\alpha}R + C$, one has that

$$V(t,x,y) \ge 1 - \sigma \ge \sigma \ge u_{\tau}(t,x,y), \text{ for all } (t,x,y) \in \omega_{\tau}^+ \cap \Omega_R^-.$$

Thus, it follows that

$$u_{\tau}(t, x, y) \le V(t, x, y)$$
, for any $\tau \ge \tau_1$ and all $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2 \setminus (\omega_{\tau}^- \cup \Omega_R^+)$. (2.44)

Also notice that

$$u_{\tau}(t,x,y), V(t,x,y) \geq 1 - \sigma \text{ in } \omega_{\tau}^{-} \text{ and } u_{\tau}(t,x,y), V(t,x,y) \leq \sigma \text{ in } \Omega_{R}^{+},$$

and f(x, y, u) is nonincreasing in $u \in (-\infty, \sigma]$ and $u \in [1 - \sigma, +\infty)$ for any $(x, y) \in \mathbb{T}^2$ by (1.2). By following similar proof as the proof of [3, Lemma 4.2] which mainly applied the sliding method and the linear parabolic estimates, one can get that

$$V(t, x, y) \ge u_{\tau}(t, x, y)$$
, in ω_{τ}^- and Ω_R^+ .

Combined with (2.44), one has that

$$V(t, x, y) \ge u_{\tau}(t, x, y)$$
, for any $\tau \ge \tau_1$ and all $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$.

Let

$$\tau_* = \inf\{\tau \in \mathbb{R}; u_\tau(t, x, y) \le V(t, x, y) \text{ for all } (t, x, y) \in \mathbb{R} \times \mathbb{R}^2\}.$$

By above arguments, one knows that $\tau_* < +\infty$. On the other hand, for any fixed (t, x, y), $u_{\tau}(t, x, y) = U_{\theta}(x \cos \theta + y \sin \theta - c_{\theta}t + \tau, x, y) \rightarrow 1$ as $\tau \rightarrow -\infty$ and V(t, x, y) < 1 by the maximum principle. By the definition of τ_* , one also has that $\tau_* > -\infty$. Thus, $|\tau_*|$ is bounded. If

$$\inf\{V(t,x,y)-u_{\tau_*};(t,x,y)\in\mathbb{R}\times\mathbb{R}^2\setminus(\omega_{\tau_*}^-\cup\Omega_R^+)\}>0,$$

there is $\eta > 0$ such that

$$V(t, x, y) \ge u_{\tau_* - \eta}(t, x, y), \text{ for } (t, x, y) \in \mathbb{R} \times \mathbb{R}^2 \setminus (\omega_{\tau_*}^- \cup \Omega_R^+).$$

Then, one can follow above arguments again to get that

$$u_{\tau_*-\eta}(t, x, y) \le V(t, x, y)$$
, for $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$,

which contradicts the definition of τ_* . Thus,

$$\inf\{V(t,x,y)-u_{\tau_*};(t,x,y)\in\mathbb{R}\times\mathbb{R}^2\setminus(\omega_{\tau_*}^-\cup\Omega_R^+)\}=0.$$

Since $V(t, x, y) \ge \sigma'$ in $\mathbb{R} \times \mathbb{R}^2 \setminus (\omega_{\tau_*}^- \cup \Omega_R^+)$ and $u_{\tau_*}(t, x, y) = U_{\theta}(x \cos \theta + y \sin \theta - c_{\theta}t + \tau_*, x, y) \to 0$ as $x \cos \theta + y \sin \theta - c_{\theta}t \to +\infty$, there is $R_1 > 0$ and there is a sequence $\{(t_n, x_n, y_n)\}_{n \in \mathbb{N}}$ in $\mathbb{R} \times \mathbb{R}^2 \setminus (\omega_{\tau_*}^- \cup \Omega_R^+)$ such that

$$-C - \tau_* \le x_n \cos \theta + y_n \sin \theta - c_\theta t_n \le R_1 \tag{2.45}$$

and

$$V(t_n, x_n, y_n) - U_{\theta}(x_n \cos \theta + y_n \sin \theta - c_{\theta}t_n + \tau_*, x_n, y_n) \to 0, \text{ as } n \to +\infty.$$
 (2.46)

Notice that x_n is bounded. Otherwise, if $x_n \to -\infty$ as $n \to +\infty$, then it follows from (2.45) and $\theta < \alpha$ that

$$x_n \cos \alpha + y_n \sin \alpha - c_{\alpha} t_n = x_n \cos \alpha + \sin \alpha \left(y_n - \frac{c_{\alpha}}{\sin \alpha} t_n \right)$$

$$= x_n \cos \alpha + \sin \alpha \left(y_n - \frac{c_{\theta}}{\sin \theta} t_n \right)$$

$$= \left(\cos \alpha - \frac{\sin \alpha \cos \theta}{\sin \theta} \right) x_n + \sin \alpha \sin \theta \left(\cos \theta x_n + \sin \theta y_n - c_{\theta} t_n \right)$$

$$\to +\infty, \qquad \text{as } n \to +\infty,$$

and $x_n^2 + (y_n - c_{\alpha\beta}t_n)^2 \to +\infty$ as $n \to +\infty$. It implies that $V(t_n, x_n, y_n) \to U^-_{\alpha\beta}(t_n, x_n, y_n) \to 1$ as $n \to +\infty$ which contradicts $u_{\tau_*}(t, x, y) \le 1 - \sigma'$ in $\mathbb{R} \times \mathbb{R}^2 \setminus \omega^-_{\tau_*}$ and (2.46). Similarly, it is not possible that $x_n \to +\infty$ as $n \to +\infty$. Thus, there is $x_* \in \mathbb{R}$ such that $x_n \to x_*$ as $n \to +\infty$. Let

 $w(t, x, y) = V(t, x, y) - u_{\tau_*}(t, x, y)$. Then, by (2.46), $w(t_n, x_n, y_n) \to 0$ as $n \to +\infty$. Consider the point $(t_n - 1, x_n - R', y_n - c_{\theta}/\sin\theta + R'\cos\theta/\sin\theta)$ for some constant R'. Notice that by (2.45),

$$(x_n - R')\cos\theta + (y_n - c_\theta/\sin\theta + R'\cos\theta/\sin\theta)\sin\theta - c_\theta(t_n - 1) \in [-C - \tau_*, R_1], \text{ for any } n$$

and

$$(x_n - R')\cos\alpha + (y_n - c_\theta/\sin\theta + R'\cos\theta/\sin\theta)\sin\alpha - c_\alpha(t_n - 1) \to +\infty$$
, as $R' \to +\infty$,

for any n. By taking R' large enough, one can let

$$V(t_n - 1, x_n - R', y_n - c_\theta + R' \cos \theta / \sin \theta) > 1 - \sigma'$$
, for any n .

Then, by noticing that $(t_n - 1, x_n - R, y_n - c_\theta + R\cos\theta/\sin\theta)$ satisfies (2.45) and hence $u_{\tau_*}(t_n - 1, x_n - R, y_n - c_\theta + R\cos\theta/\sin\theta) \le 1 - \sigma'$, one has that

$$w(t_n - 1, x_n - R, y_n - c_\theta + R\cos\theta/\sin\theta) > 0. \tag{2.47}$$

However, since V(t, x, y) and $u_{\tau_{\tau}}(t, x, y)$ are solutions of (1.1), we have that w(t, x, y) satisfies

$$w_t - \Delta w \ge -b(x, y)w$$
, for $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$,

where b(x, y) is bounded. By the linear parabolic estimates, one can get that

$$w(t_n-1,x_n-R,y_n-c_\theta+R\cos\theta/\sin\theta)\to 0$$
, as $n\to+\infty$,

which contradicts (2.47). Therefore, case (ii) is ruled out.

In conclusion,
$$c_{\theta}/\sin\theta < c_{\alpha\beta}$$
 for any $\theta \in (\alpha, \beta)$.

We finish this section by proving Claim 2.10.

Proof of Claim 2.10. Take $\delta > 0$ such that

$$\delta \leq \min\left\{\frac{\sigma}{2}, \lambda\right\},\,$$

where σ and λ are defined in (F3). Since $U_{\theta}(-\infty, x, y) = 1$ and $U_{\theta}(+\infty, x, y) = 0$ for $(x, y) \in \mathbb{T}^2$, there is C > 0 such that

$$\begin{cases} 0 < U_{\theta}(\xi, x, y) \le \delta, & \text{for } \xi \ge C \text{ and } (x, y) \in \mathbb{T}^2 \\ 1 - \delta \le U_{\theta}(\xi, x, y) < 1, & \text{for } \xi \le -C \text{ and } (x, y) \in \mathbb{T}^2. \end{cases}$$

From Lemma 2.5, there is k > 0 such that $-\partial_{\xi}U_{\theta}(\xi, x, y) \ge k$ for $-C \le \xi \le C$ and $(x, y) \in \mathbb{T}^2$. Take $\omega > 0$ such that

$$k\omega \ge \delta + M,$$
 (2.48)

where $M = \max_{(x,y,u) \in \mathbb{T}^2 \times \mathbb{R}} |f_u(x,y,u)|$. It follows from (2.42) and the definition of $U^-_{\alpha\beta}$ that there is $R_{\delta} > 0$ such that

$$V(0, x, y) \ge 1 - \delta$$
, for $(x, y) \in \Omega$

where

$$\Omega := \{x \le 0, y \in \mathbb{R}; x \cos \alpha + y \sin \alpha \le -R_{\delta}\} \cup \{x \ge 0, y \in \mathbb{R}; x \cos \beta + y \sin \beta \le -R_{\delta}\}.$$

Define

$$v^{-}(t, x, y) = U_{\theta}(\xi^{-}(t, x, y), x, y) - \delta e^{-\delta t},$$

where

$$\xi^{-}(t, x, y) = x \cos \theta + y \sin \theta - c_{\theta}t - \omega e^{-\delta t} + \omega + \hat{R}_{\delta} + C,$$

and $\hat{R}_{\delta} = R_{\delta} \sin \theta \max\{1/\sin \alpha, 1/\sin \beta\}$. We prove that $v^{-}(t, x, y)$ is a subsolution of the problem satisfied by V(t, x, y) for $t \ge 0$ and $(x, y) \in \mathbb{R}^{2}$.

Firstly, we check the initial data. Since $\alpha < \theta < \beta$, one has that

$$\{(x,y)\in\mathbb{R}^2;\xi^-(0,x,y)\leq C\}\subset\Omega.$$

Then,

$$v^-(0, x, y) \le 1 - \delta \le V(0, x, y)$$
, for $(x, y) \in \mathbb{R}^2$ such that $\xi(0, x, y) \le C$.

For $(x, y) \in \mathbb{R}^2$ such that $\xi(0, x, y) \ge C$, one has that

$$v^{-}(0, x, y) \le \delta - \delta = 0 \le V(0, x, y).$$

Thus, $v^{-}(0, x, y) \le V(0, x, y)$ for all $(x, y) \in \mathbb{R}^{2}$.

We then check that

$$Nv := v_t^- - \Delta v^- - f(x, y, v^-) \le 0$$
, for $t \ge 0$ and $(x, y) \in \mathbb{R}^2$.

By some computation and (2.2), one has that

$$Nv = \omega \delta e^{-\delta t} \partial_{\xi} U_{\theta} + \delta^{2} e^{-\delta t} + f(x, y, U_{\theta}) - f(x, y, v^{-}).$$
 (2.49)

For $t \ge 0$ and $(x, y) \in \mathbb{R}^2$ such that $\xi(t, x, y) \ge C$, one has that $0 < U_{\theta}(\xi(t, x, y), x, y) \le \delta$ and hence $v^-(t, x, y) \le 2\delta \le \sigma$. Thus, by (1.2), it follows that

$$f(x, y, U_{\theta}) - f(x, y, v^{-}) \le -\lambda \delta e^{-\delta t}. \tag{2.50}$$

Since $\partial_{\xi}U_{\theta} < 0$, it follows from (2.49) and (2.50) that

$$Nv \le \delta^2 e^{-\delta t} - \lambda \delta e^{-\delta t} \le 0.$$

Similarly, one can prove that $Nv \le 0$ for $t \ge 0$ and $(x, y) \in \mathbb{R}^2$ such that $\xi(t, x, y) \le -C$. Finally, for $t \ge 0$ and $(x, y) \in \mathbb{R}^2$ such that $-C \le \xi(t, x, y) \le C$, one has that $-\partial_{\xi} U_{\theta}(\xi(t, x, y), x, y) \ge k$ and

$$f(x, y, U_{\theta}) - f(x, y, v^{-}) \le M\delta e^{-\delta t}, \tag{2.51}$$

where $M = \max_{(x,y,u) \in \mathbb{T}^2 \times \mathbb{R}} |f_u(x,y,u)|$. Then, it follows from (2.48), (2.49) and (2.51) that

$$Nv \leq -k\omega\delta e^{-\delta t} + \delta^2 e^{-\delta t} + M\delta e^{-\delta t} \leq 0.$$

By the comparison principle, one gets that

$$V(t, x, y) \ge v^{-}(t, x, y)$$
, for $t \ge 0$ and $x \in \mathbb{R}^2$.

Then, the conclusion of Claim 2.10 follows immediately.

3 Uniqueness and stability of the curved front

This section is devoted to the proofs of uniqueness and stability of the curved front in Theorem 1.2, that is, Theorems 1.8 and 1.9.

3.1 Proof of Theorem 1.8

The idea of the proof of the uniqueness is inspired by Berestycki and Hamel [3] who proved that for any two almost-planar fronts $u_1(t, x, y)$ and $u_2(t, x, y)$, there is $T \in \mathbb{R}$ such that either $u_1(t + T, x, y) > u_2(t, x, y)$ or $u_1(t + T, x, y) = u_2(t, x, y)$.

Proof of Theorem 1.8. Assume that there is another curved front $V^*(t, x, y)$ satisfying (1.8). By (1.8), there is R > 0 large enough such that

$$V(t, x, y), \ V^*(t, x, y) \le \sigma \text{ for } (t, x, y) \in \omega^+ \text{ and } V(t, x, y), \ V^*(t, x, y) \ge 1 - \sigma \text{ for } (t, x, y) \in \omega^-,$$
 where σ is defined in (F3),

$$\omega_t^+ := \{ (t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \le 0 \text{ and } x \cos \alpha + y \sin \alpha - c_\alpha t \ge c_\alpha R \} \cup \{ (t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x > 0 \text{ and } x \cos \beta + y \sin \beta - c_\beta t \ge c_\beta R \},$$

and

$$\omega_t^- := \{ (t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x \le 0 \text{ and } x \cos \alpha + y \sin \alpha - c_\alpha t \le -c_\alpha R \} \cup \{ (t, x, y) \in \mathbb{R} \times \mathbb{R}^2; x > 0 \text{ and } x \cos \beta + y \sin \beta - c_\beta t \le -c_\beta R \}.$$

Since $c_{\alpha}/\sin \alpha = c_{\beta}/\sin \beta$, one knows that ω_t^+ and ω_t^- are connected. By a similar proof as of Lemma 2.5, there is $0 < \sigma' \le \sigma$ such that

$$\sigma' \leq V(t, x, y), \ V^*(t, x, y) \leq 1 - \sigma', \ \text{in } \mathbb{R} \times \mathbb{R}^2 \setminus (\omega_t^+ \cup \omega_t^-).$$

Then, by taking τ large enough, one has

$$V^*(t-\tau, x, y) \le \sigma' \le V(t, x, y)$$
, for $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2 \setminus (\omega_t^+ \cup \omega_t^-)$,

and

$$V(t, x, y) \ge 1 - \sigma' \ge V^*(t - \tau, x, y)$$
, for $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2 \setminus (\omega_{t-\tau}^+ \cup \omega_{t-\tau}^-)$,

It means that

$$V^*(t-\tau, x, y) \le V(t, x, y)$$
, in $\mathbb{R} \times \mathbb{R}^2 \setminus (\omega_{t-\tau}^+ \cup \omega_t^-)$.

Since f(x, y, u) is nonincreasing in $u \in (-\infty, \sigma]$ and $u \in [1 - \sigma, +\infty)$ for $(x, y) \in \mathbb{T}^2$ and by the same line of the proof of [3, Lemma 4.2], one can get that

$$V^*(t-\tau,x,y) \le V(t,x,y)$$
, for all $(t,x,y) \in \omega_{t-\tau}^-$ and $(t,x,y) \in \omega_t^+$.

and hence,

$$V^*(t-\tau, x, y) \le V(t, x, y)$$
, for all $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$ and τ large enough.

Now, we decrease τ and let

$$\tau_* = \inf\{\tau \in \mathbb{R}; V^*(t-\tau, x, y) \le V(t, x, y), \text{ for } (t, x, y) \in \mathbb{R} \times \mathbb{R}^2\}.$$

Since both V(t, x, y) and $V^*(t, x, y)$ satisfy (1.8), one knows that $\tau_* \ge 0$. Assume that $\tau_* > 0$. If

$$\inf\{V(t, x, y) - V^*(t - \tau_*, x, y); (t, x, y) \in \mathbb{R} \times \mathbb{R}^2 \setminus (\omega_{t - \tau_*}^+ \cup \omega_t^-)\} > 0,$$

then there is $\eta > 0$ such that

$$V^*(t-(\tau_*-\eta),x,y) \leq V(t,x,y), \text{ for } (t,x,y) \in \mathbb{R} \times \mathbb{R}^2 \setminus (\omega_{t-\tau_*}^+ \cup \omega_t^-).$$

By applying above arguments again, one can get that

$$V^*(t-(\tau_*-\eta),x,y) \leq V(t,x,y)$$
, for $(t,x,y) \in \mathbb{R} \times \mathbb{R}^2$,

which contradicts the definition of τ_* . Thus,

$$\inf\{V(t,x,y)-V^*(t-\tau_*,x,y);(t,x,y)\in\mathbb{R}\times\mathbb{R}^2\setminus(\omega_{t-\tau_*}^+\cup\omega_t^-)\}=0,$$

and there is a sequence $\{(t_n, x_n, y_n)\}_{n \in \mathbb{N}}$ such that

$$V(t_n, x_n, y_n) - V^*(t_n - \tau_*, x_n, y_n) \rightarrow 0$$
, as $n \rightarrow +\infty$.

Then, by following similar arguments as Step 3 of the proof of Lemma 2.8, one can get a contradiction. Thus, $\tau_* = 0$.

Therefore,

$$V(t, x, y) \ge V^*(t, x, y)$$
, for all $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$.

The same arguments can be applied by changing positions of V(t, x, y) and $V^*(t, x, y)$, and then, we can get that

$$V^*(t, x, y) \ge V(t, x, y)$$
, for all $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$.

In conclusion, $V^*(t, x, y) \equiv V(t, x, y)$.

3.2 Stability of the curved front

Take any $0 < \alpha < \beta < \pi$ such that Theorem 1.2 holds. Since $g'(\alpha) < 0$, one can take $\alpha_1 \in (0, \alpha)$ such that

$$\frac{c_{\theta}}{\sin \theta} > \frac{c_{\alpha}}{\sin \alpha}$$
, for $\theta \in [\alpha_1, \alpha]$.

Similar as Lemma 2.2, there is a smooth function $\varphi_1(x)$ with $y = -x \cot \alpha$ and $y = -x \cot \alpha_1$ being its asymptotic lines and there are positive constant k_3 , k_4 and K_4 such that

$$\begin{cases} \varphi_1''(x) < 0, & \text{for all } x \in \mathbb{R}, \\ -\cot \alpha > \varphi_1'(x) > -\cot \alpha_1, & \text{for all } x \in \mathbb{R}, \\ k_1 \operatorname{sech}(x) \le -\cot \alpha - \varphi_1'(x) \le K_4 \operatorname{sech}(x), & \text{for } x < 0, \\ k_2 \operatorname{sech}(x) \le \varphi_1'(x) + \cot \alpha_1 \le K_4 \operatorname{sech}(x), & \text{for } x \ge 0, \\ \max(|\varphi_1''(x)|, |\varphi_1'''(x)|) \le K_4 \operatorname{sech}(x), & \text{for all } x \in \mathbb{R}. \end{cases}$$

$$(3.1)$$

Take a constant ϱ which will be determined later. For every point (x, y) on the curve $y = \varphi_1(\varrho x)/\varrho$, there is a unit normal

$$e(x) = (e_1(x), e_2(x)) = \left(-\frac{\varphi_1'(\varrho x)}{\sqrt{\varphi_1'^2(\varrho x) + 1}}, \frac{1}{\sqrt{\varphi_1'^2(\varrho x) + 1}}\right).$$

For $(x, y) \in \mathbb{R}^2$ and $t \in \mathbb{R}$, take a constant ε and we define

$$U_1^-(t,x,y) = U_{e(x)}(\xi(t,x,y),x,y) + \varepsilon \mathrm{sech}(\varrho x),$$

where

$$\underline{\xi}(t,x,y) = \frac{y - c_{\alpha\beta}t - \varphi_1(\varrho x)/\varrho}{\sqrt{\varphi_1'^2(\varrho x) + 1}}.$$

Lemma 3.1 There exist ε_0 and $\varrho(\varepsilon_0)$ such that for any $0 < \varepsilon \le \varepsilon_0$ and $0 < \varrho \le \varrho(\varepsilon_0)$, the function $U_1^-(t, x, y)$ is a subsolution of (1.1). Moreover, it satisfies

$$\lim_{R \to +\infty} \sup_{x < -R} \left| U_1^-(t, x, y) - U_\alpha(x \cos \alpha + y \sin \alpha - c_\alpha t, x, y) \right| \le 2\varepsilon, \tag{3.2}$$

and

$$U_1^-(t, x, y) \le U_\alpha(x \cos \alpha + y \sin \alpha - c_\alpha t, x, y), \text{ for all } t \in \mathbb{R} \text{ and } (x, y) \in \mathbb{R}^2.$$
 (3.3)

Proof. Assume that

$$\varepsilon_0 \leq \frac{\sigma}{2}$$
,

where $\sigma > 0$ is defined in (F3). More restrictions on ε_0 will be given later. It follows from similar computation as Step 1 of the proof of Lemma 2.8 that

$$\begin{split} NU_{1}^{-} := & (U_{1}^{-})_{t} - \Delta_{x,y}U_{1}^{-} - f(x,y,U_{1}^{-}) \\ &= \partial_{\xi}U_{e(x)}\underline{\xi}_{t} - \partial_{\xi\xi}U_{e(x)}(\underline{\xi}_{x}^{2} + \underline{\xi}_{y}^{2}) - 2\nabla_{x,y}\partial_{\xi}U_{e(x)} \cdot (\underline{\xi}_{x},\underline{\xi}_{y}) - \Delta_{x,y}U_{e(x)} - \partial_{\xi}U_{e(x)}\underline{\xi}_{xx} \\ &- U_{e(x)}^{"} \cdot e'(x) \cdot e'(x) - U_{e(x)}^{'} \cdot e''(x) - 2\partial_{\xi}U_{e(x)}^{'} \cdot e'(x)\underline{\xi}_{x} - 2\partial_{x}U_{e(x)}^{'} \cdot e'(x) \\ &- \varepsilon\varrho^{2}\mathrm{sech}^{"}(\varrho x) - f(x,y,U_{1}^{-}) \end{split}$$

$$=(c_{e(x)} - \underline{\xi}_{t})\partial_{\xi}U_{e(x)} - \partial_{\xi\xi}U_{e(x)}(\underline{\xi}_{x}^{2} + \underline{\xi}_{y}^{2} - 1) - 2\partial_{x}\partial_{\xi}U_{e(x)}(\underline{\xi}_{x} - e_{1}(x)) - \partial_{\xi}U_{e(x)}\underline{\xi}_{xx}$$
$$- U_{e(x)}^{"} \cdot e'(x) \cdot e'(x) - U_{e(x)}^{'} \cdot e''(x) - 2\partial_{\xi}U_{e(x)}^{'} \cdot e'(x)\underline{\xi}_{x} - 2\partial_{x}U_{e(x)}^{'} \cdot e'(x)$$
$$- \varepsilon\rho^{2} \operatorname{sech}^{"}(\rho x) + f(x, y, U_{e(x)}) - f(x, y, U_{1}^{-}),$$

where $U_{e(x)}$, $\partial_{\xi}U_{e(x)}$, $\partial_{\xi\xi}U_{e(x)}$, $\nabla_{x,y}\partial_{\xi}U_{e(x)}$, $\Delta_{x,y}U_{e(x)}$, $U''_{e(x)}$, $e'(x)\cdot e'(x)$, $U'_{e(x)}\cdot e''(x)$, $\partial_{\xi}U'(e(x))\cdot e'(x)$, ∂

$$|\partial_{\xi\xi}U_{e(x)}(\xi_x^2 + \xi_y^2 - 1)| + 2|\partial_x\partial_\xi U_{e(x)}(\xi_x - e_1(x))| + |\partial_\xi U_{e(x)}\xi_{xx}| \le C_5\varrho \operatorname{sech}(\varrho x), \tag{3.4}$$

and

$$|U_{e(x)}'' \cdot e'(x) \cdot e'(x)| + |U_{e(x)}' \cdot e''(x)| + 2|\partial_{\xi}U_{e(x)}' \cdot e'(x)\xi_{x}| + 2|\partial_{x}U_{e(x)}' \cdot e'(x)| \le C_{6}\varrho \operatorname{sech}(\varrho x).$$
(3.5)

By a similar proof as of Claim 2.9, we can easily get that

$$C_{e(x)} - \underline{\xi}_t > 0$$
, for $x \in \mathbb{R}$.

and there is $C_7 > 0$ such that

$$c_{e(x)} - \underline{\xi}_t = c_{e(x)} - \frac{c_{\alpha\beta}}{\sqrt{\varphi_1'^2(\varrho x) + 1}} \ge C_7 \operatorname{sech}(\varrho x) > 0$$
, for x being negative enough.

Since $\varphi_1'(x)(\varrho x) \to -\cot \alpha_1$, $e(x) \to (\cos \alpha_1, \sin \alpha_1)$ as $x \to +\infty$ and $c_{\alpha_1}/\sin \alpha_1 > c_{\alpha}/\sin \alpha$, there is a constant c > 0 such that

$$c_{e(x)} - \underline{\xi}_t = c_{e(x)} - \frac{c_{\alpha\beta}}{\sqrt{\varphi_1'^2(\varrho x) + 1}} \ge c > 0, \text{ for all } x \ge 0.$$
 (3.6)

For x < 0, one can make similar arguments as in the proof of Lemma 2.8 to get that $NU_1^- \le 0$. For $x \ge 0$, one can get from (3.4), (3.5), (3.6) and Lemma 2.1 that

$$NU_{1}^{-} \le c\partial_{\xi}U_{e(x)} + (C_{5} + C_{6})\varrho \operatorname{sech}(\varrho x) + 2\varepsilon\varrho^{2}\operatorname{sech}(\varrho x) + f(x, y, U_{e(x)}) - f(x, y, U_{1}^{-}).$$
 (3.7)

For $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$ such that $\underline{\xi}(t, x, y) \ge C$ and $\underline{\xi}(t, x, y) \le -C$ where C is defined by (2.22), it follows from (F3) and $\varepsilon \le \varepsilon_0 \le \sigma/2$ that

$$f(x, y, U_{e(x)}) - f(x, y, U_1^-) \le -\lambda \varepsilon \operatorname{sech}(\varrho x).$$

Since $\partial_{\varepsilon}U_{e} < 0$, one has that

$$NU_1^- \le ((C_5 + C_6)\varrho + 2\varepsilon\varrho^2 - \lambda\varepsilon)\operatorname{sech}(\varrho x) \le 0,$$

by taking $\varrho(\varepsilon) > 0$ small enough such that

$$(C_5 + C_6)\rho + 2\varepsilon\rho^2 - \lambda\varepsilon < 0, (3.8)$$

and $0 < \varrho \le \varrho(\varepsilon)$. Finally, for $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$ such that $-C \le \xi(t, x, y) \le C$, there is k > 0 such that

$$-\partial_{\xi}U_{e}(\xi, x, y) \ge k \text{ for all } e \in \mathbb{S}.$$

Notice that

$$f(x, y, U_{\rho(x)}) - f(x, y, U^{+}) \leq M \varepsilon \operatorname{sech}(\rho x),$$

where $M := \max_{(x,y,u) \in \mathbb{T}^2 \times \mathbb{R}} |f_u(x,y,u)|$. Thus, it follows from (3.7) and (3.8) that

$$NU_1^- \le -kc + ((C_5 + C_6)\varrho + 2\varepsilon\varrho^2 + M\varepsilon)\operatorname{sech}(\varrho x) \le -kc + (\lambda + M)\varepsilon\operatorname{sech}(\varrho x) \le 0,$$

by taking $\varepsilon_0 = \min\{\sigma/2, kc/(\lambda + M)\}\$ and $0 < \varepsilon \le \varepsilon_0$.

By similar arguments as in Step 2 of the proof of Lemma 2.8, one gets that (3.2) holds. The inequality (3.3) can be gotten by comparing $U_1^-(t, x, y)$ with $U_\alpha(x \cos \alpha + y \sin \alpha - c_\alpha t, x, y)$ through similar arguments as in Step 3 of the proof of Lemma 2.8. This completes the proof. \square

Similarly, since $g'(\beta) > 0$, one can take $\beta_1 \in (\beta, \pi)$ such that

$$\frac{c_{\beta}}{\sin \beta} < \frac{c_{\theta}}{\sin \theta}$$
, for all $\theta \in (\beta, \beta_1]$.

Similar as Lemma 2.2, there is a smooth function $\varphi_2(x)$ with $y = -x \cot \beta$ and $y = -x \cot \beta_1$ being its asymptotic lines and there are positive constant k_5 , k_6 and K_5 such that

$$\begin{cases} \varphi_2''(x) < 0, & \text{for all } x \in \mathbb{R}, \\ -\cot \beta_1 < \psi_2'(x) > -\cot \beta, & \text{for all } x \in \mathbb{R}, \\ k_5 \mathrm{sech}(x) \le -\cot \beta_1 - \varphi_2'(x) \le K_5 \mathrm{sech}(x), & \text{for } x < 0, \\ k_6 \mathrm{sech}(x) \le \varphi_2'(x) + \cot \beta \le K_5 \mathrm{sech}(x), & \text{for } x \ge 0, \\ \max(|\varphi_2''(x)|, |\varphi_2'''(x)|) \le K_5 \mathrm{sech}(x), & \text{for all } x \in \mathbb{R}. \end{cases}$$

$$(3.9)$$

Take a constant ϱ which will be determined later. For every point (x, y) on the curve $y = \varphi_2(\varrho x)/\varrho$, there is a unit normal

$$e(x) = (e_1(x), e_2(x)) = \Big(-\frac{\varphi_2'(\varrho x)}{\sqrt{\varphi_2'^2(\varrho x) + 1}}, \frac{1}{\sqrt{\varphi_2'^2(\varrho x) + 1}} \Big).$$

For $(x, y) \in \mathbb{R}^2$ and $t \in \mathbb{R}$, take a constant ε and we define

$$U_2^-(t,x,y) = U_{e(x)}(\xi(t,x,y),x,y) + \varepsilon \mathrm{sech}(\varrho x),$$

where

$$\underline{\xi}(t,x,y) = \frac{y - c_{\alpha\beta}t - \varphi_2(\varrho x)/\varrho}{\sqrt{\varphi_2'^2(\varrho x) + 1}}.$$

Similar as Lemma 3.1, we can prove the following lemma.

Lemma 3.2 There exist ε_0 and $\varrho(\varepsilon_0)$ such that for any $0 < \varepsilon \le \varepsilon_0$ and $0 < \varrho \le \varrho(\varepsilon_0)$, the function $U_2^-(t, x, y)$ is a subsolution of (1.1). Moreover, it satisfies

$$\lim_{R \to +\infty} \sup_{t > R} \left| U_2^-(t, x, y) - U_{\beta}(x \cos \beta + y \sin \beta - c_{\beta}t, x, y) \right| \le 2\varepsilon,$$

and

$$U_2^-(t, x, y) \le U_\beta(x \cos \beta + y \sin \beta - c_\beta t, x, y), \text{ for all } t \in \mathbb{R} \text{ and } (x, y) \in \mathbb{R}^2.$$

Then, we need the following sub and supersolutions for the Cauchy problems of (1.1).

Lemma 3.3 For any function $u(t, x, y) \in C^{1,2}(\mathbb{R} \times \mathbb{R}^2)$, if it is a subsolution of (1.1) for $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$ with $u_t > 0$ and for any $0 < \sigma_1 < 1/2$ there is a positive constant k such that

$$u_t \ge k$$
, for $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$ such that $\sigma_1 \le u(t, x, y) \le 1 - \sigma_1$, (3.10)

then for any $0 < \delta < \sigma/2$ where σ is defined in (F3), there exist positive constants ω and λ such that

$$u(t, x, y) = u(t + \omega \delta e^{-\lambda t} - \omega \delta, x, y) - \delta e^{-\lambda t},$$

is a subsolution of (1.1) for $t \ge 0$ and $(x, y) \in \mathbb{R}^2$. Similarly, if u(t, x, y) is a smooth supersolution satisfying (3.10), then for any $0 < \delta < \sigma/2$, there exist positive constants ω and λ such that

$$\overline{u}(t, x, y) = u(t - \omega \delta e^{-\lambda t} + \omega \delta, x, y) + \delta e^{-\lambda t}$$

is a supersolution of (1.1) for $t \ge 0$ and $(x, y) \in \mathbb{R}^2$.

Proof. We only prove for the subsolution. Similar arguments can be applied for the supersolution. Take any $0 < \delta < \sigma/2$ where σ is defined in (F3). Let k > 0 such that $u_t \ge k$ for $(t, x, y) \in \mathbb{R} \times \mathbb{R}^2$ such that $\sigma \le u \le 1 - \sigma/2$. Take $\omega > 0$ such that

$$k\omega \geq \frac{\lambda + M}{\lambda}$$
,

where λ is defined in (F3) and $M := \max_{(x,y,u) \in \mathbb{T}^2 \times \mathbb{R}} |f_u(x,y,u)|$.

We then check that

$$N\underline{u} := \underline{u}_t - \Delta_{x,y}\underline{u} - f(x, y, \underline{u}) \le 0$$
, for $t > 0$ and $(x, y) \in \mathbb{R}^2$.

By computation, one can get that

$$N\underline{u} = -\omega\delta\lambda e^{-\lambda t}u_t + \delta\lambda e^{-\lambda t} + f(x, y, u(t + \omega\delta e^{-\lambda t} - \omega\delta, x, y)) - f(x, y, \underline{u}).$$

For t > 0 and $(x, y) \in \mathbb{R}^2$ such that $1 - \sigma/2 \le u(t + \omega \delta e^{-\lambda t}, x, y) \le 1$ and $0 \le u(t + \omega \delta e^{-\lambda t}, x, y) \le \sigma/2$ respectively, one has that $\underline{u}(t, x, y) \ge 1 - \sigma$ and $\underline{u}(t, x, y) \le \sigma$ respectively. Then, by (1.2), it follows that

$$f(x, y, u(t + \omega \delta e^{-\lambda t} - \omega \delta, x, y)) - f(x, y, \underline{u}) \le -\lambda \delta e^{-\lambda t}.$$

Thus, by $u_t > 0$, we have

$$Nu \le \delta \lambda e^{-\lambda t} - \lambda \delta e^{-\lambda t} \le 0.$$

For t > 0 and $(x, y) \in \mathbb{R}^2$ such that $\delta \le u(t + \omega \delta e^{-\lambda t}, x, y) \le 1 - \sigma/2$, one has that

$$Nu \leq -k\omega\delta\lambda e^{-\lambda t} + \delta\lambda e^{-\lambda t} + M\delta e^{-\lambda t} \leq 0$$
,

by the definition of ω .

This completes the proof.

Now, we are ready to prove the stability of the curved front of Theorem 1.2.

Proof of Theorem 1.9. Let

$$\underline{U}(t, x, y) = \max\{U_1^-(t + \omega \delta e^{-\lambda t} - \omega \delta, x, y) - \delta e^{-\lambda t}, U_2^-(t + \omega \delta e^{-\lambda t} - \omega \delta, x, y) - \delta e^{-\lambda t}\},$$

and

$$\overline{U}(t, x, y) = U^{+}(t - \omega \delta e^{-\lambda t} + \omega \delta, x, y) + \delta e^{-\lambda t},$$

where ω , δ and λ are defined in Lemma 3.3.

By (1.11), there is $R_{\delta} > 0$ such that

$$U_{\alpha\beta}^{-}(0,x,y) - \delta \le u_0(x,y) \le U_{\alpha\beta}^{-}(0,x,y) + \delta$$
, for $x^2 + y^2 > R_{\delta}^2$.

Then, it follows from Lemma 2.8, Lemma 3.1 and Lemma 3.2 that

$$\underline{U}(0, x, y) \le u_0(x, y) \le \overline{U}(0, x, y), \text{ for } x^2 + y^2 > R_{\delta}^2.$$

By the definition of $\psi(x)$, one has that $\psi(0) > 0$. Thus,

$$\xi(0, x, y) = \frac{y - \psi(\varrho x)/\varrho}{\sqrt{\psi'^2(\varrho x)} + 1} \to -\infty \text{ as } \varrho \to 0 \text{ for } x^2 + y^2 \le R_\delta^2,$$

which implies $U_{e(x)}(\xi(0,x,y),x,y) \to 1$ as $\varrho \to 0$ for $x^2 + y^2 \le R_\delta^2$. By taking ϱ small enough, one can make that

$$u_0(x, y) \le 1 \le U^+(0, x, y) + \delta = \overline{U}(0, x, y), \text{ for } x^2 + y^2 \le R_\delta^2.$$

Similarly, since $\varphi_1(0) < 0$ and $\varphi_2(0) < 0$, one can make that

$$u_0(x, y) \ge 0 \ge \underline{U}(0, x, y), \text{ for } x^2 + y^2 \le R_\delta^2,$$

by taking ϱ small enough. Notice that by taking ϱ small enough, δ can be arbitrary small. Thus, we have that

$$\underline{U}(0, x, y) \le u_0(x, y) \le \overline{U}(0, x, y)$$
, for all $(x, y) \in \mathbb{R}^2$.

On the other hand, by Lemma 2.5, one knows that $U_1^-(t, x, y)$, $U_2^-(t, x, y)$ and $U^+(t, x, y)$ satisfy (3.10). By Lemma 3.3 and the comparison principle, one can get that

$$\underline{U}(t, x, y) \le u(t, x, y) \le \overline{U}(t, x, y)$$
, for $t \ge 0$ and $(x, y) \in \mathbb{R}^2$.

Take a sequence $t_n = L_2 n/c_{\alpha\beta}$ where L_2 is the period of y. Then, $t_n \to +\infty$ as $n \to +\infty$. By parabolic estimates, the sequence $u_n(t,x,y) := u(t+t_n,x,y+L_2n)$ converges, locally uniformly in $\mathbb{R} \times \mathbb{R}^2$, to a solution $u_{\infty}(t,x,y)$ of (1.1). Since $U_1^-(t+t_n,x,y+L_2n) = U_1^-(t,x,y)$, $U_2^-(t+t_n,x,y+L_2n) = U_2^-(t,x,y)$ and $U^+(t+t_n,x,y+L_2n) = U^+(t,x,y)$, one has that

$$\max\{U_{1}^{-}(t + \omega \delta e^{-\lambda(t+t_{n})} - \omega \delta, x, y) - \delta e^{-\lambda(t+t_{n})}, U_{2}^{-}(t + \omega \delta e^{-\lambda(t+t_{n})} - \omega \delta, x, y) - \delta e^{-\lambda(t+t_{n})}\}$$

$$\leq u_{n}(t, x, y) \leq U^{+}(t - \omega \delta e^{-\lambda(t+t_{n})} + \omega \delta, x, y) + \delta e^{-\lambda(t+t_{n})},$$
(3.11)

and by passing $n \to +\infty$, $u_{\infty}(t, x, y)$ satisfies

$$\max\{U_1^-(t-\omega\delta,x,y),U_2^-(t-\omega\delta,x,y)\} \le u_\infty(t,x,y) \le U^+(t+\omega\delta,x,y), \text{ for } (t,x,y) \in \mathbb{R} \times \mathbb{R}^2.$$

Let $U_{12}^-(t, x, y) = \max\{U_1^-(t, x, y), U_2^-(t, x, y)\}$. Then, by Lemmas 3.1, 3.2 and similar arguments as Step 2 of the proof of Lemma 2.8, one can get that

$$\lim_{R \to +\infty} \sup_{x^2 + (y - c_{-\alpha}t)^2 > R^2} \left| U_{12}^{-}(t, x, y) - U_{\alpha\beta}^{-}(t, x, y) \right| \le 2\varepsilon.$$
 (3.12)

By Lemma 2.8, (3.12) and δ , ε can be taken arbitrary small, we have that

$$\lim_{R \to +\infty} \sup_{x^2 + (y - c_{\infty}t)^2 > R^2} \left| u_{\infty}(t, x, y) - U_{\alpha\beta}^{-}(t, x, y) \right| = 0,$$

By uniqueness of the curved front, we have $u_{\infty}(t, x, y) \equiv V(t, x, y)$. Thus, for any $\varepsilon > 0$, it follows from (1.8), (3.11), (3.12), Lemma 2.8 and taking δ small enough that there is $t_0 > 0$ large enough such that

$$|u(t_0, x, y) - V(t_0, x, y)| \le \varepsilon$$
, for all $(x, y) \in \mathbb{R}^2$.

On the other hand, by $V_t > 0$ and a similar proof as of Lemma 2.5, one knows that V(t, x, y) satisfies (3.10). By Lemma 3.3 again and the comparison principle, one gets that

$$V(t_0 + t + \omega \varepsilon e^{-\lambda t} - \omega \varepsilon, x, y) - \varepsilon e^{-\lambda t} \le u(t_0 + t, x, y) \le V(t_0 - \omega \varepsilon e^{-\lambda t} + \omega \varepsilon, x, y) + \varepsilon e^{-\lambda t},$$

for $t \ge 0$ and $(x, y) \in \mathbb{R}^2$. Then, since ε can be arbitrary small, one finally has that

$$u(t, x, y) \to V(t, x, y)$$
, as $t \to +\infty$ uniformly in $\mathbb{R} \times \mathbb{R}^2$.

This completes the proof.

4 A curved front with varying interfaces

In this section, we construct a curved front with varying interfaces. It behaves as three pulsating fronts as $t \to -\infty$ and as two pulsating fronts as $t \to +\infty$. We can not apply the idea of Hamel [16] by considering a Neumann boundary problem in the half plane x < 0 since our problem is not orthogonal symmetric with respect to y-axis in general.

Let α , β and θ satisfy Theorem 1.10. We will need the following properties.

Lemma 4.1 It holds that

$$c_{\alpha\theta}e_{\alpha\theta} = \left(\frac{c_{\alpha}\sin\theta - c_{\theta}\sin\alpha}{\sin(\theta - \alpha)}, \frac{c_{\alpha}\cos\theta - c_{\theta}\cos\alpha}{\sin(\alpha - \theta)}\right) := (c_{1}, c_{2}),$$

and

$$c_{\beta\theta}e_{\beta\theta} = \left(\frac{c_{\beta}\sin\theta - c_{\theta}\sin\beta}{\sin(\theta - \beta)}, \frac{c_{\beta}\cos\theta - c_{\theta}\cos\beta}{\sin(\beta - \theta)}\right) := (\hat{c}_{1}, \hat{c}_{2}),$$

with $c_1 > 0$ and $\hat{c}_1 < 0$. Moreover,

$$\frac{c_{\alpha}}{e_{\alpha\theta} \cdot (\cos\alpha, \sin\alpha)} = \frac{c_{\theta}}{e_{\alpha\theta} \cdot (\cos\theta, \sin\theta)} = c_{\alpha\theta} > \frac{c_{\theta_1}}{e_{\alpha\theta} \cdot (\cos\theta_1, \sin\theta_1)}, \text{ for any } \theta_1 \in (\alpha, \theta),$$

and

$$\frac{c_{\beta}}{e_{\beta\theta} \cdot (\cos\beta, \sin\beta)} = \frac{c_{\theta}}{e_{\beta\theta} \cdot (\cos\theta, \sin\theta)} = c_{\beta\theta} > \frac{c_{\theta_2}}{e_{\beta\theta} \cdot (\cos\theta_2, \sin\theta_2)}, \text{ for any } \theta_2 \in (\theta, \beta).$$

Proof. Assume by contradiction that $c_{\alpha\theta}e_{\alpha\theta} \neq (c_1, c_2)$. Take a sequence $\{t_n\}_{n\in\mathbb{N}}$ such that $t_n \to +\infty$. Then, for the sequence

$$(x_n, y_n) = (c_1, c_2)t_n$$

one has that $((x_n, y_n) - c_{\alpha\theta}e_{\alpha\theta}t_n)^2 \to +\infty$ as $n \to +\infty$ since $c_{\alpha\theta}e_{\alpha\theta} \neq (c_1, c_2)$. Notice that for any n, there are $k_n^1, k_n^2 \in \mathbb{Z}$ and $x_n', y_n' \in [0, L_2)$ such that $x_n = k_n^1 L_1 + x_n'$ and $y_n = k_n^2 L_2 + y_n'$. Moreover, up to extract subsequences of x_n and y_n , there are $x_n' \in [0, L_1]$ and $y_n' \in [0, L_2]$ such that $x_n' \to x_n'$ and $y_n' \to y_n'$ as $n \to +\infty$. Since $f(x, y, \cdot)$ is L-periodic in (x, y), one has $f(x + x_n, y + y_n, \cdot) \to f(x + x_n', y + y_n', \cdot)$ as $n \to +\infty$. Let $v_n(t, x, y) = V_{\alpha\theta}(t + t_n, x + x_n, y + y_n)$. By standard parabolic estimates, $v_n(t, x, y)$, up to extract of a subsequence, converges to a solution $v_\infty(t, x, y)$ of

$$v_t - \Delta v = f(x + x'_*, y + y'_*, v), \text{ for } t \in \mathbb{R} \text{ and } (x, y) \in \mathbb{R}^2.$$
 (4.1)

By definitions of x_n , y_n , c_1 and c_2 , one can also have that

$$U_{\alpha\theta}^-(t+t_n,x+x_n,y+y_n)\to \hat{U}_{\alpha\theta}^-(t,x,y)$$
, as $n\to +\infty$ uniformly in $\mathbb{R}\times\mathbb{R}^2$,

where

$$\hat{U}_{\alpha\theta}^{-}(t,x,y) := \max\{U_{\alpha}(x\cos\alpha + y\sin\alpha - c_{\alpha}t, x + x'_{\ast}, y + y'_{\ast}), U_{\theta}(x\cos\theta + y\sin\theta - c_{\theta}t, x + x'_{\ast}, y + y'_{\ast})\}.$$

Moreover, since $V_{\alpha\theta}(t, x, y)$ satisfies

$$\lim_{R\to+\infty} \sup_{((x,y)-c_{e_1e},te_{\alpha\theta})^2>R^2} \left| V_{\alpha\theta}(t,x,y) - U_{\alpha\theta}^-(t,x,y) \right| = 0,$$

one then gets that

$$v_n(t, x, y) \to \hat{U}_{\alpha\theta}^-(t, x, y)$$
 as $n \to +\infty$ locally uniformly in \mathbb{R}^2 .

In implies that $v_{\infty}(t, x, y) = \hat{U}_{\alpha\theta}^{-}(t, x, y)$ which is impossible since $\hat{U}_{\alpha\theta}^{-}(t, x, y)$ is not a solution of (4.1). Thus, $c_{\alpha\theta}e_{\alpha\theta} = (c_1, c_2)$. Similarly, one can prove that $c_{\beta\theta}e_{\beta\theta} = (\hat{c}_1, \hat{c}_2)$.

The signs of c_1 and \hat{c}_1 can be easily gotten from the facts $\alpha < \theta < \beta$ and $c_{\alpha}/\sin \alpha = c_{\beta}/\sin \beta > c_{\theta}/\sin \theta$.

By rotating the coordinate such that $e_{\alpha\theta}$ being the y-axis, then the speed of the pulsating front $U_{\theta_1}(x\cos\theta_1 + y\sin\theta_1 - c_{\theta_1}t, x, y)$ in direction $e_{\alpha\theta}$ can be denoted by

$$\frac{c_{\theta_1}}{e_{\alpha\theta}\cdot(\cos\theta_1,\sin\theta_1)}.$$

By Theorem 1.7, one has that

$$\frac{c_{\alpha}}{e_{\alpha\theta} \cdot (\cos\alpha, \sin\alpha)} = \frac{c_{\theta}}{e_{\alpha\theta} \cdot (\cos\theta, \sin\theta)} = c_{\alpha\theta} \text{ and } \frac{c_{\theta_1}}{e_{\alpha\theta} \cdot (\cos\theta_1, \sin\theta_1)} < c_{\alpha\theta}, \text{ for any } \theta_1 \in (\alpha, \theta).$$

This completes the proof.

Let $\varphi_1(x)$ be a smooth function such that there exist $a_1 < 0 < b_1$ such that

$$\varphi_1(x) = -x \cot \alpha$$
, for $x \le a_1, \varphi_1(x) = -x \cot \theta$, for $x \ge b_1$ and $\varphi_1''(x) > 0$ for $x \in (a_1, b_1)$.

Let $\varphi_2(x)$ be a smooth function such that there exist $a_2 < 0 < b_2$ such that

$$\varphi_2(x) = -x \cot \theta$$
, for $x \le a_2, \varphi_2(x) = -x \cot \beta$, for $x \ge b_2$ and $\varphi_2''(x) > 0$ for $x \in (a_2, b_2)$.

Let

$$\psi_1(t, x) = \varphi_1(x - c_1 t) + \rho \operatorname{sech}(x - c_1 t) + \rho \operatorname{sech}(x - \hat{c}_1 t),$$

and

$$\psi_2(t, x) = \varphi_2(x - \hat{c}_1 t) + \rho \operatorname{sech}(x - c_1 t) + \rho \operatorname{sech}(x - \hat{c}_1 t).$$

By $c_1 > 0$, $\hat{c}_1 < 0$ and making $|a_1|$, $|a_2|$, b_1 , b_2 large enough and ρ small enough, one can let $(\psi_1)_{xx} > 0$ for t negative enough and $x \le (c_1 + \hat{c}_1)t/2$ and $(\psi_2)_{xx} > 0$ for t negative enough and $x \ge (c_1 + \hat{c}_1)t/2$. Let

$$\psi(t,x) = \begin{cases} \psi_1(t,x), & \text{for } x \le (c_1 + \hat{c}_1)t/2, \\ \psi_2(t,x), & \text{for } x > (c_1 + \hat{c}_1)t/2. \end{cases}$$
(4.2)

Take a constant ϱ to be determined. For every point on the curve $y = \psi_1(\varrho t, \varrho x)$, there is a unit normal

$$e(t,x) = (e_1(t,x), e_2(t,x)) = \left(-\frac{(\psi_1)_x(\varrho t, \varrho x)}{\sqrt{(\psi_1)_x^2(\varrho t, \varrho x) + 1}}, \frac{1}{\sqrt{(\psi_1)_x^2(\varrho t, \varrho x) + 1}}\right).$$

For every point on the curve $y = \psi_2(\varrho t, \varrho x)$, there is a unit normal

$$\eta(t,x) = \left(-\frac{(\psi_2)_x(\varrho t,\varrho x)}{\sqrt{(\psi_2)_x^2(\varrho t,\varrho x)+1}}, \frac{1}{\sqrt{(\psi_2)_x^2(\varrho t,\varrho x)+1}}\right).$$

Take $\varepsilon > 0$ to be determined. For $t \in \mathbb{R}$ and $(x, y) \in \mathbb{R}^2$, define

$$\widetilde{U}^+(t,x,y) := \begin{cases} U_{e(t,x)}\left(\xi_1(t,x,y),x,y\right) + \varepsilon \mathrm{sech}(\varrho(x-c_1t)) + \varepsilon \mathrm{sech}(\varrho(x-\hat{c}_1t)), \text{ for } x \leq \frac{c_1+\hat{c}_1}{2}t, \\ U_{\eta(t,x)}\left(\xi_2(t,x,y),x,y\right) + \varepsilon \mathrm{sech}(\varrho(x-c_1t)) + \varepsilon \mathrm{sech}(\varrho(x-\hat{c}_1t)), \text{ for } x > \frac{c_1+\hat{c}_1}{2}t, \end{cases}$$

where

$$\xi_1(t, x, y) := \frac{y - c_2 t - \psi_1(\varrho t, \varrho x)/\varrho}{\sqrt{(\psi_1)_x^2 (\varrho t, \varrho x) + 1}} \text{ and } \xi_2(t, x, y) := \frac{y - \hat{c}_2 t - \psi_2(\varrho t, \varrho x)/\varrho}{\sqrt{(\psi_2)_x^2 (\varrho t, \varrho x) + 1}}.$$

By the definition of $\psi_1, \psi_2, c_1, c_2, \hat{c}_1$ and \hat{c}_2 , one can easily check that around $x = (c_1 + \hat{c}_1)t/2$,

$$U_{e(t,x)}(\xi_1(t, x, y), x, y) = U_{\theta}(x \cos \theta + y \sin \theta - c_{\theta}t, x, y) = U_{\eta(t,x)}(\xi_2(t, x, y), x, y),$$

for t negative enough. Thus, $\widetilde{U}^+(t, x, y)$ is smooth for t negative enough and $(x, y) \in \mathbb{R}^2$.

Lemma 4.2 There exist ε_0 and $\varrho(\varepsilon_0)$ such that for any $0 < \varepsilon \le \varepsilon_0$ and $0 < \varrho \le \varrho(\varepsilon_0)$, the function $\widetilde{U}^+(t,x,y)$ is a supersolution of (1.1) for t negative enough. Moreover, it satisfies

$$\lim_{R \to +\infty} \sup_{x \le 0, ((x,y) - c_{\alpha\theta} e_{\alpha\theta} t)^2 > R^2} \left| \widetilde{U}^+(t,x,y) - U^-_{\alpha\theta}(t,x,y) \right| \le 2\varepsilon, \tag{4.3}$$

$$\lim_{R \to +\infty} \sup_{x > 0, ((x, y) - \epsilon_{R\theta} e_{\theta\theta} t)^2 > R^2} \left| \widetilde{U}^+(t, x, y) - U^-_{\theta\beta}(t, x, y) \right| \le 2\varepsilon, \tag{4.4}$$

and

$$\widetilde{U}^{+}(t, x, y) \geq \max\{U_{\alpha}(x \cos \alpha + y \sin \alpha - c_{\alpha}t, x, y), U_{\theta}(x \cos \theta + y \sin \theta - c_{\theta}t, x, y), U_{\beta}(x \cos \beta + y \sin \beta - c_{\beta}t, x, y)\}, \text{ for } t \text{ negative enough and } (x, y) \in \mathbb{R}^{2}.$$

$$(4.5)$$

Proof. We only prove for the part $x \le (c_1 + \hat{c}_1)t/2$. Take $0 < \varepsilon_0 \le \sigma/2$ and more restrictions on ε_0 will be given later. Change variables $X = x - c_1 t$ and $Y = y - c_2 t$. Then,

$$\psi_1(t, X) := \varphi_1(X) + \rho \operatorname{sech}(X) + \rho \operatorname{sech}(X + (c_1 - \hat{c}_1)t),$$

and

$$e(t,X) = (e_1(t,X), e_2(t,X)) = \left(-\frac{(\psi_1)_X(\varrho t, \varrho X)}{\sqrt{(\psi_1)_X^2(\varrho t, \varrho X) + 1}}, \frac{1}{\sqrt{(\psi_1)_X^2(\varrho t, \varrho X) + 1}}\right),$$

where $(\psi_1)_X$ is taking values at $(\varrho t, \varrho X)$. One can compute that

$$(\psi_1)_t(t,X) = (c_1 - \hat{c}_1)\operatorname{sech}(X + (c_1 - \hat{c}_1)t),$$

$$(\psi_1)_X(t,X) = \varphi_1'(X) + \operatorname{sech}'(X) + \operatorname{sech}'(X + \varrho(c_1 - \hat{c}_1)t),$$

$$(\psi_1)_{tX}(t,X) = (c_1 - \hat{c}_1)\operatorname{sech}'(X + (c_1 - \hat{c}_1)t),$$

$$(\psi_1)_{XX}(t,X) = \varphi_1''(X) + \operatorname{sech}''(X) + \operatorname{sech}''(X + (c_1 - \hat{c}_1)t),$$

$$(\psi_1)_{XXX}(t,X) = \varphi_1'''(X) + \operatorname{sech}'''(X) + \operatorname{sech}'''(X + (c_1 - \hat{c}_1)t).$$

$$e_{t} = \left(-\frac{\varrho(\psi_{1})_{tX}}{((\psi_{1})_{X}^{2}+1)^{3/2}}, -\frac{\varrho(\psi_{1})_{X}(\psi_{1})_{tX}}{((\psi_{1})_{X}^{2}+1)^{3/2}}\right),$$

$$e_{X} = \left(-\frac{\varrho(\psi_{1})_{XX}}{((\psi_{1})_{X}^{2}+1)^{\frac{3}{2}}}, -\frac{\varrho(\psi_{1})_{X}(\psi_{1})_{XX}}{((\psi_{1})_{X}^{2}+1)^{\frac{3}{2}}}\right),$$

and

$$e_{XX} = \left(-\frac{\varrho^2(\psi_1)_{XXX}}{((\psi_1)_X^2+1)^{\frac{3}{2}}} + \frac{3\varrho^2(\psi_1)_X(\psi_1)_{XX}^2}{((\psi_1)_X^2+1)^{\frac{5}{2}}}, -\frac{\varrho^2(\psi_1)_{XX}^2}{((\psi_1)_X^2+1)^{\frac{3}{2}}} - \frac{\varrho^2(\psi_1)_X(\psi_1)_{XXX}}{((\psi_1)_X^2+1)^{\frac{3}{2}}} + \frac{3\varrho^2(\psi_1)_X^2(\psi_1)_{XX}^2}{((\psi_1)_X^2+1)^{\frac{5}{2}}}\right),$$

where $(\psi_1)_X$, $(\psi_1)_{XX}$, $(\psi_1)_{XXX}$, $(\psi_1)_{tX}$ are taking values at $(\varrho t, \varrho X)$ in e_t , e_X , e_{XX} . Let

$$\widetilde{U}^+(t,X,Y) = U_{e(t,X)}\left(\xi_1(t,X,Y),X + c_1t,Y + c_2t\right) + \varepsilon \mathrm{sech}(\varrho X) + \varepsilon \mathrm{sech}(\varrho X - \varrho(c_1 - \hat{c}_1)t),$$

where

$$\xi_1(t, X, Y) = \frac{Y - \psi_1(\varrho t, \varrho X)/\varrho}{\sqrt{(\psi_1)_x^2(\varrho t, \varrho X) + 1}}.$$

We need to verify that

$$N\widetilde{U}^{+} := \widetilde{U}_{t}^{+} - \Delta_{X,Y}\widetilde{U}^{+} - c_{1}\widetilde{U}_{X}^{+} - c_{2}\widetilde{U}_{Y}^{+} - f(X + c_{1}t, Y + c_{2}t, \widetilde{U}^{+}) \ge 0,$$

for t negative enough and $(x, y) \in \mathbb{R}^2$. By (2.2) and after some computation, one can get that

$$\begin{split} N\widetilde{U}^{+} = & \partial_{\xi} U_{e(t,X)}((\xi_{1})_{t} - c_{1}(\xi_{1})_{X} - c_{2}(\xi_{1})_{Y} + c_{e(t,X)}) + U'_{e(t,X)} \cdot e_{t} - U''_{e(t,X)} \cdot e_{X} \cdot e_{X} - U'_{e(t,X)} \cdot e_{XX} \\ & - 2\partial_{\xi} U'_{e(t,X)} \cdot e_{X}(\xi_{1})_{X} - 2\partial_{X} U'_{e(t,X)} \cdot e_{X} - \partial_{\xi\xi} U_{e(t,X)}((\xi_{1})_{X}^{2} + (\xi_{1})_{Y}^{2} - 1) \\ & - 2\partial_{\xi} \partial_{X} U_{e(t,X)}((\xi_{1})_{X} - e_{1}(t,X)) - 2\partial_{\xi} \partial_{Y} U_{e(t,X)}((\xi_{1})_{Y} - e_{2}(t,X)) - \partial_{\xi} U_{e(t,X)}\xi_{XX} \\ & - c_{1} U'_{e(t,X)} \cdot e_{X} - \varepsilon \varrho^{2} \mathrm{sech}''(\varrho X) - \varepsilon \varrho^{2} \mathrm{sech}''(\varrho X - \varrho(c_{1} - \hat{c}_{1})t) - c_{1}\varepsilon\varrho \mathrm{sech}'(\varrho X) \\ & - c_{1}\varepsilon\varrho'(\varrho X - \varrho(c_{1} - \hat{c}_{1})t) + f(X + c_{1}t, Y + c_{2}t, U_{e(t,X)}) - f(X + c_{1}t, Y + c_{2}t, \widetilde{U}^{+}), \end{split}$$

where $\partial_{\xi}U_{e(t,X)}$, $\partial_{\xi\xi}U_{e(t,X)}$, $\nabla_{X,Y}\partial_{\xi}U_{e(t,X)}$, $U'_{e(t,X)} \cdot e_t$, $U''_{e(t,X)} \cdot e_X \cdot e_X$, $U'_{e(t,X)} \cdot e_{XX}$, $\partial_{\xi}U'_{e(t,X)} \cdot e_X$, $\partial_{\xi}U'_{e(t,X)} \cdot e_X$, $U'_{e(t,X)} \cdot e_X$,

$$(\xi_{1})_{t} = -\frac{\varrho(\psi_{1})_{X}(\psi_{1})_{tX}}{(\psi_{1})_{X}^{2} + 1} \xi_{1} - \frac{(\psi_{1})_{t}}{\sqrt{(\psi_{1})_{X}^{2} + 1}},$$

$$(\xi_{1})_{X} = -\frac{\varrho(\psi_{1})_{X}(\psi_{1})_{XX}}{((\psi_{1})_{X}^{2} + 1)^{\frac{1}{2}}} \xi_{1} - \frac{(\psi_{1})_{X}}{\sqrt{(\psi_{1})_{X}^{2} + 1}},$$

$$(\xi_{1})_{Y} = \frac{1}{\sqrt{(\psi_{1})_{X}^{2} + 1}},$$

$$(\xi_{1})_{XX} = -\frac{\varrho^{2}(\psi_{1})_{XX}^{2} + \varrho^{2}(\psi_{1})_{X}(\psi_{1})_{XX}}{(\psi_{1})_{X}^{2} + 1} \xi_{1} + \frac{3\varrho^{2}(\psi_{1})_{X}^{2}(\psi_{1})_{XX}^{2}}{((\psi_{1})_{X}^{2} + 1)^{2}} \xi_{1} + \frac{\varrho((\psi_{1})_{X}^{2} - 1)(\psi_{1})_{XX}}{((\psi_{1})_{X}^{2} + 1)^{\frac{3}{2}}},$$

$$(\xi_{1})_{X}^{2} + (\xi_{1})_{Y}^{2} - 1 = \left(\frac{\varrho(\psi_{1})_{X}(\psi_{1})_{XX}}{((\psi_{1})_{X}^{2} + 1)^{\frac{3}{2}}} \xi_{1}^{2} + 2\left(\frac{\varrho(\psi_{1})_{X}(\psi_{1})_{XX}}{((\psi_{1})_{X}^{2} + 1)^{\frac{3}{2}}}\right) \frac{(\psi_{1})_{X}}{\sqrt{(\psi_{1})_{X}^{2} + 1}} \xi_{1}\right),$$

where $(\psi_1)_X$, $(\psi_1)_t$ $(\psi_1)_{XX}$, $(\psi_1)_{tX}$ are taking values at $(\varrho t, \varrho X)$. By Lemma 2.1, Lemma 2.4, Lemma 2.7, boundedness of $||U'_e||$, $||U''_e||$, $||\partial_{\xi}U'_e||$, $||\partial_{x}U'_e||$ and above formulas, there are constants $C_8 > 0$ and $C_9 > 0$ such that

$$\begin{aligned} |\partial_{\xi\xi} U_{e(t,X)}((\xi_1)_X^2 + (\xi_1)_Y^2 - 1)| + 2|\partial_X \partial_{\xi} U_{e(t,X)}((\xi_1)_X - e_1(t,X))| + |\partial_{\xi} U_{e(t,X)}(\xi_1)_{XX}| \\ &\leq C_8 \rho(\operatorname{sech}(\rho X) + \operatorname{sech}(\rho X - \rho(c_1 - \hat{c}_1)t)), \end{aligned}$$

and

$$\begin{split} |U'_{e(t,X)} \cdot e_t| + |U''_{e(t,X)} \cdot e_X \cdot e_X| + |U'_{e(t,X)} \cdot e_{XX}| + 2|\partial_\xi U'_{e(t,X)} \cdot e_X(\xi_1)_X| + 2|\partial_x U'_{e(t,X)} \cdot e_X| \\ + c_1|U'_{e(t,X)} \cdot e_X| &\leq C_9 \varrho(\text{sech}(\varrho X) + \text{sech}(\varrho X - \varrho(c_1 - \hat{c}_1)t)), \end{split}$$

Therefore, it follows that

$$\begin{split} N\widetilde{U}^{+} \geq & \partial_{\xi} U_{e(t,X)}((\xi_{1})_{t} - c_{1}(\xi_{1})_{X} - c_{2}(\xi_{1})_{Y} + c_{e(t,X)}) - (C_{8} + C_{9})\varrho(\operatorname{sech}(\varrho X) \\ & + \varrho \operatorname{sech}(\varrho X + \varrho(c_{1} - \hat{c}_{1})t)) - (1 + c_{1})\varepsilon\varrho^{2}(\operatorname{sech}(\varrho X) + \varrho \operatorname{sech}(\varrho X + \varrho(c_{1} - \hat{c}_{1})t)) \\ & + f(X + c_{1}t, Y + c_{2}t, U_{e(t,X)}) - f(X + c_{1}t, Y + c_{2}t, \widetilde{U}^{+}) \end{split}$$

We claim that

Claim 4.3 There exist positive constants C_{10} and C_{11} such that

$$c_{1}(\xi_{1})_{X} + c_{2}(\xi_{1})_{Y} - (\xi_{1})_{t} - c_{e(t,X)} \ge -C_{10}\varrho(\operatorname{sech}(\varrho X) + \operatorname{sech}(\varrho X + \varrho(c_{1} - \hat{c}_{1})t))|\xi_{1}| - C_{10}\varrho(\operatorname{sech}(\varrho X + \varrho(c_{1} - \hat{c}_{1})t)) + C_{11}(\operatorname{sech}(\varrho X) + \operatorname{sech}(\varrho X + \varrho(c_{1} - \hat{c}_{1})t)).$$

In order to not lengthen the proof, we postpone the proof of Claim 4.3 after the proof of this lemma.

For $\xi_1(t, X, Y) \ge C$ and $\xi_1(t, X, Y) \le -C$ where C is defined by (2.22), it follows from (1.2) that

$$f(X + c_1t, Y + c_2t, U_{e(t,x)}) - f(X + c_1t, Y + c_2t, \widetilde{U}^+) \ge \lambda \varepsilon (\operatorname{sech}(\varrho x) + \operatorname{sech}(\varrho x + \varrho(c_1 - \hat{c}t)))$$

Then, by $\partial_{\varepsilon}U_{e} < 0$, Lemma 2.4 and Claim 4.3, it follows that

$$\begin{split} N\widetilde{U}^{+} &\geq -B_1C_{10}\varrho(\operatorname{sech}(\varrho X) + \operatorname{sech}(\varrho X + \varrho(c_1 - \hat{c}_1)t)) - B_2C_{10}\varrho\operatorname{sech}(\varrho X + \varrho(c_1 - \hat{c}_1)t)) \\ &- \Big((C_8 + C_9)\varrho + (1 + c_1)\varepsilon\varrho^2 \Big) (\operatorname{sech}(\varrho X) + \varrho\operatorname{sech}(\varrho X + \varrho(c_1 - \hat{c}_1)t)) \\ &+ \lambda\varepsilon(\operatorname{sech}(\varrho X) + \operatorname{sech}(\varrho X + \varrho(c_1 - \hat{c}t))) \geq 0 \end{split}$$

where $B_1 = \sup_{e \in \mathbb{S}} \|\partial_{\xi} U_e \xi\|_{L^{\infty}}$ and $B_2 = \sup_{e \in \mathbb{S}} \|\partial_{\xi} U_e\|_{L^{\infty}}$, by taking $0 < \varrho \leq \varrho(\varepsilon)$ where $\varrho(\varepsilon)$ is small enough such that

$$-B_1C_{10}\varrho - B_2C_{10}\varrho - \left((C_8 + C_9)\varrho + (1 + c_1)\varepsilon\varrho^2\right) + \lambda\varepsilon > 0, \text{ for all } 0 < \varrho \le \varrho(\varepsilon). \tag{4.7}$$

For $-C \le \xi_1(t, x, y) \le C$, there is k > 0 such that $-\partial_{\xi} U_{e(t, x)}(\xi_1(t, x, y), x, y) \ge k$. Then, it follows from Claim 4.3 that

$$\begin{split} N\widetilde{U}^{+} \geq & kC_{11}(\operatorname{sech}(\varrho X) + \operatorname{sech}(\varrho X + \varrho(c_{1} - \hat{c}_{1})t)) - B_{1}C_{10}\varrho(\operatorname{sech}(\varrho X) + \operatorname{sech}(\varrho X + \varrho(c_{1} - \hat{c}_{1})t)) \\ & - B_{2}C_{10}\varrho\operatorname{sech}(\varrho X + \varrho(c_{1} - \hat{c}_{1})t)) - \Big((C_{8} + C_{9})\varrho + (1 + c_{1})\varepsilon\varrho^{2}\Big)(\operatorname{sech}(\varrho X) \\ & + \varrho\operatorname{sech}(\varrho X + \varrho(c_{1} - \hat{c}_{1})t)) - M\varepsilon(\operatorname{sech}(\varrho X) + \operatorname{sech}(\varrho X + \varrho(c_{1} - \hat{c}t))) \geq 0 \end{split}$$

where $M = \max_{(x,y,u)\in\mathbb{R}^2\times\mathbb{R}} |f_u(x,y,u)|$, by (4.7), taking $0 < \varepsilon \le \varepsilon_0$ and $\varepsilon_0 = \max\{\sigma/2, kC_{11}/(\lambda + M)\}$.

By the comparison principle, $\widetilde{U}^+(t, x, y)$ is a supersolution of (1.1).

By the definition of $\psi_1(x)$, $\psi_2(x)$ and Lemma 4.1, one has that

$$\xi_1(t, X, Y) \to X \cos \alpha + Y \sin \alpha = x \cos \alpha + y \sin \alpha - c_{\alpha} t \text{ as } X \to -\infty$$

and

$$\xi_1(t, X, Y) \to X \cos \theta + Y \sin \theta = x \cos \theta + y \sin \theta - c_{\theta} t \text{ as } X \to +\infty.$$

Then, by similar arguments as in Step 2 of the proof of Lemma 2.8, one can get (4.3) and (4.4). The inequality (4.5) can be gotten by comparing $\widetilde{U}^+(t, x, y)$ with $U_{\alpha}(x \cos \alpha + y \sin \alpha - c_{\alpha}t, x, y)$, $U_{\beta}(x \cos \beta + y \sin \beta - c_{\beta}t, x, y)$, $U_{\theta}(x \cos \theta + y \sin \theta - c_{\theta}t, x, y)$ respectively for t negative enough through similar arguments as in Step 3 of the proof of Lemma 2.8. This completes the proof. \square

We then prove Claim 4.3.

Proof of Claim 4.3. From (4.6), one has that

$$\begin{split} c_1(\xi_1)_X + c_2(\xi_1)_Y - (\xi_1)_t - c_{e(t,X)} &= -c_1 \frac{\varrho(\psi_1)_X(\psi_1)_{XX}}{((\psi_1)_X^2 + 1)^{\frac{1}{2}}} \xi_1 - c_1 \frac{(\psi_1)_X}{\sqrt{(\psi_1)_X^2 + 1}} + c_2 \frac{1}{\sqrt{(\psi_1)_X^2 + 1}} \\ &+ \frac{\varrho(\psi_1)_X(\psi_1)_{tX}}{(\psi_1)_X^2 + 1} \xi_1 + \frac{(\psi_1)_t}{\sqrt{(\psi_1)_X^2 + 1}} - c_{e(t,X)}. \end{split}$$

Then, by Lemma 2.1 and the definition of ψ_1 , there is $C_{10} > 0$ such that

$$\left| -c_1 \frac{\varrho(\psi_1)_X(\psi_1)_{XX}}{((\psi_1)_Y^2 + 1)^{\frac{1}{2}}} \xi_1 + \frac{\varrho(\psi_1)_X(\psi_1)_{tX}}{(\psi_1)_X^2 + 1} \xi_1 \right| \le C_{10}\varrho(\operatorname{sech}(\varrho X) + \operatorname{sech}(\varrho X + \varrho(c_1 - \hat{c}_1)t))|\xi_1|, \quad (4.8)$$

and

$$\left| \frac{(\psi_1)_t}{\sqrt{(\psi_1)_X^2 + 1}} \right| \le C_{10} \varrho \operatorname{sech}(\varrho X + \varrho(c_1 - \hat{c}_1)t). \tag{4.9}$$

Let $\theta(t, X) = \arccos(e_1(t, X))$. Then, $e(t, X) = (\cos \theta(t, X), \sin \theta(t, X))$. By the definition of $\psi_1(t, X)$, one has $\alpha < \theta(t, X) < \theta$. It follows from Lemma 4.1 that

$$-c_{1} \frac{(\psi_{1})_{X}}{\sqrt{(\psi_{1})_{X}^{2}+1}} + c_{2} \frac{1}{\sqrt{(\psi_{1})_{X}^{2}+1}} - c_{e(t,X)} = (c_{1}, c_{2})(\cos\theta(t, X), \sin\theta(t, X)) - c_{\theta(t,X)}$$

$$= c_{\alpha\theta}e_{\alpha\theta}(\cos\theta(t, X), \sin\theta(t, X)) - c_{\theta(t,X)} > 0.$$
(4.10)

Notice that $c_e > 0$ for all $e \in \mathbb{S}$. By Lemma 4.1, one has that

 $e_{\alpha\theta} \cdot (\cos \theta(t, X), \sin \theta(t, X)) > 0$, for all $X \in \mathbb{R}$.

Let

$$h(s) = \frac{c_s}{e_{col} \cdot (\cos s, \sin s)}.$$

Notice that $h(\alpha) = c_{\alpha\beta}$. Also notice that $e_1(t, X) \to \cos \alpha$ as $X \to -\infty$ and $\theta(t, X) \to \alpha$ as $X \to -\infty$ for X being very negative, one has that

$$c_{\alpha\theta}e_{\alpha\theta}(\cos\theta(t,X),\sin\theta(t,X)) - c_{\theta(t,X)}$$

$$=e_{\alpha\theta} \cdot (\cos\theta(t,X),\sin\theta(t,X))(h(\alpha) - h(\theta(t,X)))$$

$$=e_{\alpha\theta} \cdot (\cos\theta(t,X),\sin\theta(t,X))(h'(\alpha)(\alpha - \theta(t,X)) + o(|\alpha - \theta(t,X)|))$$
(4.11)

Remember that $h'(\alpha) < 0$ by the assumptions of Theorem 1.10. Moreover, by the formulas in the proof of Lemma 4.2, there is $C_{11} > 0$ such that

$$\alpha - \theta(t, X) = \int_{-\infty}^{X} \theta_X(t, s) ds = \int_{-\infty}^{X} \frac{\varrho(\psi_1)_{XX}(\varrho t, \varrho s)}{(\psi_1)_X^2(\varrho t, \varrho s) + 1} ds$$

$$\geq \frac{1}{\|(\psi_1)_X\|_{L^{\infty}}^2 + 1} ((\psi_1)_X(\varrho t, \varrho X) + \cot \alpha) \geq C_{11}(\operatorname{sech}(\varrho X) + \operatorname{sech}(\varrho X + \varrho(c_1 - \hat{c}t))). \tag{4.12}$$

By (4.8)-(4.12), we have our conclusion.

Now, we turn to prove Theorem 1.10.

Proof of Theorem 1.10. Let $u_n(t, x, y)$ be the solution of (1.1) for $t \ge -n$ with initial data

$$u_n(-n, x, y) = U_{\alpha\theta\beta}^-(-n, x, y),$$

where

$$U_{\alpha\theta\beta}^{-}(t,x,y) = \max\{U_{\alpha}(x\cos\alpha + y\sin\alpha - c_{\alpha}t, x, y), U_{\theta}(x\cos\theta + y\sin\theta - c_{\theta}t, x, y), U_{\theta}(x\cos\beta + y\sin\beta - c_{\theta}t, x, y)\}.$$

$$U_{\beta}(x\cos\beta + y\sin\beta - c_{\beta}t, x, y)\}.$$

By Lemma 4.2, it follows from the comparison principle that

$$U_{\alpha\beta\beta}^{-}(t,x,y) \le u_n(t,x,y) \le \widetilde{U}^{+}(t,x,y), \text{ for } -n \le t \le T \text{ and } (x,y) \in \mathbb{R}^2, \tag{4.13}$$

wherer T is a negative constant such that Lemma 4.2 holds for $-\infty < t \le T$. Since $U_{\alpha\theta\beta}^-(t,x,y)$ is a subsolution, the sequence $u_n(t,x,y)$ is increasing in n. Letting $n \to +\infty$ and by parabolic estimates, the sequence $u_n(t,x,y)$ converges to an entire solution u(t,x,y) of (1.1).

By (4.13), u(t, x, y) satisfies

$$U_{\alpha\theta\beta}^{-}(t,x,y) \le u(t,x,y) \le \widetilde{U}^{+}(t,x,y), \text{ for } t \le T \text{ and } (x,y) \in \mathbb{R}^{2}.$$

$$(4.14)$$

Moreover, by (4.3), (4.4) and since ε can be arbitrary small, one can get that u(t, x, y) satisfies

$$\lim_{R \to +\infty} \sup_{x \le 0, ((x,y) - c_{\alpha\theta} e_{\alpha\theta} t)^2 > R^2} |u(t,x,y) - U_{\alpha\theta}^{-}(t,x,y)| = 0, \tag{4.15}$$

and

$$\lim_{R \to +\infty} \sup_{x > 0, ((x,y) - c_{\theta\theta} e_{\theta\theta} t)^2 > R^2} |u(t,x,y) - U_{\beta\theta}^{-}(t,x,y)| = 0, \tag{4.16}$$

for t negative enough. Now, we consider the half plane $H:=\{(x,y)\in\mathbb{R}^2;x<0\}$. Take any sequence $\{t_n\}_{n\in\mathbb{N}}$ of \mathbb{R} such that $t_n\to-\infty$ as $n\to+\infty$. Notice that for any n, there are $k_n^1,\,k_n^2\in\mathbb{Z}$ and $x_n'\in[0,L_1),\,y_n'\in[0,L_2)$ such that $c_1t_n=k_n^1L_1+x_n'$ and $c_2t_n=k_n^2L_2+y_n'$. Moreover, up to extract subsequences of c_1t_n and c_2t_n , there are $x_*'\in[0,L_1]$ and $y_*'\in[0,L_2]$ such that $x_n'\to x_*'$ and $y_n'\to y_*'$ as $n\to+\infty$. Let $v_n(t,x,y)=u(t+t_n,x+c_1t_n,y+c_2t_n)$ and $H_n=H-c_1t_n$. Then, $H_n\to\mathbb{R}^2$ as $n\to+\infty$. Since $f(x,y,\cdot)$ is L-periodic in (x,y), one has that $f(x+c_1t_n,y+c_2t_n,\cdot)\to f(x+x_*',y+y_*',\cdot)$ By parabolic estimates, $v_n(t,x,y)$, up to extract of a subsequence, converges to a solution $v_\infty(t,x,y)$ of

$$v_t - \Delta v = f(x + x'_*, y + y'_*, v), \quad (t, x, y) \in \mathbb{R} \times \mathbb{R}^2.$$
 (4.17)

By the definitions of c_1 and c_2 , one can easily check that

$$U_{\alpha\theta}^-(t+t_n, x+x_n, y+y_n) \to \hat{U}_{\alpha\theta}^-(t, x, y), \text{ as } n \to +\infty \text{ uniformly in } \mathbb{R} \times \mathbb{R}^2,$$
 (4.18)

where

 $\hat{U}_{\alpha\theta}^{-}(t, x, y) := \max\{U_{\alpha}(x\cos\alpha + y\sin\alpha - c_{\alpha}t, x + x'_{*}, y + y'_{*}), U_{\theta}(x\cos\theta + y\sin\theta - c_{\theta}t, x + x'_{*}, y + y'_{*})\}.$

By (4.15), it follows that

$$\lim_{R \to +\infty} \sup_{((x,y)-c_{\alpha\theta}e_{\alpha\theta}t)^2 > R^2} \left| v_{\infty}(t,x,y) - \hat{U}_{\alpha\theta}^-(t,x,y) \right| = 0.$$

By uniqueness of the curved front, one then has that $v_{\infty}(t, x, y) \equiv \hat{V}_{\alpha\theta}(t, x, y)$ where $\hat{V}_{\alpha\theta}(t, x, y)$ is the curved front of (4.17) satisfying

$$\lim_{R \to +\infty} \sup_{((x,y) - c_{\alpha\theta} e_{\alpha\theta} t)^2 > R^2} \left| \hat{V}_{\alpha\theta}(t,x,y) - \hat{U}_{\alpha\theta}^{-}(t,x,y) \right| = 0.$$
 (4.19)

Thus, for any fixed t,

$$v_n(t, x, y) \to \hat{V}_{\alpha\theta}(t, x, y)$$
, as $n \to +\infty$ locally uniformly in H_n .

By (4.15), (4.18) and (4.19), the above convergence is uniform in $\overline{H_n}$. Thus, for any fixed t,

$$u(t+t_n, x+c_1t_n, y+c_2t_n) \to \hat{V}_{\alpha\theta}(t, x, y)$$
, as $n \to +\infty$ uniformly in \overline{H}_n .

That implies

$$u(t+t_n, x, y) \to \hat{V}_{\alpha\theta}(t, x-c_1t_n, y-c_2t_n)$$
, as $n \to +\infty$ uniformly in \overline{H} . (4.20)

By above arguments applied to $\hat{V}_{\alpha\theta}(t-t_n+t_0,x-c_1t_n,y-c_2t_n)$ for arbitrary $t_0 \in \mathbb{R}$, one can get that

$$\hat{V}_{\alpha\theta}(t-t_n+t_0,x-c_1t_n,y-c_2t_n) \to V_{\alpha\theta}(t+t_0,x,y)$$
, as $n \to +\infty$ locally uniformly for $t \in \mathbb{R}$ and uniformly for $(x,y) \in \mathbb{R}^2$.

Since t_0 is arbitrary, the above convergence is also uniform for $t \in \mathbb{R}$. Thus, by (4.20), one gets that

$$u(t, x, y) \to V_{\alpha\theta}(t, x, y)$$
, as $t \to -\infty$ uniformly in \overline{H} .

Similarly, one can prove that $u(t, x, y) \to V_{\beta\theta}(t, x, y)$ as $t \to -\infty$ uniformly in $\mathbb{R}^2 \setminus H$. On the other hand, for fixed T < 0 such that Lemma 4.2 holds, one can easily check that

$$\lim_{R\to +\infty}\sup_{x^2+y^2>R^2}\left|U^-_{\alpha\theta\beta}(T,x,y)-U^-_{\alpha\beta}(T,x,y)\right|=0,$$

and

$$\lim_{R \to +\infty} \sup_{x^2 + y^2 > R^2} \left| \widetilde{U}^+(T,x,y) - U^-_{\alpha\beta}(T,x,y) \right| \leq 2\varepsilon.$$

Since ε can be arbitrary small and by (4.14), one has that

$$\lim_{R \to +\infty} \sup_{x^2 + v^2 > R^2} \left| u(T, x, y) - U_{\alpha\beta}^{-}(T, x, y) \right| = 0.$$

By stability of the curved front, that is, Theorem 1.9, one has that

$$u(t, x, y) \to V_{\alpha\beta}(t, x, y)$$
, as $t \to +\infty$ uniformly in \mathbb{R}^2 .

This completes the proof of Theorem 1.10.

Finally, we prove Corollary 1.12 which implies that Theorem 1.10 is not empty.

Proof of Corollary 1.12. Assume without loss of generality that $e_* = (0, 1)$. Since $c_{e_*} = \min_{e \in \mathbb{S}} \{c_e\}$ and c'_e is bounded, there exist $\alpha_1 \in (0, \pi/2)$ and $\beta_1 \in (\pi/2, \pi)$ such that

$$\frac{dc_{\theta}}{d\theta}\Big|_{\theta=\alpha} = c_{\alpha}' \cdot (-\sin\alpha, \cos\alpha) \le 0 \text{ for } \alpha \in [\alpha_1, \frac{\pi}{2}],$$

and

$$\left. \frac{dc_{\theta}}{d\theta} \right|_{\theta=\beta} = c_{\beta}' \cdot (-\sin\beta, \cos\beta) \ge 0 \text{ for } \beta \in \left[\frac{\pi}{2}, \beta_1\right].$$

Let $g(\theta) = c_{\theta} / \sin \theta$. Then,

$$g'(\theta) = \frac{c'_{\theta} \cdot (-\sin\theta, \cos\theta)}{\sin\theta} - \frac{c_{\theta}\cos\theta}{\sin^2\theta}.$$

One can make α_1 , β_1 close to $\pi/2$ such that

$$g'(\alpha) < 0$$
 for all $\alpha \in [\alpha_1, \frac{\pi}{2})$ and $g'(\beta) > 0$ for all $\beta \in [\frac{\pi}{2}, \beta_1]$.

Thus, $g(\theta)$ is decreasing from $g(\alpha)$ to $g(\pi/2)$ as θ varying from α_1 to $\pi/2$, and is increasing from $g(\pi/2)$ to $g(\beta)$ as θ varying from $\pi/2$ to β_1 . By continuity, one can pick $\alpha \in [\alpha_1, \pi/2)$ and $\beta \in (\pi/2, \beta_1]$ such that

$$g(\alpha) = g(\beta), g'(\alpha) < 0 \text{ and } g'(\beta) > 0.$$

Let $e_1 = (\cos \alpha, \sin \alpha)$ and $e_2 = (\cos \beta, \sin \beta)$. By Theorem 1.2, there is a curved front $V_{e_1e_2}(t, x, y)$ of (1.1) satisfying (1.10) with $e_0 = e_*$.

Now, by rotating the coordinate, we can assume that e_* is denoted by $(\cos \theta_*, \sin \theta_*)$ where $\theta_* \in (0, \pi/2)$ is small enough. Let e_1 and e_2 be denoted by $(\cos \theta_1, \sin \theta_1)$ and $\cos \theta_2, \sin \theta_2$ respectively, where θ_1 and θ_2 are close to θ_* . By Corollary 1.5 and since θ_* is small enough which means that θ_1 is small enough, there is $\theta_3 \in (\pi/2, \pi)$ such that

$$\frac{c_{\theta_1}}{\sin \theta_1} = \frac{c_{\theta_3}}{\sin \theta_3} := c_{\theta_1 \theta_3},$$

and there is a curved front $V_{\theta_1\theta_3}$ of (1.1) satisfying (1.8) with $\alpha = \theta_1$, $\beta = \theta_3$ and $c_{\alpha\beta} = c_{\theta_1\theta_3}$. On the other hand, since θ_1 is small enough, it implies that θ_3 is close to π enough. Then, since

 θ_2 is also small enough, one has that $\theta_3 - \theta_2$ is close to π enough and hence, $(\cos \theta_2, \sin \theta_2) \cdot (\cos \theta_3, \sin \theta_3) = \cos(\theta_3 - \theta_2)$ is close to -1 enough. By Corollary 1.6, there is e_{**} such that (1.9) holds for $e_1 = (\cos \theta_2, \sin \theta_2)$, $e_2 = (\cos \theta_3, \sin \theta_3)$, $e_0 = e_{**}$ and there is a curved front $V_{\theta_2\theta_3}$ of (1.1) satisfying (1.10).

Then, by Theorem 1.10, there is an entire solution u(t, x, y) of (1.1) satisfying (1.12) and (1.13) with $\alpha = \theta_1$, $\theta = \theta_2$, $\beta = \theta_3$.

References

- [1] S. Allen and J.W. Cahn, A microscopic theory for antiphase boundary motion and its application to antiphase domain coarsening, *Acta. Mettall.* **27** (1979), 1084-1095.
- [2] H. Berestycki, F. Hamel, Front propagation in periodic excitable media, *Comm. Pure Appl. Math.* **55** (2002), 949-1032.
- [3] H. Berestycki, F. Hamel, Generalized transition waves and their properties, *Comm. Pure Appl. Math.* **65** (2012), 592-648.
- [4] Z.-H. Bu, Z.-C. Wang, Curved fronts of monostable reaction-advection-diffusion equations in space-time periodic media, *Comm. Pure Appl. Anal.* **15** (2016), 139-160.
- [5] X. Chen, J.-S. Guo, F. Hamel, H. Ninomiya, J.-M. Roquejoffre, Traveling waves with paraboloid like interfaces for balanced bistable dynamics, *Ann. Inst. H. Poincaré*, *Non Linear Anal.* **24** (2007), 369-393.
- [6] W. Ding, F. Hamel, X. Zhao, Bistable pulsating fronts for reaction-diffusion equations in a periodic habitat, *Indiana Univ. Math. J.*, **66** (2017), 1189-1265.
- [7] A. Ducrot, A multi-dimensional bistable nonlinear diffusion equation in a periodic medium, *Math. Ann.* **366** (2016), 783-818.
- [8] A. Ducrot, T. Giletti, H. Matano, Existence and convergence to a propagating terrace in one-dimensional reaction-diffusion equations, *Trans. Amer. Math. Soc.* **366** (2014), 5541-5566.
- [9] M. EI Smaily, Curved fronts in a shear flow: case of combustion nonlinearities, *Nonlinearity*, **31** (2018), 5643-5663.
- [10] M. El Smaily, F. Hamel, R. Huang, Two-dimensional curved fronts in a periodic shear flow, Nonlinear Analysis 74 (2011), 6469-6486.
- [11] J. Fang, X.-Q. Zhao, Bistable traveling waves for monotone semiflows with applications, *J. Europe. Math. Soc.* **17** (2015), 2243-2288.
- [12] P.C. Fife, J.B. McLeod, The approach of solutions of nonlinear diffusion equations to traveling front solutions, *Arch. Ration. Mech. Anal.* **65** (1977), 335-361.
- [13] R.A. Fisher, The wave of advance of advantageous genes, Ann. of Eugenics 7 (1937), 335-369.
- [14] H. Guo, Propagating speeds of bistable transition fronts in spatially periodic media, *Calc. Var. Part. Diff. Equations* (2018), 57:47.

- [15] H. Guo, F. Hamel, W.-J. Sheng, On the mean speed of bistable transition fronts in unbounded domains, *J. Math. Pures Appl.* **136** (2020), 92-157.
- [16] F. Hamel, Bistable transition fronts in \mathbb{R}^N , Adv. Math. 289 (2016), 279-344.
- [17] F. Hamel, R. Monneau, Solutions of semilinear elliptic equations in \mathbb{R}^N with conical-shaped level sets, *Comm. Part. Diff. Equations* **25** (2000), 769-819.
- [18] F. Hamel, R. Monneau, J.-M. Roquejoffre, Existence and qualitative properties of multidimensional conical bistable fronts, *Disc. Cont. Dyn. Syst. A* **13** (2005), 1069-1096.
- [19] A.N. Kolmogorov, I.G. Petrovskii, S.N. Piskunov, Étude de l'equation de la diffusion avec croissance de la quantité de matière et son application à un problème biologique, *Bull. Univ. État Moscou Ser. Intern. A* **1** (1937), 1-26.
- [20] H. Ninomiya, M. Taniguchi, Existence and global stability of traveling curved fronts in the Allen-Cahn equations, *J. Diff. Equations* **213** (2005), 204-233.
- [21] J. Nolen, L. Ryzhik, Traveling waves in a one-dimensional heterogeneous medium, *Ann. Inst. H. Poincaré*, *Analyse Non Linéaire* **26** (2009), 1021-1047.
- [22] N. Shigesada, K. Kawasaki, E. Teramoto, Traveling periodic waves in heterogeneous environments, Theor. Pop. Bio. 30 (1986), 143-160.
- [23] M. Taniguchi, Traveling fronts of pyramidal shapes in the Allen-Cahn equation, SIAM J. Math. Anal. 39 (2007), 319-344.
- [24] M. Taniguchi, The uniqueness and asymptotic stability of pyramidal traveling fronts in the Allen-Cahn equations, *J. Diff. Equations* **246** (2009), 2103-2130.
- [25] M. Taniguchi, Axially asymmetric traveling fronts in balanced bistable reaction-diffusion equations, *Ann. Inst. H. Poincaré*, *Analyse Non Linéaire*, forthcoming.
- [26] M. Taniguchi, Axisymmetric traveling fronts in balanced bistable reaction-diffusion equations, *Disc. Cont. Dyn. Syst. A* **40** (2020), 3981-3995.
- [27] X. Xin, Existence and uniqueness of travelling waves in a reaction-diffusion equation with combustion nonlinearity, *Indiana Univ. Math. J.* **40** (1991), 985-1008.
- [28] X. Xin, Existence and stability of travelling waves in periodic media governed by a bistable nonlinearity, *J. Dyn. Diff. Eq.* **3** (1991), 541-573.
- [29] J.X. Xin, Existence of planar flame fronts in convective-diffusive periodic media, *Arch. Ration. Mech. Anal.* **121** (1992), 205-233.
- [30] J.X. Xin, Existence and nonexistence of traveling waves and reaction-diffusion front propagation in periodic media, *J. Stat. Phys.* **73** (1993), 893-926.
- [31] J.X. Xin, J. Zhu, Quenching and propagation of bistable reaction-diffusion fronts in multidimensional periodic media, *Physica D* **81** (1995), 94-110.
- [32] A. Zlatoš, Existence and non-existence of transition fronts for bistable and ignition reactions, *Ann. Inst. H. Poincaré, Analyse Non Linéaire*, **34** (2017), 1687-1705.