

The poset of King permutations on a cylinder

Eli Bagno, Estrella Eisenberg, Shulamit Reches and Moriah Sigron

June 9, 2020

Abstract

A permutation $\sigma = [\sigma_1, \dots, \sigma_n] \in S_n$ is called a *cylindrical king permutation* if $|\sigma_{i+1} - \sigma_i| > 1$ for each $1 \leq i \leq n-1$ and $|\sigma_1 - \sigma_n| > 1$. The name comes from the the way one can see these permutations as describing locations of n kings on a chessboard of order $n \times n$ in such a way that (each row and each column contains exactly one king and) no two kings are attacking each other, with the additional condition that a king can move off a certain row and reappear at the beginning of that row.

In a recent paper, we dealt with the more general set of 'king permutations' i.e. the ones which satisfy only the first of the two conditions above. This set constitutes a poest under the well known containment relation on permutations.

In this article we investigate the sub-poset of the cylindrical king permutations and its structure. We examine those cylindrical king permutations whose downset is as large as possible in the upper ranks. We use a modification of Manhattan distance of the plot of a permutation and some of its applications to the cylindrical context to find a criterion for such a permutation to be k -prolific. One of our main results is that the maximal gap between two permutations in the poset of cylindrical permutations is 4.

1 Introduction

A *king permutation* is a permutation $\sigma = [\sigma_1, \dots, \sigma_n] \in S_n$ such that for each $1 \leq i \leq n-1$, $|\sigma_{i+1} - \sigma_i| > 1$. We denote the set of the king permutations of order n by K_n . For example: $K_1 = S_1, K_2 = K_3 = \emptyset$, $K_4 = \{[3142], [2413]\}$. The name "king permutations" refers to the way one can see these permutations as describing locations of n kings on a chessboard of order $n \times n$ in such a way that (each row and each column contains exactly one king and) no two kings are attacking each other. In [2] we investigated the structure of the poset of king permutations with respect to containment.

In our recent paper [1], we considered a celebrated variant of the chess game, namely the cylindrical chess, in which the right and left edges of the board are imagined to be joined in such a way that a piece can move off one edge and reappear at the opposite edge. Let us call this board a *cylindrical board*.

In the language of permutations, non-attacking kings that may move on a $n \times n$ cylindrical board can be described by the following subset of K_n :

Definition 1.1. Let CK_n be defined as

$$CK_n = \{\sigma = [\sigma_1, \dots, \sigma_n] \in S_n : |\sigma_i - \sigma_{i+1}| > 1, |\sigma_1 - \sigma_n| > 1, 1 \leq i \leq n-1\}.$$

An element of CK_n will be called a cylindrical king permutation.

For example: $CK_1 = S_1, CK_2 = CK_3 = CK_4 = \emptyset$, and

$$CK_5 = \{[31425], [14253], [42531], [25314], [53142], [24135], [41352], [13524], [35241], [52413]\}.$$

For $\sigma \in S_n$ and $\pi \in S_k$, ($k < n$), we say that σ *contains* π if there is a subsequence of (the one line notation of) σ that is order-isomorphic to that of π . If this is the case, we write $\pi \prec \sigma$. In this study we are interested in the sub-poset $\mathbb{CK} = \bigcup_{n \geq 1} CK_n$ of the poset $\mathbb{K} = \bigcup_{n \geq 1} K_n$ with respect to the containment relation.

A *bond* in a permutation $\pi \in S_n$ is a length 2 consecutive sub-sequence of adjacent numbers (i.e. interval of length 2 in the terminology of [4]). Note that a king permutation is actually a permutation without bonds. This point of view can be used also to describe the set of cylindrical king permutations. To this aim we extended in [1] the definition of bonds to cyclic bonds.


Definition 1.2. Let $\sigma = [\sigma_1, \dots, \sigma_n] \in S_n$ and let $i \in [n-1]$. We say that the pair (σ_i, σ_{i+1}) is a (regular) bond in σ if $\sigma_i - \sigma_{i+1} = \pm 1$. If $\sigma_n - \sigma_1 = \pm 1$ then we say that the pair (σ_n, σ_1) is an edge bond of σ . In general, adopting the convention that $\sigma_{n+1} = \sigma_1$, we say that (σ_i, σ_{i+1}) is a cyclic bond if it is a regular or an edge bond.

According to this new definition, a permutation is a cylindrical king permutation if and only if it has no cyclic bonds.

1.1 Overview and main results

In this paper we investigate the poset \mathbb{CK} of the cylindrical king permutations under the containment relation and its structure. In addition, we examine those cylindrical king permutations whose downset is as large as possible in the upper ranks.

Here is an example of the downset of cylindrical king permutations of the permutation: [5246173]



Recall that one can view a permutation via its graphical representation, and let us identify a permutation $\sigma = [\sigma_1, \dots, \sigma_n] \in S_n$ with its plot, i.e. the set of all lattice points of the form (i, σ_i) where $1 \leq i \leq n$. From this point of view, we can recognize king permutations, as well as cylindrical king permutations, by considering the distances between the points of their plots. The *Manhattan* or *taxicab* distance is defined as follows:

Definition 1.3. Let $\sigma \in S_n$ and let $i, j \in [n]$. The (Manhattan) distance between the i -th and the j -th entries is defined to be the L_1 distance between the corresponding points in the plot of σ :

$$d_\sigma(i, j) = \|(i, \sigma_i) - (j, \sigma_j)\|_1 = |i - j| + |\sigma_i - \sigma_j|.$$

In [5], the concept of *breadth* of a permutation $\sigma \in S_n$ is defined to be the minimal distance between any two distinct entries of σ . Explicitly:

$$br(\sigma) = \min_{i,j \in [n], i \neq j} d_\sigma(i, j).$$

Example 1.4. For $\sigma = [5371426]$ we have $d_\sigma(2, 5) = 4$, $d_\sigma(1, 4) = 7$, $d_\sigma(1, 2) = 3$ and its breadth is $br(\sigma) = 3$

Using the above definition we can characterize a king permutation as follows:

Observation 1.5. A permutation σ of order $n > 1$ is a king permutation if and only if its breadth is at least 3.

In order to characterize cylindrical king permutations in a similar way, we modify in section 2 the concepts of (Manhattan) distance and breadth to cyclic (Manhattan) distance and cyclic breadth respectively.

For a permutation $\sigma \in S_n$ and $1 \leq k \leq n - 1$, the maximal possible number of distinct permutations $\pi \in S_{n-k}$ which are contained in σ is $\binom{n}{k}$. Indeed, such a permutation π , is obtained from σ by deletion of k entries from the one-line notation of σ . Following [5], our interest is in those permutations $\sigma \in CK_n$ which contain maximally many patterns $\pi \in CK_{n-k}$.

To this aim, we use and modify the following concept of *k-prolific permutation* which was defined in [5]:

Definition 1.6. A permutation $\sigma \in S_n$ is called *k-prolific* if $|\{\pi \in S_{n-k} | \pi \prec \sigma\}| = \binom{n}{k}$.

In other words, a permutation $\sigma = [\sigma_1, \dots, \sigma_n]$ is *k-prolific* if each $(n - k)$ -subset of entries of σ forms a unique pattern in S_{n-k} . It is known (see [2]) that $\sigma \in S_n$ is 1-prolific if and only if σ is a king permutation, and this happens exactly when $br(\sigma) \geq 3$. We are interested in characterizing the cylindrical king permutations which are *k-prolific* in CK_n . I.e the cylindrical king permutations which contain a maximal number of distinct cylindrical king patterns.

In Section 2 we first prove that omitting a single entry from a permutation may decrease the cyclic breadth by at most one, and that for $\sigma \in CK_n$, there are n distinct permutations $\sigma' \in CK_{n-1}$ such that $\sigma' \prec \sigma$ if and only if the cyclic breadth of σ is greater than 3 (See Proposition 2.6 and Proposition 2.8 respectively). Then, using these propositions and our modifications to the concept of *k-prolific*, we get the two main results of this section, (which are generalizations of Theorem 2.25 in [5] to king permutation and cylindrical king permutations respectively):

1. A permutation σ is *k-prolific* in CK_n if and only if its **breadth** is greater or equal to $k + 3$. (See Theorem 2.10)
2. A permutation σ is *k-prolific* in CK_n if and only if its **cyclic breadth** is greater or equal to $k + 3$. (See Theorem 2.11).

In section 3 we investigate the structure of the poset of cylindrical king permutations. This section is a direct continuation of our previous paper [2], which discusses the structure of the poset of king permutations. Since the cylindrical king permutations is a distinguished subset of the king permutations, we examine which features of the poset of the cylindrical king are inherited from the kings poset and which features are different. One of our results is identifying the building blocks of the poset of cylindrical king permutations:

- For every $\sigma \in CK_n$ ($n \geq 5$) there is some $\pi \in CK_5$ such that $\pi \preceq \sigma$. (See Theorem 3.4).

Our main result in this section is that in contrast to the case of king permutations in which the maximal gap between two permutations is 3, in \mathbb{CK} the maximal gap is 4:

- Let $\sigma, \pi \in \mathbb{CK}$ be such that $\pi \prec \sigma$ and $|\sigma| - |\pi| > 5$. Then there is some $\tau \in \mathbb{CK}$ such that $\pi \prec \tau \prec \sigma$ and $|\sigma| - |\tau| \leq 4$. (See Theorem 3.6).

Finally, in section 4 we present directions for further research in this area.

2 Cyclic breadth and k -prolific cylindrical king permutations

We are interested in characterizing the cylindrical king permutations and analyzing some properties of their downsets in \mathbb{CK} . In particular, in this section, we focus on those cylindrical king permutations which contain as many distinct patterns as possible in the poset \mathbb{CK} . There is a close relationship between the breadth of a permutation and the number of distinct patterns it contains. Moreover, as mentioned above, one way to identify a king permutation is by verifying that its breadth is greater than or equal to 3. In order to be able to identify cylindrical king permutations in a similar way, and especially those whose downsets in \mathbb{CK} have maximal size, we modify the definitions of the *(Manhattan) distance* and the *breadth*, introducing the *cyclic (Manhattan) distance* and the *cyclic breadth* of a permutation $\sigma \in S_n$.

Along this section we consider a permutation $\sigma = [\sigma_1, \dots, \sigma_n]$ as if it was located on a circle in such a way that σ_1 occupies the north pole (See figure 1).

Definition 2.1. For $1 \leq i < j \leq n$, we denote by $((i, j))$ the shorter path on the circle leading from i to j which its length is the minimum of $j - i$ and $n - j + i$. We also set $||j - i|| = |((i, j))|$, i.e. $||j - i|| = \min(j - i, n - j + i)$

Definition 2.2. Let $\sigma \in S_n$ and let $i < j \in [n]$. Let the cyclic (Manhattan) distance between the i -th and the j -th entries be defined as:

$$cd_\sigma(i, j) = |\sigma_j - \sigma_i| + ||j - i||.$$

The cyclic breadth of $\sigma \in S_n$ is then defined to be:

$$cbr(\sigma) = \min_{i, j \in [n], i \neq j} cd_\sigma(i, j).$$

Example 2.3. Let $\sigma = [724915836] \in S_9$. Then $cd_\sigma(1, 2) = (7-2) + ||2-1|| = 6$, $cd_\sigma(2, 5) = (2-1) + ||5-2|| = 4$ and $cd_\sigma(1, 9) = (7-6) + ||9-1|| = 2$. The breath of σ is $cbr(\sigma) = 2$

We compare now between the regular (Manhattan) distance and the cyclic (Manhattan) distance as follows:

Observation 2.4. • For each $\sigma \in S_n$ and for each $i, j \in [n]$, $cd_\sigma(i, j) \leq d_\sigma(i, j)$ and so $cbr(\sigma) \leq br(\sigma)$.

- For every $i, j \in [n]$, we have $d_\sigma(i, j) = d_{\sigma^{-1}}(\sigma_i, \sigma_j)$ and thus $br(\sigma) = br(\sigma^{-1})$ which means that the regular (Manhattan) breadth is closed under inverse.
- The cyclic breadth $cbr(\sigma)$ is not always equal to $cbr(\sigma^{-1})$. For example, if $\sigma = [72415836] \in S_8$ then $\sigma^{-1} = [42735816]$, so that $cbr(\sigma) = cd_\sigma(1, 8) = 2$ but $cbr(\sigma^{-1}) = cd_{\sigma^{-1}}(1, 2) = 3$.

Using our new definitions, we can identify the cylindrical king permutation as follows:

Observation 2.5. A permutation σ of order $n > 1$ is a cylindrical king permutation if and only if its cyclic breadth is greater than or equal to 3.

Omitting a single entry from a permutation $\sigma \in S_n$ (and then standardising) produces a permutation $\sigma' \in S_{n-1}$ s.t. $\sigma' \prec \sigma$. In Proposition 2.24 of [5], the authors proved a proposition which partially controls the change in breadth of permutations in S_n : omitting a single entry from a permutation decreases the breadth by at most one. We now present and prove a proposition that extends the above result, considering the cyclic breadth rather than the regular breadth. This enables us to characterize the cyclic breadth of permutations σ' which are contained in a certain permutation σ .

Proposition 2.6. Omitting a single entry from a permutation may decrease the cyclic breadth by at most one.




Figure 1: $\sigma = [26415837]$ located on a circle in such a way that σ_1 occupies the north pole. $cd_\sigma(2, 7) = |6 - 3| + ||7 - 2|| = 6$ where $||7 - 2|| = \min\{7 - 2, 8 + 2 - 7\}$.

Proof. Let $\sigma = [\sigma_1, \sigma_2, \dots, \sigma_n] \in S_n$ and let $1 \leq i < j \leq n$ be such that without loss of generality $\sigma_i < \sigma_j$. Recall that $cd_\sigma(i, j)$ is composed of the *position part* $||j - i||$ and the *value part* $|\sigma_j - \sigma_i|$. Let σ' be the permutation obtained from σ by omitting an element σ_k for some $k \notin \{i, j\}$. We discuss the effect of omitting σ_k on each one of the parts of $cd_\sigma(i, j)$.

Observe that if $\sigma_k > \sigma_j$ or $\sigma_k < \sigma_i$ then this omission does not affect the value part of $cd_\sigma(i, j)$. On the other hand, if $\sigma_i < \sigma_k < \sigma_j$ then, its omission decreases the value part of $cd_\sigma(i, j)$ by 1. Similarly, the omission of σ_k affects the position part if and only if $k \in ((i, j))$. In case there is an effect, the difference is 1.

We conclude, that the omission of σ_k decrease $cd_\sigma(i, j)$ of σ by 2 if and only if $\sigma_i < \sigma_k < \sigma_j$ and $k \in ((i, j))$, in all other cases the difference is 0 or 1.

Now, let $i < j$ such that $cbr(\sigma) = cd_\sigma(i, j)$. Assume to the contrary that there is $k \in [n]$ such that $cbr(\sigma') = cbr(\sigma) - 2$, where σ' is obtained from σ by omitting σ_k . This means that $k \in ((i, j))$ and $\sigma_i < \sigma_k < \sigma_j$. We have now:

$$cbr(\sigma) \leq cd_\sigma(i, k) = \sigma_k - \sigma_i + ||k - i|| < \sigma_j - \sigma_i + ||j - i|| = cd_\sigma(i, j),$$

a contradiction.

We showed that if $cbr(\sigma) = cd_\sigma(i, j)$, then the omission of any element that is not σ_i or σ_j decreases the breadth by at most one. Finally, we show that for $cbr(\sigma) = cd_\sigma(i, j)$, an omission of one of the elements σ_i or σ_j also decreases the cyclic breadth by at most one. Without loss of generality let us consider the omission of σ_j .

Indeed, for every $m, l \neq j$ we have $cbr(\sigma) = cd_\sigma(i, j) \leq cd_\sigma(m, l)$. If $cd_\sigma(i, j) < cd_\sigma(m, l)$ then the omission of σ_j decreases $cd_\sigma(m, l)$ by at most 2 to a value that is greater or equal to $cd_\sigma(i, j) - 1$. If $cd_\sigma(i, j) = cd_\sigma(m, l)$ then $cbr(\sigma) = cd_\sigma(m, l)$, and due to the previous discussion, the omission of σ_j decreases $cd_\sigma(m, l)$ by at most one. Therefore if σ' is obtained from σ by omitting σ_j then also $cbr(\sigma') \geq cbr(\sigma) - 1$. \square

Remark 2.7. Note that omitting an element σ_k might also **increase** the breadth of a permutation σ . For example if $\sigma = [351246]$ then $cbr(\sigma) = cd_\sigma(3, 4) = 2$. If we omit $\sigma_3 = 1$, we get $\sigma' = [24135]$ with $cbr(\sigma') = 3$.

In [6], Coleman demonstrates that maximising the distance between every two elements in a permutation tends to increase the number of distinct sub-permutations (patterns).

In the proof of the next proposition and of several propositions and theorems in the next section, we use the concept of cyclic separators and the notation $\nabla_i^*(\sigma)$.

Let $\nabla_i^*(\sigma)$ be the permutation obtained by omitting the value i from σ and standardize. For example if $\sigma = [31425]$ then $\nabla_2^*(\sigma) = [2134]$ and $\nabla_3^*(\sigma) = [1324]$.

The definition of cyclic separators appears in the Appendix of this paper.

The next proposition characterizes the permutations $\sigma \in CK_n$ which contain n different patterns $\sigma' \prec \sigma$ s.t. $\sigma' \in CK_{n-1}$.

Proposition 2.8. *For $\sigma \in CK_n$, there are n distinct permutations $\sigma' \in CK_{n-1}$ such that $\sigma' \prec \sigma$ if and only if $cbr(\sigma) > 3$.*

Proof. Let $\sigma \in CK_n$ and suppose that there are n distinct permutations $\sigma' \in CK_{n-1}$ such that $\sigma' \prec \sigma$, which means that by omitting any entry of σ , we get a cylindrical king permutation. Since $\sigma \in CK_n$, according to observation 2.5 $cbr(\sigma) \geq 3$. Assume to the contrary that $cbr(\sigma) = 3$, thus there are entries i and j , $i < j$ such that $cbr(\sigma) = cd_\sigma(i, j) = 3$, which means that $||j - i|| = 1$ and $|\sigma_j - \sigma_i| = 2$ or vice versa. Assume without loss of generality that $\sigma_i < \sigma_j$.

- If $||j - i|| = 1$ and $\sigma_j - \sigma_i = 2$ then $\sigma_i + 1$ is a horizontal cyclic separator, so that $\sigma' = \nabla_{\sigma_i+1}^*(\sigma) \notin CK_{n-1}$, a contradiction.
- If $||j - i|| = 2$ and $\sigma_j - \sigma_i = 1$ then σ_{i+1} is a vertical cyclic separator and again $\sigma' = \nabla_{\sigma_{i+1}}^*(\sigma) \notin CK_{n-1}$, a contradiction.

Now, suppose that $cbr(\sigma) > 3$. By corollary 7 of [7], a permutation $\sigma \in S_n$ has maximum number of patterns $\sigma' \prec \sigma$ with $\sigma' \in S_{n-1}$, if and only if it has no bonds. The permutation $\sigma \in CK_n$ has no bonds as a cylindrical king, thus there are n distinct permutations $\sigma' \in S_{n-1}$ such that $\sigma' \prec \sigma$. We show now that each one of them is a cylindrical king permutation. According to Proposition 2.6, omitting a single entry from a permutation decreases the cyclic breadth by at most one. Thus, each $\sigma' \in S_{n-1}$ such that $\sigma' \prec \sigma$ satisfies $cbr(\sigma') \geq 3$ and by observation 2.5, $\sigma' \in CK_{n-1}$. \square

Recall that a permutation $\sigma \in S_n$ is k -prolific if each $(n - k)$ -subset of entries of σ forms a unique pattern. In other words, a permutation $\sigma \in S_n$ is k -prolific if it contains the maximal possible number of distinct permutations $\pi \in S_{n-k}$ such that $\pi \prec \sigma$. The breadth of a permutation affects its being a k -prolific as shown in Theorem 2.25 in [5]: σ is k -prolific if and only if $br(\sigma) \geq k + 2$. This idea can be extended to the poset of king permutations as well as to the poset of cylindrical king permutations. This is the context of the following definition.

Definition 2.9. *We say that a permutation $\sigma \in K_n, (CK_n)$ is k -prolific in $K_n, (CK_n)$ if each $(n - k)$ -subset of the entries of $\sigma = [\sigma_1, \dots, \sigma_n]$ forms a unique pattern in $K_{n-k}, (CK_{n-k})$.*

Using the above definition, we prove the following theorem:

Theorem 2.10. *A permutation σ is k -prolific in K_n if and only if $br(\sigma) \geq k + 3$.*

Proof. Assume to the contrary that σ is k -prolific but $br(\sigma) < k + 3$. This means that there are $1 \leq i, j \leq n$ such that $d_\sigma(i, j) < k + 3$. For the sake of convenience, we assume that $i < j$ and $\sigma_i < \sigma_j$.

Now, let

$$S = \{\sigma_m | i < m < j \text{ or } \sigma_i < \sigma_m < \sigma_j\}.$$

Obviously, $|S| \leq j - i - 1 + \sigma_j - \sigma_i - 1$ so that $|S| + 2 \leq d_\sigma(i, j) < k + 3$ and thus $|S| \leq k$. Omitting the elements of S from σ , we get a permutation $\sigma' \in S_{n-|S|}$ such that $\sigma' \prec \sigma$ and σ' contains a pair $u < v$ such that $d_{\sigma'}(u, v) = 2$, i.e. (σ_u, σ_v) is a bond. For example, let $\sigma = [972415836]$ and let $i = 1$ and $j = 4$. Then we have $S = \{7, 2, 5, 6, 8\}$ and after omitting the elements of S , we get $\sigma' = [4312]$ which indeed has a bond.

Now, if $|S| = k$, this means that $\sigma' \notin K_{n-k}$ so that σ can not be k -prolific in K_n . If $|S| < k$ then we can omit from σ' a set of $k - |S|$ elements which does not contain the bond mentioned above and get an $(n - k)$ -subset of the entries of σ which creates a descendant of σ which is not a member of K_{n-k} , so again σ can not be k -prolific.

On the other hand, assume that $br(\sigma) \geq k+3$ and we prove that σ is k -prolific in K_n . First, by Theorem 2.25 of [5], $br(\sigma) \geq k+2$ if and only if σ is k -prolific in S_n , so that each omission of a k -set from σ yields a different permutation in S_{n-k} . To show that each such $\pi \in S_{n-k}$ is indeed a king permutation, note that by Proposition 2.24 of [5], omitting a single entry from a permutation decreases the breadth by at most one. Thus, omitting each k -set of elements forms a permutation π with $br(\pi) \geq 3$ which implies that $\pi \in K_{n-k}$. \square

The following theorem is the generalization of Theorem 2.10 for the poset of cylindrical king permutations.

Theorem 2.11. *A permutation σ is k -prolific in CK_n if and only if $cbr(\sigma) \geq k+3$.*

Sketch of proof. The proof is almost identical to the proof of Theorem 2.10 except for the following necessary changes:

- Instead of the set S , we define the set:

$$S_{cyclic} = \{\sigma_m \mid m \in ((i, j)) \text{ or } \sigma_i < \sigma_m < \sigma_j\}.$$

- Replace the breadth br by its cyclic parallel breadth cbr .
- At the end of the proof, use Proposition 2.6 of this article instead of Proposition 2.24 of [5].

\square

3 The structure of the poset of cylindrical king permutation

It is easy to observe that for $\sigma \in S_n$, and a regular bond $(a, a+1)$ in σ , omitting a or $a+1$ amounts to the same permutation. In other words

$$\nabla_a^*(\sigma) = \nabla_{a+1}^*(\sigma). \quad (1)$$

Example 3.1. For $\sigma = [523641]$, $\nabla_2^*(\sigma) = \nabla_3^*(\sigma) = [42531]$.

This enables one to count for each permutation $\sigma \in S_n$, the number of permutations $\pi \in S_{n-1}$ such that $\pi \prec \sigma$.

Unfortunately, Equation (1) does not hold anymore when we deal with edge bonds. For example, if $\sigma = [246351]$, then $\nabla_2^*(\sigma) = [35241]$ while $\nabla_1^*(\sigma) = [13524]$. In order to overcome this problem, we define an equivalence relation based on the view of each permutation as if it was written on a circle and not on a line. For example, according to this new view, the permutations $[35241]$ and $[13524]$ are considered identical. Formally, we define:

Definition 3.2. Let C_n be the cyclic sub-group of S_n , generated by the cycle $(12 \cdots n)$. Define an equivalence relation on S_n by $\sigma \sim \tau$ if and only if there is some $\pi \in C_n$ such that $\tau = \sigma \cdot \pi$ and let S_n/C_n be the quotient space. Also, let us denote by $\text{orb}(\sigma)$ the equivalence class of σ .

Example 3.3. Let $\sigma = [23154] \in S_5$. Then

$$\text{orb}(\sigma) = \{[23154], [31542], [15423], [54231], [42315]\}.$$

The following theorem characterizes the permutations that serve as building blocks for the poset of cylindrical king permutations. Note that, if $\sigma \sim \tau$ then σ and τ have the same number of cyclic bonds. Therefore, $\sigma \in CK_n$ iff $\tau \in CK_n$.

Theorem 3.4. For every $\sigma \in CK_n$ ($n \geq 5$) there is some $\pi \in \text{orb}([31425]) \cup \text{orb}([24135])$ such that $\pi \preceq \sigma$.

Proof. We prove by induction on n , the case $n = 5$ being trivial, since $CK_5 = \text{orb}([31425]) \cup \text{orb}([24135])$. We assume that each cylindrical king permutation of order $n - 1$ contains at least one of the permutations of $\text{orb}([31425]) \cup \text{orb}([24135])$ and prove that each cylindrical king permutation of order n contains at least one element of this set.

Let $n > 5$ and assume to the contrary that there is $\sigma = [\sigma_1, \dots, \sigma_n] \in CK_n$ which does not contain any of the permutations in $\text{orb}([31425]) \cup \text{orb}([24135])$. Then for every $1 \leq i \leq n$, the permutation $\nabla_i^*(\sigma)$, contains neither of them as well. By the induction hypothesis, $\nabla_i^*(\sigma) \notin CK_{n-1}$, which implies, that every digit of σ is a cyclic separator. In particular, 1 is a cyclic vertical separator. Hence, we have: $\sigma \in \text{orb}([\dots, a, 1, a+1, \dots])$ or $\sigma \in \text{orb}([\dots, a+1, 1, a, \dots])$. Since $\sigma \in CK_n$, we know that $a > 2$. Without loss of generality, assume that $\sigma \in \text{orb}([\dots, a, 1, a+1, \dots])$. By the above, the element $a+1$ is a cyclic separator of σ . We have now two options as $a+1$ is horizontal or vertical :

- If $a+1$ is horizontal then $\sigma \in \text{orb}([\dots, a+2, a, 1, a+1, \dots])$. For the same reasons as above, a is also a cyclic horizontal separator of σ . Thus $\sigma \in \text{orb}([\dots, a+2, a, 1, a+1, a-1, \dots])$. But the sub sequence $a+2, a, 1, a+1, a-1$ is a pattern which is isomorphic to $[53142]$, so that σ contains an element of $\text{orb}([31425])$.
- If $a+1$ is vertical then $\sigma \in \text{orb}([\dots, a, 1, a+1, 2, \dots])$. As we saw above, a is also a cyclic separator of σ which must be horizontal so that $a = 3$. This implies that $\sigma \in \text{orb}([\dots, 3, 1, 4, 2, \dots])$. Since the element 5 has to be somewhere in σ thus $\sigma \in \text{orb}([\dots, 3, 1, 4, 2, \dots, 5, \dots])$ which contains the pattern $[31425]$.

□

In proposition 3.14 of [2], it is proven that for each two king permutations σ and π such that $\pi \prec \sigma$ and $|\sigma| - |\pi| = 2$ there is a king permutation τ such that $\pi \prec \tau \prec \sigma$. This does not hold for cylindrical king permutations. For example, for $\sigma = [18463527] \in CK_8$ and $\pi = [635241] \in CK_6$, even though $\pi \prec \sigma$, there is no $\tau \in CK_7$ such that $\pi \prec \tau \prec \sigma$.

Moreover, in Theorem 3.15 of [2] it is claimed that if σ, π are king permutations such that $\pi \prec \sigma$ and $|\sigma| - |\pi| > 3$ then there exists a king permutation τ such that $\pi \prec \tau \prec \sigma$ and $|\sigma| - |\tau| \in \{1, 3\}$. I.e. the maximal gap between any two king permutations in the poset \mathbb{K} is 3. This property does not hold true anymore in the poset \mathbb{CK} as the following example verifies:

Example 3.5. Let $\sigma = [579683142]$ and $\pi = [13524]$ which is obtained from σ by removing the elements 3, 1, 4, 2. It is easy to observe that there is no $\tau \in \mathbb{CK}$ such that $\pi \prec \tau \prec \sigma$.

We claim now that the maximal gap between two permutations in \mathbb{CK} is 4.

Theorem 3.6. Let $\sigma, \pi \in \mathbb{CK}$ be such that $\pi \prec \sigma$ and $|\sigma| - |\pi| > 4$. Then there is some $\tau \in \mathbb{CK}$ such that $\pi \prec \tau \prec \sigma$ and $|\sigma| - |\tau| \leq 4$.

The proof of Theorem 3.6 is very subtle and messy. It consists of many similar sub-cases which recur over and over. In order not to be too exhaustive, we present some basic observations and lemmas and then show how to deal with few typical sub-cases and leave the rest of the proof for the reader. For the sake of simplifying the proof, we present a series of claims:

Observation 3.7. Let σ be a permutation, let $\pi \in \mathbb{CK}$, and let i be such that $\pi \prec \nabla_i^*(\sigma)$. If $\nabla_i^*(\sigma)$ contains the cyclic bond consisting of the elements j and k (of σ) then $\pi \prec \nabla_k^*(\sigma)$ or $\pi \prec \nabla_j^*(\sigma)$,

Moreover, if the bond is not an edge bond then both $\pi \prec \nabla_k^*(\sigma)$ and $\pi \prec \nabla_j^*(\sigma)$.

Example 3.8. In the left picture of Figure 2, $\sigma = [264159837] \in S_9$, $\pi = [241635] \in CK_6$, such that $\pi \prec \sigma$, $\pi \prec \nabla_5^*(\sigma)$. In $\nabla_5^*(\sigma) = [25418736]$ there is a regular bond $(5, 4)$ (the digits 4 and 6 in σ), we get that $\pi \prec \nabla_4^*(\sigma)$ and $\pi \prec \nabla_6^*(\sigma)$.

In the right picture $\sigma = [264159837] \in S_9$, $\pi = [253146] \in CK_6$, such that $\pi \prec \sigma$, $\pi \prec \nabla_7^*(\sigma)$. In $\nabla_7^*(\sigma) = [26415873]$ there is an edge bond $(2, 3)$ (the digits 2 and 3 in σ), we get that $\pi \prec \nabla_3^*(\sigma)$ but $\pi \not\prec \nabla_2^*(\sigma)$.




Figure 2:

Definition 3.9. Let us denote by $\nabla_A^*(\sigma)$ the permutation $\nabla_{i_1}^*(\nabla_{i_2}^*(\dots(\nabla_{i_n}^*(\sigma)))$ where $A = \{i_1, \dots, i_n\}$ and $i_1 < \dots < i_n$. For example $\sigma = [264159837] \in S_9$, $\pi = [241635] \in CK_6$, so that $\pi \prec \sigma$. For $A_1 = \{4, 5, 8\}$, $A_2 = \{4, 5, 9\}$, $A_3 = \{5, 6, 8\}$, $A_4 = \{5, 6, 9\}$, we get $\pi = \nabla_{A_i}^*(\sigma)$ for $1 \leq i \leq 4$.

In the following observation, when we write 'adjacent', we take it in the cyclic meaning, i.e. in $[a_1, \dots, a_n]$, the elements a_1 and a_n are considered here as adjacent.

Observation 3.10. Let $\sigma \in \mathbb{CK}$ such that σ contains the (cyclic) block $(a+2, a+4, a+1, a+3)$ or its reverse, i.e. $\sigma \in \text{orb}([\dots, a+2, a+4, a+1, a+3, \dots])$ or $\sigma \in \text{orb}([\dots, a+3, a+1, a+4, a+2, \dots])$.

1. If we remove any single element i from the block $(a+2, a+4, a+1, a+3)$ in σ then $\nabla_i^*(\sigma)$ will contain a cyclic bond. Moreover, we will not obtain a cylindrical king permutation out of σ unless we omit at least 3 elements from the block, in which case, it contracts to the element $a+1$.
2. If the elements a and $a+5$ are not placed adjacent to the block $(a+2, a+4, a+1, a+3)$ or its reverse in σ , then by omitting any 3 elements of the block we get a cylindrical king permutation.

For example, let $\sigma = [39142\mathbf{6857}]$ with the block $(6, 8, 5, 7)$. Note that $\tau = \nabla_6^*(\sigma) = [38142756]$ which contains the bond $(5, 6)$, $\nabla_7^*(\tau) = [3714256]$ with the bond $(5, 6)$, but $\nabla_{\{5,6,7\}}^*(\sigma) = [361425]$ which is a cylindrical king permutation.

3. If a or $a+5$ is placed adjacent to the block $(a+2, a+4, a+1, a+3)$ or its reverse in σ , then removing it produces a cylindrical king permutation. This is due to the fact that in this case a and $a+5$ can not be cyclic separators of any type in σ . For example, let $\sigma = [3192\mathbf{46857}]$ with the block $(6, 8, 5, 7)$. Note that $\tau = \nabla_4^*(\sigma) = [31825746] \in CK_8$.

Lemma 3.11. Let $\sigma, \pi \in \mathbb{CK}$ be such that $\pi \prec \sigma$ and $|\sigma| - |\pi| > 4$. Assume that

- There is some element $a \in \mathbb{N}$ such that $\sigma \in \text{orb}([\dots, a+2, a+4, a+1, a+3, \dots])$ or $\sigma \in \text{orb}([\dots, a+3, a+1, a+4, a+2, \dots])$.
- There is some $1 \leq i \leq 4$ such that $\pi \prec \nabla_{a+i}^*(\sigma)$.

Then there is some $\tau \in \mathbb{CK}$ such that $\pi \prec \tau \prec \sigma$ and $|\sigma| - |\tau| \leq 4$.

Proof. Assume without loss of generality that $\pi \prec \nabla_{a+2}^*(\sigma)$ and that σ contains the block $(a+2, a+4, a+1, a+3)$. The proof depends on the location of this block in σ . The possibilities are: $\sigma = [\dots, a+2, a+4, a+1, a+3]$

or $\sigma = [a+2, a+4, a+1, a+3, \dots]$ or $\sigma = [\dots, a+2, a+4, a+1, a+3, \dots]$ or $\sigma = [a+1, a+3, \dots, a+2, a+4]$ or $\sigma = [a+3, \dots, a+2, a+4, a+1]$ or $\sigma = [a+4, a+1, a+3, \dots, a+2]$.

Here we show only the first case: $\sigma = [\dots, a+2, a+4, a+1, a+3]$ which seems to be the most general case in the sense that it contains all the arguments we need. The other cases can be proved using the same method.

According to Observations 3.7 and 3.10.1, $\pi \prec \nabla_A^*(\sigma)$ for $A = \{a+2, a+3, a+4\}$. If $\nabla_A^*(\sigma) \in \mathbb{CK}$ then we are done (just take $\tau = \nabla_A^*(\sigma) = [\dots, a+1]$). Otherwise, according to 3.10.2, either a or $a+5$ is adjacent to the block $(a+2, a+4, a+1, a+3)$. Assume w.l.o.g. that a is adjacent to the above block. We have 2 possibilities.

- If $\sigma = [\dots, a, a+2, a+4, a+1, a+3]$ then $\nabla_A^*(\sigma) = [\dots, a, a+1]$ and according to Observation 3.7, $\pi \prec \nabla_a^*(\nabla_A^*(\sigma)) = \nabla_{A'}^*(\nabla_A^*(\sigma)) \prec \nabla_a^*(\sigma)$ for $A' = \{a+1, a+2, a+3\}$ and so by 3.10.3 $\tau = \nabla_a^*(\sigma)$ and we are done.
- If $\sigma = [a, \dots, b, a+2, a+4, a+1, a+3]$ (where $b \neq a+5$) then $\nabla_A^*(\sigma) = [a, \dots, a+1]$. By Observation 3.7, $\pi \prec \nabla_a^*(\nabla_A^*(\sigma)) \prec \nabla_a^*(\sigma)$ or $\pi \prec \nabla_B^*(\sigma)$ for $B = A \cup \{a+1\}$.
 1. If $\pi \prec \nabla_a^*(\sigma)$ then we are done since according to 3.10.3., $\tau = \nabla_a^*(\sigma)$ is a cylindrical king permutation.
 2. If $\pi \not\prec \nabla_a^*(\sigma)$ and $\nabla_B^*(\sigma)$ is a cylindrical king permutation then we are done by taking $\tau = \nabla_B^*(\sigma)$. (Note that this is the single case in which $|\sigma| - |\tau| = 4$). Otherwise, the permutation $\nabla_B^*(\sigma)$ contains a cyclic bond, leading to one of the following two possibilities:
 - (a) $\sigma = [a, a+5, \dots, a+2, a+4, a+1, a+3]$ which is impossible since then $\nabla_B^*(\sigma)$ is of the form $[a, a+1, \dots]$ and by 3.7 we get $\pi \prec \nabla_a^*(\sigma)$ which contradicts the assumption.
 - (b) $b = a-1$, i.e. $\sigma = [a, \dots, a-1, a+3, a+1, a+4, a+2]$ and according to Observation 3.7, $\pi \prec \nabla_{a-1}^*(\nabla_A^*(\sigma)) \prec \nabla_{a-1}^*(\sigma)$. We claim that $\nabla_{a-1}^*(\sigma)$ is a cylindrical king permutation. Indeed, otherwise $\sigma = [a, a-2, \dots, a-1, a+2, a+4, a+1, a+3]$ and $\nabla_{a-1}^*(\sigma)$ contains a regular bond consisting of the elements a and $a-2$ of σ . Hence, due to 3.7 $\pi \prec \nabla_a^*(\sigma)$ which contradicts the assumption. Now, we can take $\tau = \nabla_{a-1}^*(\sigma)$ and we are done.

□

Now we return to the proof of Theorem 3.6. Let σ, π be cylindrical king permutations such that $\pi \prec \sigma$, and $|\sigma| - |\pi| > 4$ and let us prove that there is $\tau \in \mathbb{CK}$ such that $\pi \prec \tau \prec \sigma$ and $|\sigma| - |\tau| \leq 4$.

Proof. of Theorem 3.6

Since $\mathbb{CK} \subset \mathbb{K}$, we have that π and σ are king permutations. Thus, according to Theorem 3.15 in [2], there is $\tau \in \mathbb{K}$ such that $\pi \prec \tau \prec \sigma$ and $|\sigma| - |\tau| \in \{1, 3\}$. If $|\sigma| - |\tau| = 3$ then according to the proof of Theorem 3.15 in [2], σ contains the block $(a+2, a+4, a+1, a+3)$ or its reverse and τ is obtained from σ by omitting 3 elements from this block. Then we can proceed according to Lemma 3.11. Assume now that $|\sigma| - |\tau| = 1$. If $\tau \in \mathbb{CK}$ then we are done. Otherwise, τ is of the form: $\tau = [a, \dots, a+1]$ and there exists some cyclic separator b such that $\tau = \nabla_b^*(\sigma)$. If b is vertical then we have $\sigma = [a, \dots, a+1, b]$, otherwise b is horizontal i.e. $b = a+1$ and so $\sigma = [a, \dots, a+1, \dots, a+2]$ (or their reverses). Here we consider only the vertical case: $\sigma = [a, \dots, a+1, b]$. Since $\pi \prec \nabla_b^*(\sigma) = [a, \dots, a+1]$, by Observation 3.7, $\pi \prec \nabla_a^*(\sigma)$ or $\pi \prec \nabla_{a+1}^*(\sigma)$.

We consider here only the case $\pi \prec \nabla_{a+1}^*(\sigma)$ but $\pi \not\prec \nabla_a^*(\sigma)$. If $\nabla_{a+1}^*(\sigma) \in \mathbb{CK}$ we are done. Otherwise $\nabla_{a+1}^*(\sigma)$ contains a cyclic bond. Thus we have for σ one of the following options: either $\sigma = [a, \dots, b \pm 1, a+1, b]$ or $\sigma = [a, a+2, \dots, a+1, b]$. Let us consider again the first case, $\sigma = [a, \dots, b+1, a+1, b]$. In this case, $\nabla_{a+1}^*(\sigma)$ has a regular bond consisting of the elements b and $b+1$ of σ and so by Observation 3.7, $\pi \prec \nabla_b^*(\sigma)$ and $\pi \prec \nabla_{b+1}^*(\sigma)$. If $\nabla_{b+1}^*(\sigma)$ is a cylindrical king permutation, we are done. Otherwise, the omission of the element $b+1$ produces a cyclic bond. Again, there are two options for the structure of σ . The first option is that $a = b+2$ and thus $\sigma = [a, \dots, a-1, a+1, a-2]$ and thus according to Lemma 3.11 we are done. The second option (if $a \neq b+2$) is that $\sigma = [a, \dots, a+2, b+1, a+1, b]$, in this case $\nabla_A^*(\sigma) = [a, \dots, a+1, b]$ for

$A = \{a+2, b+1\}$ is a cylindrical king permutation. (Indeed, the pair $(a+1, b)$ is not a bond since $a+1$ and b are adjacent in σ . The elements $a+1$ or $a-1$ can not appear beside the element a since σ is a cylindrical king permutation). This completes the proof. \square

4 Directions for further research

- In [5], Bevan, Homberger and Tenner proved that a k -prolific permutation in S_n must be at least of size $\frac{k^2}{2} + 2k + 1$. A challenge may be to find a lower bound for the size of a k -prolific permutation in K_n and in CK_n .
- In the chess game, one can further enable the pieces to go off a column and reappear at the beginning of that columns, thus obtaining a 'chess on the torus'. The permutations describing non-attacking kings on this board are called torical king permutations and constitute a sub-poset of their own. The work on these permutations is in progress.

5 Appendix

In a recent paper [3], the authors defined a new concept, called *separator*. A separator in a permutation $\sigma = [\sigma_1, \dots, \sigma_n] \in S_n$ is an element σ_i that its omission forms a new bond. In particular, if $\sigma \in K_n$ is a king permutation, then the permutation obtained by the omission of σ_i is a king permutation if and only if σ_i is not a separator in σ . In order to characterize the structure of the poset of cylindrical kings, we need a cyclic version of that concept. We start by recalling the definition of the (regular) separator (see Definition 2.1 in [3]).

Definition 5.1. For $\sigma = [\sigma_1, \dots, \sigma_n] \in S_n$ we say that $\sigma_i = \mathbf{a}$ separates σ_{j_1} from σ_{j_2} in σ if by omitting the element a from σ we get a **new** bond. This happens if and only if one of the following cases holds:

1. j_1, i, j_2 are subsequent numbers and $|\sigma_{j_1} - \sigma_{j_2}| = 1$, i.e

$$\sigma = [\dots, b, \mathbf{a} = \sigma_i, b \pm 1, \dots]$$

In this case we say that a is a **vertical separator**.

2. $\sigma_{j_1}, \sigma_i = a, \sigma_{j_2}$ is an increasing or decreasing sub sequence of subsequent numbers, and $|j_1 - j_2| = 1$, i.e,

$$\sigma = [\dots, \mathbf{a}, \dots, a \pm 1, a \mp 1, \dots]$$

or

$$\sigma = [\dots, a \pm 1, a \mp 1, \dots, \mathbf{a}, \dots].$$

In this case we say that a is a **horizontal separator**.

Cylindrical kings are actually permutations having no **cyclic** bonds. In order to be able to deal with edge bonds, we have to modify the definition 5.1 of the separator.

Definition 5.2. Let $\sigma = [\sigma_1, \dots, \sigma_n] \in S_n$. An element σ_i will be called an **edge separator** if omitting it produces a new edge bond. This happens when one of the following holds:

1. If $|\sigma_1 - \sigma_{n-1}| = 1$ then σ_n is a (vertical) edge separator.
2. If $|\sigma_n - \sigma_2| = 1$ then σ_1 is a (vertical) edge separator.
3. If for some $1 < i < n$ we have that $\sigma_1, \sigma_i, \sigma_n$ is an increasing or decreasing sub sequence of subsequent numbers, then we say that σ_i is a (horizontal) edge separator.

We say that an element σ_i is a **cyclic separator** if it is a regular or an edge separator. In this case we say that σ_i cyclically separates some element from another.

Example 5.3. Let $\sigma = [52341]$, then $\sigma_5 = 1$ is a (vertical) edge separator in σ . Let $\sigma = [6257134]$, then $\sigma_3 = 5$ is a (horizontal) edge separator in σ . If $\sigma = [246135]$, then $\sigma_6 = 5$ is a vertical edge separator and a horizontal (regular) separator in σ .

References

- [1] E. Bagno E. Eisenberg, S. Reches, and M. Sigron. *Counting King Permutations on the Cylinder*. Jan. 2020. arXiv: 2001.02948v1.
- [2] E. Bagno E. Eisenberg, S. Reches, and M. Sigron. “On the poset of King-Non-Attacking permutations”. In: *European J. Combin* 87.103119 (2020). DOI: <https://doi.org/10.1016/j.ejc.2020.103119>.
- [3] E. Bagno E. Eisenberg, S. Reches, and M. Sigron. *Separators - a new statistic for permutations*. May 2019. arXiv: 1905.12364.
- [4] D. Bevan. *Permutation patterns: basic definitions and notation*. arXiv: 1506.06673.
- [5] D. Bevan, C. Homberger, and B. Tenner. “Prolific permutations and permuted packings: Downsets containing many large patterns”. In: *Journal of Combinatorial Theory, Series A* 153 (Jan. 2018), pp. 98–121. DOI: <https://10.1016/j.jcta.2017.08.006>.
- [6] M. Coleman. “An Answer to a Question by Wilf on Packing Distinct Patterns in a Permutation”. In: *Electr. J. Comb.* 11 (May 2004). DOI: <https://10.37236/1849>.
- [7] C. Homberger. “Counting Fixed-Length Permutation Patterns”. In: *Online Journal of Analytic Combinatorics* 7 (2012).