Microscopic derivation of the extended Gross-Pitaevskii equation for quantum droplets in binary Bose mixtures

Hui Hu and Xia-Ji Liu

Centre for Quantum Technology Theory, Swinburne University of Technology, Melbourne, Victoria 3122, Australia

(Dated: September 18, 2020)

An ultradilute quantum droplet is a self-bound liquid-like state recently observed in ultracold Bose-Einstein condensates. In most previous theoretical studies, it is described by a phenomenological low-energy effective theory, termed as the extended Gross–Pitaevskii equation. Here, we microscopically derive the Gross–Pitaevskii equation for the condensate and also for a pairing field in an inhomogeneous quantum droplet realized by Bose-Bose mixtures with attractive inter-species interaction. We show that the inclusion of the pairing field is essential, in order to have a consistent description of the droplet state. We clarify that, the extended Gross–Pitaevskii equation used earlier should be understood as the equation of motion for the pairing field, rather than the condensate. The fluctuations of the pairing field give rise to low-energy collective excitations of the droplet. We also present the Bogoliubov equations for gapless phonon modes and gapped modes due to pairing in real space, which characterizes single-particle-like excitations of the droplet. The equations of motion derived in this work for the condensate and the pairing field serve an ideal starting point to understand the structure and collective excitations of non-uniform ultradilute quantum droplets in on-going cold-atom experiments.

I. INTRODUCTION

One of the recent breakthroughs in ultracold atomic physics is the realization of a self-bound liquid-like droplet state [\[1](#page-5-0)[–9\]](#page-5-1). In three spatial dimensions, this new quantum state of matter arises from a delicate balance [\[10](#page-5-2)] between the attractive mean-field potential $\partial E_{\rm MF}/\partial n$ and the repulsive potential $\partial E_{\rm LHY}/\partial n$ provided by the celebrated Lee-Huang-Yang (LHY) term E_{LHY} from quantum fluctuations [\[11](#page-5-3)], which scale like n and $n^{3/2}$, respectively, as a function of the total density n of the system. In almost all the theoretical studies of quantum droplets [\[10,](#page-5-2) [12](#page-5-4)[–26\]](#page-5-5), the extended Gross–Pitaevskii (GP) equation $(E = E_{\text{MF}} + E_{\text{LHY}})$ [\[10\]](#page-5-2),

$$
i\hbar \frac{\partial}{\partial t} \psi = \left[-\frac{\hbar^2}{2m} \nabla^2 + \frac{\partial E}{\partial n} \left(n = |\psi|^2 \right) \right] \psi, \qquad (1)
$$

has been extensively used to describe the structure and dynamics of the condensate wave ψ of the droplet. This phenomenological low-energy effective theory is often thought to be unavoidable, since the LHY term derived from the standard Bogoliubov theory [\[27](#page-5-6)] becomes complex due to an unstable, softening phonon mode and therefore should be amended in an empirical way [\[10\]](#page-5-2). As a priori assumption, the extended GP equation gives a useful description of the low-lying softening phonon mode, in view of the density functional theory.

It turns out that an amended LHY term is not the only choice. As demonstrated in our recent work [\[28,](#page-5-7) [29\]](#page-5-8), the inconsistency of a complex LHY term in the Bogoliubov theory of quantum droplets could be alternatively removed by the inclusion of pairing between two bosons in different species. The bosonic pairing changes the unstable softening mode into stable gapped mode and leads to much more accurate ground-state energy for the droplet state, as benchmarked by the state-of-the-art diffusion

Monte Carlo simulations [\[20](#page-5-9), [21,](#page-5-10) [24](#page-5-11)]. Roughly speaking, however, the gapped mode should be considered as single-particle excitations with relatively high energy, as we learn from the conventional fermionic pairing theories [\[30](#page-5-12)[–33\]](#page-6-0). Therefore, we lose the track of the low-lying collective excitations. Immediate questions then are, how can we find those collective excitations and how to physically interpret the phenomenological extended GP equation, Eq. [\(1\)](#page-0-0), within the pairing theory?

In this work, we derive and discuss the equations of motion for the condensate, the pairing field and their fluctuations in a non-uniform quantum droplet realized with attractive Bose-Bose mixtures, following the framework of the previously developed microscopic pairing theory [\[28,](#page-5-7) [29](#page-5-8)]. We show that the existence of the pairing field implies additional $U(1)$ symmetry breaking for the manybody paired bosons, i.e., the bosonic Cooper pairs. Analogous to a Bardeen–Cooper-Schrieffer (BCS) fermionic superfluid [\[31](#page-6-1), [32](#page-6-2)], the low-lying collective excitations of the droplet state then should be characterized by the pair fluctuations around the mean-field saddle-point for the pairing field. We confirm this picture, by microscopically derive the extended GP equation for a large droplet, where the local density approximation could be applied. We also present the Bogoliubov equations for the gapless phonon mode associated with the $U(1)$ symmetry breaking of the condensate and for the gapped mode due to the pairing. We interpret these two relatively high-lying modes as single-particle excitations of the droplet state.

In brief, we have extended our previous work [\[28,](#page-5-7) [29](#page-5-8)] to consider a finite-size quantum droplet with inhomogeneous density distribution. Non-uniform Bogoliubov equations and extended GP equation have been derived to describe single-particle excitations and collective excitations of the droplet state, respectively. A selfconsistent numerical solution of these equations could provide us a more in-depth understanding of the structure and collective oscillations of a self-bound quantum droplet in free space.

II. MODEL HAMILTONIAN

We start by considering a three-dimensional homonuclear Bose-Bose mixture such as a cloud of ³⁹K atoms in two hyperfine states (i.e., a 39 K- 39 K mixture), as in recent experiments [\[4](#page-5-13), [6\]](#page-5-14). For simplicity, we assume equal repulsive intra-species interactions with strengths $g_{11} = g_{22} = g$ and attractive inter-species interactions $g_{12} = g_{21}$, and also equal population in each species. The grand canonical Hamiltonian of the system can then be written as,

$$
\hat{K} = \int d\mathbf{x} \left[\mathcal{H}_0 + \mathcal{H}_{\text{intra}} + \mathcal{H}_{\text{inter}} \right],\tag{2}
$$

$$
\mathscr{H}_0 = \sum_{i=1,2} \hat{\phi}_i^{\dagger}(\mathbf{x}) \left[-\frac{\hbar^2 \nabla^2}{2m} + V_T(\mathbf{x}) - \mu \right] \hat{\phi}_i(\mathbf{x}),
$$
(3)

$$
\mathscr{H}_{\text{intra}} = \frac{g}{2} \sum_{i=1,2} \hat{\phi}_i^{\dagger}(\mathbf{x}) \hat{\phi}_i^{\dagger}(\mathbf{x}) \hat{\phi}_i(\mathbf{x}) \hat{\phi}_i(\mathbf{x}), \qquad (4)
$$

$$
\mathcal{H}_{\text{inter}} = -\frac{\hat{\Delta}^{\dagger}\hat{\Delta}}{g_{12}} - \left[\hat{\Delta}\hat{\phi}_{1}^{\dagger}\left(\mathbf{x}\right)\hat{\phi}_{2}^{\dagger}\left(\mathbf{x}\right) + \text{H.c.}\right] \tag{5}
$$

where $\hat{\phi}_i^{\dagger}(\mathbf{x})$ and $\hat{\phi}_i(\mathbf{x})$ ($i = 1, 2$) are creation and annihilation field operators for the i-species bosons with mass $m_1 = m_2 = m$ and with chemical potential $\mu_1 = \mu_2 = \mu$. We have explicitly included an external harmonic trap $V_T(\mathbf{x}) = m\omega^2 \mathbf{x}^2/2$, in order to account for a possible residual weak potential in experiments. In three dimensions, the bare interaction strengths g_{ij} are to be regularized, due to the well-known ultraviolet divergence of the contact inter-particle interaction. We shall rewrite them in terms of the three-dimensional s-wave scattering lengths $a_{11} = a_{22} = a$ and a_{12} ,

$$
\frac{1}{g_{ij}} = \frac{m}{4\pi\hbar^2 a_{ij}} - \frac{1}{\mathcal{V}} \sum_{\mathbf{k}} \frac{m}{\hbar^2 \mathbf{k}^2},\tag{6}
$$

where V is the volume of the system. We note that in $\mathcal{H}_{\text{inter}}$ we have taken the Hubbard-Stratonovich transformation and have introduced a pairing field $\Delta(\mathbf{x})$ to decouple the inter-species interaction Hamiltonian [\[29\]](#page-5-8). In the weakly interacting regime, it suffices to take a static saddle-point solution for the pairing field. Hence, we treat $\Delta(\mathbf{x}) = \Delta(\mathbf{x})$ as a variational c-number function. The dynamics of the pairing field can be added back later, when we consider the fluctuations around the saddle point.

III. BOGOLIUBOV THEORY WITH PAIRING

We use the standard Bogoliubov theory to solve the model Hamiltonian, in the presence of a static pairing field $\Delta(\mathbf{x})$. Following Refs. [\[34,](#page-6-3) [35\]](#page-6-4), in the Bogoliubov approximation we rewrite the bosonic field operators,

$$
\hat{\phi}_i(\mathbf{x}) = \phi_c(\mathbf{x}) + \delta \hat{\phi}_i(\mathbf{x}), \qquad (7)
$$

where $\delta \hat{\phi}_i$ is considered as a small correction to the condensate wave-function $\phi_c(\mathbf{x}) = e^{i\theta(\mathbf{x})} |\phi_c(\mathbf{x})|$ and $\theta(\mathbf{x})$ is the phase of the condensate. For the ground state, we take $\theta(\mathbf{x}) = 0$; while for a vortex state, we set $\theta(\mathbf{x}) = \varphi$ with the polar coordinate φ . Our model Hamiltonian may then be expanded through second order in $\delta \hat{\phi}_i^{\dagger}$ and $\delta \hat{\phi}_i$, and the linear term vanish identically if $\phi_c(\mathbf{x})$ satisfies the GP equation,

$$
\left[-\frac{\hbar^2 \nabla^2}{2m} + V_T(\mathbf{x}) - \mu + g |\phi_c|^2\right] \phi_c - \tilde{\Delta}(\mathbf{x}) \phi_c^* = 0. \tag{8}
$$

By introducing the notations $C(\mathbf{x}) = g |\phi_c|^2$ and

$$
\hat{T}(\mathbf{x}) \equiv -\frac{\hbar^2 \left[\nabla + i\nabla\theta(\mathbf{x})\right]^2}{2m} + V_T(\mathbf{x}) - \mu,
$$
 (9)

we may rewrite the GP equation into the relation for $\Delta(\mathbf{x}) \equiv e^{-2i\theta} \tilde{\Delta}(\mathbf{x}),$

$$
\Delta\left(\mathbf{x}\right) = C\left(\mathbf{x}\right) + \left[C\left(\mathbf{x}\right)\right]^{-1/2} \hat{T}\left(\mathbf{x}\right) \left[C\left(\mathbf{x}\right)\right]^{1/2}.\tag{10}
$$

We then obtain the truncated Bogoliubov Hamiltonian,

$$
\hat{K}_B = \sum_{i=1,2} \int d\mathbf{x} \left[\delta \hat{\phi}_i^\dagger \mathcal{L}_0 \delta \hat{\phi}_i + \left(\frac{C}{2} e^{2i\theta} \delta \hat{\phi}_i^\dagger \delta \hat{\phi}_i^\dagger + \text{H.c.} \right) \right] \n- \int d\mathbf{x} \left[\left(\tilde{\Delta} \delta \hat{\phi}_1^\dagger \delta \hat{\phi}_2^\dagger + \text{H.c.} \right) + \frac{C^2}{g} + \frac{|\Delta|^2}{g_{12}} \right], \tag{11}
$$

where $\mathscr{L}_0 \equiv -\hbar^2 \nabla^2/(2m) + V_T(\mathbf{x}) - \mu + 2C(\mathbf{x})$. The Bogoliubov Hamiltonian consists of a c-number condensate part and a quadratic form in in $\delta \hat{\phi}_i$ and $\delta \hat{\phi}_i^{\dagger}$. This quadratic form could be diagonalized with the linear Bogoliubov transformation,

$$
\delta\hat{\phi}_i(\mathbf{x}) = e^{+i\theta(\mathbf{x})} \sum_n \left[u_{ni}(\mathbf{x}) \hat{\alpha}_n + v_{ni}^*(\mathbf{x}) \hat{\alpha}_n^{\dagger} \right], \tag{12}
$$

$$
\delta \hat{\phi}_{i}^{\dagger}(\mathbf{x}) = e^{-i\theta(\mathbf{x})} \sum_{n} \left[u_{ni}^{*}(\mathbf{x}) \hat{\alpha}_{n}^{\dagger} + v_{ni}(\mathbf{x}) \hat{\alpha}_{n} \right], (13)
$$

where $\hat{\alpha}_n^{\dagger}$ and $\hat{\alpha}_n$ are creation and annihilation field operators of Bogoliubov quasiparticles satisfying the usual Bose commutation relations. We can show that the truncated Bogoliubov Hamiltonian reduces to

$$
\hat{K}_B = -\int d\mathbf{x} \left[\frac{C^2 (\mathbf{x})}{g} + \frac{|\Delta (\mathbf{x})|^2}{g_{12}} \right] - \int d\mathbf{x} \sum_{ni} E_n |v_{ni} (\mathbf{x})|^2 + \sum_n E_n \hat{\alpha}_n^{\dagger} \hat{\alpha}_n, (14)
$$

provided that the quasiparticle wave-functions $u_{ni}(\mathbf{x})$ and $v_{ni}(\mathbf{x})$ obey the coupled Bogoliubov eigenvalue equations $(i = 1, 2)$,

$$
\mathcal{L}u_{ni} + C(\mathbf{x})v_{ni} - \Delta(\mathbf{x})v_{n,3-i} = +E_n u_{ni}, \quad (15)
$$

$$
\mathcal{L}^*v_{ni} + C(\mathbf{x})u_{ni} - \Delta^*(\mathbf{x})u_{n,3-i} = -E_n v_{ni}, \quad (16)
$$

where $\mathscr{L} \equiv \hat{T}(\mathbf{x}) + 2C(\mathbf{x})$ is a Hermitian operator satisfying $\int d\mathbf{x} u^* \mathscr{L} v = \int d\mathbf{x} v \mathscr{L}^* u^*$ [\[34\]](#page-6-3). It is easy to check that the wave-functions satisfy the normalization and orthogonality conditions,

$$
\sum_{i=1,2} \int d\mathbf{x} \left[u_{ni}^* u_{mi} - v_{ni}^* v_{mi} \right] = \delta_{nm},\tag{17}
$$

$$
\sum_{i=1,2} \int d\mathbf{x} \left[u_{ni} v_{mi} - v_{ni} u_{mi} \right] = 0.
$$
 (18)

The Bogoliubov equations [\(15\)](#page-1-0) and [\(16\)](#page-1-0) have a wellknown particle-hole symmetry: if u_{ni} and v_{ni} are a solution with energy E_n , then there is always another solution v_{ni}^* and u_{ni}^* with energy $-E_n$. In the diagonalized Bogoliubov Hamiltonian Eq. [\(14\)](#page-1-1), therefore, we choose the positive eigenvalues $E_n \geq 0$. As a result, the thermodynamic potential at zero temperature takes the form,

$$
\Omega = -\int d\mathbf{x} \left[\frac{C^2}{g} + \frac{|\Delta|^2}{g_{12}} + \sum_{ni} E_n |v_{ni}(\mathbf{x})|^2 \right], \quad (19)
$$

from which, we may determine the variational pairing field $\Delta(\mathbf{x})$ through the *functional* minimization, i.e.,

$$
\frac{\delta\Omega\left[\mu,\Delta\left(\mathbf{x}\right)\right]}{\delta\Delta\left(\mathbf{x}\right)} = 0.\tag{20}
$$

It is worth noting that the LHY contribution $\Omega_{\rm LHY}$ = $-\int d\mathbf{x} \sum_{ni} E_n |v_{ni}(\mathbf{x})|^2$ is formally divergent. This divergence can be removed by regularizing the bare interaction strengths g and g_{12} . In practice, we introduce a high-energy cut-off energy E_c , above which the discreteness of the energy spectrum E_n is no longer important and we semi-classically solve the Bogoliubov equations under the local density approximation to obtain $u_{ki}(\mathbf{x})$, $v_{\mathbf{k}i}(\mathbf{x})$ and $E_{\mathbf{k}}$ [\[36](#page-6-5)]. We then rewrite the thermodynamic potential into two parts, $\Omega = \int d\mathbf{x} [\Omega_d(\mathbf{x}) + \Omega_c(\mathbf{x})]$, where

$$
\Omega_d = -\frac{m}{4\pi\hbar^2} \left(\frac{C^2}{a} + \frac{|\Delta|^2}{a_{12}} \right) - \sum_{i, E_n < E_c} E_n \left| v_{ni} \right|^2, \text{(21)}
$$

$$
\Omega_c = \sum_{\mathbf{k}} \frac{m\left(C^2 + |\Delta|^2\right)}{\hbar^2 \mathbf{k}^2} - \sum_{i, E_{\mathbf{k}} \ge E_c} E_{\mathbf{k}} \left|v_{\mathbf{k}i}\right|^2. \tag{22}
$$

For a given chemical potential μ , the GP equation [\(8\)](#page-1-2), the Bogoliubov equations [\(15\)](#page-1-0) and [\(16\)](#page-1-0), and the thermodynamic potential Eqs. [\(21\)](#page-2-0) and [\(22\)](#page-2-0) form a closed set of equations to determine $C(\mathbf{x})$ and $\Delta(\mathbf{x})$, and consequently the condensate wave-function $|\phi_c(\mathbf{x})| \propto \sqrt{C(\mathbf{x})}$ and the Bogoliubov spectrum E_n . This is the first key result of our work. The total number of atoms can be calculated from the thermodynamic potential relation $N = -\partial\Omega/\partial\mu$, which provides the normalization to $\phi_c(\mathbf{x})$. We note that, if we neglect the quantum depletion, which is small in the weakly interacting regime, we may write down directly $|\phi_c(\mathbf{x})| = \sqrt{mC(\mathbf{x})/(4\pi\hbar^2 a)}$.

For simplicity, from now on we focus on the ground state with a phase $\theta(\mathbf{x}) = 0$. Our discussion given below can be easily extended to the general case with a nonzero phase factor $\theta(\mathbf{x}) \neq 0$ and we will put $\theta(\mathbf{x})$ back when it is needed.

IV. BULK PROPERTIES OF QUANTUM DROPLETS

For a large droplet in the absence of any external potential and in the ground state, the function $C(\mathbf{x})$ and the pairing field $\Delta(\mathbf{x})$ are essentially real constant, except at the edge of the droplet. Thus, if we neglect the edge effect, the GP equation [\(8\)](#page-1-2) gives the relation $C = \mu + \Delta$. The Bogoliubov equations [\(15\)](#page-1-0) and [\(16\)](#page-1-0) in momentum space take the form,

$$
\begin{bmatrix}\nB_{\mathbf{k}} & 0 & C & -\Delta \\
0 & B_{\mathbf{k}} & -\Delta & C \\
C & -\Delta & B_{\mathbf{k}} & 0 \\
-\Delta & C & 0 & B_{\mathbf{k}}\n\end{bmatrix}\n\begin{bmatrix}\nu_{\mathbf{k}1} \\
u_{\mathbf{k}2} \\
v_{\mathbf{k}1} \\
v_{\mathbf{k}2}\n\end{bmatrix} = E_{\mathbf{k}}\n\begin{bmatrix}\n+u_{\mathbf{k}1} \\
+u_{\mathbf{k}2} \\
-v_{\mathbf{k}1} \\
-v_{\mathbf{k}2}\n\end{bmatrix}, (23)
$$

where $B_{\mathbf{k}} \equiv \varepsilon_{\mathbf{k}} + C + \Delta$ with $\varepsilon_{\mathbf{k}} = \hbar^2 \mathbf{k}^2/(2m)$. We then obtain two Bogoliubov spectra,

$$
E_{-}(\mathbf{k}) = \sqrt{\varepsilon_{\mathbf{k}} \left(\varepsilon_{\mathbf{k}} + 2C + 2\Delta\right)},\tag{24}
$$

$$
E_{+}(\mathbf{k}) = \sqrt{(\varepsilon_{\mathbf{k}} + 2C)(\varepsilon_{\mathbf{k}} + 2\Delta)}.
$$
 (25)

For both dispersions, the wave-functions $u_{\mathbf{k}i}(\mathbf{x})$ = $u_{ki}e^{i\mathbf{k}\cdot\mathbf{x}}/\sqrt{\mathcal{V}}$ and $v_{ki}(\mathbf{x}) = v_{ki}e^{i\mathbf{k}\cdot\mathbf{x}}/\sqrt{\mathcal{V}}$ are given by,

$$
u_{\mathbf{k}1}^2 = u_{\mathbf{k}2}^2 = \frac{1}{4} \left(\frac{B_{\mathbf{k}}}{E_{\mathbf{k}}} + 1 \right),\tag{26}
$$

$$
v_{\mathbf{k}1}^2 = v_{\mathbf{k}2}^2 = \frac{1}{4} \left(\frac{B_{\mathbf{k}}}{E_{\mathbf{k}}} - 1 \right),\tag{27}
$$

from which, we determine the thermodynamic potential,

$$
\frac{\Omega}{\mathcal{V}} = -\frac{m}{4\pi\hbar^2} \left[\frac{C^2}{a} + \frac{\Delta^2}{a_{12}} \right] + \frac{1}{2} \sum_{\mathbf{k}} \left[E_+ \left(\mathbf{k} \right) + E_- \left(\mathbf{k} \right) \right]
$$

$$
-2 \left(\varepsilon_{\mathbf{k}} + C + \Delta \right) + \frac{C^2 + \Delta^2}{\varepsilon_{\mathbf{k}}} \right]. \tag{28}
$$

This result was obtained in our previous works using a path-integral functional approach [\[28,](#page-5-7) [29\]](#page-5-8). By integrating over the momentum k , we arrive at the expression,

$$
\frac{\Omega}{\mathcal{V}} = -\frac{m}{4\pi\hbar^2} \left[\frac{C^2}{a} + \frac{\Delta^2}{a_{12}} \right] + \frac{8m^{3/2}C^{5/2}}{15\pi^2\hbar^3} \mathcal{G}_3\left(\frac{\Delta}{C}\right), \tag{29}
$$

where $\mathcal{G}_3(\alpha) \equiv (1 + \alpha)^{5/2} + h_3(\alpha)$ with $h_3(\alpha) \equiv$ $(15/4)\int_0^\infty dt\sqrt{t}[\sqrt{(t+1)(t+\alpha)}-(t+1/2+\alpha/2)+(1-\alpha)^2]$ α ²/(8t)]. Near the equilibrium density of quantum droplets, the chemical potential $|\mu|$ is typically much smaller than C and Δ [\[28](#page-5-7), [29\]](#page-5-8). We may then expand $\Omega(\mu)$ in powers of μ , $\Omega(\mu) = \Omega^{(0)} + \mu \Omega^{(1)} + \cdots$, where

$$
\frac{\Omega^{(0)}}{\mathcal{V}} = -\frac{m}{4\pi\hbar^2} \left(\frac{1}{a} + \frac{1}{a_{12}} \right) \Delta^2 + \frac{32\sqrt{2}m^{3/2}}{15\pi^2\hbar^3} \Delta^{5/2} (30)
$$

$$
\frac{\Omega^{(1)}}{\mathcal{V}} = -\frac{m}{2\pi\hbar^2 a} \Delta + \frac{8\sqrt{2}m^{3/2}}{3\pi^2\hbar^3} \Delta^{3/2}.
$$
 (31)

By taking the derivative $-\partial\Omega/\partial\mu$ [\[37\]](#page-6-6), we obtain the pairing gap

$$
\Delta \simeq \frac{2\pi\hbar^2 a}{m} n \left[1 + \eta\right],\tag{32}
$$

and the total energy per unit volume $E/\mathcal{V} = \mu n + \Omega/\mathcal{V}$.

$$
\frac{E}{\mathcal{V}} \simeq -\frac{\pi \hbar^2}{m} \left(a + \frac{a^2}{a_{12}} \right) n^2 (1 + 2\eta) \n+ \frac{256\sqrt{\pi}}{15} \frac{\hbar^2 a^{5/2}}{m} n^{5/2} \left(1 + \frac{5}{2} \eta \right),
$$
\n(33)

where the correction factor $\eta = 32\sqrt{na^3}/(3\sqrt{\pi}) \ll 1$ comes from the second term in $\Omega^{(1)}/\mathcal{V}$. This small correction is absent if we approximate $C \simeq \Delta$ in the LHY thermodynamic potential (i.e., the second term on the right-hand-side of Eq. [\(29\)](#page-2-1)).

V. LARGE DROPLETS WITHIN THE LOCAL DENSITY APPROXIMATION

To take into account the edge effect for a large quantum droplet, we may take the local density approximation, by assuming very slowly varying $C(\mathbf{x})$ and $\Delta(\mathbf{x})$ in real space. That is, $C(\mathbf{x})$ and $\Delta(\mathbf{x})$ change very slowly at the scale of the healing length $\xi_c = (8\pi n a)^{-1/2}$, which is set by the condition $\hbar^2/(2m\xi_c^2) \sim 4\pi\hbar^2 a n/m$ [\[38\]](#page-6-7). This amounts to neglecting all the discrete energy levels in the summation of Eq. [\(21\)](#page-2-0). In other words, we treat all the wave-functions classically as plane-waves in a locally uniform cell (located at the position x) with a well-defined wave-vector **k**. The quasi-particle amplitudes u_{ki} and $v_{\mathbf{k}i}$ and the corresponding energy level $E_{\mathbf{k}}$ (including the two branches $E_-(\mathbf{k})$ and $E_+(\mathbf{k})$ can then be obtained from Eqs. (26) , (27) , (24) and (25) , respectively, with the position dependence is explicitly kept by using the slowly spatially varying function $C(\mathbf{x})$ and the pairing field $\Delta(\mathbf{x})$. The neglect of the discrete energy levels is equivalent to setting the cut-off energy $E_c = 0$ in Eqs. (21) and (22) , and consequently at each position **x** we have,

$$
\Omega_d = -\frac{m}{4\pi\hbar^2} \left[\frac{C^2 \left(\mathbf{x} \right)}{a} + \frac{\Delta^2 \left(\mathbf{x} \right)}{a_{12}} \right],\tag{34}
$$

and

$$
\Omega_c = \sum_{\mathbf{k}} \frac{C^2(\mathbf{x}) + \Delta^2(\mathbf{x})}{2\varepsilon_{\mathbf{k}}} - \sum_{E_{\mathbf{k}}} \frac{B_{\mathbf{k}}(\mathbf{x}) - E_{\mathbf{k}}(\mathbf{x})}{2}, \tag{35}
$$

where we have treated $\Delta(\mathbf{x})$ as a *real* and non-negative function, since it only differs slightly from $C(\mathbf{x})$. We note that, $B_{\mathbf{k}}(\mathbf{x})$ and $E_{\mathbf{k}}(\mathbf{x})$ in the expression of Ω_c should be understood as the local dispersion relations at the position **x** with $C(\mathbf{x})$ and $\Delta(\mathbf{x})$, and we also need to consider both dispersions $E_-(\mathbf{k}; \mathbf{x})$ and $E_+(\mathbf{k}; \mathbf{x})$. The summation in Ω_c over the momentum **k** at the position x is easy to carry out. Following the derivation in Eq. [\(29\)](#page-2-1), we obtain

$$
\Omega_c = \frac{8m^{3/2}}{15\pi^2\hbar^3} \left[C\left(\mathbf{x} \right) + \Delta \left(\mathbf{x} \right) \right]^{5/2},\tag{36}
$$

where we have used the fact that $C(\mathbf{x}) \simeq \Delta(\mathbf{x})$, so the function $h_3(C/\Delta) \propto [C(\mathbf{x}) - \Delta(\mathbf{x})]^2 / \Delta^2(\mathbf{x}) \ll 1$ can be safely neglected [\[28](#page-5-7)]. By adding the contribution Ω_d in Eq. [\(34\)](#page-3-0), we find that

$$
\Omega = -\int d\mathbf{x} \frac{m}{4\pi\hbar^2} \left[\frac{C^2(\mathbf{x})}{a} + \frac{\Delta^2(\mathbf{x})}{a_{12}} \right] + \int d\mathbf{x} \frac{8m^{3/2}}{15\pi^2\hbar^3} \left[C(\mathbf{x}) + \Delta(\mathbf{x}) \right]^{5/2}.
$$
 (37)

By taking the functional derivative $\delta\Omega[\Delta(\mathbf{x})]/\delta\Delta = 0$, we find that,

$$
\left[\frac{C}{a}\frac{\delta C}{\delta \Delta} + \frac{\Delta}{a_{12}}\right] - \frac{16\sqrt{m}}{3\pi\hbar} \left[C + \Delta\right]^{3/2} \left(\frac{\delta C}{\delta \Delta} + 1\right) = 0.
$$
\n(38)

From the GP equation [\(10\)](#page-1-3), the function $C(\mathbf{x})$ is related to the pairing field $\Delta(\mathbf{x})$,

$$
C(\mathbf{x}) \simeq \Delta(\mathbf{x}) - [\Delta(\mathbf{x})]^{-1/2} \hat{T}(\mathbf{x}) [\Delta(\mathbf{x})]^{1/2}. \qquad (39)
$$

As $\Delta(x)$ is a very slowly varying function for a large droplet, to a good approximation we may take $\delta C/\delta \Delta =$ 1 and also set $C(\mathbf{x}) \simeq \Delta(\mathbf{x})$ in the second term of Eq. [\(38\)](#page-3-1). The latter is equivalent to neglecting the small correction factor η in Eq. [\(32\)](#page-3-2) and Eq. [\(33\)](#page-3-3), as we discussed earlier. By further inspired by the relation [\(32\)](#page-3-2) to introduce $\Phi^2(\mathbf{x}) \equiv [m/(2\pi\hbar^2 a)]e^{2i\theta(\mathbf{x})}\Delta(\mathbf{x})$ = $[m/(2\pi\hbar^2a)]\tilde{\Delta}(\mathbf{x})$, we rewrite Eq. [\(38\)](#page-3-1) into the form,

$$
\left[\hat{T} - \frac{2\pi\hbar^2}{m}\left(a + \frac{a^2}{a_{12}}\right)|\Phi|^2 + \frac{128\sqrt{\pi}}{3}\frac{\hbar^2 a^{5/2}}{m}|\Phi|^3\right]\Phi = 0.
$$
\n(40)

By recalling $\hat{T} = -\hbar^2 \nabla^2/(2m) - \mu = -\hbar^2 \nabla^2/(2m) - i\hbar \partial_t$ in the absence of external potential, the above equation is precisely the extended GP equation [\(1\)](#page-0-0), once we take Eq. [\(33\)](#page-3-3) with $\eta = 0$ as the density functional $E(n)$. Hence, we have microscopically derived the extended GP equation, under the condition (i.e., within the local density approximation) that it is applicable.

To check the self-consistency of the local density approximation, it is useful to note that, from Eq. [\(40\)](#page-3-4) the droplet typically changes at the length scale [\[10](#page-5-2)]

$$
\xi = \frac{64\sqrt{6}}{5\pi} \left(1 + \frac{a}{a_{12}} \right)^{-3/2} a \tag{41}
$$

with an equilibrium density

$$
n = \frac{25\pi}{16384} \left(1 + \frac{a}{a_{12}} \right)^2 a^{-3}.
$$
 (42)

Therefore, we find the ratio,

$$
\frac{\xi}{\xi_c} = 2\sqrt{3} \left(1 + \frac{a}{a_{12}} \right)^{-1/2},\tag{43}
$$

which is about 11.5 under the typical experimental condition $a_{12} \simeq -1.1a$ [\[4,](#page-5-13) [6\]](#page-5-14). As a result of $\xi \gg \xi_c$, the local density approximation is well satisfied for the experimentally realized Bose droplets [\[4](#page-5-13), [6](#page-5-14)].

Our derivation clearly shows that the wave-function in the extended GP equation [\(1\)](#page-0-0) represents the pairing field $\tilde{\Delta}(\mathbf{x})$, rather than the condensate wave-function $\phi_c(\mathbf{x})$, as one may na¨ıvely anticipate. The latter satisfies instead the ordinary GP equation, as given in Eq. [\(8\)](#page-1-2). This clarification is another key result of our work.

In our microscopic pairing theory, the quantum droplet could be viewed as a mixture of bosonic atoms and loosely-bound many-body bosonic Cooper pairs, analogous to a two-component Fermi superfluid where fermions and loosely-bound fermionic Cooper pairs coexist [\[30](#page-5-12)[–32\]](#page-6-2). Eq. [\(40\)](#page-3-4) derived here can therefore be regarded as the bosonic counterpart of the well-known Ginzburg–Landau equation for the BCS pairing order pa-rameter [\[39\]](#page-6-8). Physically, there are two $U(1)$ symmetry breakings, one is associated with the condensate wavefunction $\phi_c(\mathbf{x})$ and another with the pairing field $\Delta(\mathbf{x})$. As in a fermionic superfluid, low-energy collective excitations correspond to the fluctuations around the static saddle-point solution $\Delta(\mathbf{x})$ and can be studied by linearizing the extended GP equations [\(1\)](#page-0-0) or [\(40\)](#page-3-4) with

$$
\Phi(t) = e^{-i\mu t/\hbar} \left[\Phi_0 + \sum_n \left(U_n e^{-i\omega_n t} + V_n^* e^{+i\omega_n t} \right) \right],\tag{44}
$$

to the first order in fluctuations $U_n(\mathbf{x})$ and $V_n(\mathbf{x})$. This leads to the Bogoliubov equations for pairing fluctuations. In contrast, the quasi-particles described by the Bogoliubov equations in [\(15\)](#page-1-0) and [\(16\)](#page-1-0) should be understood as single-particle excitations of bosonic atoms.

It is worth noting that, in a scalar weakly-interacting Bose gas collective excitations and single-particle excitations are strongly correlated, due to the existence of the condensate, so the Bogoliubov quasi-particles are generally treated as collective excitations [\[40](#page-6-9), [41\]](#page-6-10). The situation in quantum droplets is less clear, as a result of the bosonic pairing. A careful identification of collective excitations and single-particle excitations is therefore needed, by calculating of density-density correlation functions, presumably within the random-phaseapproximation [\[41](#page-6-10)].

VI. CONCLUSIONS

In summary, we have constructed a microscopic pairing theory for describing the non-uniform states of quantum droplets realized with an attractive Bose-Bose mixture in three dimensions. We have pointed out that the droplet can be viewed as a coherent mixture of bosonic atoms and loosely-bound bosonic Cooper pairs, both of which are Bose-condensed. We have presented the equations of motion for the atomic condensate and the pairing field of Cooper pairs, as well as the fluctuations around them. The resulting closed set of equations (8) , (15) , (16) , (21) and [\(22\)](#page-2-0) forms the basis to investigate the structure and dynamics of quantum droplets in future studies.

By using these equations for a large quantum droplet, where the spatial variation in the condensate wavefunction and the pairing field is small, we have microscopically derived the extended Gross-Pitaevskii equation, which has been previously used a phenomenological low-energy effective theory. We have clarified that the extended Gross-Pitaevskii equation describes the pairing field, instead of the condensate wave-function as one may naïvely expect. This clarification would be important for properly distinguishing the collective excitations and single-particle behaviors in the intriguing new quantum state of self-bound liquid-like droplets.

We note finally that, in practice the functional minimization of the thermodynamic potential with respect to the variational pairing field is difficult to carry out numerically. For the most relevant case of a self-bound spherical quantum droplet in free space, therefore it is useful to parameterize the pairing field $\Delta(r)$ by using a few variational parameters. For example, for a large droplet we may consider a flat-top distributed droplet with the form

$$
\Delta(r) = \Delta_0 \left[1 + \frac{\cosh(\kappa r)}{\cosh(\kappa R)} \right]^{-1},\tag{45}
$$

where the parameter κ controls the shape of the droplet at the edge $r \sim R$. The initial input for the variational parameters Δ_0 , κ and R can be obtained by solving the extended Gross-Pitaevskii Eq. [\(40\)](#page-3-4). For the given pairing field $\Delta(r)$ in Eq. [\(45\)](#page-4-0) and chemical potential $\mu < 0$ (for the droplet state), we first solve the nonlinear Gross-Pitaevskii Eq. (8) for $C(r)$ by using the standard iterative algorithm [\[42\]](#page-6-11). As the effective potential in the Gross-Pitaevskii equation, i.e., $V_{\text{eff}} = -\mu - \Delta(r)$ always curves upwards, we could numerically find a stable solution for the function $C(r)$, which acquires a flat-top distribution with $C(r = 0) = \mu + \Delta_0$ and vanishes at $r \sim R$. In the second step, we solve the coupled Bogoliubov equations, Eqs. [\(15\)](#page-1-0) and [\(16\)](#page-1-0), for the quasi-particle energies and wave-functions, following the work by Hutchinson, Zaremba and Griffin [\[43\]](#page-6-12). We then calculate the total thermodynamic potential Ω by using Eqs. [\(21\)](#page-2-0) and [\(22\)](#page-2-0). In this way, we obtain Ω as functions of Δ_0 , κ and R . The minimization then leads to an optimum set of the

variational parameters, improving the initial guess from the solution of Eq. [\(40\)](#page-3-4).

ACKNOWLEDGMENTS

This research was supported by the Australian Research Council's (ARC) Discovery Program, Grant No. DP170104008 (H.H.) and Grant No. DP180102018 (X.- J.L).

- [1] I. Ferrier-Barbut, H. Kadau, M. Schmitt, M. Wenzel, and T. Pfau, Observation of Quantum Droplets in a Strongly Dipolar Bose Gas, Phys. Rev. Lett. 116, 215301 (2016).
- [2] M. Schmitt, M. Wenzel, F. Böttcher, I. Ferrier-Barbut, and T. Pfau, Self-bound droplets of a dilute magnetic quantum liquid, Nature (London) 539, 259 (2016).
- [3] L. Chomaz, S. Baier, D. Petter, M. J. Mark, F. Wächtler, L. Santos, and F. Ferlaino, Quantum-Fluctuation-Driven Crossover from a Dilute Bose-Einstein Condensate to a Macrodroplet in a Dipolar Quantum Fluid, Phys. Rev. X 6, 041039 (2016).
- [4] C. Cabrera, L. Tanzi, J. Sanz, B. Naylor, P. Thomas, P. Cheiney, and L. Tarruell, Quantum liquid droplets in a mixture of Bose-Einstein condensates, Science 359, 301 (2018).
- [5] P. Cheiney, C. R. Cabrera, J. Sanz, B. Naylor, L. Tanzi, and L. Tarruell, Bright Soliton to Quantum Droplet Transition in a Mixture of Bose-Einstein Condensates, Phys. Rev. Lett. 120, 135301 (2018).
- [6] G. Semeghini, G. Ferioli, L. Masi, C. Mazzinghi, L. Wolswijk, F. Minardi, M. Modugno, G. Modugno, M. Inguscio, and M. Fattori, Self-Bound Quantum Droplets of Atomic Mixtures in Free Space, Phys. Rev. Lett. 120, 235301 (2018).
- [7] G. Ferioli, G. Semeghini, L. Masi, G. Giusti, G. Modugno, M. Inguscio, A. Gallemi, A. Recati, and M. Fattori, Collisions of Self-Bound Quantum Droplets, Phys. Rev. Lett. 122, 090401 (2019).
- [8] F. Böttcher, M. Wenzel, J.-N. Schmidt, M. Guo, T. Langen, I. Ferrier-Barbut, T. Pfau, R. Bombín, J. Sánchez-Baena, J. Boronat, and F. Mazzanti, Dilute dipolar quantum droplets beyond the extended Gross-Pitaevskii equation, Phys. Rev. Research 1, 033088 (2019).
- [9] C. D'Errico, A. Burchianti, M. Prevedelli, L. Salasnich, F. Ancilotto, M. Modugno, F. Minardi, and C. Fort, Observation of quantum droplets in a heteronuclear bosonic mixture, Phys. Rev. Research 1, 033155 (2019).
- [10] D. S. Petrov, Quantum Mechanical Stabilization of a Collapsing Bose-Bose Mixture, Phys. Rev. Lett. 115, 155302 (2015) .
- [11] T. D. Lee, K. Huang, and C. N. Yang, Eigenvalues and Eigenfunctions of a Bose System of Hard Spheres and Its Low-Temperature Properties, Phys. Rev. 106, 1135 (1957).
- [12] D. S. Petrov and G. E. Astrakharchik, Ultradilute Low-Dimensional Liquids, Phys. Rev. Lett. 117, 100401 (2016)
- [13] D. Baillie, R. M. Wilson, R. N. Bisset, and P. B. Blakie, Self-bound dipolar droplet: A localized matter wave in free space, Phys. Rev. A 94 , $021602(R)$ (2016).
- [14] F. Wächtler and L. Santos, Ground-state properties and elementary excitations of quantum droplets in dipolar Bose-Einstein condensates, Phys. Rev. A 94, 043618

(2016).

- [15] D. Baillie, R.M. Wilson, and P. B. Blakie, Collective Excitations of Self-Bound Droplets of a Dipolar Quantum Fluid, Phys. Rev. Lett. 119, 255302 (2017).
- [16] Y. Li, Z. Luo, Y. Liu, Z. Chen, C. Huang, S. Fu, H. Tan, and B. A. Malomed, Two-dimensional solitons and quantum droplets supported by competing self- and crossinteractions in spin-orbit-coupled condensates, New J. Phys. 19, 113043 (2017).
- [17] A. Cappellaro, T. Macrì, G. F. Bertacco, and L. Salasnich, Equation of state and self-bound droplet in Rabicoupled Bose mixtures, Sci. Rep. 7, 13358 (2017).
- [18] X. Cui, Spin-orbit-coupling-induced quantum droplet in ultracold Bose-Fermi mixtures, Phys. Rev. A 98, 023630 (2018).
- [19] C. Staudinger, F. Mazzanti and R. E. Zillich, Self-bound Bose mixtures, Phys. Rev. A 98, 023633 (2018).
- [20] L. Parisi, G. E. Astrakharchik, and S. Giorgini, Liquid State of One-Dimensional Bose Mixtures: A Quantum Monte Carlo Study, Phys. Rev. Lett. 122, 105302 (2019).
- [21] V. Cikojević, L. Vranješ Markic, G. E. Astrakharchik, and J. Boronat, Universality in ultradilute liquid Bose-Bose mixtures, Phys. Rev. A 99, 023618 (2019).
- [22] Y. Kartashov, G. Astrakharchik, B. Malomed, and L. Torner, Frontiers in multidimensional self-trapping of nonlinear fields and matter, Nat. Rev. Phys. 1, 185 (2019).
- [23] M. Tylutki, G. E. Astrakharchik, B. A. Malomed, and D. S. Petrov, Collective excitations of a one-dimensional quantum droplet, Phys. Rev. A 101, 051601(R) (2020).
- [24] V. Cikojević, L. Vranješ Markić, and J. Boronat, Finiterange effects in ultradilute quantum drops, New J. Phys. 22, 053045 (2020).
- [25] Y. Wang, L. Guo, S. Yi, and T. Shi, Theory for Self-Bound States of Dipolar Bose-Einstein Condensates, [arXiv:2002.11298](http://arxiv.org/abs/2002.11298) (2020).
- [26] M. Ota and G. E. Astrakharchik, Beyond Lee-Huang-Yang description of self-bound Bose mixtures, SciPost Phys. 9, 020 (2020).
- [27] D. M. Larsen, Binary mixtures of dilute Bose gases with repulsive interactions at low temperature, Ann. Phys. (N.Y.) 24, 89 (1963).
- [28] H. Hu and X.-J. Liu, Consistent theory of self-bound quantum droplets with bosonic pairing, [arXiv:2005.08581](http://arxiv.org/abs/2005.08581) (2020).
- [29] H. Hu, Jia Wang, and X.-J. Liu, Microscopic pairing theory of a binary Bose mixture with interspecies attractions: bosonic BEC-BCS crossover and ultradilute low-dimensional quantum droplets, [arXiv:2006.00434v](http://arxiv.org/abs/2006.00434)2 (2020); to appear in Physical Review A.
- [30] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, Microscopic Theory of Superconductivity, Phys. Rev. 106, 162 (1957).
- [31] H. Hu, X.-J. Liu, and P. D. Drummond, Equation of state of a superfluid Fermi gas in the BCS-BEC crossover, Europhys. Lett. 74, 574 (2006).
- [32] R. B. Diener, R. Sensarma, and M. Randeria, Quantum fluctuations in the superfluid state of the BCS-BEC crossover, Phys. Rev. A 77, 023626 (2008).
- [33] H. Hu, X.-J. Liu, and P. D. Drummond, Comparative study of strong-coupling theories of a trapped Fermi gas at unitarity, Phys. Rev. A 77, 061605(R) (2008).
- [34] A. L. Fetter, Nonuniform States of an Imperfect Bose Gas, Ann. Phys. (N.Y.) 70, 67 (1972).
- [35] A. Griffin, Conserving and gapless approximations for an inhomogeneous Bose gas at finite temperatures, Phys. Rev. B 53, 9341 (1996).
- [36] X.-J. Liu, H. Hu, and P. D. Drummond, Mean-field thermodynamics of a spin-polarized spherically trapped Fermi gas at unitarity, Phys. Rev. A 75, 023614 (2007).
- [37] We note that, if we minimize $\Omega(\mu, \Delta)$ with respect to Δ , we obtain the condition, $\mu \simeq (1 + a/a_{12})\Delta +$ $[32\sqrt{2}m^{1/2}a/(3\pi\hbar)]\Delta^{3/2}$, which is bound from below. The low bound $\mu_c = -(\pi^2/768)(1 + a/a_{12})^3 [\hbar^2/(2ma^2)]$ is the critical chemical potential, below which one can

no longer find the local minimum in $\Omega(\mu, \Delta)$. Therefore, μ_c is the critical chemical potential for the droplet-to-gas transition, which happens at sufficiently low density.

- [38] F. Dalfovo, S. Giorgini, L. P. Pitaevskii, and S. Stringari, Theory of Bose-Einstein condensation in trapped gases, Rev. Mod. Phys. 71, 463 (1999).
- [39] C. A. R. Sá de Melo, M. Randeria, and J. R. Engelbrecht, Crossover from BCS to Bose superconductivity: Transition temperature and time-dependent Ginzburg-Landau theory, Phys. Rev. Lett. 71, 3202 (1993).
- [40] A. Griffin, Excitations in a Bose-Condensed Liquid (Cambridge University Press, New York, 1993).
- [41] X.-J. Liu, H. Hu, A. Minguzzi, and M. P. Tosi, Collective oscillations of a confined Bose gas at finite temperature in the random-phase approximation, Phys. Rev. A 69, 043605 (2004).
- [42] H. Pu and N. P. Bigelow, Properties of Two-Species Bose Condensates, Phys. Rev. Lett. 80, 1130 (1998).
- [43] D. A. W. Hutchinson, E. Zaremba, and A. Griffin, Finite Temperature Excitations of a Trapped Bose Gas, Phys. Rev. Lett. 78, 1842 (1997).