

Lattice walk area combinatorics, some remarkable trigonometric sums and Apéry-like numbers

Stéphane Ouvry* and Alexios P. Polychronakos†

June 12, 2020

Abstract

Explicit algebraic area enumeration formulae are derived for various lattice walks generalizing the canonical square lattice walk, and in particular for the triangular lattice chiral walk recently introduced by the authors. A key element in the enumeration is the derivation of some remarkable identities involving trigonometric sums –which are also important building blocks of non trivial quantum models such as the Hofstadter model– and their explicit rewriting in terms of multiple binomial sums. An intriguing connection is also made with number theory and some classes of Apéry-like numbers, the cousins of the Apéry numbers which play a central role in irrationality considerations for $\zeta(2)$ and $\zeta(3)$.

* LPTMS, CNRS, Université Paris-Sud, Université Paris-Saclay, 91405 Orsay Cedex, France; *stephane.ouvry@u-psud.fr*

† Department of Physics, City College of New York, NY10031 and the Graduate Center of CUNY, New York, NY 10016, USA; *apolychronakos@ccny.cuny.edu*

1 Introduction

The enumeration of random walks of given algebraic area on a two-dimensional lattice is a hard and challenging problem. The algebraic area is defined as the oriented area spanned by the walk as it traces the lattice. A unit lattice cell enclosed in the counterclockwise (positive) way has an area $+1$, whereas when enclosed in the clockwise (negative) way it has an area -1 . The total algebraic area is the area enclosed by the walk weighted by the winding number: if the walk winds around more than once, the area is counted with multiplicity. The combinatorics of such walks depend on the exact rule generating them and on the lattice geometry. The canonical example is closed random walks on a square lattice. This problem can be mapped to the famous Hofstadter model [2] of a particle hopping on a square lattice pierced by a constant magnetic field, with the value of the magnetic field playing a role analogous to the chemical potential for the area of the walk. Indeed, algebraic area enumerations are mapped on quantum mechanical models since in quantum mechanics a magnetic field couples to the area spanned by the particle.

An exact formula for the number of square lattice walks of given length and algebraic area was only recently obtained in the form of nested binomial sums [1]. The analysis revealed some remarkable trigonometric sums to be key ingredients for the algebraic area enumeration. They are defined for p and q coprime positive integers as

$$\frac{1}{q} \sum_{k=1}^q b_{p/q}^{l_1}(k) b_{p/q}^{l_2}(k+1) \dots b_{p/q}^{l_j}(k+j-1) \quad (1)$$

where $b_{p/q}(k)$ is a trigonometric function called spectral function which depends on the rational number p/q , and l_1, l_2, \dots, l_j is a set of positive or null integers. In the algebraic area enumeration for square lattice walks these integers are the parts in the compositions of the integer n , i.e., $n = l_1 + \dots + l_j$ and all l_i positive, with n fixing the length of the walk. But we will consider more general lattice walks where some of the l_i 's can be null, in a way to be specified below.

In [1] the focus was on the spectral function

$$b_{p/q}(k) = (2 \sin(\pi k p/q))^2 \quad (2)$$

which encodes the Hofstadter dynamics. The algebraic area enumeration was obtained in part thanks to an explicit rewriting of the trigonometric sum (1), when evaluated for the Hofstadter spectral function (2), in terms of the binomial multiple sums

$$\frac{1}{q} \sum_{k=1}^q b_{p/q}^{l_1}(k) b_{p/q}^{l_2}(k+1) \dots b_{p/q}^{l_j}(k+j-1) = \sum_{\substack{A=-\infty \\ A \text{ even}}}^{\infty} e^{i\pi A p/q} \quad (3)$$

$$\sum_{k_3=-l_3}^{l_3} \dots \sum_{k_j=-l_j}^{l_j} \binom{2l_1}{l_1 + A/2 + \sum_{i=3}^j (i-2)k_i} \binom{2l_2}{l_2 - A/2 - \sum_{i=3}^j (i-1)k_i} \prod_{i=3}^j \binom{2l_i}{l_i + k_i}$$

Eq. (3) is valid for any set of positive or null integers l_i with an A -summation range finite due to the first two binomials, where A appears. In the specific case where the l_i 's are all positive –as is the case for the square lattice walks algebraic area enumeration– A is restricted in the interval $[-2\lfloor(l_1 + \dots + l_j)^2/4\rfloor, 2\lfloor(l_1 + \dots + l_j)^2/4\rfloor]$. When some of the l_i 's are null these bounds can be generalized (see, e.g., the bounds in eq. (11)).

We note that when we replace $e^{i\pi Ap/q}$ by 1 in (3) we get the binomial identity

$$\binom{2(l_1 + \dots + l_j)}{l_1 + \dots + l_j} = \sum_{\substack{A=-\infty \\ A \text{ even}}}^{\infty} \quad (4)$$

$$\sum_{k_3=-l_3}^{l_3} \cdots \sum_{k_j=-l_j}^{l_j} \binom{2l_1}{l_1 + A/2 + \sum_{i=3}^j (i-2)k_i} \binom{2l_2}{l_2 - A/2 - \sum_{i=3}^j (i-1)k_i} \prod_{i=3}^j \binom{2l_i}{l_i + k_i}$$

where the resulting binomial in the LHS¹ will be interpreted later on as a factor contributing to the counting of lattice walks. Again, formula (4) is valid for any set of positive or null integers l_i ; if the l_i 's are all positive the bounds on A are as specified above.

We remark here that the trigonometric sum (1) reduces to the binomial multiple sum given in (3) in the case $b_{p/q}(k) = (2 \sin(\pi kp/q))^2$ only when $l_1 + \dots + l_j < q$, i.e., for large enough values of q . In view of the algebraic area enumeration of square lattice walks, where the algebraic area counting ends up being the coefficient of $e^{i\pi Ap/q}$ (see (9) below), this constraint on q eliminates open walks which could be confused with closed ones by periodicity².

In [3] we revisited the algebraic area enumeration of [1] and noted that it admits a statistical mechanical interpretation in terms of particles obeying generalized *exclusion statistics* [4] with exclusion parameter $g = 2$ ($g = 0$ for bosons, $g = 1$ for fermions, higher g means a stronger exclusion beyond Fermi). Other lattice walks admit a similar interpretation with higher integer values of g . We also introduced the notion of g -compositions where some zeros can be inserted at will inside the set of the l_i 's with the restriction that no more than $g - 2$ zeros lay in succession. The integer n admits g^{n-1} such compositions. In particular, $g = 1$ -exclusion refers to the unique composition $n = n$, whereas $g = 2$ -exclusion corresponds to the standard compositions with no zeros at all. We also constructed triangular lattice chiral walks realizing $g = 3$ -exclusion with spectral function

$$b_{p/q}(k) = (2 \sin(2\pi kp/q)) (2 \sin(2\pi(k+1)p/q)) \quad (5)$$

We finally hinted at other walks corresponding to statistics with higher values of the exclusion parameter g and to other spectral functions. However, for the triangular lattice chiral walks, as well as for other cases, an explicit algebraic area enumeration formula

¹This binomial counting can be easily checked by first summing over A and subsequently over the k_i 's, redefining them appropriately; see [1].

²Extrapolating (3) as such to any value of $q \geq 1$ would amount to enforcing, for any given integer l , the identity $\sum_{k=1}^q e^{2ik\pi l/q} = 0$ even though this is valid only when l is not a multiple of q (when l is a multiple of q the sum is actually equal to q).

was missing due to the lack of binomial expressions analogous to (3) for the triangular spectral function (5).

In the present work we focus on filling this gap by uncovering such expressions for entire classes of trigonometric spectral functions generalizing (2) and (5). Namely, we consider, on the one hand

$$b_{p/q}(k) = (2 \sin(\pi k p/q))^r \quad (6)$$

and on the other hand

$$b_{p/q}(k) = (2 \sin(\pi k p/q))(2 \sin(\pi(k+1)p/q)) \dots (2 \sin(\pi(k+r-1)p/q)) \quad (7)$$

where in both instances r can be even or odd. The case $r = 2$ reproduces³ (2) and (5) respectively. We will see that the basic structure of the binomial multiple sum (3) naturally generalizes to these cases. In the Appendix we will also derive the relevant generalization for the spectral function

$$b_{p/q}(k) = (2 \sin(\pi k p/q))^{r/2} (2 \sin(\pi(k+1)p/q))^{r/2} \quad (8)$$

where r is even, yet another possible generalization of (5).

Turning to the algebraic area combinatorics *per se*, these expressions, as already mentioned, will allow for explicit enumeration formulae analogous to the square lattice walks formula obtained in [1] for $g = 2$ and the Hofstadter spectral function (2). This requires introducing an appropriate weighting coefficient in the summation over compositions of the integer n . We refer to [1] for detailed explanations of how this procedure unfolds and to [3] for the connection to g -exclusion statistics and the resulting generalizations. With the g -exclusion statistics weighting coefficients [3]

$$\begin{aligned} c_g(l_1, l_2, \dots, l_j) &= \frac{(l_1 + \dots + l_{g-1} - 1)!}{l_1! \dots l_{g-1}!} \prod_{i=1}^{j-g+1} \binom{l_i + \dots + l_{i+g-1} - 1}{l_{i+g-1}} \\ &= \frac{\prod_{i=1}^{j-g+1} (l_i + \dots + l_{i+g-1} - 1)!}{\prod_{i=1}^{j-g} (l_{i+1} + \dots + l_{i+g-1} - 1)!} \prod_{i=1}^j \frac{1}{l_i!} \end{aligned}$$

we can express the lattice walks algebraic area enumeration for $g \geq 2$ -exclusion and a general periodic spectral function $b_{p/q}(k)$ by means of the g -cluster coefficient⁴

$$b(n) = gn \sum_{\substack{l_1, l_2, \dots, l_j \\ g\text{-composition of } n}} c_g(l_1, l_2, \dots, l_j) \frac{1}{q} \sum_{k=1}^q b_{p/q}^{l_1}(k) b_{p/q}^{l_2}(k-1) \dots b_{p/q}^{l_j}(k-j+1) \quad (9)$$

As already stressed, (9) yields the algebraic area combinatorics provided that an expression analogous to (3) is known for the specific $b_{p/q}(k)$. Indeed, the summation index A

³The actual spectral function (5) for triangular lattice chiral walks has a factor 2 in front of the π 's which we omit here to stay in line with (6); it anyway amounts to a trivial redefinition of $p/q \rightarrow 2p/q$.

⁴For statistical mechanics considerations the g -cluster coefficient introduced in [3] is the expression in (9) multiplied by $(-1)^{n-1}q/(gn)$.

in (3) has to be interpreted in (9) as the algebraic area, and the coefficient multiplying the exponential factor $e^{i\pi Ap/q}$ is the sought for algebraic area counting number. It will, in particular, yield the triangular lattice chiral walk counting described by $g = 3$ -exclusion and spectral function (5).

Finally, we will discuss the unexpected occurrence of Apéry-like numbers in the cluster coefficient (9) evaluated at particular values of p/q for certain g -exclusions and spectral functions. Apéry-like numbers are interesting *per se* since they are cousins of the celebrated Apéry numbers which allow for a proof of the irrationality of $\zeta(2)$ and $\zeta(3)$. One key characteristic of these numbers is that they are integer solutions of second order recursion relations. As we will see, some of the $\zeta(2)$ Apéry-like numbers fascinatingly emerge in the algebraic enumeration formula (9).

2 Trigonometric sums $\sum_{k=1}^q b_{p/q}^{l_1}(k) b_{p/q}^{l_2}(k+1) \cdots b_{p/q}^{l_j}(k+j-1)$

We aim at uncovering explicit binomial multiple sums analogous to (3) for the spectral functions (6) and (7). In fact, the form of (3) is quite robust and suggestive, and allows deducing such generalizations by simple deformations while preserving its overall structure. We stress that, from now on, some l_i 's can be null according to the g -composition structure discussed previously, i.e., no more than $g - 2$ zeros in succession inside the set. The A -summation bounds, when specified, will explicitly depend on the parameter g .

2.1 Square lattice walks generalization: $b_{p/q}(k) = (2 \sin(\pi kp/q))^r$

We first list two basic facts:

- When $q \rightarrow \infty$ one obtains the overall counting

$$\int_0^1 (2 \sin(\pi s))^{r l_1 + l_2 + \dots + r l_j} ds = \binom{r(l_1 + l_2 + \dots + l_j)}{r(l_1 + l_2 + \dots + l_j)/2} \quad (10)$$

so we focus on $(l_1 + l_2 + \dots + l_j)$ such that $r(l_1 + l_2 + \dots + l_j)$ be even. It means that for r even any set l_1, l_2, \dots, l_j is admissible, whereas for r odd the l_i 's have to be such that their sum be even.

- It is obvious that for a given r

$$\frac{1}{q} \sum_{k=1}^q ((2 \sin(\pi kp/q))^r)^{l_1} ((2 \sin(\pi(k+1)p/q))^r)^{l_2} \dots ((2 \sin(\pi(k+j-1)p/q))^r)^{l_j}$$

amounts to

$$\frac{1}{q} \sum_{k=1}^q \left((2 \sin(\pi k p/q))^2 \right)^{rl_1/2} \left((2 \sin(\pi(k+1)p/q))^2 \right)^{rl_2/2} \dots \left((2 \sin(\pi(k+j-1)p/q))^2 \right)^{rl_j/2}$$

which is essentially the Hofstadter case $r = 2$, i.e., for the spectral function $(2 \sin(\pi k p/q))^2$, but now with $l_i \rightarrow rl_i/2$.

Based on the above observations, the binomial multiple sum in (3) for the $r = 2$ Hofstadter case becomes, for $b_{p/q}(k) = (2 \sin(\pi k p/q))^r$ with r even⁵,

$$\frac{1}{q} \sum_{k=1}^q b_{p/q}^{l_1}(k) b_{p/q}^{l_2}(k+1) \dots b_{p/q}^{l_j}(k+j-1) = \sum_{\substack{A=-(g-1)r[(l_1+\dots+l_j)^2/4] \\ A \text{ even}}}^{(g-1)r[(l_1+\dots+l_j)^2/4]} e^{i\pi A p/q} \quad (11)$$

$$\sum_{k_3=-rl_3/2}^{rl_3/2} \dots \sum_{k_j=-rl_j/2}^{rl_j/2} \binom{rl_1}{rl_1/2 + A/2 + \sum_{i=3}^j (i-2)k_i} \binom{rl_2}{rl_2/2 - A/2 - \sum_{i=3}^j (i-1)k_i} \prod_{i=3}^j \binom{rl_i}{rl_i/2 + k_i}$$

which is valid when $r(l_1 + \dots + l_j)/2 < q$ holds, and where we have specified the range $[-(g-1)r[(l_1 + \dots + l_j)^2/4], (g-1)r[(l_1 + \dots + l_j)^2/4]]$ in which A needs to be restricted.

In the r odd case we expect a binomial multiple sum analogous to (11). To see this in full generality, and to give a full proof of the original formula with even r , let us first recall the Poisson summation formula for any q -periodic function $f(x) = f(x+q)$

$$\sum_{k=1}^q f(k) = \sum_{n=-\infty}^{\infty} \tilde{f}(nq) \quad (12)$$

where \tilde{f} is the Fourier transform of f defined as

$$\tilde{f}(k) = \int_0^q f(x) e^{-2i\pi k x/q} dx, \quad f(x) = \frac{1}{q} \sum_{k=-\infty}^{\infty} \tilde{f}(k) e^{2i\pi k x/q}$$

⁵The overall counting, found by replacing $e^{i\pi A p/q}$ by 1 is

$$\binom{r(l_1 + l_2 + \dots + l_j)}{r(l_1 + l_2 + \dots + l_j)/2} = \sum_{\substack{A=-(g-1)r[(l_1+\dots+l_j)^2/4] \\ A \text{ even}}}^{(g-1)r[(l_1+\dots+l_j)^2/4]}$$

$$\sum_{k_3=-rl_3/2}^{rl_3/2} \dots \sum_{k_j=-rl_j/2}^{rl_j/2} \binom{rl_1}{rl_1/2 + A/2 + \sum_{i=3}^j (i-2)k_i} \binom{rl_2}{rl_2/2 - A/2 - \sum_{i=3}^j (i-1)k_i} \prod_{i=3}^j \binom{rl_i}{rl_i/2 + k_i}$$

Let us consider the function $f(x) = \frac{1}{q} b_{p/q}^{l_1}(x) b_{p/q}^{l_2}(x+1) \dots b_{p/q}^{l_j}(x+j-1)$ which is indeed q -periodic due to $r(l_1 + l_2 + \dots + l_j)$ being always assumed even. We have

$$\begin{aligned}
\tilde{f}(nq) &= \int_0^q f(k) e^{-2i\pi kn} dk \\
&= \frac{1}{q} \int_0^q b_{p/q}^{l_1}(k) b_{p/q}^{l_2}(k+1) \dots b_{p/q}^{l_j}(k+j-1) e^{-2i\pi kn} dk \\
&= \frac{1}{q} \int_0^q \prod_{i=1}^j \frac{1}{i^{rl_i}} \left(e^{i\pi(k+i-1)p/q} - e^{-i\pi(k+i-1)p/q} \right)^{rl_i} e^{-2i\pi kn} dk \\
&= \frac{1}{q} \int_0^q \prod_{i=1}^j \frac{1}{i^{rl_i}} \sum_{k_i=-rl_i/2}^{rl_i/2} \binom{rl_i}{rl_i/2 + k_i} e^{2i\pi(k+i-1)k_i p/q} (-1)^{rl_i/2 - k_i} e^{-2i\pi kn} dk \\
&= \sum_{k_1=-rl_1/2}^{rl_1/2} \dots \sum_{k_j=-rl_j/2}^{rl_j/2} \prod_{i=1}^j \binom{rl_i}{rl_i/2 + k_i} \frac{(-1)^{rl_i/2 - k_i}}{i^{rl_i}} \int_0^1 e^{2i\pi \sum_{i=1}^j k_i s p + 2i\pi \sum_{i=1}^j (i-1)k_i p/q} e^{-2i\pi s q n} ds \\
&= \sum_{k_1=-rl_1/2}^{rl_1/2} \dots \sum_{k_j=-rl_j/2}^{rl_j/2} \prod_{i=1}^j \binom{rl_i}{rl_i/2 + k_i} \frac{(-1)^{rl_i/2 - k_i}}{i^{rl_i}} e^{2i\pi \sum_{i=1}^j (i-1)k_i p/q} \delta\left(\sum_{i=1}^j k_i p - nq\right)
\end{aligned} \tag{13}$$

As stressed above, $r(l_1 \dots + l_j)$ is even and thus the sum of the k_i is an integer. Further, p and q are coprime. These facts imply that the Kronecker- δ in (13) enforces

$$p \sum_{i=1}^j k_i = qn \quad \text{and thus} \quad \sum_{i=1}^j k_i = tq \quad \text{and} \quad n = tp$$

for some integer t . Now $|\sum_{i=1}^j k_i| \leq r(l_1 + \dots + l_j)/2$ and thus, under the condition $r(l_1 + \dots + l_j)/2 < q$, t is necessarily equal to 0, implying that $\sum_{i=1}^j k_i = 0$ and $n = 0$. From the Poisson summation formula (12) then we infer $\sum_{k=1}^q f(k) = \tilde{f}(0) = \int_0^q f(x) dx$; that is, for $b_{p/q}(k) = (2 \sin(\pi k p/q))^r$,

$$\frac{1}{q} \sum_{k=1}^q b_{p/q}^{l_1}(k) b_{p/q}^{l_2}(k+1) \dots b_{p/q}^{l_j}(k+j-1) = \frac{1}{q} \int_0^q b_{p/q}^{l_1}(k) b_{p/q}^{l_2}(k+1) \dots b_{p/q}^{l_j}(k+j-1) dk \tag{14}$$

What has been achieved in (14) is the trading of the original sum over k from 1 to q in the LHS for the integral over k from 0 to q in the RHS, which is valid provided that $r(l_1 + \dots + l_j)/2 < q$.

We can easily check that the trigonometric integral yields the binomial multiple sum (3) in the $r = 2$ case, or more generally (11) in the r even case. To do so let us proceed

from the last line of (13): enforcing the Kronecker δ in the summand we obtain

$$\begin{aligned}
& \frac{1}{q} \int_0^q b_{p/q}^{l_1}(k) b_{p/q}^{l_2}(k+1) \dots b_{p/q}^{l_j}(k+j-1) dk \\
&= \frac{1}{q} \int_0^q dk \prod_{i=1}^j \left(2 \sin(\pi k p/q + \pi(i-1)p/q) \right)^{rl_i} = \int_0^1 dt \prod_{i=1}^j \left(2 \sin(\pi t + \pi(i-1)p/q) \right)^{rl_i} \\
&= \sum_{k_1=-rl_1/2}^{rl_1/2} \dots \sum_{k_j=-rl_j/2}^{rl_j/2} \prod_{i=1}^j \binom{rl_i}{rl_i/2 + k_i} e^{2i\pi \sum_{i=1}^j (i-1)k_i p/q} \tag{15}
\end{aligned}$$

The change of integration from $(1/q) \int_0^q dk$ to $\int_0^1 dt$ in the variable $t = kp/q$ in the second line is justified since $r(l_1 + \dots + l_j)$ is even and the integrand has period 1 in t . We still need to enforce the constraint $\sum_{i=1}^j k_i = 0$ in the summation variables k_i . To reproduce the A -expansion with exponential factors $e^{i\pi A p/q}$ in the binomial multiple sums (3) and (11), we denote by A the coefficient $2 \sum_{i=1}^j (i-1)k_i$ of $i\pi p/q$ appearing in the exponential of the last line in (15). The resulting system of two equations, $\sum_{i=1}^j k_i = 0$ and $A = 2 \sum_{i=1}^j (i-1)k_i$, can be readily solved for, e.g., the first two variables k_1 and k_2 , to yield

$$k_1 = -A/2 + \sum_{i=3}^j (i-2)k_i \quad , \quad k_2 = A/2 - \sum_{i=3}^j (i-1)k_i$$

Finally, changing summation variables from k_i to $-k_i$ and noting that each binomial is invariant under changing the sign of k_i , we obtain

$$\begin{aligned}
& \frac{1}{q} \sum_{k=0}^q b_{p/q}^{l_1}(k) b_{p/q}^{l_2}(k+1) \dots b_{p/q}^{l_j}(k+j-1) \\
&= \int_0^1 dt \prod_{i=1}^j \left(2 \sin(\pi t + \pi(i-1)p/q) \right)^{rl_i} \\
&= \sum_{\substack{A=-(g-1)r[(l_1+\dots+l_j)^2/4] \\ \text{in steps of 2}}}^{(g-1)r[(l_1+\dots+l_j)^2/4]} e^{i\pi A p/q} \sum_{k_3=-rl_3/2}^{rl_3/2} \dots \sum_{k_j=-rl_j/2}^{rl_j/2} \tag{16} \\
& \left(\binom{rl_1}{rl_1/2 + A/2 + \sum_{i=3}^j (i-2)k_i} \right) \left(\binom{rl_2}{rl_2/2 - A/2 - \sum_{i=3}^j (i-1)k_i} \right) \prod_{i=3}^j \binom{rl_i}{rl_i/2 + k_i}
\end{aligned}$$

i.e., precisely (11) but now valid for r even and r odd, with a specific A -summation dictated by the condition that in (16) the first two binomial entries $rl_1/2 + A/2 + \sum_{i=3}^j (i-2)k_i$ and $rl_2/2 - A/2 - \sum_{i=3}^j (i-1)k_i$ still take integer values for all $k_i \in [-rl_i/2, rl_i/2]$, $i = 3, \dots, j$, as was the case in (15) for the first two binomial entries $rl_1/2 + k_1$ and $rl_2/2 + k_2$ for all $k_1 \in [-rl_1/2, rl_1/2]$ and $k_2 \in [-rl_2/2, rl_2/2]$. It follows that in the case r even, where the k_i 's are all integers, A has to be even, and in the case r odd, where the k_i 's are either integers or half integers, $l_1 + l_2 + \dots + l_j$ has to be even and

A of the same parity as $l_1 + l_3 + \dots$ (or $l_2 + l_4 + \dots$). In both cases this boils down to $A \in [-(g-1)r[(l_1 + \dots + l_j)^2/4], (g-1)r[(l_1 + \dots + l_j)^2/4]]$ in steps of 2. We also note that, in this and all subsequent formulae, we follow the convention that the sum of all the lower entries in the binomials in (16) be zero, which fixes the form of such expressions among various equivalent parametrizations.

We can express the A -binomial block in (16) in an integral form by augmenting the LHS to the double integral $\frac{1}{2} \int_0^1 dt \int_0^2 dt' \prod_{i=1}^j \left(2 \sin(\pi t + \pi(i-1)t') \right)^{r l_i} \delta(p/q - t')$ and using $2 \sum_{n=-\infty}^{\infty} \delta(p/q - t' - 2n) = \sum_{A=-\infty}^{\infty} e^{i\pi A(p/q - t')}$ to get

$$\begin{aligned} & \frac{1}{2} \int_0^2 dt' \int_0^1 dt \prod_{i=1}^j \left(2 \sin(\pi t + \pi(i-1)t') \right)^{r l_i} e^{i\pi A t'} \tag{17} \\ &= \sum_{k_3=-r l_3/2}^{r l_3/2} \cdots \sum_{k_j=-r l_j/2}^{r l_j/2} \binom{r l_1}{r l_1/2 + A/2 + \sum_{i=3}^j (i-2)k_i} \binom{r l_2}{r l_2/2 - A/2 - \sum_{i=3}^j (i-1)k_i} \prod_{i=3}^j \binom{r l_i}{r l_i/2 + k_i} \end{aligned}$$

In the multiple sum of the RHS A is constrained as above, depending on r being even or odd. However, the integral in the LHS is valid for all integer values of A , yielding zero for the values that do not appear in the RHS.

2.2 Triangular generalization: $b_{p/q}(k) = (2 \sin(\pi k p/q)) (2 \sin(\pi(k+1)p/q)) \dots (2 \sin(\pi(k+r-1)p/q))$

We can proceed in exactly the same way for triangular-like spectral functions of the type $b_{p/q}(k) = (2 \sin(\pi k p/q)) (2 \sin(\pi(k+1)p/q)) \dots (2 \sin(\pi(k+r-1)p/q))$. Again

- $q \rightarrow \infty$ recovers the overall counting

$$\int_0^1 (2 \sin(\pi s))^{r l_1 + r l_2 + \dots + r l_j} ds = \binom{r(l_1 + l_2 + \dots + l_j)}{r(l_1 + l_2 + \dots + l_j)/2}$$

as in (10), so we still focus on sets of l_i 's such that $r(l_1 + l_2 + \dots + l_j)$ is even, again ensuring the q -periodicity of the functions at hand

- The rewriting of the trigonometric sum as a trigonometric integral follows the same lines as in (13) under the same condition $r(l_1 + \dots + l_j)/2 < q$ since the sole input in this condition is the highest power of $e^{i\pi k p/q}$ that appears in $b_{p/q}(k)$ given by (7), which happens to be again r

2.2.1 Triangular chiral walks $r = 2$: $b_{p/q}(k) = (2 \sin(\pi kp/q))(2 \sin(\pi(k+1)p/q))$

Following the same steps as in **2.1**, we can rewrite the trigonometric sum corresponding to $b_{p/q}(k) = (2 \sin(\pi kp/q))(2 \sin(\pi(k+1)p/q))$ as the simple integral

$$\begin{aligned} & \frac{1}{q} \sum_{k=1}^q b_{p/q}^{l_1}(k) b_{p/q}^{l_2}(k+1) \dots b_{p/q}^{l_j}(k+j-1) = \\ & \int_0^1 dt \left(2 \sin(\pi t) \right)^{l_1} \prod_{i=2}^j \left(2 \sin(\pi t + \pi(i-1)p/q) \right)^{l_{i-1}+l_i} \left(2 \sin(\pi t + \pi j p/q) \right)^{l_j} \end{aligned} \quad (18)$$

provided that $l_1 + \dots + l_j < q$.

Integrating (18) leads to the appropriate deformation of the binomial multiple sum (3) for the spectral function $b_{p/q}(k) = (2 \sin(\pi kp/q))(2 \sin(\pi(k+1)p/q))$, a deformation which could also have been directly guessed by simple manipulations: in (1) the integer l_1 is associated with the index k , $l_1 + l_2$ with $k+1$, $l_2 + l_3$ with $k+2$, etc. This leads to⁶

$$\begin{aligned} & \frac{1}{q} \sum_{k=1}^q b_{p/q}^{l_1}(k) b_{p/q}^{l_2}(k+1) \dots b_{p/q}^{l_j}(k+j-1) \\ & = \int_0^1 dt \left(2 \sin(\pi t) \right)^{l_1} \prod_{i=2}^j \left(2 \sin(\pi t + \pi(i-1)p/q) \right)^{l_{i-1}+l_i} \left(2 \sin(\pi t + \pi j p/q) \right)^{l_j} \\ & = \sum_{\substack{A = -\lceil (l_1 + \dots + l_j)^2/2 \rceil - (g-2)\lfloor (l_1 + \dots + l_j)^2/2 \rfloor \\ \text{A same parity as } l_1 + l_2 + \dots + l_j}} e^{i\pi A p/q} \sum_{k_3 = -(l_2 + l_3)/2}^{(l_2 + l_3)/2} \dots \sum_{k_j = -(l_{j-1} + l_j)/2}^{(l_{j-1} + l_j)/2} \sum_{k_{j+1} = -l_j/2}^{l_j/2} \\ & \left(\binom{l_1}{l_1/2 + A/2 + \sum_{i=3}^{j+1} (i-2)k_i} \right) \left(\binom{l_1 + l_2}{(l_1 + l_2)/2 - A/2 - \sum_{i=3}^{j+1} (i-1)k_i} \right) \\ & \times \prod_{i=3}^j \left(\binom{l_{i-1} + l_i}{(l_{i-1} + l_i)/2 + k_i} \right) \left(\binom{l_j}{l_j/2 + k_{j+1}} \right) \end{aligned} \quad (19)$$

We note that A in the summation (19) spans the interval $[-\lceil (l_1 + \dots + l_j)^2/2 \rceil - (g-2)\lfloor (l_1 + \dots + l_j)^2/2 \rfloor, \lceil (l_1 + \dots + l_j)^2/2 \rceil + (g-2)\lfloor (l_1 + \dots + l_j)^2/2 \rfloor]$ increasing by steps of 2, which in particular implies that A is of the same parity as $l_1 + l_2 + \dots + l_j$.

⁶With overall counting, obtained in the $q \rightarrow \infty$ limit by replacing $e^{i\pi A p/q}$ by 1:

$$\binom{2(l_1 + l_2 + \dots + l_j)}{l_1 + l_2 + \dots + l_j}$$

2.2.2 $r = 3$: $b_{p/q}(k) = (2 \sin(\pi k p/q)) (2 \sin(\pi(k+1)p/q)) (2 \sin(\pi(k+2)p/q))$
with $l_1 + \dots + l_j$ even

Similarly to the previous cases one can rewrite the $r = 3$ triangular trigonometric sum as the simple integral

$$\begin{aligned} & \frac{1}{q} \sum_{k=1}^q b_{p/q}^{l_1}(k) b_{p/q}^{l_2}(k+1) \dots b_{p/q}^{l_j}(k+j-1) = \\ & \int_0^1 dt \left(2 \sin(\pi t) \right)^{l_1} \left(2 \sin(\pi t + \pi p/q) \right)^{l_1+l_2} \prod_{i=3}^j \left(2 \sin(\pi t + \pi(i-1)p/q) \right)^{l_{i-2}+l_{i-1}+l_i} \\ & \times \left(2 \sin(\pi t + \pi j p/q) \right)^{l_{j-1}+l_j} \left(2 \sin(\pi t + \pi(j+1)p/q) \right)^{l_j} \end{aligned}$$

provided that $3(l_1 + \dots + l_j)/2 < q$.

Likewise one obtains the binomial multiple sum⁷

$$\begin{aligned} & \frac{1}{q} \sum_{k=1}^q b_{p/q}^{l_1}(k) b_{p/q}^{l_2}(k+1) \dots b_{p/q}^{l_j}(k+j-1) = \sum_{\substack{A=-\infty \\ \text{A same parity as } l_1+l_3+\dots \text{ or } l_2+l_4+\dots}}^{\infty} e^{i\pi A p/q} \\ & \sum_{k_3=-(l_1+l_2+l_3)/2}^{(l_1+l_2+l_3)/2} \dots \sum_{k_j=-(l_{j-2}+l_{j-1}+l_j)/2}^{(l_{j-2}+l_{j-1}+l_j)/2} \sum_{k_{j+1}=-(l_{j-1}+l_j)/2}^{(l_{j-1}+l_j)/2} \sum_{k_{j+2}=-l_j/2}^{l_j/2} \\ & \left(\binom{l_1}{l_1/2 + A/2 + \sum_{i=3}^{j+2} (i-2)k_i} \right) \left(\binom{l_1+l_2}{(l_1+l_2)/2 - A/2 - \sum_{i=3}^{j+2} (i-1)k_i} \right) \\ & \times \prod_{i=3}^j \left(\binom{l_{i-2}+l_{i-1}+l_i}{(l_{i-2}+l_{i-1}+l_i)/2 + k_i} \right) \left(\binom{l_{j-1}+l_j}{(l_{j-1}+l_j)/2 + k_{j+1}} \right) \left(\binom{l_j}{l_j/2 + k_{j+2}} \right) \end{aligned} \quad (20)$$

where A has to be of the same parity as $l_1 + l_3 + \dots$ (or $l_2 + l_4 + \dots$) and obviously a finite range. The cases $r = 4$ and beyond are treated in the Appendix.

3 Algebraic area enumeration and Apéry-like numbers

3.1 Algebraic area enumeration

We can retrieve from the cluster coefficient (9) algebraic area enumeration formulae for various random lattice walks. For example, from (16) for $b_{p/q}(k) = (2 \sin(\pi k p/q))^r$ with

⁷With overall counting, obtained by replacing $e^{i\pi A p/q}$ by 1:

$$\left(\begin{array}{c} 3(l_1 + l_2 + \dots + l_j) \\ 3(l_1 + l_2 + \dots + l_j)/2 \end{array} \right)$$

r even and g -exclusion, (9) becomes

$$b(n) = gn \sum_{\substack{A=-(g-1)r\lfloor(l_1+\dots+l_j)^2/4\rfloor \\ A \text{ even}}}^{(g-1)r\lfloor(l_1+\dots+l_j)^2/4\rfloor} e^{i\pi Ap/q} \sum_{\substack{l_1, l_2, \dots, l_j \\ g\text{-composition of } n}} c_g(l_1, l_2, \dots, l_j) \quad (21)$$

$$\sum_{k_3=-rl_3/2}^{rl_3/2} \cdots \sum_{k_j=-rl_j/2}^{rl_j/2} \binom{rl_1}{rl_1/2 + A/2 + \sum_{i=3}^j (i-2)k_i} \binom{rl_2}{rl_2/2 - A/2 - \sum_{i=3}^j (i-1)k_i} \prod_{i=3}^j \binom{rl_i}{rl_i/2 + k_i}$$

with overall counting, given by replacing $e^{i\pi Ap/q}$ by 1

$$\binom{gn}{n} \binom{rn}{rn/2} \quad (22)$$

The second binomial in (22), as initially discussed in (4) and displayed in the various overall counting cases of subsection (2.1), results from the trigonometric sums replacing $e^{i\pi Ap/q}$ by 1 in the limit $q \rightarrow \infty$, whereas the first one results from the summation of the exclusion weight coefficients c_g over all g -compositions of the integer n .

3.1.1 Square lattice walks: $b_{p/q}(k) = (2 \sin(\pi kp/q))^2$

As already stated, the standard square lattice walks are specifically $g = 2$ and $r = 2$ and are defined in terms of the Hamiltonian [3]

$$H = (1 - u)v + v^{-1}(1 - u^{-1})$$

where u and v respectively stand for the *right* and *up* hopping operators on the lattice, with commutation $vu = Q uv$, where $Q = e^{i\Phi} = e^{i2\pi p/q}$ is the noncommutativity parameter encoding the presence of the magnetic field perpendicular to the lattice, with Φ the magnetic flux per plaquette. We recover the Hofstadter spectral function as

$$b_{p/q}(k) = (1 - Q^{-k})(1 - Q^k) = (2 \sin(\pi kp/q))^2$$

The Hamiltonian describes a random walk with elementary steps *up*, *right* followed by *up*, *down*, and *down* followed by *left*. It means that starting from the origin $(0, 0)$ it reaches after one step the lattice points $(0, 1)$, $(1, 1)$, $(0, -1)$ or $(-1, -1)$ with equal probability. This generates deformed walks on the square lattice (see Fig.1) which are equivalent through a modular transformation to the usual square lattice walks. (This modular transformation amounts to the transformation $u \rightarrow -uv$, which leaves the u, v commutation relation unchanged and turns H into $u + v + u^{-1} + v^{-1}$.) $b(n)$ in (21) then yields the desired algebraic area counting [1]

$$b(n) = \sum_{\substack{A=2\lfloor n^2/4\rfloor \\ A \text{ even}}}^{2\lfloor n^2/4\rfloor} e^{i\pi Ap/q} C'_{2n}(A)$$

where

$$C_{2n}(A) = 2n \sum_{\substack{l_1, l_2, \dots, l_j \\ \text{2-composition of } n}} c_2(l_1, l_2, \dots, l_j) \quad (23)$$

$$\sum_{k_3=-l_3}^{l_3} \cdots \sum_{k_j=-l_j}^{l_j} \binom{2l_1}{l_1 + A/2 + \sum_{i=3}^j (i-2)k_i} \binom{2l_2}{l_2 - A/2 - \sum_{i=3}^j (i-1)k_i} \prod_{i=3}^j \binom{2l_i}{l_i + k_i}$$

with A even in the interval $[-2\lfloor n^2/4 \rfloor, 2\lfloor n^2/4 \rfloor]$. $C_{2n}(A)$ counts the number of closed square lattice walks of length $2n$ —there are overall $\binom{2n}{n}^2$ of them, see (22)—enclosing an algebraic area $A/2$ in the interval⁸ $[-\lfloor n^2/4 \rfloor, \lfloor n^2/4 \rfloor]$: indeed the mapping of random walk algebraic area to the Hofstadter model [1] is via the weighting factor $Q^{\text{algebraic area}}$, where $Q = e^{2i\pi p/q}$, so here, with $e^{i\pi Ap/q}$ appearing in (21), the algebraic area is $A/2$.

3.1.2 Square lattice walks: $b_{p/q}(k) = (2 \sin(\pi kp/q))^4$

Let us now look at square lattice walks with $g = 2$ and $r = 4$ which are defined in terms of the Hamiltonian

$$H = (u + u^{-1})^2 v + v^{-1} (u + u^{-1})^2 \quad (24)$$

The corresponding spectral function

$$b_{p/q}(k) = (Q^k + Q^{-k})^4 = (2 \cos(2\pi kp/q))^4$$

can be put in the standard form (6) for $r = 4$ by redefining $u \rightarrow iu$ and $Q \rightarrow \sqrt{Q}$, which does not affect the counting of walks nor the area weighting.

The Hamiltonian (24) describes a random walk with elementary steps in groups of one random step *up* or *down* and two independent random steps *right* or *left*. It means that starting from the origin $(0,0)$ it reaches after one step the lattice points $(2,1)$, $(-2,1)$, $(2,-1)$ or $(-2,-1)$ with probability $1/8$, or the lattice points $(1,0)$ or $(-1,0)$ with probability $1/4$. The same walk can be described as a particle hopping on an *even* or *odd* square sublattice, where even points are those with x and y coordinates adding to an even integer, the remaining being odd. The walk proceeds randomly on one of the sublattices but at each step it has the option to move to the nearest up or down point of the opposite sublattice, with each such jump contributing a factor of two in the weight of the walk. The Hamiltonian (24) counts the weighted number of such closed walks of a given total area.

There are $\binom{2n}{n} \binom{4n}{2n}$ such closed walks of length $2n$, as in (22). The enumeration of such walks enclosing a given algebraic area, with the proper weight, is given by (21):

$$b(n) = \sum_{\substack{A=-4\lfloor n^2/4 \rfloor \\ A \text{ even}}^{4\lfloor n^2/4 \rfloor}} e^{i\pi Ap/q} C'_{2n}(A)$$

⁸This can be easily seen geometrically for lattice walks of length $2n$ with n even, which have largest possible area $\pm(n/2)^2$: this is the walk circling a square of side $n/2$ anti-clockwise or clockwise.

where

$$C'_{2n}(A) = 2n \sum_{\substack{l_1, l_2, \dots, l_j \\ \text{2-composition of } n}} c_2(l_1, l_2, \dots, l_j) \\ \sum_{k_3=-2l_3}^{2l_3} \cdots \sum_{k_j=-2l_j}^{2l_j} \binom{4l_1}{2l_1+A/2 + \sum_{i=3}^j (i-2)k_i} \binom{4l_2}{2l_2-A/2 - \sum_{i=3}^j (i-1)k_i} \prod_{i=3}^j \binom{4l_i}{2l_i + k_i}$$

with A even in the interval $[-4\lfloor n^2/4 \rfloor, 4\lfloor n^2/4 \rfloor]$. $C'_{2n}(A)$ counts the number of closed square lattice walks described above of length $2n$ and enclosing an algebraic area $A/2$.

3.1.3 Square lattice walks: $b_{p/q}(k) = \left((2 \sin(\pi k p/q)) (2 \sin(\pi(k+1)p/q)) \right)^2$

Now consider square lattice walks with $g = 2$ and $r = 4$ defined by the Hamiltonian

$$H = (u + u^{-1})v(u + u^{-1}) + (u + u^{-1})v^{-1}(u + u^{-1}) \quad (25)$$

The spectral function can be brought to the standard form (8) for $r = 4$ by an appropriate redefinition of $u \rightarrow -iu$

$$b_{p/q}(k) = (2 \sin(\pi k p/q))^2 (2 \sin(\pi(k+1)p/q))^2$$

Its treatment is given in the subsection 5.2 of the Appendix.

This walk proceeds with sets of one step left or right, one step up or down and another step left or right. With an appropriate redefinition of u and v (modular transformation) this walk can also be mapped to a walk proceeding on odd or even square sublattices, as in the last subsection, but now the weight of jumping on the opposite sublattice is not 2, as before, but rather $Q + Q^{-1}$. So in this description the weight of the walks depends explicitly on Q , unlike any other walk we encountered before.

There are again $\binom{2n}{n} \binom{4n}{2n}$ such closed walks of length $2n$. The enumeration of such walks enclosing a given algebraic area, with the proper weight, is given by

$$b(n) = \sum_{\substack{A=-\infty \\ A \text{ even}}}^{\infty} e^{i\pi A p/q} C''_{2n}(A)$$

where

$$C''_{2n}(A) = 2n \sum_{\substack{l_1, l_2, \dots, l_j \\ \text{2-composition of } n}} c_2(l_1, l_2, \dots, l_j) \\ \sum_{k_3=-(l_2+l_3)}^{l_2+l_3} \cdots \sum_{k_j=-(l_{j-1}+l_j)}^{l_{j-1}+l_j} \sum_{k_{j+1}=-l_j}^{l_j} \binom{2l_1}{l_1+A/2 + \sum_{i=3}^{j+1} (i-2)k_i} \binom{2(l_1+l_2)}{l_1+l_2-A/2 - \sum_{i=3}^{j+1} (i-1)k_i} \\ \times \prod_{i=3}^j \binom{2(l_{i-1}+l_i)}{l_{i-1}+l_i+k_i} \binom{2l_j}{l_j+k_{j+1}} \quad (26)$$

$C''_{2n}(A)$ counts again the weighted number of closed square lattice walks described above of length $2n$ enclosing an algebraic area $A/2$. It differs from the corresponding number (25) only in the weighting factor when jumping sublattices.

3.1.4 Triangular lattice chiral walks: $b_{p/q}(k) = (2 \sin(\pi k p/q))(2 \sin(\pi(k+1)p/q))$

From (19) for the triangular spectral function (7) with $r = 2$ and g -exclusion we obtain

$$\begin{aligned}
b(n) = gn & \sum_{\substack{A = -\lceil n^2/2 \rceil - (g-2)\lfloor n^2/2 \rfloor \\ A \text{ same parity as } n}}^{\lceil n^2/2 \rceil + (g-2)\lfloor n^2/2 \rfloor} e^{i\pi A p/q} \sum_{\substack{l_1, l_2, \dots, l_j \\ g\text{-composition of } n}} c_g(l_1, l_2, \dots, l_j) \\
& \sum_{k_3 = -(l_2+l_3)/2}^{(l_2+l_3)/2} \cdots \sum_{k_j = -(l_{j-1}+l_j)/2}^{(l_{j-1}+l_j)/2} \sum_{k_{j+1} = -l_j/2}^{l_j/2} \\
& \binom{l_1}{l_1/2 + A/2 + \sum_{i=3}^{j+1} (i-2)k_i} \binom{l_1 + l_2}{(l_1 + l_2)/2 - A/2 - \sum_{i=3}^{j+1} (i-1)k_i} \\
& \times \prod_{i=3}^j \binom{l_{i-1} + l_i}{(l_{i-1} + l_i)/2 + k_i} \binom{l_j}{l_j/2 + k_{j+1}} \tag{27}
\end{aligned}$$

with overall counting given by replacing $e^{i\pi A p/q}$ by 1

$$\binom{gn}{n} \binom{2n}{n}$$

Triangular $g = 3$ lattice chiral walks correspond to the quantum Hamiltonian

$$H = i(-u + u^{-1})v + v^{-2}$$

with spectral function

$$b_{p/q}(k) = \left(2 \sin \frac{2\pi p k}{q}\right) \left(2 \sin \frac{2\pi p(k+1)}{q}\right)$$

as already given in (5). They are depicted in Figs.2–4 (see [3] for more details; these walks are the generalization to four quadrants of the Kreweras walks [5]). Since the exclusion parameter is $g = 3$ the counting above reduces to

$$\binom{3n}{n, n, n}$$

which is the number of closed triangular lattice chiral walks of length $3n$. The cluster coefficient (27) then yields the triangular lattice chiral walks algebraic area counting

$$b(n) = \sum_{\substack{A = -n^2 \\ A \text{ in steps of } 2}}^{n^2} e^{i\pi A p/q} C_{3n}(A)$$

where

$$\begin{aligned}
C_{3n}(A) = & 3n \sum_{\substack{l_1, l_2, \dots, l_j \\ \text{3-compositions of } n}} c_3(l_1, l_2, \dots, l_j) \sum_{k_3=-(l_2+l_3)/2}^{(l_2+l_3)/2} \dots \sum_{k_j=-(l_{j-1}+l_j)/2}^{(l_{j-1}+l_j)/2} \sum_{k_{j+1}=-l_j/2}^{l_j/2} \\
& \binom{l_1}{l_1/2 + A/2 + \sum_{i=3}^{j+1} (i-2)k_i} \binom{l_1 + l_2}{(l_1 + l_2)/2 - A/2 - \sum_{i=3}^{j+1} (i-1)k_i} \\
& \times \prod_{i=3}^j \binom{l_{i-1} + l_i}{(l_{i-1} + l_i)/2 + k_i} \binom{l_j}{l_j/2 + k_{j+1}} \tag{28}
\end{aligned}$$

with A in the interval $[-n^2, n^2]$ with same parity as n .

$C_{3n}(A)$ counts the number of closed triangular lattice chiral walks of length $3n$ enclosing an algebraic area A . Indeed, the mapping of triangular algebraic area-quantum triangular Hamiltonian discussed in [3] is via $\mathcal{Q}^{\text{algebraic area}}$ where $\mathcal{Q} = e^{2i\pi p/q}$. Since in $b_{p/q}(k)$ of (7) the building block $2 \sin(\pi k p/q)$ is used, rather than $2 \sin(2\pi k p/q)$ as in (5), we end up with $e^{i\pi A p/q}$ in (27) in place of $e^{2i\pi A p/q}$, so that the algebraic area is A . One can directly check by explicit enumeration that when n is odd A is also odd (see, e.g., $n = 1$ with 3 walks of algebraic area 1 and 3 walks of algebraic area -1) and when n is even A is also even (as in $n = 2$, with algebraic areas $0, \pm 2$ and ± 4).

We conclude our discussion of algebraic area counting by remarking that it was possible to extract explicit expressions in terms of binomial sums for $C_{2n}(A)$ in (23), $C'_{2n}(A)$ in (25) and $C_{3n}(A)$ in (28) from the cluster coefficients (21) or (27) because the summation constraints over A in the relevant binomial multiple sums (16) with $r = 2, 4$ (A even) or (19) with $r = 2$ (A same parity as n), as well as the summation ranges, depend only on n and not on the l_i 's themselves. Similar expressions would apply for walks deriving from odd r binomial sums, like (16) or (20), provided that the binomials appearing in the expressions are understood to vanish for values of A leading to noninteger entries, as discussed after (16).

It is a curious fact that if, in the binomial multiple sums or the cluster coefficients, we sum over *all* integer values of A without restrictions, and analytically continue the binomials to fractional values using Gamma functions, the resulting infinite sums are closely related to the finite ones over the allowed values of A . This point is detailed and explained in the subsection 5.3 of the Appendix. It means, considering for example the binomial multiple sum (16), that for even r and any set of l_i 's, the cumulative sum of the infinite sequence of coefficients of *odd* A , which are rational numbers times $1/\pi^2$, converges to the standard binomial counting $\binom{r(l_1+l_2+\dots+l_j)}{r(l_1+l_2+\dots+l_j)/2}$.

3.2 Apéry-like numbers

We finally turn to the occurrence of Apéry-like numbers in cluster coefficients (9) when evaluated at certain values of p/q . We stress that we no more view $b(n)$ as generating

algebraic area enumerations of actual lattice walks, but instead consider it as a stand-alone mathematical entity that happens to lead to such occurrences.

3.2.1 Apéry-like numbers $g = 2$ and $r = 2$: $b_{p/q}(k) = (2 \sin(\pi k p/q))^2$

Let us consider⁹ $b(n)$ in (21). For $g = 2$ and $r = 2$ it gives, for $n = 1, 2, 3, \dots$

$$\begin{aligned} p/q = 1 &\Rightarrow b(n) = \binom{2n}{n}^2 \Leftrightarrow \text{closed square lattice walks counting} \\ p/q = 1/2 &\Rightarrow b(n) = 4, 20, 112, 676, 4304, 28496, \dots \end{aligned}$$

These are the Apéry-like numbers $\zeta(2)$ sequence OEIS A081085

$$\sum_{k=0}^n \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k} = \sum_{k=0}^{\lfloor n/2 \rfloor} 4^{n-2k} \binom{n}{2k} \binom{2k}{k}^2$$

with recurrence relation

$$(n+1)^2 b(n+1) - (12n(n+1) + 4)b(n) + 32n^2 b(n-1) = 0$$

3.2.2 Apéry-like numbers $g = 2$ and $r = 1$: $b_{p/q}(k) = 2 \sin(\pi k p/q)$

Let us still focus on (21) but now for $g = 2$ and $r = 1$, with n necessarily even¹⁰. We find, for $n = 2, 4, 6, \dots$

$$\begin{aligned} p/q = 1 &\Rightarrow b(n) = (-1)^{n/2} \binom{n}{n/2}^2 \\ p/q = 1/2 &\Rightarrow b(n) = 4, 20, 112, 676, 4304, 28496, \dots \end{aligned}$$

These are the same Apéry-like numbers as above

$$\sum_{k=0}^{n/2} \binom{n/2}{k} \binom{2k}{k} \binom{n-2k}{n/2-k}$$

now occurring for even n 's. Indeed, cases $r = 2$ and ($r = 1$, n even) are essentially equivalent: calling $n = 2n'$ for $r = 1$, then $(2 \sin(\pi k p/q))^{n=l_1+l_2+\dots+l_j}$ with l_1, l_2, \dots, l_j a composition of n , is in fact $((2 \sin(\pi k p/q))^2)^{l'_1+l'_2+\dots+l'_j=n'}$ with l'_1, l'_2, \dots, l'_j a composition of n' , which is the $r = 2$ result.

⁹Or equivalently, using (14)

$$b(n) = gn \sum_{\substack{l_1, l_2, \dots, l_j \\ g\text{-composition of } n}} c_g(l_1, l_2, \dots, l_j) \int_0^1 dt \prod_{i=1}^j \left(2 \sin(\pi t + \pi(i-1)p/q) \right)^{rl_i}$$

¹⁰ n is necessarily even because $l_1 + l_2 + \dots + l_j$ (which is equal to n) has to be even.

3.2.3 Apéry-like numbers $g = 2$ and $r = 4$: $b_{p/q}(k) = (2 \sin(\pi k p/q))^4$

Let us again focus on $b(n)$ in (21) but now for $g = 2$ and $r = 4$: we find, for $n = 1, 2, 3, \dots$

$$\begin{aligned} p/q = 1 &\Rightarrow b(n) = \binom{2n}{n} \binom{4n}{2n} \\ p/q = 1/2 &\Rightarrow b(n) = 12, 164, 2352, 34596, 516912, 7806224, \dots \end{aligned}$$

These are the Apéry-like numbers $\zeta(2)$ sequence OEIS A143583

$$\sum_{k=0}^n \binom{2k}{k} \binom{4k}{2k} \binom{2n-2k}{n-k} \binom{4n-4k}{2n-2k} / \binom{2n}{n} = \sum_{k=0}^n 4^{n-k} \binom{2n-2k}{n-k} \binom{2k}{k}^2$$

with recurrence relation

$$(n+1)^2 b(n+1) - (32n(n+1) + 12)b(n) + 256n^2 b(n-1) = 0$$

3.2.4 Apéry-like numbers $g = 3$ and $r = 2$: $b_{p/q}(k) = (2 \sin(\pi k p/q))(2 \sin(\pi(k+1)p/q))$

Finally we focus ¹¹ on $b(n)$ in (27). We find, for $g = 3$ and $n = 1, 2, 3, \dots$

$$\begin{aligned} p/q = 1 &\Rightarrow b(n) = (-1)^n \binom{3n}{n} \binom{2n}{n} \Leftrightarrow \text{triangular lattice chiral walks counting} \\ p/q = 1/2 &\Rightarrow b(n) = \binom{3n/2}{n/2} \binom{n}{n/2} \quad \text{if } n \text{ multiple of 2 and 0 otherwise} \\ p/q = 1/3 &\Rightarrow b(n) = 3, 9, 21, 9, -297, -2421, \dots \end{aligned}$$

These are Apéry-like numbers $\zeta(2)$ sequence OEIS A006077

$$\sum_{k=0}^{\lfloor n/3 \rfloor} (-1)^k 3^{n-3k} \binom{n}{3k} \binom{2k}{k} \binom{3k}{k} = \sum_{k=0}^{\lfloor n/3 \rfloor} (-1)^k 3^{n-3k} \binom{n}{n-3k, k, k, k}$$

with recurrence relation

$$(n+1)^2 b(n+1) + (9n(n+1) + 3)b(n) + 27n^2 b(n-1) = 0$$

¹¹Or equivalently, using (18), on

$$\begin{aligned} b(n) = gn &\sum_{\substack{l_1, l_2, \dots, l_j \\ \text{g-composition of } n}} c_g(l_1, l_2, \dots, l_j) \\ &\int_0^1 dt \left(2 \sin(\pi t) \right)^{l_1} \prod_{i=2}^j \left(2 \sin(\pi t + \pi(i-1)p/q) \right)^{l_{i-1} + l_i} \left(2 \sin(\pi t + \pi j p/q) \right)^{l_j} \end{aligned}$$

4 Conclusions

The trigonometric identities analyzed in this work, as well as their generalizations to other spectral functions that can be derived along the lines presented here, allow us to obtain expressions for the algebraic area counting of a broad set of random walks on two-dimensional lattices. The only requirement is that these walks be described by a Hamiltonian of the general form introduced in [3], admitting an interpretation as systems of generalized exclusion statistics with specific spectral functions. A wide class of lattice walk models can be embedded into this framework, and we gave a few examples in the present work, most notably the triangular chiral walk introduced originally in [3].

The most obvious and interesting extension of our results would be in obtaining the area counting of other, more general types of walks. From the algebraic point of view, an immediate choice presents itself: the Hamiltonian

$$H_m = (u + u^{-1})^m v + v^{-1}(u + u^{-1})^m, \quad m = 1, 2, \dots$$

describes a class of Hofstadter-like models representing generalized random walks on the square lattice, with $m = 1$ the standard (Hofstadter) random walk and $m = 2$ the walk studied in subsection **3.1.2**. The model for general m represents a walk that proceeds in groups of one random step up or down and then m independent random steps left or right, but other representations are possible by performing modular transformations to the lattice (or redefinitions of the u, v operators in the Hamiltonian). All these walks belong to the class of $g = 2$ exclusion statistics and their area counting is readily given by the relevant $g = 2$ cluster coefficients and generalized trigonometric sums.

Clearly this is just the tip of a large iceberg as far as lattice walk models are concerned. For instance, another class of walks at $g = 2$ would be described by the Hamiltonian

$$\tilde{H}_m = (u^m + u^{m-1} + \dots + u^{-m})v + v^{-1}(u^m + u^{m-1} + \dots + u^{-m})$$

This represents walks proceeding with a random step up or down to one of the $2m + 1$ neighboring points in the left-right direction of distance up to m from the original horizontal position with equal probability. Again, the combinatorics of these walks are readily obtained with our methods. Yet other walks can be constructed, with asymmetrical propagation rules and belonging to higher g statistics. The only limitation, or criterion, is the potential relevance and physical significance of these walks, and this remains an open field of investigation.

The emergence of Apéry-like numbers within the mathematical structure of these walks is another intriguing but obscure issue. At the present level of our understanding this is something of a mystery, or curiosity. It would be satisfying to have a better understanding of the relation between random walks and Apéry numbers, with an eye to possible applications in the mathematics of ζ -functions and/or statistical models.

Finally, the Hamiltonians H_m and \tilde{H}_m presented above are all Hermitian and thus have a real spectrum, generalizing the corresponding spectrum of the Hofstadter model

that leads to the celebrated “butterfly” fractal structure. It is expected that the spectrum of all the above models will have a similarly fractal structure. The shape and eigenvalue statistics of the spectrum of these generalized models is an intriguing topic for further research.

Acknowledgments

S.O. acknowledges interesting discussions with Olivier Giraud, in particular regarding (16) and (17). He also thanks Stephan Wagner for mentioning the relation of the triangular lattice chiral walks of subsection (3.1.4) to Kreweras walks. A.P. acknowledges the hospitality of LPTMS, CNRS at Université Paris-Saclay (Faculté des Sciences d’Orsay), where this work was initiated. A.P.’s research was partially supported by NSF under grant 1519449 and by an “Aide Investissements d’Avenir” LabEx PALM grant (ANR-10-LABX-0039-PALM).

5 Appendix

5.1 Triangular $r = 4$: $b_{p/q}(k) = (2 \sin(\pi k p/q)) (2 \sin(\pi(k+1)p/q)) (2 \sin(\pi(k+2)p/q)) (2 \sin(\pi(k+3)p/q))$

Likewise

$$\begin{aligned} \frac{1}{q} \sum_{k=1}^q b_{p/q}^{l_1}(k) b_{p/q}^{l_2}(k+1) \dots b_{p/q}^{l_j}(k+j-1) &= \sum_{\substack{A=-\infty \\ A \text{ even}}}^{\infty} e^{i\pi A p/q} \\ &\sum_{k_3=-(l_1+l_2+l_3)/2}^{(l_1+l_2+l_3)/2} \sum_{k_4=-(l_1+l_2+l_3+l_4)/2}^{(l_1+l_2+l_3+l_4)/2} \dots \sum_{k_j=-(l_{j-3}+l_{j-2}+l_{j-1}+l_j)/2}^{(l_{j-3}+l_{j-2}+l_{j-1}+l_j)/2} \sum_{k_{j+1}=-(l_{j-2}+l_{j-1}+l_j)/2}^{(l_{j-2}+l_{j-1}+l_j)/2} \sum_{k_{j+2}=-(l_{j-1}+l_j)/2}^{(l_{j-1}+l_j)/2} \sum_{k_{j+3}=-l_j/2}^{l_j/2} \\ &\binom{l_1}{l_1/2 + A/2 + \sum_{i=3}^{j+3} (i-2)k_i} \binom{l_1+l_2}{(l_1+l_2)/2 - A/2 - \sum_{i=3}^{j+3} (i-1)k_i} \binom{l_1+l_2+l_3}{(l_1+l_2+l_3)/2 + k_3} \\ &\prod_{i=4}^j \binom{l_{i-3}+l_{i-2}+l_{i-1}+l_i}{(l_{i-3}+l_{i-2}+l_{i-1}+l_i)/2 + k_i} \binom{l_{j-2}+l_{j-1}+l_j}{(l_{j-2}+l_{j-1}+l_j)/2 + k_{j+1}} \binom{l_{j-1}+l_j}{(l_{j-1}+l_j)/2 + k_{j+2}} \binom{l_j}{l_j/2 + k_{j+3}} \end{aligned} \quad (29)$$

with overall counting

$$\binom{4(l_1+l_2+\dots+l_j)}{2(l_1+l_2+\dots+l_j)}$$

One notes that as in previous cases the binomial multiple sum (29) is nothing but the trigonometric integral

$$\begin{aligned} & \frac{1}{q} \sum_{k=1}^q b_{p/q}^{l_1}(k) b_{p/q}^{l_2}(k+1) \dots b_{p/q}^{l_j}(k+j-1) = \\ & \int_0^1 dt \left(2 \sin(\pi t) \right)^{l_1} \left(2 \sin(\pi t + \pi p/q) \right)^{l_1+l_2} \left(2 \sin(\pi t + \pi 2p/q) \right)^{l_1+l_2+l_3} \\ & \prod_{i=4}^j \left(2 \sin(\pi t + \pi(i-1)p/q) \right)^{l_{i-3}+l_{i-2}+l_{i-1}+l_i} \left(2 \sin(\pi t + \pi j p/q) \right)^{l_{j-2}+l_{j-1}+l_j} \\ & \left(2 \sin(\pi t + \pi(j+1)p/q) \right)^{l_{j-1}+l_j} \left(2 \sin(\pi t + \pi(j+2)p/q) \right)^{l_j} \end{aligned}$$

under the provision that $2(l_1 + \dots + l_j) < q$.

Clearly for a general r the spectral function $b_{p/q}(k) = (2 \sin(\pi k p/q)) (2 \sin(\pi(k+1)p/q)) \dots (2 \sin(\pi(k+r-1)p/q))$ can be treated along the same lines as in subsections (2.2.1) and (2.2.2) and above.

5.2 Another triangular chiral walks generalization:

$$b_{p/q}(k) = (2 \sin(\pi k p/q))^{r/2} (2 \sin(\pi(k+1)p/q))^{r/2} \quad \text{with } r \text{ even}$$

When $b_{p/q}(k) = (2 \sin(\pi k p/q))^2$ we have seen that (3), rewritten as

$$\begin{aligned} & \frac{1}{q} \sum_{k=1}^q b_{p/q}^{l_1}(k) b_{p/q}^{l_2}(k+1) \dots b_{p/q}^{l_j}(k+j-1) = \sum_{\substack{A=-\infty \\ A \text{ even}}}^{\infty} e^{i\pi A p/q} \\ & \sum_{k_3=-l_3}^{l_3} \dots \sum_{k_j=-l_j}^{l_j} \binom{2l_1}{l_1 + A/2 + \sum_{i=3}^j (i-2)k_i} \binom{2l_2}{l_2 - A/2 - \sum_{i=3}^j (i-1)k_i} \prod_{i=3}^j \binom{2l_i}{l_i + k_i} \end{aligned}$$

generalizes for $b_{p/q}(k) = (2 \sin(\pi k p/q))^r$ and r is even to

$$\begin{aligned} & \frac{1}{q} \sum_{k=1}^q b_{p/q}^{l_1}(k) b_{p/q}^{l_2}(k+1) \dots b_{p/q}^{l_j}(k+j-1) = \sum_{\substack{A=-\infty \\ A \text{ even}}}^{\infty} e^{i\pi A p/q} \\ & \sum_{k_3=-rl_3/2}^{rl_3/2} \dots \sum_{k_j=-rl_j/2}^{rl_j/2} \binom{rl_1}{rl_1/2 + A/2 + \sum_{i=3}^j (i-2)k_i} \binom{rl_2}{rl_2/2 - A/2 - \sum_{i=3}^j (i-1)k_i} \prod_{i=3}^j \binom{rl_i}{rl_i/2 + k_i} \end{aligned}$$

Likewise, when $b_{p/q}(k) = (2 \sin(\pi kp/q))(2 \sin(\pi(k+1)p/q))$, equation (19),

$$\begin{aligned}
& \frac{1}{q} \sum_{k=1}^q b_{p/q}^{l_1}(k) b_{p/q}^{l_2}(k+1) \dots b_{p/q}^{l_j}(k+j-1) \\
&= \sum_{\substack{A=-\infty \\ \text{A same parity } l_1+l_2+\dots+l_j}}^{\infty} e^{i\pi Ap/q} \sum_{k_3=-(l_2+l_3)/2}^{(l_2+l_3)/2} \dots \sum_{k_j=-(l_{j-1}+l_j)/2}^{(l_{j-1}+l_j)/2} \sum_{k_{j+1}=-l_j/2}^{l_j/2} \\
& \quad \binom{l_1}{l_1/2 + A/2 + \sum_{i=3}^{j+1} (i-2)k_i} \binom{l_1+l_2}{(l_1+l_2)/2 - A/2 - \sum_{i=3}^{j+1} (i-1)k_i} \\
& \quad \prod_{i=3}^j \binom{l_{i-1}+l_i}{(l_{i-1}+l_i)/2 + k_i} \binom{l_j}{l_j/2 + k_{j+1}}
\end{aligned}$$

generalizes for $b_{p/q}(k) = ((2 \sin(\pi kp/q))^{r/2} (2 \sin(\pi(k+1)p/q))^{r/2})$ and r even to

$$\begin{aligned}
& \frac{1}{q} \sum_{k=1}^q b_{p/q}^{l_1}(k) b_{p/q}^{l_2}(k+1) \dots b_{p/q}^{l_j}(k+j-1) \\
&= \sum_{\substack{A=-\infty \\ \text{A same parity } r(l_1+l_2+\dots+l_j)/2}}^{\infty} e^{i\pi Ap/q} \sum_{k_3=-r(l_2+l_3)/4}^{r(l_2+l_3)/4} \dots \sum_{k_j=-r(l_{j-1}+l_j)/4}^{r(l_{j-1}+l_j)/4} \sum_{k_{j+1}=-rl_j/4}^{rl_j/4} \\
& \quad \binom{rl_1/2}{rl_1/4 + A/2 + \sum_{i=3}^{j+1} (i-2)k_i} \binom{r(l_1+l_2)/2}{r(l_1+l_2)/4 - A/2 - \sum_{i=3}^{j+1} (i-1)k_i} \\
& \quad \prod_{i=3}^j \binom{r(l_{i-1}+l_i)/2}{r(l_{i-1}+l_i)/4 + k_i} \binom{rl_j/2}{rl_j/4 + k_{j+1}}
\end{aligned}$$

5.3 Regarding (16): summing over A odd when r is even

So far one has considered the $r(l_1 + l_2 + \dots + l_j)$ even cases so that the $q \rightarrow \infty$ limit in the trigonometric sum (1) yields an overall binomial counting which is an integer and contributes as such to the overall counting of closed lattice walks. We have seen that this trigonometric sum can be rewritten as a multiple binomial sum of the type (16) or (19) with some constraints on the evenness or oddness of the A 's (and additionally of $l_1 + l_2 + \dots + l_j$ in the case r odd). In the $r(l_1 + l_2 + \dots + l_j)$ odd cases, on the other hand, (1) would not rewrite anymore as a multiple binomial sum.

Still, and quite generally, one could take the binomial multiple sums (16) (and likewise (19)) at face value for all possible entries A even or odd and $l_1 + l_2 + \dots + l_j$ even or odd.

In the r even case we already know that the A even summation in (16) has a finite range and yields exactly the overall integer counting binomial. The A odd summation happens to yield again the same overall binomial but with each term in the sum a rational number times $1/\pi^2$ and an infinite summation range. The $1/\pi^2$ factor comes from the first two binomials in (16) due the relaxation of the constraint that their entries be integers (since A is now odd). Likewise in the r odd case, when $l_1 + l_2 + \dots + l_j$ is even, we already know that A even or odd summations, depending on the parity of $l_1 + l_3 + \dots$, have a finite range and yield the usual overall integer counting binomial; it is still true that summing over A even with $l_1 + l_3 + \dots$ odd or on A odd with $l_1 + l_3 + \dots$ even would yield the same overall counting binomial with again terms $1/\pi^2$ times rational numbers and an infinite summation range. Finally when both r and $l_1 + l_2 + \dots + l_j$ are odd, A even and odd summations have finite range to yield the overall binomial which is in this case $1/\pi$ times a rational number. In all these instances the coefficients sum up to $\binom{r(l_1+l_2+\dots+l_j)}{r(l_1+l_2+\dots+l_j)/2}$ for both A even or odd summations, with finite or infinite ranges depending on the situation.

To better understand these weird A -summations, let us first focus on the regular A -summations and consider the LHS of (17) i.e., the binomial multiple sum

$$\sum_{k_3=-r l_3/2}^{r l_3/2} \cdots \sum_{k_j=-r l_j/2}^{r l_j/2} \binom{r l_1}{r l_1/2 + A/2 + \sum_{i=3}^j (i-2)k_i} \binom{r l_2}{r l_2/2 - A/2 - \sum_{i=3}^j (i-1)k_i} \prod_{i=3}^j \binom{r l_i}{r l_i/2 + k_i}$$

One wishes to go backward and get the double integral in the RHS of (17), which, when summed over A , directly yield the overall counting binomial

$$\binom{r(l_1 + l_2 + \dots + l_j)}{r(l_1 + l_2 + \dots + l_j)/2}$$

For simplicity let us consider the case r even: since r is even, all the k_i 's $i = 3, \dots, j$ are integers, and since we know that A has then to be even (see below (16)), in the first two binomials both $r l_1/2 + A/2 + \sum_{i=3}^j (i-2)k_i$ and $r l_2/2 - A/2 - \sum_{i=3}^j (i-1)k_i$ are integers. Using that for an integer n

$$\int_0^1 dt e^{2i\pi(k-n)t}$$

is the Kronecker $\delta(k, n)$ meaning

$$\sum_{\substack{k=-\infty \\ k \text{ integer}}}^{\infty} \delta(k, n) f(k) = f(n)$$

we can rewrite these binomials as

$$\binom{r l_1}{r l_1/2 + A/2 + \sum_{i=3}^j (i-2)k_i} = \sum_{\substack{k_1=-r l_1/2 \\ k_1 \text{ integer}}}^{r l_1/2} \int_0^1 dt e^{2i\pi(k_1 - (A/2 + \sum_{i=3}^j (i-2)k_i))t} \binom{r l_1}{r l_1/2 + k_1}$$

$$\binom{rl_2}{rl_2/2 - A/2 - \sum_{i=3}^j (i-1)k_i} = \sum_{\substack{k_2=-rl_2/2 \\ k_2 \text{ integer}}}^{rl_2/2} \int_0^1 dt' e^{2i\pi(k_2 + A/2 + \sum_{i=3}^j (i-1)k_i)t'} \binom{rl_2}{rl_2/2 + k_2}$$

where the summations are restricted to $[-rl_1/2, rl_1/2]$ and $[-rl_2/2, rl_2/2]$ since there is no point to sum outside these intervals where the binomials trivially vanish. So the LHS of (17) becomes

$$\sum_{\substack{k_1=-rl_1/2 \\ k_1 \text{ integer}}}^{rl_1/2} \cdots \sum_{\substack{k_j=-rl_j/2 \\ k_j \text{ integer}}}^{rl_j/2} \int_0^1 dt e^{2i\pi(k_1 - (A/2 + \sum_{i=3}^j (i-2)k_i))t} \int_0^1 dt' e^{2i\pi(k_2 + A/2 + \sum_{i=3}^j (i-1)k_i)t'} \prod_{i=1}^j \binom{rl_i}{rl_i/2 + k_i}$$

which is

$$\sum_{\substack{k_1=-rl_1/2 \\ k_1 \text{ integer}}}^{rl_1/2} \cdots \sum_{\substack{k_j=-rl_j/2 \\ k_j \text{ integer}}}^{rl_j/2} \int_0^1 dt \int_0^1 dt' e^{i\pi A(t'-t)} \prod_{i=1}^j \binom{rl_i}{rl_i/2 + k_i} e^{2i\pi k_i((i-1)t' - (i-2)t)}$$

i.e., since obviously

$$\sum_{\substack{k_i=-rl_i/2 \\ k_i \text{ integer}}}^{rl_i/2} \binom{rl_i}{rl_i/2 + k_i} e^{2i\pi k_i((i-1)t' - (i-2)t)} = \left(2 \cos(\pi((i-1)t' - (i-2)t)) \right)^{rl_i}$$

and calling $t' - t = t''$, we obtain¹²

$$\int_0^1 dt \int_0^1 dt'' e^{i\pi A t''} \prod_{i=1}^j \left(2 \cos(\pi((i-1)t'' + t)) \right)^{rl_i}$$

We have to sum over A even: since $\sum_{A \text{ even}} e^{i\pi A t''} = \sum_{n=-\infty}^{\infty} \delta(t'', n)$

$$\begin{aligned} \sum_{A \text{ even}} \int_0^1 dt \int_0^1 dt'' e^{i\pi A t''} \prod_{i=1}^j \left(2 \cos(\pi((i-1)t'' + t)) \right)^{rl_i} &= \int_0^1 dt (2 \cos(\pi t))^{r(l_1 + l_2 + \dots + l_j)} \\ &= \binom{r(l_1 + l_2 + \dots + l_j)}{r(l_1 + l_2 + \dots + l_j)/2} \end{aligned} \quad (30)$$

where the overall binomial counting has been obtained as expected.

¹²Or equivalently as in the RHS of (17)

$$\frac{1}{2} \int_0^1 dt \int_0^2 dt'' e^{i\pi A t''} \prod_{i=1}^j \left(2 \sin(\pi((i-1)t'' + t)) \right)^{rl_i}$$

Now still assuming r being even, so that all the k_i 's $i = 3, \dots, j$ are integers, let us insist that the summation over A be on A odd so that both $rl_1/2 + A/2 + \sum_{i=3}^j (i-2)k_i$ and $rl_2/2 - A/2 - \sum_{i=3}^j (i-1)k_i$ are half-integers. Using that for an half-integer $n/2$

$$\int_0^1 dt e^{2i\pi(k-n/2)t}$$

is the Kronecker $\delta(k, n/2)$ meaning

$$\sum_{\substack{k=-\infty \\ k \text{ half integer}}}^{\infty} \delta(k, n/2) f(k) = f(n/2)$$

we rewrite the same two binomials as

$$\binom{rl_1}{rl_1/2 + A/2 + \sum_{i=3}^j (i-2)k_i} = \sum_{\substack{k_1=-\infty \\ k_1 \text{ half integer}}}^{\infty} \int_0^1 dt e^{2i\pi(k_1 - (A/2 + \sum_{i=3}^j (i-2)k_i))t} \binom{rl_1}{rl_1/2 + k_1}$$

$$\binom{rl_2}{rl_2/2 - A/2 - \sum_{i=3}^j (i-1)k_i} = \sum_{\substack{k_2=-\infty \\ k_2 \text{ half integer}}}^{\infty} \int_0^1 dt' e^{2i\pi(k_2 + A/2 + \sum_{i=3}^j (i-1)k_i)t'} \binom{rl_2}{rl_2/2 + k_2}$$

Doing the same manipulations as above except for the first two binomials the LHS of (17) then becomes

$$\int_0^1 dt \int_0^1 dt'' \sum_{\substack{k_1=-\infty \\ k_1 \text{ half integer}}}^{\infty} \sum_{\substack{k_2=-\infty \\ k_2 \text{ half integer}}}^{\infty} \binom{rl_1}{rl_1/2 + k_1} \binom{rl_2}{rl_2/2 + k_2} e^{i\pi A t''} e^{2i\pi(k_1+k_2)t} e^{2i\pi k_2 t''} \prod_{i=3}^j \left(2 \cos(\pi((i-1)t'' + t)) \right)^{r l_i}$$

Summing over all A odd i.e., over $A + 2k_2$ even –since k_2 is an half integer– yields again a Kronecker enforcing $t'' = 0$ so that after summation one obtains

$$\int_0^1 dt \sum_{\substack{k_1=-\infty \\ k_1 \text{ half integer}}}^{\infty} \sum_{\substack{k_2=-\infty \\ k_2 \text{ half integer}}}^{\infty} \binom{rl_1}{rl_1/2 + k_1} \binom{rl_2}{rl_2/2 + k_2} e^{2i\pi(k_1+k_2)t} (2 \cos(\pi t))^{r(l_3+\dots+l_j)}$$

Comparing with (30) we see that in order to get the same overall binomial counting everything boils down to showing that in the same way that obviously

$$\sum_{\substack{k_1=-r l_1/2 \\ k_1 \text{ integer}}}^{r l_1/2} \sum_{\substack{k_2=-r l_2/2 \\ k_2 \text{ integer}}}^{r l_2/2} \binom{rl_1}{rl_1/2 + k_1} \binom{rl_2}{rl_2/2 + k_2} e^{2i\pi(k_1+k_2)t} = (2 \cos(\pi t))^{r(l_1+l_2)} \quad (31)$$

holds,

$$\begin{aligned} & \sum_{\substack{k_1=-\infty \\ k_1 \text{ half integer}}}^{\infty} \sum_{\substack{k_2=-\infty \\ k_2 \text{ half integer}}}^{\infty} \binom{rl_1}{rl_1/2 + k_1} \binom{rl_2}{rl_2/2 + k_2} e^{2i\pi(k_1+k_2)t} \\ &= (2 \cos(\pi t))^{r(l_1+l_2)} \end{aligned}$$

should also hold.

To show this let us focus on the trivial identity (31) which is nothing but

$$\sum_{\substack{k_1=-r_1/2 \\ k_1 \text{ integer}}}^{r_1/2} \sum_{\substack{k_2=-r_2/2 \\ k_2 \text{ integer}}}^{r_2/2} \binom{rl_1}{rl_1/2 + k_1} \binom{rl_2}{rl_2/2 + k_2} e^{2i\pi(k_1+k_2)t} = \sum_{\substack{k=-r(l_1+l_2)/2 \\ k \text{ integer}}}^{r(l_1+l_2)/2} \binom{r(l_1+l_2)}{r(l_1+l_2)/2 + k} e^{2i\pi kt}$$

or equivalently, harmlessly relaxing the range of k_1, k_2 and k summations,

$$\sum_{\substack{k_1=-\infty \\ k_1 \text{ integer}}}^{\infty} \sum_{\substack{k_2=-\infty \\ k_2 \text{ integer}}}^{\infty} \binom{rl_1}{rl_1/2 + k_1} \binom{rl_2}{rl_2/2 + k_2} e^{2i\pi(k_1+k_2)t} = \sum_{\substack{k=-\infty \\ k \text{ integer}}}^{\infty} \binom{r(l_1+l_2)}{r(l_1+l_2)/2 + k} e^{2i\pi kt} \quad (32)$$

Let us to rederive it in an other way : defining $k = k_1 + k_2$ we can rewrite

$$\begin{aligned} & \sum_{\substack{k_1=-\infty \\ k_1 \text{ integer}}}^{\infty} \sum_{\substack{k_2=-\infty \\ k_2 \text{ integer}}}^{\infty} \binom{rl_1}{rl_1/2 + k_1} \binom{rl_2}{rl_2/2 + k_2} e^{2i\pi(k_1+k_2)t} \\ &= \sum_{\substack{k_+=-\infty \\ k \text{ integer}}}^{\infty} \sum_{\substack{k_1=-\infty \\ k_1 \text{ integer}}}^{\infty} \binom{rl_1}{rl_1/2 + k_1} \binom{rl_2}{rl_2/2 + k - k_1} e^{2i\pi kt} \end{aligned}$$

Thanks to the Chu-Vandermonde identity

$$\binom{l_1 + l_2}{l'_1 + l'_2} = \sum_{\substack{k_1=-\max(l'_1, l'_2) \\ k_1 \text{ integer}}}^{\max(l'_1, l'_2)} \binom{l_1}{l'_1 + k_1} \binom{l_2}{l'_2 - k_1}$$

we conclude that we indeed recover (32).

It is clear that the same conclusion can be reached when k_1 and k_2 are now both half integers namely

$$\begin{aligned} & \sum_{\substack{k_1=-\infty \\ k_1 \text{ half integer}}}^{\infty} \sum_{\substack{k_2=-\infty \\ k_2 \text{ half integer}}}^{\infty} \binom{rl_1}{rl_1/2 + k_1} \binom{rl_2}{rl_2/2 + k_2} e^{2i\pi(k_1+k_2)t} \\ &= \sum_{\substack{k=-\infty \\ k \text{ integer}}}^{\infty} \binom{r(l_1+l_2)}{r(l_1+l_2)/2 + k} e^{2i\pi kt} \quad (33) \end{aligned}$$

Indeed k_1 and k_2 being both half integers then $k = k_1 + k_2$ is again an integer so we can write

$$\begin{aligned} & \sum_{\substack{k_1=-\infty \\ k_1 \text{ half integer}}}^{\infty} \sum_{\substack{k_2=-\infty \\ k_2 \text{ half integer}}}^{\infty} \binom{rl_1}{rl_1/2 + k_1} \binom{rl_2}{rl_2/2 + k_2} e^{2i\pi(k_1+k_2)t} \\ &= \sum_{\substack{k=-\infty \\ k \text{ integer}}}^{\infty} \sum_{\substack{k_1=-\infty \\ k_1 \text{ half integer}}}^{\infty} \binom{rl_1}{rl_1/2 + k_1} \binom{rl_2}{rl_2/2 + k - k_1} e^{2i\pi kt} \end{aligned}$$

Thanks to the generalized Chu-Vandermonde identity

$$\binom{l_1 + l_2}{l'_1 + l'_2} = \sum_{\substack{k_1=-\infty \\ k_1 \text{ half integer}}}^{\infty} \binom{l_1}{l'_1 + k_1} \binom{l_2}{l'_2 - k_1}$$

we reach indeed the identity (33) for the half integers summations. From which it directly follows that in the presence of the additional $(2 \cos(\pi t))^{r(l_3+\dots+l_j)}$ term integrating over t from 0 to 1 one ends up getting again the same overall binomial counting, as desired.

References

- [1] S. Ouvry and S. Wu, “The algebraic area of closed lattice random walks”, *Journal of Physics A: Mathematical and Theoretical*, Volume 52, Number 25 (2019).
- [2] D.R. Hofstadter, “Energy levels and wave functions of Bloch electrons in rational and irrational magnetic fields”, *Phys. Rev. B* **14** (1976) 2239.
- [3] S. Ouvry and A. Polychronakos, “Exclusion statistics and lattice random walks”, *NPB[FS]* 948 (2019)114731.
- [4] F.D.M. Haldane, “Fractional statistics in arbitrary dimensions: A generalization of the Pauli principle”, *Phys. Rev. Lett.* **67** (1991) 937940; see also Y.S. Wu, “Statistical distribution for generalized ideal gas of fractional-statistics particles”, *Phys. Rev. Lett.* **73** (1994) 922925; A.P. Polychronakos, “Nonrelativistic bosonization and fractional statistics”, *Nucl. Phys.* **B324** (1989) 597; A. Dasnières de Veigy and S. Ouvry, “Equation of State of an Anyon gas in a Strong Magnetic Field”, *Phys. Rev. Lett.* **72** (1994) 600.
- [5] see e.g., O. Bernardi, “Bijective counting of Kreweras walks and loopless triangulations” *Journal of Combinatorial Theory - Series A*, Vol 114(5) (2007) 931-956.




Figure 1: The lattice in (3.1.1).




Figure 2: U , V and W are the three possible hoppings on the triangular lattice. As an illustration two chiral walks going around *up-vertex* and *down-vertex* triangular cells starting from the black bullet lattice site.






Figure 3: Three of the 6 possible chiral walks starting from the same black bullet lattice site. Only the 3 outgoing arrows represent possible motions from the original site.



$UWUWV^2$



$U^2W^2V^2$

Figure 4: $UWUWV^2$ and $U^2W^2V^2$ walks.