

# SPECTRAL ORDERING AND 2-SWITCH TRANSFORMATIONS

ELISMAR OLIVEIRA, VICTOR N. SCHVÖLLNER, AND VILMAR TREVISAN

ABSTRACT. We address the problem of ordering trees with the same degree sequence by their spectral radii. To achieve that, we consider 2-switch transformations which preserve the degree sequence and establish when the index decreases. Our main contribution is to determine a total ordering of a particular family by their indices according to a given parameter related to sizes in the tree.

## 1. INTRODUCTION

Giving a graphical degree sequence, a general, natural, and well studied problem is to determine, with respect to a given parameter, the extremal members in the family of graphs satisfying this degree sequence. A 2-switch transformation (see definition below) is a convenient way to study this problem since it is a well known fact that for two graphs with the same degree sequence, one can be obtained from the other by applying successive 2-switches.

The main purpose of this note is to address the problem of finding extremal members in families of graphs having the same degree sequence, with respect to the *spectral radius* (or *index*), which is the largest eigenvalue of the adjacency matrix. We remark that this problem has been studied in great generality in the celebre paper by T. Biyikoğlu and J. Leydold [5], where the authors show that, in the maximum element, the degree sequence is non-increasing with respect to an ordering of the vertices induced by breadth-first search that is consistent with the eigenvector associated with the index. In [1], the authors determined the tree having maximum spectral radius among all caterpillar with a fixed degree sequence. We observe that in both papers 2-switch transformations is used to analyse the variation of the index.

In order to explain our results, we need a few definitions. For a graph  $G = (V, E)$  having four distinct vertices  $a, b, c, d \in V$  such that  $ab, cd \in E$  and  $ac, bd \notin E$ , the removal of the edges  $ab$  and  $cd$  from  $G$  and the addition of  $ac$  and  $bd$  to  $G$  is referred to as a *2-switch* in  $G$ . This is a well studied classical operation (see, for example [4, 8]). It is straightforward to check that 2-switch operations preserve the degree sequence.

As a way to illustrate how this class of problems may be approached, we study how the spectral radius varies upon 2-switch transformations in the family  $\mathfrak{F}(n)$  of trees given in Figure 1. The technique we use is a powerful algorithmic tool that allows one to compare the indices of two trees without computing them. Our main result is a total ordering in this family and, as a consequence, we obtain the extremal members.

---

1991 *Mathematics Subject Classification.* 05C50, 05C05, 15A18.

*Key words and phrases.* spectral radius; tree; 2-switch; ordering;

We believe that this result is remarkable, since it is quite unusual to obtain a total order by any graph parameter. Spectral parameters have been used to classify many families, however it is rare that a total order is obtained. As examples, we refer to the papers [2, 3, 6, 10, 13, 14, 15], where the ordering of graphs by the index is studied.




FIGURE 1. The family of trees  $\mathfrak{F}(n)$ .

The remaining of the paper is as follows. In order to explain that the family we study is not arbitrary, we devote the rest of this introduction to justify our choice. In Section 2, we define the family  $\mathfrak{F}(n)$  and the 2-switch transformations we perform. Moreover, we explain our powerful technique to obtain the order, that is based on an algorithmic tool, allowing to compare indices of trees without computing them. In Section 3 we obtain necessary analytical properties of some recurrence relations that appear in our comparison method. In Section 4 we show how the spectral radius varies upon 2-switches transformations. In Section 5, we use the 2-switches to obtain a total ordering in  $\mathfrak{F}(n)$ . Finally, in Section 6 we reason how 2-switches may be useful for the class of problems proposed here.

**1.1. Motivation for choosing the family.** We start by introducing some notation from [12] which is specially useful to represent the trees in the family  $\mathfrak{F}(n)$ . In that paper, it was proven that the number of Laplacian eigenvalues less than the average degree  $2 - \frac{2}{n}$  of a tree having  $n$  vertices is at least  $\lceil \frac{n}{2} \rceil$ . We remark that pendant paths of length 2 play an important role there and serve as a motivation for our choice.

Let  $T$  be a tree with  $n$  vertices, and let  $u$  be a vertex of degree at least  $\ell$  of  $T$  having  $\ell \geq 1$  pendant paths attached at  $u$ . We denote the *sum* of pendant paths attached at  $u$  by  $P(u) = P_{q_1} \oplus \dots \oplus P_{q_\ell}$ , as illustrated in Figure 2. The number of edges in each path is denoted by  $\#P_q = q$ . A subgraph obtained by a vertex  $u$  attached to  $r \geq 1$  paths of length 2, is called a *sun with  $r$  rays* and denoted by  $S_r$ . To simplify the representation, we use the concatenation symbol and write  $P_q * S_r$ . To further simplify the graphic part of the representation, we will use a black square  $\blacksquare$  to represent a pendant sun  $S_r$  attached to a vertex, and a single edge to represent the entire path  $P_q$ , omitting the  $r$  pendant  $P_2$ 's and the  $q$  vertices. We will refer to this as the  $(P_q, S_r)$  *representation* of this generalized pendant path  $P_q * S_r$ , as shown in Figure 2 (right). We can consider  $q = 0$  for paths  $P_q$  of length 0, as well as  $r = 0$  for no pendant  $S_r$ . However, we do *not* allow both  $r = q = 0$  simultaneously.

We say a vertex  $u$  is a *starlike vertex* if it has degree  $\geq 3$  and has at least two generalized




FIGURE 2. Vertex  $u$  with  $P(u) = P_1 \oplus 2P_2 \oplus P_4 \oplus P_5$  (left) and  $(P_q, S_r)$  representation of a generalized pendant path (right).

pendant paths attached to it. Using this notation, we can write any tree  $T$  having a single starlike vertex  $u$  as  $T = u + P_{q_1} * S_{r_1} \oplus \cdots \oplus P_{q_\ell} * S_{r_\ell}$  where  $\ell \geq 1$ .

In particular, any member of  $\mathfrak{F}(n)$ , as in the Figure 1, has a single starlike vertex  $u$  and is represented as

$$T = u + P_h * S_0 \oplus P_{q_1} * S_{r_1} \oplus P_{q_2} * S_{r_2},$$

for  $h, q_1, q_2 \geq 2$ .

Our choice of the family is related to the fact that we would like to study the spectral radius ordering of 2-switches on trees having a single starlike vertex. We notice that the general family of trees  $\mathfrak{H}$  with just one starlike vertex in this notation is given by,

$$\mathfrak{H} := \{T \mid T = u + P_{q_0} * S_{r_0} \oplus P_{q_1} * S_{r_1} \oplus P_{q_2} * S_{r_2}, q_0, q_1, q_2 \geq 0, r_0, r_1, r_2 \geq 0\}.$$

In this generality, our tool becomes too involved and, hence, in order to simplify the computations and to obtain symmetry, we have chosen to study a special case of this family  $\mathfrak{H}$ , where we replace  $S_{r_0}$  by  $S_0$  (no pendant  $P_2$ 's on the path  $P_h$ ).

## 2. A FAMILY AND OUR TOOL

In this section we define the family which we will determine a spectral radius ordering when performing 2-switch operation, as well as an algorithmic tool that we believe it is powerful for this class of problems.

**2.1. Our Family.** We consider the family  $\mathfrak{F}(n)$  of all trees  $T$  given by Figure 1, with the following constraints:

- 1)  $q_1, q_2 \geq 2$ ;
- 2)  $r_1, r_2 \geq 2$  and  $r_1 < r_2$ ;
- 3)  $h \geq 2$ .

Notice that, in this case, the number of vertices is  $n = |T| = 1 + h + q_1 + q_2 + 2(r_1 + r_2) \geq 7 + 2(r_1 + r_2) \geq 17$ .

We are interested in ordering by the index (spectral radius) the trees with a fixed degree sequence, in  $\mathfrak{F}(n)$ . As a particular case, we obtain the extreme members of the family, that is, the trees in  $\mathfrak{F}(n)$  having largest and minimum spectral radius. The degree sequence a  $T \in \mathfrak{F}(n)$  is given by

$$d := [r_1 + 1, r_2 + 1, 3, 2^{n-(r_1+r_2)-2}, 1^{r_1+r_2+1}].$$

In order to keep the degree sequence we fix  $r_2 > r_1 \geq 2$ . In this way each element in  $\mathfrak{F}(n)$ , with  $n$  vertices and degree sequence  $d$  is uniquely determined by the 3-uple  $[h, q_1, q_2]$  that is,

$$\mathfrak{F}(n) := \{T = [h, q_1, q_2] \mid h + q_1 + q_2 = n - 1 - 2(r_1 + r_2)\}.$$

Now we consider two types of 2-switches on  $T = [h, q_1, q_2] \in \mathfrak{F}(n)$ . Later, in Section 4, we determine how to order these operations by their spectral radii.

Type I - We switch the vertices between the central path and the right (or the left) branch of  $T$  obtaining a new tree  $T'$  (see Figure 3). More precisely, we have a new member of  $\mathfrak{F}(n)$ , where the parameters are changed by  $q'_1 = h - s + t$  (the closest to  $r_1 P_2$ ) and  $h' = q_1 - t + s$  (the new central path after some relabeling). Let us call that a  $(s, t)$ -2-switch of type II (see Figure 3). In order to do the 2-switch, we disconnect the edges  $[u_1, u_2]$  and  $[w_1, w_2]$  and reconnect  $[u_1, w_2]$  and  $[w_1, u_2]$ .




FIGURE 3. The 2-switch of type I.

Type II - We switch the vertices between the right and the left branches of  $T$  obtaining a new tree  $T'$ ; Notice that this transformation is indeed a 2-switch preserving the degree sequence ( $d := [r_1 + 1, r_2 + 1, 3, 2^{n-(r_1+r_2)-2}, 1^{r_1+r_2+1}]$ ). More precisely, we

have a new member of  $\mathfrak{F}(n)$ , where the parameters are changed by  $q'_1 = q_2 + t - s$  (the closest to  $r_1 P_2$ ) and  $q'_2 = q_1 - t + s$  (the closest to  $r_1 P_2$ ). Let us call that a  $(s, t)$ -2-switch of type II (see Figure 4). In order to do the 2-switch, we disconnect the edges  $[u_1, u_2]$  and  $[v_1, v_2]$  and reconnect  $[u_1, v_2]$  and  $[v_1, u_2]$ .





FIGURE 4. The 2-switch of type II.

We can always assume that  $\lambda = \rho(T) > \sqrt{r_2 + 2}$  for all  $T$  in  $\mathfrak{F}(n)$  because the sun  $S_{r_2+1}$ , whose spectral radius is  $\sqrt{r_2 + 2}$ , is a proper subgraph of  $T$ , since  $q_2 \geq 2$ . But in fact, we need a larger lower bound for  $\lambda = \rho(T)$  for our results.

Let  $T_j := \underbrace{[2, 2, \dots, 2, j]}_{r_2 \text{ times}}$  be the starlike tree composed by  $r_2$  legs of  $P_2$  and a path of length  $j$ , with  $j \geq 3$  as Figure 5 illustrates.

FIGURE 5. The graph  $T_j$ .

**Theorem 2.1.** *Let  $T \in \mathfrak{F}(n)$ . If  $j \geq 3$ , then*

$$\rho(T) > \rho(T_j).$$

This result may be proven by our comparison method that will be explained next, but there is a simpler proof, for which we need the following definitions and known result of Lemma 2.2.

An internal path in a graph  $G$ , denoted by  $v_1 v_2, \dots, v_{r-1} v_r$ , is a path beginning at  $v_1$  and ending at  $v_r$ , where  $v_1$  and  $v_r$  both have degree bigger than two, while all other vertices have degree two. The vertices  $v_1$  and  $v_r$  are not necessarily distinct. We denote by  $C_n$ , the cycle on  $n$  vertices and by  $W_n$  the tree with  $n$  vertices where two vertices have degree three and the distance between them is  $n - 5$ . The following result appears in the work by Hoffman and Smith [9].

**Lemma 2.2.** *Let  $G$  be a graph with  $n$  vertices,  $G \neq C_n, W_n$ . Let  $G'$  be the graph with  $n + 1$  vertices obtained from  $G$  by inserting a new vertex of degree two in an edge  $e$ . Then*

- (a) *if  $e$  lies on an internal path then  $\lambda(G') < \lambda(G)$ ;*
- (b) *if  $e$  does not lie on an internal path then  $\lambda(G') > \lambda(G)$ .*

*Proof.* (Theorem 2.1) Let  $T = [h, q_1, q_2] \in \mathfrak{F}(n)$  and  $3 \leq j$ .

Let  $H'_1$  be the tree obtained by adding an edge in the internal path starting from  $u$  to the vertex  $r_1 P_2$ . We notice that  $T \neq W_n$ , as it has a vertex  $u$  of degree  $r_2 + 1 > 3$ , and hence, by Lemma 2.2,  $\rho(H'_1) < \rho(T)$ . We now remove the pendant vertex of the path  $P_h$  in  $T$ , obtaining a tree  $H_1$ , a proper subtree of  $H'_1$ . It follows that  $\rho(H_1) < \rho(T)$ . We apply this process successively  $h$  times, obtaining a tree  $H_h$  composed by the starlikes  $r_2 P_2$  and  $r_1 P_2$  linked in their centers by a path of length  $q_2 + h + q_1 + 1$  such that  $\rho(H_h) < \rho(T)$ .

Now, if  $3 \leq j \leq q_2 + h + q_1 + 1$ , we see that  $T_j$  is a proper subtree of  $H_h$  and therefore,  $\rho(T_j) < \rho(H_h) < \rho(T)$ .

For  $j > q_2 + h + q_1 + 1$ , we keep adding edges in the internal path starting at  $r_2 P_2$  and ending at  $r_1 P_2$  until the length of the path is at least  $j$ , obtaining a tree  $H_j$ . This operation, according the Lemma 2.2, decreases the spectral radius. As  $T_j$  is a proper subtree of  $H_j$ , it follows that  $\rho(T_j) < \rho(H_j) < \rho(T)$ . □

2.2. **Our tool.** We would like to recall the algorithm **Diagonalize** $(T, \alpha)$ . For a tree  $T$  and a real number  $\alpha$  this algorithm outputs a sequence  $(d_v)_{v \in V(T)}$ .

**Algorithm Diagonalize** $(T, \alpha)$

1. List the vertices of  $T$  in postorder as  $v_1, \dots, v_n$ .
2. For each  $i = 1, \dots, n$  set  $d_{v_i} \leftarrow \alpha$ .
3. For each  $i = 1, \dots, n$ :
4. If  $v_i$  has a child  $v_j$  such that  $d_{v_j} = 0$ , then
  - set  $d_{v_i} \leftarrow -\frac{1}{2}$  and  $d_{v_j} \leftarrow 2$ .
  - Further, if  $v_i$  has a parent  $v_p$ , remove the edge  $v_p v_i$  from  $T$ .
5. Otherwise, set  $d_{v_i} \leftarrow d_{v_i} - \sum d_{v_j}^{-1}$ , summing over all children  $v_j$  of  $v_i$ .

The above algorithm of Jacobs and Trevisan [11] can be used to estimate eigenvalues of a given tree. It is based on diagonalization of the matrix  $A(T) + \alpha I$ , where  $A(T)$  is the adjacency matrix of  $T$  and  $\alpha$  is a real number. Its nice feature is that it can be easily executed manually directly on the drawing of a tree. The authors proved that this algorithm diagonalizes  $A(T) + \alpha I$  and, additionally, the following result holds.

**Theorem 2.3.** *For a tree  $T$ , let  $(d_v)_{v \in V(T)}$  be the values produced by **Diagonalize** $(T, -\alpha)$ . Then the diagonal matrix  $D = \text{diag}(d_v)_{v \in V(T)}$  is congruent to  $A(T) + \alpha I$ , hence the number of ( positive | negative | zero ) entries in  $(d_v)_{v \in V(T)}$  is equal to the number of eigenvalues of  $A(T)$  that are ( greater than  $\alpha$  | smaller than  $\alpha$  | equal to  $\alpha$  ).*




FIGURE 6. The algorithm **Diagonalize** $(T, -\lambda)$  applied to a generic member of trees  $\mathfrak{F}(n)$ .

Let  $(d_v)_{v \in V}$  be the sequence obtained by executing **Diagonalize** $(T, -\lambda)$ , when  $T \in \mathfrak{F}(n)$  (see Figure 6). Since we are going to use this algorithm in different trees, it is useful to adopt a new notation, recording the tree we are using and the vertex where

we are applying, which is

$$f_T(v) := d_v, \quad \forall v \in V(T).$$

Taking  $\lambda = \rho(T)$  we will reason that  $f_T(v_n) = 0$  and  $f_T(v_i) < 0$  for  $i \neq n$ .

For notation simplicity, let us rename the values in the extremity of each leaf  $v_1$  by  $a_1 = -\lambda$ , then  $f_T(v_{j+1}) = a_{j+1} = -\lambda - \frac{1}{a_j}$  for  $j \geq 1$ . In particular,  $a_1 \in (-\infty, -2)$  and  $a_2 = -\lambda - \frac{1}{a_1} = -\lambda + \frac{1}{\lambda} \in (-\infty, -\frac{3}{2})$ .

On the vertex with  $r_1 P_2$  and  $r_2 P_2$  the algorithm produces, respectively

$$b_1 = -\lambda - \frac{r_1}{a_2}, \quad \text{and} \quad c_1 = -\lambda - \frac{r_2}{a_2}.$$

From these vertices towards the root  $u$  we obtain two sequences  $b_j$  and  $c_j$ , obeying the same relation as  $a_j$ , that is,  $b_{j+1} = -\lambda - \frac{1}{b_j}$  and  $c_{j+1} = -\lambda - \frac{1}{c_j}$ .

The vertices labeled with the values of the numeric sequences generated by the application of the algorithm **Diagonalize**  $(T, -\lambda)$  appear in Figure 6.

By the application of the algorithm, we know that  $a_j, b_j, c_j < 0$  for all indices appearing in the picture and

$$(1) \quad f_T(u) = -\lambda - \frac{1}{a_h} - \frac{1}{b_{q_1}} - \frac{1}{c_{q_2}} = 0,$$

otherwise, if some previous vertex produces zero then, the step (4) of the algorithm will produce a positive value which is not possible because  $\lambda$  is the index of  $T$ . Now suppose that we have a new tree  $\tilde{T}$  with new parameters  $[h', q'_1, q'_2]$  (same number of vertices and with the same  $r_1$  and  $r_2$ ). We now execute **Diagonalize** $(\tilde{T}, -\lambda)$ .

Since the tree  $\tilde{T}$  has the same properties of  $T$  we obtain the same sequences and the same formula at the root  $u$ . More precisely, the execution of **Diagonalize** $(\tilde{T}, -\lambda)$  produces the same sequences,  $a_j, b_j, c_j$  and

$$(2) \quad f_{\tilde{T}}(u) = -\lambda - \frac{1}{a_h} - \frac{1}{b_{q'_1}} - \frac{1}{c_{q'_2}}$$

From Theorem 2.3 it follows that  $\rho(T) > \rho(\tilde{T})$  if and only if  $-\lambda - \frac{1}{a_h} - \frac{1}{b_{q'_1}} - \frac{1}{c_{q'_2}} < 0$  and all  $a_j, b_j, c_j < 0$ .

We will see that in order to determine the sign of Equation (2), we need to deal in great detail with the recurrences appearing when the algorithm **Diagonalize** $(\tilde{T}, -\lambda)$  is implemented. For that we will determine some analytical properties of the recurrences  $a_j, b_j$  and  $c_j$ .

### 3. ANALYTICAL PROPERTIES OF RECURRENCE SEQUENCES

As we observed in the previous section, the only information needed is the sign of the numeric sequences generated. All the recurrence relations are of the same kind, differing only by the initial value. More precisely, they are of the form

$$(3) \quad z_{j+1} = \varphi(z_j) \quad \text{where} \quad \varphi(t) = -\lambda - \frac{1}{t}$$



for  $t \neq 0$  and  $\lambda = \rho(T) > \rho(T_j) > 2$ . Hence it depends only on the analytical behavior of the function  $\varphi(t)$ .

In [3] this sequence was extensively studied and its behaviour can be summarized by the following result whose proof is a combination of the results found in [3].

**Theorem 3.1.** *Let  $z_j$  be the recurrence in formula (3), then*

- (a)  $\varphi(t) = t$  has two fixed points  $\theta = \frac{-\lambda - \sqrt{\lambda^2 - 4}}{2} < -1$  and  $\theta^{-1}$ ;
- (b)  $\theta(\lambda)$  is decreasing as a function of  $\lambda$ ;
- (c)  $z_j = \theta + \frac{\theta^{-1} - \theta}{\beta(\theta^2)^j + 1}$ , where the constant  $\beta \in \mathbb{R}$  is obtained by choosing the value  $z_1$ ;
- (d) The sequence  $a_j$  obtained from  $z_j$  by taking  $a_1 = -\lambda$  is given by  $a_j = \theta - \frac{\sqrt{\lambda^2 - 4}}{(\theta^2)^j - 1} < 0$ , for any  $j \geq 1$ . In particular,  $a_j$  is increasing and  $\lim_{j \rightarrow \infty} a_j = \theta$ .

We also need to understand the sequences  $b_j$  and  $c_j$  obtained from  $z_j$  by considering  $b_1 = -\lambda - \frac{r_1}{a_2}$  and  $c_1 = -\lambda - \frac{r_2}{a_2}$ ,  $2 \leq r_1 < r_2$ , respectively.

We are going to study both at the same time by considering a parametric sequence

$$(4) \quad z_{j+1}(r) = \varphi(z_j(r))$$

with  $z_1(r) := -\lambda - \frac{r}{a_2}$  for some  $r \geq 2$ .

The main facts can be summarized in the following result.

**Theorem 3.2.** *Let  $z_j$  be the recurrence in formula (4), then*

- (a)  $z_j(r) = \theta + \frac{\theta^{-1} - \theta}{\beta(\theta^2)^j + 1}$ , where the constant  $\beta := \beta(r) \in \mathbb{R}$  is given by

$$\beta := \frac{r - a_2\theta}{a_2\theta - r\theta^2};$$

- (b)  $\beta(r)$  is a continuous function of  $r$  in  $(2, \infty) \setminus r_*$ , where  $r_* := \frac{a_2}{\theta}$ . Moreover  $\beta(r)$  has a single root in  $r^* = a_2\theta$  and  $\beta(r) > 0$ , for  $r \in (r_*, r^*)$
- (c) The sequence  $z_j(r) < 0$ , for  $j \geq 1$ , is decreasing and  $\lim_{j \rightarrow \infty} z_j = \theta$ .

*Proof.* **(a)** Using Theorem 3.1 (c) and the fact that  $-\lambda = \theta + \theta^{-1}$  and  $\theta - \theta^{-1} = -\sqrt{\lambda^2 - 4}$  we obtain

$$z_1 = -\lambda - \frac{r}{a_2} = \theta + \theta^{-1} - \frac{r}{a_2} \text{ and } z_1 = \theta + \frac{\theta^{-1} - \theta}{\beta(\theta^2)^1 + 1},$$

producing

$$\beta := \beta(r) = \frac{\theta^{-1} \frac{r}{a_2} - 1}{1 - \theta \frac{r}{a_2}} = \frac{r\theta^{-1} - a_2}{a_2 - r\theta} = \frac{r - a_2\theta}{a_2\theta - r\theta^2}.$$

**(b)** We notice that  $\beta(r)$  is a rational function, hence continuous, except for the roots of the denominator. Thus, the discontinuity occurs at  $r_* := \frac{a_2}{\theta}$ . Also,  $\beta(r)$  has only one possible root in  $r^* = a_2\theta$ .

Additionally,  $\lim_{r \rightarrow r_*^+} \beta(r) = +\infty$ . To see that we just take  $r := r_* + \delta$  for  $\delta > 0$ , then

$$\beta(r_* + \delta) = \frac{r_* + \delta - a_2\theta}{a_2\theta - (r_* + \delta)\theta^2} = \frac{\delta + (r_* - a_2\theta)}{(a_2\theta - r_*\theta^2) - \delta\theta^2} = \frac{\delta + (r_* - a_2\theta)}{-\delta\theta^2} \xrightarrow{\delta \rightarrow 0^+} +\infty$$

because  $r_* - a_2\theta = \frac{a_2}{\theta} - a_2\theta = a_2\left(\frac{1-\theta^2}{\theta}\right) < 0$ .




FIGURE 7. The behaviour of  $\beta(r)$ .

As  $r^* - r_* = a_2(\theta - \theta^{-1}) = a_2(-\sqrt{\lambda^2 - 4}) > 0$ , we see that  $r_* < r^*$ . Also, differentiating with respect to  $r$  we conclude that  $\beta(r)$  is decreasing and take the value zero only for  $r^* := a_2\theta > r_*$ . Thus we conclude that  $\beta(r) > 0$  for  $r \in (r_*, r^*)$ . Figure 7 illustrates a typical behaviour of the function  $\beta(r)$ .

(c) We recall that  $r^* = a_2\theta$ , moreover, from Theorem 3.1 (d), we know that  $\lim_{j \rightarrow \infty} a_j = \theta$  hence

$$(5) \quad r^* = \lim_{j \rightarrow \infty} a_2 a_j.$$

By Theorem 2.1, we know that  $\rho(T_j) < \rho(T)$ , where  $T_j$  is the tree of Figure 5. Now, we apply **Diagonalize**( $T_j, -\lambda$ ) with the root at  $S_{r_2}$ , for  $\lambda = \rho(T)$ . By our comparison method, we see that

$$f_{T_j}(u) = -\lambda - \frac{1}{a_j} - \frac{r_2}{a_2} < 0$$

or, equivalently,  $a_{j+1} < \frac{r_2}{a_2}$ , and since  $a_2 < 0$ , we get  $r_2 < a_2 a_{j+1}$ .

Taking the limit on both sides and using Equation (5), we obtain

$$r_2 \leq \lim_{j \rightarrow \infty} a_2 a_{j+1} = \lim_{j \rightarrow \infty} a_2 a_j = r^*.$$

We also observe that  $r_* < r_1$ , because  $r_* = \frac{a_2}{\theta} = 2 \frac{\lambda^2 - 1}{\lambda(\lambda + \sqrt{\lambda^2 - 4})} < 2 \leq r_1$ .

Since  $z_j = \theta + \frac{\theta^{-1} - \theta}{\beta(r)(\theta^2)^{j+1}}$ , for  $\theta^{-1} - \theta = \sqrt{\lambda^2 - 4} > 0$  and  $\theta < 0$ , we see that  $z_j$  is always negative and decreasing, as long as  $\beta(r) > 0$ . Now, because  $r_* < r_1 < r_2 \leq r^*$ , we see from item (b), that  $\beta(r) > 0$ . Moreover  $z_j$  tends to  $\theta$  as  $j \rightarrow \infty$ .  $\square$

#### 4. ORDERING THE 2-SWITCHES OF $\mathfrak{F}(n)$

We show in this section how the spectral radius varies in each case for all the possible 2-switching positions in the appropriate interval. We notice that  $\mathfrak{F}(n)$  is preserved by both types of  $(s, t)$ -2-switch, only changing  $[h, q_1, q_2]$  (see Figure 6).

**4.1. Warmup: Ordering 2-switches of Type I.** Given a 2-switch of Type I such that  $T = [h, q_1, q_2] \rightarrow \tilde{T} = [h', q'_1, q'_2]$  (see Figure 3), we observe that the actual result of the operation in the tree is an increment (decrement) of the length  $h$  with a decrement (increment) of the length  $q_1$ , while  $q_2$  remains unchanged.

In order to study the behavior of the spectral radius of members of this family, it is enough to study the 2-switch  $T = [h, q_1, q_2] \rightarrow \tilde{T} = [h' = h - 1, q'_1 = q_1 + 1, q'_2 = q_2]$ , since this will cover all possible positions.

We will prove that  $\rho(T) > \rho(\tilde{T})$  using our comparison method, hence we need to prove that if  $-\lambda - \frac{1}{a_h} - \frac{1}{b_{q_1}} - \frac{1}{c_{q_2}} = 0$  then  $-\lambda - \frac{1}{a_{h'}} - \frac{1}{b_{q'_1}} - \frac{1}{c_{q_2}} = -\lambda - \frac{1}{a_{h-1}} - \frac{1}{b_{q_1+1}} - \frac{1}{c_{q_2}} < 0$ . Notice that, from the first equation, we obtain  $-\lambda - \frac{1}{c_{q_2}} = \frac{1}{a_h} + \frac{1}{b_{q_1}}$  and substituting in the second one it is equivalent to  $\frac{1}{a_h} + \frac{1}{b_{q_1}} - \frac{1}{a_{h-1}} - \frac{1}{b_{q_1+1}} < 0$ , which in turn is equivalent to

$$(6) \quad (a_h - a_{h+1}) + (b_{q_1+2} - b_{q_1+1}) < 0.$$

From Theorem 3.1, the sequence  $a_j$  is increasing thus  $a_h - a_{h+1} < 0$ . From Theorem 3.2, the sequence  $b_j$  is decreasing thus  $b_{q_1+2} - b_{q_1+1} < 0$ .

We remark that the transformation  $T = [h, q_1, q_2] \rightarrow \tilde{T} = [h' = h - 1, q'_1 = q_1, q'_2 = q_2 + 1]$  is also of Type-I and using a similar argument, we can show that the index decreases as well. This proves the following theorem.

**Theorem 4.1.** *Let  $T = [h, q_1, q_2]$  be a tree in  $\mathfrak{F}(n)$  and  $\tilde{T} = [h', q'_1, q'_2]$  be the graph obtained by a 2-switch in Figure 3. If  $h' = h - 1, q'_1 = q_1 + 1$  and  $q'_2 = q_2$  or if  $h' = h - 1, q'_1 = q_1$  and  $q'_2 = q_2 + 1$  then  $\rho(T) > \rho(\tilde{T})$ .*

We remark that this result may be obtained also by using Lemma 2.2 due to Hoffman & Smith [9]. We add an edge on the internal path from  $u$  to  $r_1 P_2$  (or from  $u$  to  $r_2 P_2$ ) and then erase the pendant vertex from  $P_h$ , so that the spectral radius decreases even more, keeping both with the same number of vertices.

Our method, after we obtained that the sequence  $b_j$  is decreasing, is simple enough to provide the alternative proof. We observe, however, that for 2-switching of Type II, we are not aware of a known result that apply. Additionally, or perhaps because of that, the application of our method requires to overcome quite a few technical difficulties.

**4.2. Ordering 2-switches of Type II.** We observe that a 2-switch of type II can be seen as a displacement of the central path of length  $h$  from the position closest to  $r_1 P_2$  to the closest to  $r_2 P_2$  (or vice-versa). Indeed, if we take  $s = q_2$  then  $q'_1 = q_2 + t - s = t$  and  $q'_2 = q_1 - t + q_2 = q_1 + q_2 - t$ , for  $1 \leq t \leq q_1 - 1$ . For instance, taking  $t = 1$  we can apply the 2-switch sequentially.

Since every configuration  $[h, q_1, q_2]$  can be obtained by successive changes by 1, we only need to consider the case where  $T = [h, q_1, q_2] \rightarrow \tilde{T} = [h, q'_1 = q_1 - 1, q'_2 = q_2 + 1]$ . We will prove that this operation decreases the spectral radius

By using our method of Section 2, given a 2-switch of Type II we need to prove that if  $-\lambda - \frac{1}{a_h} - \frac{1}{b_{q_1}} - \frac{1}{c_{q_2}} = 0$  then  $-\lambda - \frac{1}{a_h} - \frac{1}{b_{q_1-1}} - \frac{1}{c_{q_2+1}} < 0$ .

**Theorem 4.2.** *Let  $T = [h, q_1, q_2]$  be a tree obtained by a 2-switch in Figure 4. If  $r_2 > r_1 \geq 2, q'_1 = q_1 - 1$  and  $q'_2 = q_2 + 1$  then  $\rho(T) > \rho(T')$ .*

*Proof.* We recall that we need to prove that, given a 2-switch of Type II, we need to prove that if  $-\lambda - \frac{1}{a_h} - \frac{1}{b_{q_1}} - \frac{1}{c_{q_2}} = 0$  then  $-\lambda - \frac{1}{a_h} - \frac{1}{b_{q_1-1}} - \frac{1}{c_{q_2+1}} < 0$ . We define

$$(7) \quad \mathcal{I} := -\frac{1}{a_h} + \left(-\lambda - \frac{1}{b_{q_1-1}}\right) - \frac{1}{c_{q_2+1}} = -\frac{1}{a_h} + b_{q_1} - \frac{1}{c_{q_2+1}}$$

and, from the first equation,  $-\frac{1}{a_h} - \frac{1}{b_{q_1}} + \left(-\lambda - \frac{1}{c_{q_2}}\right) = 0$  or  $c_{q_2+1} = \frac{1}{a_h} + \frac{1}{b_{q_1}}$ . Substituting that in (7) we obtain

$$(8) \quad \mathcal{I} = -\frac{1}{a_h} + b_{q_1} - \frac{1}{\frac{1}{a_h} + \frac{1}{b_{q_1}}}.$$

Using the fact that  $a_h < 0$  and  $b_{q_1} < 0$  conclude that  $\mathcal{I} < 0$  if and only if

$$(9) \quad a_h(b_{q_1}^2 - 1) < b_{q_1}.$$

We already know that  $a_h < \theta < b_{q_1}$  but we do not know the sign of  $b_{q_1}^2 - 1$ . We claim that  $b_{q_1}^2 - 1 > 0$ . To see that, we recall that  $b_j$  is decreasing and  $b_1$  is given by  $b_1 := -\lambda - \frac{r_1}{-\lambda + \frac{1}{\lambda}} = -\lambda - \frac{2}{-\lambda + \frac{1}{\lambda}} = \phi(\lambda, r_1)$  where the auxiliary function  $\phi : A \rightarrow \mathbb{R}$  is given by

$$\phi(t, r) := -t - \frac{r}{-t + \frac{1}{t}},$$

defined on the set  $A = \{(t, r) \mid t \geq 2, r \geq 2\} \subset \mathbb{R}^2$ .




FIGURE 8. The graph of  $\phi(t, r)^2 - 1$  plotted in  $A$ .

It is easy to see that  $\phi(t, r)^2 - 1 > 0$  for  $(t, r) \in A$ , see Figure 8. In other words,  $b_1^2 - 1 > 0$  and  $b_1 < -1$  because it is negative. For any  $T \in \mathfrak{F}(n)$  we conclude that  $\theta < b_{q_1} < -1 = b_1$ , for any  $T \in \mathfrak{F}(n)$ . In particular  $b_{q_1}^2 - 1 > 0$ .

From these facts, we can rewrite equation (9) in the equivalent form

$$(10) \quad a_h < \frac{b_{q_1}}{b_{q_1}^2 - 1}.$$

In order to conclude our proof, it is sufficient to prove that  $\theta < \frac{b_{q_1}}{b_{q_1}^2 - 1}$  because  $a_h < \theta$  for all  $h$ . At this point, it is useful to introduce a second auxiliary function

$$\psi(t) := \frac{t}{t^2 - 1}, t < 0.$$

This function is obviously decreasing in the interval  $(-\infty, -1)$ . As the correspondence  $j \rightarrow b_j$  is also decreasing, we conclude that the correspondence  $j \rightarrow \psi(b_j)$  is increasing and, as a consequence,  $\frac{b_1}{b_1^2 - 1} < \frac{b_{q_1}}{b_{q_1}^2 - 1}$ .

We claim that  $\frac{b_1}{b_1^2 - 1} > \theta$  or equivalently

$$(11) \quad \frac{\left(-\lambda - \frac{r_1}{-\lambda + \frac{1}{\lambda}}\right)}{\left(-\lambda - \frac{r_1}{-\lambda + \frac{1}{\lambda}}\right)^2 - 1} > \frac{-\lambda - \sqrt{\lambda^2 - 4}}{2},$$

for  $\lambda = \rho(T)$ .

The inequality (11) is equivalent to  $g(x, r) > 0$  for  $x \geq \rho(T) \geq \sqrt{r_2 + 2} \geq \sqrt{r_1 + 3}$  and  $r = r_1 \geq 2$ , where  $g : B \rightarrow \mathbb{R}$  is given by

$$g(x, r) = \frac{\left(-x - \frac{r}{-x + \frac{1}{x}}\right)}{\left(-x - \frac{r}{-x + \frac{1}{x}}\right)^2 - 1} + \frac{x + \sqrt{x^2 - 4}}{2},$$

defined on  $B = \{(x, r) \mid x \geq \sqrt{r + 3}, r \geq 2\} \subset \mathbb{R}^2$ .




FIGURE 9. The graph of  $g$  on  $B$ .

As we can see in Figure 9, the function is always positive in this set concluding our proof.  $\square$

## 5. SPECTRAL RADIUS ORDERING IN $\mathfrak{F}(n)$

In this section we provide a total ordering by the spectral radius in the family  $\mathfrak{F}(n)$ . In order to recall our notation, we notice that  $r_1 > r_2 \geq 2$  and the number  $n$  of vertices

are fixed, hence each element in  $\mathfrak{F}(n)$  is uniquely determined by the 3-uple  $[h, q_1, q_2]$ , that is,

$$\mathfrak{F}(n) := \{T = [h, q_1, q_2] \mid h + q_1 + q_2 = n - 1 - 2(r_1 + r_2), h, q_1, q_2 \geq 2\}.$$

The degree sequence of an element  $T \in \mathfrak{F}(n)$  is given by  $d := [r_1+1, r_2+1, 3, 2^{n-(r_1+r_2)-2}, 1^{r_1+r_2+1}]$ , for a fixed pair  $r_1 > r_2 \geq 2$ . We recall the well known result from [8] (also [4]):

**Theorem 5.1.** *If  $G$  e  $H$  have the same degree sequence then, there exists a 2-switch sequence transforming  $G$  into  $H$ .*

As a consequence, each two elements in  $\mathfrak{F}(n)$  are transformed into another by a sequence of 2-switch transformations.

The remarkable fact is that, as we will see next, we can reach any element from another using only two 2-switches. Additionally, the sequence of 2-switches is closed in the family, that is, any intermediate member is also a member. Given  $T = [h, q_1, q_2] \in \mathfrak{F}(n)$  we define the following operations.

- $\alpha : \mathfrak{F}(n) \rightarrow \mathfrak{F}(n)$ , given by  $\alpha([h, q_1, q_2]) = [h - 1, q_1 + 1, q_2]$ , for  $h \geq 3$ ;
- $\beta : \mathfrak{F}(n) \rightarrow \mathfrak{F}(n)$ , given by  $\beta([h, q_1, q_2]) = [h, q_1 - 1, q_2 + 1]$ , for  $q_1 \geq 3$ .
- $\gamma : \mathfrak{F}(n) \rightarrow \mathfrak{F}(n)$ , given by  $\beta([h, q_1, q_2]) = [h - 1, q_1, q_2 + 1]$ , for  $h \geq 3$ .

We remark that the transformations make sense for  $h, q_1, q_2 \geq 1$ , and the results from Section 4 do apply. However, we observe that the original  $T = [h, q_1, q_2]$  and the transformed tree  $T' = [h', q'_1, q'_2]$  must have parameters  $h, h', q_1, q'_1, q_2, q'_2 \geq 2$ , otherwise they will not be 2-switches, because the degree sequence changes.

**Theorem 5.2.** *Let  $\alpha, \beta$  and  $\gamma$  be the transformations defined in  $\mathfrak{F}(n)$ . Let  $T^* = [h_0, 2, 2]$ , where  $h_0 := n - 5 - 2(r_1 + r_2) \geq 2$ . The following facts are true*

- (a)  $\alpha, \beta$  and  $\gamma$  are 2-switch transformations;
- (b)  $\alpha, \beta$  and  $\gamma$  are index decreasing transformations;
- (c) Any  $T \in \mathfrak{F}(n)$  can be obtained from  $T^*$  by a sequence of  $\alpha$  and  $\beta$  transformations;
- (d) Any  $T \in \mathfrak{F}(n)$  can be obtained from  $T^*$  by a sequence of  $\alpha$  and  $\gamma$  transformations.

*Proof.* From our definition of Section 2, we see that  $\alpha$  and  $\gamma$  are Type-I 2-switches, while  $\beta$  is a Type-II 2-switch and our results of Theorem 4.1 and Theorem 4.2 apply, hence they decrease the index. This proves (a) and (b).

(c) To see that, consider the tree  $T^* = [h_0, 2, 2]$ . As  $T^* = u + P_{h_0} * S_0 \oplus P_2 * S_{r_1} \oplus P_2 * S_{r_2}$ , and  $h_0 = n - 5 - 2(r_1 + r_2) \geq 2$ , because  $n \geq 7 + 2(r_1 + r_2)$ , we see that  $T^* \in \mathfrak{F}(n)$ . Let  $T = [h', q'_1, q'_2]$  be any tree in  $\mathfrak{F}(n)$  ( $h', q'_1, q'_2 \geq 2$ ). Then we have the following  $\alpha([h_0, 2, 2]) = [h_0 - 1, 2 + 1, 2]$ ,  $\alpha([h_0 - 1, 2 + 1, 2]) = [h_0 - 2, 2 + 2, 2]$ , and so on, until we obtain  $h_0 - k = h'$ , that is  $\alpha^k([h_0, 2, 2]) = [h', 2 + (h_0 - h'), 2]$ .

Now we apply  $\beta$  transformation  $j$  times obtaining  $\beta^j \alpha^k([h_0, 2, 2]) = [h', 2 + (h_0 - h') - j, 2 + j]$ .

At this point we claim that, making  $2 + (n - 5 - 2(r_1 + r_2) - h') - j = q'_1$  we get  $2 + j = q'_2$ . Indeed,  $2 + (n - 5 - 2(r_1 + r_2) - h') - j = q'_1$  means that  $j = 2 + (n - 5 - 2(r_1 + r_2) - h' - q'_1) - q'_2 + q'_2 = q'_2 - 2 + (n - (1 + h' + q'_1 + q'_2 + 2(r_1 + r_2))) = q'_2 - 2$ . Hence  $2 + j = q'_2$ . Thus we conclude that there exist  $k, j \in \mathbb{N}$  such that  $\beta^j \alpha^k(T^*) = T$  for any  $T \in \mathfrak{F}(n)$ .

(d) A similar reasoning as in (c) shows that  $T^* \in \mathfrak{F}(n)$ . We now apply  $k' = q'_1 - 2$  times the transformation  $\alpha$ , arriving at  $[h_0 - k', q'_1, 2]$ . Reasoning as above we show that there is an integer  $j'$  so that  $h_0 - k' - j' = h'$  and  $q'_2 = 2 + j'$ . Hence applying now  $j'$  times the transformation  $\gamma$ , shows that  $\gamma^{j'} \alpha^{k'}(T^*) = T$ .  $\square$

We observe that  $T^* = [h_0, 2, 2]$  has the configuration with largest possible  $h_0$ . From Theorem 5.2 (b) and (c) (or from (b) and (d)), it follows that  $T^*$  is the extremal element of  $\mathfrak{F}(n)$ : it has the maximum spectral radius.

The next result is quite remarkable in the sense that it provides a complete ordering of  $\mathfrak{F}(n)$  using only  $\alpha$  and  $\gamma$  transformations, allowing us to find also the element of minimum index.

**Theorem 5.3.** *Consider the family  $\mathfrak{F}(n)$ , where  $2 \leq r_1 < r_2$ ,  $h_0 = n - 5 - 2(r_1 + r_2) \geq 2$ ,  $T^* = [h_0, 2, 2] \in \mathfrak{F}(n)$  and  $T_* = [2, 2, h_0] \in \mathfrak{F}(n)$ . The following claims are true.*

(a) *The ordered sequence*

$$\mathcal{A} := \{T^*, \alpha(T^*), \dots, \alpha^{h_0-2}(T^*), \gamma(T^*), \alpha(\gamma(T^*)), \dots, \alpha^{h_0-3}(\gamma(T^*)), \dots, \gamma^{h_0-2}(T^*) = T_*\}$$

*is equal to  $\mathfrak{F}(n)$ ;*

(b) *The sequence  $\mathcal{A}$  is ordered by the inverse lexicographic order:*

*$[x, y, z] \succ [x', y', z']$  iff  $z' > z$  or  $z' = z$  but  $y' > y$ .*

(c) *If  $[x, y, z] \succ [x', y', z']$  then  $\rho([x, y, z]) > \rho([x', y', z'])$ ;*

(d) *The maximum (resp. minimum) index in  $\mathfrak{F}(n)$  is  $\rho(T^*)$  (resp.  $\rho(T_*)$ ).*

Before we prove Theorem 5.3 we need a technical lemma.

**Lemma 5.4.** *If  $T^* = [h_0, 2, 2] \in \mathfrak{F}(n)$ , then  $\rho(\alpha^{h_0-j-2}(\gamma^j(T^*))) > \rho(\gamma^{j+1}(T^*))$ , for  $j = 0, 1, 2, \dots, h_0 - 2$ , that is,*

$$\rho([2, h_0 - j, 2]) > \rho([h_0 - (j + 1), 2, 2 + (j + 1)]).$$

*Proof.* We will consider the first case  $j = 0$ . The rest of the cases are identical. Thus, we must prove that

$$\rho([2, h_0, 2]) > \rho([h_0 - 1, 2, 3]).$$

Taking  $\lambda = \rho([2, h_0, 2])$  and applying  $\text{Diagonalize}([2, h_0, 2], -\lambda)$ , we have

$$-\lambda - \frac{1}{a_2} - \frac{1}{b_{h_0}} - \frac{1}{c_2} = 0.$$

From this, we obtain  $c_3 = \frac{1}{\frac{1}{a_2} + \frac{1}{b_{h_0}}}$ .

Analogously, applying  $\text{Diagonalize}([h_0 - 1, 2, 3], -\lambda)$ , we have

$$-\lambda - \frac{1}{a_{h_0-1}} - \frac{1}{b_2} - \frac{1}{c_3} =: \mathcal{I}.$$

We need to show that  $\mathcal{I} < 0$ . Writing  $\mathcal{I} = a_{h_0} - \frac{1}{b_2} - \frac{1}{\frac{1}{a_2} + \frac{1}{b_{h_0}}}$ , we conclude that  $\mathcal{I} < 0$  if the function

$$g(\lambda, r_1, h_0) := a_{h_0} - \frac{1}{b_2} - \frac{1}{\frac{1}{a_2} + \frac{1}{b_{h_0}}}$$

is always negative for  $r_1 \geq 2$ ,  $h_0 \geq 2$  and  $\lambda \geq \sqrt{r_1 + 3}$ .

At a first glance, we can not plot a graph as we did before because we have three variables. However, if we consider the variation of the variable  $h_0$  we observe that the correspondence  $h_0 \rightarrow g(\lambda, r_1, h_0)$  is monotonously increasing because

$$\frac{d}{dh_0}g(\lambda, r_1, h_0) = \frac{da_{h_0}}{dh_0} - \frac{1}{\left(\frac{1}{a_2} + \frac{1}{b_{h_0}}\right)^2} \frac{1}{b_{h_0}^2} \frac{db_{h_0}}{dh_0} > 0,$$

since  $\frac{da_{h_0}}{dh_0} > 0$  and  $\frac{db_{h_0}}{dh_0} < 0$ .

Hence we just need to show that the limit function

$$f(\lambda, r_1) := \lim_{h_0 \rightarrow \infty} g(\lambda, r_1, h_0) = \theta - \frac{1}{b_2} - \frac{1}{\frac{1}{a_2} + \frac{1}{\theta}}$$

is always negative for  $C := \{(\lambda, r_1) \mid r_1 \geq 2, \lambda \geq \sqrt{r_1 + 3}\}$ , as shown on Figure 10, concluding our proof.




FIGURE 10. The graph of  $f$  on  $C$ .

□

*Proof. (of Theorem 5.3)*

To see (a) we first see that both transformations  $\alpha$  and  $\gamma$  decrease the index. By Theorem 5.2 (d), we can reach any configuration from  $T^*$ , showing that every configuration appears in the sequence  $\mathcal{A}$ .

For (b), we notice that the inverse lexicographic order:

$[x, y, z] \succ [x', y', z']$  if and only if  $z' > z$  or  $z' = z$  but  $y' > y$ , is naturally produced in  $\mathcal{A}$ . For instance comparing  $[x, y, z]$  with  $[x', y', z'] = \alpha([x, y, z]) = [x - 1, y + 1, z]$  we obtain  $z' = z$  but  $y' > y$ . The only possible difficulty is to compare, for example,  $[x, y, z] = \alpha^{h_0-2}(T^*) = [2, h_0 - 2, 2]$  with  $[x', y', z'] = \gamma(T^*) = [h_0 - 1, 2, 3]$ , in this case we have  $z' > z$ .

For (c) we can use the same reasoning, if the next element is obtained from the previous one by  $\alpha$  the index decreases by Theorem 4.1. Again, it remains to analyse the case  $[x, y, z] = \alpha^{h_0-2}(T^*) = [2, h_0 - 2, 2]$  and  $[x', y', z'] = \gamma(T^*) = [h_0 - 1, 2, 3]$ . In this case the index decrease by Lemma 5.4.

The item (d) is a direct consequence of the previous items. □



**Example 5.5.** We consider  $\mathfrak{F}(23)$ , with  $r_1 = 2$ ,  $r_2 = 3$  and  $h_0 = n - 5 - 2(r_1 + r_2) = 8$ . In this case  $T^* = [h_0, 2, 2] = [8, 2, 2]$  and  $T_* = [2, 2, h_0] = [2, 2, 8]$ . We will use the same procedure described in the proof of Theorem 5.3 to build a table where we show the spectral radius of each intermediary tree:

Transf. / $\mathcal{A}$	Tree	Index
$T^*$	[8, 2, 2]	2.31431268823172996316982502630
$\alpha(T^*)$	[7, 3, 2]	2.30752321205156788164155922354
$\alpha^2(T^*)$	[6, 4, 2]	2.30509257122848263666229555974
$\alpha^3(T^*)$	[5, 5, 2]	2.30414417603593895847264293478
$\alpha^4(T^*)$	[4, 6, 2]	2.30348720654135784657134525755
$\alpha^5(T^*)$	[3, 7, 2]	2.30226165044440472718571097461
$\alpha^6(T^*)$	[2, 8, 2]	2.29881642949995094980856594643
$\gamma(T^*)$	[7, 2, 3]	2.28520768467980257500859073365
$\alpha(\gamma(T^*))$	[6, 3, 3]	2.28076523286917478390041282633
$\alpha^2(\gamma(T^*))$	[5, 4, 3]	2.27913084342903308996825366690
$\alpha^3(\gamma(T^*))$	[4, 5, 3]	2.27834791245706879712729095155
$\alpha^4(\gamma(T^*))$	[3, 6, 3]	2.27748824925244285093685838480
$\alpha^5(\gamma(T^*))$	[2, 7, 3]	2.27554403106324050144208754160
$\gamma^2(T^*)$	[6, 2, 4]	2.27010998510725135104117051475
$\alpha(\gamma^2(T^*))$	[5, 3, 4]	2.26762484634172519636930335282
$\alpha^2(\gamma^2(T^*))$	[4, 4, 4]	2.26667762008239070931668388638
$\alpha^3(\gamma^2(T^*))$	[3, 5, 4]	2.26605728367174815669677409819
$\alpha^4(\gamma^2(T^*))$	[2, 6, 4]	2.26506821261118740374393886088
$\gamma^3(T^*)$	[5, 2, 5]	2.26290253458453744084697620016
$\alpha(\gamma^3(T^*))$	[4, 3, 5]	2.26171078345443097808224085587
$\alpha^2(\gamma^3(T^*))$	[3, 4, 5]	2.26119844487818869745804831320
$\alpha^3(\gamma^3(T^*))$	[2, 5, 5]	2.26069897200749878293447592468
$\gamma^4(T^*)$	[4, 2, 6]	2.25980268994372236598891968054
$\alpha(\gamma^4(T^*))$	[3, 3, 6]	2.25927957177517211016191460326
$\alpha^2(\gamma^4(T^*))$	[2, 4, 6]	2.25898741243972985580277387992
$\gamma^5(T^*)$	[3, 2, 7]	2.25857320563154910353684549335
$\alpha(\gamma^5(T^*))$	[2, 3, 7]	2.25834278165321357168906331906
$\gamma^6(T^*) = T_*$	[2, 2, 8]	2.25810972712429442797185863240

As expected, the index ordering in  $\mathfrak{F}(23)$  is total.

*Remark 1.* Theorem 5.3 has a powerful application, not only we can obtain the extremal indices in  $\mathfrak{F}(n)$ , but given any  $[x, y, z] \in \mathfrak{F}(n)$  and a 2-switch  $F$ , such that  $F([x, y, z]) = [x', y', z']$  we can immediately say if  $F$  increases or decreases the index by using the order relation in Theorem 5.3 (c).

## 6. ADDITIONAL CONSIDERATIONS ON THE 2-SWITCH SPECTRAL ORDERING PROBLEM

We believe that transformations by 2-switches which are monotonous with respect to the spectral radius consist of a powerful tool for spectral graph theory problems. We

just recall that, following [13] we have the following operation which was a central piece to prove that all the starlike trees are completely ordered by their index in our recent work [14].

**Lemma 6.1.** [7, 13]. *Let  $u$  be a vertex of a non-trivial connected graph  $G$ , and let  $G_{k,l}^0$  denote the graph obtained from  $G$  by adding pendant paths of length  $k$  and  $l$  at  $u$ . If  $k \geq l \geq 1$ , then  $\lambda_1(G_{k,l}^0) > \lambda_1(G_{k+1,l-1}^0)$ .*

We can rewrite this result as a 2-switch spectral ordering problem. Define  $\mathfrak{G}$  the family of all trees  $T$  given by Figure 11,




FIGURE 11. The family of trees  $\mathfrak{G}$ .

with the following constraints.

- (1)  $q_1, q_2 \geq 2$ ;
- (2)  $T_0$  is any fixed tree.

Notice that, in this case,  $n = |T| = |T_0| + q_1 + q_2$ .


If we are interested in to the maximization/minimization of the index (spectral radius) of trees with a fixed degree sequence, in  $\mathfrak{F}(n)$  we get the degree sequence

$$d := [d_{T_0}, 2^{(q_1+q_2)-2}, 1^2],$$

where  $d_{T_0}$  is the sequence of degrees in  $T_0$  including the root  $u$ . In this way each element in  $\mathfrak{G}$ , with  $n$  vertices and degree sequence  $d$  is uniquely determined by the pair  $[q_1, q_2]$  that is,

$$\mathfrak{G}(n) := \{[q_1, q_2] \mid n = |T_0| + q_1 + q_2\}.$$

To preserve the degree sequence we can do a 2-switch as in Figure 12, obtaining a new tree  $T'$ .

FIGURE 12. 2-Switch from  $T$  to  $T'$ .

Taking  $\lambda$  as the index of  $T$  and applying the Diagonalize algorithm to the trees  $T$  and  $T'$  we get  $-\lambda - \xi - \frac{1}{a_{q_1}} - \frac{1}{a_{q_2}} = 0$ , where  $\xi$  is the contribution of  $T_0$  and,  $-\lambda - \xi - \frac{1}{a_{q'_1}} - \frac{1}{a_{q'_2}} < 0$  if and only if  $\rho(T) > \rho(T')$ .

Assuming,  $q'_1 = q_1 - 1$ ,  $q'_2 = q_2 + 1$  and  $q_1 < q_2 + 1$  we get, substituting  $-\lambda - \xi = \frac{1}{a_{q_1}} + \frac{1}{a_{q_2}}$  in the inequality, the equivalency

$$\frac{1}{a_{q_1}} + \frac{1}{a_{q_2}} - \frac{1}{a_{q_1-1}} - \frac{1}{a_{q_2+1}} < 0.$$

Adding and subtracting  $-\lambda$  and using the formula (4) we obtain that  $\rho(T) > \rho(T')$  if and only if

$$(12) \quad a_{q_1} + \frac{1}{a_{q_1}} < a_{q_2+1} + \frac{1}{a_{q_2+1}}.$$

We notice that  $a_j$  is increasing and  $a_j < \theta < -1$ . The function  $\phi(x) = x + \frac{1}{x}$  is increasing in the interval  $(-\infty, -1)$ . Also,  $a_{q_1} < a_{q_2+1}$  because we assume  $q_1 < q_2 + 1$ . Thus  $\phi(a_{q_1}) < \phi(a_{q_2+1})$ , which is exactly the claim in the equation (12). We have proved the following result which is equivalent to Lemma 6.1.

**Theorem 6.2.** *Let  $T$  be the graph in Figure 11 and  $T'$  be the graph obtained by a 2-switch in Figure 12. If  $q_1 < q_2 + 1$ ,  $q'_1 = q_1 - 1$  and  $q'_2 = q_2 + 1$  then  $\rho(T) > \rho(T')$ .*

#### ACKNOWLEDGMENTS

This work had the financial support of project MATHAmSud 88881.143281/2017-01. Victor N. Schvllöner was partially supported by PROICO 03-0918 UNSL. V. Trevisan acknowledges partial support of CNPq grants 409746/2016-9 and 303334/2016-9, CAPES-Print 88887.467572/2019-00 and FAPERGS project PqG 17/2551-0001.

#### REFERENCES

- [1] F. Belardo, E.M. Li Marzi, S.K. Simić, On the index of caterpillars, *Discrete Mathematics* 308 (2008) 324–330.

- [2] F. Belardo, E.M. Li Marzi, S.K. Simič, Ordering graphs with index in the interval  $(2, \sqrt{2 + \sqrt{5}})$ , *Discrete Appl. Math.* 156 (2008) 1670 – 1682.
- [3] F. Belardo, E. R. Oliveira, V. Trevisan, Spectral ordering of trees with small index, *Linear Algebra and its Applications*, Volume 575, 2019, Pages 250-272, ISSN 0024-3795, <https://doi.org/10.1016/j.laa.2019.04.012>.
- [4] C. Berge, “Graphes et hypergraphes”, *Monographies Universitaires de Mathématiques*, Number 37, Dunod, Paris, 1970.
- [5] T. Biyikoglu, J. Leydold, Graphs with given degree sequence and maximal spectral radius, *Electronic Journal of Combinatorics*, 15 #R119 (2008).
- [6] A. Chang, Q. Huang, Ordering trees by their largest eigenvalues, *Linear Algebra Appl.* 370 (2003) 1751–84.
- [7] S. Simić D. Cvetković, P. Rowlinson. *Eigenspaces of Graphs*. Cambridge University Press, 1997.
- [8] D.R. Fulkerson, A.J. Hoffman, M.H. McAndrew, Some properties of graphs with multiple edges, *Canad. J. Math.* 17 (1965) 166-177.
- [9] A.J. Hoffman, J.H. Smith, On the spectral radii of topologically equivalent graphs, M. Fiedler (Ed.), *Recent Advances in Graph theory*, Academia Praha, Prague (1975), pp. 273-281
- [10] M. Hofmeister, On the two largest eigenvalues of trees, *Linear Algebra Appl.* 260 (1997) 43–59.
- [11] David P. Jacobs, Vilmar Trevisan, Locating the eigenvalues of trees, *Linear Algebra Appl.* 434 (2011) 81–88.
- [12] David P. Jacobs, Elismar R. Oliveira, Vilmar Trevisan, Most Laplacian eigenvalues of a tree are small, preprint(2020).
- [13] Wenshui Lin and Xiaofeng Guo. Ordering trees by their largest eigenvalues. *Linear Algebra and its Applications*, 418(2):450 – 456, 2006.
- [14] Elismar Oliveira, Dragan Stavonović, Vilmar Trevisan, Spectral radius ordering of starlike trees, *Linear Multilinear Algebra* (2018), in press, doi:10.1080/03081087.2018.1524435.
- [15] F. Zhang, Z. Chen, Ordering graphs with small index, *Linear Algebra Appl.* 121 (2002) 295–306.

INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UFRGS, PORTO ALEGRE, BRAZIL  
*E-mail address:* `elismar.oliveira@ufrgs.br`

INSTITUTO DE MATEMÁTICA APLICADA SAN LUIS (UNSL-CONICET), UNIVERSIDAD NACIONAL DE SAN LUIS  
*E-mail address:* `victor.schvollner.tag@gmail.com`

INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UFRGS, PORTO ALEGRE, BRAZIL AND DEPARTMENT OF MATHEMATICS AND APPLICATIONS, UNIVERSITY OF NAPLES FEDERICO II, ITALY  
*E-mail address:* `trevisan@mat.ufrgs.br`