

“On-the-fly” calculation of the Vibrational Sum-frequency Generation Spectrum at the Air-water Interface

Deepak Ojha

*Dynamics of Condensed Matter and Center for Sustainable Systems Design,
Chair of Theoretical Chemistry, Department of Chemistry,
Paderborn University, Warburger Str. 100, 33098 Paderborn, Germany*

Thomas D. Kühne*

*Dynamics of Condensed Matter and Center for Sustainable Systems Design,
Chair of Theoretical Chemistry, Department of Chemistry,
Paderborn University, Warburger Str. 100, 33098 Paderborn, Germany and
Paderborn Center for Parallel Computing and Institute for Lightweight Design,
Paderborn University, Warburger Str. 100, D-33098 Paderborn, Germany*

(Dated: June 26, 2020)

In the present work, we provide an electronic structure based method for the “on-the-fly” determination of vibrational sum frequency generation (v-SFG) spectra. The predictive power of this scheme is demonstrated at the air-water interface. While the instantaneous fluctuations in dipole moment are obtained using the maximally localized Wannier functions, the fluctuations in polarizability are approximated to be proportional to the second moment of Wannier functions. The spectrum henceforth obtained captures the signatures of hydrogen bond stretching, bending, as well as low-frequency librational modes.

PACS numbers: 31.15.-p, 31.15.Ew, 71.15.-m, 71.15.Pd

I. INTRODUCTION

Vibrational spectroscopy provides microscopic fingerprints of the structure and dynamics at the molecular level in condensed phase systems.^{1–3} However, theoretical interpretation and peak characterization of vibrational spectra predominantly relies on molecular dynamics simulations.^{4–9} Nevertheless, the success of simulations also depends largely on the forcefield employed to describe the interatomic interactions. In this regard, ab-initio molecular dynamics (AIMD) has proven to be extremely useful as the interatomic forces are obtained from accurate electronic structure calculations.^{10,11} For periodic systems, the overall electronic state within the AIMD framework is generally expressed in the terms of Bloch orbitals

$$\Psi(r, k) = e^{(ik \cdot r)} u_i(r, k), \quad (1a)$$

where

$$u_i(r, k) = u_i(r + R, k), \quad (1b)$$

with $\Psi(r, k)$ being the electronic wavefunction, $u_i(r, k)$ the Bloch function and R a translational lattice parameter.¹² An alternative representation, which is more suited for chemical problems, is provided by so-called maximally localized Wannier functions (MLWFs), i.e. $w_n(r - R)$ that are obtained by a unitary transformation of the Bloch orbitals.^{13,14} The construction of this

Wannier representation enables to split the continuously varying total electronic density into contributions originating from localized fragments of the system. Mathematically, MLWFs are expressed as

$$w_n(r - R) = \frac{V}{2\pi^3} \int_{BZ} d\mathbf{k} e^{-i\mathbf{k} \cdot R} \sum_{m=1}^J U_{mn}^{(k)} \psi_{m\mathbf{k}}(r), \quad (2)$$

where R is the lattice vector of the unit cell and V is the real-space primitive cell volume. The $J \times J$ matrix $U_{mn}^{(k)}$ is the unitary transformation matrix and $\psi_{m\mathbf{k}}(\mathbf{r})$ are the eigenstates of the system computed by density function theory (DFT). The corresponding MLWFs are then obtained by the unitary transformation $U_{mn}^{(k)}$ that minimizes the spread functional

$$S = \sum_n S_n = \sum_n (\langle w_n | r^2 | w_n \rangle - \langle w_n | r | w_n \rangle^2). \quad (3)$$

Therein, $\langle r^2 \rangle$ is the second moment, whereas $\langle r \rangle^2$ is the squared first moment of the Wannier centers. This unitary transformation based localization can be readily implemented on the position operator \hat{r} within the Wannier representation to obtain localized orbitals for a given periodic system of arbitrary symmetry.^{15–17} As a result, the scheme can be used to compute the electronic contributions to the polarization of a system. Moreover, it also allows to calculate instantaneous fluctuations in the molecular dipole moment and within the linear-response regime, obtain the linear as well as nonlinear infrared spectrum using time-correlation function formalism.^{18–24} In this regard, Raman and higher nonlinear analogs like v-SFG, 2D-vSFG and 2D-Raman can also be computed

* td kuehne@mail.upb.de

by applying a constant periodic electric field using the Berry phase formalism²⁵⁻²⁷, or by calculating the polarizability tensor A

$$A_{ij} = -\frac{\delta M_i(\mathbf{E})}{\delta E_j}, \quad (4)$$

where M is the total dipole moment and E is an externally applied electric field. This scheme of computing the polarizability tensor has been utilized to obtain isotropic Raman spectrum by means of density functional perturbation theory.²⁸⁻³¹

In this paper, we present a novel computational method to obtain the v-SFG spectrum of the air-water interface. This anisotropic Wannier Polarizability (WP) method is based on a technique of computing the fluctuations within the dipole moment and polarizability "on-the-fly" during an AIMD simulation without any additional computational cost.³² For that purpose, the fluctuations in the dipole moment are obtained using the Wannier centers, whereas the components of the polarizability tensor are approximated using the second moment of the Wannier centers. However, it is noteworthy to mention that several other computational studies have obtained the vSFG spectrum using empirical maps³³⁻⁴⁰, velocity correlations⁴¹⁻⁴⁴, as well as directly from AIMD simulations⁴⁵⁻⁴⁸.

The rest of the paper is structured as follows. In section 2, we outline our anisotropic WP method, whereas in section 3 the computational details to compute the v-SFG of the air-water interface are described. The corresponding numerical results are shown in section 4, before concluding remarks are provided in section 5.

II. ANISOTROPIC WANNIER POLARIZABILITY METHOD

The original isotropic WP method has been implemented to compute the isotropic Raman spectrum of isolated gas-phase molecules, as well as aqueous solutions.^{32,49} The underlying principle of the method is that the polarization induced by an externally applied perturbation is directly proportional to the molecular volume of the system.^{63,64} Since, Wannier centers provide a picture, where the total electronic density is partitioned into the localized electronic densities of different fragments of the system, the fluctuations in the electronic polarizability can be connected to the fluctuations of the volume of the Wannier centers instead of the overall molecular volume. As a result, the net isotropic polarizability can be expressed as

$$\bar{A} = \frac{1}{3} \sum_{i=1}^{N_{WF}} A_i = \frac{\beta}{3} \sum_i^{N_{WF}} S_i^3, \quad (5)$$

where S_i is the spread of the i^{th} Wannier center, N_{WF} is the number of MLWFs and β is a proportionality

constant. The isotropic Raman spectrum is then obtained as the Fourier transform of the polarizability time-correlation function.

On similar lines, the v-SFG spectrum of a non-centrosymmetric system is given as

$$\chi_{abc}^2(\omega) = \int_0^\infty dt e^{i\omega t} \langle \dot{A}_{ab}(0) \cdot \dot{M}_c(t) \rangle, \quad (6)$$

where χ_{abc}^2 is the second order susceptibility, whereas A_{ab} is the ab^{th} component of the polarizability tensor and M_c is c^{th} component of the dipole moment.^{33,36,45} In contrast to Raman spectroscopy, the computation of v-SFG spectra requires the diagonal elements of the polarizability tensor. In this regard, we note that the second moment, i.e. $\langle w_n | r^2 | w_n \rangle$ and the polarizability are tensors of same size. Accordingly, we have approximated that the component specific fluctuations in the polarizability are proportional to the second moment of the Wannier centers, i.e.

$$A_{ab} \propto \langle w_n | r_{ab}^2 | w_n \rangle. \quad (7)$$

The strength of the anisotropic WP method is that for each set of an electron pair, we have a unique Wannier center and its corresponding moments. As a result, the method can be used to specifically study the contributions from the different fragments of the system. Moreover, it is also computationally less expensive as the polarizability is determined "on-the-fly" from the second moments of Wannier centers, which is in contrast with existing approaches, where the polarizability is obtained by numerical differentiation of the total dipole moment with respect to an externally applied electric field. This is to say that a simple minimization of the spread functional provides the Wannier centers and their corresponding moments, which are used to obtain the dipole and the polarizability, respectively. Thus, a single AIMD-based Wannier center calculation is sufficient to obtain the dipole moment, as well as the polarizability.

III. COMPUTATIONAL DETAILS

Ab initio molecular dynamics simulations were performed by using the method of Car and Parrinello,^{10,50} as implemented in the CPMD code⁵¹. Simulations of the air-water interface comprising of 80 H_2O molecules were performed at 300 K in a cubic box of edge length 12.43 Å corresponding to the density at ambient conditions.⁵² The air-water interface was generated by increasing the edge length of the box to 37.2 Å in the z-direction. The Kohn-Sham formulation of density functional theory was applied to represent the electronic structure of the system within a plane wave basis set.¹¹ In order to represent the core-shell electrons, Vanderbilt ultra-soft pseudopotentials were used and the plane wave expansion of Kohn-Sham orbitals was truncated at a kinetic energy cutoff of 25 Ry.⁵³ The electronic orbitals were assigned a fictitious




Figure 1. Distribution of molecular dipole moment (μ) of water molecules at ambient conditions, as computed using the maximally localized Wannier centers.

mass of 400 a.u. and equations of motion were integrated with a time step of 4 a.u.

In the present work, we have used the dispersion corrected BYLP-D exchange and correlation (XC) functional,^{54–56} since previous AIMD studies have shown that inclusion of London dispersion interactions not only improves the structure, but also predicts the dynamics, spectroscopy and phase diagram of "*ab initio*" water and aqueous solutions in better agreement with experiment.^{48,57–60} The initial configuration was generated using classical molecular dynamics simulations. Subsequently, the production run was carried out in the canonical NVT ensemble using Nose-Hoover thermostats for 50 ps.

The identification of interfacial water molecules at the air-water system was conducted using the algorithm for the identification of truly interfacial molecules ITIM.^{61,62} This scheme uses a probe sphere to detect the molecules at the surface. The radius of the probe sphere was set to 2 Å which has been proven to be a good value for water.⁶² A cutoff-based cluster search was also performed using 3.5 Å as a cut-off, which corresponds to the first minimum of the O···O radial distribution function in liquid water.

IV. RESULTS AND DISCUSSION

To demonstrate the predictive power of the present anisotropic WP method, we have computed the v-SFG of at the air-water interface. For the sake of simplicity, we have assumed that the contributions originating from Wannier centers, which are associated with the lone pair of electrons, to the overall polarizability can be safely neglected. The spectral dynamics is predominantly governed by the dynamical evolution of the Wannier centers corresponding to the bonded electron




Figure 2. The dipole-polarizability cross-correlation function, as obtained by the present anisotropic WP method.

pairs. The average molecular dipole moment of the water molecules obtained using the Wannier centers, whose distribution is shown in Fig. 1, was found to be 2.46 Debye. The dipole-polarizability cross-correlation function and the v-SFG spectrum computed based on the fluctuations within the dipole moments obtained by using the Wannier centers and polarizabilities by means of the second moment are shown in Figs. 2 and 3, respectively. We find that the v-SFG spectrum obeys char-




Figure 3. The v-SFG spectrum of interfacial water molecules computed by the present anisotropic WP method.

acteristic peaks corresponding to librational, bending, OH stretching, as well as free OH modes. Since there are various previous experimental and simulation based studies analysing the stretching, bending and librational modes within the v-SFG spectrum at the air-water interface, we only briefly highlight our findings in light of the existing literature. First, we will focus on the spectral region of 3000–3800 cm^{-1} , which is predominantly attributed to OH stretching modes. More precisely, earlier simulation studies have reported a broad negative peak between 3000 to 3600 cm^{-1} and a sharp positive peak around 3700 cm^{-1} .^{33,34,36,37,45} Using our anisotropic WP

method, we also find a broad negative peak at 2900-3500 cm^{-1} and sharp positive peak around $\sim 3600 cm^{-1}$. The former broad negative peak contribution originates from hydrogen-bonded water molecules with the overall dipole aligned towards the bulk, whereas the latter sharp positive peak is connected with the free and dangling OH modes of the interfacial water molecules. The observed red-shift within the peak positions can be most likely attributed to the choice of XC functional and employed pseudopotentials in the present study. Earlier experimental and simulation studies of the bending mode have reported a broad negative peak around 1650 cm^{-1} and a positive shoulder around 1750 cm^{-1} .^{43,46} Here, using the anisotropic WP method, we also observe a broad negative peak between 1400 and 1650 cm^{-1} , which is governed by the free and dangling OH modes. However, at variance to these earlier studies,^{43,46} we can not confirm any positive shoulder in our calculations. Finally, we observe a negative peak at around 450-650 cm^{-1} that is governed by the librational motion of water molecules. Apart from a consistent red-shift within the peak positions, our results are in good agreement with earlier results that have also reported a negative peak in the region of 700-800 cm^{-1} .⁴²

V. SUMMARY

To summarize, we have proposed a computationally efficient “on-the-fly” method to determine the v-SFG spectrum for interfacial systems. This anisotropic WP

method utilizes the second moment of the Wannier centers to estimate the polarizability fluctuations. The major strength of this method is that it captures the spectral signatures of the system for the collective, as well as highly localized modes. Furthermore, it can be directly applied to spectral decomposition by computing fragment-specific contributions from the Wannier centers and their second moment to assist the interpretation of the experimental measurements. Moreover, the algorithm employed here can be easily extended to other spectroscopic techniques like two-dimensional v-SFG,⁶⁵ time-dependent v-SFG,⁶⁶ 2D-Raman-Thz,⁶⁷ pump-probe Thz⁶⁸ and 2D-Raman,⁶⁹ to name just a few. From the application perspective, interfacial reactivity, “on-water” catalysis, and other interfacial chemical processes can also be studied using our anisotropic WP-based method. Nevertheless, for greater agreement with the experiment, it would be important to better understand the role of simulation protocols and the approximations made, which we propose as an extensions for future works.

ACKNOWLEDGMENTS

The authors would like to thank the Paderborn Center for Parallel Computing (PC²) for the generous allocation of computing time on FPGA-based supercomputer “Noctua”. This project has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No 716142).

-
- [1] S. Nihonyanagi, S. Yamaguchi, T. Tahara, *Chem. Rev.* **16**, 10665 (2017).
 - [2] Q. Du, R. Superfine, E. Freysz, and Y. R. Shen, *Phys. Rev. Lett.* **70**, 2313 (1993).
 - [3] Y. R. Shen and V. Ostroverkhov, *Chem. Rev.* **106**, 1140 (2006).
 - [4] H. J. Bakker and J. L. Skinner, *Chem. Rev.* **110**, 1498 (2010).
 - [5] G. L. Richmond, *Chem. Rev.* **102**, 2693 (2002).
 - [6] G. L. Richmond, *Annu. Rev. Phys. Chem.* **52**, 357 (2001).
 - [7] F. Perakis, L. D. Marco, A. Shalit, F. Tang, Z. R. Kann, T. D. Kühne, R. Torre, M. Bonn, and Y. Nagata *Chem. Rev.*, **116**, 7590 (2016).
 - [8] D. Ojha, K. Karhan, and T. D. Kühne, *Sci. Rep.* **8**, 16888 (2018).
 - [9] C. John, T. Spura, S. Habershon, and T. D. Kühne, *Phys. Rev. E* **93**, 043305 (2016).
 - [10] D. Marx and J. Hutter, *Ab Initio Molecular Dynamics: Basic Theory and Advanced Methods*, Cambridge University Press, Cambridge, 2009.
 - [11] T. D. Kühne, *WIREs Comput. Mol. Sci.* **4**, 391 (2014).
 - [12] I. Souza, N. Marzari and D. Vanderbilt, *Phys. Rev. B* **65**, 035109 (2001).
 - [13] N. Marzari and D. Vanderbilt, *Phys. Rev. B* **56**, 12847 (1997).
 - [14] N. Marzari, A. A. Mostofi, J. R. Yates, I. Souza, D. Vanderbilt, *Rev. Mod. Phys.* **84**, 1419 (2012).
 - [15] P. L. Silvestrelli, *Phys. Rev. B* **59**, 9703 (1999).
 - [16] G. Berghold, C. J. Mundy, A. H. Romero, J. Hutter, and M. Parrinello, *Phys. Rev. B* **61**, 10040 (2000).
 - [17] R. Resta, *Phys. Rev. Lett.* **80**, 1800 (1998).
 - [18] W. Chen, M. Sharma, R. Resta, G. Galli, and R. Car *Phys. Rev. B* **77**, 245114 (2008).
 - [19] Marie-Pierre Gaigeot and M. Sprik, *J. Phys. Chem. B* **107**, 10344 (2003).
 - [20] M. Heyden, J. Sun, S. Funkner, G. Mathias, H. Forbert, M. Havenith and D. Marx, *Proc. Nat. Acad. Sci.* **107**, 12068 (2010).
 - [21] C. Zhang, R. Z. Khaliullin, D. Bovi, L. Guidoni, and T. D. Kühne, *J. Phys. Chem. Lett.* **4**, 3245 (2013).
 - [22] D. Ojha, A. Henao, and T. D. Kühne, *J. Chem. Phys.* **148**, 102328 (2018).
 - [23] D. Ojha and A. Chandra, *Phys. Chem. Chem. Phys.*, **21**, 6485 (2019).
 - [24] D. Ojha and A. Chandra, *Chem. Phys. Lett.*, **751**, 137493 (2020).
 - [25] R. Resta, *Rev. Mod. Phys.* **66**, 899 (1994).

- [26] R. D. King-Smith and D. Vanderbilt, *Phys. Rev. B* **47**, 1651 (1993).
- [27] P. Umari and A. Pasquarello *Phys. Rev. Lett.* **89**, 157602 (2002).
- [28] S. Baroni, S. De Gironcoli, A. Dal Corso, and P. Gianozzi, *Rev. Mod. Phys.* **73**, 515 (2001).
- [29] A. Putrino and M. Parrinello, *Phys. Rev. Lett.* **88**, 176401 (2002).
- [30] S. Luber, M. Iannuzzi, and J. Hutter, *J. Chem. Phys.* **141**, 094503 (2014).
- [31] T. D. Kühne et al., *J. Chem. Phys.* **152**, 194103 (2020).
- [32] P. Partovi Azar and T. D. Kühne, *J. Comput. Chem.* **36**, 2188 (2015).
- [33] A. Morita and J. T. Hynes, *Chem. Phys.* **258**, 371 (2000).
- [34] A. Morita and J. T. Hynes, *J. Phys. Chem. B* **106**, 673 (2002).
- [35] T. Ishiyama, T. Imamura, and A. Morita *Chem. Rev.* **114**, 8447 (2014).
- [36] P. A. Pieniazek, C. J. Tainter, and J. L. Skinner, *J. Amer. Chem. Soc.* **133**, 10360 (2011).
- [37] Y. Li and J. L. Skinner, *J. Chem. Phys.* **2016**, **145**, 124509.
- [38] Y. Li and J. L. Skinner, *J. Chem. Phys.* **2016**, **145**, 031103.
- [39] P. A. Pieniazek, C. J. Tainter, and J. L. Skinner, *J. Chem. Phys.* **135**, 044701 (2011).
- [40] Y. Li, M. Gruenbaum and J. L. Skinner, *Proc. Nat. Acad. Sci.* **110**, 1992 (2013).
- [41] T. Ohta, K. Usui, T. Hasegawa, M. Bonn and Y. Nagata, *J. Chem. Phys.* **143**, 124702 (2015).
- [42] R. Khatib, T. Hasegawa, M. Sulpizi, E. H. G. Backus, M. Bonn, and Y. Nagata, *J. Phys. Chem. C* **120**, 18665 (2016).
- [43] Y. Nagata, C-S Hsieh, T. Hasegawa, J. Voll, E. H. G. Backus, and M. Bonn, *J. Phys. Chem. Lett.* **11**, 1872 (2013).
- [44] N. K. Kaliannan, A. Henao, H. Wiebeler, F. Zysk, T. Ohto, Y. Nagata, and T. D. Kühne, *Mol. Phys.* **118**, 1620358 (2020).
- [45] M. Sulpizi, M. Salanne, M. Sprik, and M. Pierre Gaiageot *J. Phys. Chem. Lett.* **4**, 83 (2013).
- [46] R. Khatib and M. Sulpizi *J. Phys. Chem. Lett.* **8**, 1310 (2017).
- [47] C. Liang, J. Jeon, and M. Cho, *J. Phys. Chem. Lett.* **10**, 1153 (2019).
- [48] T. Ohto, M. Dodia, J. Xu, S. Imoto, F. Tang, F. Zysk, T. D. Kühne, Y. Shigeta, M. Bonn, X. Wu, and Y. Nagata, *J. Phys. Chem. Lett.* **10**, 4914 (2019).
- [49] P. Partovi-Azar, T. D. Kühne, and P. Kaghazchi, *Phys. Chem. Chem. Phys.* **17**, 22009 (2015).
- [50] R. Car and M. Parrinello, *Phys. Rev. Lett.* **1985**, **55**, 2471–2474.
- [51] J. Hutter, A. Alavi, T. Deutsch, M. Bernasconi, S. Goedecker, D. Marx, M. Tuckerman, and M. Parrinello, CPMD Program, MPI für Festkörperforschung and IBM Zurich Research Laboratory. See www.cpm.org.
- [52] D. R. Lide, *CRC Handbook of Chemistry and Physics*, *84th Edition*. Taylor & Francis, 2003.
- [53] D. Vanderbilt, *Phys. Rev. B*, **41**, 7892 (1990).
- [54] A. D. Becke, *Phys. Rev. A* **1988**, **38**, 3098–3100.
- [55] C. Lee, W. Yang, and R. G. Parr, *Phys. Rev. B* **1988**, **37**, 785–789.
- [56] S. Grimme, *J. Comput. Chem.* **2006**, **27**, 1787–1799.
- [57] M. J. McGrath, I. F. Kuo, and J. I. Siepmann, *Phys. Chem. Chem. Phys.* **2011**, **13**, 19943–19950.
- [58] T. D. Kühne, T. A. Pascal, E. Kaxiras, and Y. Jung, *J. Phys. Chem. Lett.* **2**, 105 (2011).
- [59] M. J. McGrath, I. F. Kuo, J. N. Ghogomu, C. J. Mundy, and J. I. Siepmann, *J. Phys. Chem. B* **2011**, **115**, 11688–11692.
- [60] Y. Nagata, T. Ohto, M. Bonn, and T. D. Kühne, *J. Chem. Phys.* **144**, 204705 (2016).
- [61] L. B. Pártay, G. Hantal, P. Jedlovsky, Á. Vincze, and G. Horvai, *J. Comp. Chem.* **29**, 945 (2008).
- [62] M. Sega, S. S. Kantorovich, P. Jedlovsky, and M. Jorge, *J. Chem. Phys.* **138**, 044110 (2013).
- [63] K. E. Laidig and R. F. W. Bader *J. Chem. Phys.* **93**, 7213 (1990).
- [64] R. F. W. Bader, *Atoms in Molecules: A Quantum Theory*, Oxford University Press:Oxford (1990)
- [65] J. Bredenbeck, A. Ghosh, H-K Nienhuys, and M. Bonn, *Acc. Chem. Res.* **42**, 1332 (2009).
- [66] D. Ojha, N. K. Kaliannan, and T. D. Kühne, *Commun. Chem.* **2**, 116 (2019).
- [67] J. Savolainen, S. Ahmed, and P. Hamm, *Proc. Nat. Acad. Sci.* **110**, 20402 (2013).
- [68] H. Elgabarty, T. Kampfrath, D. J. Bonthuis, V. Balos, N. K. Kaliannan, P. Loche, R. R. Netz, M. Wolf, T. D. Kühne, and M. Sajadi, *Sci. Adv.* **6**, eaay7074 (2020).
- [69] Y. Tanimura and S. Mukamel, *J. Chem. Phys.*, **99** 9496 (1993).