A NEW CLASS OF MINIMAL ASYMPTOTIC BASES

MELVYN B. NATHANSON

ABSTRACT. A set A of nonnegative integers is an asymptotic basis of order h if every sufficiently large integer can be represented as the sum of h not necessarily distinct elements of A. An asymptotic basis A is minimal if removing any element of A destroys every representation of infinitely many integers. In this paper, a new class of minimal asymptotic bases is constructed.

1. G-adic asymptotic bases

Let $N_0 = \{0, 1, 2, 3, \ldots\}$ be the set of nonnegative integers and let h be a positive integer. Let A, A_1, \ldots, A_h be subsets of \mathbf{N}_0 . We define the *sumset*

 $A_1 + \cdots + A_h = \{a_1 + \cdots + a_h : a_i \in A_i \text{ for all } i = 1, 2, \ldots, h\}$

and the h-fold sumset

$$
hA = \underbrace{A + \dots + A}_{h \text{ summands}} = \{a_1 + \dots + a_h : a_i \in A \text{ for all } i = 1, 2, \dots, h\}.
$$

The set A is a basis of order h if every nonnegative integer can be represented as the sum of h not necessarily distinct elements of A, that is, if $hA = N_0$. The set A is an *asymptotic basis of order h* if hA contains all sufficiently large integers. An asymptotic basis of order h is minimal if no proper subset of A is an asymptotic basis of order h. Thus, if A is a minimal asymptotic basis of order h, then, for all $a \in A$, there are infinitely many integers n such that removing a from A destroys every representation of n as a sum of h elements of A .

Minimal asymptotic bases are extremal objects in additive number theory, and are related to the conjecture of Erdős and Turán $[4]$ that the representation function of an asymptotic basis of order h must be unbounded.

At this time there are few explicit constructions of minimal asymptotic bases. Nathanson [\[11,](#page-9-1) [12\]](#page-9-2) used a 2-adic construction to produce the first examples of minimal asymptotic bases. This method was extended to g-adically defined sets by Chen [\[2\]](#page-9-3), Chen and Chen [\[1\]](#page-9-4), Chen and Tang [\[3\]](#page-9-5), Jia [\[5\]](#page-9-6), Jia and Nathanson [\[6\]](#page-9-7), Lee $[7]$, Li and Li $[8]$, Ling and Tang $[9, 10]$ $[9, 10]$, Sun $[15, 16]$ $[15, 16]$, and Sun and Tao $[17]$. This paper constructs a new class of minimal asymptotic bases.

An interval of integers of length t is a set of t consecutive integers. For $u, v \in \mathbb{N}_0$ with $u \leq v$, the set

 $[u, v] = \{x \in \mathbb{N}_0 : u \leq x \leq v\}$

is an interval of integers of length $v - u + 1$. Thus, $[1, h] = \{1, 2, ..., h\}$.

Date: May 6, 2022.

²⁰¹⁰ Mathematics Subject Classification. 11B13, 11B05, 11B34, 11B75.

Key words and phrases. Additive number theory, additive basis, asymptotic basis, minimal asymptotic basis, Erdős-Turán conjecture.

Supported in part by a grant from the PSC-CUNY Research Awards Program.

A G-adic sequence is a strictly increasing sequence of positive integers $\mathcal{G} = (g_i)_{i=0}^{\infty}$ such that $g_0 = 1$ and g_{i-1} divides g_i for all $i \geq 1$. Let $(d_i)_{i=1}^{\infty}$ be the sequence of positive integers defined by

$$
d_i = \frac{g_i}{g_{i-1}}.
$$

For all $i \geq 1$, we have

$$
(1) \t\t d_i \ge 2
$$

and

$$
(2) \t\t g_i = d_1 d_2 \cdots d_i.
$$

For all i and j with $0 \leq i \leq j$, we have

(3)
$$
\frac{g_{i+j}}{g_i} = d_{i+1}d_{i+2}\cdots d_{i+j}.
$$

Every positive integer n has a unique $\mathcal{G}\text{-}adic representation$

$$
n = \sum_{j=0}^{\infty} x_j g_j
$$

where

$$
x_j \in [0, d_{j+1} - 1]
$$

for all $j \in \mathbb{N}_0$ and $x_j = 0$ for all sufficiently large j (Nathanson [\[13,](#page-9-15) [14\]](#page-9-16)). Equivalently, for every positive integer n, there is a unique nonempty finite set $F \subseteq \mathbb{N}_0$ and a unique set $\{x_j : j \in F\}$ such that

$$
(4) \t\t n = \sum_{j \in F} x_j g_j
$$

where $x_j \in [1, d_{j+1} - 1]$ for all $j \in F$.

For every integer $g \geq 2$, the usual g-adic representation uses the G-adic sequence $G = (g^i)_{i=0}^{\infty}$ with quotients $d_i = g$ for all $i \geq 1$.

Lemma 1. Let $\mathcal{G} = (g_i)_{i=0}^{\infty}$ be a \mathcal{G} -adic system. If $n = \sum_{j \in F} x_j g_j$ is a positive integer with $x_j \in [1, d_{j+1} - 1]$ for all $j \in F$, then

$$
g_M\leq n < g_{M+1}
$$

if and only if

$$
\max(F) = M.
$$

Proof. If $\max(F) = M$, then $F \subseteq [0, M]$ and

$$
g_M \le x_M g_M \le n \le \sum_{j \in F} (d_{j+1} - 1)g_j
$$

$$
\le \sum_{j=0}^M (d_{j+1} - 1)g_j = \sum_{j=0}^M (g_{j+1} - g_j)
$$

$$
= g_{M+1} - 1 < g_{M+1}.
$$

Conversely, if $g_M \le n < g_{M+1}$ and $\max(F) = M'$, then the inequality

$$
g_{M'} \le x_{M'} g_{M'} \le n < g_{M+1}
$$

implies $M' \leq M$. If $M' \leq M - 1$, then

$$
n \le \sum_{j=0}^{M'} (d_{j+1} - 1)g_j < g_{M'+1} \le g_M
$$

which is absurd. Therefore, $g_M \le n < g_{M+1}$ implies $\max(F) = M$. This completes the proof. \Box

Let W be a nonempty set of nonnegative integers, and let $\mathcal{F}^*(W)$ be the set of all nonempty finite subsets of W. Let $\mathcal{G} = (g_i)_{i=0}^{\infty}$ be a \mathcal{G} -adic sequence. We define the set of positive integers

$$
A_{\mathcal{G}}(W) = \left\{ \sum_{j \in F} x_j g_j : F \in \mathcal{F}^*(W) \text{ and } x_j \in [1, d_{j+1} - 1] \right\}.
$$

Note that $0 \notin A_{\mathcal{G}}(W)$ because $\emptyset \notin \mathcal{F}^*(W)$.

Let $h \geq 2$. A partition of N_0 is a sequence $W = (W_i)_{i=0}^{h-1}$ of nonempty pairwise disjoint sets such that

$$
\mathbf{N}_0 = W_0 \cup W_1 \cup \cdots \cup W_{h-1}.
$$

Theorem 1. Let $h \geq 2$ and let $W = (W_i)_{i=0}^{h-1}$ be a partition of \mathbf{N}_0 . Let $\mathcal{G} = (g_i)_{i=0}^{\infty}$ be a G-adic sequence. The set

$$
A_{\mathcal{G}}(\mathcal{W}) = \bigcup_{i=0}^{h-1} A_{\mathcal{G}}(W_i)
$$

is an asymptotic basis of order h with h-fold sumset

$$
hA_{\mathcal{G}}(\mathcal{W}) = \{ n \in \mathbf{N}_0 : n \ge h \}.
$$

Proof. The smallest integer in the set $A_{\mathcal{G}}(\mathcal{W})$ is $1 = 1 \cdot g_0$. It follows that $h \in$ $hA_G(\mathcal{W})$ but $[0, h-1] \cap hA_G(\mathcal{W}) = \emptyset$.

Every positive integer *n* has a unique G-adic representation $n = \sum_{j \in F} x_j g_j$, where F is a nonempty finite set of nonnegative integers and $x_j \in [0, d_{j+1} - 1]$. For $i \in [0, h - 1]$, let $F_i = F \cap W_i$

and

$$
\overline{a}
$$

$$
n_i = \sum_{j \in F_i} x_j g_j.
$$

If $F_i = \emptyset$, then $n_i = 0 \notin A_G(\mathcal{W})$. If $F_i \neq \emptyset$, then

$$
n_i = \sum_{j \in F_i} x_j g_j \in A_{\mathcal{G}}(W_i) \subseteq A_{\mathcal{G}}(\mathcal{W}).
$$

Let

$$
L = \{i \in [0, h-1] : F_i \neq \emptyset\} = \{i \in [0, h-1] : n_i \ge 1\}
$$

and

$$
|L|=\ell_0.
$$

We have $\ell_0 \in [1, h]$ and

$$
n = \sum_{i \in L} n_i \in \ell_0 A_{\mathcal{G}}(\mathcal{W}).
$$

Let ℓ be the largest integer such that $\ell \leq h$ and $n \in \ell A_G(\mathcal{W})$. We must prove that $\ell = h$.

If $n \in \ell A_{\mathcal{G}}(\mathcal{W})$, then there exist integers $n_1, \ldots, n_\ell \in A_{\mathcal{G}}(\mathcal{W})$ such that

$$
n = n_1 + \dots + n_{k-1} + n_k + n_{k+1} + \dots + n_\ell.
$$

For each $i \in [1, \ell]$ there is an integer $s_i \in [0, h-1]$ and a set $F_{s_i} \in \mathcal{F}^*(W_{s_i})$ such that n_i has the G-adic representation

$$
n_i = \sum_{j \in F_{s_i}} x_j g_j \in A_{\mathcal{G}}(W_{s_i}).
$$

Suppose that $\ell < h$. If $|F_{s_k}| \geq 2$ for some $k \in [1, \ell]$, then there are nonempty sets F'_{s_i} and F''_{s_i} such that

$$
F_{s_k} = F'_{s_k} \cup F''_{s_k} \quad \text{and} \quad F'_{s_k} \cap F''_{s_k} = \emptyset.
$$

The integers

$$
n'_k = \sum_{j \in F'_{s_k}} x_j g_j \in A_{\mathcal{G}}(W_{s_k}) \quad \text{and} \quad n''_k = \sum_{j \in F''_{s_k}} x_j g_j \in A_{\mathcal{G}}(W_{s_k})
$$

satisfy

$$
n_k = n'_k + n''_k
$$

and so

$$
n = n_1 + \cdots + n_{k-1} + n'_k + n''_k + n_{k+1} + \cdots + n_\ell \in (\ell+1)A_{\mathcal{G}}(\mathcal{W}).
$$

This contradicts the maximality of ℓ , and so $|F_{s_i}| = 1$ for all $i \in [1, \ell]$ and

$$
n_i = x_{j_i} g_{j_i}
$$

for some $g_{j_i} \in W_{s_i}$ and $x_{j_i} \in [0, d_{j_i+1}-1]$. If $x_{j_k} \geq 2$ for some $k \in [1, \ell],$ then

$$
g_{j_k} \in A_{\mathcal{G}}(W_{s_k})
$$
 and $(x_{j_k} - 1)g_{j_k} \in A_{\mathcal{G}}(W_{s_k})$

and

$$
n_k = g_{j_k} + (x_{j_k} - 1)g_{j_k}.
$$

It follows that

$$
n = n_1 + \cdots + n_{k-1} + g_{j_k} + (x_{j_k} - 1)g_{j_k} + n_{k+1} + \cdots + n_\ell \in (\ell+1)A_{\mathcal{G}}(\mathcal{W}),
$$

which again contradicts the maximality of ℓ . Therefore, $x_{j_i} = 1$ for all $i \in [1, \ell],$ and

$$
n = g_{j_1} + \cdots + g_{j_k} + \cdots + g_{j_\ell}.
$$

If $j_k \geq 1$ for some $k \in [1, \ell]$, then

$$
g_{j_k} = g_{j_k - 1} + (d_{j_k} - 1)g_{j_k - 1}
$$

and

$$
n = g_{j_1} + \cdots + g_{j_k-1} + (d_{j_k} - 1)g_{j_k-1} + \cdots + g_{j_\ell} \in (\ell+1)A_{\mathcal{G}}(\mathcal{W}),
$$

which also contradicts the maximality of ℓ . Therefore, $j_k = 0$ and $g_{j_k} = g_0 = 1$ for all $k \in [1, \ell]$, and so

$$
n = \underbrace{g_0 + \dots + g_0}_{\ell \text{ summands}} = \underbrace{1 + \dots + 1 + \dots + 1}_{\ell \text{ summands}} = \ell \leq h - 1,
$$

which is absurd. This completes the proof. \Box

Theorem 2. Let $h \geq 2$ and let $W = (W_i)_{i=0}^{h-1}$ be a partition of \mathbf{N}_0 . The set

$$
A = \{0\} \cup A_{\mathcal{G}}(\mathcal{W})
$$

is a basis of order h but not a minimal asymptotic basis of order h.

Proof. Because $0 \in A$ and $1 \in A_{\mathcal{G}}(\mathcal{W}) \subseteq A$, we have

$$
\ell = (h - \ell) \cdot 0 + \ell \cdot 1 \in hA
$$

for all $\ell \in [0, h-1]$. By Theorem [1,](#page-2-0)

$$
\{n \in \mathbf{N}_0 : n \ge h\} = hA_{\mathcal{G}}(\mathcal{W}) \subseteq hA
$$

and so A is a basis of order h.

The set A is not a minimal asymptotic basis of order h because $0 \in A$ and the removal of 0 from A gives the set $A \setminus \{0\} = A_{\mathcal{G}}(\mathcal{W})$, which is still an asymptotic basis of order h. This completes the proof. \Box

2. Minimal asymptotic bases

The following lemma generalizes a result of Jia [\[5\]](#page-9-6).

Lemma 2. Let $\mathcal{G} = (g_i)_{i=0}^{\infty}$ be a \mathcal{G} -adic sequence. Let $(u_i)_{i=1}^p$ be a strictly increasing finite sequence of nonnegative integers, and let $(v_j)_{j=1}^q$ be a finite sequence of not necessarily distinct nonnegative integers. Let

$$
x_i \in [1, d_{u_i+1} - 1]
$$

for all $i \in [1, p]$ and

$$
y_j \in [1, d_{v_j+1}-1]
$$

for all $j \in [1, q]$. If

(5)
$$
n = \sum_{i=1}^{p} x_i g_{u_i} = \sum_{j=1}^{q} y_j g_{v_j}
$$

then

(6)
$$
\sum_{u_i \leq u_k} x_i g_{u_i} \leq \sum_{v_j \leq u_k} y_j g_{v_j}
$$

for all $k \in [1, p]$.

Proof. Because the sequence $(u_i)_{i=1}^p$ is strictly increasing, Lemma [1](#page-1-0) implies

$$
\sum_{u_i \le u_k} x_i g_{u_i} = \sum_{i=1}^k x_i g_{u_i} < g_{u_{k+1}}
$$

for all $k \in [1, p]$. Choosing $k = p$ gives

$$
n = \sum_{i=1}^{p} x_i g_{u_i} < g_{u_{p+1}}.
$$

Relation [\(5\)](#page-4-0) implies

$$
g_{v_j} \leq n < g_{u_p+1}
$$

an so $v_j \le u_p$ for all $j \in [1, q]$. This implies [\(6\)](#page-4-1) for $k = p$.

In the sequence $G = (g_i)_{i=0}^{\infty}$, the integer g_i divides g_j for all $i \leq j$. Let $k \in$ [1, p – 1]. Because the sequence $(u_i)_{i=1}^p$ is strictly increasing, for all $i \in [k+1,p]$ we have

$$
u_k < u_k + 1 \le u_{k+1} \le u_i
$$

and g_{u_k+1} divides g_{u_i} , that is,

$$
g_{u_i} \equiv 0 \pmod{g_{u_k+1}}.
$$

If $v_j \geq u_k + 1$, then

$$
g_{v_j} \equiv 0 \pmod{g_{u_k+1}}.
$$

Rearranging [\(5\)](#page-4-0), we obtain

(7)
$$
\sum_{u_i \le u_k} x_i g_{u_i} - \sum_{v_j \le u_k} y_j g_{v_j} = \sum_{v_j \ge u_k + 1} y_j g_{v_j} - \sum_{u_i > u_k} x_i g_{u_i}
$$

$$
\equiv 0 \pmod{g_{u_k + 1}}.
$$

If

$$
\sum_{u_i \le u_k} x_i g_{u_i} > \sum_{v_j \le u_k} y_j g_{v_j}
$$

then Lemma [1](#page-1-0) gives

$$
0 < \sum_{i=1}^{k} x_i g_{u_i} - \sum_{v_j \le u_k} y_j g_{v_j} \le \sum_{i=1}^{k} x_i g_{u_i} < g_{u_{k+1}}.
$$

This inequality contradicts congruence (7) . This completes the proof. \Box

Theorem 3. Let $h \geq 2$ and let t be an integer such that

$$
(8) \t t \ge 1 + \frac{\log h}{\log 2}.
$$

Let $W = (W_i)_{i=0}^{h-1}$ be a partition of \mathbf{N}_0 such that, for all $i \in [0, h-1]$, there is an infinite set \mathcal{M}_i of positive integers such that

$$
(9) \t\t\t\t[M_i - t + 1, M_i] \subseteq W_i
$$

for all $M_i \in \mathcal{M}_i$. Let $\mathcal{G} = (g_i)_{i=0}^{\infty}$ be a \mathcal{G} -adic sequence. The set

$$
A_{\mathcal{G}}(\mathcal{W}) = \bigcup_{i=0}^{h-1} A_{\mathcal{G}}(W_i)
$$

is a minimal asymptotic basis of order h.

Proof. By Theorem [1,](#page-2-0) the set $A_{\mathcal{G}}(W)$ is an asymptotic basis of order h.

Let $a \in A_{\mathcal{G}}(\mathcal{W})$. Without loss of generality, we can assume that $a = a_0 \in A_{\mathcal{G}}(W_0)$ and

(10)
$$
a_0 = \sum_{j \in F_0} x_{0,j} g_j
$$

where $F_0 \in \mathcal{F}^*(W_0)$ and $x_{0,j} \in [1, d_{j+1} - 1]$ for all $j \in F_0$. Let $M_0 = \max(F_0)$. By Lemma [1,](#page-1-0)

$$
g_{M_0} \le a_0 < g_{M_0+1}.
$$

For all $i \in [1, h-1]$, choose an integer M_i in the infinite set \mathcal{M}_i such that

$$
(11) \t\t\t M_i \ge M_0 + t
$$

and let

(12)
$$
a_i = \sum_{\substack{j \in W_i \\ j < M_0}} (d_{j+1} - 1) g_j + g_{M_i} \in A_{\mathcal{G}}(W_i).
$$

This is the $\mathcal{G}\text{-}\text{adic representation of }a_i$. Let

(13)
$$
n = \sum_{i=0}^{h-1} a_i = a_0 + \sum_{i=1}^{h-1} \sum_{\substack{j \in W_i \\ j < M_0}} (d_{j+1} - 1) g_j + \sum_{i=1}^{h-1} g_{M_i}.
$$

This is the $\mathcal{G}\text{-}\text{adic representation of }n.$

Let

$$
n = \sum_{i=0}^{h-1} b_{k_i}
$$

be any representation of n as the sum of h elements of $A_{\mathcal{G}}(\mathcal{W})$, where $k_i \in [0, h-1]$ for all $i \in [0, h-1]$ and $b_{k_i} \in A_{\mathcal{G}}(W_{k_i})$. We must prove that $b_{k_i} = a_0$ for some $i \in [0, h - 1].$

Each integer b_{k_i} is of the form

$$
b_{k_i} = \sum_{j \in E_{k_i}} y_{i,j} g_j \in A_{\mathcal{G}}(W_{k_i})
$$

where $E_{k_i} \in \mathcal{F}^*(W_{k_i})$ and $y_{i,j} \in [1, g_{j+1} - 1]$ for all $j \in E_{k_i}$. The uniqueness of the G-adic representation implies that if $\{k_0, k_1, \ldots, k_{h-1}\} = [0, h-1]$, then, after rearrangement, $k_i = i$ and $a_i = b_i$ for all $i \in [0, h - 1]$.

If ${k_0, k_1, \ldots, k_{h-1}} \neq [0, h-1]$, then there exists $s \in [0, h-1]$ such that $s \notin$ ${k_0, k_1, \ldots, k_{h-1}}$. Suppose that $s \neq 0$. Recall that

$$
M_s \ge M_0 + t
$$

and

$$
[M_s - t + 1, M_s] \subseteq W_s.
$$

Because $k_i \neq s$ for all $i \in [0, h-1]$, we have

$$
[M_s - t + 1, M_s] \cap E_{k_i} \subseteq W_s \cap W_{k_i} = \emptyset.
$$

We construct the partition

$$
E_{k_i} = E'_{k_i} \cup E''_{k_i}
$$

with

$$
E'_{k_i} = \{j \in E_{k_i} : j \leq M_s - t\}
$$

and

$$
E_{k_i}'' = \{ j \in E_{k_i} : j \ge M_s + 1 \}.
$$

The sets E'_{k_i} and E''_{k_i} are not necessarily nonempty. Let

$$
b_{k_i}=b_{k_i}^\prime+b_{k_i}^{\prime\prime}
$$

where

$$
b'_{k_i} = \sum_{j \in E'_{k_i}} y_{i,j} g_j \quad \text{and} \quad b''_{k_i} = \sum_{j \in E''_{k_i}} y_{i,j} g_j.
$$

Note that $b'_{k_i} = 0$ if $E'_{k_i} = \emptyset$ and $b''_{k_i} = 0$ if $E''_{k_i} = \emptyset$. By Lemma [1,](#page-1-0)

$$
(14) \t\t b'_{k_i} < g_{M_s - t + 1}
$$

and

$$
b_{k_i}'' = 0 \qquad \text{or} \qquad b_{k_i}'' \ge g_{M_s + 1}.
$$

Let

(15)
$$
n' = \sum_{i=0}^{h-1} b'_{k_i} \quad \text{and} \quad n'' = \sum_{i=0}^{h-1} b''_{k_i}.
$$

Recall inequalities [\(1\)](#page-1-1) and [\(8\)](#page-5-1):

$$
d_i \ge 2 \qquad \text{and} \qquad h \le 2^{t-1}.
$$

From (15) , (14) , and (3) , we obtain

$$
n' < hg_{M_s-t+1} \le 2^{t-1}g_{M_s-t+1} \le g_{M_s-t+1} \prod_{i=1}^{t-1} d_{M_s-t+i} = g_{M_s}.
$$

Therefore, the $\mathcal{G}\text{-}\text{adic representation of } n'$ is of the form

$$
n' = \sum_{j=0}^{M_s - 1} z_j g_j
$$

with $z_j \in [0, d_{j+1} - 1]$. Because

$$
n'' = 0 \qquad \text{or} \qquad n'' \ge g_{M_s + 1},
$$

the $\mathcal{G}\text{-}\text{adic representation of }n^{\prime\prime}$ is of the form

$$
n'' = \sum_{j=M_s+1}^{\infty} z_j g_j
$$

with $z_j \in [0, d_{j+1} - 1]$ and $z_j \ge 1$ for only finitely many j. Therefore,

$$
n = n' + n'' = \sum_{i=0}^{M_s - 1} z_i g_i + \sum_{i=M_s + 1}^{\infty} z_i g_i
$$

is the G-adic representation of n. In this representation, the coefficient of g_{M_s} is 0, which contradicts the construction of n . It follows that

$$
[1, h-1] \subseteq \{k_0, k_1, \ldots, k_{h-1}\}.
$$

Renumbering the integers b_i , we can assume that $k_i = i$ and $b_i \in A_{\mathcal{G}}(W_i)$ for all $i \in [1, h-1].$

We must prove that $k_0 = 0$, or, equivalently, that $b_0 \in A_{\mathcal{G}}(W_0)$. If not, then $b_0 \in A_{\mathcal{G}}(W_r)$ for some $r \in [1, h-1]$. Because

$$
M_0 = \max(F_0) \in F_0 \subseteq W_0
$$

we have

$$
M_0 \notin \bigcup_{i=0}^{h-1} E_{k_i} \subseteq \bigcup_{i=1}^{h-1} W_i.
$$

Identity [\(10\)](#page-5-2) and Lemma [1](#page-1-0) give

$$
g_{M_0} \le a_0 = \sum_{j \in F_0} x_{0,j} g_j < g_{M_0+1}.
$$

From [\(12\)](#page-6-1) we have

$$
n = a_0 + \sum_{i=1}^{h-1} \sum_{\substack{j \in W_i \\ j < M_0}} (d_{j+1} - 1) g_j + \sum_{i=1}^{h-1} g_{M_i}
$$

$$
= \sum_{i=0}^{h-1} \sum_{j \in E_i} y_{i,j} g_j.
$$

Summing only over terms g_j with $j \leq M_0$ and applying Lemma [2,](#page-4-2) we obtain

$$
a_0 + \sum_{i=1}^{h-1} \sum_{\substack{j \in W_i \\ j < M_0}} (d_{j+1} - 1)g_j \le \sum_{i=0}^{h-1} \sum_{\substack{j \in E_i \\ j \le M_0}} y_{i,j}g_j = \sum_{i=0}^{h-1} \sum_{\substack{j \in E_i \\ j < M_0}} y_{i,j}g_j
$$

$$
= \sum_{\substack{j \in E_0 \\ j < M_0}} y_{0,j}g_j + \sum_{i=1}^{h-1} \sum_{\substack{j \in E_i \\ j < M_0}} y_{i,j}g_j
$$

$$
< g_{M_0} + \sum_{i=1}^{h-1} \sum_{\substack{j \in W_i \\ j < M_0}} (d_{j+1} - 1)g_j
$$

$$
\le a_0 + \sum_{i=1}^{h-1} \sum_{\substack{j \in W_i \\ j < M_0}} (d_{j+1} - 1)g_j.
$$

This is absurd, and so $b_0 \in A_G(W_0)$. It follows that the integer n defined by [\(13\)](#page-6-2) has a unique representation as the sum of h elements of $A_{\mathcal{G}}(\mathcal{W})$. Therefore,

(16)
$$
n \notin h(A_{\mathcal{G}}(\mathcal{W}) \setminus \{a\}).
$$

For all $i \in [1, h-1]$, there are infinitely many integers $M_i \in \mathcal{M}_i$ with $M_i \geq M_0+t$, and so infinitely many positive integers n satisfying [\(16\)](#page-8-0). It follows that $A_{\mathcal{G}}(\mathcal{W})\backslash\{a\}$ is not an asymptotic basis of order h for all $a \in A_{\mathcal{G}}(\mathcal{W})$. Equivalently, $A_{\mathcal{G}}(\mathcal{W})$ is a minimal asymptotic basis of order h. This completes the proof. \Box

Corollary 1. Let $W = W_0 \cup W_1$ be a partition of \mathbf{N}_0 such that both W_0 and W_1 contain infinitely many pairs of consecutive integers. Let $\mathcal{G} = (g_i)_{i=0}^{\infty}$ be a \mathcal{G} -adic sequence. The set

$$
A_{\mathcal{G}}(\mathcal{W}) = A_{\mathcal{G}}(W_1) \cup A_{\mathcal{G}}(W_2)
$$

is a minimal asymptotic basis of order 2.

Proof. This is the case $h = 2$ of Theorem [3.](#page-5-3)

3. Open problems

(1) Let $h \geq 2$. By Theorem [1,](#page-2-0) for every partition $\mathcal{W} = (W_i)_{i=0}^{h-1}$ of \mathbf{N}_0 and every G-adic sequence, the set

$$
A = \{0\} \cup A_{\mathcal{G}}(\mathcal{W})
$$

is a basis of order h that is not a minimal asymptotic basis of order h .

(a) Determine the set of all integers $a \in A$ such that $A \setminus \{a\}$ is an asymptotic basis of order h.

10 MELVYN B. NATHANSON

- (b) Determine the set of all integers $a \in A$ such that $A \setminus \{a\}$ is a minimal asymptotic basis.
- (2) Let $\mathcal{G} = (g_i)_{i=0}^{\infty}$ be a \mathcal{G} -adic sequence. Let $h \geq 2$.
	- (a) Construct partitions $W = (W_i)_{i=0}^{h-1}$ of \mathbf{N}_0 such that the set $A_{\mathcal{G}}(W)$ is a minimal asymptotic basis of order h.
	- (b) Construct partitions $W = (W_i)_{i=0}^{h-1}$ of \mathbf{N}_0 such that the set $A_{\mathcal{G}}(W)$ is not a minimal asymptotic basis of order h.
- (3) Let $(d_i)_{i=1}^{\infty}$ be a sequence of 2s and 3s. Let $\mathcal{G} = (g_i)_{i=0}^{\infty}$ be the \mathcal{G} -adic sequence defined by $g_0 = 1$ and $g_i = \prod_{j=1}^i d_i$ for $i \ge 1$. Consider problem [\(2\)](#page-9-17) with respect to this G -adic sequence. Of particular interest are the infinitely many G-adic sequences $G = (g_i)_{i=0}^{\infty}$ with quotients $\{d_{2i-1}, d_{2i}\} = \{2, 3\}$ for all $i = 1, 2, 3, ...$ In this case, $g_{2i} = 6^i$ for all *i*.

REFERENCES

- [1] F.-J. Chen and Y.-G. Chen, On minimal asymptotic bases, European J. Combin. 32 (2011), no. 8, 1329–1335.
- [2] Y.-G. Chen, On the Erdős-Turán conjecture, C. R. Math. Acad. Sci. Paris 350 (2012), no. 21-22, 933–935.
- [3] Y.-G. Chen and M. Tang, On a problem of Nathanson, Acta Arith. 185 (2018), no. 3, 275– 280.
- [4] P. Erdős and P. Turán, On a problem of Sidon in additive number theory, and on some related problems, J. London Math. Soc. 16 (1941), 212-215.
- [5] X.-D. Jia, Minimal bases and g-adic representations of integers, Number Theory: New York Seminar 1991-1995 (New York), Springer-Verlag, 1996, pp. 201–209.
- [6] X.-D. Jia and M. B. Nathanson, A simple construction of minimal asymptotic bases, Acta Arith. 52 (1989), no. 2, 95–101.
- [7] J. B. Lee, A construction of minimal asymptotic bases, Periodica Math. Hungar. 26 (1993), 211–218.
- [8] J. Li and J. Li, On minimal asymptotic basis of order 4, J. Math. Res. Appl. 36 (2016), no. 6,16, 651–658.
- [9] D. Ling and M. Tang, On minimal asymptotic g-adic bases, Bull. Aust. Math. Soc. 92 (2015), no. 3, 374–379.
- [10] , On minimal asymptotic bases of order three, Colloq. Math. 151 (2018), no. 1, 9–18.
- [11] M. B. Nathanson, *Minimal bases and maximal nonbases in additive number theory*, J. Number Theory 6 (1974), 324–333. [12] $\frac{1}{\text{Mean of 2}}$, Minimal bases and powers of 2, Acta Arith. 49 (1988), no. 5, 525–532.
-
- [13] $\qquad \qquad$, Additive systems and a theorem of de Bruijn, Amer. Math. Monthly 121 (2014), no. 1, 5–17.
- [14] $____\,,$ Limits and decomposition of de Bruijn's additive systems, Combinatorial and additive number theory. II, Springer Proc. Math. Stat., vol. 220, Springer, Cham, 2017, pp. 255– 267.
- [15] C.-F. Sun, On minimal asymptotic basis of order $g-1$, Indag. Math. (N.S.) 30 (2019), no. 1, 128–135.
- [16] , On a problem of Nathanson on minimal asymptotic bases, Journal of Number Theory 218 (2021), 152–160.
- [17] C.-F. Sun and T.-T. Tao, On g-adic minimal asymptotic bases of order h, Int. J. Number Theory 15 (2019), no. 2, 389–406.

Lehman College (CUNY), Bronx, NY 10468 Email address: melvyn.nathanson@lehman.cuny.edu