

ON THE STRINGY HODGE NUMBERS OF MIRRORS OF QUASI-SMOOTH CALABI-YAU HYPERSURFACES

VICTOR V. BATYREV

ABSTRACT. Mirrors X^\vee of quasi-smooth Calabi-Yau hypersurfaces X in weighted projective spaces $\mathbb{P}(w_0, \dots, w_d)$ can be obtained as Calabi-Yau compactifications of non-degenerate affine toric hypersurfaces defined by Laurent polynomials whose Newton polytope is the lattice simplex spanned by $d+1$ lattice vectors v_i satisfying the relation $\sum_i w_i v_i = 0$. In this paper, we compute the stringy E -function of mirrors X^\vee and compare it with the Vafa's orbifold E -function of quasi-smooth Calabi-Yau hypersurfaces X . As a result, we prove the equalities of Hodge numbers $h_{\text{str}}^{p,q}(X^\vee) = h_{\text{orb}}^{d-1-p,q}(X)$ for all p, q and d as it is expected in mirror symmetry.

1. INTRODUCTION

Let X be a d -dimensional Calabi-Yau variety, i.e., a normal irreducible projective variety with at worst canonical Gorenstein singularities satisfying the conditions: $\Omega_X^d \cong \mathcal{O}_X$ and $h^i(X, \mathcal{O}_X) = 0$ ($0 < i < d$). We take a desingularization $\rho : Y \rightarrow X$ of X such that the exceptional locus of ρ is a union of smooth irreducible divisors D_1, \dots, D_s with only normal crossings and write

$$K_Y = \rho^* K_X + \sum_{i=1}^s a_i D_i$$

with non-negative integers a_i . We set $D_\emptyset := Y$ and consider for any subset $J \subseteq I = \{1, \dots, s\}$ the smooth projective variety $D_J := \bigcap_{j \in J} D_j \subseteq Y$ together with its open subvariety $D_J^\circ := D_J \setminus \bigcup_{j \notin J} D_j$. We denote by $E(D_J; u, v) := \sum_{p,q} (-1)^{p+q} h^{p,q}(D_J) u^p v^q$ the usual Poincaré polynomial of Hodge numbers of smooth projective varieties D_J and extend them by additivity to E -polynomials $E(D_J^\circ; u, v) := \sum_{p,q} e^{p,q}(D_J^\circ) u^p v^q$ defined by the Hodge-Deligne numbers $e^{p,q}(D_J^\circ)$ of quasi-projective varieties D_J° .

Definition 1.1. [Bat98] The stringy E -function of a Calabi-Yau variety X is the rational function in two variables u, v defined by two equivalent formulas:

$$\begin{aligned} E_{\text{str}}(X; u, v) &:= \sum_{\emptyset \subseteq J \subseteq I} \left(\prod_{j \in J} \frac{uv - 1}{(uv)^{a_j+1} - 1} - 1 \right) E(D_J; u, v) \\ &= \sum_{\emptyset \subseteq J \subseteq I} \left(\prod_{j \in J} \frac{uv - 1}{(uv)^{a_j+1} - 1} \right) E(D_J^\circ; u, v). \end{aligned}$$

If $E_{\text{str}}(X; u, v)$ is a polynomial, then the *stringy Hodge numbers* $h_{\text{str}}^{p,q}(X)$ are defined via its coefficients:

$$E_{\text{str}}(X; u, v) = \sum_{p,q} (-1)^{p+q} h_{\text{str}}^{p,q}(X) u^p v^q.$$

The stringy Hodge numbers are very useful for testing Mirror Symmetry. If two d -dimensional Calabi-Yau varieties X and X^\vee are mirror symmetric to each other, then

$$(1.1) \quad h_{\text{str}}^{p,q}(X) = h_{\text{str}}^{d-p,q}(X^\vee), \quad \forall p, q.$$

Equations (1.1) can be equivalently reformulated in the form

$$(1.2) \quad E_{\text{str}}(X^\vee; u, v) = (-u)^d E_{\text{str}}(X; u^{-1}, v).$$

Equation (1.2) has been checked for all pairs of Calabi-Yau hypersurfaces in Gorenstein toric Fano varieties defined by a pair (Δ, Δ^*) of dual to each other reflexive polytopes [Bat94, BB96]. Note that in general the stringy E -function $E_{\text{str}}(X; u, v)$ is a rational function whose denominator is a product of cyclotomic polynomials $\Phi_k(uv)$ in uv . Therefore, the equation (1.2) can be satisfied only provided $E_{\text{str}}(X; u, v)$ is a polynomial.

In this paper we consider Calabi-Yau hypersurfaces $X = X_w$ of degree $w = \sum_{i=0}^d w_i$ in weighted projective space $\mathbb{P}(\bar{w}) = \mathbb{P}(w_0, w_1, \dots, w_d)$ which is not assumed to be Gorenstein. Such a weighted projective space $\mathbb{P}(\bar{w})$ is defined by a *weight vector* $\bar{w} \in \mathbb{Z}^{d+1}$ having *IP-property*. Recall that $\bar{w} \in \mathbb{Z}^{d+1}$ has *IP-property* if the convex hull of all non-negative lattice points $(u_0, u_1, \dots, u_d) \in \mathbb{Z}_{\geq 0}^{d+1}$ satisfying the relation $\sum_{i=0}^d w_i u_i = w$ is a d -dimensional lattice polytope containing $\mathbf{1} := (1, \dots, 1)$ in its interior [Ska96]. This lattice polytope is just the Newton polytope of a general weighted homogeneous polynomial W defining X_w in $\mathbb{P}(\bar{w})$. If $0 \in \mathbb{C}^{d+1}$ is the only singularity of the affine hypersurface $\{W = 0\} \subset \mathbb{C}^{d+1}$, then the corresponding Calabi-Yau hypersurface $X_w \subset \mathbb{P}(\bar{w})$ is called *quasi-smooth* and the weight vector \bar{w} is called *transverse*. It is known that any transverse weight vector \bar{w} has *IP-property* [Ska96]. We consider the weighted projective space $\mathbb{P}(\bar{w})$ as a toric variety with respect to the natural regular action of the d -dimensional algebraic torus $\mathbb{T}_{\bar{w}} := (\mathbb{C}^*)^{d+1}/H$, where

$$H = \{(\lambda^{w_0}, \dots, \lambda^{w_d}) \in (\mathbb{C}^*)^{d+1} : \lambda \in \mathbb{C}^*\}.$$

The lattice of characters of $\mathbb{T}_{\bar{w}}$ is the sublattice

$$N_{\bar{w}} := \{(u_0, u_1, \dots, u_d) \in \mathbb{Z}^{d+1} \mid \sum_{i=0}^d w_i u_i = 0\} \subset \mathbb{Z}^{d+1}.$$

We use the following common notations:

$$(q_0, q_1, \dots, q_d) := \left(\frac{w_0}{w}, \frac{w_1}{w}, \dots, \frac{w_d}{w} \right) \in \mathbb{Q}_{>0}^{d+1},$$

$$\theta(l) := (\theta_0(l), \theta_1(l), \dots, \theta_d(l)) = (lq_0, lq_1, \dots, lq_d), \quad 0 \leq l < w;$$

$$\tilde{\theta}(l) := (\tilde{\theta}_0(l), \tilde{\theta}_1(l), \dots, \tilde{\theta}_d(l)) \in [0, 1)^{d+1}, \quad 0 \leq l < w,$$

where $\tilde{\theta}(l)$ is the canonical representative of $\theta(l)$ in $(\mathbb{R}/\mathbb{Z})^{d+1}$.

Let $\mathbb{T}_{\bar{w}}^* \cong (\mathbb{C}^*)^d$ be the dual to $\mathbb{T}_{\bar{w}}$ torus with the lattice of characters $M_{\bar{w}} = \mathbb{Z}^{d+1}/\mathbb{Z}\bar{w}$. Denote by $\Delta_{\bar{w}}^*$ the d -dimensional lattice simplex in $M_{\bar{w}} \otimes \mathbb{R}$ whose vertices v_0, v_1, \dots, v_d are primitive lattice vectors spanning $M_{\bar{w}}$ and satisfying the relation $\sum_{i=0}^d w_i v_i = 0$. We claim that mirrors X_w^\vee of quasi-smooth Calabi-Yau hypersurfaces $X_w \subset \mathbb{P}(\bar{w})$ can be obtained as Calabi-Yau compactifications of non-degenerate

affine hypersurfaces in $Z_{\bar{w}} \subset \mathbb{T}_{\bar{w}}^*$ defined by Laurent polynomials with the Newton polytope $\Delta_{\bar{w}}^*$. Using a formula of Vafa for the orbifold Euler number of a quasi-smooth Calabi-Yau hypersurface $X_w \subset \mathbb{P}(\bar{w})$ [Vaf89]:

$$\chi_{\text{orb}}(X_w) = \frac{1}{w} \sum_{l,r=0}^{w-1} \prod_{\substack{0 \leq i \leq d \\ lq_i, rq_i \in \mathbb{Z}}} \left(1 - \frac{1}{q_i}\right),$$

the topological mirror symmetry duality of the stringy Euler numbers

$$E_{\text{str}}(X_{\bar{w}}^{\vee}; 1, 1) = (-1)^{d-1} E_{\text{str}}(X_w; 1, 1)$$

was checked for this mirror construction in [BS20, Theorem 1.6]. We note that the quasi-smooth Calabi-Yau hypersurfaces $X_w \subset \mathbb{P}(\bar{w})$ are orbifolds and, by a theorem of Yasuda [Yas04], the stringy Hodge numbers $h_{\text{str}}^{p,q}(X_w)$ are equal to the orbifold Hodge numbers $h_{\text{orb}}^{p,q}(X_w)$ introduced by Chen and Ruan [CR04].

In order to compute the orbifold Hodge numbers $h_{\text{orb}}^{p,q}(X_w)$ by the method of Vafa and to compare them with the stringy Hodge numbers of mirrors $X_{\bar{w}}^{\vee}$ we need some integers associated with elements l in the cyclic group $G_0 := \mathbb{Z}/w\mathbb{Z}$:

- $\text{age}(l) := \sum_{i=0}^d \tilde{\theta}_i(l) = \sum_{\tilde{\theta}_i(l) \neq 0} \tilde{\theta}_i(l)$, ($l \in \mathbb{Z}/w\mathbb{Z}$);
- $\text{size}(l) := \text{age}(l) + \text{age}(w-l) = \text{Card}\{i \in \{0, 1, \dots, d\} : \tilde{\theta}_i(l) \neq 0\}$.

If $P(t) = \sum_i b_i t^{r_i}$ is a polynomial or a power series with exponents $r_i \in \frac{1}{w}\mathbb{Z}$ we denote by $[P(t)]_{\text{int}}$ the projector dropping all monomials with fractional exponents in $P(t)$ and leaving only monomials $b_i t^{r_i}$ with exponents $r_i \in \mathbb{Z}$. Using the action of G_0 by w -th root of unity $e^{2\pi i k/w}$ on $t^{k/w}$, the projector $[*]_{\text{int}}$ can be expressed as a standard Reynolds operator

$$[P]_{\text{int}} = \frac{1}{|G_0|} \sum_{g \in G_0} P^g.$$

In particular, one has

$$(1.3) \quad [P \cdot Q]_{\text{int}} = P \cdot [Q]_{\text{int}},$$

if P contains only monomials with integral exponents.

Our main result that extends [BS20, Theorem 1.6] is the following:

Theorem 1.2. *Let $\bar{w} \in \mathbb{Z}^{d+1}$ be a weight vector with IP-property. Denote by $Z_{\bar{w}} \subset \mathbb{T}_{\bar{w}}^*$ a non-degenerate affine hypersurface defined by a Laurent polynomial with the Newton polytope $\Delta_{\bar{w}}^* = \text{conv}(v_0, \dots, v_d)$. Then $Z_{\bar{w}}$ admits a Calabi-Yau compactification $X_{\bar{w}}^{\vee}$ and the stringy E -function of $X_{\bar{w}}^{\vee}$ can be computed by the formula*

$$E_{\text{str}}(X_{\bar{w}}^{\vee}; u, v) = \frac{1}{uv} \sum_{0 \leq l < w} \left[\prod_{\tilde{\theta}_i(l)=0} \frac{(uv)^{q_i} - uv}{1 - (uv)^{q_i}} \right]_{\text{int}} \cdot (-u)^{\text{size}(l)} \left(\frac{v}{u}\right)^{\text{age}(l)}.$$

Moreover, if the weight vector \bar{w} is transverse, then $E(X_{\bar{w}}^{\vee}; u, v)$ is a polynomial and one has

$$E_{\text{str}}(X_{\bar{w}}^{\vee}; u, v) = (-u)^{d-1} E_{\text{orb}}(X_w; u^{-1}, v)$$

where $X_w \subset \mathbb{P}(\bar{w})$ is a quasi-smooth Calabi-Yau hypersurface of degree $w = \sum_{i=0}^d w_i$.

We remark that the considered mirror construction for arbitrary quasi-smooth Calabi-Yau hypersurfaces $X_w \subset \mathbb{P}(\overline{w})$ extends the well-known Berglund-Hübsch-Krawitz mirror construction [BH93, Kra10, CR18]. In this construction, one considers a transverse polynomial W of special form

$$W = \sum_{i=0}^d c_i \prod_{j=0}^d z_i^{a_{ij}}, \quad c_0, c_1, \dots, c_d \in \mathbb{C}^*.$$

Such a polynomial W is called *invertible*. Consider the action of $l \in G_0 = \mathbb{Z}/w\mathbb{Z}$ on \mathbb{C}^{d+1} by diagonal matrices

$$\text{diag}(e^{2\pi i l q_0}, e^{2\pi i l q_1}, \dots, e^{2\pi i l q_d}).$$

One associates with the quasi-smooth Calabi-Yau hypersurface $X_w = \{W = 0\} \subset \mathbb{P}^d(\overline{w})$ an orbifold Landau-Ginzburg model with the G_0 -invariant superpotential W . We set $\widetilde{G}_0 := SL(d+1, \mathbb{C}) \cap \widetilde{G}_{\widetilde{W}}$, where $\widetilde{G}_{\widetilde{W}} \cong G_W$ is a maximal abelian diagonal symmetry group of the dual invertible polynomial

$$\widetilde{W} := \sum_{j=0}^d c_j \prod_{i=0}^d \widetilde{z}_i^{a_{ij}}.$$

Since every monomial W_i in W has degree w , we obtain the equations

$$\sum_{j=0}^d a_{ij} w_j = w \quad \forall i \in \{0, 1, \dots, d\}$$

which imply that the Newton polytope of the Laurent polynomial

$$\widetilde{W} / \left(\prod_{i=0}^d \widetilde{z}_i \right)^w$$

with respect to the \widetilde{G}_0 -invariant lattice $M_{\widetilde{w}} = \mathbb{Z}^{d+1} / \mathbb{Z}\widetilde{w}$ is a d -dimensional simplex $\Delta_{\widetilde{w}}^*$ whose vertices v_0, v_1, \dots, v_d generate $M_{\widetilde{w}}$ and satisfy the relation $\sum_{i=0}^d w_i v_i = 0$.

Remark 1.3. Note that the mirror construction for quasi-smooth hypersurfaces $X_w \subset \mathbb{P}(\overline{w})$ and the proof of Theorem 1.2 naturally extend to the case of Calabi-Yau quotients $X_w/G \subset \mathbb{P}(\overline{w})/G$, where $G \subset SL(d+1, \mathbb{C})$ ($G_0 \subset G$) is a finite abelian group of diagonal symmetries of the defining transverse polynomial W . In this case, mirrors of X_w/G are Calabi-Yau compactifications of non-degenerate affine hypersurfaces $Z_w \subset \mathbb{T}_{\overline{w}}^*(G)$ whose Newton polytope is the same lattice simplex $\Delta_{\overline{w}}^*$, but considered with respect to a larger lattice $M_{\overline{w}}(G)$ such that $M_{\overline{w}}(G)/M_{\overline{w}}$ is isomorphic to G/G_0 .

Remark 1.4. A combinatorial formula for the orbifold E -function of pair (W, G) , in spirit of Vafa [Vaf89], has been obtained by Ebeling and Takahashi in [ET13, Theorem 6]. The topological mirror duality between the orbifold E -functions of (W, G) and $(\widetilde{W}, \widetilde{G})$ was shown by Ebeling, Gusein-Zade, and Takahashi in [EGZT16, Theorem 9]. The isomorphism of the orbifold Hodge spaces in Berglund-Hübsch-Krawitz mirror construction was shown by Chiodo-Ruan [CR10] and Borisov [Bo13].

2. E -POLYNOMIALS OF NON-DEGENERATE HYPERSURFACES

In this section we review some results of Danilov and Khovanskii [DKh86] (see also [Bat93]).

Let X be a quasi-projective variety over \mathbb{C} . Then the cohomology space with compact supports $H_c^k(X, \mathbb{C})$ have natural mixed Hodge structure and one defines *Hodge-Deligne numbers* of X as

$$e^{p,q}(X) := \sum_k (-1)^k h^{p,q}(H_c^k(X)).$$

Then

$$E(X; u, v) := \sum_{p,q} e^{p,q}(X) u^p v^q.$$

is called E -polynomial of X .

Let M be a free abelian group of rank d . We consider M as the lattice of characters of d -dimensional algebraic torus $\mathbb{T} := \text{Hom}(M, \mathbb{C}^*)$, i.e., $\mathbb{T} = \text{Spec } \mathbb{C}[M]$. A Laurent polynomial $f \in \mathbb{C}[M]$ defines an affine hypersurface $Z_f = \{f = 0\} \subset \mathbb{T}$. Assume that the Newton polytope Δ of f has dimension d . By choosing a \mathbb{Z} -basis of M , we obtain an isomorphism $\mathbb{C}[M] \cong \mathbb{C}[t_1^{\pm 1}, \dots, t_d^{\pm 1}]$. Denote by $M_\Delta := C_\Delta \cap (\mathbb{Z} \oplus M)$ the graded monoid of all lattice points in the $(d+1)$ -dimensional cone $C_\Delta := \mathbb{R}_{\geq 0}(1, \Delta) \subset \mathbb{R} \oplus M_{\mathbb{R}}$, where $\deg(k, m) := k$ ($\forall k \in \mathbb{Z}, m \in M$). Consider the graded semigroup ring

$$S_\Delta := \mathbb{C}[M_\Delta] = \bigoplus_{k \geq 0} S_\Delta^{(k)} \subset \mathbb{C}[t_0, t_1^{\pm 1}, \dots, t_d^{\pm 1}],$$

whose homogeneous components are \mathbb{C} -spaces of dimension $|k\Delta \cap M|$. Note that $S_\Delta^{(1)} \subset t_0 \mathbb{C}[t_1^{\pm 1}, \dots, t_d^{\pm 1}]$ can be identified with the \mathbb{C} -vector space $L(\Delta) \subset \mathbb{C}[t_1^{\pm 1}, \dots, t_d^{\pm 1}]$ spanned by all Laurent polynomials with the Newton polytope Δ .

Definition 2.1. A Laurent polynomial $f \in L(\Delta) \subset \mathbb{C}[t_1^{\pm 1}, \dots, t_d^{\pm 1}]$ is called *non-degenerate* if

$$t_0 f, t_0 t_1 \frac{\partial f}{\partial t_1}, \dots, t_0 t_d \frac{\partial f}{\partial t_d} \in t_0 L(\Delta) = S_\Delta^{(1)}$$

form a regular sequence in S_Δ .

Remark 2.2. Let $A \subset \Delta \cap M$ be a finite subset which includes all vertices of the lattice polytope Δ . Then non-degenerate Laurent polynomials

$$f(\mathbf{t}) = \sum_{m \in A} a_m \mathbf{t}^m, \quad a_m \in \mathbb{C},$$

form a Zariski dense open subset U_A in the affine space $\mathbb{C}^{|A|}$ of all coefficients $\{a_m\}_{m \in A}$. The subset U_A can be explicitly defined by the non-vanishing of the principal A -determinant introduced by Gelfand, Kapranov and Zelevinski [GKZ94].

Remark 2.3. The Poincaré series of the graded ring S_Δ

$$P(\Delta, t) := \sum_{k \geq 0} |k\Delta \cap M| t^k$$

is a rational function of the form

$$P(\Delta, t) = \frac{\psi_0(\Delta) + \psi_1(\Delta)t + \cdots + \psi_d(\Delta)t^d}{(1-t)^{d+1}},$$

where $\psi_0(\Delta) = 0$ and $\sum_{i=0}^d \psi_i(\Delta) = \text{Vol}_d(\Delta) = d! \text{vol}(\Delta) \in \mathbb{Z}$. The coefficient $\psi_i(\Delta)$ is equal to the dimension of the i -th homogeneous component of the artinian ring

$$S_\Delta / (t_0 f, t_0 t_1 \frac{\partial f}{\partial t_1}, \dots, t_0 t_d \frac{\partial f}{\partial t_d}).$$

Theorem 2.4. [DKh86, Remark 4.6] *Let $f \in L(\Delta)$ be a non-degenerate Laurent polynomial and let $Z_{f,\Delta} \subset \mathbb{T}$ be the non-degenerate affine hypersurface defined by the equation $f = 0$. Then the E -polynomial $E(Z_{f,\Delta}; u, 1)$ has the following form:*

$$E(Z_{f,\Delta}; u, 1) = \frac{(u-1)^d - (-1)^d}{u} + (-1)^{d-1} \sum_{i=1}^d \psi_i(\Delta) u^{i-1}.$$

Remark 2.5. Explicit formulas for all Hodge-Deligne numbers $e^{p,q}(Z_f)$ as coefficients in E -polynomial $E(Z_f; u, v)$ were obtained by Danilov and Khovanskii only for simple d -dimensional lattice polytopes Δ [DKh86, Theorem 5.6]. The computations for arbitrary d -dimensional lattice polytopes Δ are algorithmic more complicated (see [BB96, Sa20]).

In this paper we need explicit formulas for E -polynomials only in case when $\Delta \subset M_{\mathbb{R}}$ is a d -dimensional lattice simplex with vertices v_0, v_1, \dots, v_d . We need also formulas for E -polynomials of affine hypersurfaces Z_{f,Δ_J} corresponding to faces D_J of Δ parametrized by nonempty subsets $J \subseteq I := \{0, 1, \dots, d\}$. If $J = \{j_1, \dots, j_k\} \subset I$ is a subset, we denote $\Delta_J := \text{conv}\{v_{j_1}, \dots, v_{j_k}\}$. It is convenient to associate with a simplex Δ a finite abelian group $G(\Delta)$ defined as follows:

Definition 2.6. Let Δ be a d -dimensional lattice simplex with vertices v_0, v_1, \dots, v_d . We set $\tilde{v}_i := (1, v_i) \in \mathbb{Z} \oplus M$ ($0 \leq i \leq d$) and denote by $M' \subset \mathbb{Z} \oplus M$ the sublattice of rank $d+1$ spanned by linearly independent lattice vectors $\{\tilde{v}_i\}_{i \in I}$. Then

$$G(\Delta) := (\mathbb{Z} \oplus M) / M'$$

is a finite abelian group of order $\text{Vol}_d(\Delta) = d! \text{vol}(\Delta)$ and one can write an element $\tilde{m} = (k, m) \in (\mathbb{Z} \oplus M) / M'$ as a unique \mathbb{Q} -linear combination

$$\tilde{m} = (k, m) = \sum_{i=0}^d \tilde{\theta}_i(m) \tilde{v}_i,$$

where

$$(\tilde{\theta}_0(m), \tilde{\theta}_1(m), \dots, \tilde{\theta}_d(m)) \in [0, 1]^{d+1}.$$

We set

$$\text{age}(\tilde{m}) = \sum_{i=0}^d \tilde{\theta}_i(m) = k, \quad \text{size}(\tilde{m}) := \text{age}(\tilde{m}) + \text{age}(-\tilde{m}).$$

For any subset $J \subseteq I$ we define

$$G(\Delta_J) := \{\tilde{m} = (\tilde{\theta}_0(m), \tilde{\theta}_1(m), \dots, \tilde{\theta}_d(m)) : \tilde{\theta}_j(m) = 0 \forall j \notin J\}.$$

Theorem 2.7. *Let $\Delta \subset M_{\mathbb{R}}$ be a d -dimensional lattice simplex with $d + 1$ lattice vertices v_i ($i \in I$), and let $J \subseteq I$ be a subset. Then*

$$E(Z_{f,\Delta_J}; u, v) = \frac{(uv - 1)^{|J|-1} - (-1)^{|J|-1}}{uv} + \frac{(-1)^{|J|-2}}{uv} \sum_{0 \neq \tilde{m} \in G(\Delta_J)} v^{\text{size}(\tilde{m})} \left(\frac{u}{v}\right)^{\text{age}(\tilde{m})}.$$

Proof. First of all we note that

$$\psi_i(\Delta) = |\{\tilde{m} \in G(\Delta) : \text{age}(\tilde{m}) = i\}| \quad \forall i \in \{0, 1, \dots, d\},$$

since one can choose a maximal regular sequence $s_0, s_1, \dots, s_d \in S_{\Delta}^{(1)}$ corresponding to $d + 1$ monomials $\tilde{v}_0, \tilde{v}_1, \dots, \tilde{v}_d \in M_{\Delta}$ and obtain a monomial \mathbb{C} -basis of $S_{\Delta}/(s_0, s_1, \dots, s_d)$ that corresponds to elements of $G(\Delta)$. Therefore, by 2.4, we obtain

$$E(Z_{f,\Delta}; u, 1) = \frac{(u - 1)^d - (-1)^d}{u} + \frac{(-1)^{d-1}}{u} \sum_{0 \neq \tilde{m} \in G(\Delta)} u^{\text{age}(\tilde{m})}.$$

Now we use the algorithm of Danilov and Khovanskii for simple polytopes Δ (see [DKh86, Sections 4 and 5]). We apply induction on dimension d and assume that the formula in Theorem 2.7 holds for all proper faces Δ_J of Δ . Consider the Zariski closure $\overline{Z}_{f,\Delta}$ of the affine hypersurface $Z_{f,\Delta}$ in the projective toric variety \mathbb{P}_{Δ} . Then $\overline{Z}_{f,\Delta}$ is quasi-smooth and the Hodge structure in $H_c^*(\overline{Z}_{f,\Delta})$ is pure, i.e., only elements $\tilde{m} \in G(\Delta)$ with the maximal size $\text{size}(\tilde{m}) = d + 1$ can contribute to $H_c^{d-1}(\overline{Z}_{f,\Delta})$. It remains to apply the additivity and the induction hypothesis to the equality

$$E(\overline{Z}_{f,\Delta}; u, v) = \sum_{J \subset I} E(Z_{f,\Delta_J}; u, v).$$

□

Corollary 2.8. *Let $\bar{w} = (w_0, w_1, \dots, w_d) \in \mathbb{Z}_{>0}^{d+1}$ be a well-formed weight vector and let $\Delta \subset M_{\mathbb{R}}$ be the d -dimensional lattice simplex whose vertices v_0, v_1, \dots, v_d generate M and satisfy the relation $\sum_{i=0}^d w_i v_i = 0$. For any subset $J \subset I = \{0, 1, \dots, d\}$ consider the subgroup*

$$G_J := \{l \in \mathbb{Z}/w\mathbb{Z} : \tilde{\theta}_j(l) = 0 \quad \forall j \notin J\} \subseteq G_I = G_0 = \mathbb{Z}/w\mathbb{Z}.$$

Then

$$E(Z_{f,\Delta_J}; u, v) = \frac{(uv - 1)^{|J|-1} - (-1)^{|J|-1}}{uv} + \frac{(-1)^{|J|-2}}{uv} \sum_{0 \neq l \in G_J} v^{\text{size}(l)} \left(\frac{u}{v}\right)^{\text{age}(l)}.$$

Proof. The statement follows immediately from Theorem 2.7. Note that this particular case was considered by Corti and Golyshev in [CG11, Theorem 1.3]. □

3. STRINGY E -FUNCTIONS OF CALABI-YAU HYPERSURFACES

Let $M \cong \mathbb{Z}^d$ be a lattice of rank d , $N := \text{Hom}(M, \mathbb{Z})$ the dual lattice, and $\langle *, * \rangle : M \times N \rightarrow \mathbb{Z}$ the natural pairing. We set $M_{\mathbb{R}} := M \otimes \mathbb{R}$ and $N_{\mathbb{R}} := N \otimes \mathbb{R}$. By a *lattice polytope* $\Delta \subset M_{\mathbb{R}}$ we mean a convex hull of finitely many lattice points

$A \subset M$, i.e., $\Delta = \text{conv}(A)$. For a d -dimensional lattice polytope Δ we consider the piecewise linear function

$$\text{ord}_\Delta : N_{\mathbb{R}} \rightarrow \mathbb{R}, \quad \text{ord}_\Delta(y) := \min_{x \in \Delta} \langle x, y \rangle.$$

The domains of linearity of ord_Δ form a complete rational polyhedral fan Σ_Δ in $N_{\mathbb{R}}$ which is called the *normal fan of Δ* [CLS11]. The normal fan Σ_Δ consists of cones σ_Θ which 1-to-1 correspond to faces $\Theta \preceq \Delta$:

$$\sigma_\Theta := \{y \in N_{\mathbb{R}} : \langle x, y \rangle = \text{ord}_\Delta(y) \quad \forall x \in \Theta\}.$$

Definition 3.1. Let $\Delta \subset M_{\mathbb{R}}$ be a d -dimensional lattice polytope. The subset

$$F(\Delta) := \{x \in M_{\mathbb{R}} : \langle x, n \rangle \geq \text{ord}_\Delta(n) + 1 \quad \forall n \in N \setminus \{0\}\} \subset \Delta.$$

is called the *Fine interior* of Δ .

Remark 3.2. The Fine interior of a lattice polytope Δ is either empty, or a rational polytope of dimension $\leq d$. The convex hull $\text{conv}(\Delta^\circ \cap M)$ is always contained in $F(\Delta)$. If $d = 2$, then $F(\Delta) = \text{conv}(\Delta^\circ \cap M)$ [Bat17, Prop.2.9]. There exist exactly 9 lattice polytopes Δ of dimension $d = 3$ such that $F(\Delta) \neq \emptyset$, but $\Delta^\circ \cap M = \emptyset$ [BKS19, Appendix B].

Theorem 3.3. [Bat17, Theorem 2.23] *A non-degenerate affine hypersurface $Z_\Delta \subset \mathbb{T} := \text{Hom}(M, \mathbb{C}^*)$ defined by a Laurent polynomial $f \in \mathbb{C}[M]$ with the Newton polytope Δ admits a Calabi-Yau compactification X_Δ if and only if the Fine interior of Δ is a single lattice point.*

Definition 3.4. A d -dimensional lattice polytope $\Delta \subset M_{\mathbb{R}}$ containing $0 \in M$ in its interior is called *canonical Fano polytope* if $0 = \Delta^\circ \cap M$, i.e., 0 is the unique interior lattice point in Δ .

Remark 3.5. If $F(\Delta) = \{0\}$, then Δ is a canonical Fano polytope, but the converse is not true if $\dim \Delta \geq 3$. All 3-dimensional canonical Fano polytopes are classified by Kasprzyk [Kas10]. Among them exist exactly 9089 canonical Fano polytopes Δ such that $\dim F(\Delta) \geq 1$ [BKS19].

It is elementary to show the following:

Proposition 3.6. *Let Δ be a d -dimensional canonical Fano polytope. Denote by $\Delta^* \subset N_{\mathbb{R}}$ the dual rational polytope*

$$\Delta^* := \{y \in N_{\mathbb{R}} : \langle x, y \rangle \geq -1 \quad \forall x \in \Delta\}.$$

Then $F(\Delta) = \{0\}$ if and only if

$$[\Delta^*] := \text{conv}(\Delta^* \cap N) \subset N_{\mathbb{R}}$$

is also a d -dimensional canonical Fano polytope.

Corollary 3.7. *Let $\bar{w} \in \mathbb{Z}_{>0}^{d+1}$ be a weight vector. A general hypersurface $X_w \subset \mathbb{P}(\bar{w})$ of degree $w = \sum_{i=0}^d w_i$ is a Calabi-Yau variety if and only if \bar{w} has IP-property.*

Corollary 3.8. *Let $\bar{w} \in \mathbb{Z}_{>0}^{d+1}$ be a weight vector. Consider d -dimensional simplex $\Delta_{\bar{w}}^* = \text{conv}(v_0, v_1, \dots, v_d) \subset M_{\mathbb{R}}$ such that $M = \sum_i \mathbb{Z}v_i$ and $\sum_i w_i v_i = 0$. Then $F(\Delta_{\bar{w}}^*) = \{0\}$ if and only if \bar{w} has IP-property.*

Remark 3.9. Assume that Δ and $[\Delta^*]$ are canonical Fano polytopes, i.e., $F(\Delta) = \{0\}$. There exist many ways to obtain a Calabi-Yau compactifications of a non-degenerate affine hypersurface $Z_\Delta \subset \mathbb{T}$. For constructing a Calabi-Yau compactification one has to choose a finite subset $B \subset [\Delta^*] \cap N$ such that $\nabla_B := \text{conv}(B)$ is a canonical Fano polytope and consider the fan Σ^B in $N_{\mathbb{R}}$ with cones $\mathbb{R}_{\geq 0}\nabla$ spanned by proper faces $\nabla \prec \nabla_B$ and $\{0\} \in N_{\mathbb{R}}$. Then a Calabi-Yau compactification X_Δ^B of Z_Δ is the Zariski closure of Z_Δ in \mathbb{Q} -Gorenstein Fano toric variety \mathbb{P}_B associated with the fan Σ^B [ACG16, Theorem 1].

Theorem 3.10. [Bat17, Theorem 4.10] *Let $\Delta \subset M_{\mathbb{R}}$ be a d -dimensional lattice polytope with $F(\Delta) = \{0\}$. Then the stringy E -function of a Calabi-Yau compactification X_Δ of a non-degenerate affine hypersurface $Z_\Delta \subset \mathbb{T}$ with the Newton polytope Δ can be computed by the formula*

$$E_{\text{str}}(X_\Delta; u, v) = \sum_{\substack{\Theta \preceq \Delta \\ \dim \Theta \geq 1}} E(Z_\Theta; u, v) (uv - 1)^{d - \dim \Theta} \sum_{n \in \sigma_\Theta^\circ} (uv)^{\text{ord}_\Delta(n)},$$

where $E(Z_\Theta; u, v)$ denotes the E -polynomial of the affine hypersurface $Z_\Theta \subset \mathbb{T}_\Theta$ corresponding to a face $\Theta \preceq \Delta$, and σ_Θ° is the relative interior of the $(d - \dim \Theta)$ -dimensional cone $\sigma_\Theta \in \Sigma_\Delta$.

Remark 3.11. We note that the combinatorial formula in Theorem 3.10 depends only on Δ and it does not depend on the choice of a Calabi-Yau compactification X_Δ^B .

We specialize the formula in Theorem 3.10 to the case when

$$\Delta = \Delta_{\bar{w}}^* = \text{conv}(v_0, v_1, \dots, v_d) \subset M_{\mathbb{R}}$$

for a weight vector \bar{w} with IP -property. In this case we need the following standard combinatorial fact about the set $L^{\bar{w}}(k)$ of nonnegative integral solutions $\bar{u} = (u_0, u_1, \dots, u_d) \in \mathbb{Z}_{\geq 0}^{d+1}$ to the linear diophantine equation:

$$\sum_{i=0}^d w_i u_i = \langle \bar{w}, \bar{u} \rangle = kw, \quad k \in \mathbb{Z}_{>0}.$$

Proposition 3.12. *For any subset $J \subseteq I = \{0, 1, \dots, d\}$ denote by $L_J^{\bar{w}}(k)$ the subset in $L^{\bar{w}}(k)$:*

$$L_J^{\bar{w}}(k) := \{\bar{u} \in L^{\bar{w}}(k) : u_j = 0 \Leftrightarrow j \in J\}.$$

Then

$$\sum_{k>0} |L_J^{\bar{w}}(k)| t^{-k} = \left[\prod_{j \notin J} \frac{1}{t^{q_j} - 1} \right]_{\text{int}}.$$

Proof. Consider $d + 1$ variables t_0, t_1, \dots, t_d and associate with a lattice point $\bar{u} \in \mathbb{Z}_{\geq 0}^{d+1}$ the monomial $\mathbf{t}^{\bar{u}} = \prod_i t_i^{u_i}$. A lattice point $\bar{u} \in \mathbb{Z}_{\geq 0}^{d+1}$ belongs to $L^{\bar{w}}(k)$ if and only if by setting $t_i = t^{-q_i}$ ($0 \leq i \leq d$) we get a monomial in t with the integral exponent $-k$, i. e.,

$$\prod_{i=0}^d (t^{-q_i})^{u_i} = t^{-k}.$$

Therefore,

$$\sum_{k>0} |L_{\bar{w}}^{\bar{w}}(k)| t^{-k} = \left[\prod_{j \notin J} (t^{-q_j} + t^{-2q_j} + \dots) \right]_{int} = \left[\prod_{j \notin J} \frac{1}{t^{q_j} - 1} \right]_{int}.$$

□

Corollary 3.13. *Let $\Delta = \Delta_{\bar{w}}^* = \text{conv}(v_0, v_1, \dots, v_d)$ be d -dimensional simplex whose vertices a primitive vectors satisfying the relation $\sum_{i=0}^d w_i v_i = 0$. Then for any $J \subset I$ one has*

$$\sum_{n \in \sigma_J^{\circ}} (uv)^{\text{ord}_{\Delta}(n)} = \left[\prod_{j \notin J} \frac{1}{(uv)^{q_j} - 1} \right]_{int}.$$

Proof. The function ord_{Δ} is linear on the $(d+1-|J|)$ -dimensional simplicial cone σ_J° and it has value (-1) on every vertex of the rational dual simplex $\Delta^* = \Delta_{\bar{w}}^*$. The set of lattice points $n \in \sigma_J^{\circ}$ such that $\text{ord}_{\Delta}(n) = -k$ can be identified with the subset $L_{\bar{w}}^{\bar{w}}(k)$ of nonnegative integral solutions of the linear diophantine equation $\sum_i w_i u_i = w$. Now the statement follows from 3.12. □

Theorem 3.14. *Let $\bar{w} \in \mathbb{Z}^{d+1}$ be a weight vector with IP-property. Then a Calabi-Yau compactification $X_{\bar{w}}^{\vee}$ of a non-degenerate hypersurface $Z_{\bar{w}} \subset \mathbb{T}_{\bar{w}}^*$ defined by a Laurent polynomial with Newton polytope $\Delta_{\bar{w}}^*$ has the following stringy E -function*

$$E_{\text{str}}(X_{\bar{w}}^{\vee}; u, v) = \sum_{\substack{J \subset I \\ |J| \geq 2}} E(Z_{\bar{w}, J}; u, v) (uv - 1)^{d+1-|J|} \left[\prod_{j \notin J} \frac{1}{(uv)^{q_j} - 1} \right]_{int},$$

where

$$E(Z_{\bar{w}, J}; u, v) = \frac{(uv - 1)^{|J|-1} - (-1)^{|J|-1}}{uv} + \frac{(-1)^{|J|-2}}{uv} \sum_{0 \neq l \in G_J} v^{\text{size}(l)} \left(\frac{u}{v} \right)^{\text{age}(l)}.$$

Proof. The statement follows immediately from Theorem 3.10 using 3.13 and 2.8. □

In Example 3.10 below we illustrate the combinatorial formula from Theorem 3.14.

Example 3.15. Take the weight vector $\bar{w} = (1, 5, 12, 18)$. It has IP-property. The affine hypersurface $Z_f \subset (\mathbb{C}^*)^3$ defined by the equation

$$f_{\bar{w}}^0(\mathbf{t}) = \frac{1}{t_1^5 t_2^{12} t_3^{18}} + t_1 + t_2 + t_3 = 0$$

is non-degenerate. The Newton polytope of $f_{\bar{w}}^0(\mathbf{t})$ is a 3-dimensional lattice simplex Δ with $F(\Delta) = \{0\}$ and $\text{Vol}_3(\Delta) = 36 = w$. By 3.8, Z_f admits a Calabi-Yau compactification X_{Δ} . We compute its stringy E -function $E_{\text{str}}(X_{\Delta}; u, v)$ using Theorem 3.10.

Note that the sum over faces $\Delta_J \preceq \Delta$ ($\dim \Delta_J \geq 1$) in the formula 3.14 consists of $1 + 4 + 6 = 11$ terms ($\dim \Delta_J \in \{3, 2, 1\}$). By 2.8, we obtain

$$E(Z_f; u, v) = ((uv)^2 - 3uv + 3) + (u^2 + 10uv + v^2) + 3(u + v) + 6(u + v) + 5 =$$

$$= (uv)^2 + u^2 + 7uv + v^2 + 9(u + v) + 8.$$

The terms corresponding to four 2-dimensional faces $\Delta_J \prec \Delta$ are

$$(uv - w), (uv - 2)\frac{uv - 1}{(uv)^5 - 1}, (uv - 3(u + v) - 7), (uv - 6(u + v) - 7).$$

The terms corresponding to six 1-dimensional faces $\Delta_J \prec \Delta$ are

$$\frac{(1 + 7uv + 7(uv)^2 + 7(uv)^3 + 7(uv)^4 + 7(uv)^5)(uv - 1)}{(uv)^5 - 1}, (1 + 2uv), (1 + uv),$$

$$6, \frac{(1 + (uv)^3)(uv - 1)}{(uv)^5 - 1}, \frac{(1 + (uv)^2 + (uv)^4)(uv - 1)}{(uv)^5 - 1}.$$

The stringy Euler number $E_{\text{str}}(X_\Delta; 1, 1)$ equals

$$36 - \frac{1}{5} - 1 - 12 - 18 + 1 \cdot \frac{36}{5} + 1 \cdot 2 + 1 \cdot 3 + 6 \cdot 1 + 1 \cdot \frac{2}{5} + 1 \cdot \frac{3}{5} = 24.$$

Note that the sum of all four terms with the denominator $(uv)^5 - 1$ equals $1 + 7uv$. Thus, we obtain

$$E_{\text{str}}(X_\Delta; u, v) = (uv)^2 + u^2 + 7uv + v^2 + 9(u + v) + 8 + (7uv + 1) +$$

$$+(uv - 2) + (uv - 3(u + v) - 7) + (uv - 6(u + v) - 7) + 6 + (2 + 3uv) =$$

$$= (uv)^2 + u^2 + 20uv + v^2 + 1.$$

This is E -polynomial of a $K3$ -surface.

4. THE FORMULA OF VAFA

Let $W \in R := \mathbb{C}[z_0, z_1, \dots, z_d]$ be a non-degenerate weighted homogeneous polynomial with weights $w_i := \deg z_i$ ($0 \leq i \leq d$) and $\deg W = w = \sum_{i=0}^d w_i$, i.e.,

$$W(\lambda^{w_0} z_0, \dots, \lambda^{w_d} z_d) = \lambda^w W(z_0, \dots, z_d) \quad \forall \lambda \in \mathbb{C}^*$$

and the common zero of all partial derivatives

$$W'_i := \frac{\partial W}{\partial z_i}, \quad (0 \leq i \leq d)$$

is only the origin $0 \in \mathbb{C}^{n+1}$. Then $d + 1$ homogeneous polynomials W_i of degree $w - w_i$ ($0 \leq i \leq d$) form a regular sequence in the graded ring R and the quotient $\overline{R} := R/\langle W'_0, \dots, W'_d \rangle$ is a graded artinian ring with homogeneous components \overline{R}_m such that

$$P(W, t) = \sum_{m \geq 0} \dim_{\mathbb{C}} \overline{R}_m t^m = \prod_{j=0}^d \left(\frac{1 - t^{w-w_i}}{1 - t^{w_i}} \right)$$

and

$$\mu := \dim_{\mathbb{C}} \overline{R} = \prod_{j=0}^d \left(\frac{w - w_i}{w_i} \right)$$

is the *Milnor number* of W . As above, we set

$$(q_0, q_1, \dots, q_d) := \frac{1}{w}(w_0, w_1, \dots, w_d)$$

and consider the cyclic group $G_0 = \mathbb{Z}/w\mathbb{Z}$ that acts on \mathbb{C}^{n+1} by diagonal matrices

$$\exp(2\pi i\theta(l)) = \text{diag}(e^{2\pi i\theta_0(l)}, \dots, e^{2\pi i\theta_d(l)}), \quad l \in \mathbb{Z}/w\mathbb{Z}.$$

For any element $l \in \mathbb{Z}/l\mathbb{Z}$ we define the subspace

$$U^{(l)} := \{(z_0, z_1, \dots, z_n) \in \mathbb{C}^{n+1} : z_j = 0 \ \forall lq_j \in \mathbb{Z}\}.$$

Then the restriction $W^{(l)}$ of the polynomial W to the subspace $U^{(l)} \subset \mathbb{C}^{d+1}$ is also non-degenerate and therefore we obtain another polynomial

$$P(W^{(l)}, t) := \prod_{lq_j \in \mathbb{Z}} \left(\frac{1 - t^{w-w_j}}{1 - t^{w_j}} \right).$$

An explicit method for computing the orbifold Hodge numbers $h_{\text{orb}}^{p,q}(X_w)$ of quasi-smooth Calabi-Yau hypersurfaces X_w was first suggested by Vafa [Vaf89]. He gave an explicit formula for the orbifold Euler

$$(4.1) \quad \chi_{\text{orb}}(X_w) = \frac{1}{w} \sum_{l,r=0}^{w-1} \prod_{\substack{0 \leq i \leq d \\ lq_i, r q_i \in \mathbb{Z}}} \left(1 - \frac{1}{q_i} \right).$$

The method for computing the orbifold Hodge numbers of X_w was illustrated by Vafa in some examples [Vaf89, p.1182-1183], but an exact combinatorial mathematical formula based on his ideas appeared later in the paper of Kreuzer, Schimmrigk and Skarke [KSS92, Formula (6)](see also Klemm, Lian, Roan and Yau [KLRY98, Formula (3.2)]):

$$(4.2) \quad P(t, \bar{t}) = \sum_{0 \leq l < w} \left[\prod_{\tilde{\theta}_i(l)=0} \frac{1 - (t\bar{t})^{1-q_i}}{1 - (t\bar{t})^{q_i}} \prod_{\tilde{\theta}_i(l) \neq 0} (t\bar{t})^{\frac{1}{2}-q_i} \left(\frac{t}{\bar{t}} \right)^{\tilde{\theta}_i(l) - \frac{1}{2}} \right]_{\text{int}},$$

where

$$\theta(l) = l(q_0, \dots, q_d), \quad \tilde{\theta}(l) = (\tilde{\theta}_0(l), \dots, \tilde{\theta}_d(l)) \in [0, 1]^{d+1},$$

Remark 4.1. It is important to note that the polynomial $P(t, \bar{t}) = \sum_{p,q} h^{p,q} t^p \bar{t}^q$ in the formula (4.2) has nonnegative integral coefficients $h^{p,q}$ which are not the orbifold Hodge numbers $h_{\text{orb}}^{p,q}(X_w)$ of the quasi-smooth Calabi-Yau hypersurface, but the Hodge numbers $h^{d-1-p,q}$ of its mirror. This can be seen more explicitly in examples below. In particular, for Calabi-Yau 3-folds X one obtains

$$P(t, \bar{t}) = (1 + t^3)(1 + \bar{t}^3) + h^{2,1}(X)(t\bar{t} + (t\bar{t})^2) + h^{1,1}(X)(t(\bar{t})^2 + t^2\bar{t}).$$

Example 4.2. Let $d = 4$, $w = 5$ and $\bar{w} := (1, 1, 1, 1, 1)$. Then for $l = 0$ we have

$$P_0(t, \bar{t}) = \left[\prod_{j=0}^4 \left(\frac{1 - (t\bar{t})^{4/5}}{1 - (t\bar{t})^{1/5}} \right) \right]_{\text{int}} = [(1 + (t\bar{t})^{1/5} + (t\bar{t})^{2/5} + (t\bar{t})^{3/5} + (t\bar{t})^4)^5]_{\text{int}} =$$

$$1 + 101t\bar{t} + 101(t\bar{t})^2 + (t\bar{t})^3.$$

For the remaining values $l \in \{1, 2, 3, 4\}$, one obtains four monomials

$$t^3, t^2\bar{t}, t\bar{t}^2, \bar{t}^3.$$

Since a Calabi-Yau quintic 3-fold $X_5 \subset P^4$ has Hodge numbers $h^{1,1}(X_5) = 1$ and $h^{2,1}(X_5) = 101$, the formula (4.2) produces the Hodge numbers of the mirror of X_5 .

We need a modified version of the formula (4.2):

Proposition 4.3. *Let*

$$Q(u, v) := \left[\sum_{0 \leq l < w} \left(\prod_{\theta_i(l) \in \mathbb{Z}} \frac{1 - (uv)^{1-q_i}}{1 - (uv)^{q_i}} \right) \cdot \prod_{\tilde{\theta}_i(l) \notin \mathbb{Z}} (uv)^{\frac{1}{2}-q_i} \left(\frac{u}{v} \right)^{\tilde{\theta}_i(l) - \frac{1}{2}} \right]_{int}.$$

Then one has

$$Q(u, v) = \frac{1}{uv} \sum_{0 \leq l < w} \left[\prod_{\tilde{\theta}_i(l) \in \mathbb{Z}} \frac{(uv)^{q_i} - uv}{1 - (uv)^{q_i}} \right]_{int} \cdot v^{\text{size}(l)} \left(\frac{u}{v} \right)^{\text{age}(l)}.$$

Proof. By $\sum_{i=0}^d q_i = 1$, we have

$$\prod_{\tilde{\theta}_i(l) \notin \mathbb{Z}} (uv)^{-q_i} = \frac{1}{uv} \prod_{\tilde{\theta}_i(l) \in \mathbb{Z}} (uv)^{q_i}.$$

Using

$$\prod_{\tilde{\theta}_i(l) \notin \mathbb{Z}} (uv)^{1/2-q_i} = (uv)^{\frac{\text{size}(l)}{2}} \prod_{\tilde{\theta}_i(l) \notin \mathbb{Z}} (uv)^{-q_i},$$

we obtain

$$\begin{aligned} Q(u, v) &= \left[\sum_{0 \leq l < w} \left(\prod_{\theta_i(l) \in \mathbb{Z}} \frac{1 - (uv)^{1-q_i}}{1 - (uv)^{q_i}} \right) \cdot \prod_{\tilde{\theta}_i(l) \notin \mathbb{Z}} (uv)^{\frac{1}{2}-q_i} \left(\frac{u}{v} \right)^{\tilde{\theta}_i(l) - \frac{1}{2}} \right]_{int} = \\ &= \sum_{0 \leq l \leq w-1} \frac{1}{uv} \left[\left(\prod_{\tilde{\theta}_i(l) \in \mathbb{Z}} \frac{(uv)^{q_i} - uv}{1 - (uv)^{q_i}} \right) (uv)^{\frac{\text{size}(l)}{2}} \left(\frac{u}{v} \right)^{-\frac{\text{size}(l)}{2}} \cdot \prod_{\tilde{\theta}_i(l) \notin \mathbb{Z}} \left(\frac{u}{v} \right)^{\tilde{\theta}_i(l)} \right]_{int} = \\ &= \frac{1}{uv} \sum_{0 \leq l < w} \left[\prod_{\tilde{\theta}_i(l) \in \mathbb{Z}} \frac{(uv)^{q_i} - uv}{1 - (uv)^{q_i}} \right]_{int} \cdot \left(v^{\text{size}(l)} \prod_{\tilde{\theta}_i(l) \notin \mathbb{Z}} \left(\frac{u}{v} \right)^{\tilde{\theta}_i(l)} \right) = \\ &= \frac{1}{uv} \sum_{0 \leq l < w} \left[\prod_{\tilde{\theta}_i(l) \in \mathbb{Z}} \frac{(uv)^{q_i} - uv}{1 - (uv)^{q_i}} \right]_{int} \cdot v^{\text{size}(l)} \left(\frac{u}{v} \right)^{\text{age}(l)}. \end{aligned}$$

□

Corollary 4.4. *Assum that $\bar{w} \in \mathbb{Z}_{>0}^{d+1}$ is a transverse weight vector and $X_w \subset \mathbb{P}(\bar{w})$ is a quasi-smooth Calabi-Yau hypersurface. Then for the mirror stringy E -polynomial of X_w we have*

$$(-u)^{d-1} E_{\text{orb}}(X_w; u^{-1}, v) = \frac{1}{uv} \sum_{0 \leq l < w} \left[\prod_{\tilde{\theta}_i(l) \in \mathbb{Z}} \frac{(uv)^{q_i} - uv}{1 - (uv)^{q_i}} \right]_{int} \cdot (-v)^{\text{size}(l)} \left(\frac{u}{v} \right)^{\text{age}(l)}.$$

Example 4.5. Consider the transverse weight vector $\bar{w} = (1, 5, 12, 18)$. Then $w = \sum_i w_i = 36$. There are twelve elements l in $\mathbb{Z}/36\mathbb{Z}$ acting on \mathbb{C}^4 with size 4: one element of age 1, ten elements of age 2, and one element of age 3. This gives polynomial $u^2 + 10uv + v^2$. The element $l = 0$ determines the term

$$\left[\left(\frac{1 - (uv)^{\frac{35}{36}}}{1 - (uv)^{\frac{1}{36}}} \right) \cdot \left(\frac{1 - (uv)^{\frac{31}{36}}}{1 - (uv)^{\frac{5}{36}}} \right) \left(\frac{1 - (uv)^{\frac{2}{3}}}{1 - (uv)^{\frac{1}{3}}} \right) \left(\frac{1 - (uv)^{\frac{1}{2}}}{1 - (uv)^{\frac{1}{2}}} \right) \right]_{int} = 1 + 10uv + (uv)^2.$$

There exist five elements $l \in \mathbb{Z}/36\mathbb{Z}$ of size 2 and age 1 with the corresponding term

$$\left[(uv)^{\frac{5}{6}} \left(\frac{1 - (uv)^{\frac{2}{3}}}{1 - (uv)^{\frac{1}{3}}} \right) \left(\frac{1 - (uv)^{\frac{1}{2}}}{1 - (uv)^{\frac{1}{2}}} \right) \right]_{int} = \left[(uv)^{\frac{5}{6}} (1 + (uv)^{\frac{1}{3}}) \right]_{int} = 0.$$

There exist twelve elements $l \in \mathbb{Z}/36\mathbb{Z}$ of size 3 and age 1 or 2 with the corresponding term

$$\left[(uv)^{\frac{1}{2}} \left(\frac{1 - (uv)^{\frac{1}{2}}}{1 - (uv)^{\frac{1}{2}}} \right) \right]_{int} \cdot (-v)u^{\text{age}(l)-1} = \left[(uv)^{\frac{1}{2}} \cdot 1 \right]_{int} \cdot (-v)u^{\text{age}(l)-1} = 0.$$

There exist six elements $l \in \mathbb{Z}/36\mathbb{Z}$ of size 3 and age 1 or 2 with the corresponding term

$$\left[(uv)^{\frac{1}{3}} \left(\frac{1 - (uv)^{\frac{2}{3}}}{1 - (uv)^{\frac{1}{3}}} \right) \right]_{int} \cdot (-v)u^{\text{age}(l)-1} = \left[(uv)^{\frac{1}{3}} (1 + (uv)^{\frac{1}{3}}) \right]_{int} \cdot (-v)u^{\text{age}(l)-1} = 0.$$

By summing all 36 terms, we obtain

$$(-u)^2 E_{\text{orb}}(X_w; u^{-1}, v) = 1 + 10uv + (uv)^2 + u^2 + 10uv + v^2 = 1 + u^2 + 20uv + v^2 + (uv)^2.$$

Example 4.6. Take the transverse weight vector $\bar{w} = (1, 1, 2, 2, 2)$. Then $w = 8$ and we obtain the action of $\mathbb{Z}/8\mathbb{Z}$ on \mathbb{C}^5 with charges $(1/8, 1/8, 1/4, 1/4, 1/4)$. There exist 6 elements $l \in \mathbb{Z}/8\mathbb{Z}$ of size 5 (one of age 1, two of age 2, two of age 3 and one of age 4). This gives a polynomial $-v^3 - 2v^2u - 2vu^2 - u^3$. The element $l = 0$ in $\mathbb{Z}/8\mathbb{Z}$ determines the term

$$\left[\left(\frac{1 - (uv)^{\frac{7}{8}}}{1 - (uv)^{\frac{1}{8}}} \right)^2 \cdot \left(\frac{1 - (uv)^{\frac{3}{4}}}{1 - (uv)^{\frac{1}{4}}} \right)^3 \right]_{int} = 1 + 83uv + 83(uv)^2 + (uv)^3.$$

It remains to consider the element of order 2 in $l \in \mathbb{Z}/8\mathbb{Z}$. It has size 2 and age 1. By putting $t = (uv)^{\frac{1}{4}}$ in the equality

$$(1 + t + t^2)^3 = 1 + 3t + 6t^2 + 7t^3 + 6t^4 + 3t^5 + t^6,$$

we obtain the corresponding term

$$\left[(uv)^{\frac{3}{4}} \cdot \left(\frac{1 - (uv)^{\frac{3}{4}}}{1 - (uv)^{\frac{1}{4}}} \right)^3 \right]_{int} = 3uv + 3(uv)^2.$$

Therefore, we obtain

$$(-u)^3 E_{\text{orb}}(X_w; u^{-1}, v) = 1 + 86uv - v^3 - 2v^2u - 2vu^2 - u^3 + 86(uv)^2 + (uv)^3.$$

In particular, the Euler number of $X_8 \subset \mathbb{P}(1, 1, 2, 2, 2)$ equals -168 .

5. THE PROOF

Proof of Theorem 2.1 In order to prove the required equality

$$E_{\text{str}}(X_{\bar{w}}^{\vee}; u, v) = \frac{1}{uv} \sum_{0 \leq l < w} \left[\prod_{\tilde{\theta}_i(l)=0} \frac{(uv)^{q_i} - uv}{1 - (uv)^{q_i}} \right]_{\text{int}} \cdot (-u)^{\text{size}(l)} \left(\frac{v}{u} \right)^{\text{age}(l)}$$

we split its left hand side and right hand side into a sum over elements $l \in G_0 = \mathbb{Z}/w\mathbb{Z}$. We set

$$P_{\bar{w}}^{(l)}(u, v) := \frac{1}{uv} \left[\prod_{\tilde{\theta}_i(l)=0} \frac{(uv)^{q_i} - uv}{1 - (uv)^{q_i}} \right]_{\text{int}} \cdot (-u)^{\text{size}(l)} \left(\frac{v}{u} \right)^{\text{age}(l)}, \quad l \in \mathbb{Z}/w\mathbb{Z},$$

and use the formula in Theorem 3.14 to write

$$E_{\text{str}}(X_{\bar{w}}^{\vee}; u, v) = \sum_{0 \leq l < w} E_{\text{str}}^{(l)}(X_{\bar{w}}^{\vee}; u, v),$$

where

$$E_{\text{str}}^{(0)}(X_{\bar{w}}^{\vee}; u, v) := \sum_{\substack{J \subseteq I \\ |J| \geq 2}} \frac{(uv - 1)^{|J|-1} - (-1)^{|J|-1}}{uv} \cdot (uv - 1)^{d+1-|J|} \left[\prod_{j \notin J} \frac{1}{(uv)^{q_j} - 1} \right]_{\text{int}}$$

and

$$E_{\text{str}}^{(l)}(X_{\bar{w}}^{\vee}; u, v) := \sum_{\substack{J \subseteq I \\ l \in \bar{G}_J}} \frac{(-1)^{|J|} (uv)^{\text{size}(l)}}{uv} \left(\frac{u}{v} \right)^{\text{age}(l)} \cdot (uv - 1)^{d+1-|J|} \left[\prod_{j \notin J} \frac{1}{(uv)^{q_j} - 1} \right]_{\text{int}}, \quad l \neq 0.$$

Case 1. Let $l = 0$. Then $\text{age}(l) = \text{size}(l) = 0$ and the untwisted term equals

$$\begin{aligned} P_{\bar{w}}^{(0)}(u, v) &= \frac{1}{uv} \left[\prod_{i=0}^d \frac{(uv)^{q_i} - uv}{1 - (uv)^{q_i}} \right]_{\text{int}} = \frac{1}{uv} \left[\prod_{i=0}^d \left(\frac{uv - 1}{(uv)^{q_i} - 1} - 1 \right) \right]_{\text{int}} = \\ &= \frac{1}{uv} \sum_{\emptyset \subseteq J \subseteq I} (uv - 1)^{d+1-|J|} (-1)^{|J|} \left[\prod_{j \notin J} \frac{1}{(uv)^{q_j} - 1} \right]_{\text{int}} = \\ &= \frac{(uv - 1)^{d+1}}{uv} \left[\prod_{i=0}^d \frac{1}{(uv)^{q_i} - 1} \right]_{\text{int}} - \frac{(uv - 1)^d}{uv} \sum_{i=0}^d \left[\prod_{j \neq i} \frac{1}{(uv)^{q_j} - 1} \right]_{\text{int}} + \\ &\quad + \frac{1}{uv} \sum_{\substack{J \subseteq I \\ |J| \geq 2}} (uv - 1)^{d+1-|J|} (-1)^{|J|} \left[\prod_{j \notin J} \frac{1}{(uv)^{q_j} - 1} \right]_{\text{int}}. \end{aligned}$$

On the other hand, we have

$$E_{\text{str}}^{(0)}(X_{\bar{w}}^{\vee}; u, v) = \sum_{\substack{J \subseteq I \\ |J| \geq 2}} \frac{(uv - 1)^{|J|-1} - (-1)^{|J|-1}}{uv} \cdot (uv - 1)^{d+1-|J|} \left[\prod_{j \notin J} \frac{1}{(uv)^{q_j} - 1} \right]_{\text{int}} =$$

$$\frac{(uv-1)^{d+1}}{uv} \sum_{\substack{J \subseteq I \\ |J| \geq 2}} \left[\prod_{j \notin J} \frac{1}{(uv)^{q_j} - 1} \right]_{int} + \frac{(uv-1)^{d+1-|J|}(-1)^{|J|}}{uv} \sum_{\substack{J \subseteq I \\ |J| \geq 2}} \left[\prod_{j \notin J} \frac{1}{(uv)^{q_j} - 1} \right]_{int}$$

Now we use the equality $\sum_{i=0}^d q_i = 1$ to obtain $uv = \prod_{i=0}^d (uv)^{q_i}$ and apply it in

$$\begin{aligned} uv \left[\prod_{i=0}^d \left(\frac{1}{(uv)^{q_i} - 1} \right) \right]_{int} &= \left[uv \prod_{i=0}^d \left(\frac{1}{(uv)^{q_i} - 1} \right) \right]_{int} = \left[\prod_{i=0}^d \left(\frac{(uv)^{q_i}}{(uv)^{q_i} - 1} \right) \right]_{int} = \\ &= \left[\prod_{i=0}^d \left(\frac{1}{(uv)^{q_i} - 1} + 1 \right) \right]_{int} = \sum_{\emptyset \subseteq J \subseteq I} \left[\prod_{j \notin J} \frac{1}{(uv)^{q_j} - 1} \right]_{int}. \end{aligned}$$

Therefore

$$\sum_{\substack{J \subseteq I \\ |J| \geq 2}} \left[\prod_{j \notin J} \frac{1}{(uv)^{q_j} - 1} \right]_{int} = (uv-1) \left[\prod_{i=0}^d \frac{1}{(uv)^{q_i} - 1} \right]_{int} - \sum_{i=0}^d \left[\prod_{j \neq i} \frac{1}{(uv)^{q_j} - 1} \right]_{int}.$$

If we multiply the last equality by $(uv-1)^d/uv$ and add to both sides of the obtained equality the sum

$$\frac{(uv-1)^{d+1-|J|}(-1)^{|J|}}{uv} \sum_{\substack{J \subseteq I \\ |J| \geq 2}} \left[\prod_{j \notin J} \frac{1}{(uv)^{q_j} - 1} \right]_{int},$$

Then the left hand side becomes $E_{\text{str}}^{(0)}(X_{\Delta}; u, v)$ and the right hand side becomes $P^{(0)}(u, v)$. Thus,

$$E_{\text{str}}^{(0)}(X_{\Delta}; u, v) = P_{\bar{w}}^{(0)}(u, v).$$

Case 2. Let $l \neq 0$. We write $I = \{0, 1, \dots, d\}$ as disjoint union of two subsets:

$$I = J(l) \cup \overline{J(l)}, \quad J(l) := \{j \in I : \tilde{\theta}_j(l) \neq 0\}, \quad \overline{J(l)} := \{j \in I : \tilde{\theta}_j(l) = 0\}.$$

Now we have

$$\begin{aligned} P^{(l)}(X_w; u, v) &= \frac{1}{uv} \left[\prod_{j \in \overline{J(l)}} \frac{(uv) - (uv)^{q_j}}{(uv)^{q_j} - 1} \right]_{int} (-v)^{\text{size}(l)} \left(\frac{u}{v} \right)^{\text{age}(l)} = \\ &= \frac{1}{uv} \left[\prod_{j \in \overline{J(l)}} \left(\frac{(uv) - 1}{(uv)^{q_j} - 1} - 1 \right) \right]_{int} (-v)^{\text{size}(l)} \left(\frac{u}{v} \right)^{\text{age}(l)} = \\ &= \frac{1}{uv} \sum_{\overline{J} \subseteq \overline{J(l)}} (uv-1)^{|\overline{J}|} (-1)^{|\overline{J(l)}| - |\overline{J}|} \left[\prod_{j \in \overline{J}} \frac{1}{(uv)^{q_j} - 1} \right]_{int} \cdot (-v)^{\text{size}(l)} \left(\frac{u}{v} \right)^{\text{age}(l)}. \end{aligned}$$

On the other hand, we have

$$E_{\text{str}}^{(l)}(X_{\bar{w}}^{\vee}; u, v) = \sum_{\substack{J \subseteq I \\ l \in G_J}} \frac{(-1)^{|J|} v^{\text{size}(l)}}{uv} \left(\frac{u}{v} \right)^{\text{age}(l)} \cdot (uv-1)^{d+1-|J|} \left[\prod_{j \notin J} \frac{1}{(uv)^{q_j} - 1} \right]_{int}.$$

Note that $l \in G_J \Leftrightarrow J(l) \subseteq J$ and $|\overline{J(l)}| = d + 1 - \text{size}(l)$. Therefore, we obtain for $E_{\text{str}}^{(l)}(X_{\overline{w}}^{\vee}; u, v)$ and $P^{(l)}(X_w; u, v)$ the same sums: the first sum runs over $J \supseteq J(l)$, the second sum runs over the complements $\overline{J} \subseteq \overline{J(l)}$. Thus, we obtain

$$E_{\text{str}}^{(l)}(X_{\overline{w}}^{\vee}; u, v) = P^{(l)}(X_w; u, v) \quad \forall l \in \mathbb{Z}/w\mathbb{Z}.$$

If \overline{w} is transverse, by 4.4, this implies

$$E_{\text{str}}(X_{\overline{w}}^{\vee}; u, v) = (-u)^{d-1} E_{\text{orb}}(X_w; u^{-1}, v)$$

as it is expected in mirror symmetry. \square

Theorem 1.2 can be used for constructing examples of Calabi-Yau varieties X which do not have mirrors:

Example 5.1. Take the weight vector with *IP*-property $\overline{w} = (1, 1, 2, 4, 5)$. Then $w = 13$. We note that this weight vector is not transverse. By Theorem 1.2, we can compute the stringy E -function of a Calabi-Yau compactification $X_{\overline{w}}^*$ of the affine hypersurface $Z_{\overline{w}} \subset (\mathbb{C}^*)^5$ defined by the Laurent polynomial

$$f_{\overline{w}}^0(\mathbf{t}) = \frac{1}{t_1 t_2^2 t_3^4 t_4^5} + t_1 + t_2 + t_3 + t_4.$$

All nonzero elements $l \in \mathbb{Z}/13\mathbb{Z}$ have size 5 and determine twelve terms that give rise to polynomial $-v^3 - 5uv^2 - 5u^2v - u^3$. The untwisted term ($l = 0$)

$$\left[\left(\frac{1 - (uv)^{\frac{12}{13}}}{1 - (uv)^{\frac{1}{13}}} \right)^2 \cdot \left(\frac{1 - (uv)^{\frac{11}{13}}}{1 - (uv)^{\frac{2}{13}}} \right) \cdot \left(\frac{1 - (uv)^{\frac{9}{13}}}{1 - (uv)^{\frac{4}{13}}} \right) \cdot \left(\frac{1 - (uv)^{\frac{8}{13}}}{1 - (uv)^{\frac{5}{13}}} \right) \right]_{\text{int}}$$

is not a polynomial, because its value for $u = v = 1$ is $\frac{1092}{5} \notin \mathbb{Z}$ (cf. [BS20, Example 1.13]). This is explained by existence of terminal singularities on minimal models of $Z_{\overline{w}}$ [DR01]. Therefore, $E_{\text{str}}(X_{\overline{w}}^*; u, v)$ is not a polynomial, and $X_{\overline{w}}^*$ has no mirror.

REFERENCES

- [ACG16] M. Artebani, P. Comparin, R. Guillot, *Families of Calabi-Yau hypersurfaces in \mathbb{Q} -Fano toric varieties*, J. Math. Pures Appl. (9) **106** (2016), no. 2, 319–341.
- [Bat93] V. V. Batyrev, *Variations of the mixed Hodge structure of affine hypersurfaces in algebraic tori*, Duke Math. J. Volume 69, Number 2 (1993), 349–409.
- [Bat94] V. V. Batyrev, *Dual Polyhedra and Mirror Symmetry for Calabi-Yau Hypersurfaces in Toric Varieties*, Journal of Algebraic Geometry **3** (1994), no. 3, 493–535.
- [BB96] V. V. Batyrev and L. A. Borisov, *Mirror duality and string-theoretic Hodge numbers*, Invent. Math. 126 (1) (1996) 183–203,
- [Bat98] V. V. Batyrev, *Stringy Hodge numbers of varieties with Gorenstein canonical singularities*, Integrable systems and algebraic geometry (Kobe/Kyoto 1997), World Sci. Publ., River Edge, NJ (1998), 1–31.
- [Bat17] V. V. Batyrev, *The stringy Euler number of Calabi-Yau hypersurfaces in toric varieties and the Mavlyutov duality*, Pure Appl. Math. Q. **13** (2017), no. 1, 1–47.
- [BKS19] V. V. Batyrev, A. M. Kasprzyk, and K. Schaller, *On the Fine Interior of Three-dimensional Canonical Fano Polytopes*, arXiv:1911.12048.
- [BS20] V. V. Batyrev and K. Schaller, *Mirror symmetry for quasi-smooth Calabi-Yau hypersurfaces in weighted projective spaces*, arXiv:2006.04465
- [BH93] P. Berglund, T. Hübsch, *A generalized construction of mirror manifolds*, Nucl. Phys. **B 393** (1993), 377–391.

- [Bo13] L. A. Borisov, *Berglund–Hübsch mirror symmetry via vertex algebras*, Commun. Math. Phys. **320** (1) (2013) 73–99.
- [CR04] W. Chen and Y. Ruan, *A new cohomology theory of orbifold*, Comm. Math. Phys. **248** (2004), no. 1, 1–31.
- [CR18] E. Clader and Y. Ruan, *Mirror symmetry constructions*, B-model Gromov-Witten theory, Trends Math., Birkhäuser/Springer, Cham, 2018, pp. 1–77.
- [CR10] A. Chiodo and Y. Ruan, *LG/CY correspondence: the state space isomorphism*, Adv. Math. **227** (2010).
- [CG11] A. Corti, V. Golyshev, *Hypergeometric equations and weighted projective spaces*, Sci. China Math. **54** (8) (2011) 1577–1590.
- [CLS11] D. A. Cox, J. B. Little and H. K. Schenck, *Toric varieties*, Graduate Studies in Mathematics, 124, Amer. Math. Soc., Providence, RI, 2011.
- [DR01] D. Dais, M. Roczen, *On the String-Theoretic Euler Number of 3-dimensional A-D-E Singularities*, Advances in Geometry **1** (2001), 373–426.
- [DKh86] V. I. Danilov, A. G. Khovanski, *Newton polyhedra and an algorithm for calculating Hodge-Deligne numbers*, Izv. Akad. Nauk SSSR Ser. Mat. **50** (1986), 925–945.
- [ET13] W. Ebeling, A. Takahashi, *Variance of the exponents of orbifold Landau-Ginzburg models*, Math. Res. Lett. **20**(1) (2013), 51–65.
- [EGZT16] W. Ebeling, S.M. Gusein-Zade, A. Takahashi, *Orbifold E-function of dual invertible polynomials*, J. Geom. Phys. **106** (2016), 184–191.
- [GKZ94] I.M. Gelfand, M. M. Kapranov, A. V. Zelevinsky. *Discriminants, Resultants and Multidimensional Determinants*, 1994.
- [Kas10] A. M. Kasprzyk, *Canonical toric Fano threefolds*, Canadian Journal of Mathematics **62** (2010), no. 6, 1293–1309.
- [KLR98] A. Klemm, B. Lian, S.-S. Roan and S.-T. Yau *Calabi-Yau fourfolds for M- and F-Theory compactifications*, Nucl. Phys. **B 518** (1998), 515–574.
- [Kra10] M. Krawitz, *FJRW rings and Landau-Ginzburg mirror symmetry*, ProQuest LLC, Ann Arbor, MI, 2010, Thesis (Ph.D.)—University of Michigan.
- [KSS92] M. Kreuzer, R. Schimmrigk, K. Skarke, *Abelian Landau-Ginzburg orbifolds and mirror symmetry*, Nucl. Phys. **B 372** (1992), 61–86.
- [Sa20] M. Saito, *Intersection complexes of toric varieties and mixed Hodge modules*, arXiv:2006.04081.
- [Ska96] H. Skarke, *Weight systems for toric Calabi-Yau varieties and reflexivity of Newton polyhedra*, Modern Phys. Lett. A **11** (1996), no. 20, 1637–1652.
- [Vaf89] C. Vafa, *String vacua and orbifolded LG models*, Modern Phys. Lett. A **4** (1989), no. 12, 1169–1185.
- [Yas04] T. Yasuda, *Twisted jets, motivic measure and orbifold cohomology*, Comp. Math. **140** (2004), 396–422.

FACHBEREICH MATHEMATIK, UNIVERSITÄT TÜBINGEN, AUF DER MORGENSTELLE 10, 72076 TÜBINGEN, GERMANY

E-mail address: batyrev@math.uni-tuebingen.de