SYSTEMS OF PARAMETERS AND THE COHEN–MACAULAY PROPERTY

JÜRGEN HERZOG AND SOMAYEH MORADI

ABSTRACT. We recall a numerical criteria for Cohen–Macaulayness related to system of parameters, and introduce monomial ideals of König type which include the edge ideals of König graphs. We show that a monomial ideal is of König type if and only if its corresponding residue class ring admits a system of parameters whose elements are of the form $x_i - x_j$. This provides an algebraic characterization of König graphs. We use this special parameter systems for the study of the edge ideal of König graphs and the study of the order complex of a certain family of posets. Finally, for any simplicial complex Δ we introduce a system of parameters for $K[\Delta]$ with a universal construction principle, independent of the base field and only dependent on the faces of Δ . This system of parameters is an efficient tool to test Cohen–Macaulayness of the Stanley–Reisner ring of a simplicial complex.

INTRODUCTION

Systems of parameters play an important role in dimension theory. As a consequence of Krull's generalized principal ideal theorem it can be seen that in a Noetherian local ring (R, \mathfrak{m}) with dim R = d, there exist elements $f_1, \ldots, f_d \in \mathfrak{m}$ with dim $R/(f_1, \ldots, f_d) = 0$. Such a sequence of elements of R is called a system of parameters, or sop for short. A similar statement holds for standard graded K-algebras with K a field. In our applications we mainly consider such algebras.

One of the central problems in Combinatorial Commutative Algebra is to show that a certain K-algebra attached to a combinatorial object is Cohen-Macaulay. Usually the Cohen-Macaulay property has a nice combinatorial interpretation. In the case that the defining ideal of the algebra is a monomial ideal, Hochster's formula [10] and its extension by Takayama [17] are powerful tools to investigate the homological properties of the algebra. In the case that the defining ideal is a binomial prime ideal, one may use the squarefree divisor complex [3] or one may use Gröbner basis theory to reduce the problem to the case of monomial ideals.

In this paper we propose another approach which is based on the basic fact that R is Cohen–Macaulay if and only if one (equivalently all) of the sop's of R form(s) a regular sequence. This approach confronts us with two problems. The first problem is to find a suitable sop, the second is to decide whether the given sop forms a regular sequence. Regarding the first problem, Stanley [15, Proposition 4.3] finds an explicit special sop for the Stanley–Reisner ring of any balanced simplicial complex. This

²⁰¹⁰ Mathematics Subject Classification. Primary 13H10; Secondary 05C25.

Key words and phrases. System of parameters, Cohen-Macaulay, König graph.

also includes the order complexes. In the cases considered here we also use special sop's.

In the first section of this paper however we first deal with the second problem. Based on results of Serre [13], see also [2, Theorem 4.6.10] one has a numerical condition for when a sop is a regular sequence. Indeed, let $f_1, \ldots, f_d \in \mathfrak{m}$ be a sop of Rand let $\overline{R} = R/(f_1, \ldots, f_d)$. Then, denoting by e(M) the multiplicity of an R-module M, one has $e(\overline{R}) \ge e(R)$, and if $e(\overline{R}) = e(R)$ then f_1, \ldots, f_d is a regular sequence (equivalently, R is Cohen-Macaulay). Moreover, if (f_1, \ldots, f_d) is a reduction ideal of \mathfrak{m} and f_1, \ldots, f_d is a regular sequence, then $e(\overline{R}) = e(R)$, see Proposition 1.1. There is also a graded version of this criterion, see Proposition 1.2. In the case that R is a standard graded K-algebra and the sop f_1, \ldots, f_d is homogeneous with deg $f_i = a_i$, then this sop is a regular sequence if and only if $e(\overline{R}) = a_1a_2 \cdots a_de(R)$. By a lack of good references we provided the detailed proofs of these results.

In Proposition 1.3 we give in the graded case a measure for the difference $e(\overline{R}) - e(R)$. As a consequence we obtain in Corollary 1.4 the result that if the sop f_1, \ldots, f_d is a superficial sequence, and $R/(f_1, \ldots, f_r)$ is Cohen–Macaulay for some r < d, then R is Cohen–Macaulay.

In Section 2 we study a class of posets and their order complexes as well as König graphs by means of sop's. We consider a poset P which as a set is the disjoint union of two sets C_1 and C_2 , where $C_1 : x_1 < x_2 < \cdots < x_n$ and $C_2 : y_1 < y_2 < \cdots < y_n$ are maximal chains in P. For such a poset the sequence $x_1 - y_1, \ldots, x_n - y_n$ is a sop of the Stanley–Reisner ring $K[\Delta(P)]$, where $\Delta(P)$ denotes the order complex of P. The covering relations $x_i < y_j$ in P we call the diagonals of P. The Cohen– Macaulay property of $K[\Delta(P)]$ can be expressed in terms of the diagonals of P. Indeed, in Theorem 2.1 it is shown $K[\Delta(P)]$ is Cohen–Macaulay if and only if it is pure shellable, and that this is equivalent to the condition that the diagonals of Psatisfy the following conditions: (i) if $x_i < y_j$ or $y_i < x_j$, then j = i + 1, and (ii) $\{x_i, y_{i+1}\} \notin \Delta(P)$ implies that $\{x_{i+1}, y_i\} \in \Delta(P)$. In a similar fashion it can be characterized when $I_{\Delta(P)}$ has a linear resolution, see Proposition 2.2.

Note that $I_{\Delta(P)}$ may be viewed as the edge ideal I(G) of a suitable bipartite graph G. So the question arises for which graphs G can we find a sop f_1, \ldots, f_d of K[V(G)]/I(G), where each f_i is just a difference of two variables, like we have it for $K[\Delta(P)]$. The advantage of such sop's is that after reduction they preserve the monomial structure and just identify vertices. The surprising answer to the above question is that a graph G admits such a special sop if and only if G is a König graph. In fact, this is a corollary of a more general theorem. Let $I \subset S$ be a monomial ideal in the polynomial ring $S = K[x_1, \ldots, x_n]$ over the field K in n variables. We denote by m-grade(I) the maximal length of a regular sequence of monomials in I, and call this number the monomial grade of I. One has m-grade(I) \leq grade(I) = height(I). We call I a monomial ideal of König type if $I \neq 0$ and m-grade(I) = height(I). The naming is justified by the fact that if I = I(G) for some graph G, then height(I) = $\tau(G)$ and m-grade(I) = $\nu(G)$, so that the edge ideal of a graph G is a monomial ideal of König type if and only if G is a König graph. Now our Theorem 2.3 says that a monomial ideal $I \subset S = K[x_1, \ldots, x_n]$ is a monomial ideal of König type if and only if S/I admits a sop f_1, \ldots, f_d , where each f_k is of the form $x_i - x_j$ for suitable *i* and *j*.

Applied to graphs this result reads a follows: let G be a graph without isolated vertices, S = K[V(G)] and for any edge $e = \{x, y\} \in E(G)$, let $f_e = x - y$ be an element in S. Then G is a König graph if and only if there exists a subset $\{e_1, \ldots, e_d\}$ of edges of G such that f_{e_1}, \ldots, f_{e_d} is a sop for R = S/I(G). This sop has the nice property that $\operatorname{reg}(R/(f_{e_1}, \ldots, f_{e_d})R) \leq \operatorname{reg}(R)$, as shown in Theorem 2.6.

For a graph G, we denote by $\operatorname{mi}(G)$ the number of maximal independent sets of G. It is an important problem in graph theory to give upper bounds for $\operatorname{mi}(G)$. For a König graph it was shown in [9, Corollary 3.4] that $2^{\nu(G)}$ is an upper bound for $\operatorname{mi}(G)$ where $\nu(G)$ denotes the maximum size of matchings of G, and in [1, Theorem 1] it was proved that $\operatorname{mi}(G) \leq M(G) + 1$, where M(G) is the number of induced matchings in G. By using our special sop for unmixed König graphs we give a stronger bound for $\operatorname{mi}(G)$ and at the same time provide a combinatorial criterion for the Cohen–Macaulay property for unmixed König graphs. A different combinatorial characterization of Cohen-Macaulay König graphs is known from [4, Proposition 28]. Our result (Theorem 2.7) is a follows: Let G be a König graph and $\{e_1, \ldots, e_m\}$ be a maximal matching of G with $\nu(G) = m$, and let k be the number of induced matchings of G contained in $\{e_1, \ldots, e_m\}$. Then $\operatorname{mi}(G) \leq k+1$ and equality holds if and only if G is a Cohen-Macaulay graph.

In the last section we introduce a sop for the Stanley–Reisner ring of any simplicial complex Δ . We call it the universal sop of $K[\Delta]$ because it is built in a uniform way for all simplicial complexes, and its construction does not depend on the base field K. The price we have to pay for this, is that this is not a sop of linear forms, instead it is defined as follows: $p_i(\Delta) = \sum_{\substack{F \in \Delta \\ |F|=i}} \prod_{j \in F} x_j$ for $i = 1, \ldots, \dim \Delta + 1$, see Theorem 3.1. By using Proposition 1.2 we obtain a Cohen–Macaulay criterion for $K[\Delta]$ in terms of this sop. This turns out to be a useful computational tool to check Cohen-Macaulayness, as we demonstrate at the example of a chessboard complex.

1. CRITERIA OF COHEN–MACAULAYNESS IN TERMS OF SYSTEMS OF PARAMETERS

In this section we collect some results on sop's which all are based on results of Serre (see [2, Theorem 4.6.10]) and which in terms of multiplicities allow to check whether a ring or a module is Cohen–Macaulay. One of the first efficient applications of these criteria was given by the first author of this paper in order to study the conormal module and the module of differentials of a K-algebra, see [7].

Proposition 1.1. Let R be a Noetherian local ring (or a standard graded K-algebra) with (graded) maximal ideal \mathfrak{m} , and let $I \subset R$ be an ideal generated by a (homogeneous) sop of R. Then

- (a) $e(R/I) = \ell(R/I) \ge e(R)$.
- (b) If e(R/I) = e(R), then R is Cohen-Macaulay.
- (c) If I is a reduction ideal of \mathfrak{m} and R is Cohen-Macaulay, then e(R/I) = e(R).

If (b) holds, then the sop which generates I is a regular sequence. In particular, r(R) = r(R/I), and so R is Gorenstein if and only if R/I is Gorenstein. (Here we denote by r(M) the (Cohen-Macaulay) type of a Cohen-Macaulay module M).

Proof. For the proof we recall a few facts: Let $M \neq 0$ be a finitely generated R-module of dimension d and $I \subseteq \mathfrak{m}$ be an ideal with dim M/IM = 0. Then

$$e(I,M) = \lim_{k \to \infty} (d!/k^d) \,\ell(M/I^{k+1}M)$$

is called the multiplicity of M with respect to I. The multiplicity of M, denoted e(M), is the multiplicity of M with respect to \mathfrak{m} .

Obviously, if $I \subseteq J \subseteq \mathfrak{m}$, then $e(I, M) \ge e(J, M)$. In particular,

(1)
$$e(I,M) \ge e(M).$$

On the other hand, if I is a reduction ideal of \mathfrak{m} with respect to M, that is, if $I\mathfrak{m}^k M = \mathfrak{m}^{k+1}M$ for some k, then equality holds in (1), see [2, Lemma 4.5.5].

We also need the following result ([2, Corollary 4.6.11] or [7] where it first appeared: let I be generated by a sop and assume that M has positive rank. Then

(i) $\ell(M/IM) \ge e(I, M) \operatorname{rank} M$.

(ii) M is Cohen-Macaulay if and only if $\ell(M/IM) = e(I, M)$ rank M.

Now we apply these results to the case that M = R. We first notice that $e(R/I) = \ell(R/I)$, since dim R/I = 0. Next (1) and (i) imply

(2)
$$\ell(R/I) \ge e(I,R) \ge e(R).$$

This proves (a). If $\ell(R/I) = e(R)$, then (2) implies $\ell(R/I) = e(I, R)$, and then (ii) yields (b). Finally, if I is a reduction ideal of \mathfrak{m} , then e(I, R) = e(R), and (2) together with (ii) implies (c).

If R is Cohen–Macaulay, then each sop is a regular sequence.

Now we turn to a graded version of Proposition 1.1.

Proposition 1.2. Let R be a standard graded K-algebra with graded maximal ideal \mathfrak{m} , and let I be generated by the homogeneous sop f_1, \ldots, f_d with deg $f_i = a_i$ for $i = 1, \ldots, d$. Then

(a) $e(R/I) = \ell(R/I) \ge a_1 a_2 \cdots a_d e(R)$.

(b) R is Cohen–Macaulay if and only if $e(R/I) = a_1 a_2 \cdots a_d e(R)$.

If the equivalent conditions given in (b) hold, then f_1, \ldots, f_d is a regular sequence. In particular r(R) = r(R/I), and R is Gorenstein if and only if R/I is Gorenstein.

Proof. Let $a = a_1 a_2 \cdots a_m$ and set $b_i = a/a_i$ for $i = 1, \ldots, d$. Then

$$e(f_1^{b_1}, \dots, f_d^{b_d}, R) = b_1 b_2 \cdots b_d e(f_1, \dots, f_d, R)$$

see [16, Proposition 11.2.9]. Moreover, deg $f_i^{b_i} = a$ for $i = 1, \ldots, d$. Therefore,

$$(f_1^{b_1},\ldots,f_d^{b_d})\subseteq \mathfrak{m}^a.$$

Thus, since $e(\mathfrak{m}^a, R) = a^d e(R)$ ([16, Proposition 11.2.9]), we obtain that

(3)
$$b_1 b_2 \cdots b_d e(f_1, \dots, f_d, R) = e(f_1^{b_1}, \dots, f_d^{b_d}, R) \ge e(\mathfrak{m}^a, R) = a^d e(R),$$

which together with (i) in the proof of Proposition 1.1 imply the inequality in (a).

(b) Assuming that $\ell(R/I) = a_1 a_2 \cdots a_d e(R)$, we obtain together with (3) that

$$a_1a_2\cdots a_de(R) = \ell(R/I) \ge e(f_1,\ldots,f_d,R) \ge a_1a_2\cdots a_de(R),$$

and hence $\ell(R/I) = e(f_1, \ldots, f_d, R)$. Thus (ii) in the proof of Proposition 1.1 implies that R is Cohen–Macaulay. Conversely, suppose that R is Cohen–Macaulay. Then f_1, \ldots, f_d is a regular sequence. Let $\operatorname{Hilb}_R(t) = Q_R(t)/(1-t)^d$ be the Hilbert series of R. Then

$$Q_{R/I}(t) = \operatorname{Hilb}_{R/I}(t) = \operatorname{Hilb}_{R}(t) \prod_{i=1}^{d} (1 - t^{a_i}) = Q_R(t) (\prod_{i=1}^{d} (\sum_{j=0}^{a_{i-1}} t^j).$$

It follows that

$$\ell(R/I) = Q_{R/I}(1) = Q_R(1)a_1a_2\cdots a_d = e(R)a_1a_2\cdots a_d$$

The next result is a certain refinement of the statements given in Proposition 1.2. We first recall the following fact (see for example ([2, Proposition A.4.]): Let (R, \mathfrak{m}) be a Noetherian local ring and f_1, \ldots, f_m a sequence of elements in \mathfrak{m} . Then

- (α) dim $R \ge \dim R/(f_1, \ldots, f_m) \ge \dim R m$, and
- (β) dim $R/(f_1, \ldots, f_m)$ = dim R m if and only f_1, \ldots, f_m can be completed to a sop of R.

A similar statement holds for graded K-algebras.

Proposition 1.3. With the assumptions and notation of Proposition 1.2 let

$$U_i = \operatorname{Ker}(R/(f_1, \dots, f_{i-1}) \xrightarrow{J_i} R/(f_1, \dots, f_{i-1}))$$

Then

(a) dim
$$U_i \leq d - i$$
 for all i .

(b) Set dim $U_i = -1$ if $U_i = 0$. Then

$$e(R/I) = a_1 \cdots a_d e(R) + \sum_{\substack{i=1\\\dim U_i = d-i}}^d a_{i+1} \cdots a_d e(U_i).$$

In particular, if deg $f_i = 1$ for all *i*, then

$$e(R/I) = e(R) + \sum_{\substack{i=1 \ \dim U_i = d-i}}^{a} e(U_i).$$

Proof. (a) U_i is a submodule of $R/(f_1, \ldots, f_{i-1})$ with $f_iU_i = 0$. Thus U_i is a $R/(f_1, \ldots, f_i)$ -module and hence dim $U_i \leq \dim R/(f_1, \ldots, f_i) = d - i$, where the equation follows from (β) .

(b) From the following exact sequence

$$0 \to U_i \to (R/(f_1, \dots, f_{i-1}))(-a_i) \xrightarrow{f_i} R/(f_1, \dots, f_{i-1}) \to R/(f_1, \dots, f_i) \to 0$$

we deduce the equality

(4)
$$\operatorname{Hilb}_{R/(f_1,\dots,f_i)}(t) = (1 - t^{a_i}) \operatorname{Hilb}_{R/(f_1,\dots,f_{i-1})}(t) + \operatorname{Hilb}_{U_i}(t).$$

We have $\operatorname{Hilb}_{R/(f_1,\ldots,f_i)}(t) = Q_i(t)/(1-t^{d-i})$ with $Q_i(1) = e(R/(f_1,\ldots,f_i))$, similarly, $\operatorname{Hilb}_{R/(f_1,\ldots,f_{i-1})}(t) = Q_{i-1}(t)/(1-t^{d-(i-1)})$ with $Q_{i-1}(1) = e(R/(f_1,\ldots,f_{i-1}))$ and $\operatorname{Hilb}_{U_i}(t) = P_i(t)/(1-t^{\delta_i})$ with $P_i(1) = e(U_i)$ and $\delta_i \leq d-i$.

Thus (4) implies that

$$Q_i(t)/(1-t^{d-i}) = (1-t^{a_i})(Q_{i-1}(t)/(1-t^{d-(i-1)})) + P_i(t)/(1-t^{\delta_i}),$$

from which we deduce that

$$Q_i(t) = Q_{i-1}(t) \left(\sum_{j=0}^{a_i-1} t^j\right) + (1-t)^{d-i-\delta_i} P_i(t).$$

Substituting t by 1, we get

$$e(R/(f_1,\ldots,f_i)) = \begin{cases} a_i e(R/(f_1,\ldots,f_{i-1})), & \text{if } \dim U_i < d-i, \\ a_i e(R/(f_1,\ldots,f_{i-1})) + e(U_i), & \text{if } \dim U_i = d-i. \end{cases}$$

These formulas together with induction on i complete the proof.

Proposition 1.3 together with Proposition 1.2 has the following a surprising consequence

Corollary 1.4. With the assumptions and notation of Proposition 1.3 let r < dbe an integer with the property that dim $U_i < d - i$ for i = 1, ..., r and that $R/(f_1, ..., f_r)$ is Cohen-Macaulay. Then R is Cohen-Macaulay and $U_i = 0$ for all i. In particular, if $f_1, ..., f_d$ is a superficial sequence and $R/(f_1, ..., f_i)$ is Cohen-Macaulay for some i < d, then R is Cohen-Macaulay.

Proof. Proposition 1.3 implies that $e(R/(f_1, \ldots, f_r)) = a_1 \cdots a_r e(R)$. Since by assumption $R/(f_1, \ldots, f_r)$ is Cohen–Macaulay, it follows that f_{r+1}, \cdots, f_d is a regular $R/(f_1, \ldots, f_r)$ -sequence. Hence, $e(R/(f_1, \ldots, f_d)) = a_{r+1} \cdots a_d e(R/(f_1, \ldots, f_r))$, and we deduce that $e(R/(f_1, \ldots, f_d)) = a_1 \cdots a_d e(R)$. Thus the desired result follows from Proposition 1.2.

We close this section with a remark and a question. Let $S = K[x_1, \ldots, x_n]$ be the polynomial ring over the field K, R = S/I with $I \subset (x_1, \ldots, x_n)^2$ a graded ideal. Let f_1, \ldots, f_d be linear forms of S which form a sop for R. Let $\overline{S} = S/(f_1, \ldots, f_d)$. Then \overline{S} is isomorphic to a polynomial ring in n - d variables, and $R/(f_1, \ldots, f_d) = \overline{S}/\overline{I}$, where $\overline{I} = I\overline{S}$.

Remark 1.5. With the notation introduced we have $\operatorname{proj} \dim_{\overline{S}} \overline{I} \leq \operatorname{proj} \dim_{S} I$ and $\mu(\overline{I}) \leq \mu(I)$. Equality holds in both inequalities, if S/I is Cohen–Macaulay.

Proof. By the Auslander-Buchsbaum formula,

$$\operatorname{proj\,dim}_S S/I = n - \operatorname{depth} S/I \ge n - \operatorname{dim} S/I = n - d = \operatorname{proj\,dim}_{\overline{S}} S/I.$$

The last equation holds since $\dim \overline{S}/\overline{I} = 0$. This implies the first assertion. It is obvious that $\mu(I) \ge \mu(I\overline{S})$. Finally, if S/I is Cohen–Macaulay, then I is generated by a regular sequence and the desired equalities hold. \Box

Remark 1.5 implies in particular that if $\mu(I) < \mu(I)$, then R cannot be Cohen-Macaulay. On the other hand, $\mu(\overline{I}) = \mu(I)$, does not necessarily imply that R is Cohen-Macaulay. For example for the cycle graph $C_6 : x_1, \ldots, x_6$, the sequence $x_1 - x_2, x_3 - x_4, x_5 - x_6$ is a sop for $R = K[x_1, \ldots, x_6]/I(C_6)$ and $\overline{I(C_6)} = (x_1^2, x_3^2, x_5^2, x_1x_3, x_3x_5, x_1x_5)$. Then $\mu(\overline{I(C_6)}) = \mu(I(C_6)) = 6$, while R is not Cohen-Macaulay.

In view of these inequalities one is tempted to ask whether under the assumptions of Remark 1.5 we have $\operatorname{reg}(\overline{I}) \leq \operatorname{reg}(I)$. In the next section we show that this inequality for the regularity indeed holds for the edge ideal of König graphs and suitable natural sop's.

2. Special systems of parameters applied to order complexes and König graphs

In this section we define monomial ideals of König type which include edge ideals of König graphs and give a characterization for these ideals in terms some sop's for their quotient rings. Also we apply Proposition 1.1 to the Stanley-Reisner ring of two families of simplicial complexes, namely the order complex of a finite poset and the independence complex of a König graph and give combinatorial descriptions for the Cohen-Macaulay property of these rings.

For a poset P, a nonempty subposet C of P which is totally ordered is called a *chain* in P. The *order complex* of P denoted by $\Delta(P)$ is the simplicial complex whose faces are the chains in P. The *length* of a chain C in P is defined to be |C| - 1. The *height* of an element x in P is defined to be the maximal length of a chain descending from x. For elements x and y in a poset P, it is said that y covers x, denoted x < y, if x < y and there exists no $z \in P$ such that x < z < y. Also for a monomial ideal I, the cardinality of any minimal generating set of monomials of I is denoted by $\mu(I)$.

Theorem 2.1. Let P be a poset which as a set is the disjoint union of two sets C_1 and C_2 , where $C_1 : x_1 < x_2 < \cdots < x_n$ and $C_2 : y_1 < y_2 < \cdots < y_n$ are maximal chains in P and let $\Delta = \Delta(P)$. Then the following conditions are equivalent:

- (a) Δ is Cohen-Macaulay.
- (b) Δ is pure shellable.
- (c) Δ satisfies the following conditions:
 - (1) If $x_i \lt y_j$ or $y_i \lt x_j$, then j = i + 1, and
 - (2) $\{x_i, y_{i+1}\} \notin \Delta$ implies that $\{x_{i+1}, y_i\} \in \Delta$.

Proof. (b) \Rightarrow (a): By [8, Theorem 8.2.6] the assertion holds.

(a) \Rightarrow (c): Suppose that Δ is Cohen-Macaulay. Then Δ is pure. Suppose $x_i \leq y_j$ for some *i* and *j*. If $j \leq i$, then the chain $x_1 < \cdots < x_i < y_j < y_{j+1} < \cdots < y_n$ is a chain of cardinality at least n + 1, which is included in some maximal chain of *P*. But by purity of Δ any maximal chain should have cardinality $|C_1| = n$, which

gives a contradiction. Thus i < j. Similarly if $y_i < x_j$ for some i and j, then i < j. Now assume that $x_i < y_j$ and by contradiction let $j \neq i + 1$. Since j > i, one should have j > i + 1. Then $x_1 < \cdots < x_i < y_j < y_{j+1} < \cdots < y_n$ is a maximal chain of cardinality at most n - 1 in P, which is again a contradiction to purity of Δ . So j = i + 1. The argument for the case $y_i < x_j$ is similar.

To prove (2), first we show that the sequence $x_1 - y_1, x_2 - y_2, \ldots, x_n - y_n$ is a sop for the ring $R = S/I_{\Delta}$, where $S = K[x_1, \ldots, x_n, y_1, \ldots, y_n]$. Indeed by (1), $x_i y_i \in I_{\Delta}$ for any $1 \leq i \leq n$. So we have $x_i^2, y_i^2 \in (I_{\Delta}, x_1 - y_1, x_2 - y_2, \ldots, x_n - y_n)$ for all *i* and then dim $R/(x_1 - y_1, x_2 - y_2, \ldots, x_n - y_n)R = 0$. Also dim R =dim $\Delta + 1 = n$. Now, by contradiction suppose that for some *i*, $\{x_i, y_{i+1}\} \notin$ Δ and $\{x_{i+1}, y_i\} \notin \Delta$. This means that $x_i y_{i+1}$ and $x_{i+1} y_i$ belong to the set of minimal generators $\mathcal{G}(I_{\Delta})$ of I_{Δ} . One has $R/(x_1 - y_1, x_2 - y_2, \ldots, x_n - y_n)R \cong$ $K[x_1, \ldots, x_n]/I'$, where $I' = (x_i x_j : x_i \text{ and } y_j \text{ are non-comparable in } P)$. Since R is Cohen-Macaulay, by Remark 1.5, one should have $\mu(I_{\Delta}) = \mu(I')$. But since $\mathcal{G}(I_{\Delta}) = \{x_i y_j : x_i \text{ and } y_j \text{ are non-comparable in } P\}$, and $x_i y_{i+1}, x_{i+1} y_i \in \mathcal{G}(I_{\Delta})$ correspond to just one element in $\mathcal{G}(I')$ that is $x_i x_{i+1}$, we have $\mu(I') < \mu(I_{\Delta})$, a contradiction. Thus $\{x_i, y_{i+1}\} \in \Delta$ or $\{x_{i+1}, y_i\} \in \Delta$.

(c) \Rightarrow (b): Let P be a poset satisfying the assumptions of (c). Let $F = \{z_1 < z_2 < \cdots < z_k\}$ be an arbitrary facet of Δ . First note that $z_1 < z_2 < \cdots < z_k$. Also assumption (1) of (c) implies that $\{x_i, y_i\} \notin \Delta$ for any $1 \leq i \leq n$, and then $|\{x_i, y_i\} \cap F| \leq 1$. We claim that for each facet F of Δ , $|\{x_i, y_i\} \cap F| = 1$ for any $1 \leq i \leq n$. One can easily see that for i = 1 the claim holds true. Indeed if $z_1 \in C_1$, then $z_1 = x_1$, because otherwise $F \subsetneq F \cup \{x_1\} \in \Delta$, a contradiction. Similarly if $z_1 \in C_2$, then $z_1 = y_1$. So $|\{x_1, y_1\} \cap F| = 1$. Assume inductively that for any $i = 1, \ldots, m-1$, $|\{x_i, y_i\} \cap F| = 1$. We show that $|\{x_m, y_m\} \cap F| = 1$. We have $x_{m-1} \in F$ or $y_{m-1} \in F$. Without loss of generality suppose $x_{m-1} \in F$. If $x_m \in F$, we are done. So assume that $x_m \notin F$. Note that $x_{m-1} \neq z_k$, since otherwise $F \subsetneq F \cup \{x_m\} \in \Delta$, a contradiction. So there exists $z_t \in F$ such that $x_{m-1} < z_t$. If $z_t = x_j$ for some j, then j > m and hence $x_{m-1} < x_m < z_t$, which contradicts to $x_{m-1} < z_t$. Thus $z_t = y_j$ for some j and by (1), j = m. Thus $z_t = y_m \in F$. So $|\{x_m, y_m\} \cap F| = 1$. Therefore any facet F of Δ has cardinality n such that for any $1 \leq i \leq n$, either $x_i \in F$ or $y_i \in F$. Thus Δ is pure.

Let $\mathcal{F}(\Delta)$ denotes the set of facets of Δ . To prove the shellability, consider the ordering on $\mathcal{F}(\Delta)$ as follows. For the facets F_i and F_j of Δ , we set $F_i \prec F_j$ if there exists $1 \leq t \leq n$ such that $y_t \in F_j \setminus F_i$ and for any d < t, the elements of height d in F_i and F_j are the same. Now, let $F_i \prec F_j$ and $1 \leq t \leq n$ be such that $y_t \in F_j \setminus F_i$ and for any d < t, the elements of height d in F_i and F_j are the same. If $\{x_t, y_{t+1}\} \in \Delta$, then we set $F_k = (F_j \setminus \{y_t\}) \cup \{x_t\}$. Then $F_k \in \mathcal{F}(\Delta)$, $F_k \prec F_j$ and $F_j \setminus F_k = \{y_t\}$. So we may assume that $\{x_t, y_{t+1}\} \notin \Delta$. Thus by assumption, $\{x_{t+1}, y_t\} \in \Delta$. Let r be the greatest integer with $y_t, y_{t+1}, \ldots, y_r \in F_j \setminus F_i$ and $\{x_{t+1}, y_t\} \in \Delta$. Two cases may happen:

(1) Suppose that $y_{r+1} \in F_j \setminus F_i$. Then by the maximality of r, $\{x_{r+2}, y_{r+1}\} \notin \Delta$. So by assumption $\{x_{r+1}, y_{r+2}\} \in \Delta$. Then $F_k = (F_j \setminus \{y_{r+1}\}) \cup \{x_{r+1}\} \in \mathcal{F}(\Delta)$, $F_k \prec F_j$ and $F_j \setminus F_k = \{y_{r+1}\}$.

(2) Suppose that $y_{r+1} \notin F_j \setminus F_i$. Then either $y_{r+1} \notin F_j$ or $y_{r+1} \in F_i \cap F_j$. In both cases, we set $F_k = (F_j \setminus \{y_r\}) \cup \{x_r\} \in \mathcal{F}(\Delta)$ and F_k is the facet which fulfils the desired condition for shellability.

The next result shows that the linear resolution property for $I_{\Delta(P)}$ can be again expressed in terms of conditions on chains of the form $x_i < y_j$ in P. For a graph Gby V(G) and E(G) we mean the vertex set and the edge set of G, respectively. Also for a subset $S \subseteq V(G)$, the induced subgraph of G on the set S is denoted by G_S .

Proposition 2.2. Let P be a poset with the assumptions of Theorem 2.1 and $\Delta = \Delta(P)$. Then I_{Δ} has a linear resolution if and only if whenever $\{x_i, y_j\}, \{x_r, y_s\} \in \Delta$, then $\{x_i, y_s\} \in \Delta$ or $\{x_r, y_j\} \in \Delta$.

Proof. We have $I_{\Delta} = I(G)$, where G is a bipartite graph with bipartition $X = \{x_1, x_2, \ldots, x_n\}, Y = \{y_1, y_2, \ldots, y_n\}$ and the edge set $E(G) = \{\{x_i, y_j\} : \{x_i, y_j\} \notin \Delta\}$. Thus by [6, Theorem 1], I_{Δ} has a linear resolution if and only if G^c is a chordal graph. Any cycle C of length $m \geq 5$, has a chord in G^c , because C has at least 3 vertices from X or from Y, and G_X^c and G_Y^c are complete graphs. So G^c is chordal if and only if any cycle of length 4 in G^c has a chord. Note that $C : x_i, x_r, y_s, y_j$ is a cycle in G^c if and only if $\{x_i, y_j\}, \{x_r, y_s\} \in \Delta$ and it has a chord if and only if $\{x_i, y_j\}, \{x_r, y_s\} \in \Delta$. Thus G^c is a chordal graph if and only if whenever $\{x_i, y_j\}, \{x_r, y_s\} \in \Delta$, then $\{x_i, y_s\} \in \Delta$ or $\{x_r, y_j\} \in \Delta$. \Box

For a graph G, let $\tau(G)$ be the minimum cardinality of a vertex cover of G and $\nu(G)$ denotes the maximum cardinality of a matching of G. One can see that for any graph G, $\tau(G) \geq \nu(G)$. Recall that a graph G is called a König graph, when this inequality becomes an equality. Let $I \subset S$ be a monomial ideal in the polynomial ring $S = K[x_1, \ldots, x_n]$ over the field K in n variables. We denote by m-grade(I) the maximal length of a regular sequence of monomials in I, and call this number the monomial grade of I. One has m-grade $(I) \leq \operatorname{grade}(I) = \operatorname{height}(I)$. We call I a monomial ideal of König type if $I \neq 0$ and m-grade $(I) = \operatorname{height}(I)$. The naming is justified by the fact that if I = I(G) for some graph G, then $\operatorname{height}(I) = \tau(G)$ and m-grade $(I) = \nu(G)$, so that the edge ideal of König graphs are the monomial ideals of König type among edge ideals.

The following theorem characterizes monomial ideals of König type in terms of existence of some forms of sop's for their quotient rings.

Theorem 2.3. Let $I \subset S = K[x_1, ..., x_n]$ be a monomial ideal. Then the following conditions are equivalent:

- (a) I is a monomial ideal of König type.
- (b) S/I admits a sop f_1, \ldots, f_d , where each f_k is of the form $x_i x_j$ for suitable *i* and *j*.

Proof. (a) \Rightarrow (b): Let h = height(I). Since I is a monomial ideal of König type, there exists a regular sequence of monomials u_1, \ldots, u_h with $u_i \in I$ for $i = 1, \ldots, h$.

We may assume that each u_i belongs to the unique minimal set of monomial generators $\mathcal{G}(I)$ of I. Indeed, if $u_i = wv$ with $v \in \mathcal{G}(I)$, then we may replace u_i by v in the above regular sequence.

For a monomial $u \in S$ we set $\operatorname{supp}(u) = \{i: x_i | u\}$. We may assume that $\bigcup_{u \in \mathcal{G}(I)} \operatorname{supp}(u) = [n]$. Indeed, suppose this is not the case. Then for simplicity we may assume that $\bigcup_{u \in \mathcal{G}(I)} \operatorname{supp}(u) = \{1, \ldots, r\}$. Note that $r \geq 1$, since $I \neq 0$. Let $R = K[x_1, \ldots, x_r]/(u: u \in \mathcal{G}(I))$. Then $S/I = R[x_{r+1}, \ldots, x_n]$, and $x_1 - x_{r+1}, \ldots, x_1 - x_n$ is part of a sop of S/I and $(S/I)/(x_1 - x_{r+1}, \ldots, x_1 - x_n) \cong R$. Thus if R has the desired sop, then so does S/I.

We proceed by induction on $d = \dim S/I$. If d = 0, then there is nothing to show. Suppose now that d > 0. Then at least one u_i is not a pure power. Assume this is not the case. After a relabeling of the variables we may then assume that $u_i = x_i^{a_i}$ with $a_i > 1$ for $i = 1, \ldots, h$. Since d > 0 and $\bigcup_{u \in \mathcal{G}(I)} \operatorname{supp}(u) = [n]$, there must exist $u \in \mathcal{G}(I)$, with $\operatorname{supp}(u) \notin \{1, \ldots, h\}$. Then $\operatorname{height}(I) \ge \operatorname{height}(u_1, \ldots, u_h, u) \ge h+1$, a contradiction.

We may assume that u_1 is not a pure power, say $u_1 = x_1^{a_1} x_2^{a_2} \cdots x_n^{a_n}$ with $a_1, a_2 > 0$. Let $f_1 = x_1 - x_2$. We claim that f_1 is a part of a sop. In other words, f_1 is not contained in any minimal prime ideal P of I with height(P) = height(I) = h. Indeed, let P be a minimal prime ideal containing f_1 . Since I is a monomial ideal, P is a monomial prime ideal. Therefore, if $f_1 \in P$, then $x_1, x_2 \in P$. For each i there exists j_i such that x_{j_i} divides u_i and $x_{j_i} \in P$. Thus, $Q = (x_1, x_2, x_{j_2}, \cdots, x_{j_d}) \subset P$. Since u_1, \ldots, u_h is a regular sequence, the supports of the u_i are pairwise disjoint. This implies that the variables generating Q are pairwise distinct. It follows that height $(P) \ge h + 1$, and f_1 is a part of a sop.

Identifying x_1 with x_2 , we see that $(S/I)/(f_1) \cong K[x_2, \ldots, x_n]/\overline{I}$, where $\overline{I} = (x_2^{a_1+a_2}\cdots x_n^{a_n}, u_2, \ldots, u_h, \cdots)$. This shows that m-grade $(\overline{I}) \ge h$. Since f_1 is a parameter element of S/I it follows that height $(\overline{I}) = \text{height}(I) = h$ and since in general m-grade $(\overline{I}) \le \text{height}(\overline{I})$, we must have m-grade $(\overline{I}) = \text{height}(\overline{I})$. This means that \overline{I} is a again monomial ideal of König type. Since $\dim((S/I)/(f_1)) = d - 1$, we may apply our induction hypothesis, and find a sop f_2, \ldots, f_d of $(S/I)/(f_1)$ as required in (b), Then f_1, f_2, \ldots, f_d is the desired sop for S/I,

(b) \Rightarrow (a): Let R = S/I and $\overline{R} = R/(f_1, \ldots, f_d)R$. Then $\overline{R} \cong \overline{S}/\overline{I}$, where \overline{I} is a monomial ideal, because reduction modulo f_1, \ldots, f_d simply identifies variables. For simplicity we may assume that $\overline{S} = K[x_1, \ldots, x_h]$. Since dim $(\overline{R}) = 0$ and since \overline{I} is a monomial ideal, it follows that $\mathcal{G}(\overline{I})$ contains a pure power $x_i^{a_i}$ of each the variables x_1, \ldots, x_h . Let u_1, \ldots, u_h be generators of I with the property that u_i specializes to $x_i^{a_i}$ under the reduction modulo f_1, \ldots, f_d . Suppose u_i and u_j have a common factor for $i \neq j$. Then this is also the case for $x_i^{a_i}$ and $x_j^{a_j}$, a contradiction. Therefore, u_1, \ldots, u_h is a regular sequence, and so m-grade(I) = height(I).

Applying Theorem 2.3 to the edge ideal of König graphs, we have the following algebraic characterization for a König graph G in terms of special sop's for R = S/I(G). Note that by the proof of Theorem 2.3, each elements $x_i - x_j$ of the sop for R = S/I(G) corresponds to an edge $e = \{x_i, x_j\}$ of G.

Corollary 2.4. Let G be a graph without isolated vertices, S = K[V(G)] and for any edge $e = \{x, y\} \in E(G)$, let $f_e = x - y$ be an element in S. Then G is a König graph if and only if there exists a subset $\{e_1, \ldots, e_d\}$ of edges of G such that f_{e_1}, \ldots, f_{e_d} is a sop for R = S/I(G).

Let R be a graded ring and M a finitely generated graded R-module. It is known that for an M-superficial sequence f_1, \ldots, f_m of linear forms $\operatorname{reg}(R/(f_1, \ldots, f_m)R) \leq \operatorname{reg}(R)$, see [5, Proposition 20.20]. In view of this fact it is natural to ask the following

Question 2.5. Let R be a graded ring, M be a finitely generated graded R-module and f_1, \ldots, f_m be a sop of linear forms for M. Does the inequality

$$\operatorname{reg}(M/(f_1,\ldots,f_m)M) \le \operatorname{reg}(M)$$

hold?

In the following theorem we prove the expected inequality for the S-module R = S/I(G), when G is a König graph and the sop is of a natural special form. First we recall some definitions. Two edges e and e' of a graph G form a gap, when no endpoint of e and e' are adjacent in G. Otherwise we say that e and e' are adjacent. Moreover, a subset A of edges of G is called an *induced matching* if any two edges in A form a gap. By a(G) we mean the maximum cardinality of an induced matching of G. The maximum cardinality of an independent set of G is denoted by $\alpha(G)$. For a graph G, with the vertex set $\{x_1, \ldots, x_n\}$ the whiskered graph of G is a graph which is obtained by adding new vertices $\{y_1, \ldots, y_n\}$ and edges $\{\{x_i, y_i\} : 1 \le i \le n\}$ to G. This new graph is denoted by $G \cup W(G)$ and the edges $\{x_i, y_i\}$ are called whiskers.

Theorem 2.6. Let G be a graph, S = K[V(G)] and $\{e_1, \ldots, e_m\} \subseteq E(G)$ such that f_{e_1}, \ldots, f_{e_m} is a sop for R = S/I(G). Then

$$\operatorname{reg}(R/(f_{e_1},\ldots,f_{e_m})R) \le \operatorname{reg}(R).$$

Proof. Let $e_i = \{x_i, y_i\}$ for any $1 \le i \le m$. Then by the proof of Theorem 2.3, $\bigcup_{i=1}^m e_i = V(G)$ and without loss of generality we may assume that for some $r \le m$, $C = \{x_1, x_2, \ldots, x_r\}$ is a minimal vertex cover of G and $\{e_1, \ldots, e_r\}$ is a maximal matching of G. Set $\overline{R} = R/(f_{e_1}, \ldots, f_{e_m})R$. Then $\overline{R} \cong K[x_1, x_2, \ldots, x_r]/L$ where

$$L = (x_1^2, \dots, x_r^2, x_i x_j)$$
: $i < j \le r$, e_i and e_j are adjacent in G).

Thus $\operatorname{reg}(\overline{R}) = \operatorname{reg}(K[x_1, \ldots, x_r]/L)$. Since polarization does not change the regularity, one has $\operatorname{reg}(K[x_1, \ldots, x_r]/L) = \operatorname{reg}(T/I(H'))$, where $T = K[x_1, \ldots, x_r, x'_1, \ldots, x'_r]$ for new variables x'_1, \ldots, x'_r and $H' = H \cup W(H)$, where H is a graph on $\{x_1, \ldots, x_r\}$ with

$$E(H) = \{\{x_i, x_j\}: i < j \le r, e_i \text{ and } e_j \text{ are adjacent in } G\}$$

and $\{x_i, x'_i\}$ is a whisker of H' for any $1 \le i \le r$. Since any whiskered graph is very well-covered, by [12, Theorem 1.3], $\operatorname{reg}(T/I(H')) = \operatorname{a}(H')$. One can easily see that $\operatorname{a}(H')$ is precisely the maximum size of independent sets of vertices in H. Moreover, by the definition of H, a set $\{x_{s_1}, \ldots, x_{s_t}\}$ of vertices of H is an independent set of H if and only if $\{e_{s_1}, \ldots, e_{s_t}\} \subseteq \{e_1, \ldots, e_m\}$ is an induced matching of G. Therefore

 $a(H') \leq a(G)$. Thus $reg(\overline{R}) = reg(K[x_1, \dots, x_r]/L) = reg(T/I(H')) = a(H') \leq a(G) \leq reg(R)$. The last inequality holds by [11, Lemma 2.2]. \Box

In general the inequality of Theorem 2.6 may be strict. Let $G = C_n$ denotes the cycle graph with n vertices. Suppose $E(G) = \{e_i = \{x_i, x_{i+1}\} : 1 \leq i \leq n-1\} \cup \{e_n = \{x_1, x_n\}\}$ and $n \equiv 2 \mod 4$. Let n = 4m + 2 for some $m \geq 1$. Then $\{f_{e_{2i-1}} : 1 \leq i \leq 2m+1\}$ is a sop for R = S/I(G). With the notation used in the proof of Theorem 2.6, $H' = C_{2m+1} \cup W(C_{2m+1})$ and $\operatorname{reg}(\overline{R}) = a(H') = \alpha(H) = m$. Since $\{e_{3i+1} : 0 \leq i \leq m\}$ is an induced matching of G of cardinality m + 1, $\operatorname{reg}(R) \geq a(G) \geq m+1 > m = \operatorname{reg}(\overline{R})$.

For a graph G, let $\operatorname{mi}(G)$ denote the number of maximal independent sets of G. After Erdös and Moser considered the problem of determining the largest value of $\operatorname{mi}(G)$ in terms of the number of vertices of G, investigating this number and upper bounds for it has been studied for various classes of graphs. In [9, Corollary 3.4] it was shown that for a König graph G, $2^{\nu(G)}$ is an upper bound for $\operatorname{mi}(G)$. Also in [1, Theorem 1] it was proved that $\operatorname{mi}(G) \leq M(G) + 1$, where M(G) is the number of induced matchings in G. In the following for an unmixed König graph G, we use a sop of the form f_{e_1}, \ldots, f_{e_d} for S/I(G) and improve the upper bound for $\operatorname{mi}(G)$. Moreover, we give a combinatorial description of the Cohen-Macaulay property for unmixed König graphs. A different combinatorial characterization of Cohen-Macaulay König graphs is presented in [4, Proposition 28]. Recall that a graph G is called unmixed if all the minimal vertex covers (maximal independent sets) of G are of the same cardinality.

Theorem 2.7. Let G be an unmixed König graph, $\{e_1, \ldots, e_m\}$ be a maximal matching of G with $\tau(G) = \nu(G) = m$ and k be the number of induced matchings of G contained in $\{e_1, \ldots, e_m\}$. Then

- (a) $\operatorname{mi}(G) \leq k+1$, and
- (b) G is a Cohen-Macaulay graph if and only if mi(G) = k + 1.

Proof. Let S = K[V(G)], R = S/I(G) and $\dim(R) = d$. By the proof of Theorem 2.4, there exist $e_{m+1}, \ldots, e_d \in E(G)$ such that f_{e_1}, \ldots, f_{e_d} is a sop for R. We may assume that $\{x_1, x_2, \ldots, x_m\}$ is a minimal vertex cover of G, where $x_i \in e_i$ for all $1 \leq i \leq m$. Then $R/(f_{e_1}, \ldots, f_{e_d})R \cong K[x_1, x_2, \ldots, x_m]/L$, where

$$L = (x_1^2, x_2^2, \dots, x_m^2, x_i x_j : i < j \le m, e_i \text{ and } e_j \text{ are adjacent in } G).$$

So $\ell(R/(f_{e_1},\ldots,f_{e_d})R) = \ell(K[x_1,x_2,\ldots,x_m]/L)$. Since $x_1^2, x_2^2,\ldots,x_m^2 \in L$, any basis element of $K[x_1,x_2,\ldots,x_m]/L$ other than 1, is an squarefree monomial of the form $x_{i_1}\cdots x_{i_r}$, where $\{e_{i_1},\ldots,e_{i_r}\} \subseteq \{e_1,\ldots,e_m\}$ is an induced matching of G. So $\ell(K[x_1,x_2,\ldots,x_m]/L) = k + 1$. Note that R is the Stanley-Reisner ring of the independence complex Δ_G of G. Since G is unmixed, e(R) is the number of facets of Δ_G which is equal to mi(G). Now, by using Proposition 1.2,

$$\min(G) = e(R) \le \ell(R/(f_{e_1}, \dots, f_{e_d})R) = \ell(K[x_1, x_2, \dots, x_m]/L) = k + 1$$

and equality holds if and only if G is Cohen-Macaulay.

Theorem 2.7 shows that in a Cohen-Macaulay König graph G, no matter which maximal matching M of G we choose, if $|M| = \nu(G)$, then the number of induced matchings of G contained in M is equal to mi(G).

3. A UNIVERSAL SYSTEM OF PARAMETERS FOR STANLEY–REISNER RINGS

Let K be a field and Δ an arbitrary simplicial complex on [n] of dimension d-1, and let $S = K[x_1, \ldots, x_n]$ be the polynomial ring in n variables. We show that there exists a universal standard sop for $K[\Delta] = S/I_{\Delta}$. Using this sop we present a criterion for the Cohen-Macaulayness of $K[\Delta]$.

For $i = 1, \ldots, d$, we set

$$p_i(\Delta) = \sum_{F \in \Delta \ |F|=i} \mathbf{x}_F,$$

where $\mathbf{x}_F = \prod_{i \in F} x_i$.

Theorem 3.1. The residue classes of the elements $p_1(\Delta), \ldots, p_d(\Delta)$ in $K[\Delta]$ form a sop of $K[\Delta]$.

Proof. We first consider the case that Δ is the *n*-simplex Γ_n . In that case $K[\Delta] = S$, and we have to show that dim $S/(p_1(\Delta), \ldots, p_n(\Delta)) = 0$. In order to simplify notation we write p_i for $p_i(\Delta)$ and all *i*.

Note that

(5)
$$p_i = \sum_{|F|=i} \mathbf{x}_F.$$

Let < denote the reverse lexicographical order, and let $J = (p_1, \ldots, p_n)$. We claim that $x_i^i \in in_{\leq}(J)$ for $i = 1, \ldots, n$. Then this shows that indeed dim S/J = 0.

For $i = 1, \ldots, n$, let

$$g_i = \sum_{j=1}^{i} (-1)^{j+1} x_i^{i-j} p_j.$$

We will show that $x_i^i = in_{\leq}(g_i)$ for $i = 1, \ldots, n$.

Observe first that g_i is homogeneous of degree i and that $x_i^i \in \text{supp}(g_i)$. In order to complete the proof of the claim, we have to show that if u is a monomial of degree i with $u > x_i^i$, then $u \notin \text{supp}(g_i)$.

Let v be a monomial in the support of g_i , $v = x_1^{a_1} \cdots x_n^{a_n}$. Then $a_j \leq 1$ for all $j \neq i$. Hence if x_j^a divides u for $j \neq i$ and a > 1, then $u \notin \operatorname{supp}(g_i)$. Therefore, we may assume that $u = x_G x_i^k$ with $0 \leq k \leq i$, $i \notin G$ and |G| = i - k. Moreover, since $u > x_i^i$ in the reverse lexicographic order, it follows that $G \subset [i-1]$. Hence k > 0, because u is of degree i. Then $u \in \operatorname{supp}(x_i^k p_{i-k})$ with coefficients 1, because $x_G \in \operatorname{supp}(p_{i-k})$, and we also have that $u \in \operatorname{supp}(x_i^{k-1} p_{i-k+1})$ with coefficient 1, because $x_G x_i = x_{G \cup \{i\}}$ belongs to $\operatorname{supp}(p_{i-k+1})$.

Since u does not belong to the support of any other summand $x_i^{i-j}p_j$ of g_i , and since the summands in which u appears in the support, have different signs, we conclude that $u \notin \operatorname{supp}(g_i)$, as desired.

In order to deal with the general case we observe that

$$S/(I_{\Delta}, p_1(\Delta), \dots, p_d(\Delta)) = S/(I_{\Delta}, p_1(\Gamma_n), \dots, p_n(\Gamma_n)).$$

In particular it follows that dim $S/(I_{\Delta}, p_1(\Delta), \dots, p_d(\Delta)) = 0$. This yields the desired conclusion.

Remark 3.2. Let $p_i \in S$ be defined as in (5), and let $J = (p_1, \ldots, p_n)$. Then for the reverse lexicographical order < we have

$$in_{\leq}(J) = (x_1, x_2^2, \dots, x_i^i, \dots, x_n^n).$$

Indeed, Theorem 3.1 implies that p_1, \ldots, p_n is a regular sequence, and since $x_1, x_2^2, \ldots, x_n^n$ is also a regular sequence and since deg $x_i^i = \deg p_i$ for $i = 1, \ldots, n$, we see that

$$\ell(S/J) = \ell(S/(x_1, x_2^2, \dots, x_n^n) \ge \ell(S/\inf_{\le}(J)) = \ell(S/J)$$

Hence, $\ell(S/(x_1, x_2^2, \dots, x_n^n)) = \ell(S/\operatorname{in}_{<}(J))$, which yields the desired conclusion.

Corollary 3.3. Let e be the number of facets F of Δ with |F| = d. Then $K[\Delta]$ is Cohen-Macaulay if and only if

$$\ell(S/(I_{\Delta}, p_1(\Delta), \dots, p_d(\Delta))) = d!e.$$

Proof. We note that e = e(R) is the multiplicity of $K[\Delta]$. Thus the assertion follows from Theorem 3.1 and Proposition 1.2.

Let Δ be a simplicial complex of dimension d-1. While d!e only depends on the simplicial complex Δ , in general the number $\ell(S/(I_{\Delta}, p_1(\Delta), \dots, p_d(\Delta)))$ also depends on the characteristic of the field K. This is not surprising since this is also the case for the Cohen–Macaulay property of $K[\Delta]$.

Corollary 3.3 can also be used as a computational tool to determine the depth of a Stanley–Reisner ring. We demonstrate this with the following example: consider the chessboard \mathcal{P}_n of size $n \times n$. The set of non-attacking rooks on \mathcal{P}_n is a simplicial complex which we denote $\Delta(\mathcal{P}_n)$. Fix a field K. For n > 1, the Stanley Reisner ring $K[\Delta(\mathcal{P}_n)]$ is not Cohen–Macaulay. Indeed if n = 2, then $\Delta(\mathcal{P}_n)$ is not connected and hence not Cohen–Macaulay and if n > 2, then for any face F with |F| = n - 2, $\operatorname{link}_{\Delta(\mathcal{P}_n)}(F) = \Delta(\mathcal{P}_2)$. Hence $\Delta(\mathcal{P}_n)$ can not be Cohen–Macaulay. But what is the depth of $K[\Delta(\mathcal{P}_n)]$? For n = 2, 3 the depth can be computed by using the depth command implemented in CoCoA. But already for n = 4, depth $K[\Delta(\mathcal{P}_4)]$ can not be computed by CoCoA. Instead we use the fact, first shown by Smith [14, Theorem 4.8], that for any simplicial complex Δ one has

depth
$$K[\Delta] = \max\{i \colon K[\Delta^{(i)}] \text{ is Cohen-Macaulay}\},\$$

where $\Delta^{(i)} = \langle F \in \Delta : |F| = i \rangle$ is the *i*th skeleton of Δ .

In order to obtain the Stanley–Reisner ideal of the (n-1)-skeleton of \mathcal{P}_n we have to add to $I_{\Delta(\mathcal{P}_n)}$ the monomials corresponding to the facets of \mathcal{P}_n . Then we use Corollary 3.3 to check the Cohen–Macaulayness of the (n-1)-skeleton. For example when n = 4, the calculation for $I_{\Delta(\mathcal{P}_4)^{(3)}}$ gives

$$\ell(S/(I_{\Delta(\mathcal{P}_4)^{(3)}}, p_1(\Delta(\mathcal{P}_4)^{(3)}), p_2(\Delta(\mathcal{P}_4)^{(3)}), p_3(\Delta(\mathcal{P}_4)^{(3)}))) = 6e(K[\Delta(\mathcal{P}_4)^{(3)}]).$$

The length on the left hand side can be computed by means of the multiplicity command of CoCoA. The output comes almost immediately. Hence, $K[\Delta(\mathcal{P}_4)^{(3)}]$ is Cohen–Macaulay and so depth $K[\Delta(\mathcal{P}_4)] = 3$.

References

- Alekseev, V.E.: An upper bound for the number of maximal independent sets in a graph. Discrete Math. Appl. 17, No. 4, 355—359 (2007)
- Bruns, W., Herzog, J.: Cohen-Macaulay rings. Cambridge studies in advanced mathematics.
 39. Cambridge University Press, London, Cambridge, New York (1993)
- [3] Bruns, W., Herzog, J.: Semigroup rings and simplicial complexes. J. Pure Appl. Algebra 122, 185–208 (1997)
- [4] Castrillòn, I.D., Cruz, R., Reyes, E.: On well-covered, vertex decomposable and Cohen-Macaulay graphs. Electron J. Combin. 23(2) (2016)
- [5] Eisenbud, D.: Commutative Algebra: with a view toward algebraic geometry. Vol. 150. Springer Science and Business Media (2013)
- [6] Fröberg, R.: On Stanley-Reisner rings. Banach Center Publications 26, 57–70 (1990)
- [7] Herzog, J.: Ein Cohen-Macaulay-Kriterium mit Anwendungen auf den Konormalenmodul und den Differentialmodul. Mathematische Zeitschrift. 163, 149–162 (1978)
- [8] Herzog, J., Hibi, T.: Monomial ideals. Graduate Texts in Mathematics. 260, Springer-Verlag London, Ltd., London (2011)
- [9] Hoang, D.T. and Trung, T. N.: Coverings, matchings and the number of maximal independent sets of graphs. Australasian Journal of Combinatorics. 73(3), 424—431 (2019)
- [10] Hochster, M.: Cohen-Macaulay rings, combinatorics, and simplicial complexes. Ring Theory II. Lect. Notes in Pure Appl. Math. 26, 171–223 (1977)
- [11] Katzmann, M.: Characteristic-independence of Betti numbers of graph ideals. J. Combin. Theory Ser. A 113, no. 3, 435-454 (2006)
- [12] Mahmoudi, M., Mousivand, A., Crupi, M., Rinaldo, G., Terai, N., and Yassemi, S.: Vertex decomposability and regularity of very well-covered graphs. Journal of Pure and Applied Algebra. 215(10), 2473–2480 (2011)
- [13] Serre, J.P.: Algèbre locale, multiplicités: cours au Collège de France, 1957-1958. 11, Springer Science & Business Media (1997)
- [14] Smith D.E.: On the Cohen-Macaulay property in commutative algebra and simplicial topology. Pacific J. Math. 141, 165–196 (1990)
- [15] Stanley, R.P.: Combinatorics and Commutative Algebra. Volume 41, Springer (1996)
- [16] Swanson, I., Huneke, C.: Integral closure of ideals, rings, and modules. 13, Cambridge University Press (2006)
- [17] Takayama, Y.: Combinatorial characterizations of generalized Cohen-Macaulay monomial ideals. Bulletin mathématique de la Société des Sciences Mathématiques de Roumanie, 327– 344 (2005)

Jürgen Herzog, Fachbereich Mathematik, Universität Duisburg-Essen, Campus Essen, 45117 Essen, Germany

E-mail address: juergen.herzog@uni-essen.de

Somayeh Moradi, Department of Mathematics, School of Science, Ilam University, P.O.Box 69315-516, Ilam, Iran

E-mail address: so.moradi@ilam.ac.ir