

Main Manuscript for

Reply to Zhang et al.: Slowing the rate of spread of COVID-19

Angeline G. Pendergrass^{1,2‡*}, Kristie L. Ebi³, and Micah B. Hahn⁴

¹ Institute for Atmospheric and Climate Science, ETH-Zurich, Zurich, Switzerland

² Climate and Global Dynamics Laboratory, National Center for Atmospheric Research, Boulder, CO, USA.

³ Department of Global Health, University of Washington, Seattle, WA, USA

⁴ Institute for Circumpolar Health Studies, University of Alaska-Anchorage, Anchorage, AK, USA

* Angeline G. Pendergrass

Email: apgrass@ucar.edu

AGP: https://orcid.org/0000-0003-2542-1461

KLE: https://orcid.org/0000-0003-4746-8236

MBH: https://orcid.org/0000-0002-7022-3918

This PDF file includes:

Main Text Figure 1

⁺ This work was done during personal time as it is outside the of her funded projects.

Main Text

Zhang et al. (1) reported that mandating face coverings in public is necessary to decrease the rate of new COVID-19 infections. A counterexample disproves this finding.

We agree with Zhang et al. that masks can be an important element in reducing virus transmission (2) and that widespread use may be essential for returning to full activity. However, misattributing the impacts of interventions during a pandemic can jeopardize lives.

Zhang et al.'s analyses assumed the number of cases increased at an approximately constant rate after lockdown (their Figs. 2b,c) as a function of government-mandated non-pharmaceutical interventions (NPIs). The conclusion drawn was that interventions that do not include requiring masks, "are insufficient by themselves" and will not decrease the rate of new cases, and that "mandated face covering represents the determinant in shaping the trends of the pandemic worldwide."

Counterexamples disprove these conclusions. Denmark, Norway, and Switzerland experienced decreases in the rate of new cases without requiring masks in public. The Swiss federal government implemented NPIs similar to those in Italy and New York City approximately two weeks after local transmission began (3, 4). However, unlike Italy and New York City, face coverings were not mandated for the general public. Interventions included closing schools and non-essential public-facing businesses and banning public and private events. The number of people who could gather was also limited to 1000 people starting February 28, and to 5 starting March 21 (4).


The rate of new cases in Switzerland peaked in late March and began to decline by mid-April (Fig. 1b), despite the lack of any requirement for or widespread use of masks in public (5, 6). If the assumptions underlying Zhang et al. were valid, then the number of cases should have continued to increase at a steady rate.

The decreasing rate is likely attributable to a number of factors, including the incubation period of the virus, the length of time between the onset of symptoms, when tests are taken and confirmed, and adherence to NPIs. Variation in the rate of new infections in response to an intervention can be quantified with SEIR and other equations (7). Shortfalls in the Zhang et al. analyses could have been avoided if the author team had included expertise in COVID-19 transmission dynamics.

Attributing the reduction in the rate of new cases entirely to mask requirements implies that the other measures had no effect and neglects the potential importance of physical distancing for limiting COVID-19 transmission. The evolution of COVID-19 in Switzerland in the spring of 2020 illustrates that decreasing the rate of new cases is achievable in some settings without mandating face coverings, disproving the conclusion reached by Zhang et al. A full suite of epidemiologic tools is necessary in these challenging times.

Acknowledgments

Dennis Hartmann, Reto Knutti, Paulo Ceppi, Flavio Lehner, Brian Smoliak and Stephen Po-Chedley provided feedback. Weekly smoother from James Annan. **Figure 1.** Evolution of COVID-19 in Switzerland in spring of 2020. (a) Cumulative confirmed cases (c.f. (1) Fig. 2b,c). (b) Daily new confirmed cases (c.f. (1) Fig. 3). Raw data are shown in bars, the same data with the weekly cycle removed and a 1-2-1 filter applied is shown by the line. Data from (8).



References

- R. Zhang, Y. Li, A. L. Zhang, Y. Wang, M. J. Molina, Identifying airborne transmission as the dominant route for the spread of COVID-19. *PNAS* (2020) https://doi.org/10.1073/pnas.2009637117 (June 15, 2020).
- 2. D. K. Chu, *et al.*, Physical distancing, face masks, and eye protection to prevent person-toperson transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. *The Lancet* (2020) https://doi.org/10.1016/S0140-6736(20)31142-9 (June 19, 2020).
- World Health Organization, "Coronavirus disease 2019 (COVID-19): Situation Report 42" (2020) (June 21, 2020).
- 4. Federal Office of Public Health, "COVID-19: Epidemiological interim report" (2020) (June 21, 2020).
- 5. D. Balmer, 94 Prozent der Pendler tragen keine Maske und die Fallzahlen steigen. *Tages- Anzeiger* (July 2, 2020).
- 6. swissinfo.ch/dos, Up to 94% of Swiss train travellers going without mask. *swissinfo.ch* (July 2, 2020).
- 7. H. W. Hethcote, The Mathematics of Infectious Diseases. SIAM Rev. 42, 599-653 (2000).
- 8. Federal Office of Public Health, New coronavirus: Situation in Switzerland Confirmed coronavirus infections, deaths, epidemiological situation (2020) (June 18, 2020).