

A Two-Timescale Stochastic Algorithm Framework for Bilevel Optimization: Complexity Analysis and Application to Actor-Critic

Mingyi Hong, Hoi-To Wai, Zhaoran Wang and Zhuoran Yang

December 22, 2020

Abstract

This paper analyzes a two-timescale stochastic algorithm framework for bilevel optimization. Bilevel optimization is a class of problems which exhibit a two-level structure, and its goal is to minimize an outer objective function with variables which are constrained to be the optimal solution to an (inner) optimization problem. We consider the case when the inner problem is unconstrained and strongly convex, while the outer problem is constrained and has a smooth objective function. We propose a two-timescale stochastic approximation (TTSA) algorithm for tackling such a bilevel problem. In the algorithm, a stochastic gradient update with a larger step size is used for the inner problem, while a projected stochastic gradient update with a smaller step size is used for the outer problem. We analyze the convergence rates for the TTSA algorithm under various settings: when the outer problem is strongly convex (resp. weakly convex), the TTSA algorithm finds an $\mathcal{O}(K^{-2/3})$ -optimal (resp. $\mathcal{O}(K^{-2/5})$ -stationary) solution, where K is the total iteration number. As an application, we show that a two-timescale natural actor-critic proximal policy optimization algorithm can be viewed as a special case of our TTSA framework. Importantly, the natural actor-critic algorithm is shown to converge at a rate of $\mathcal{O}(K^{-1/4})$ in terms of the gap in expected discounted reward compared to a global optimal policy.

1 Introduction

Consider bilevel optimization problems of the form:

$$\min_{x \in X \subseteq \mathbb{R}^{d_1}} \ell(x) := f(x, y^*(x)) \quad \text{subject to} \quad y^*(x) \in \arg \min_{y \in \mathbb{R}^{d_2}} g(x, y), \quad (1)$$

where $d_1, d_2 \geq 1$ are integers; X is a closed and convex subset of \mathbb{R}^{d_1} , $f : X \times \mathbb{R}^{d_2} \rightarrow \mathbb{R}$ and $g : X \times \mathbb{R}^{d_2} \rightarrow \mathbb{R}$ are continuously differentiable functions with respect to (w.r.t.) x, y . Problem (1) involves two optimization problems following a two-level structure. We refer to $\min_{y \in \mathbb{R}^{d_2}} g(x, y)$ as the *inner problem* whose solution depends on x , and $g(x, y)$ is called the *inner objective function*; $\min_{x \in X} \ell(x)$ is referred as the *outer problem*, which represents the outer objective function that we wish to minimize and $\ell(x) \equiv f(x, y^*(x))$ is called the *outer objective function*. Moreover, both f, g can be stochastic functions whose gradient may be difficult to compute. Despite being a non-convex stochastic problem in general, (1) has a wide range of applications, e.g., reinforcement learning, hyperparameter optimization, game theory, etc..

Table 1: Summary of the main results. SC stands for strongly convex, WC for weakly convex, C for convex; k is the iteration counter, K is the total number of iterations.

$\ell(x)$	CONSTRAINT	STEP SIZE (α_k, β_k)	RATE (OUTER)	RATE (INNER)
SC	$X \subseteq \mathbb{R}^{d_1}$	$\mathcal{O}(k^{-1}), \mathcal{O}(k^{-2/3})$	$\mathcal{O}(K^{-2/3})^\dagger$	$\mathcal{O}(K^{-2/3})^*$
C	$X \subseteq \mathbb{R}^{d_1}$	$\mathcal{O}(K^{-3/4}), \mathcal{O}(K^{-1/2})$	$\mathcal{O}(K^{-1/4})^\P$	$\mathcal{O}(K^{-1/2})^*$
WC	$X \subseteq \mathbb{R}^{d_1}$	$\mathcal{O}(K^{-3/5}), \mathcal{O}(K^{-2/5})$	$\mathcal{O}(K^{-2/5})^\#$	$\mathcal{O}(K^{-2/5})^*$

† in terms of $\|x^K - x^*\|^2$, where x^* is the optimal solution; * in terms of $\|y^K - y^*(x^{K-1})\|^2$, where $y^*(x^{K-1})$ is the optimal inner solution for fixed x^{K-1} ; ¶ measured using $\ell(x) - \ell(x^*)$; $^\#$ measured using distance to a fixed point with the Moreau proximal map $\hat{x}(\cdot)$; see (13).

Tackling (1) is challenging as it involves solving the inner and outer optimization problems *simultaneously*. Even in the simplest case when $\ell(x)$ and $g(x, y)$ are strongly convex in x, y , respectively, solving (1) is difficult. For instance, if we aim to minimize $\ell(x)$ via a gradient method, at any iterate $x^{\text{cur}} \in \mathbb{R}^{d_1}$ – applying the gradient method for (1) involves a double-loop algorithm that (a) solves the inner optimization problem $y^*(x^{\text{cur}}) = \arg \min_{y \in \mathbb{R}^{d_2}} g(x^{\text{cur}}, y)$ and then (b) evaluates the gradient as $\nabla \ell(x^{\text{cur}})$ based on the solution $y^*(x^{\text{cur}})$. Depending on the application, step (a) is usually accomplished by applying yet another gradient method for solving the inner problem (unless a closed form solution for $y^*(x^{\text{cur}})$ exists). In this way, the resulting algorithm necessitates a double-loop structure.

To this end, [22] and the references therein proposed a stochastic algorithm for (1) involving a *double-loop* update. During the iterations, the inner problem $\min_{y \in \mathbb{R}^{d_2}} g(x^{\text{cur}}, y)$ is solved using a stochastic gradient (SGD) method, with the solution denoted by $\hat{y}^*(x^{\text{cur}})$. Then, the outer problem is optimized with an SGD update using estimates of $\nabla f(x^{\text{cur}}, \hat{y}^*(x^{\text{cur}}))$. Such a double-loop algorithm is proven to converge to a stationary solution, yet a few practical issues lingers: *How to select step size and termination criterion for the inner loop? What if the (stochastic) gradients of the inner and outer problems are only revealed sequentially, or these problems are required to be updated at the same time (e.g., in sequential games)?*

To address the above issues, this paper investigates a *single-loop* stochastic algorithm for (1). Focusing on a class of the bilevel optimization (1) where the inner problem is unconstrained and strongly convex, and the outer objective function is smooth, our contributions are three-fold:

- We study a two-timescale stochastic approximation (TTSA) algorithm [7] for the concerned class of bilevel optimization. The TTSA algorithm updates both outer and inner solutions simultaneously, by using some cheap estimates of stochastic gradients of both inner and outer objectives. The algorithm guarantees convergence by improving the inner (resp., outer) solution with a larger (resp., smaller) step size, also known as using a faster (resp., slower) timescale.
- We analyze the expected convergence rates of the TTSA algorithm. Our results are summarized in Table 1. Our analysis is accomplished by building a set of *coupled inequalities* for the one-step update in TTSA. For strongly convex outer function, we show inequalities that couple between the outer and inner optimality gaps. For convex or weakly convex outer functions, we establish inequalities coupling between the difference of outer iterates, the optimality of function values, and the inner optimality gap. We also provide new and generic results for solving coupled

inequalities. The distinction of timescales between step sizes of the inner and outer updates plays a crucial role in our convergence analysis.

- As an application, we illustrate the application of our analysis results on a two-timescale natural actor-critic policy optimization algorithm with linear function approximation [27, 40]. The natural actor-critic algorithm converges at the rate $\mathcal{O}(K^{-1/4})$ to an optimal policy, which is comparable to the state-of-the-art results.

The rest of this paper is organized as follows. §2 formally describes the problem setting of bilevel optimization and specify the problem class of interest. In addition, the TTSA algorithm is introduced and some application examples are discussed. §3 presents the main convergence results for the generic bilevel optimization problem (1). The convergence analysis is also presented where we highlight the main proof techniques used. Lastly, §4 discusses the application to reinforcement learning. Notice that technical details of the proof have been relegated to the online appendix [23].

1.1 Related Works

The study of bilevel optimization problem (1) can be traced to that of Stackelberg games [46], where the outer (resp. inner) problem optimizes the action taken by a leader (resp. the follower). In the optimization literature, bilevel optimization was introduced in [10] for resource allocation problems, and later studied in [9]. Furthermore, bilevel optimization is a special case of the broader class problem of Mathematical Programming with Equilibrium Constraints [35].

Many related algorithms have been proposed for bilevel optimization. To list a few, this includes approximate descent methods [18, 50], and penalty-based method [24, 52]. The approximate descent methods deal with a subclass of problem where the outer problem possesses certain (local) differentiability property, while the penalty-based methods approximate the inner problems and/or the outer problems with an appropriate penalty functions. It is noted in [12] that descent based method has relatively strong assumptions about the inner problem (such as non-degeneracy), while the penalty based methods are typically slow. Moreover, these works typically focus on asymptotic convergence analysis, without characterizing the convergence rates; see [12] for a comprehensive survey.

In [13, 22, 25], the authors considered bi-level problems in the (stochastic) unconstrained setting, when the outer problem is non-convex and the inner problem is strongly (or strictly) convex. These works are more related to the algorithms and results to be developed in the current paper. In this case, the (stochastic) gradient of the outer problem may be computed using the chain rule. However, to obtain an accurate estimate, one has to either use *double loop* structure where the inner loop solves the inner sub-problem to a high accuracy [13, 22], or use a large batch-size (e.g., $\mathcal{O}(1/\epsilon)$) [25]. Both of these methods could be difficult to implement in practice. In reference [43], the authors analyzed a special bi-level problem where there is a single optimization variable in both outer and inner levels. The authors proposed a Sequential Averaging Method (SAM) algorithm which can provably solve a problem with strongly convex outer problem, and convex inner problems. Building upon the SAM, [32, 34] developed first-order algorithms for bi-level problem, without requiring that for each fixed outer-level variable, the inner-level solution must be a singleton.

In a different line of recent works, references [33, 45] proposed and analyzed different versions of the so-called truncated back-propagation approach for approximating the (stochastic) gradient

of the outer-problem, and established convergence for the respective algorithms. The idea is to use a dynamical system to model an optimization algorithm that solves the inner problem, and then replace the optimal solution of the inner problem by unrolling a few iterations of the updates. However, computing the hyper-gradient of the outer problem requires using back-propagation through the optimization algorithm, and can be computationally very expensive. It is important to note that none of the methods discussed above have considered *single-loop* stochastic algorithms, in which a small batch of samples are used to approximate the inner and outer gradients at each iteration. Later we will see that the ability of being able to update using a small number of samples for both outer and inner problems is critical in a number of applications.

In contrast to the above mentioned works, this paper considers an TTSA algorithm for stochastic bilevel optimization, which is a single-loop algorithm employing cheap stochastic estimates of the gradient. Notice that TTSA [7] is a class of algorithms designed to solve coupled system of (nonlinear) equations. While its asymptotic convergence property has been well understood, e.g., [7, 8, 29], the convergence rate analysis have been focused on *linear* cases, e.g., [14, 28, 31]. In general, the bilevel optimization problem (1) requires a nonlinear TTSA algorithm. For this case, an asymptotic convergence rate is analyzed in [38] under a restricted form of nonlinearity. For convergence rate analysis, [43] considered a single-loop algorithm for deterministic bilevel optimization with only one variable, and [17] studied the convergence rate when the expected updates are strongly monotone.

Finally, it is worthwhile to mention that various forms of TTSA have been applied to tackle compositional stochastic optimization [51], policy evaluation methods [6, 49], and actor-critic methods [5, 30, 36]. Notice that some of these optimization problems can be cast as a bilevel optimization, as we will demonstrate next.

Notations Unless otherwise noted, $\|\cdot\|$ is the Euclidean norm on finite dimensional Euclidean space. For a twice differentiable function $f : X \times Y \rightarrow \mathbb{R}$, $\nabla_x f(x, y)$ (resp. $\nabla_y f(x, y)$) denotes its partial gradient taken w.r.t. x (resp. y), and $\nabla_{yx}^2 f(x, y)$ (resp. $\nabla_{xy}^2 f(x, y)$) denotes the Jacobian of $\nabla_y f(x, y)$ at y (resp. $\nabla_x f(x, y)$ at x).

2 Two-Timescale Stochastic Approximation Algorithm for (1)

To formally define the problem class of interest, we state the following conditions on the bilevel optimization problem (1).

Assumption 1. *The outer functions $f(x, y)$ and $\ell(x) := f(x, y^*(x))$ satisfy:*

1. *For any $x \in \mathbb{R}^{d_1}$, $\nabla_x f(x, \cdot)$ and $\nabla_y f(x, \cdot)$ are Lipschitz continuous with respect to (w.r.t.) $y \in \mathbb{R}^{d_2}$, and with constants L_{fx} and L_{fy} , respectively.*
2. *For any $y \in \mathbb{R}^{d_2}$, $\nabla_y f(\cdot, y)$ is Lipschitz continuous w.r.t. $x \in X$, and with constants \bar{L}_{fy} .*
3. *For any $x \in X, y \in \mathbb{R}^{d_2}$, we have $\|\nabla_y f(x, y)\| \leq C_{fy}$, for some $C_{fy} > 0$.*
4. *The function $\ell(\cdot)$ is weakly convex with modulus μ_ℓ , i.e.,*

$$\ell(w) \geq \ell(v) + \langle \nabla \ell(v), w - v \rangle + \mu_\ell \|w - v\|^2, \quad \forall w, v \in X. \quad (2)$$

Assumption 2. *The inner function $g(x, y)$ satisfies:*

1. For any $x \in X$ and $y \in \mathbb{R}^{d_2}$, $g(x, y)$ is twice continuously differentiable in (x, y) ;
2. For any $x \in X$, $\nabla_y g(x, \cdot)$ is Lipschitz continuous w.r.t. $y \in \mathbb{R}^{d_2}$, and with constants L_g .
3. For any $x \in X$, $g(x, \cdot)$ is strongly convex in y , and with modulus $\mu_g > 0$.
4. For any $x \in X$, $\nabla_{xy}^2 g(x, \cdot)$ and $\nabla_{yy}^2 g(x, \cdot)$ are Lipschitz continuous w.r.t. $y \in \mathbb{R}^{d_2}$, and with constants $L_{gxy} > 0$ and $L_{gyy} > 0$, respectively.
5. For any $x \in X$ and $y \in \mathbb{R}^{d_2}$, we have $\|\nabla_{xy}^2 g(x, y)\| \leq C_{gxy}$ for some $C_{gxy} > 0$.
6. For any $y \in \mathbb{R}^m$, $\nabla_{xy}^2 g(\cdot, y)$ and $\nabla_{yy}^2 g(\cdot, y)$ are Lipschitz continuous w.r.t. $x \in X$, and with constants $\bar{L}_{gxy} > 0$ and $\bar{L}_{gyy} > 0$, respectively.

Basically, Assumption 1, 2 require that the inner and outer functions f, g are well-behaved. In particular, $\nabla_x f$, $\nabla_y f$, $\nabla_{xy}^2 g$, and $\nabla_{yy}^2 g$ are Lipschitz continuous w.r.t. x when y is fixed, and Lipschitz continuous w.r.t. y when x is fixed.

Our first endeavor is to develop a single-loop stochastic algorithm for tackling (1). Focusing on solutions which satisfy the first-order stationary condition of (1), we aim at finding a pair of solution (x^*, y^*) such that

$$\nabla_y g(x^*, y^*) = 0, \quad \langle \nabla \ell(x^*), x - x^* \rangle \geq 0, \quad \forall x \in X. \quad (3)$$

Given x^* , a solution y^* satisfying the first condition in (3) may be found by a cheap stochastic gradient recursion such as $y \leftarrow y - \beta h_g$ with $\mathbb{E}[h_g] = \nabla_y g(x^*, y)$. On the other hand, given $y^*(x)$ and suppose that we can obtain a cheap stochastic gradient estimate h_f with $\mathbb{E}[h_f] = \nabla \ell(x) = \bar{\nabla}_x f(x, y^*(x))$, where $\bar{\nabla}_x f(x, y)$ is a surrogate for $\nabla \ell(x)$ (to be described later), then the second condition can be satisfied by a simple projected stochastic gradient recursion as $x \leftarrow P_X(x - \alpha h_f)$, where $P_X(\cdot)$ denotes the Euclidean projection onto X .

A challenge in designing a *single-loop* algorithm for satisfying (3) is to ensure that the outer function's gradient $\bar{\nabla}_x f(x, y)$ is evaluated at an inner solution y that is close to $y^*(x)$. This led us to develop a *two-timescale stochastic approximation* (TTSA) [7] framework, as summarized in Algorithm 1. An important feature is that the algorithm utilizes two step sizes α_k, β_k for the outer (x^k), inner (y^k) solution, respectively, designed with different timescales as $\alpha_k/\beta_k \rightarrow 0$. As a larger step size is taken to optimize y^k , the latter shall stay close to $y^*(x^k)$. Using this strategy, it is expected that y^k will converge to $y^*(x^k)$ asymptotically.

Inspired by [22], we provide a method for computing a surrogate of $\nabla \ell(x)$ given y with general objective functions satisfying Assumption 1, 2. Given $y^*(x)$, we observe that using chain rule, the gradient of $\ell(x)$ can be derived as

$$\nabla \ell(x) = \nabla_x f(x, y^*(x)) - \nabla_{xy}^2 g(x, y^*(x)) [\nabla_{yy}^2 g(x, y^*(x))]^{-1} \nabla_y f(x, y^*(x)). \quad (4)$$

We may now define $\bar{\nabla}_x f(x, y)$ as a surrogate of $\nabla \ell(x)$ by replacing $y^*(x)$ with $y \in \mathbb{R}^{d_2}$:

$$\bar{\nabla}_x f(x, y) := \nabla_x f(x, y) - \nabla_{xy}^2 g(x, y) [\nabla_{yy}^2 g(x, y)]^{-1} \nabla_y f(x, y). \quad (5)$$

Notice that $\nabla \ell(x) = \bar{\nabla}_x f(x, y^*(x))$. Eq. (5) is a surrogate for $\nabla \ell(x)$ that may be used in TTSA. We emphasize that (5) is not the only construction and the TTSA can accommodate other forms of gradient surrogate.

Algorithm 1. Two-Timescale Stochastic Approximation (TTSA)

S0) Initialize the variable $(x^0, y^0) \in X \times \mathbb{R}^{d_2}$ and the step size sequence $\{\alpha_k, \beta_k\}_{k \geq 0}$;
S1) For iteration $k = 0, \dots, K$,

$$y^{k+1} = y^k - \beta_k \cdot h_g^k, \quad (6a)$$

$$x^{k+1} = P_X(x^k - \alpha_k \cdot h_f^k), \quad (6b)$$

where h_g^k, h_f^k are stochastic estimates of $\nabla_y g(x^k, y^k), \bar{\nabla}_x f(x^k, y^{k+1})$, respectively, and $P_X(\cdot)$ is the Euclidean projection operator onto the convex set X .

Let $\mathcal{F}_k := \sigma\{y^0, x^0, \dots, y^k, x^k\}$, $\mathcal{F}'_k := \sigma\{y^0, x^0, \dots, y^k, x^k, y^{k+1}\}$ be the filtration of the random variables up to iteration k , where $\sigma\{\cdot\}$ denotes the σ -algebra generated by the random variables. We consider the following assumption regarding h_f^k, h_g^k :

Assumption 3. For any $k \geq 0$, there exist constants σ_g, σ_f , and a nonincreasing sequence $\{b_k\}_{k \geq 0}$ such that:

$$\mathbb{E}[h_g^k | \mathcal{F}_k] = \nabla_y g(x^k, y^k), \quad \mathbb{E}[h_f^k | \mathcal{F}'_k] = \bar{\nabla}_x f(x^k, y^{k+1}) + B_k, \quad \|B_k\| \leq b_k, \quad (7a)$$

$$\mathbb{E}[\|h_g^k - \nabla_y g(x^k, y^k)\|^2 | \mathcal{F}_k] \leq \sigma_g^2 \cdot \{1 + \|\nabla_y g(x^k, y^k)\|^2\}, \quad (7b)$$

$$\mathbb{E}[\|h_f^k - B_k - \bar{\nabla}_x f(x^k, y^{k+1})\|^2 | \mathcal{F}'_k] \leq \sigma_f^2. \quad (7c)$$

Notice that the conditions on h_g^k are standard when the latter is taken as a stochastic gradient of $g(x^k, y^k)$, while h_f^k is a potentially biased estimate of $\bar{\nabla}_x f(x^k, y^{k+1})$. As we will see in our convergence analysis, the bias shall decay polynomially to zero. In light of (5), it is possible to design a practical scheme for constructing h_f^k with $\mathcal{O}(1 + \log k)$ stochastic oracle calls to satisfy such requirements; we relegate the details to §E.4 of our online version [23].

To further understand the property of the TTSA algorithm with (5), we borrow the following results from [22] on the Lipschitz continuity of the maps $\nabla \ell(x), y^*(x)$:

Lemma 1. [22, Lemma 2.2] Under Assumption 1, 2. It holds

$$\|\bar{\nabla}_x f(x, y) - \nabla \ell(x)\| \leq L\|y^*(x) - y\|, \quad \|y^*(x_1) - y^*(x_2)\| \leq L_y\|x_1 - x_2\|, \quad (8a)$$

$$\|\nabla \ell(x_1) - \nabla \ell(x_2)\| = \|\nabla f(x_1, y^*(x_1)) - \nabla f(x_2, y^*(x_2))\| \leq L_f\|x_1 - x_2\|. \quad (8b)$$

for any $x, x_1, x_2 \in X$ and $y \in \mathbb{R}^{d_2}$, where we have defined

$$L := L_{f_x} + \frac{L_{f_y} C_{g_{xy}}}{\mu_g} + C_{f_y} \left(\frac{L_{g_{xy}}}{\mu_g} + \frac{L_{g_{yy}} C_{g_{xy}}}{\mu_g^2} \right), \quad (9)$$

$$L_f := L_{f_x} + \frac{(\bar{L}_{f_y} + L) C_{g_{xy}}}{\mu_g} + C_{f_y} \left(\frac{\bar{L}_{g_{xy}}}{\mu_g} + \frac{\bar{L}_{g_{yy}} C_{g_{xy}}}{\mu_g^2} \right), \quad L_y = \frac{C_g}{\mu_g}.$$

The above properties will be pivotal in establishing the convergence of TTSA. Notice that assumption (7c) combined with Lemma 1 leads to the following estimate:

$$\mathbb{E}[\|h_f^k\|^2 | \mathcal{F}'_k] \leq \tilde{\sigma}_f^2 + 3b_k^2 + 3L^2\|y^{k+1} - y^*(x^k)\|^2, \quad \tilde{\sigma}_f^2 := \sigma_f^2 + 3 \sup_{x \in X} \|\nabla \ell(x)\|^2. \quad (10)$$

Throughout, we assume $\tilde{\sigma}_f^2$ is bounded, e.g., it can be satisfied if X is bounded, or if $\ell(x)$ has uniformly bounded gradient.

2.1 Applications

Practical problems such as hyperparameter optimization [21, 37, 45], Stackelberg games [46] can be cast into special cases of bilevel optimization problem (1) satisfying the aforementioned assumptions. To be specific, we discuss two applications of the bilevel optimization problem (1) below.

Model-Agnostic Meta-Learning An important paradigm of machine learning is to find model that adapts to multiple training sets in order to achieve the best performance for individual tasks. Among others, a popular formulation is model-agnostic meta learning (MAML) [19] which minimizes an outer objective of empirical risk on all training sets, and the inner objective is the one-step projected gradient. Let $D^{(j)} = \{z_i^{(j)}\}_{i=1}^n$ be the j -th ($j \in [J]$) training set with sample size n , MAML can be formulated as a bilevel optimization problem [42]:

$$\begin{aligned} \min_{\theta \in \Theta} \quad & \sum_{j=1}^J \sum_{i=1}^n \bar{\ell}(\theta^{*(j)}(\theta), z_i^{(j)}) \\ \text{subject to} \quad & \theta^{*(j)}(\theta) \in \arg \min_{\theta^{(j)}} \left\{ \sum_{i=1}^n \langle \theta^{(j)}, \nabla_{\theta} \bar{\ell}(\theta, z_i^{(j)}) \rangle + \frac{\lambda}{2} \|\theta^{(j)} - \theta\|^2 \right\}. \end{aligned} \tag{11}$$

Here θ is the shared model parameter, $\theta^{(j)}$ is the adaptation of θ to the j th training set, and $\bar{\ell}$ is the loss function. It can be checked that the inner problem is strongly convex. Moreover, [20] proved that, assuming λ is sufficiently large and $\bar{\ell}$ is strongly convex, the outer problem is also strongly convex (H1). We have H1, H2.(b), and H3 for stochastic gradient updates, assuming $\bar{\ell}$ is sufficiently regular, and the losses are the logistic loss. In fact, [41] demonstrated that an algorithm with no inner loop achieves a comparable performance to [20].

Policy Optimization Another application of the bilevel optimization problem is the policy optimization problem, particularly when combined with an actor-critic scheme. The optimization involved is to find an optimal policy to maximize the expected (discounted) reward. Here, the ‘actor’ serves as the outer problem and the ‘critic’ serves as the inner problem which evaluates the performance of the ‘actor’ (current policy). To avoid redundancy, we refer our readers to §4 where we present a detailed case study. The latter will also shed lights on the generality of our proof techniques for TTSA algorithms.

3 Main Results

This section presents the convergence results for TTSA algorithm for (1). We first discuss a few concepts pivotal to our analysis.

Tracking Error TTSA tackles the inner and outer problems simultaneously using single-loop updates. Due to the coupled nature of the inner and outer problems, in order to obtain an upper

bound on the *optimality gap* $\Delta_x^k := \mathbb{E}[\|x^k - x^*\|^2]$, where x^* is an optimal solution to (1), we need to estimate the *tracking error* defined as

$$\Delta_y^k := \mathbb{E}[\|y^k - y^*(x^{k-1})\|^2] \quad \text{where} \quad y^*(x) = \arg \min_{y \in \mathbb{R}^{d_2}} g(x, y). \quad (12)$$

For any $x \in X$, $y^*(x)$ is well defined since the inner problem is strongly convex due to Assumption 2. By definition, Δ_y^k quantifies how close y^k is from the optimal solution to inner problem given x^{k-1} .

Moreau Envelop Fix $\rho > 0$, define the Moreau envelop and proximal map as

$$\Phi_{1/\rho}(z) := \min_{x \in X} \{\ell(x) + (\rho/2)\|x - z\|^2\}, \quad \hat{x}(z) := \arg \min_{x \in X} \{\ell(x) + (\rho/2)\|x - z\|^2\}. \quad (13)$$

For any $\epsilon > 0$, $x^k \in X$ is said to be an ϵ -*nearly stationary solution* [16] if x^k is an approximate fixed point of $\{\hat{x} - \text{I}\}(\cdot)$, where

$$\tilde{\Delta}_x^k := \mathbb{E}[\|\hat{x}(x^k) - x^k\|^2] \leq \rho^2 \cdot \epsilon. \quad (14)$$

We observe that if $\epsilon = 0$, then $x^k \in X$ is a stationary solution to (1) satisfying the second condition in (3). As we will demonstrate next, the *near-stationarity* condition (14) provides an apparatus to quantify the finite-time convergence of TTSA in the case when $\ell(x)$ is non-convex.

3.1 Strongly Convex Outer Objective Function

Our first result considers the instance of (1) where $\ell(x)$ is strongly convex. We obtain:

Theorem 1. *Under Assumption 1 with $\mu_\ell > 0$, Assumption 2, 3. Assume that the step sizes satisfy*

$$\alpha_k \leq c_0 \beta_k^{3/2}, \quad \beta_k \leq c_1 \alpha_k^{2/3}, \quad \frac{\beta_{k-1}}{\beta_k} \leq 1 + \beta_k \mu_g / 8, \quad \frac{\alpha_{k-1}}{\alpha_k} \leq 1 + 3\alpha_k \mu_\ell / 4, \quad (15a)$$

$$\alpha_k \leq \frac{1}{\mu_\ell}, \quad \beta_k \leq \min \left\{ \frac{1}{\mu_g}, \frac{\mu_g}{L_g^2(1 + \sigma_g^2)}, \frac{\mu_g^2}{48c_0^2 L^2 L_y^2} \right\}, \quad 8\mu_\ell \alpha_k \leq \mu_g \beta_k, \quad \forall k \geq 0, \quad (15b)$$

where the constants L, L_y were defined in Lemma 1 and we have $c_0, c_1 > 0$. The bias satisfies $b_k^2 \leq \tilde{c}_b \alpha_{k+1}$. For any $k \geq 1$, the TTSA iterates satisfy

$$\begin{aligned} \Delta_x^k &\lesssim \prod_{i=0}^{k-1} (1 - \alpha_i \mu_\ell) \left[\Delta_x^0 + \frac{L^2}{\mu_\ell^2} \Delta_y^0 \right] + \frac{c_1 L^2}{\mu_\ell^2} \left[\frac{\sigma_g^2}{\mu_g} + \frac{c_0^2 L_y^2}{\mu_g^2} \tilde{\sigma}_f^2 \right] \alpha_{k-1}^{2/3}, \\ \Delta_y^k &\lesssim \prod_{i=0}^{k-1} (1 - \beta_i \mu_g / 4) \Delta_y^0 + \left[\frac{\sigma_g^2}{\mu_g} + \frac{c_0^2 L_y^2}{\mu_g^2} \tilde{\sigma}_f^2 \right] \beta_{k-1}, \end{aligned} \quad (16)$$

where the symbol \lesssim denotes that the numerical constants are omitted.

We emphasize that the conditions in (15) are satisfied by both *constant* and *diminishing* step sizes. For example, we set $\alpha_k = c_\alpha / (k + k_\alpha), \beta_k = c_\beta / (k + k_\beta)^{2/3}$ with

$$k_\alpha = \max \left\{ 35 \left(\frac{L_g}{\mu_g} \right)^3 (1 + \sigma_g^2)^{\frac{3}{2}}, \frac{(512)^{\frac{3}{2}} L^2 L_y^2}{\mu_\ell^2} \right\}, \quad c_\alpha = \frac{8}{3\mu_\ell}, \quad k_\beta = \frac{1}{4} k_\alpha, \quad c_\beta = \frac{32}{3\mu_g}. \quad (17)$$

Then, the step sizes satisfy (15) with $c_0 = \frac{\mu_g^{3/2}}{\mu_\ell}$, $c_1 = 10\frac{\mu_\ell^{2/3}}{\mu_g}$. Theorem 1 then shows the last iterate convergence rate for the optimality gap and the tracking error to be $\mathcal{O}(k^{-2/3})$. To compute an ϵ -optimal solution with $\Delta_x^k \leq \epsilon$, the TTSA algorithm requires $\mathcal{O}(\log(1/\epsilon)/\epsilon^{3/2})$ calls of stochastic oracles of the inner and outer gradients¹.

While this is arguably the easiest case of (1), we notice that the double-loop algorithm in [22] requires $\mathcal{O}(1/\epsilon)$, $\mathcal{O}(1/\epsilon^2)$ stochastic oracles for the outer, inner functions, respectively. As such, the TTSA algorithm requires less number of stochastic oracles for the inner function.

3.2 Smooth (Possibly Non-convex) Outer Objective Function

We focus on the case where $\ell(x)$ is weakly convex. We obtain

Theorem 2. *Under Assumption 1, 2, 3. Let $K \geq 1$ be the maximum iteration number and set*

$$\alpha = \min \left\{ \frac{\mu_g^2}{8L_y L L_g^2(1 + \sigma_g^2)}, \frac{1}{4L_y L} K^{-3/5} \right\}, \quad \beta = \min \left\{ \frac{\mu_g}{L_g^2(1 + \sigma_g^2)}, \frac{2}{\mu_g} K^{-2/5} \right\}. \quad (18)$$

Assume $b_k^2 \leq \alpha$. For any $K \geq 1$, the iterates from the TTSA algorithm satisfy

$$\mathbb{E}[\tilde{\Delta}_x^K] = \mathcal{O}(K^{-2/5}), \quad \mathbb{E}[\Delta_y^{K+1}] = \mathcal{O}(K^{-2/5}), \quad (19)$$

where K is an independent uniformly distributed random variable on $\{0, \dots, K-1\}$; and we recall $\tilde{\Delta}_x^k := \|\hat{x}(x^k) - x^k\|^2$. When K is large and $\mu_\ell < 0$, setting $\rho = 2|\mu_\ell|$ yields

$$\mathbb{E}[\tilde{\Delta}_x^K] \lesssim \left[L^2 \left(\Delta^0 + \frac{\sigma_g^2}{\mu_g^2} \right) + \mu_g \tilde{\sigma}_f^2 \right] \frac{K^{-2/5}}{|\mu_\ell|^2}, \quad \mathbb{E}[\Delta_y^{K+1}] \lesssim \left[\frac{\Delta^0}{\mu_g} + \frac{\sigma_g^2}{\mu_g^2} + \frac{\mu_g \sigma_f^2}{L^2} \right] K^{-2/5}, \quad (20)$$

where we defined $\Delta^0 := \max\{\Phi_{1/\rho}(x^0), \text{OPT}^0, \Delta_y^0\}$, and used the approximations $L \approx L_y$, $\alpha < 1$, $\mu_g \ll 1$; the symbol \lesssim denotes that the numerical constants are omitted.

The above result uses constant step sizes determined by the maximum number of iterations, K . Here we set the step sizes as $\alpha_k \asymp K^{-3/5}$ and $\beta_k \asymp K^{-2/5}$. Similar to the previous case, α_k/β_k converges to zero as K goes to infinity. Nevertheless, Theorem 2 shows that TTSA requires $\mathcal{O}(\epsilon^{-5/2} \log(1/\epsilon))$ calls of stochastic oracle for sampled gradient/Hessian to find an ϵ -nearly stationary solution. In addition, it is worth noting that when $X = \mathbb{R}^{d_1}$, Theorem 2 implies that TTSA achieves $\mathbb{E}[\|\nabla \ell(x^K)\|^2] = \mathcal{O}(K^{-2/5})$, i.e., x^K is an $\mathcal{O}(K^{-2/5})$ -approximate stationary point of $\ell(x)$ in expectation.

Let us compare our sampling complexity bounds to the double loop algorithm in [22], which requires $\mathcal{O}(\epsilon^{-3})$ (resp. $\mathcal{O}(\epsilon^{-2})$) stochastic oracle calls for the inner problem (resp. outer problem), to reach an ϵ -stationary solution. The sample complexity of TTSA yields a tradeoff for the inner and outer stochastic oracles. We also observe that a trivial extension to a single-loop algorithm results in a *constant* error bound². Finally, we can extend Theorem 2 to the case where $\ell(\cdot)$ is a convex function.

¹Each stochastic oracle call requires a stochastic gradient/Hessian sample. The polynomial bias decay requires $\mathcal{O}(1 + \log(k))$ calls to the oracle, justifying the log factor in the bound.

²To see this, the readers are referred to [22, Theorem 3.1]. If a single inner iteration is performed, $t_k = 1$, so $\bar{A}^k \geq \|y^0 - y^*(x^k)\|$ which is a constant. Then the r.h.s. of (3.70), (3.73), (3.74) in [22, Theorem 3.1] will all have a constant term.

Corollary 1. Under Assumption 1 with $\mu_\ell \geq 0$, Assumption 2, 3. Consider (1) with $X \subseteq \mathbb{R}^{d_1}$ such that $D_x = \sup_{x, x' \in X} \|x - x'\| < \infty$. Let $K \geq 1$ be the maximum iteration number and set

$$\alpha = \min \left\{ \frac{\mu_g^2}{8L_y L L_g^2 (1 + \sigma_g^2)}, \frac{1}{4L_y L} K^{-3/4} \right\}, \quad \beta = \min \left\{ \frac{\mu_g}{L_g^2 (1 + \sigma_g^2)}, \frac{2}{\mu_g} K^{-1/2} \right\}. \quad (21)$$

Assume that $b_k \leq c_b K^{-1/4}$. For large K , the TTSA algorithm satisfies

$$\mathbb{E}[\ell(x^K) - \ell(x^*)] = \mathcal{O}(K^{-1/4}), \quad \mathbb{E}[\Delta_y^{K+1}] = \mathcal{O}(K^{-1/2}), \quad (22)$$

where K is an independent uniform random variable on $\{0, \dots, K-1\}$. By convexity, the above implies $\mathbb{E}[\ell(\frac{1}{K} \sum_{k=1}^K x^k) - \ell(x^*)] = \mathcal{O}(K^{-1/4})$.

3.3 Convergence Analysis

We now present the proofs for Theorem 1, 2. The proof for Corollary 1 is similar to that of Theorem 2. Due to the space limitation, we refer the readers to [23]. We highlight that the proofs of both theorems rely on the similar ideas of tackling coupled inequalities.

Proof of Theorem 1 Our proof relies on bounding the optimality gap and tracking error *coupled with each other*. First we derive the convergence of the inner problem.

Lemma 2. Under Assumption 1, 2, 3. Suppose that the step size satisfies (15a), (15b). For any $k \geq 1$, it holds that

$$\Delta_y^{k+1} \leq \prod_{\ell=0}^k (1 - \beta_\ell \mu_g / 2) \Delta_y^0 + \frac{8}{\mu_g} \left\{ \sigma_g^2 + \frac{4c_0^2 L_y^2}{\mu_g} [\tilde{\sigma}_f^2 + 3b_0^2] \right\} \beta_k. \quad (23)$$

Notice that the bound in (23) relies on the strong convexity of the inner problem and the Lipschitz properties established in Lemma 1 for $y^*(x)$; see §A.1. We emphasize that the step size condition $\alpha_k \leq c_0 \beta_k^{3/2}$ is crucial in establishing the above bound. As the second step, we bound the convergence of the outer problem.

Lemma 3. Under Assumption 1 with $\mu_\ell > 0$, Assumption 2, 3. Assume that the bias satisfies $b_k^2 \leq \tilde{c}_b \alpha_{k+1}$. With (15a), (15b), for any $k \geq 1$, it holds that

$$\begin{aligned} \Delta_x^{k+1} &\leq \prod_{\ell=0}^k (1 - \alpha_\ell \mu_\ell) \Delta_x^0 + \left[\frac{4\tilde{c}_b}{\mu_\ell^2} + \frac{2\tilde{\sigma}_f^2 + 6b_0^2}{\mu_\ell} \right] \alpha_k \\ &\quad + \left[\frac{2L^2}{\mu_\ell} + 3\alpha_0 L^2 \right] \sum_{j=0}^k \alpha_j \prod_{\ell=j+1}^k (1 - \alpha_\ell \mu_\ell) \Delta_y^{j+1}. \end{aligned} \quad (24)$$

see §A.2. We observe that (23), (24) lead to a pair of coupled inequalities. To compute the final bound in the theorem, we substitute (23) into (24). As $\Delta_y^{j+1} = \mathcal{O}(\beta_j) = \mathcal{O}(\alpha_j^{2/3})$, the dominating term in (24) can be estimated as

$$\sum_{j=0}^k \alpha_j \prod_{\ell=j+1}^k (1 - \alpha_\ell \mu_\ell) \Delta_y^{j+1} = \sum_{j=0}^k \mathcal{O}(\alpha_j^{5/3}) \prod_{\ell=j+1}^k (1 - \alpha_\ell \mu_\ell) = \mathcal{O}(\alpha_k^{2/3}), \quad (25)$$

yielding the desirable rates in the theorem. See §A.3 for details.

Proof of Theorem 2 Without strong convexity in the outer problem, the analysis becomes more challenging. To this end, we first develop the following lemma on coupled inequalities with numerical sequences, which will be pivotal to our analysis:

Lemma 4. *Let $K \geq 1$ be an integer. Consider sequences of non-negative scalars $\{\Omega^k\}_{k=0}^K$, $\{\Upsilon^k\}_{k=0}^K$, $\{\Theta^k\}_{k=0}^K$. Let $c_0, c_1, c_2, d_0, d_1, d_2$ be some positive constants. If the recursion holds*

$$\Omega^{k+1} \leq \Omega^k - c_0 \Theta^{k+1} + c_1 \Upsilon^{k+1} + c_2, \quad \Upsilon^{k+1} \leq (1 - d_0) \Upsilon^k + d_1 \Theta^k + d_2, \quad (26)$$

for any $k \geq 0$. Then provided that $c_0 - c_1 d_1 (d_0)^{-1} > 0, d_0 - d_1 c_1 (c_0)^{-1} > 0$, it holds

$$\begin{aligned} \frac{1}{K} \sum_{k=1}^K \Theta^k &\leq \frac{\Omega^0 + \frac{c_1}{d_0} (\Upsilon^0 + d_1 \Theta^0 + d_2)}{(c_0 - c_1 d_1 (d_0)^{-1}) K} + \frac{c_2 + c_1 d_2 (d_0)^{-1}}{c_0 - c_1 d_1 (d_0)^{-1}} \\ \frac{1}{K} \sum_{k=1}^K \Upsilon^k &\leq \frac{\Upsilon^0 + d_1 \Theta^0 + d_2 + \frac{d_1}{c_0} \Omega^0}{(d_0 - d_1 c_1 (c_0)^{-1}) K} + \frac{d_2 + d_1 c_2 (c_0)^{-1}}{d_0 - d_1 c_1 (c_0)^{-1}}. \end{aligned} \quad (27)$$

The proof of the above lemma is simple and is relegated to §B.1.

We demonstrate that stationarity measures of the TTSA iterates satisfy (26). The conditions $c_0 - c_1 d_1 (d_0)^{-1} > 0, d_0 - d_1 c_1 (c_0)^{-1} > 0$ impose constraints on the step sizes and (27) leads to a finite-time bound on the convergence of TTSA. To begin our derivation of Theorem 2, we observe the following coupled descent lemma:

Lemma 5. *Under Assumption 1, 2, 3. If $\mu_g \beta / 2 < 1, \beta L_g^2 (1 + \sigma_g^2) \leq \mu_g$, then the following inequalities hold for any $k \geq 0$:*

$$\text{OPT}^{k+1} \leq \text{OPT}^k - \frac{1 - \alpha L_f}{2\alpha} \mathbb{E}[\|x^{k+1} - x^k\|^2] + \alpha [2L^2 \Delta_y^{k+1} + 2b_0^2 + \sigma_f^2] \quad (28a)$$

$$\Delta_y^{k+1} \leq (1 - \mu_g \beta / 2) \Delta_y^k + \left(\frac{2}{\mu_g \beta} - 1 \right) L_y^2 \cdot \mathbb{E}[\|x^k - x^{k-1}\|^2] + \beta^2 \sigma_g^2. \quad (28b)$$

The proof of (28a) is due to the smoothness of outer function $\ell(\cdot)$ established in Lemma 1, while (28b) follows from the strong convexity of the inner problem. See the details in §B.2. Note that (28a), (28b) together is a special case of (26) with:

$$\begin{aligned} \Omega^k &= \text{OPT}^k, \quad \Theta^k = \mathbb{E}[\|x^k - x^{k-1}\|^2], \quad c_0 = \frac{1}{2\alpha} - \frac{L_f}{2}, \quad c_1 = 2\alpha L^2, \quad c_2 = \alpha(2b_0^2 + \sigma_f^2), \\ \Upsilon^k &= \Delta_y^k, \quad d_0 = \mu_g \beta / 2, \quad d_1 = \left(\frac{2}{\mu_g \beta} - 1 \right) L_y^2, \quad d_2 = \beta^2 \sigma_g^2. \end{aligned}$$

Notice that $\Theta^0 = 0$. Assuming that $\alpha \leq 1/2L_f$, we notice the following implications:

$$\frac{\alpha}{\beta} \leq \frac{\mu_g}{8L_y L} \implies c_0 - c_1 \frac{d_1}{d_0} \geq \frac{1}{8\alpha} > 0, \quad d_0 - d_1 \frac{c_1}{c_0} \geq \frac{\mu_g \beta}{4} > 0, \quad (29)$$

i.e., if (29) holds, then the conclusion (27) can be applied. It can be shown that the step sizes in

(18) satisfy (29). Applying Lemma 4 shows that

$$\begin{aligned} \frac{1}{K} \sum_{k=1}^K \mathbb{E}[\|x^k - x^{k-1}\|^2] &\leq \frac{2\text{OPT}^0 + \frac{L}{L_y}(\Delta_y^0 + 4\sigma_g^2/\mu_g^2)}{L_y L \cdot K^{8/5}} + \frac{(2b_0^2 + \sigma_f^2) + 8\frac{\sigma_g^2 L^2}{\mu_g^2}}{2L_y^2 L^2 \cdot K^{6/5}}, \\ \frac{1}{K} \sum_{k=1}^K \Delta_y^k &\leq \frac{2\Delta_y^0 + \frac{8\sigma_g^2}{\mu_g^2} + \frac{4L_y}{\mu_g L} \text{OPT}^0}{K^{3/5}} + \frac{8\frac{\sigma_g^2}{\mu_g^2} + \frac{\mu_g(2b_0^2 + \sigma_f^2)}{2L^2}}{K^{2/5}}. \end{aligned}$$

Again, we emphasize that the two timescales step size design is crucial to establishing the above upper bounds. Now, recalling the properties of the Moreau envelop in (13), we obtain the following descent estimate:

Lemma 6. *Under Assumption 1, 2, 3. Set $\rho > -\mu_\ell$. For any $k \geq 0$,*

$$\begin{aligned} \mathbb{E}[\Phi_{1/\rho}(x^{k+1}) - \Phi_{1/\rho}(x^k)] &\leq \frac{5\rho}{2} \mathbb{E}[\|x^{k+1} - x^k\|^2] + \left[\frac{2\alpha\rho L^2}{\rho + \mu_\ell} + 3\alpha^2 \rho L^2 \right] \Delta_y^{k+1} \\ &\quad - \frac{(\rho + \mu_\ell)\rho\alpha}{4} \mathbb{E}[\|\hat{x}^k - x^k\|^2] + \left[\frac{2\rho}{\rho + \mu_\ell} + \rho(\tilde{\sigma}_f^2 + 3\alpha) \right] \alpha^2. \end{aligned} \quad (30)$$

See details in §B.3. Summing up the inequality (30) from $k = 0$ to $k = K - 1$ gives the following upper bound:

$$\begin{aligned} \frac{1}{K} \sum_{k=0}^{K-1} \mathbb{E}[\|\hat{x}(x^k) - x^k\|^2] &\leq \frac{4}{(\rho + \mu_\ell)\rho} \left[\frac{\Phi_{1/\rho}(x^0)}{\alpha K} + \frac{5\rho}{2\alpha K} \sum_{k=1}^K \mathbb{E}[\|x^k - x^{k-1}\|^2] \right] \\ &\quad + \frac{4}{\rho + \mu_\ell} \left[\frac{\frac{L^2}{\rho + \mu_\ell} + 3\alpha L^2}{K} \sum_{k=1}^K \Delta_y^k + \left[\frac{2}{\rho + \mu_\ell} + \tilde{\sigma}_f^2 + 3\alpha \right] \alpha \right]. \end{aligned}$$

Combining the above and $\alpha \asymp K^{-3/5}$ yields the desired rate of $\frac{1}{K} \sum_{k=0}^{K-1} \mathbb{E}[\|\hat{x}(x^k) - x^k\|^2] = \mathcal{O}(K^{-2/5})$. In particular, the asymptotic bound is given by setting $\rho = 2|\mu_\ell|$.

Proof of Corollary 1 We observe that Lemma 5 can be applied directly in this setting since convex functions are also weakly convex. With the step size choice (21), similar conclusions hold as:

$$\begin{aligned} \frac{1}{K} \sum_{k=1}^K \mathbb{E}[\|x^k - x^{k-1}\|^2] &\leq \frac{2\text{OPT}^0 + \frac{L}{L_y}(\Delta_y^0 + 4\sigma_g^2/\mu_g^2)}{L_y L \cdot K^{7/4}} + \frac{(2b_0^2 + \sigma_f^2) + 8\frac{\sigma_g^2 L^2}{\mu_g^2}}{2L_y^2 L^2 \cdot K^{6/4}}, \\ \frac{1}{K} \sum_{k=1}^K \Delta_y^k &\leq \frac{2\Delta_y^0 + \frac{8\sigma_g^2}{\mu_g^2} + \frac{4L_y}{\mu_g L} \text{OPT}^0}{K^{1/2}} + \frac{8\frac{\sigma_g^2}{\mu_g^2} + \frac{\mu_g(2b_0^2 + \sigma_f^2)}{2L^2}}{K^{1/2}}. \end{aligned}$$

With the additional property $\mu_\ell \geq 0$, in §E.1 we further derive an alternative descent estimate to (28a) that leads to the desired bound of $K^{-1} \sum_{k=1}^K \text{OPT}^k$.

4 Application to Reinforcement Learning

Consider a Markov decision process (MDP) (S, A, γ, P, r) , where S and A are the state and action spaces, respectively, $\gamma \in [0, 1)$ is the discount factor, $P(s'|s, a)$ is the transition kernel to the next state s' given the current state s and action a , and $r(s, a) \in [0, 1]$ is the reward at (s, a) . Furthermore, the initial state s_0 is drawn from a fixed distribution ρ_0 . We follow a stationary policy $\pi : S \times A \rightarrow \mathbb{R}$. For any $(s, a) \in S \times A$, $\pi(a|s)$ is the probability of the agent choosing action $a \in A$ at state $s \in S$. Note that a policy $\pi \in X$ induces a Markov chain on S . Denote the induced Markov transition kernel as P^π such that $s_{t+1} \sim P^\pi(\cdot|s_t)$. For any $s, s' \in S$, we have $P^\pi(s'|s) = \sum_{a \in A} \pi(a|s)P(s'|s, a)$. For any $\pi \in X$, P^π is assumed to induce a stationary distribution over S , denoted by μ^π . We assume that $|A| < \infty$ while $|S|$ is possibly infinite (but countable). To simplify our notations, for any distribution ρ on S , we let $\langle \cdot, \cdot \rangle_\rho$ be the inner product with respect to ρ , and $\|\cdot\|_{\mu^\pi \otimes \pi}$ be the weighted ℓ_2 -norm with respect to the probability measure $\mu^\pi \otimes \pi$ over $S \times A$ (where f, g are measurable functions on $S \times A$)

$$\langle f, g \rangle_\rho = \sum_{s \in S} \langle f(s, \cdot), g(s, \cdot) \rangle \rho(s), \quad \|f\|_{\mu^\pi \otimes \pi} = \sqrt{\sum_{s \in S} \left\{ \sum_{a \in A} \pi(a|s) \cdot [f(s, a)]^2 \right\} \mu^\pi(s)}.$$

In policy optimization, our objective is to maximize the expected total discounted reward received by the agent with respect to the policy π , i.e.,

$$\max_{\pi \in X \subseteq \mathbb{R}^{|S| \times |A|}} -\ell(\pi) = \mathbb{E}_\pi \left[\sum_{t \geq 0} \gamma^t \cdot r(s_t, a_t) \mid s_0 \sim \rho_0 \right], \quad (31)$$

where \mathbb{E}_π is the expectation with the actions taken according to policy π . To see that (31) is approximated as a bilevel problem, set P^π as the Markov operator under the policy π . We let Q^π be the unique solution to the following Bellman equation [48]:

$$Q(s, a) = r(s, a) + \gamma(P^\pi Q)(s, a), \quad \forall s, a \in S \times A. \quad (32)$$

Notice that the following holds:

$$Q^\pi(s, a) = \mathbb{E}_\pi \left[\sum_{t \geq 0} \gamma^t r(s_t, a_t) \mid s_0 = s, a_0 = a \right], \quad \mathbb{E}_{a \sim \pi(\cdot|s)} [Q^\pi(s, a)] = \langle Q^\pi(s, \cdot), \pi(\cdot|s) \rangle.$$

Further, we parameterize Q using a linear approximation $Q(s, a) \approx Q_\theta(s, a) := \phi^\top(s, a)\theta$, where $\phi : S \times A \rightarrow \mathbb{R}^d$ is a known feature mapping and $\theta \in \mathbb{R}^d$ is a finite-dimensional parameter. Using the fact that $\ell(\pi) = -\mathbb{E}_\pi [Q^\pi(s, a)]$, problem (31) can be approximated as a bilevel optimization problem such that:

$$\begin{aligned} \min_{\pi \in X \subseteq \mathbb{R}^{|S| \times |A|}} \quad & \ell(\pi) = -\langle Q_{\theta^*(\pi)}, \pi \rangle_{\rho_0} \\ \text{subject to} \quad & \theta^*(\pi) \in \arg \min_{\theta \in \mathbb{R}^d} \frac{1}{2} \|Q_\theta - r - \gamma P^\pi Q_\theta\|_{\mu^\pi \otimes \pi}^2. \end{aligned} \quad (33)$$

Solving Policy Optimization Problem We illustrate how to adopt the TTSA algorithm to solve (33). First, the inner problem is the policy evaluation (a.k.a. ‘critic’) which minimizes the mean squared Bellman error (MSBE). A standard approach is TD learning [47]. We draw two consecutive state-action pairs (s, a, s', a') satisfying $s \sim \mu^{\pi^k}$, $a \sim \pi^k(\cdot|s)$, $s' \sim P(\cdot|s, a)$, and $a' \sim \pi^k(\cdot|s')$, and

update the critic via

$$\theta^{k+1} = \theta^k - \beta h_g^k \quad \text{with} \quad h_g^k = [\phi^\top(s, a)\theta^k - r(s, a) - \gamma\phi^\top(s', a')\theta^k]\phi(s, a), \quad (34)$$

where β is the step size. This step resembles (6a) of TTSA except that the mean field $\mathbb{E}[h_g^k|\mathcal{F}_k]$ is a semigradient of the MSBE function.

Secondly, the outer problem searches for the policy (a.k.a. ‘actor’) that maximizes the expected discounted reward. To develop this step, let us define the following visitation measure and the Bregman divergence as:

$$\rho^{\pi^k}(s) := (1 - \gamma)^{-1} \sum_{t \geq 0} \gamma^t \mathbb{P}(s_t = s), \quad \bar{D}_{\psi, \rho^{\pi^k}}(\pi, \pi^k) := \sum_{s \in S} D_\psi(\pi(\cdot|s), \pi^k(\cdot|s)) \rho^{\pi^k}(s),$$

such that $\{s_t\}_{t \geq 0}$ is a trajectory of states obtained by drawing $s_0 \sim \rho_0$ and following the policy π^k , and D_ψ is the Kullback-Leibler (KL) divergence between probability distributions over A . We also define the following gradient surrogate:

$$[\bar{\nabla}_\pi f(\pi^k, \theta^{k+1})](s, a) = -(1 - \gamma)^{-1} Q_{\theta^{k+1}}(s, a) \rho^{\pi^k}(s), \quad \forall (s, a).$$

Similar to (5) and under the additional assumption that the linear approximation is exact, i.e., $Q_{\theta^*(\pi^k)} = Q^{\pi^k}$, we can show $\bar{\nabla}_\pi f(\pi^k, \theta^*(\pi^k)) = \nabla \ell(\pi^k)$ using the policy gradient theorem [48]. In a similar vein as (6b) in TTSA, we consider the mirror descent step for improving the policy (cf. proximal policy optimization in [44]):

$$\pi^{k+1} = \arg \min_{\pi \in X} \left\{ -(1 - \gamma)^{-1} \langle Q_{\theta^{k+1}}, \pi - \pi^k \rangle_{\rho^{\pi^k}} + \frac{1}{\alpha} \bar{D}_{\psi, \rho^{\pi^k}}(\pi, \pi^k) \right\}, \quad (35)$$

where α is the step size. Note that the above update can be performed as:

$$\pi^{k+1}(\cdot|s) \propto \pi^k(\cdot|s) \exp \left[\alpha_k (1 - \gamma)^{-1} Q_{\theta^{k+1}}(s, \cdot) \right] = \pi^0(\cdot|s) \exp \left[(1 - \gamma)^{-1} \phi(s, \cdot)^\top \sum_{i=0}^k \alpha \theta^{i+1} \right].$$

In other words, π^{k+1} can be represented using the running sum of critic $\sum_{i=0}^k \alpha \theta^{i+1}$. This is similar to the natural policy gradient method [27], and the algorithm requires a low memory footprint. Finally, the recursions (34), (35) give the two-timescale natural actor critic (TT-NAC) algorithm.

4.1 Convergence Analysis of TT-NAC

Consider the following assumptions on the MDP model of interest.

Assumption 4. *The reward function is uniformly bounded by a constant \bar{r} . That is, $|r(s, a)| \leq \bar{r}$ for all $(s, a) \in S \times A$.*

Assumption 5. *The feature map $\phi: S \times A \rightarrow \mathbb{R}^d$ satisfies $\|\phi(s, a)\|_2 \leq 1$ for all $(s, a) \in S \times A$. The action-value function associated with each policy is a linear function of ϕ . That is, for any policy $\pi \in X$, there exists $\theta^*(\pi) \in \mathbb{R}^d$ such that $Q^\pi(\cdot, \cdot) = \phi(\cdot, \cdot)^\top \theta^*(\pi) = Q_{\theta^*(\pi)}(\cdot, \cdot)$.*

Assumption 6. *For each policy $\pi \in X$, the induced Markov chain P^π admits a unique stationary distribution μ^π for all $\pi \in X$. Let there exists $\mu_\phi > 0$ such that $\mathbb{E}_{s \sim \mu^\pi, a \sim \pi(\cdot|s)}[\phi(s, a)\phi^\top(s, a)] \succeq \mu_\phi^2 \cdot I_d$ for all $\pi \in X$.*

Assumption 7. For any $(s, a) \in S \times A$ and any $\pi \in X$, let $\varrho(s, a, \pi)$ be a probability measure over S , defined by

$$[\varrho(s, a, \pi)](s') = (1 - \gamma)^{-1} \sum_{t \geq 0} \gamma^t \cdot \mathbb{P}(s_t = s'), \quad \forall s' \in S. \quad (36)$$

That is, $\varrho(s, a, \pi)$ is the visitation measure induced by the Markov chain starting from $(s_0, a_0) = (s, a)$ and follows π afterwards. For any π^* , there exists $C_\rho > 0$ such that

$$\mathbb{E}_{s' \sim \rho^{\pi^*}} \left[\left| \frac{\varrho(s, a, \pi)}{\rho^{\pi^*}}(s') \right|^2 \right] \leq C_\rho^2, \quad \forall (s, a) \in S \times A, \pi \in X.$$

In Assumption 5, we assume that each Q^π is linear which implies that the linear function approximation is exact. The assumption that the stationary distribution μ^π exists for any policy π is a common property for the MDP analyzed in TD learning, e.g., [4, 15]. Assumption 6 further assumes that the smallest eigenvalue of Σ_π is bounded uniformly away from zero. Finally, Assumption 7 postulates that ρ^* is regular such that the density ratio between $\varrho(s, a, \pi)$ and ρ^* has uniformly bounded second-order moments under ρ^* . Such an assumption is closely related to the concentratability coefficient [1, 2, 39], which characterizes the distribution shift incurred by policy updates and is conjectured essential for the sample complexity analysis of reinforcement learning methods [11].

To state our main convergence results, let us define the quantities of interest:

$$\Delta_Q^{k+1} := \mathbb{E}[\|\theta^{k+1} - \theta^*(\pi^k)\|_2^2], \quad \text{OPT}^k := \mathbb{E}[\ell(\pi^k) - \ell(\pi^*)], \quad (37)$$

where the expectations above are taken with respect to the i.i.d. draws of state-action pairs in (34) for TT-NAC. We remark that Δ_Q^k , analogous to Δ_y^k used in TTSA, is the tracking error that characterizes the performance of TD learning when the target value function, Q^{π^k} , is time-varying due to policy updates. We obtain:

Theorem 3. Consider the TT-NAC algorithm (34)-(35) for the policy optimization problem (33). Let $K \geq 32^2$ be the maximum number of iterations. Under Assumption 4 - 7, and we set the step sizes as

$$\alpha = \frac{(1 - \gamma)^3 \mu_\phi}{\sqrt{\bar{r}} \cdot C_\rho^2} \min \left\{ \frac{(1 - \gamma)^2}{128 \mu_\phi^{-2}}, K^{-3/4} \right\}, \quad \beta = \min \left\{ \frac{(1 - \gamma) \mu_\phi^2}{8}, \frac{16}{(1 - \gamma) \mu_\phi^2} K^{-1/2} \right\}. \quad (38)$$

Then the following holds

$$\mathbb{E}[\text{OPT}^K] = \mathcal{O}(K^{-1/4}), \quad \mathbb{E}[\Delta_Q^{K+1}] = \mathcal{O}(K^{-1/2}), \quad (39)$$

where K is an independent random variable uniformly distributed over $\{0, \dots, K - 1\}$.

To shed lights on our analysis, we first observe the following performance difference lemma proven in [26, Lemma 6.1]:

$$\ell(\pi) - \ell(\pi^*) = (1 - \gamma)^{-1} \langle Q^\pi, \pi^* - \pi \rangle_{\rho^{\pi^*}}, \quad \forall \pi \in X, \quad (40)$$

where π^* is an optimal policy solving (33). The above implies a restricted form of convexity, and our analysis uses the insight that (40) plays a similar role as (2) [with $\mu_\ell \geq 0$] and characterizes the loss geometry of the outer problem.

Our result shows that the TT-NAC algorithm finds an optimal policy at the rate of $\mathcal{O}(K^{-1/4})$ in terms of the objective value. This rate is comparable to another variant of the TT-NAC algorithm

in [54], which provided a customized analysis for TT-NAC. In contrast, the analysis for our TT-NAC algorithm is rooted in the general TTSA framework developed in §3.3 for tackling bilevel optimization problems. Notice that analysis for the two-timescale actor-critic algorithm can also be found in [53], which provides an $\mathcal{O}(K^{-2/5})$ convergence rate to a stationary solution.

5 Conclusion

This paper develops efficient two-timescale stochastic approximation algorithms for a class of bilevel optimization problems where the inner problem is unconstrained and strongly convex. We show the convergence rates of the proposed TTSA algorithm under the settings where the outer objective function is either strongly convex, convex, or non-convex. Additionally, we show how our theory and analysis can be customized to a two-timescale actor-critic proximal policy optimization algorithm in reinforcement learning, and obtain a comparable convergence rate to existing literature.

A Omitted Proofs of Theorem 1

To simplify notations, for any $n, m \in \mathbb{N}$, we define the following quantities for brevity of notations.

$$G_{m:n}^{(1)} = \prod_{i=m}^n (1 - \beta_i \mu_g / 4), \quad G_{m:n}^{(2)} = \prod_{i=m}^n (1 - \alpha_i \mu_\ell). \quad (41)$$

A.1 Proof of Lemma 2

Following a direct expansion of the updating rule for y^{k+1} and taking the conditional expectation given filtration \mathcal{F}_k yield that

$$\mathbb{E}[\|y^{k+1} - y^*(x^k)\|^2 | \mathcal{F}_k] \leq (1 - 2\beta_k \mu_g) \|y^k - y^*(x^k)\|^2 + \beta_k^2 \mathbb{E}[\|h_g^k\|^2 | \mathcal{F}_k],$$

where we used the unbiasedness of h_g^k [cf. Assumption 3] and the strong convexity of g . By direct computation and (7b) in Assumption 3, we have

$$\begin{aligned} \mathbb{E}[\|h_g^k\|^2 | \mathcal{F}_k] &= \mathbb{E}[\|h_g^k - \nabla g(x^k, y^k)\|^2 | \mathcal{F}_k] + \|\nabla g(x^k, y^k)\|^2 \\ &\leq \sigma_g^2 + (1 + \sigma_g^2) \|\nabla g(x^k, y^k)\|^2 \leq \sigma_g^2 + (1 + \sigma_g^2) \cdot L_g^2 \|y^k - y^*(x^k)\|^2, \end{aligned} \quad (42)$$

where the last inequality uses Assumption 2 and the optimality of the inner problem $\nabla_y g(x^k, y^*(x^k)) = 0$. As $\beta_k L_g^2 (1 + \sigma_g^2) \leq \mu_g$, we have

$$\mathbb{E}[\|y^{k+1} - y^*(x^k)\|^2 | \mathcal{F}_k] \leq (1 - \beta_k \mu_g) \cdot \|y^k - y^*(x^k)\|^2 + \beta_k^2 \sigma_g^2. \quad (43)$$

Using the basic inequality $2ab \leq 1/c \cdot a^2 + c \cdot b^2$ for all $c \geq 0$ and $a, b \in \mathbb{R}$, we have

$$\|y^k - y^*(x^k)\|^2 \leq (1 + 1/c) \cdot \|y^k - y^*(x^{k-1})\|^2 + (1 + c) \|y^*(x^{k-1}) - y^*(x^k)\|^2. \quad (44)$$

Note we have taken the convention that $x^{-1} = x^0$. Furthermore, we observe that

$$\|y^*(x^{k-1}) - y^*(x^k)\|^2 \leq L_y^2 \|x^k - x^{k-1}\|^2 \leq \alpha_{k-1}^2 L_y^2 \cdot \|h_f^{k-1}\|^2, \quad (45)$$

where the first inequality follows from Lemma 1, and the second inequality follows from the non-expansive property of projection. We have set $h_f^{-1} = 0$ as convention.

Through setting $c = \frac{2(1-\beta_k\mu_g)}{\beta_k\mu_g}$, we have $(1+1/c)(1-\beta_k\mu_g) = 1 - \frac{\mu_g}{2}\beta_k$. Substituting the above quantity c into (44) and combining with (43) show that

$$\begin{aligned} & \mathbb{E}[\|y^{k+1} - y^*(x^k)\|^2 | \mathcal{F}_k] \\ & \leq \left(1 - \frac{\beta_k\mu_g}{2}\right) \cdot \|y^k - y^*(x^{k-1})\|^2 + \beta_k^2 \cdot \sigma_g^2 + \frac{2 - \mu_g\beta_k}{\mu_g\beta_k} \cdot \alpha_{k-1}^2 L_y^2 \cdot \|h_f^{k-1}\|^2. \end{aligned} \quad (46)$$

Taking the total expectation and using (10), we have

$$\Delta_y^{k+1} \leq (1 - \beta_k\mu_g/2) \cdot \Delta_y^k + \beta_k^2 \sigma_g^2 + \frac{2 - \mu_g\beta_k}{\mu_g\beta_k} \alpha_{k-1}^2 L_y^2 [\tilde{\sigma}_f^2 + 3b_{k-1}^2 + 3L^2 \Delta_y^k],$$

with the convention $\alpha_{-1} = 0$. Using $\alpha_{k-1} \leq 2\alpha_k$, $\alpha_k \leq c_0\beta_k^{3/2}$, we have

$$\begin{aligned} \Delta_y^{k+1} & \leq \left[1 - \beta_k\mu_g/2 + \frac{12c_0^2 L_y^2 L^2}{\mu_g} \beta_k^2\right] \cdot \Delta_y^k + \beta_k^2 \sigma_g^2 + \frac{4c_0^2 L_y^2}{\mu_g} \beta_k^2 \cdot [\tilde{\sigma}_f^2 + 3b_0^2] \\ & \leq \left[1 - \beta_k\mu_g/4\right] \cdot \Delta_y^k + \beta_k^2 \sigma_g^2 + \frac{4c_0^2 L_y^2}{\mu_g} \beta_k^2 \cdot [\tilde{\sigma}_f^2 + 3b_0^2], \end{aligned}$$

where the last inequality is due to (15a). Solving the recursion leads to

$$\Delta_y^{k+1} \leq G_{0:k}^{(1)} \Delta_y^0 + \sum_{j=0}^k \beta_j^2 G_{j+1:k}^{(1)} \left\{ \sigma_g^2 + \frac{4c_0^2 L_y^2}{\mu_g} [\tilde{\sigma}_f^2 + 3b_0^2] \right\}. \quad (47)$$

Since $\beta_{k-1}/\beta_k \leq 1 + \beta_k \cdot (\mu_g/8)$, applying Lemma 9 to $\{\beta_k\}_{k \geq 0}$ with $a = \mu_g/4$ and $q = 2$, we have $\sum_{j=0}^k \beta_j^2 G_{j+1:k}^{(1)} \leq \frac{8\beta_k}{\mu_g}$. Finally, we can simplify (47) as

$$\Delta_y^{k+1} \leq G_{0:k}^{(1)} \Delta_y^0 + C_y^{(1)} \beta_k, \quad \text{where } C_y^{(1)} := \frac{8}{\mu_g} \left\{ \sigma_g^2 + \frac{4c_0^2 L_y^2}{\mu_g} [\tilde{\sigma}_f^2 + 3b_0^2] \right\}. \quad (48)$$

A.2 Proof of Lemma 3

Due to the projection property, we get

$$\|x^{k+1} - x^*\|^2 \leq \|x^k - \alpha_k h_f^k - x^*\|^2 = \|x^k - x^*\|^2 - 2\alpha_k \langle h_f^k, x^k - x^* \rangle + \alpha_k^2 \|h_f^k\|^2.$$

Taking the conditional expectation given \mathcal{F}'_k gives

$$\begin{aligned} \mathbb{E}[\|x^{k+1} - x^*\|^2 | \mathcal{F}'_k] & \leq \|x^k - x^*\|^2 - 2\alpha_k \langle \nabla \ell(x^k), x^k - x^* \rangle + \alpha_k^2 \mathbb{E}[\|h_f^k\|^2 | \mathcal{F}'_k] \\ & \quad - 2\alpha_k \langle \bar{\nabla}_x f(x^k, y^{k+1}) - \nabla \ell(x^k) + B_k, x^k - x^* \rangle, \end{aligned} \quad (49)$$

where the inequality follows from (7a). The strong convexity implies $\langle \nabla \ell(x^k), x^k - x^* \rangle \geq \mu_\ell \|x^k - x^*\|^2$, we further bound the r.h.s. of (49) via

$$\begin{aligned} & \mathbb{E}[\|x^{k+1} - x^*\|^2 | \mathcal{F}'_k] \\ & \leq (1 - 2\alpha_k \mu_\ell) \|x^k - x^*\|^2 - 2\alpha_k \langle \bar{\nabla}_x f(x^k, y^{k+1}) - \nabla \ell(x^k) + B_k, x^k - x^* \rangle + \alpha_k^2 \mathbb{E}[\|h_f^k\|^2 | \mathcal{F}'_k] \\ & \leq (1 - \alpha_k \mu_\ell) \|x^k - x^*\|^2 + \frac{\alpha_k}{\mu_\ell} \|\bar{\nabla}_x f(x^k, y^{k+1}) - \nabla \ell(x^k) + B_k\|^2 + \alpha_k^2 \mathbb{E}[\|h_f^k\|^2 | \mathcal{F}'_k] \\ & \leq (1 - \alpha_k \mu_\ell) \cdot \|x^k - x^*\|^2 + (2\alpha_k / \mu_\ell) \cdot \{L^2 \|y^{k+1} - y^*(x^k)\|^2 + b_k^2\} + \alpha_k^2 \cdot \mathbb{E}[\|h_f^k\|^2 | \mathcal{F}'_k], \end{aligned}$$

where the last inequality is from Lemma 1. Using (10) and taking total expectation:

$$\begin{aligned}\Delta_x^{k+1} &\leq [1 - \alpha_k \mu_\ell] \Delta_x^k + [2\alpha_k / \mu_\ell] L^2 \Delta_y^{k+1} + 2\alpha_k b_k^2 / \mu_\ell + \alpha_k^2 [\tilde{\sigma}_f^2 + 3b_0^2 + 3L^2 \Delta_y^{k+1}] \\ &\leq [1 - \alpha_k \mu_\ell] \cdot \Delta_x^k + [2\alpha_k / \mu_\ell + 3\alpha_k^2] \cdot L^2 \cdot \Delta_y^{k+1} + \alpha_k^2 [2\tilde{c}_b / \mu_\ell + \tilde{\sigma}_f^2 + 3b_0^2],\end{aligned}$$

where we have used $b_k^2 \leq \tilde{c}_b \alpha_k$. Solving the recursion above leads to

$$\begin{aligned}\Delta_x^{k+1} &\leq G_{0:k}^{(2)} \Delta_x^0 + \sum_{j=0}^k \left\{ \left[\frac{2\tilde{c}_b}{\mu_\ell} + \tilde{\sigma}_f^2 + 3b_0^2 \right] \alpha_j^2 G_{j+1:k}^{(2)} + \left[\frac{2L^2}{\mu_\ell} + 3\alpha_0 L^2 \right] \alpha_j G_{j+1:k}^{(2)} \Delta_y^{j+1} \right\} \\ &\leq G_{0:k}^{(2)} \Delta_x^0 + \frac{2}{\mu_\ell} \left[\frac{2\tilde{c}_b}{\mu_\ell} + \tilde{\sigma}_f^2 + 3b_0^2 \right] \cdot \alpha_k + \left(\frac{2L^2}{\mu_\ell} + 3\alpha_0 L^2 \right) \sum_{j=0}^k \alpha_j G_{j+1:k}^{(2)} \Delta_y^{j+1}.\end{aligned}$$

The last inequality follows from applying Lemma 9 with $q = 2$ and $a = \mu_\ell$.

A.3 Bounding Δ_x^k by coupling with Δ_y^k

Using (48), we observe that

$$\sum_{j=0}^k \alpha_j G_{j+1:k}^{(2)} \Delta_y^{j+1} \leq \sum_{j=0}^k \alpha_j G_{j+1:k}^{(2)} \left\{ G_{0:j}^{(1)} \Delta_y^0 + C_y^{(1)} \beta_j \right\}. \quad (50)$$

We bound each term on the right-hand side of (50). For the first term, as $\mu_\ell \alpha_i \leq \mu_g \beta_i / 8$ [cf. (15b)], applying Lemma 10 with $a = \mu_\ell$, $b = \mu_g / 4$, $\gamma_i = \alpha_i$, $\rho_i = \beta_i$ gives

$$\sum_{j=0}^k \alpha_j G_{j+1:k}^{(2)} G_{0:j}^{(1)} \Delta_y^0 \leq \frac{1}{\mu_\ell} G_{0:k}^{(2)} \Delta_y^0. \quad (51)$$

Recall that $\beta_j \leq c_1 \cdot \alpha_j^{2/3}$. Applying Lemma 9 with $q = 5/3$, $a = \mu_\ell$ yields

$$\sum_{j=0}^k \alpha_j \beta_j G_{j+1:k}^{(2)} \leq c_1 \sum_{j=0}^k \alpha_j^{5/3} G_{j+1:k}^{(2)} \leq c_1 \frac{2}{\mu_\ell} \alpha_k^{2/3}, \quad (52)$$

We obtain a bound on the optimality gap as

$$\Delta_x^{k+1} \leq G_{0:k}^{(2)} \left\{ \Delta_x^0 + \left[\frac{2L^2}{\mu_\ell^2} + \frac{3\alpha_0 L^2}{\mu_\ell} \right] \Delta_y^0 \right\} + \frac{2}{\mu_\ell} \left[\frac{2\tilde{c}_b}{\mu_\ell} + \tilde{\sigma}_f^2 + 3b_0^2 \right] \alpha_k + \frac{2c_1}{\mu_\ell} \left[\frac{2L^2}{\mu_\ell} + 3\alpha_0 L^2 \right] C_y^{(1)} \alpha_k^{2/3}.$$

To simplify the notation, we define the constants

$$C_x^{(0)} = \Delta_x^0 + \left[\frac{2L^2}{\mu_\ell^2} + \frac{3\alpha_0 L^2}{\mu_\ell} \right] \Delta_y^0, C_x^{(1)} = \frac{2}{\mu_\ell} \left[\frac{2\tilde{c}_b}{\mu_\ell} + \tilde{\sigma}_f^2 + 3b_0^2 \right] + \frac{2c_1}{\mu_\ell} \left[\frac{2L^2}{\mu_\ell} + 3\alpha_0 L^2 \right] C_y^{(1)}$$

Then, as long as $\alpha_k < 1/\mu_\ell$ and we use the step size parameters in (17), we have

$$\Delta_x^{k+1} \leq G_{0:k}^{(2)} C_x^{(0)} + C_x^{(1)} \alpha_k^{2/3} = \mathcal{O} \left(\left[\frac{L^2 \sigma_g^2}{\mu_\ell^2 \mu_g^2} + \frac{L^2 L_y^2 (\sigma_f^2 + \sup_{x \in X} \|\nabla \ell(x)\|^2)}{\mu_\ell^4} \right] \frac{1}{k^{2/3}} \right).$$

B Omitted Proofs of Theorem 2 and Corollary 1

B.1 Proof of Lemma 4

We observe that summing the first and the second inequalities in (26) from $k = 0$ to $k = K - 1$ gives:

$$c_0 \sum_{k=1}^K \Theta^k \leq \Omega^0 + c_1 \sum_{k=1}^K \Upsilon^k + c_2 \cdot K. \quad (53)$$

$$d_0 \sum_{k=1}^K \Upsilon^k \leq \Upsilon^1 + d_1 \sum_{k=1}^K \Theta^k + d_2 \cdot K. \quad (54)$$

Substituting (53) into (54) gives

$$d_0 \sum_{k=1}^K \Upsilon^k \leq \Upsilon^1 + d_2 \cdot K + \frac{d_1}{c_0} \left[\Omega^0 + c_1 \sum_{k=1}^K \Upsilon^k + c_2 \cdot K \right]. \quad (55)$$

Therefore, if $d_0 - d_1 \frac{c_1}{c_0} > 0$, a simple computation yields the second inequality in (27). Similarly, we substitute (54) into (53) to yield

$$c_0 \sum_{k=1}^K \Theta^k \leq \Omega^0 + c_2 \cdot K + \frac{c_1}{d_0} \left[\Upsilon^1 + d_1 \sum_{k=1}^K \Theta^k + d_2 \cdot K \right]. \quad (56)$$

Under $c_0 - c_1 \frac{d_1}{d_0} > 0$, simple computation yields the first inequality in (27).

B.2 Proof of Lemma 5

Recall that we defined $\text{OPT}^k := \mathbb{E}[\ell(x^k) - \ell(x^*)]$ for each $k \geq 0$. To begin with, we have the following descent estimate

$$\ell(x^{k+1}) \leq \ell(x^k) + \langle \nabla \ell(x^k), x^{k+1} - x^k \rangle + (L_f/2) \|x^{k+1} - x^k\|^2. \quad (57)$$

The optimality condition of step (6b) leads to the following bound

$$\begin{aligned} \langle \nabla \ell(x^k), x^{k+1} - x^k \rangle &\leq \langle \nabla \ell(x^k) - \bar{\nabla}_x f(x^k, y^{k+1}) - B_k, x^{k+1} - x^k \rangle \\ &\quad + \langle B_k + \bar{\nabla}_x f(x^k, y^{k+1}) - h_f^k, x^{k+1} - x^k \rangle - \frac{1}{\alpha} \|x^{k+1} - x^k\|^2, \end{aligned}$$

where we obtained the inequality by adding and subtracting $B_k + \bar{\nabla}_x f(x^k, y^{k+1}) - h_f^k$. Then, taking the conditional expectation on \mathcal{F}'_k , for any $c, d > 0$, we obtain

$$\begin{aligned} &\mathbb{E}[\langle \nabla \ell(x^k), x^{k+1} - x^k \rangle | \mathcal{F}'_k] \\ &\leq \mathbb{E}[\| \nabla \ell(x^k) - \bar{\nabla}_x f(x^k, y^{k+1}) - B_k \| \cdot \|x^{k+1} - x^k\| | \mathcal{F}'_k] \\ &\quad + \mathbb{E}[\| B_k + \bar{\nabla}_x f(x^k, y^{k+1}) - h_f^k \| \|x^{k+1} - x^k\| | \mathcal{F}'_k] - \frac{1}{\alpha} \mathbb{E}[\|x^{k+1} - x^k\|^2 | \mathcal{F}'_k] \\ &\leq \frac{1}{2c} \mathbb{E}[\| \nabla \ell(x^k) - \bar{\nabla}_x f(x^k, y^{k+1}) - B_k \|^2 | \mathcal{F}'_k] + \frac{c}{2} \mathbb{E}[\|x^{k+1} - x^k\|^2 | \mathcal{F}'_k] \\ &\quad + \frac{\sigma_f^2}{2d} + \frac{d}{2} \mathbb{E}[\|x^{k+1} - x^k\|^2 | \mathcal{F}'_k] - \frac{1}{\alpha} \mathbb{E}[\|x^{k+1} - x^k\|^2 | \mathcal{F}'_k], \end{aligned}$$

where the second inequality follows from the Young's inequality and Assumption 3. Simplifying the terms above leads to

$$\begin{aligned} & \mathbb{E}[\langle \nabla \ell(x^k), x^{k+1} - x^k \rangle | \mathcal{F}'_k] \\ & \leq \frac{1}{2c} \mathbb{E}[\|\nabla \ell(x^k) - \bar{\nabla}_x f(x^k, y^{k+1}) - B_k\|^2 | \mathcal{F}'_k] + \frac{\sigma_f^2}{2d} + \left(\frac{c+d}{2} - \frac{1}{\alpha}\right) \cdot \mathbb{E}[\|x^{k+1} - x^k\|^2 | \mathcal{F}'_k]. \end{aligned}$$

Setting $d = c = \frac{1}{2\alpha}$, plugging the above to (57), and taking the full expectation:

$$\text{OPT}^{k+1} \leq \text{OPT}^k - \left(\frac{1}{2\alpha} - \frac{L_f}{2}\right) \cdot \mathbb{E}[\|x^{k+1} - x^k\|^2] + \alpha \Delta^{k+1} + \alpha \sigma_f^2, \quad (58)$$

where we have denoted Δ^{k+1} as follows

$$\Delta^{k+1} := \mathbb{E}[\|\bar{\nabla}_x f(x^k; y^{k+1}) - \nabla \ell(x^k) - B_k\|^2] \stackrel{(8a)}{\leq} 2L^2 \mathbb{E}[\|y^{k+1} - y^*(x^k)\|^2] + 2b_k^2,$$

where the last inequality follows from Lemma 1 and (7a) in Assumption 3. Next, following from the standard SGD analysis [cf. (43)] and using $\beta \leq \mu_g / (L_g^2(1 + \sigma_g^2))$, we have

$$\begin{aligned} & \mathbb{E}[\|y^{k+1} - y^*(x^k)\|^2 | \mathcal{F}_k] \leq (1 - \mu_g \beta) \mathbb{E}[\|y^k - y^*(x^k)\|^2 | \mathcal{F}_k] + \beta^2 \sigma_g^2 \\ & \leq (1+c)(1 - \mu_g \beta) \mathbb{E}[\|y^k - y^*(x^{k-1})\|^2 | \mathcal{F}_k] + (1+1/c) \mathbb{E}[\|y^*(x^k) - y^*(x^{k-1})\|^2 | \mathcal{F}_k] + \beta^2 \sigma_g^2 \\ & \leq (1 - \mu_g \beta / 2) \mathbb{E}[\|y^k - y^*(x^{k-1})\|^2 | \mathcal{F}_k] + \left(\frac{2}{\mu_g \beta} - 1\right) \cdot \mathbb{E}[\|y^*(x^k) - y^*(x^{k-1})\|^2 | \mathcal{F}_k] + \beta^2 \sigma_g^2, \\ & \leq (1 - \mu_g \beta / 2) \mathbb{E}[\|y^k - y^*(x^{k-1})\|^2 | \mathcal{F}_k] + \left(\frac{2}{\mu_g \beta} - 1\right) L_y^2 \cdot \mathbb{E}[\|x^k - x^{k-1}\|^2 | \mathcal{F}_k] + \beta^2 \sigma_g^2, \end{aligned}$$

where the last inequality is due to the Lipschitz continuity property (8a) and $\mu_g \beta < 1$. Furthermore, we have picked $c = \mu_g \beta \cdot [2(1 - \mu_g \beta)]^{-1}$, so that

$$(1+c)(1 - \mu_g \beta) = 1 - \mu_g \beta / 2, \quad 1/c + 1 = 2/(\mu_g \beta) - 1.$$

Taking a full expectation on both sides leads to the desired result.

B.3 Proof of Lemma 6

For simplicity, we let \hat{x}^{k+1} and \hat{x} denote $\hat{x}(x^{k+1})$ and $\hat{x}(x)$, respectively. For any $x \in X$, letting $x_1 = \hat{x}$ and $x_2 = x$ in (2), we get

$$\ell(\hat{x}) \geq \ell(x) + \langle \nabla \ell(x), \hat{x} - x \rangle + \frac{\mu \ell}{2} \|\hat{x} - x\|^2. \quad (59)$$

Moreover, by the definition of \hat{x} , for any $x \in X$, we have

$$\ell(x) + \frac{\rho}{2} \|x - x\|^2 - \left[\ell(\hat{x}) + \frac{\rho}{2} \|\hat{x} - x\|^2 \right] = \ell(x) - \left[\ell(\hat{x}) + \frac{\rho}{2} \|\hat{x} - x\|^2 \right] \geq 0. \quad (60)$$

Adding the two inequalities above, we obtain

$$-\frac{\mu \ell + \rho}{2} \cdot \|\hat{x} - x\|^2 \geq \langle \nabla \ell(x), \hat{x} - x \rangle. \quad (61)$$

Note that we choose ρ such that $\rho + \mu_\ell > 0$. To proceed, combining the definitions of the Moreau envelop and \hat{x} in (13), for x^{k+1} , we have

$$\begin{aligned}
\Phi_{1/\rho}(x^{k+1}) &\stackrel{(13)}{=} \ell(\hat{x}^{k+1}) + \frac{\rho}{2} \cdot \|x^{k+1} - \hat{x}^{k+1}\|^2 \leq \ell(\hat{x}^k) + \frac{\rho}{2} \cdot \|x^{k+1} - \hat{x}^k\|^2 \\
&\leq \ell(\hat{x}^k) + \frac{\rho}{2} \cdot \|x^k - \hat{x}^k\|^2 + \frac{\rho}{2} \cdot \|x^{k+1} - x^k\|^2 + \rho\alpha \cdot \langle \hat{x}^k - x^k, h_f^k \rangle \\
&\quad + \alpha\rho \langle h_f^k, x^k - x^{k+1} \rangle + \rho\|x^{k+1} - x^k\|^2 \\
&\stackrel{(13)}{=} \Phi_{1/\rho}(x^k) + \frac{5\rho}{2} \cdot \|x^{k+1} - x^k\|^2 + \rho\alpha \langle \hat{x}^k - x^k, h_f^k \rangle + \alpha^2\rho\|h_f^k\|^2, \tag{62}
\end{aligned}$$

where the first equality and the first inequality follow from the optimality of $\hat{x}^{k+1} = \hat{x}(x^{k+1})$, and the second term is from the optimality condition in (6b). For any x^* that is a global optimal solution for the original problem $\min_{x \in X} \ell(x)$, we must have

$$\Phi_{1/\rho}(x^*) = \min_{x \in X} \left\{ \ell(x) + \frac{\rho}{2} \|x - x^*\|^2 \right\} = \ell(x^*),$$

where the last equality holds because

$$\Phi_{1/\rho}(x^*) = \min_{x \in X} \left\{ \ell(x) + \frac{\rho}{2} \|x - x^*\|^2 \right\} \leq \ell(x^*) + \frac{\rho}{2} \|x^* - x^*\|^2 = \ell(x^*), \tag{63}$$

$$\Phi_{1/\rho}(z) = \min_{x \in X} \left\{ \ell(x) + \frac{\rho}{2} \|x - z\|^2 \right\} \geq \min_{x \in X} \ell(x) = \ell(x^*), \quad \forall z \in X. \tag{64}$$

Taking expectation of $\langle \hat{x}^k - x^k, h_f^k \rangle$ while conditioning on \mathcal{F}'_k , we have:

$$\begin{aligned}
&\mathbb{E}[\langle \hat{x}^k - x^k, h_f^k \rangle | \mathcal{F}'_k] \\
&= \mathbb{E}[\langle \hat{x}^k - x^k, h_f^k - \bar{\nabla}_x f(x^k, y^{k+1}) + \bar{\nabla}_x f(x^k, y^{k+1}) - \nabla \ell(x^k) + \nabla \ell(x^k) \rangle | \mathcal{F}'_k] \\
&= \langle \hat{x}^k - x^k, B_k \rangle + \mathbb{E}[\langle \hat{x}^k - x^k, \bar{\nabla}_x f(x^k, y^{k+1}) - \nabla \ell(x^k) \rangle + \langle \hat{x}^k - x^k, \nabla \ell(x^k) \rangle | \mathcal{F}'_k], \tag{65}
\end{aligned}$$

where the second equality follows from (7a) in Assumption 3. By Young's inequality, for any $c > 0$, we have

$$\begin{aligned}
\langle \hat{x}^k - x^k, B_k \rangle &\leq \frac{c}{4} \|\hat{x}^k - x^k\|^2 + \frac{1}{c} b_k^2, \tag{66} \\
\mathbb{E}[\langle \hat{x}^k - x^k, \bar{\nabla}_x f(x^k, y^{k+1}) - \nabla \ell(x^k) \rangle | \mathcal{F}'_k] &\leq \frac{1}{c} \|\bar{\nabla}_x f(x^k, y^{k+1}) - \nabla \ell(x^k)\|^2 + \frac{c}{4} \|\hat{x}^k - x^k\|^2,
\end{aligned}$$

where we also use (7a) in deriving (66). Combining (61), (65), (66), and setting $c = (\rho + \mu_\ell)/2$, we obtain that

$$\begin{aligned}
&\mathbb{E}[\langle \hat{x}^k - x^k, h_f^k \rangle | \mathcal{F}'_k] \tag{67} \\
&\leq \frac{c}{2} \|\hat{x}^k - x^k\|^2 + \frac{1}{c} b_k^2 + \frac{1}{c} \mathbb{E}[\|\bar{\nabla}_x f(x^k, y^{k+1}) - \nabla \ell(x^k)\|^2] - \frac{\rho + \mu_\ell}{2} \|\hat{x}^k - x^k\|^2 \\
&= \frac{2}{\rho + \mu_\ell} \cdot \mathbb{E}[\|\bar{\nabla}_x f(x^k, y^{k+1}) - \nabla \ell(x^k)\|^2] - \frac{(\rho + \mu_\ell)}{4} \cdot \|\hat{x}^k - x^k\|^2 + \frac{2}{\rho + \mu_\ell} \cdot b_k^2 \\
&\leq \frac{2L^2}{\rho + \mu_\ell} \cdot \mathbb{E}[\|y^{k+1} - y^*(x^k)\|^2] - \frac{(\rho + \mu_\ell)}{4} \cdot \|\hat{x}^k - x^k\|^2 + \frac{2}{\rho + \mu_\ell} \cdot b_k^2,
\end{aligned}$$

where the last step follows from the first inequality of Lemma 1. Plugging the above into (62), and taking a full expectation, we obtain

$$\begin{aligned}
& \mathbb{E}[\Phi_{1/\rho}(x^{k+1})] - \mathbb{E}[\Phi_{1/\rho}(x^k)] \\
& \leq \frac{5\rho}{2} \mathbb{E}[\|x^{k+1} - x^k\|^2] + \frac{2\rho\alpha L^2}{\rho + \mu_\ell} \Delta_y^{k+1} - \frac{(\rho + \mu_\ell)\rho\alpha}{4} \mathbb{E}[\|\hat{x}^k - x^k\|^2] + \frac{2\rho\alpha b_k^2}{\rho + \mu_\ell} + \alpha^2 \rho \mathbb{E}[\|h_f^k\|^2] \\
& \leq \frac{5\rho}{2} \mathbb{E}[\|x^{k+1} - x^k\|^2] + \frac{2\rho\alpha L^2}{\rho + \mu_\ell} \Delta_y^{k+1} - \frac{(\rho + \mu_\ell)\rho\alpha}{4} \mathbb{E}[\|\hat{x}^k - x^k\|^2] + \frac{2\rho\alpha b_k^2}{\rho + \mu_\ell} \\
& \quad + \alpha^2 \rho (\tilde{\sigma}_f^2 + 3b_k^2 + 3L^2 \Delta_y^{k+1}) \\
& \leq \frac{5\rho}{2} \mathbb{E}[\|x^{k+1} - x^k\|^2] + \left[\frac{2\alpha\rho L^2}{\rho + \mu_\ell} + 3\alpha^2 \rho L^2 \right] \Delta_y^{k+1} - \frac{(\rho + \mu_\ell)\rho\alpha}{4} \mathbb{E}[\|\hat{x}^k - x^k\|^2] \\
& \quad + \left[\frac{2\rho}{\rho + \mu_\ell} + \rho(\tilde{\sigma}_f^2 + 3b_0^2) \right] \alpha^2,
\end{aligned}$$

where the last inequality is due to the assumption $b_k^2 \leq \alpha$.

C Proof of Theorem 3

Hereafter, we let $\langle \cdot, \cdot \rangle$ and $\|\cdot\|$ denote the inner product and ℓ_1 -norm on $\mathbb{R}^{|A|}$, respectively. For any two policies π_1 and π_2 , for any $s \in S$, $\|\pi_1(\cdot|s) - \pi_2(\cdot|s)\|_1$ is the total variation distance between $\pi_1(\cdot|s)$ and $\pi_2(\cdot|s)$. For any $f, f': S \times A \rightarrow \mathbb{R}$, define the following norms:

$$\|f\|_{\rho,1} = \left[\sum_{s \in S} \|f(s, \cdot)\|_1^2 \rho(s) \right]^{1/2}, \quad \|f\|_{\rho,\infty} = \left[\sum_{s \in S} \|f(s, \cdot)\|_\infty^2 \rho(s) \right]^{1/2}$$

The following result can be derived from the Hölder's inequality:

$$|\langle f, f' \rangle_\rho| \leq \sum_{s \in S} |\langle f(s, \cdot), f'(s, \cdot) \rangle| \rho(s) \leq \|f\|_{\rho,1} \|f'\|_{\rho,\infty}. \quad (68)$$

Lastly, it can be shown that $\|\pi\|_{\rho,1} = 1$, $\|\pi\|_{\rho,\infty} \leq 1$.

Under Assumption 5, $\theta^*(\pi)$ is the solution to the inner problem with $Q^\pi(\cdot, \cdot) = \phi(\cdot, \cdot)^\top \theta^*(\pi)$. Below we first show that $\theta^*(\pi)$ and Q^π are Lipschitz continuous maps with respect to $\|\cdot\|_{\rho^*,1}$, where ρ^* is the visitation measure of an optimal policy π^* .

Lemma 7. *Under Assumption 4–7, for any two policies $\pi_1, \pi_2 \in X$,*

$$\begin{aligned}
& \|Q^{\pi_1} - Q^{\pi_2}\|_{\rho^*,\infty} \leq (1 - \gamma)^{-2} \cdot \bar{r} \cdot C_\rho \cdot \|\pi_1 - \pi_2\|_{\rho^*,1}, \\
& \|\theta^*(\pi_1) - \theta^*(\pi_2)\|_2 \leq (1 - \gamma)^{-2} \cdot \bar{r} \cdot C_\rho / \mu_\phi \cdot \|\pi_1 - \pi_2\|_{\rho^*,1},
\end{aligned} \quad (69)$$

where \bar{r} is an upper bound on the reward function, μ_ϕ is specified in Assumption 6, and C_ρ is defined in Assumption 7.

The proof of the above lemma is relegated to §C.1.

In the sequel, we first derive coupled inequalities on the non-negative sequences $\text{OPT}^k := \mathbb{E}[\ell(\pi^k) - \ell(\pi^*)]$, $\Delta_Q^k := \mathbb{E}[\|\theta^k - \theta^*(\pi^{k-1})\|^2]$, $\mathbb{E}[\|\pi^k - \pi^{k+1}\|_{\rho^*,1}^2]$, then we apply Lemma 4 to derive the convergence rates of TT-NAC. Using the performance difference lemma [cf. (40)], we obtain the

following

$$\begin{aligned}\ell(\pi^{k+1}) - \ell(\pi^k) &= -(1-\gamma)^{-1} \langle Q^{\pi^{k+1}}, \pi^{k+1} - \pi^* \rangle_{\rho^*} + (1-\gamma)^{-1} \langle Q^{\pi^k}, \pi^k - \pi^* \rangle_{\rho^*} \\ &= (1-\gamma)^{-1} \langle -Q^{\pi^k}, \pi^{k+1} - \pi^k \rangle_{\rho^*} + (1-\gamma)^{-1} \langle Q^{\pi^k} - Q^{\pi^{k+1}}, \pi^{k+1} - \pi^* \rangle_{\rho^*}.\end{aligned}\quad (70)$$

Applying the inequality (68), we further have

$$\langle Q^{\pi^k} - Q^{\pi^{k+1}}, \pi^{k+1} - \pi^* \rangle_{\rho^*} \leq \|Q^{\pi^k} - Q^{\pi^{k+1}}\|_{\rho^*, \infty} \|\pi^{k+1} - \pi^*\|_{\rho^*, 1} \quad (71)$$

$$\leq 2L_Q \|\pi^k - \pi^{k+1}\|_{\rho^*, 1} \leq \frac{1-\gamma}{4\alpha} \|\pi^{k+1} - \pi^k\|_{\rho^*, 1}^2 + \frac{4L_Q^2 \alpha}{1-\gamma}, \quad (72)$$

where $L_Q := (1-\gamma)^{-2} \bar{r} \cdot C_\rho$. The above inequality follows from $\|\pi^k - \pi^{k+1}\|_{\rho^*, 1} \leq 2$ for any $\pi_1, \pi_2 \in X$ and applying Lemma 7. Then, combining (70), (72) leads to

$$\ell(\pi^{k+1}) - \ell(\pi^k) \leq \frac{-1}{1-\gamma} \langle Q^{\pi^k}, \pi^{k+1} - \pi^k \rangle_{\rho^*} + \frac{1}{4\alpha} \|\pi^{k+1} - \pi^k\|_{\rho^*, 1}^2 + 4 \frac{\alpha L_Q^2}{(1-\gamma)^2}. \quad (73)$$

Let us bound the first term in the right-hand side of (73). To proceed, note that the policy update (35) can be implemented for each state individually as below:

$$\pi^{k+1}(\cdot|s) = \arg \min_{\nu: \sum_a \nu(a)=1, \nu(a) \geq 0} \left\{ -(1-\gamma)^{-1} \cdot \langle Q_{\theta^{k+1}}(s, \cdot), \nu \rangle + 1/\alpha_k \cdot D_\psi(\nu, \pi^k(\cdot|s)) \right\}, \quad (74)$$

for all $s \in S$. Observe that we can modify ρ^{π^k} in (35) to ρ^* without changing the optimal solution for this subproblem. Specifically, (35) can be written as

$$\pi^{k+1} = \arg \min_{\pi \in X} \left\{ -(1-\gamma)^{-1} \langle Q_{\theta^{k+1}}, \pi - \pi^k \rangle_{\rho^*} + \frac{1}{\alpha} \bar{D}_{\psi, \rho^*}(\pi, \pi^k) \right\}. \quad (75)$$

We have

$$-(1-\gamma)^{-1} \langle Q^{\pi^k}, \pi^{k+1} - \pi^k \rangle_{\rho^*} = (1-\gamma)^{-1} [\langle Q_{\theta^{k+1}} - Q^{\pi^k}, \pi^{k+1} - \pi^k \rangle_{\rho^*} - \langle Q_{\theta^{k+1}}, \pi^{k+1} - \pi^k \rangle_{\rho^*}]$$

Furthermore, from (75), we obtain

$$\frac{\langle Q_{\theta^{k+1}}, \pi^{k+1} - \pi^k \rangle_{\rho^*}}{1-\gamma} \geq \frac{1}{\alpha} \sum_{s \in S} \langle \nabla D_\psi(\pi^{k+1}(\cdot|s), \pi^k(\cdot|s)), \pi^{k+1}(\cdot|s) - \pi^k(\cdot|s) \rangle_{\rho^*}(s), \quad (76)$$

where the inequality follows from the optimality condition of the mirror descent step. Meanwhile, the 1-strong convexity of $D_\psi(\cdot, \cdot)$ implies that

$$\langle \nabla D_\psi(\pi^{k+1}(\cdot|s), \pi^k(\cdot|s)), \pi^{k+1}(\cdot|s) - \pi^k(\cdot|s) \rangle \geq \|\pi^{k+1}(\cdot|s) - \pi^k(\cdot|s)\|^2. \quad (77)$$

Thus, combining (76) and (77), and applying Young's inequality, we further have

$$\begin{aligned}& -(1-\gamma)^{-1} \langle Q^{\pi^k}, \pi^{k+1} - \pi^k \rangle_{\rho^*} \\ & \leq \frac{1}{4\alpha} \|\pi^{k+1} - \pi^k\|_{\rho^*, 1}^2 + \alpha(1-\gamma)^{-2} \cdot \|Q^{\pi^k} - Q_{\theta^{k+1}}\|_{\rho^*, \infty}^2 - \frac{1}{\alpha} \|\pi^{k+1} - \pi^k\|_{\rho^*, 1}^2 \\ & = \alpha(1-\gamma)^{-2} \cdot \|Q^{\pi^k} - Q_{\theta^{k+1}}\|_{\rho^*, \infty}^2 - \frac{3}{4\alpha} \|\pi^{k+1} - \pi^k\|_{\rho^*, 1}^2.\end{aligned}\quad (78)$$

By direct computation and using $\|\phi(s, a)\| \leq 1$ [cf. Assumption 5], we have

$$\begin{aligned} \|Q^{\pi^k} - Q_{\theta^{k+1}}\|_{\rho^*, \infty}^2 &= \sum_{s \in \mathcal{S}} \left\{ \max_{a \in A} |\phi(s, a)^\top [\theta^*(\pi^k) - \theta^{k+1}]| \right\}^2 \rho^*(s) \\ &\leq \sum_{s \in \mathcal{S}} \max_{a \in A} \{ \|\phi(s, a)\|^2 \} \|\theta^*(\pi^k) - \theta^{k+1}\|^2 \rho^*(s) \leq \|\theta^*(\pi^k) - \theta^{k+1}\|^2. \end{aligned} \quad (79)$$

Combining (73), (78), and (79), we obtain

$$\ell(\pi^{k+1}) - \ell(\pi^k) \leq \alpha(1 - \gamma)^{-2} \|\theta^*(\pi^k) - \theta^{k+1}\|^2 - \frac{1}{2\alpha} \|\pi^{k+1} - \pi^k\|_{\rho^*, 1}^2 + 4(1 - \gamma)^{-2} L_Q^2 \alpha.$$

Taking full expectation leads to

$$\text{OPT}^{k+1} - \text{OPT}^k \leq \alpha(1 - \gamma)^{-2} \Delta_Q^{k+1} - \frac{1}{2\alpha} \mathbb{E}[\|\pi^{k+1} - \pi^k\|_{\rho^*, 1}^2] + 4(1 - \gamma)^{-2} L_Q^2 \alpha. \quad (80)$$

Next, we consider bounding the convergence of Δ_Q^k . Let $\mathcal{F}_k = \sigma\{\theta^0, \pi^0, \dots, \theta^k, \pi^k\}$ be the σ -algebra generated by the first $k + 1$ actor and critic updates. Under Assumption 5, we can write the conditional expectation of h_g^k as

$$\begin{aligned} \mathbb{E}[h_g^k | \mathcal{F}_k] &= \mathbb{E}_{\mu^{\pi^k}} [\{Q_{\theta^k}(s, a) - r(s, a) - \gamma Q_{\theta^k}(s', a')\} \phi(s, a) | \mathcal{F}_k] \\ &= \mathbb{E}_{\mu^{\pi^k}} [\phi(s, a) \{ \phi(s, a) - \gamma \phi(s', a') \}^\top] [\theta^k - \theta^*(\pi^k)], \end{aligned} \quad (81)$$

where $\mathbb{E}_{\mu^{\pi^k}}[\cdot]$ denotes the expectation taken with $s \sim \mu^{\pi^k}$, $a \sim \pi^k(\cdot | s)$, $s' \sim P(\cdot | s, a)$, $a' \sim \pi^k(\cdot | s')$. Under Assumption 5 and 6, Lemma 3 of [4] shows that $\mathbb{E}[h_g^k | \mathcal{F}_k]$ is a semigradient of the MSBE function $\|Q_{\theta^k} - Q_{\theta^*(\pi^k)}\|_{\mu^{\pi^k} \otimes \pi^k}^2$. Particularly, we obtain

$$\mathbb{E}[h_g^k | \mathcal{F}_k]^\top [\theta^k - \theta^*(\pi^k)] \geq (1 - \gamma) \|Q_{\theta^k} - Q_{\theta^*(\pi^k)}\|_{\mu^{\pi^k} \otimes \pi^k}^2 \geq \mu_{\text{td}} \|\theta^k - \theta^*(\pi^k)\|_2^2, \quad (82)$$

where we have let $\mu_{\text{td}} = (1 - \gamma)\mu_\phi^2$. Moreover, Lemma 5 of [4] demonstrates that the second order moment $\mathbb{E}[\|h_g^k\|_2^2 | \mathcal{F}_k]$ is bounded as

$$\mathbb{E}[\|h_g^k\|_2^2 | \mathcal{F}_k] \leq 8 \|Q_{\theta^k} - Q_{\theta^*(\pi^k)}\|_{\mu^{\pi^k} \otimes \pi^k}^2 + \sigma_{\text{td}}^2 \leq 8 \|\theta^k - \theta^*(\pi^k)\|_2^2 + \sigma_{\text{td}}^2, \quad (83)$$

where $\sigma_{\text{td}}^2 = 4\bar{r}^2(1 - \gamma)^{-2}$. Combining (82), (83) and recall $\beta \leq \mu_{\text{td}}/8$, it holds

$$\begin{aligned} \mathbb{E}[\|\theta^{k+1} - \theta^*(\pi^k)\|_2^2 | \mathcal{F}_k] &= \|\theta^k - \theta^*(\pi^k)\|_2^2 - 2\beta \mathbb{E}[h_g^k | \mathcal{F}_k]^\top [\theta^k - \theta^*(\pi^k)] + \beta^2 \mathbb{E}[\|h_g^k\|_2^2 | \mathcal{F}_k] \\ &\leq (1 - 2\mu_{\text{td}}\beta + 8\beta^2) \cdot \|\theta^k - \theta^*(\pi^k)\|_2^2 + \beta^2 \cdot \sigma_{\text{td}}^2 \\ &\leq (1 - \mu_{\text{td}}\beta) \cdot \|\theta^k - \theta^*(\pi^k)\|_2^2 + \beta^2 \cdot \sigma_{\text{td}}^2. \end{aligned} \quad (84)$$

By Young's inequality and Lemma 7, we further have

$$\begin{aligned} &\mathbb{E}[\|\theta^{k+1} - \theta^*(\pi^k)\|_2^2 | \mathcal{F}_k] \\ &\leq (1 + c)(1 - \mu_{\text{td}}\beta) \|\theta^k - \theta^*(\pi^{k-1})\|_2^2 + (1 + 1/c) \|\theta^*(\pi^k) - \theta^*(\pi^{k-1})\|_2^2 + \beta^2 \sigma_{\text{td}}^2 \\ &\leq (1 - \mu_{\text{td}}\beta/2) \|\theta^k - \theta^*(\pi^{k-1})\|_2^2 + \left(\frac{2}{\mu_{\text{td}}\beta} - 1 \right) \bar{L}_Q \|\pi^k - \pi^{k+1}\|_{\rho^*, 1}^2 + \beta^2 \sigma_{\text{td}}^2, \end{aligned} \quad (85)$$

where we have chosen $c > 0$ such that $(1 + c)(1 - \mu_{\text{td}}\beta) = 1 - \mu_{\text{td}}\beta/2$, which implies that $1/c + 1 = 2/(\mu_{\text{td}}\beta) - 1 > 0$ [cf. (38)]; The last inequality comes from Lemma 7 with the constant $\bar{L}_Q = (1 - \gamma)^{-4} \cdot \bar{r} \cdot C_\rho^2 \cdot \mu_\phi^{-2}$.

From (80), (85), we identify that condition (26) of Lemma 4 holds with:

$$\begin{aligned}\Omega^k &= \text{OPT}^k, \quad \Theta^k = \mathbb{E}[\|\pi^k - \pi^{k-1}\|_{\rho^*,1}^2], \quad c_0 = \frac{1}{2\alpha}, \quad c_1 = \alpha(1-\gamma)^{-2}, \quad c_2 = 4(1-\gamma)^{-2}L_Q^2 \cdot \alpha, \\ \Upsilon^k &= \mathbb{E}[\|\theta^k - \theta^*(\pi^{k-1})\|^2], \quad d_0 = \mu_{\text{td}}\beta/2, \quad d_1 = \left(\frac{2}{\mu_{\text{td}}\beta} - 1\right)\bar{L}_Q > 0, \quad d_2 = \beta^2 \cdot \sigma_{\text{td}}^2.\end{aligned}$$

Selecting the step sizes as in (38), one can verify that $\frac{\alpha}{\beta} < \frac{\mu_{\text{td}}(1-\gamma)}{16\sqrt{L_Q}}$. This ensures

$$c_0 - c_1 d_1 (d_0)^{-1} > 1/(4\alpha), \quad d_0 - c_1 d_1 (c_0)^{-1} > \mu_{\text{td}}\beta/4. \quad (86)$$

Applying Lemma 4, we obtain

$$\begin{aligned}\frac{1}{K} \sum_{k=1}^K \mathbb{E}[\|\pi^k - \pi^{k+1}\|_{\rho^*,1}^2] &\leq \frac{\text{OPT}^0 \cdot 4\alpha + \frac{8\alpha^2(1-\gamma)^{-2}}{\mu_{\text{td}}\beta}(\Delta_Q^0 + \beta^2\sigma_{\text{td}}^2)}{K} + \frac{8\alpha^2(4L_Q^2 + \beta\sigma_{\text{td}}^2/\mu_{\text{td}})}{(1-\gamma)^2} \\ \frac{1}{K} \sum_{k=1}^K \mathbb{E}[\Delta_Q^{k+1}] &\leq \frac{\mathbb{E}[\Delta_Q^0] + \beta^2\sigma_{\text{td}}^2 + \frac{4\alpha}{\mu_{\text{td}}\beta}\bar{L}_Q\text{OPT}^0}{\mu_{\text{td}}\beta K/4} + \frac{\beta^2\sigma_{\text{td}}^2 + \frac{16\alpha^2}{\mu_{\text{td}}\beta}(1-\gamma)^{-2}L_Q^2\alpha^2}{\mu_{\text{td}}\beta/4}.\end{aligned}$$

Particularly, plugging in $\alpha \asymp K^{-3/4}$, $\beta \asymp K^{-1/2}$ shows that the convergence rates are $K^{-1} \sum_{k=1}^K \mathbb{E}[\|\pi^k - \pi^{k+1}\|_{\rho^*,1}^2] = \mathcal{O}(K^{-3/2})$, $K^{-1} \sum_{k=1}^K \mathbb{E}[\Delta_Q^{k+1}] = \mathcal{O}(K^{-1/2})$.

Our last step is to analyze the convergence rate of the objective value OPT^k . To this end, we observe the following three-point inequality [3]

$$\frac{-\langle Q_{\theta^{k+1}}, \pi^{k+1} - \pi^* \rangle_{\rho^*}}{1-\gamma} \leq \frac{1}{\alpha} \left[\bar{D}_{\psi, \rho^*}(\pi^*, \pi^k) - \bar{D}_{\psi, \rho^*}(\pi^*, \pi^{k+1}) - \bar{D}_{\psi, \rho^*}(\pi^{k+1}, \pi^k) \right]. \quad (87)$$

Meanwhile, by the inequalities (68), (70), (71), we have

$$\begin{aligned}\ell(\pi^{k+1}) - \ell(\pi^k) &= -(1-\gamma)^{-1} \langle Q^{\pi^k}, \pi^{k+1} - \pi^k \rangle_{\rho^*} + (1-\gamma)^{-1} \langle Q^{\pi^k} - Q^{\pi^{k+1}}, \pi^{k+1} - \pi^* \rangle_{\rho^*} \\ &\leq (1-\gamma)^{-1} [\langle -Q^{\pi^k}, \pi^{k+1} - \pi^* \rangle_{\rho^*} - \langle Q^{\pi^k}, \pi^* - \pi^k \rangle_{\rho^*} + \|\pi^{k+1} - \pi^*\|_{\rho^*,1} \|Q^{\pi^k} - Q^{\pi^{k+1}}\|_{\rho^*,\infty}] \\ &\leq (1-\gamma)^{-1} [\langle -Q^{\pi^k}, \pi^{k+1} - \pi^* \rangle_{\rho^*} - \langle Q^{\pi^k}, \pi^* - \pi^k \rangle_{\rho^*} + 2L_Q \|\pi^{k+1} - \pi^k\|_{\rho^*,1}],\end{aligned}$$

where the last inequality follows from Lemma 7. Now, with the performance difference lemma $\ell(\pi^*) - \ell(\pi^k) = (1-\gamma)^{-1} \langle -Q^{\pi^k}, \pi^* - \pi^k \rangle_{\rho^*}$, the above simplifies to

$$\ell(\pi^{k+1}) - \ell(\pi^*) \leq (1-\gamma)^{-1} [-\langle Q^{\pi^k}, \pi^{k+1} - \pi^* \rangle_{\rho^*} + 2L_Q \|\pi^{k+1} - \pi^k\|_{\rho^*,1}]$$

With $\langle Q^{\pi^k}, \pi^{k+1} - \pi^* \rangle_{\rho^*} = \langle Q^{\pi^k} - Q_{\theta^{k+1}}, \pi^{k+1} - \pi^* \rangle_{\rho^*} + \langle Q_{\theta^{k+1}}, \pi^{k+1} - \pi^* \rangle_{\rho^*}$ and applying the three-point inequality (87), we have

$$\begin{aligned}\ell(\pi^{k+1}) - \ell(\pi^*) &\leq \frac{2}{1-\gamma} [L_Q \|\pi^{k+1} - \pi^k\|_{\rho^*,1} + \|Q^{\pi^k} - Q_{\theta^{k+1}}\|_{\rho^*,\infty}] \\ &\quad + \frac{1}{\alpha} [\bar{D}_{\psi, \rho^*}(\pi^*, \pi^k) - \bar{D}_{\psi, \rho^*}(\pi^*, \pi^{k+1}) - \bar{D}_{\psi, \rho^*}(\pi^{k+1}, \pi^k)]. \\ &\leq \frac{2}{1-\gamma} [L_Q \|\pi^{k+1} - \pi^k\|_{\rho^*,1} + \|\theta^*(\pi^k) - \theta^{k+1}\|_2] + \frac{1}{\alpha} [\bar{D}_{\psi, \rho^*}(\pi^*, \pi^k) - \bar{D}_{\psi, \rho^*}(\pi^*, \pi^{k+1})]\end{aligned}$$

where the last inequality used (79) and the fact that \bar{D}_{ψ, ρ^*} is non-negative. Finally, taking the full

expectation on both sides of the inequality, we obtain

$$\begin{aligned} \text{OPT}^{k+1} &\leq 2(1-\gamma)^{-1}\mathbb{E}[\|\theta^*(\pi^k) - \theta^{k+1}\|_2] + 2(1-\gamma)^{-1}L_Q\mathbb{E}[\|\pi^{k+1} - \pi^k\|_{\rho^*,1}] \\ &\quad + \frac{1}{\alpha}\mathbb{E}[\bar{D}_{\psi,\rho^*}(\pi^*, \pi^k) - \bar{D}_{\psi,\rho^*}(\pi^*, \pi^{k+1})]. \end{aligned} \quad (88)$$

Summing up both sides from $k = 0$ to $k = K - 1$ and dividing by K yields

$$\begin{aligned} \frac{1}{K}\sum_{k=1}^K \text{OPT}^k &\leq \frac{1}{\alpha K}\{\bar{D}_{\psi,\rho^*}(\pi^*, \pi^0) - \bar{D}_{\psi,\rho^*}(\pi^*, \pi^K)\} \\ &\quad + \frac{2}{(1-\gamma)K}\sum_{k=1}^K \{\mathbb{E}[\|\theta^*(\pi^{k-1}) - \theta^k\|_2] + L_Q \cdot \mathbb{E}[\|\pi^k - \pi^{k-1}\|_{\rho^*,1}]\}. \end{aligned} \quad (89)$$

Using Cauchy-Schwarz's inequality, it can be easily seen that the right-hand side is $\mathcal{O}(K^{-1/4})$. This concludes the proof of the corollary.

C.1 Proof of Lemma 7

We first bound $|Q^{\pi_1}(s, a) - Q^{\pi_2}(s, a)|$. By the Bellman equation (32) and the performance difference lemma (40), we have

$$\begin{aligned} Q^{\pi_1}(s, a) - Q^{\pi_2}(s, a) &= \sum_{s' \in S} P(s'|s, a) \cdot [V^{\pi_1}(s') - V^{\pi_2}(s')] \\ &= (1-\gamma)^{-1} \cdot \sum_{s' \in S} P(s'|s, a) \cdot \mathbb{E}_{\tilde{s} \sim \tilde{\varrho}(s', \pi_1)} [\langle Q^{\pi_2}(\tilde{s}, \cdot), \pi_1(\cdot|\tilde{s}) - \pi_2(\cdot|\tilde{s}) \rangle], \end{aligned} \quad (90)$$

where $\tilde{\varrho}(s', \pi_1)$ is the visitation measure obtained by the Markov chain induced by π_1 with the initial state fixed to s' . Recall the definition of the visitation measure $\varrho(s, a, \pi)$ in (36). We rewrite (90) as

$$Q^{\pi_1}(s, a) - Q^{\pi_2}(s, a) = (1-\gamma)^{-1} \cdot \mathbb{E}_{\tilde{s} \sim \varrho(s, a, \pi_1)} [\langle Q^{\pi_2}(\tilde{s}, \cdot), \pi_1(\cdot|\tilde{s}) - \pi_2(\cdot|\tilde{s}) \rangle]. \quad (91)$$

Moreover, notice that $\sup_{(s,a) \in S \times A} |Q^\pi(s, a)| \leq (1-\gamma)^{-1} \cdot \bar{r}$ under Assumption 4. Then, applying Hölder's inequality to (91), we obtain

$$\begin{aligned} |Q^{\pi_1}(s, a) - Q^{\pi_2}(s, a)| &\leq (1-\gamma)^{-1} \cdot \mathbb{E}_{\tilde{s} \sim \tilde{\varrho}(s', \pi_1)} [\|Q^{\pi_2}(\tilde{s}, \cdot)\|_\infty \cdot \|\pi_1(\cdot|\tilde{s}) - \pi_2(\cdot|\tilde{s})\|_1] \\ &\leq (1-\gamma)^{-2} \cdot \bar{r} \cdot \mathbb{E}_{\tilde{s} \sim \rho^*} \left[\frac{\varrho(s, a, \pi)}{\rho^*}(\tilde{s}) \cdot \|\pi_1(\cdot|\tilde{s}) - \pi_2(\cdot|\tilde{s})\|_1 \right] \\ &\leq (1-\gamma)^{-2} \cdot \bar{r} \cdot \left\{ \mathbb{E}_{\tilde{s} \sim \rho^*} \left[\left| \frac{\varrho(s, a, \pi)}{\rho^*}(\tilde{s}) \right|^2 \right] \mathbb{E}_{\tilde{s} \sim \rho^*} [\|\pi_1(\cdot|\tilde{s}) - \pi_2(\cdot|\tilde{s})\|_1^2] \right\}^{1/2} \\ &\leq (1-\gamma)^{-2} \cdot \bar{r} \cdot C_\rho \cdot \|\pi_1 - \pi_2\|_{\rho^*,1}, \end{aligned} \quad (92)$$

where the second inequality is from the boundedness of Q^π , the third one is the Cauchy-Schwarz inequality, and the last one is from Assumption 7. Finally, we have

$$\|Q^{\pi_1} - Q^{\pi_2}\|_{\rho^*,\infty} \leq (1-\gamma)^{-2} \cdot \bar{r} \cdot C_\rho \cdot \|\pi_1 - \pi_2\|_{\rho^*,1}.$$

It remains to bound $\|\theta^*(\pi_1) - \theta^*(\pi_2)\|^2$. Under Assumption 5, we have

$$\begin{aligned}\|Q^{\pi_1} - Q^{\pi_2}\|_{\mu^{\pi^*} \otimes \pi^*}^2 &= \mathbb{E}_{s \sim \mu^{\pi^*}, a \sim \pi^*(\cdot|s)} \{ [Q^{\pi_1}(s, a) - Q^{\pi_2}(s, a)]^2 \} \\ &= \mathbb{E}_{s \sim \mu^{\pi^*}, a \sim \pi^*(\cdot|s)} \left(\{ \phi(s, a)^\top [\theta^*(\pi_1) - \theta^*(\pi_2)] \}^2 \right) \\ &= [\theta^*(\pi_1) - \theta^*(\pi_2)]^\top \Sigma_{\pi^*} [\theta^*(\pi_1) - \theta^*(\pi_2)].\end{aligned}\tag{93}$$

Then, combining Assumption 6 and (92), we have

$$\mu_\phi^2 \|\theta^*(\pi_1) - \theta^*(\pi_2)\|^2 \leq \|Q^{\pi_1} - Q^{\pi_2}\|_{\mu^{\pi^*} \otimes \pi^*}^2 \leq (1 - \gamma)^{-4} \cdot \bar{r}^2 \cdot C_\rho^2 \cdot \|\pi_1 - \pi_2\|_{\rho^*, 1}^2,\tag{94}$$

which yields the second inequality in Lemma 7. We conclude the proof.

D Auxiliary Lemmas

The proofs for the lemmas below can be found in the online appendix [23].

Lemma 8. [28, Lemma 12] *Let $\{\gamma_j\}_{j \geq 0}$ be a non-increasing, non-negative sequence such that $\gamma_0 < 1/a$, it holds for any $k \geq 0$ that*

$$\sum_{j=0}^k \gamma_j \prod_{\ell=j+1}^k (1 - \gamma_\ell a) \leq \frac{1}{a}.\tag{95}$$

Lemma 9. *Fix a real number $1 < q \leq 2$. Let $\{\gamma_j\}_{j \geq 0}$ be a non-increasing, non-negative sequence such that $\gamma_0 < 1/(2a)$. Suppose that $\frac{\gamma_{\ell-1}}{\gamma_\ell} \leq 1 + \frac{a}{2(q-1)}\gamma_\ell$. Then, it holds for any $k \geq 0$ that*

$$\sum_{j=0}^k \gamma_j^q \prod_{\ell=j+1}^k (1 - \gamma_\ell a) \leq \frac{2}{a} \gamma_k^{q-1}.\tag{96}$$

Lemma 10. *Fix the real numbers $a, b > 0$. Let $\{\gamma_j\}_{j \geq 0}, \{\rho_j\}_{j \geq 0}$ be nonincreasing, non-negative sequences such that $2a\gamma_j \leq b\rho_j$ for all j . Then, it holds that*

$$\sum_{j=0}^k \gamma_j \prod_{\ell=j+1}^k (1 - \gamma_\ell a) \prod_{i=0}^j (1 - \rho_i b) \leq \frac{1}{a} \prod_{\ell=0}^k (1 - \gamma_\ell a), \quad \forall k \geq 0.\tag{97}$$

E Technical Results Omitted from the Main Paper

E.1 Proof of Corollary 1

Our proof departs from that of Theorem 2 through manipulating the descent estimate (57) in an alternative way. The key is to observe the following three-point inequality [3]:

$$\langle h_f^k, x^{k+1} - x^* \rangle \leq \frac{1}{2\alpha} \left\{ \|x^* - x^k\|^2 - \|x^* - x^{k+1}\|^2 - \|x^k - x^{k+1}\|^2 \right\},\tag{98}$$

where x^* is an optimal solution to (1). Observe that

$$\langle \nabla \ell(x^k), x^{k+1} - x^k \rangle = \langle \nabla \ell(x^k) - h_f^k, x^{k+1} - x^* \rangle + \langle h_f^k, x^{k+1} - x^* \rangle + \langle \nabla \ell(x^k), x^* - x^k \rangle.\tag{99}$$

Notice that due to the convexity of $\ell(x)$, we have $\langle \nabla \ell(x^k), x^* - x^k \rangle \leq -\text{OPT}^k$. Furthermore,

$$\begin{aligned} & \langle \nabla \ell(x^k) - h_f^k, x^{k+1} - x^* \rangle \\ &= \langle \nabla \ell(x^k) - h_f^k + B_k + \bar{\nabla}_x f(x^k, y^{k+1}) - B_k - \bar{\nabla}_x f(x^k, y^{k+1}), x^{k+1} - x^* \rangle \\ &\leq D_x \{b_k + L\|y^{k+1} - y^*(x^k)\|\} + \langle B_k + \bar{\nabla}_x f(x^k, y^{k+1}) - h_f^k, x^{k+1} - x^k + x^k - x^* \rangle. \end{aligned}$$

We notice that $\mathbb{E}[\langle B_k + \bar{\nabla}_x f(x^k, y^{k+1}) - h_f^k, x^k - x^* \rangle | \mathcal{F}'_k] = 0$. Thus, taking the total expectation on both sides and applying Young's inequality on the last inner product lead to

$$\mathbb{E}[\langle \nabla \ell(x^k) - h_f^k, x^{k+1} - x^* \rangle] \leq D_x \{b_k + L\mathbb{E}[\|y^{k+1} - y^*(x^k)\|]\} + \frac{\alpha}{2}\sigma_f^2 + \frac{1}{2\alpha}\mathbb{E}[\|x^{k+1} - x^k\|^2]$$

Substituting the above observations into (57) and using the three-point inequality (98) give

$$\begin{aligned} \mathbb{E}[\ell(x^{k+1}) - \ell(x^k)] &\leq D_x \{b_k + L\mathbb{E}[\|y^{k+1} - y^*(x^k)\|]\} + \frac{1}{2\alpha} \left\{ \|x^* - x^k\|^2 - \|x^* - x^{k+1}\|^2 \right\} \\ &\quad + \frac{\alpha}{2}\sigma_f^2 - \text{OPT}^k + \frac{L_f}{2}\mathbb{E}[\|x^{k+1} - x^k\|^2]. \end{aligned}$$

Summing up both sides from $k = 0$ to $k = K - 1$ and dividing by K gives

$$\begin{aligned} \frac{1}{K} \sum_{k=1}^K \text{OPT}^k &\leq D_x b_0 + \frac{\alpha\sigma_f^2}{2} + \frac{\|x^* - x^0\|^2}{2\alpha K} + \frac{D_x L}{K} \sum_{k=1}^K \mathbb{E}[\|y^k - y^*(x^{k-1})\|] \\ &\quad + \frac{L_f}{2K} \sum_{k=1}^K \mathbb{E}[\|x^k - x^{k-1}\|^2]. \end{aligned}$$

Applying the Cauchy-Schwartz inequality shows that $\frac{1}{K} \sum_{k=1}^K \mathbb{E}[\|y^k - y^*(x^{k-1})\|] = \sqrt{\frac{1}{K} \sum_{k=1}^K \Delta_y^k} = \mathcal{O}(K^{-1/4})$. The proof is concluded.

E.2 Proof of Lemma 9

To derive this result, we observe that

$$\begin{aligned} \sum_{j=0}^k \gamma_j^q \prod_{\ell=j+1}^k (1 - \gamma_\ell a) &\leq \gamma_k^{q-1} \sum_{j=0}^k \gamma_j \frac{\gamma_j^{q-1}}{\gamma_k^{q-1}} \prod_{\ell=j+1}^k (1 - \gamma_\ell a) \\ &= \gamma_k^{q-1} \sum_{j=0}^k \gamma_j \prod_{\ell=j+1}^k \left(\frac{\gamma_{\ell-1}}{\gamma_\ell} \right)^{q-1} (1 - \gamma_\ell a). \end{aligned}$$

Furthermore, from the conditions on γ_ℓ ,

$$\left(\frac{\gamma_{\ell-1}}{\gamma_\ell} \right)^{q-1} (1 - \gamma_\ell a) \leq \left(1 + \frac{a}{2(q-1)} \gamma_\ell \right)^{q-1} (1 - \gamma_\ell a) \leq 1 - \frac{a}{2} \gamma_\ell.$$

Therefore,

$$\sum_{j=0}^k \gamma_j^q \prod_{\ell=j+1}^k (1 - \gamma_\ell a) \leq \gamma_k^{q-1} \sum_{j=0}^k \gamma_j \prod_{\ell=j+1}^k \left(1 - \frac{a}{2} \gamma_\ell \right) \leq \frac{2}{a} \gamma_k^{q-1}.$$

This concludes the proof.

E.3 Proof of Lemma 10

First of all, the condition $2a\gamma_j \leq b\rho_j$ implies

$$\frac{1 - \rho_i b}{1 - \gamma_i a} \leq 1 - \rho_i b/2, \quad \forall i \geq 0.$$

As such, we observe $\prod_{i=0}^j \frac{1 - \rho_i b}{1 - \gamma_i a} \leq \prod_{i=0}^j (1 - \rho_i b/2)$ and subsequently,

$$\sum_{j=0}^k \gamma_j \prod_{\ell=j+1}^k (1 - \gamma_\ell a) \prod_{i=0}^j (1 - \rho_i b) \leq \left[\prod_{\ell=0}^k (1 - \gamma_\ell a) \right] \sum_{j=0}^k \gamma_j \prod_{i=0}^j (1 - \rho_i b/2).$$

Furthermore, for any $j = 0, \dots, k$, it holds

$$\rho_j \prod_{i=0}^{j-1} (1 - \rho_i b/2) = \frac{2}{b} \left[\prod_{i=0}^{j-1} (1 - \rho_i b/2) - \prod_{i=0}^j (1 - \rho_i b/2) \right], \quad (100)$$

where we have taken the convention $\prod_{i=0}^{-1} (1 - \rho_i b/2) = 1$. We obtain that

$$\begin{aligned} \sum_{j=0}^k \gamma_j \prod_{i=0}^j (1 - \rho_i b/2) &\leq \frac{b}{2a} \sum_{j=0}^k \rho_j \prod_{i=0}^{j-1} (1 - \rho_i b/2) \\ &= \frac{1}{a} \sum_{j=0}^k \left[\prod_{i=0}^{j-1} (1 - \rho_i b/2) - \prod_{i=0}^j (1 - \rho_i b/2) \right] \leq \frac{1}{a}, \end{aligned}$$

where the last inequality follows from the bound $1 - \prod_{i=0}^k (1 - \rho_i \mu_g b/2) \leq 1$. Combining with the above inequality yields the desired results.

E.4 Computing Biased Samples of $\bar{\nabla}_x f(x^k, y^{k+1})$

We illustrate how to construct estimates h_f^k which satisfy Assumption 3, where the approximate composite gradient is give by (5), and analyze the magnitude of the bias involved. Note that for general TTSA, as analyzed in Theorem 1, Theorem 2, it is necessary for the gradient estimate h_f^k to have a bias that *decays polynomially*. To this end, we recall that h_f^k aims to estimate

$$\bar{\nabla}_x f(x^k, y^{k+1}) = \nabla_x f(x^k, y^{k+1}) - \nabla_{xy}^2 g(x^k, y^{k+1}) [\nabla_{yy}^2 g(x^k, y^{k+1})]^{-1} \nabla_y f(x^k, y^{k+1}). \quad (101)$$

Without loss of generality, we shall drop the superscripts of iteration index from now on. We declare $x \equiv x^k, y \equiv y^{k+1}$. Each of the gradient/Jacobian/Hessian in the above can be written as (where $\xi^{(i)}$ is distributed with $\mu^{(i)}$, $i = 1, 2$)

$$\begin{aligned} \nabla_x f(x, y) &= \mathbb{E}_{\mu^{(1)}} [\nabla_x f(x, y; \xi^{(1)})], & \nabla_y f(x, y) &= \mathbb{E}_{\mu^{(1)}} [\nabla_y f(x, y; \xi^{(1)})], \\ \nabla_{xy}^2 g(x, y) &= \mathbb{E}_{\mu^{(2)}} [\nabla_{xy}^2 g(x, y; \xi^{(2)})], & \nabla_{yy}^2 g(x, y) &= \mathbb{E}_{\mu^{(2)}} [\nabla_{yy}^2 g(x, y; \xi^{(2)})]. \end{aligned}$$

Inspired by [22], let $t \geq 1$ be an integer, we consider the following procedure:

1. Select $p \in \{0, \dots, t-1\}$ uniformly at random.
2. Draw $2 + p$ samples as $\xi^{(1)} \sim \mu^{(1)}, \xi_0^{(2)}, \dots, \xi_p^{(2)} \sim \mu^{(2)}$ *independently*.

3. Construct h_f^k via

$$h_f^k = \nabla_x f(x, y; \xi^{(1)}) - \nabla_{xy}^2 g(x, y; \xi_0^{(2)}) \left[\frac{t}{L_g} \prod_{i=1}^p \left(I - \frac{1}{L_g} \nabla_{yy}^2 g(x, y; \xi_i^{(2)}) \right) \right] \nabla_y f(x, y; \xi^{(1)}), \quad (102)$$

where as a convention, we have set $\prod_{i=1}^p \left(I - \frac{1}{L_g} \nabla_{yy}^2 g(x, y; \xi_i^{(2)}) \right) = I$ if $p = 0$.

The following lemma shows that h_f^k satisfies Assumption 3:

Lemma 11. *Under Assumption 1, 2. Consider h_f^k constructed in (102). For any $t \geq 1$,*

$$\left\| \bar{\nabla}_x f(x, y) - \mathbb{E}[h_f^k] \right\| \leq C_{gxy} C_{fy} \frac{1}{\mu_g} \cdot \left(1 - \frac{\mu_g}{L_g} \right)^t. \quad (103)$$

Furthermore, assume that $\mathbb{E}[\|\nabla_y f(x, y; \xi^{(1)})\|^2] \leq C_y$, $\mathbb{E}[\|\nabla_{xy}^2 g(x, y; \xi^{(2)})\|^2] \leq C_g$,

$$\begin{aligned} \mathbb{E}[\|\nabla_x f(x, y) - \nabla_x f(x, y; \xi^{(1)})\|^2] &\leq \sigma_{fx}^2, & \mathbb{E}[\|\nabla_y f(x, y) - \nabla_y f(x, y; \xi^{(1)})\|^2] &\leq \sigma_{fy}^2, \\ \mathbb{E}[\|\nabla_{xy}^2 g(x, y) - \nabla_{xy}^2 g(x, y; \xi^{(2)})\|^2] &\leq \sigma_{gxy}^2, \end{aligned}$$

then we have

$$\mathbb{E}[\|h_f^k - \mathbb{E}[h_f^k]\|^2] \leq \sigma_{fx}^2 + \frac{3}{\mu_g^2} \left[(\sigma_{fy}^2 + C_y^2) \{ \sigma_{gxy}^2 + 2C_{gxy}^2 \} + \sigma_{fy}^2 C_{gxy}^2 \right]. \quad (104)$$

Proof. Since the samples are drawn independently, the expected value of h_f^k is

$$\mathbb{E}[h_f^k] = \nabla_x f(x, y) - \nabla_{xy}^2 g(x, y) \mathbb{E} \left[\frac{t}{L_g} \prod_{i=1}^p \left(I - \frac{1}{L_g} \nabla_{yy}^2 g(x, y; \xi_i^{(2)}) \right) \right] \nabla_y f(x, y). \quad (105)$$

We have

$$\begin{aligned} &\| \bar{\nabla}_x f(x, y) - \mathbb{E}[h_f^k] \| \\ &= \left\| \nabla_{xy}^2 g(x, y) \left\{ \mathbb{E} \left[\frac{t}{L_g} \prod_{i=1}^p \left(I - \frac{1}{L_g} \nabla_{yy}^2 g(x, y; \xi_i^{(2)}) \right) \right] - [\nabla_{yy} g(x, y)]^{-1} \right\} \nabla_y f(x, y) \right\| \\ &\leq C_{gxy} C_{fy} \left\| \mathbb{E} \left[\frac{t}{L_g} \prod_{i=1}^p \left(I - \frac{1}{L_g} \nabla_{yy}^2 g(x, y; \xi_i^{(2)}) \right) \right] - [\nabla_{yy} g(x, y)]^{-1} \right\|, \end{aligned}$$

where the last inequality follows from Assumption 1-3 and Assumption 2-5. Applying [22, Lemma 3.2], the latter norm can be bounded by $\frac{1}{\mu_g} \left(1 - \frac{\mu_g}{L_g} \right)^t$. This concludes the proof for the first part.

It remains to bound the variance of h_f^k . We first let $H_{yy} = \frac{t}{L_g} \prod_{i=1}^p \left(I - \frac{1}{L_g} \nabla_{yy}^2 g(x, y; \xi_i^{(2)}) \right)$. To estimate the variance of h_f^k , using (105), we observe that

$$\begin{aligned} \mathbb{E}[\|h_f^k - \mathbb{E}[h_f^k]\|^2] &= \mathbb{E}[\|\nabla_x f(x, y; \xi^{(1)}) - \nabla_x f(x, y)\|^2] \\ &\quad + \mathbb{E} \left[\left\| \nabla_{xy}^2 g(x, y; \xi_0^{(2)}) H_{yy} \nabla_y f(x, y; \xi^{(1)}) - \nabla_{xy}^2 g(x, y) \mathbb{E}[H_{yy}] \nabla_y f(x, y) \right\|^2 \right]. \end{aligned}$$

The first term on the right hand side can be bounded by σ_{fx}^2 . Furthermore

$$\begin{aligned} & \nabla_{xy}^2 g(x, y; \xi_0^{(2)}) H_{yy} \nabla_y f(x, y; \xi^{(1)}) - \nabla_{xy}^2 g(x, y) \mathbb{E}[H_{yy}] \nabla_y f(x, y) \\ &= \{ \nabla_{xy}^2 g(x, y; \xi_0^{(2)}) - \nabla_{xy}^2 g(x, y) \} H_{yy} \nabla_y f(x, y; \xi^{(1)}) \\ & \quad + \nabla_{xy}^2 g(x, y) \{ H_{yy} - \mathbb{E}[H_{yy}] \} \nabla_y f(x, y; \xi^{(1)}) \\ & \quad + \nabla_{xy}^2 g(x, y) \mathbb{E}[H_{yy}] \{ \nabla_y f(x, y; \xi^{(1)}) - \nabla_y f(x, y) \}. \end{aligned}$$

We also observe

$$\mathbb{E}[\|\nabla_y f(x, y; \xi^{(1)})\|^2] = \mathbb{E}[\|\nabla_y f(x, y; \xi^{(1)}) - \nabla_y f(x, y)\|^2] + \mathbb{E}[\|\nabla_y f(x, y)\|^2] \leq \sigma_{fy}^2 + C_y^2.$$

Using $(a + b + c)^2 \leq 3(a^2 + b^2 + c^2)$ and the Cauchy-Schwarz's inequality, we have

$$\begin{aligned} & \mathbb{E} \left[\left\| \nabla_{xy}^2 g(x, y; \xi_0^{(2)}) H_{yy} \nabla_y f(x, y; \xi^{(1)}) - \nabla_{xy}^2 g(x, y) \mathbb{E}[H_{yy}] \nabla_y f(x, y) \right\|^2 \right] \\ & \leq 3 \left\{ (\sigma_{fy}^2 + C_y^2) \{ \sigma_{gxy}^2 \mathbb{E}[\|H_{yy}\|^2] + C_{gxy}^2 \mathbb{E}[\|H_{yy} - \mathbb{E}[H_{yy}]\|^2] \} + \sigma_{fy}^2 C_{gxy}^2 \cdot \mathbb{E}[\|H_{yy}\|^2] \right\}. \end{aligned}$$

Next, a slight modification of the proof of Lemma 3.2 in [22] gives

$$\|\mathbb{E}[H_{yy}]\|^2 \leq (\mathbb{E}[\|H_{yy}\|])^2 \leq \mu_g^{-2}. \quad (106)$$

Together, the above gives the following estimate on the variance:

$$\mathbb{E}[\|h_f^k - \mathbb{E}[h_f^k]\|^2] \leq \sigma_{fx}^2 + \frac{3}{\mu_g^2} \left\{ (\sigma_{fy}^2 + C_y^2) \{ \sigma_{gxy}^2 + 2C_{gxy}^2 \} + \sigma_{fy}^2 C_{gxy}^2 \right\}. \quad (107)$$

This concludes the proof for the second part. \square

Finally, let us discuss the consequence of Lemma 11 on the sample complexity of the TTSA algorithm under the bias assumptions in Theorem 1, 2, Corollary 1. In particular, (103) shows that

$$\mathbb{E}[h_f^k] = \bar{\nabla}_x f(x^k, y^{k+1}) + B_k \quad \text{with} \quad \|B_k\| \leq b_k = \mathcal{O}((1 - \mu_g/L_g)^t). \quad (108)$$

Recall that the theorems require $b_k \leq c_b k^{-a}$ for some $a > 0$. To satisfy this, one only needs $t = \Theta(\log k)$. As such, the k -th iteration of TTSA requires $1 + \Theta(\log k)$ queries of stochastic oracles.

References

- [1] AGARWAL, A., KAKADE, S. M., LEE, J. D., AND MAHAJAN, G. (2019). Optimality and approximation with policy gradient methods in Markov decision processes. *arXiv preprint arXiv:1908.00261*.
- [2] ANTOS, A., SZEPESVÁRI, C., AND MUNOS, R. (2008). Learning near-optimal policies with bellman-residual minimization based fitted policy iteration and a single sample path. *Machine Learning* **71**, 1, 89–129.
- [3] BECK, A. (2017). *First-order methods in optimization*. Vol. **25**. SIAM.
- [4] BHANDARI, J., RUSSO, D., AND SINGAL, R. (2018). A finite time analysis of temporal difference learning with linear function approximation. In *Conference On Learning Theory*. 1691–1692.
- [5] BHATNAGAR, S., GHAVAMZADEH, M., LEE, M., AND SUTTON, R. S. (2008). Incremental natural actor-critic algorithms. In *Advances in Neural Information Processing Systems*. 105–112.

- [6] BHATNAGAR, S., PRECUP, D., SILVER, D., SUTTON, R. S., MAEI, H. R., AND SZEPESVÁRI, C. (2009). Convergent temporal-difference learning with arbitrary smooth function approximation. In *Advances in Neural Information Processing Systems*. 1204–1212.
- [7] BORKAR, V. S. (1997). Stochastic approximation with two time scales. *Systems & Control Letters* **29**, 5, 291–294.
- [8] BORKAR, V. S. AND PATTATHIL, S. (2018). Concentration bounds for two time scale stochastic approximation. In *Allerton Conference on Communication, Control, and Computing*. 504–511.
- [9] BRACKEN, J., FALK, J. E., AND MCGILL, J. T. (1974). Technical note—the equivalence of two mathematical programs with optimization problems in the constraints. *Operations Research* **22**, 5, 1102–1104.
- [10] BRACKEN, J. AND MCGILL, J. T. (1973). Mathematical programs with optimization problems in the constraints. *Operations Research* **21**, 1, 37–44.
- [11] CHEN, J. AND JIANG, N. (2019). Information-theoretic considerations in batch reinforcement learning. *arXiv preprint arXiv:1905.00360*.
- [12] COLSON, B., MARCOTTE, P., AND SAVARD, G. (2007). An overview of bilevel optimization. *Annals of Operations Research* **153**, 1, 235–256.
- [13] COUELLAN, N. AND WANG, W. (2016). On the convergence of stochastic bi-level gradient methods.
- [14] DALAL, G., SZORENYI, B., AND THOPPE, G. (2019). A tale of two-timescale reinforcement learning with the tightest finite-time bound. *arXiv preprint arXiv:1911.09157*.
- [15] DANN, C., NEUMANN, G., PETERS, J., AND OTHERS. (2014). Policy evaluation with temporal differences: A survey and comparison. *Journal of Machine Learning Research* **15**, 809–883.
- [16] DAVIS, D. AND DRUSVYATSKIY, D. (2018). Stochastic subgradient method converges at the rate $o(k^{-1/4})$ on weakly convex functions. *arXiv preprint arXiv:1802.02988*.
- [17] DOAN, T. T. (2020). Nonlinear two-time-scale stochastic approximation: Convergence and finite-time performance. *arXiv preprint arXiv:2011.01868*.
- [18] FALK, J. E. AND LIU, J. (1995). On bilevel programming, part I: General nonlinear cases. *Mathematical Programming* **70**, 47–72.
- [19] FINN, C., ABBEEL, P., AND LEVINE, S. (2017). Model-agnostic meta-learning for fast adaptation of deep networks. In *International Conference on Machine Learning*. 1126–1135.
- [20] FINN, C., RAJESWARAN, A., KAKADE, S., AND LEVINE, S. (2019). Online meta-learning. *arXiv preprint arXiv:1902.08438*.
- [21] FRANCESCHI, L., FRASCONI, P., SALZO, S., GRAZZI, R., AND PONTIL, M. (2018). Bilevel programming for hyperparameter optimization and meta-learning. In *International Conference on Machine Learning*. 1563–1572.
- [22] GHADIMI, S. AND WANG, M. (2018). Approximation methods for bilevel programming. *arXiv preprint arXiv:1802.02246*.
- [23] HONG, M., WAI, H., WANG, Z., AND YANG, Z. (2020). A two-timescale framework for bilevel optimization: Complexity analysis and application to actor-critic. *arXiv preprint arXiv:2007.05170*.
- [24] ISHIZUKA, Y. AND AIYOSHI, E. (1992). Double penalty method for bilevel optimization problems. *Ann Oper Res* **34**, 73–88.
- [25] JI, K., YANG, J., AND LIANG, Y. (2020). Provably faster algorithms for bilevel optimization and applications to meta-learning.
- [26] KAKADE, S. AND LANGFORD, J. (2002). Approximately optimal approximate reinforcement learning. In *International Conference on Machine Learning*. 267–274.
- [27] KAKADE, S. M. (2002). A natural policy gradient. In *Advances in Neural Information Processing Systems*. 1531–1538.

- [28] KALEDIN, M., MOULINES, E., NAUMOV, A., TADIC, V., AND WAI, H.-T. (2020). Finite time analysis of linear two-timescale stochastic approximation with Markovian noise. *arXiv preprint arXiv:2002.01268*.
- [29] KARMAKAR, P. AND BHATNAGAR, S. (2018). Two time-scale stochastic approximation with controlled Markov noise and off-policy temporal-difference learning. *Mathematics of Operations Research* **43**, 1, 130–151.
- [30] KONDA, V. R. AND TSITSIKLIS, J. N. (2000). Actor-critic algorithms. In *Advances in Neural Information Processing Systems*. 1008–1014.
- [31] KONDA, V. R., TSITSIKLIS, J. N., AND OTHERS. (2004). Convergence rate of linear two-time-scale stochastic approximation. *Annals of Applied Probability* **14**, 2, 796–819.
- [32] LI, J., GU, B., AND HUANG, H. (2020). Improved bilevel model: Fast and optimal algorithm with theoretical guarantee.
- [33] LIKHOSHERSTOV, V., SONG, X., CHOROMANSKI, K., DAVIS, J., AND WELLER, A. (2020). Ufo-blo: Unbiased first-order bilevel optimization.
- [34] LIU, R., MU, P., YUAN, X., ZENG, S., AND ZHANG, J. (2020). A generic first-order algorithmic framework for bi-level programming beyond lower-level singleton.
- [35] LUO, Z.-Q., PANG, J.-S., AND RALPH, D. (1996). *Mathematical Programs with Equilibrium Constraints*. Cambridge University Press.
- [36] MAEI, H. R., SZEPEŠVÁRI, C., BHATNAGAR, S., AND SUTTON, R. S. (2010). Toward off-policy learning control with function approximation. In *International Conference on Machine Learning*.
- [37] MEHRA, A. AND HAMM, J. (2019). Penalty method for inversion-free deep bilevel optimization. *arXiv preprint arXiv:1911.03432*.
- [38] MOKKADEM, A., PELLETIER, M., AND OTHERS. (2006). Convergence rate and averaging of nonlinear two-time-scale stochastic approximation algorithms. *Annals of Applied Probability* **16**, 3, 1671–1702.
- [39] MUNOS, R. AND SZEPEŠVÁRI, C. (2008). Finite-time bounds for fitted value iteration. *Journal of Machine Learning Research* **9**, May, 815–857.
- [40] PETERS, J. AND SCHAAL, S. (2008). Natural actor-critic. *Neurocomputing* **71**, 7-9, 1180–1190.
- [41] RAGHU, A., RAGHU, M., BENGIO, S., AND VINYALS, O. (2019). Rapid learning or feature reuse? towards understanding the effectiveness of MAML. In *International Conference on Learning Representations*.
- [42] RAJESWARAN, A., FINN, C., KAKADE, S., AND LEVINE, S. (2019). Meta-learning with implicit gradients. *arXiv preprint arXiv:1909.04630*.
- [43] SABACH, S. AND SHTERN, S. (2017). A first order method for solving convex bilevel optimization problems. *SIAM Journal on Optimization* **27**, 2, 640–660.
- [44] SCHULMAN, J., WOLSKI, F., DHARIWAL, P., RADFORD, A., AND KLIMOV, O. (2017). Proximal policy optimization algorithms. *arXiv preprint arXiv:1707.06347*.
- [45] SHABAN, A., CHENG, C.-A., HATCH, N., AND BOOTS, B. (2019). Truncated back-propagation for bilevel optimization. In *Artificial Intelligence and Statistics*. 1723–1732.
- [46] STACKELBERG, H. V. (1952). *The theory of market economy*. Oxford University Press.
- [47] SUTTON, R. S. (1988). Learning to predict by the methods of temporal differences. *Machine Learning* **3**, 1, 9–44.
- [48] SUTTON, R. S. AND BARTO, A. G. (2018). *Reinforcement learning: An introduction*. MIT press.
- [49] SUTTON, R. S., MAEI, H. R., PRECUP, D., BHATNAGAR, S., SILVER, D., SZEPEŠVÁRI, C., AND WIEWIORA, E. (2009). Fast gradient-descent methods for temporal-difference learning with linear function approximation. In *International Conference on Machine Learning*. 993–1000.
- [50] VICENTE, L., SAVARD, G., AND JUDICE, J. (1994). Descent approaches for quadratic bilevel programming. *Journal of Optimization Theory and Applications* **81**, 379–399.

- [51] WANG, M., FANG, E. X., AND LIU, H. (2017). Stochastic compositional gradient descent: algorithms for minimizing compositions of expected-value functions. *Mathematical Programming* **161**, 1-2, 419–449.
- [52] WHITE, D. J. AND ANANDALINGAM, G. (1993). A penalty function approach for solving bi-level linear programs. *Journal of Global Optimization* *3*, 397–419.
- [53] WU, Y., ZHANG, W., XU, P., AND GU, Q. (2020). A finite time analysis of two time-scale actor-critic methods. *arXiv preprint arXiv:2005.01350*.
- [54] XU, T., WANG, Z., AND LIANG, Y. (2020). Non-asymptotic convergence analysis of two time-scale (natural) actor-critic algorithms. *arXiv preprint arXiv:2005.03557*.