

Analyzing ribosome remodeling in health and disease

Aleksandra A. Petelski,^{1,2,3} & Nikolai Slavov^{1,2,3,✉}

¹Department of Bioengineering, Northeastern University, Boston, MA 02115, USA

²Barnett Institute, Northeastern University, Boston, MA 02115, USA

³Department of Biology, Northeastern University, Boston, MA 02115, USA

✉ Correspondence: nslavov@alum.mit.edu or nslavov@northeastern.edu

Increasing evidence suggests that ribosomes actively regulate protein synthesis. However, much of this evidence is indirect, leaving this layer of gene regulation largely unexplored, in part due to methodological limitations. Indeed, we review evidence demonstrating that commonly used methods, such as transcriptomics, are inadequate because the variability in mRNAs coding for ribosomal proteins (RP) does not necessarily correspond to RP variability. Thus protein remodeling of ribosomes should be investigated by methods that allow direct quantification of RPs, ideally of isolated ribosomes. We review such methods, focusing on mass spectrometry and emphasizing method-specific biases and approaches to control these biases. We argue that using multiple complementary methods can help reduce the danger of interpreting reproducible systematic biases as evidence for ribosome remodeling.

Introduction

The control of gene expression is crucial for all biological processes, from developmental stages and homeostasis maintenance to regeneration processes. This regulation occurs at multiple layers, both transcriptional and post-transcriptional levels. Historically, transcription has been studied more extensively than translation, in large part because of the accessibility of technologies for nucleic acid analysis. However, gene regulation via translation was appreciated as early as the late 1960's. For example, the production of insulin was linked to the increased number of polysomes and protein synthesis [1]. Similarly, the synthesis of globin (hemoglobin polypeptides) was found to depend on heme, the oxygen-carrying iron-rich molecule in the blood [2]. In the early 1980s, different aspects of early embryonic development, such as blastocyst formation [3] and disparate patterns of protein synthesis [4] were linked to post-transcriptional regulation. Key developmental genes, such as Oscar, are regulated at the level of translation and spatial localization [5, 6]. Such post-transcriptional regulation is mediated at least in part by translational control mechanisms including RNA binding proteins, translation factors, and micro RNAs [7, 8].

In addition to these well-established mechanisms, increasing evidence suggests that ribosomes may also regulate translation of mRNAs [9–14]. Ribosomes have long been viewed as passive players in translation, with a fundamental role of catalyzing peptide-bond formation but exerting regulatory effects only based on their availability [15, 16] and translational control being exerted by a teamwork of cis- and trans-regulators in which the interaction of mRNA structures and sequences worked with translation factors, RNA binding proteins, and microRNAs; the ribosome complex was simply viewed as an effector of translation. However, ribosome-mediated regulation through the alteration of ribosomal RNA and ribosomal proteins were both hypothesized [9, 11, 17] and supported by mostly indirect evidence as recently reviewed by Emmott et al. [14]. Quantification of ribosomal proteins from isolated ribosomes has begun to provide more direct evidence, as mammalian ribosomes have been found to exhibit differential protein stoichiometry that depends on the growth conditions and on the number of ribosomes per mRNA [18]. Such observations suggest that different ribosome complexes may exist in order to fulfill disparate functions, which can consequentially have regulatory effects on translation. This model, termed ribosome specialization, challenges the notion that ribosomes are static enzymes and instead introduces them as active participants in post-transcriptional regulation. In this model, ribosomes could be specific to distinct cell population and can affect the translation of mRNAs. The hypothesis of ribosome-mediated translational regulation has been further supported by observations of differences in ribosomal protein composition occurring under stress [19–21] and cell differentiation [22–25]. Localized synthesized RPs in the axon can also contribute to ribosome remodeling [26, 27], suggesting possible roles of ribosome-mediated translational control in neuronal functions. The possibility of ribosome specialization is further supported by the observations that mutations of specific ribosomal proteins selectively affect the synthesis of specific proteins and are strongly associated with distinct phenotypes such as cancer and aging [28, 29]. Such selectivity suggests that ribosomes can regulate gene expression [10, 11, 14, 30]. Interestingly, some RPs, such as RACK1, have been observed to dynamically associate and dissociate from ribosomes and specifically affect the translation of short mRNAs [31–33].

Specialized ribosomes have been suggested to hold specific functional roles, especially in the context of immunology and cancer. The idea of ribosomes driving cancer progression could be tied to observations of disease states characterized by dysfunctional ribosomes, disorders collectively known as ribosomopathies [34, 35]. Patients with such disorders showed increased risk for diseases of uncontrolled cell growth, such as cancer, later in life [36]. More direct evidence has shown the association of RP gene mutations with numerous cancers, raising the prospect of the

existence of oncoribosomes [37, 38]. Ribosomes have been also implicated in the immunosurveillance of cancer and other types of pathogenic cells. MHC class I molecules, which are important in alerting the immune system when cells are virally infected, are believed to be derived from DRiPs (defective ribosomal products), unstable molecules that degrade much more quickly than functional proteins that are in stable conformation [39]. In order to rapidly produce such products, a subset of ribosomes could be assigned to synthesize DRiPs that have enhanced antigen presentation [40, 41]. Immunoribosomes may also serve as an efficient source of peptides that can stimulate antibody production upon the invasion of pathogenic molecules. These models of functionally specialized immunoribosomes and oncoribosomes remain insufficiently tested, and further testing would benefit from the approaches discussed in this review.

Ribosomes in eukaryotes are made of about 80 core ribosomal proteins and four ribosomal RNAs (rRNAs) [42]. The modifications of both molecular groups are important in the functionality of the ribosome complex and are potential sources of ribosome heterogeneity and specialization [43, 44]. Modifications of rRNA, such as methylation and pseudouridylation, which together span a total of 7000 nucleotides, are known to stabilize the ribosome structure. Variant rRNA alleles and rRNA methylation may contribute to ribosomal specialization. In both mice and humans, several rRNA sequence variants were identified and shown to exhibit tissue-specific expression; furthermore, at least 23 percent of rRNA nucleotides are estimated to exhibit variant alleles within the general human population [45]. On average, 32 variants were found to be expressed in single individuals, while those sequence variants were found to significantly overlap with sequences that are functionally important to ribosome function. The diversity of rRNA variants are thus suggested to have a biologically important role. Indeed, the methylation of the 16S rRNA at a specific guanosine nucleotide revealed that this modification plays a role in controlling mistranslation and could explain streptomycin-resistant phenotypes of *M. tuberculosis* [46]. Both rRNA and protein modifications may contribute to ribosome specialization; The role of rRNA has been reviewed by Mauro and Matsuda [47], and in this review we will focus on methods for investigating ribosome specialization via modifications of the core ribosomal proteins.

Post-transcriptional regulation of ribosomal proteins

While suggestive evidence for ribosome remodeling has originated from indirect methods (such as measuring transcripts coding for RPs [50, 51]), such data remain inconclusive because RP synthesis and degradation are extensively regulated [48, 49, 52, 53]. RP molecules that are not incor-




Figure 1 | Ribosomal proteins are under post-transcriptional control (a) A heatmap showing the levels of ribosomal genes at the levels of RNA, ribosome density, and protein in lymphoblastoid cell lines based on data from Battle *et al.* [48]. The data points for each gene are displayed as \log_2 fold-changes relative to their mean. The corresponding distributions of fold changes for all genes across all cell lines are shown to the right. To control for variable input amounts from different cell lines, the data from each cell line were normalized to the same total amount of RP gene products. (b) Protein to mRNA ratios (PTRs) were quantified across different human tissues, and gene sets with statistically significant shifts in the tissue-type specific relative protein to RNA ratios (rPTR) are highlighted. Of interest, ribosomal proteins have high rPTRs in the kidney and low rPTRs the stomach, indicating significant post-transcriptional regulation. This panel is based on the analysis by Franks *et al.* [49].

porated into a complex are rapidly degraded[52]. Because of this post-transcriptional regulation, analysis of ribosome remodeling in health and disease should rely on direct protein measurements. While the abundance of RP transcripts are usually the most accessible data, these measures are indirect. Consider, for example, the variability of RP transcripts, ribosome density, and ribosomal proteins across a panel of lymphoblastoid cell lines shown in Fig. 1a. The data indicate substantial transcript variability, which diminishes at the level of ribosome density and is almost absent for the ribosomal proteins, Fig. 1a. Thus, variable RP transcripts do not necessarily indicate variable abundance of RPs, Fig. 1a. More generally, the ratio between RPs and their corresponding transcripts can vary substantially as shown in Fig. 1b for two tissues and observed in many other cases [49, 54]. Therefore, the levels of mRNAs coding for RPs are rather indirect evidence to support conclusions about the protein composition of ribosomes.

To obtain more direct evidence for protein remodeling of ribosomes, one should directly measure RP abundances as a more direct approach for evaluating ribosome remodeling across different conditions, Fig. 1a. Such analyses in cell lysates have suggested changes in the protein composition of ribosomes as budding yeast undergoes the diauxic shift [55], during aging [56, 57], and upon LPS-stimulation of mouse dendritic cells[58]. In addition, RP analysis from cell lysates has shown that ribosomal transcripts exhibit slower elongation rates with decreased protein production

relative to other transcripts with similar ribosome densities[59].

Although RPs are known to degrade very quickly when not incorporated into complexes[52], RPs quantified in total cell lysates may originate in part from other extraribosomal complexes [17, 60–64]. Excluding the influence of such extraribosomal complexes requires the analysis of isolated ribosomes. The use of isolation methods, such as sucrose gradient fractionation or affinity purification[65], prior to protein measurements provides an even more direct way to assess ribosome remodeling in different conditions [18, 66, 67].

In some studies, the measurement noise is comparable to the variability of RPs across conditions [68]. In such cases, we may conclude that ribosomes do not remodel across the examined conditions or the degree of remodeling is too small to be detected by the methodology used. The smaller the measurement errors, the more confident we may be that ribosomes do not remodel across the set of studied conditions. For example, the degree of ribosome remodeling (if any at all) is very small during the aging of mouse brains [68]. As observed RP changes can be small, they may be comparable to or smaller than the measurement noise.




Figure 2 | Different methods provide different estimates of ribosomal protein abundance. A scatter plot comparing estimated abundance of ribosomal proteins from Western blots [69] and from flow cytometry analysis of RPs tagged by green fluorescent protein (GFP)[70]. The Pearson correlation between the two estimates is modest, $\rho = 0.25$, despite the fact that the measurements by each method are highly reproducible. The weak correlation may reflect shared biases from the construction of the tagged proteins or biological differences in protein abundance. The first option seems much more likely, but it remains inconclusive without additional data and analysis.

Analyzing ribosome remodeling in individual cells

Ribosome remodeling is likely to contribute to the specialized proteomes of the diverse cell types that arise during development and diverge to form different tissues [14]. Furthermore, cyclic transcription of mRNAs coding for RPs suggests that ribosome biogenesis is highly temporarily organized and metabolically coordinated during the cellular life cycle [74–76]. While these possibilities are of considerable interest, their investigation poses particular challenges for direct RP quantification by complementary methods [77, 78]. These challenges stem from the difficulty of quantifying proteins in tissues comprised of heterogeneous cells. Single-cell proteomics has approached this question in the context of embryonic stem cells differentiating to epiblast lineages in embryoid bodies [73]. In this system, the correlations among RPs (Fig. 3) revealed one large and one small cluster, suggesting that the RPs from the small cluster covary in a cell-type specific manner. However, these data have not been cross-validated by independent methods and thus remain inconclusive. Nonetheless, advances in single-cell proteomics hold the potential to increase the reliability of single-cell protein analysis and to enable cross-validation of such measurements [77–80]. Thus these technological advances may soon enable direct examination of ribosome specialization within the diverse cell types that comprise different tissues.

The need for complementary methods

The technical biases of each method type are challenging to overcome, but the use of complementary methods with divergent weaknesses can help guard against the potential influence of systematic artifacts on the final results. Thus, the use of complementary techniques allows for both the attainment of strong evidence for ribosome remodeling and the cross-validation of results that can support more confident conclusions. As an example, RPs from isolated ribosomes may be quantified separately by both mass spectrometry and Western blots and then the results can be compared as shown in Fig. 4. The measured RP differences obtained from these two disparate methods are highly similar, indicating that the results are likely to be driven by biological effects rather than the inherent biases attributed to each method. More broadly, if biases were the main force behind the results, the outcomes would have been presumably different; thus, the more different the methods, the less likely they are to share biases, and the more beneficial they are in being used together as complementary methods.

Direct quantification of RPs with complementary methods (as shown in Fig. 4) can reveal changes in the ribosome complex that may contribute to post-transcriptional regulation of gene




Figure 4 | Quantifying ribosomal proteins with complementary methods Monosomal and polysomal ribosomes from embryonic stem cells were isolated by sucrose gradients and analyzed by mass-spectrometry and by Western blots [18]. The results exemplify qualitative agreement for the polysomal enrichment of Rps29 and Rps14 as well as relatively small differences across the two methods. The differences may be due both to the different cell lines used or to the quantification biases of each method [18]. **(a)** The Polysomal enrichment of RPs was quantified by mass-spectrometry using isobaric tags. The analysis was performed by digesting the RPs either with trypsin or with Lys-C and the results provided consistent estimates that are combined in these boxplots [18]. **(b)** Polysomal enrichment of RPs quantified by Western blots [18]. RPs were quantified by Western blots in monosomes and polysomes from E14 mouse ESCs. Rpl32 was used as a loading control and the boxplots summarize data from 9 ratios for each quantified RP. The panels are reproduced from Slavov *et al.* [18].

expression [18]. Mass spectrometry (MS) offers an array of such complementary approaches for discovering and validating ribosome remodeling. These powerful methods directly measure protein or peptide abundance, either through relative or absolute quantification, and can provide information about the stoichiometry of the ribosome complex along with modifications of the ribosomal proteins. These results can be enhanced and cross-validated by structural methods, such as X-ray crystallography and cryogenic electron microscopy (cryo-EM). These latter methods can directly probe the arrangement of ribosomal proteins within the complex. In this review, we discuss the types of biases, limitations, and advantages associated with each approach type, focusing particularly on the investigation of ribosome remodeling.

Quantifying ribosomal proteins via mass spectrometry

The levels of ribosomal protein expression can be directly measured using mass spectrometry proteomics. Whole proteins, or even entire ribosome complexes can be analyzed by top-down mass spectrometry [81, 82]. However, because MS analysis of full length proteins is technologically challenging, it is more common to first digest proteins into peptides and then analyze the peptides by MS. This latter approach is known as bottom-up mass spectrometry, which involves the caveat of inferring protein levels from digested peptides. These two branches (top-down and bottom-up) of mass spectrometry can be implemented by many methodologies, each of which has systematic biases and can directly quantify RP stoichiometry across biological conditions.

Top-down and bottom-up proteomics methods provide complementary measurements, as they quantify proteins in inherently different ways and thus can help detect and mitigate biases. Technical biases that can affect the ultimate biological interpretation of any experiment can be potentially introduced at any step of the proteomics pipeline, from cell lysis and sample preparation to MS analysis. These distinct biases can be controlled by using different methods. Ultimately reliable results require cross-validating the results from different methods that share as few biases as possible.

Biases in bottom-up mass spectrometry

Bottom-up proteomics can be implemented by many methods, all of which have caveats, specifically for quantifying ribosomal proteins. The amount of ribosomes in most cells is generally ample, about 10 million in a typical mammalian cell. Yet, the sequences of ribosomal proteins are rather short, with a median length of about 133 amino acids in human cells, in stark contrast to the overall proteome sequence length, which is 375 amino acids. Thus, ribosomal proteins tend to produce fewer peptides during digestion. In addition, ribosomal proteins are likely to produce shorter peptides when using trypsin, the most commonly used protease, due to a large number of arginine (R) and lysine (K) amino acids present in the protein sequences [83]. The over representation of K and R may contribute to a higher miscleavage rate, which can complicate analyses. These downsides can be taken into account by the use of multiple alternative proteases such as Lys-C and Glu-C in order to cross-validate results with the commonly used protease of trypsin [84, 85]. These biases can be very significant when estimating absolute protein abundances (due to differences in peptide flyability, a collective term describing efficiency of ionization and detection), and these biases may cancel out in relative protein quantification [86, 87].

Quantifying protein stoichiometry by bottom-up mass spectrometry

The differences in peptide flyability and other peptide specific biases pose a major challenge to quantifying stoichiometry (ratios) between different proteins and their proteoforms [86, 88]. Indeed, a peptide might be more intense because it is delivered more efficiently to the MS analyzer rather than because it originates from a more abundant protein. These peptide specific biases cancel out when performing relative quantification of a protein across different samples. This idea of canceling out biases can be extended by first-principle models, such as HIquant, to allow quantifying stoichiometry between different proteins independent from peptide specific biases [87]. Such an approach is likely to be particularly fruitful for quantifying RP preforms originating from different prologues, alternative splicing events or post-translational modifications [14, 87, 89].

In bottom-up proteomics, cells can be lysed and proteins extracted by a variety of methods [90–92]. Then proteins are digested to peptides, which are separated via liquid chromatography (LC) and ionized through electrospray ionization (ESI) or matrix-assisted desorption/ionization (MALDI). Then, these peptides are introduced into the mass spectrometer as precursor ions, which can be used for direct quantification or can be further isolated and disintegrated into fragment ions that are then used for sequence identification and quantification. Bottom-up MS may either simultaneously isolate and fragment multiple peptides in parallel (known as DIA; data independent acquisition), or analyze a single peptide at a time (known as DDA; data dependent acquisition). DDA methods were introduced in 1990s and have been widely used for decades [93, 94]. DIA was introduced later (2004) by Yates and colleagues [95] and has matured into methods that offer the advantage of parallel analysis and afford identifying and quantifying many thousands of peptides in a single run [96, 97].

Quantification of peptides can be based directly on precursor ions (MS1-based) or fragment ions (MS2-based). Both of these peptide quantification approaches can be implemented by multiple methods. These methods differ in the way and extent to which they control technical biases stemming from the proteomics pipeline, from sample preparation procedures to LC-MS analysis. Wet-lab procedures encompass the experimental steps of isolating ribosomes from cell lysates, protein digestion into peptides, and various labeling techniques. In LC-MS analysis, biases can be introduced due to differential peptide separation, ionization, and isolation for MS2 analysis. The available bottom-up methods control for these technical biases at different stages. Due to the disparities in biases, different bottom-up approaches can be viewed as highly complementary techniques that allow us to gain more confidence in results showing biological changes, especially in ribosome remodeling where the changes may be small.

Control of biases at the MS1-Level of Peptide Quantification

Methods that use the MS1 level in order to quantify peptides can be further classified either as labeled or label-free approaches. The label-free approaches are attractive for their simplicity and fewer experimental steps. In these methods, each sample undergoes the sample preparation and mass spectrometry analysis separately. In contrast, labeled techniques allow analyzing multiple samples in parallel and thus more opportunities to control for biases that occur during the parallel stages. Labeling approaches for MS1-based quantification include: SILAC (Stable Isotope Labeling of Amino Acids in Cell Culture), dimethyl labeling, and mTRAQ. SILAC introduces sample labeling to living cells during protein synthesis through the metabolic labeling of newly synthesized proteins [98]. More specifically, cells are incubated in cell culture medium that contains stable-isotope enriched amino acids, usually arginine and lysine [99, 100], which are then incorporated into new proteins. Dimethyl labeling, on the other hand, introduces sample labeling after digestion. This chemical labeling approach relies on the reaction between formaldehyde and sodium cyanoborohydride with lysine side chains and N-terminal primary amines in order to form dimethylamines [101, 102]. mTRAQ quantification, which also introduces labels after digestion. It was designed specifically for targeted MS, in which specific peptides found in previous shotgun runs could be probed [103, 104]. All of these methods quantify peptides at the level of the precursor ions.

The introduction of biases can be controlled through mixing the samples as early as possible within the experimental pipeline. The earlier the samples are combined together, the sooner the samples can be exposed to identical conditions and the sooner biases can be controlled. Among the available MS methods, SILAC allows mixing of samples earliest, even prior to ribosome isolation and MS procedures, thus enabling the control of all biases associated with these experimental and MS steps. This is especially advantageous when studying subcellular fractions, such as the ribosome complex. Labeling methods, such as dimethyl labeling and mTRAQ, allows the mixing of samples prior to LC separation and MS analysis, which controls for LC- and MS-related biases, while label-free methods will not control for these biases, as each sample is analyzed individually.

The processes of peptide separation via liquid chromatography and subsequent MS analysis can introduce technical biases that MS1-based labeling methods can control but label-free cannot. Each run in the mass spectrometer is subject to a host of variable factors that might influence measured ion intensities, including variability in peptide separation, ionization, and instrumentation. This can cause different runs to experience drifts in retention time and m/z , leading to complicated analyses. Ion suppression, a problematic phenomenon that affects the final amount of charged ions

that ultimately reaches the detector, can also undermine quantitative accuracy. In the cases of labeling, digested peptides from each sample are separated, ionized, and analyzed together, allowing for the samples to experience the same nuances associated with each of those steps. Label-free approaches in particular suffer from the fact that large portions of peptides are not detected in every sample, which is termed as the missing value problem[105]. This problem makes it more difficult to compare RP abundance across different conditions. Advances relying on matching peptide intensity readouts, i.e. MaxLFQ[106], and on enhanced peptide identifications via DIA [95–97] or DDA methods incorporating retention time information [78, 107, 108] can mitigate the missing value problem when using label-free approaches.

While MS-1 based labeling methods allow controlling for biases during peptide separation and ionization, labeling itself can introduce biases, limitations, and artifacts. SILAC introduces heavy isotopes into live cells and animals, which can induce unintended growth changes[109, 110] and may even affect behavioral characteristics of mice[111, 112]. Also, depending on the model system used, the potential conversion of arginine to proline may lead to underestimation of heavy-labeled peptides[113]. An additional limitation when using SILAC is that the time to fully label a biological system of interest can take days to weeks. Dimethyl labeling, on the other hand, takes around 5 minutes and is highly cost-effective, as the reagents are inexpensive [114]. However, dimethyl labeling has been associated with a loss of hydrophilic peptides[115], which can lead to fewer overall peptide identifications. Overall, any type of labeling approach should be assessed for completeness of labeling to allow for confident quantitation of RP remodeling.

MS1-based quantification techniques share some common limitations. A disadvantage of MS1-based labeling methods is that the number of multiplexed samples is usually limited to 2 or 3. As the number of isotopically labeled samples increases, so does the number of precursor ions in MS1 scans. The high density of ions may lead to interference between coeluting ions with very similar m/z ratios. The use of NeuCode metabolic labeling offers higher SILAC multiplicity through the use of labels that differ in mass on the scale of milli-Daltons[116, 117]; thus, this technique requires higher resolving power of the mass analyzers [116, 117]. Because of the inherent limitations that MS1-level methods present, it is important to consider using complementary methods rather than multiple technical replicates of just one method type. MS1-based methods can be co-validated with MS2-based approaches, which still come with their own biases; however, the set of biases derived from both quantification approaches are different, allowing for increased confidence in quantifying even small RP changes.

Control of biases at the MS2-Level of Peptide Quantification

As opposed to MS-1 based methods, precursor ions are isolated and then fragmented, and the resulting (fragment) ions analyzed by another (MS2) scan. In most cases, MS2-based quantification is used with isobaric tags, such as tandem mass tags (TMT)[118] or Isobaric Tags for Relative and Absolute Quantification (iTRAQ)[119, 120]. DIA analysis can also benefit from incorporating MS2-level data in order to improve the accuracy of label-free quantification [121].

With isobaric labeling, peptides of each sample are covalently labeled with a sample-specific mass tag. Mass tags are comprised of a reporter ion, a balance group and a reactive group, which is usually an amine reactive group (NHS group) that binds to primary amines from the peptides. After labeling, the samples are mixed together, allowing the analysis multiple samples (up to 16 with TMT Pro) in a single mass spectrometry run. Thus, the multiplexed samples undergo ionization and ion selection for MS2 analysis together, greatly mitigating the missing value problem. Additionally, the multiplexing permits analyzing more samples per unit time. Another advantage is that the number of precursor ions detected during MS1 scans does not increase with the number of samples, as each isobaric tag has an identical mass which allows a particular peptide from different samples to appear as a single feature in the ion map defined by retention times and m/z ratios. Peptides from different samples are thus indistinguishable at the MS1 level. Upon isolation and fragmentation of precursor ions, they release distinct reporter ions and peptide fragments, some of which remain bound to the balance group. Since the reporter ions and the balance groups have sample-specific number of heavy isotopes, they allow for the ratiometric quantification of samples.

Since all peptide from a sample release the same reporter ion upon fragmentation, the accuracy of quantifying a peptide from its reporter ions is dependent on isolating only its precursor ion for subsequent fragmentation and MS2 analysis. In practice, the MS2 scan typically also contains coisolated peptides. The degree of coisolation can be estimated and used to remove peptides with unacceptably high coisolation. Narrower isolation windows [79] and sampling elution peaks at their apexes [122] reduce coisolation but may not completely eliminate it. When multiple peptide precursors are coisolated for fragmentation, the measured reporter ion intensities are proportional to the superposition of peptide abundances, which results in inaccurate quantification and generally compressed peptide ratios. However, ratio compression is not necessarily due to coisolation since it may be caused by other factors, such as unintentional cross-labeling or sample carry-over on the LC column. Coisolation can be reduced by subjecting fragment ions to further isolation and fragmentation steps by taking MS3 or even higher scans[123]. This MS3 approach reduces the number of analyzed ions and thus diminishes the sensitivity and the throughput of the analysis. A

second option for reducing bias due to coisolation is to use the peptide fragments with attached balance groups, which are termed complement ions [124]. These complement ions are produced during the fragmentation step as a result of the mass balance group remaining attached to the peptide or its fragments, and thus they can be specific to the analyzed peptide.

Abundant peptides, such as those originating from RPs, tend to be less affected by coisolation since the majority of the reporter ions used for quantification will be derived from the abundant peptides. However, coisolation can still contribute significant bias to RP quantification and thus necessitates quality controls. One way to benchmark data quality is through the calculation of the agreement of different peptides originating from the same protein. Ribosomal peptides that originate from a RP should indicate consistent quantification of the protein; the degree of consistency can be quantified by measures such as reliability or coefficients of variation [49].

The methods of quantifying peptides at either the MS1 or MS2 levels come with inherent biases that might complicate analyses and may present reproducible results that are actually artifacts induced from systematic biases. However, the set of biases for each method type is different, and thus different methods may complement each other. Instead of choosing just one mass spectrometry method to identify changes of ribosomal proteins, methods that differ as much as possible (such as MS1 and MS2 based quantification) or non mass-spec methods should be used in parallel (as shown in Fig. 4) so that the results can be co-validated.

Top-down mass spectrometry

With top-down proteomics [81, 82], whole ribosomal proteins or even whole ribosomes can be analyzed, offering a more intact picture of the ribosome and potential characterization of proteoforms and post-translational modifications (PTMs). These methods identify the full amino acid sequence of a protein including modifications and thus do not require protein inference from peptides. However, these techniques are more challenging because compared to peptides, proteins are more difficult to solubilize, separate, ionize and sequence [125]. These challenges result in lower sensitivity and throughput of the analysis, as well as higher technical requirements from the instruments for high resolving power at high m/z ratios [126–128]. Nonetheless, these challenges are rapidly being addressed by innovative methods for protein separation (such as capillary zone electrophoresis [129]), sensitive methods allowing the detection of individual ions [130, 131], and by community standards [132].

The biases associated with either bottom-up or top-down proteomics are very different, which presents opportunities for combining them synergistically. The hybrid combination of mass spec-

trometry techniques has been used to characterize whole ribosome complexes by quantifying RP levels along with proteoforms [127]. Such an approach has offered novel observations about cysteine modification of RPS27, and ribosome assembly sites through the characterization of ribosomes originating from human, plant, and bacterial cells [133]. Additionally, the elusive ribosomal protein SRA was successfully quantified and found to have heterogeneous stoichiometry in *E. coli* ribosomes [127].

Complementary structural biology techniques

We have focused our review on the higher throughput MS methods, but many structural biology methods can provide more detailed information about ribosome modifications and structure albeit at lower throughput. These include chemical cross linking of proteins [134, 135], X-ray diffraction, and cryo-EM. The analysis of both protein sequence in tandem with spatial information allows not only to co-validate results ribosome remodeling observations but also to start revealing its functional significance. Such a combined approach has been used to study other complexes, such as the human nuclear pore complex scaffold [136], bacteriophage portal complexes[137], 26S proteasome complex[138] and COPII vesicles that transport proteins to the Golgi apparatus from the endoplasmic reticulum [139]. In fact, such combination of structural biology and mass spectrometry has been used to study a particular subcomplex involved in ribosome biogenesis [140, 141]. The synergy between these methods can help identify specialized ribosomal structures and confirmations regulating protein synthesis.

X-ray diffraction requires the crystallization of the sample of interest prior to analysis, while cryo-EM eliminates this need. The process of crystallization can demand much optimization, which can involve the purification of many ribosomes. The obviation of crystallization allows for the study of more kinds of ribosomes without the limitation of crystallizing them. The flash-freezing of samples allows the observation of molecules in a "near-native" state; since molecules are not constrained to a crystal, they have more degrees of freedom, thus allowing for more conformational states to be studied. However, cryo-EM studies can require anywhere from hundreds to thousands single particle images, which involves long hours of image acquisition at the microscope. Furthermore, many of these images are discarded due to the phenomenon known as beam-induced motion, which produces blurred images [142]. Despite limitations, both technique types continue to reveal the ribosome protein structure. Recently, X-ray diffraction was used to discover the role of potassium ions within the ribosome on the stabilization of the protein complex

in both the initiation and elongation states of translation [143]. Cryo-EM has been nearing the sensitivity and resolution of X-ray diffraction, as seen through the report of the entire bacterial ribosome resolved at two angstroms [144].

The synergy of mass spectrometry and structural biology techniques can empower novel observations of the ribosome complexes. These method types are different in terms of sample preparation and measurement acquisition, yet are complementary in the type of information that is provided. Combining protein composition information with conformational changes can help reveal different roles of ribosomes within protein synthesis, and ultimately gene expression. Such knowledge can help decipher the degree of ribosome remodeling during normal development and physiology (e.g., immunoribosomes) and during diseases (e.g., oncoribosomes). The combination of mass spectrometry with structural biology has been used to elucidate several facets of the ribosome complex subunits, including ribosome assembly in bacteria [140, 145, 146] and the effect of dimerization on ribosomes when nutrients are scarce [147]. In addition, ribosome remodeling has been found to occur in the bacterial ribosome through the use of MS and X-ray crystallography [148].

This paper emphasized the need to use multiple complementary approaches for quantifying ribosome remodeling and briefly reviewed sources of systematic biases as well as approaches for mitigating their influence. These approaches may afford reliable quantification of ribosome remodeling, which is a starting point for investigating its functional roles in regulating mRNA translation. Identifying ribosomes associated with specific conditions – such as disease states, developmental stages, or metabolic conditions – can start to reveal different populations of ribosomes. These observations will serve as a starting basis for characterizing new principles in the regulation of RNA translation that may reshape our understanding of one of the most fundamental biological processes. In the long term, this new understanding can enable the design of therapies that specifically target translation for disease and regenerative treatments.

Acknowledgments: This work was funded by a New Innovator Award from the NIGMS from the National Institutes of Health to N.S. under Award Number DP2GM123497.

References

1. Wool I, Stirewalt WS, Kurihara K, Low RB, Bailey P, et al. (1968) Mode of action of insulin in the regulation of protein biosynthesis in muscle. *Recent progress in hormone research* 24: 139–213.

2. Levere RD, Granick S (1965) Control of hemoglobin synthesis in the cultured chick blastoderm by delta-aminolevulinic acid synthetase: increase in the rate of hemoglobin formation with delta-aminolevulinic acid. *Proceedings of the National Academy of Sciences of the United States of America* 54: 134.
3. Braude PR (1979) Control of protein synthesis during blastocyst formation in the mouse. *Developmental biology* 68: 440–452.
4. Rosenthal ET, Hunt T, Ruderman JV (1980) Selective translation of mrna controls the pattern of protein synthesis during early development of the surf clam, *spisula solidissima*. *Cell* 20: 487–494.
5. Markussen FH, Michon AM, Breitwieser W, Ephrussi A (1995) Translational control of oskar generates short OSK, the isoform that induces pole plasma assembly. *Development* 121: 3723–3732.
6. Martin KC, Ephrussi A (2009) mRNA localization: gene expression in the spatial dimension. *Cell* 136: 719–730.
7. Castello A, Fischer B, Eichelbaum K, Horos R, Beckmann BM, et al. (2012) Insights into RNA biology from an atlas of mammalian mRNA-binding proteins. *Cell* 149: 1393–1406.
8. Bartel DP (2009) Micrnas: target recognition and regulatory functions. *cell* 136: 215–233.
9. Mauro VP, Edelman GM (2002) The ribosome filter hypothesis. *Proceedings of the National Academy of Sciences* 99: 12031–12036.
10. Komili S, Farny NG, Roth FP, Silver PA (2007) Functional specificity among ribosomal proteins regulates gene expression. *Cell* 131: 557–571.
11. Gilbert WV (2011) Functional specialization of ribosomes? *Trends in biochemical sciences* 36: 127–132.
12. Slavov N (2015) From differential transcription of ribosomal proteins to differential structure of ribosomes doi:10.7287/peerj.preprints.1504v1.
13. Preiss T (2015) All ribosomes are created equal. really? *Trends in biochemical sciences* 41: 10.1016/j.tibs.2015.11.009.
14. Emmott E, Jovanovic M, Slavov N (2018) Ribosome stoichiometry: from form to function. *Trends in biochemical sciences* 44: 95–109. doi:10.1016/j.tibs.2018.10.009.
15. Lodish HF (1974) Model for the regulation of mrna translation applied to haemoglobin synthesis. *Nature* 251: 385–388.
16. Mills EW, Green R (2017) Ribosomopathies: Theres strength in numbers. *Science* 358.

17. Naora H, Naora H (1999) Involvement of ribosomal proteins in regulating cell growth and apoptosis: translational modulation or recruitment for extraribosomal activity? *Immunology and cell biology* 77: 197–205.
18. Slavov N, Semrau S, Airoidi E, Budnik B, van Oudenaarden A (2015) Differential stoichiometry among core ribosomal proteins. *Cell Reports* 13: 865 – 873. doi: 10.1016/j.celrep.2015.09.056.
19. Ferretti MB, Ghalei H, Ward EA, Potts EL, Karbstein K (2017) Rps26 directs mRNA-specific translation by recognition of kozak sequence elements. *Nat Struct Mol Biol* 24: 700–707.
20. Samir P, Browne CM, Rahul, Sun M, Shen B, et al. (2018) Identification of changing ribosome protein compositions using mass spectrometry.
21. Ghulam MM, Catala M, Abou Elela S (2020) Differential expression of duplicated ribosomal protein genes modifies ribosome composition in response to stress. *Nucleic Acids Research* 48: 1954–1968.
22. Kawasaki H, Shigenaga A, Uda M, Baba T, Ogawa H, et al. (2011) Nitration of tryptophan in ribosomal proteins is a novel post-translational modification of differentiated and naïve pc12 cells. *Nitric Oxide* 25: 176–182.
23. Akanuma G, Nanamiya H, Natori Y, Yano K, Suzuki S, et al. (2012) Inactivation of ribosomal protein genes in bacillus subtilis reveals importance of each ribosomal protein for cell proliferation and cell differentiation. *Journal of bacteriology* 194: 6282–6291.
24. Khajuria RK, Munschauer M, Ulirsch JC, Fiorini C, Ludwig LS, et al. (2018) Ribosome levels selectively regulate translation and lineage commitment in human hematopoiesis. *Cell* 173: 90–103.
25. Shi Z, Fujii K, Kovary KM, Genuth NR, Röst HL, et al. (2017) Heterogeneous ribosomes preferentially translate distinct subpools of mRNAs genome-wide. *Mol Cell* 67: 71–83.e7.
26. Cagnetta R, Frese CK, Shigeoka T, Krijgsveld J, Holt CE (2018) Rapid Cue-Specific remodeling of the nascent axonal proteome. *Neuron* 99: 29–46.e4.
27. Shigeoka T, Koppers M, Wong HHW, Lin JQ, Cagnetta R, et al. (2019) On-Site ribosome remodeling by locally synthesized ribosomal proteins in axons. *Cell Rep* 29: 3605–3619.e10.
28. Steffen KK, MacKay VL, Kerr EO, Tsuchiya M, Hu D, et al. (2008) Yeast life span extension by depletion of 60S ribosomal subunits is mediated by gcn4. *Cell* 133: 292–302.
29. Lawrence MS, Stojanov P, Mermel CH, Robinson JT, Garraway LA, et al. (2014) Discovery and saturation analysis of cancer genes across 21 tumour types. *Nature* 505: 495–501.

30. Xue S, Barna M (2012) Specialized ribosomes: a new frontier in gene regulation and organismal biology. *Nature Reviews Molecular Cell Biology* 13: 355–369.
31. Gerbasi VR, Weaver CM, Hill S, Friedman DB, Link AJ (2004) Yeast Asc1p and mammalian RACK1 are functionally orthologous core 40S ribosomal proteins that repress gene expression. *Molecular and cellular biology* 24: 8276–8287.
32. Thompson MK, Rojas-Duran MF, Gangaramani P, Gilbert WV (2016) The ribosomal protein asc1/rack1 is required for efficient translation of short mrnas. *Elife* 5: e11154.
33. Johnson AG, Lapointe CP, Wang J, Corsepius NC, Choi J, et al. (2019) Rack1 on and off the ribosome. *RNA* 25: 881–895.
34. Draptchinskaia N, Gustavsson P, Andersson B, Pettersson M, Willig TN, et al. (1999) The gene encoding ribosomal protein s19 is mutated in diamond-blackfan anaemia. *Nature genetics* 21: 169–175.
35. Liu JM, Ellis SR (2006) Ribosomes and marrow failure: coincidental association or molecular paradigm? *Blood* 107: 4583–4588.
36. De Keersmaecker K, Sulima SO, Dinman JD (2015) Ribosomopathies and the paradox of cellular hypo-to hyperproliferation. *Blood* 125: 1377–1382.
37. Sulima SO, Hofman IJ, De Keersmaecker K, Dinman JD (2017) How ribosomes translate cancer. *Cancer discovery* 7: 1069–1087.
38. Kampen KR, Sulima SO, De Keersmaecker K (2018) Rise of the specialized onco-ribosomes. *Oncotarget* 9: 35205.
39. Yewdell JW, Antón LC, Bennink JR (1996) Defective ribosomal products (drips): a major source of antigenic peptides for mhc class i molecules? *The Journal of Immunology* 157: 1823–1826.
40. Yewdell JW, Nichitta CV (2006) The drip hypothesis decennial: support, controversy, refinement and extension. *Trends in immunology* 27: 368–373.
41. Wei J, Yewdell JW (2019) Immunoribosomes: Wheres theres fire, theres fire. *Molecular immunology* 113: 38–42.
42. Warner JR (1999) The economics of ribosome biosynthesis in yeast. *Trends in biochemical sciences* 24: 437–440.
43. Sloan KE, Warda AS, Sharma S, Entian KD, Lafontaine DL, et al. (2017) Tuning the ribosome: The influence of rrna modification on eukaryotic ribosome biogenesis and function. *RNA biology* 14: 1138–1152.
44. Penzo M, Galbiati A, Treré D, Montanaro L (2016) The importance of being (slightly) modified: The role of rrna editing on gene expression control and its connections with cancer. *Biochimica et Biophysica Acta (BBA)-Reviews on Cancer* 1866: 330–338.

45. Parks MM, Kurylo CM, Dass RA, Bojmar L, Lyden D, et al. (2018) Variant ribosomal rna alleles are conserved and exhibit tissue-specific expression. *Science advances* 4: eaao0665.
46. Wong SY, Javid B, Addepalli B, Piszczek G, Strader MB, et al. (2013) Functional role of methylation of g518 of the 16s rna 530 loop by gidb in mycobacterium tuberculosis. *Antimicrobial agents and chemotherapy* 57: 6311–6318.
47. Mauro VP, Matsuda D (2016) Translation regulation by ribosomes: Increased complexity and expanded scope. *RNA biology* 13: 748–755.
48. Battle A, Khan Z, Wang SH, Mitrano A, Ford MJ, et al. (2015) Impact of regulatory variation from RNA to protein. *Science* 347: 664–667.
49. Franks A, Airoidi E, Slavov N (2017) Post-transcriptional regulation across human tissues. *PLoS computational biology* 13: e1005535.
50. Kondrashov N, Pusic A, Stumpf CR, Shimizu K, Hsieh AC, et al. (2011) Ribosome-mediated specificity in Hox mRNA translation and vertebrate tissue patterning. *Cell* 145: 383–397.
51. Slavov N, Botstein D (2011) Coupling among growth rate response, metabolic cycle, and cell division cycle in yeast. *Molecular Biology of the Cell* 22: 1997 – 2009. doi:10.1091/mbc.E11-02-0132.
52. Sung MK, Reitsma JM, Sweredoski MJ, Hess S, Deshaies RJ (2016) Ribosomal proteins produced in excess are degraded by the ubiquitin–proteasome system. *Molecular biology of the cell* 27: 2642–2652.
53. Eisenberg AR, Higdon A, Keskin A, Hodapp S, Jovanovic M, et al. (2018) Precise post-translational tuning occurs for most protein complex components during meiosis. *Cell Rep* 25: 3603–3617.e2.
54. Zhang B, Wang J, Wang X, Zhu J, Liu Q, et al. (2014) Proteogenomic characterization of human colon and rectal cancer. *Nature* 513: 382–387.
55. Slavov N, Budnik B, Schwab D, Airoidi E, van Oudenaarden A (2014) Constant Growth Rate Can Be Supported by Decreasing Energy Flux and Increasing Aerobic Glycolysis. *Cell Reports* 7: 705 – 714. doi:10.1016/j.celrep.2014.03.057.
56. Janssens GE, Meinema AC, Gonzalez J, Wolters JC, Schmidt A, et al. (2015) Protein biogenesis machinery is a driver of replicative aging in yeast. *Elife* 4: e08527.
57. Kelmer Sacramento E, Kirkpatrick JM, Mazzetto M, Baumgart M, Bartolome A, et al. (2020) Reduced proteasome activity in the aging brain results in ribosome stoichiometry loss and aggregation. *Mol Syst Biol* 16: e9596.
58. Jovanovic M, Rooney MS, Mertins P, Przybylski D, Chevrier N, et al. (2015) Dynamic profiling of the protein life cycle in response to pathogens. *Science* 347.

59. Riba A, Di Nanni N, Mittal N, Arhné E, Schmidt A, et al. (2019) Protein synthesis rates and ribosome occupancies reveal determinants of translation elongation rates. *Proceedings of the National Academy of Sciences* 116: 15023–15032.
60. Warner JR, McIntosh KB (2009) How common are extraribosomal functions of ribosomal proteins? *Molecular cell* 34: 3–11.
61. Statello L, Maugeri M, Garre E, Nawaz M, Wahlgren J, et al. (2018) Identification of rna-binding proteins in exosomes capable of interacting with different types of rna: Rbp-facilitated transport of rnas into exosomes. *PloS one* 13.
62. Weisberg RA (2008) Transcription by moonlight: structural basis of an extraribosomal activity of ribosomal protein s10. *Molecular cell* 32: 747–748.
63. Bhavsar RB, Makley LN, Tsonis PA (2010) The other lives of ribosomal proteins. *Human genomics* 4: 327.
64. Zhou X, Liao WJ, Liao JM, Liao P, Lu H (2015) Ribosomal proteins: functions beyond the ribosome. *Journal of molecular cell biology* 7: 92–104.
65. Dougherty JD (2017) The expanding toolkit of translating ribosome affinity purification. *Journal of Neuroscience* 37: 12079–12087.
66. Shigeoka T, Koppers M, Wong HHW, Lin JQ, Cagnetta R, et al. (2019) On-site ribosome remodeling by locally synthesized ribosomal proteins in axons. *Cell reports* 29: 3605–3619.
67. Hummel M, Cordewener JH, de Groot JC, Smeekens S, America AH, et al. (2012) Dynamic protein composition of a rabdopsis thaliana cytosolic ribosomes in response to sucrose feeding as revealed by label free ms e proteomics. *Proteomics* 12: 1024–1038.
68. Amirbeigiab S, Kiani P, Sanchez AV, Krisp C, Kazantsev A, et al. (2019) Invariable stoichiometry of ribosomal proteins in mouse brain tissues with aging. *Proceedings of the National Academy of Sciences* 116: 22567–22572.
69. Ghaemmaghami S, Huh WK, Bower K, Howson RW, Belle A, et al. (2003) Global analysis of protein expression in yeast. *Nature* 425: 737–741.
70. Newman JRS, Ghaemmaghami S, Ihmels J, Breslow DK, Noble M, et al. (2006) Single-cell proteomic analysis of *s. cerevisiae* reveals the architecture of biological noise. *Nature* 441: 840–846.
71. Ferretti MB, Ghalei H, Ward EA, Potts EL, Karbstein K (2017) Rps26 directs mrna-specific translation by recognition of kozak sequence elements. *Nature structural & molecular biology* 24: 700.

72. Budnik B, Levy E, Harmange G, Slavov N (2017) Mass-spectrometry of single mammalian cells quantifies proteome heterogeneity during cell differentiation. *bioRxiv* 1: DOI: 10.1101/102681.
73. Budnik B, Levy E, Harmange G, Slavov N (2018) SCoPE-MS: mass-spectrometry of single mammalian cells quantifies proteome heterogeneity during cell differentiation. *Genome Biology* 19: 161. doi:10.1186/s13059-018-1547-5.
74. Slavov N, Macinkas J, Caudy A, Botstein D (2011) Metabolic cycling without cell division cycling in respiring yeast. *Proceedings of the National Academy of Sciences* 108: 19090 – 19095. doi:10.1073/pnas.1116998108.
75. Slavov N, Botstein D, Caudy A (2014) Extensive regulation of metabolism and growth during the cell division cycle. *bioRxiv* doi:10.1101/005629.
76. Slavov N, Airoidi EM, van Oudenaarden A, Botstein D (2012) A conserved cell growth cycle can account for the environmental stress responses of divergent eukaryotes. *Molecular Biology of the Cell* 23: 1986 – 1997. doi:10.1091/mbc.E11-11-0961.
77. Slavov N (2020) Unpicking the proteome in single cells. *Science* 367: 512–513. doi: 10.1126/science.aaz6695.
78. Slavov N (2020) Single-cell protein analysis by mass spectrometry. *Current Opinion in Chemical Biology* 60: 1 – 9. doi:10.1016/j.cbpa.2020.04.018.
79. Specht H, Emmott E, Petelski AA, Gray Huffman R, Perlman DH, et al. (2019) Single-cell mass-spectrometry quantifies the emergence of macrophage heterogeneity. *bioRxiv* doi:10.1101/665307.
80. Specht H, Slavov N (2018) Transformative opportunities for single-cell proteomics. *Journal of Proteome Research* 17: 2563 – 2916. doi:10.1021/acs.jproteome.8b00257.
81. Catherman AD, Skinner OS, Kelleher NL (2014) Top down proteomics: facts and perspectives. *Biochemical and biophysical research communications* 445: 683–693.
82. Toby TK, Fornelli L, Kelleher NL (2016) Progress in top-down proteomics and the analysis of proteoforms. *Annual review of analytical chemistry* 9: 499–519.
83. Lott BB, Wang Y, Nakazato T (2013) A comparative study of ribosomal proteins: linkage between amino acid distribution and ribosomal assembly. *BMC biophysics* 6: 13.
84. Giansanti P, Tsiatsiani L, Low TY, Heck AJ (2016) Six alternative proteases for mass spectrometry–based proteomics beyond trypsin. *Nature protocols* 11: 993–1006.
85. Morsa D, Baiwir D, La Rocca R, Zimmerman TA, Hanozin E, et al. (2019) Multi-enzymatic limited digestion: The next-generation sequencing for proteomics? *Journal of Proteome Research* 18: 2501–2513. doi:10.1021/acs.jproteome.9b00044. PMID: 31046285.

86. Peng M, Taouatas N, Cappadona S, Van Breukelen B, Mohammed S, et al. (2012) Protease bias in absolute protein quantitation. *Nature methods* 9: 524.
87. Malioutov D, Chen T, Airoidi E, Jaffe J, Budnik B, et al. (2019) Quantifying homologous proteins and proteoforms. *Molecular & Cellular Proteomics* 18: 162–168. doi:10.1074/mcp.TIR118.000947.
88. Giansanti P, Aye TT, van den Toorn H, Peng M, van Breukelen B, et al. (2015) An augmented Multiple-Protease-Based human phosphopeptide atlas. *Cell Rep* 11: 1834–1843.
89. Emmott E, Jovanovic M, Slavov N (2019) Approaches for studying ribosome specialization. *Trends in Biochemical Sciences* 44: 478 – 479. doi:10.1016/j.tibs.2019.01.008.
90. Hughes CS, Foehr S, Garfield DA, Furlong EE, Steinmetz LM, et al. (2014) Ultrasensitive proteome analysis using paramagnetic bead technology. *Molecular systems biology* 10: 757.
91. Specht H, Harmange G, Perlman DH, Emmott E, Niziolek Z, et al. (2018) Automated sample preparation for high-throughput single-cell proteomics. *bioRxiv* doi:10.1101/399774.
92. Sielaff M, Kuharev J, Bohn T, Hahlbrock J, Bopp T, et al. (2017) Evaluation of fasp, sp3, and ist protocols for proteomic sample preparation in the low microgram range. *Journal of proteome research* 16: 4060–4072.
93. Cravatt BF, Simon GM, Yates Iii JR (2007) The biological impact of mass-spectrometry-based proteomics. *Nature* 450: 991.
94. Aebersold R, Mann M (2016) Mass-spectrometric exploration of proteome structure and function. *Nature* 537: 347–355. doi:10.1038/nature19949.
95. Venable JD, Dong MQ, Wohlschlegel J, Dillin A, Yates JR (2004) Automated approach for quantitative analysis of complex peptide mixtures from tandem mass spectra. *Nat Methods* 1: 39–45.
96. Demichev V, Messner CB, Vernardis SI, Lilley KS, Ralser M (2020) DIA-NN: neural networks and interference correction enable deep proteome coverage in high throughput. *Nat Methods* 17: 41–44.
97. Muntel J, Kirkpatrick J, Bruderer R, Huang T, Vitek O, et al. (2019) Comparison of protein quantification in a complex background by DIA and TMT workflows with fixed instrument time. *J Proteome Res* 18: 1340–1351.
98. Ong SE, Blagoev B, Kratchmarova I, Kristensen DB, Steen H, et al. (2002) Stable isotope labeling by amino acids in cell culture, silac, as a simple and accurate approach to expression proteomics. *Molecular & cellular proteomics* 1: 376–386.

99. Blagoev B, Ong SE, Kratchmarova I, Mann M (2004) Temporal analysis of phosphotyrosine-dependent signaling networks by quantitative proteomics. *Nature biotechnology* 22: 1139–1145.
100. Berger SJ, Lee SW, Anderson GA, Paša-Tolić L, Tolić N, et al. (2002) High-throughput global peptide proteomic analysis by combining stable isotope amino acid labeling and data-dependent multiplexed-ms/ms. *Analytical chemistry* 74: 4994–5000.
101. Hsu JL, Huang SY, Chow NH, Chen SH (2003) Stable-isotope dimethyl labeling for quantitative proteomics. *Analytical chemistry* 75: 6843–6852.
102. Boersema PJ, Raijmakers R, Lemeer S, Mohammed S, Heck AJR (2009) Multiplex peptide stable isotope dimethyl labeling for quantitative proteomics. *Nat Protoc* 4: 484–494.
103. DeSouza LV, Romaschin AD, Colgan TJ, Siu KM (2009) Absolute quantification of potential cancer markers in clinical tissue homogenates using multiple reaction monitoring on a hybrid triple quadrupole/linear ion trap tandem mass spectrometer. *Analytical chemistry* 81: 3462–3470.
104. Kang UB, Yeom J, Kim H, Lee C (2010) Quantitative analysis of mtraq-labeled proteome using full ms scans. *Journal of Proteome research* 9: 3750–3758.
105. Webb-Robertson BJM, Wiberg HK, Matzke MM, Brown JN, Wang J, et al. (2015) Review, evaluation, and discussion of the challenges of missing value imputation for mass spectrometry-based label-free global proteomics. *Journal of proteome research* 14: 1993–2001.
106. Cox J, Hein MY, Lubner CA, Paron I, Nagaraj N, et al. (2014) Accurate proteome-wide label-free quantification by delayed normalization and maximal peptide ratio extraction, termed maxlq. *Molecular & cellular proteomics* 13: 2513–2526.
107. Chen AT, Franks A, Slavov N (2019) Dart-id increases single-cell proteome coverage. *PLOS Computational Biology* 15: 1–30. doi:10.1371/journal.pcbi.1007082.
108. Yu SH, Kiriakidou P, Cox J (2020) Isobaric matching between runs and novel psm-level normalization in maxquant strongly improve reporter ion-based quantification. *bioRxiv* .
109. Crotty F, Blackshaw R, Murray PJ (2011) Differential growth of the fungus *absidia cylindrospora* on ¹³c/¹⁵n-labelled media. *Rapid Communications in Mass Spectrometry* 25: 1479–1484.
110. Filiou MD, Varadarajulu J, Teplytska L, Reckow S, Maccarrone G, et al. (2012) The ¹⁵n isotope effect in *escherichia coli*: A neutron can make the difference. *Proteomics* 12: 3121–3128.
111. Frank E, Kessler MS, Filiou MD, Zhang Y, Maccarrone G, et al. (2009) Stable isotope metabolic labeling with a novel ¹⁵n-enriched bacteria diet for improved proteomic analyses of mouse models for psychopathologies. *PloS one* 4.

112. Filiou MD, Webhofer C, Gormanns P, Zhang Y, Reckow S, et al. (2012) The 15 n isotope effect as a means for correlating phenotypic alterations and affected pathways in a trait anxiety mouse model. *Proteomics* 12: 2421–2427.
113. Van Hoof D, Pinkse MW, Ward-Van Oostwaard D, Mummery CL, Heck AJ, et al. (2007) An experimental correction for arginine-to-proline conversion artifacts in silac-based quantitative proteomics. *Nature methods* 4: 677–678.
114. Boersema PJ, Raijmakers R, Lemeer S, Mohammed S, Heck AJ (2009) Multiplex peptide stable isotope dimethyl labeling for quantitative proteomics. *Nature protocols* 4: 484.
115. Lau HT, Suh HW, Golkowski M, Ong SE (2014) Comparing silac-and stable isotope dimethyl-labeling approaches for quantitative proteomics. *Journal of proteome research* 13: 4164–4174.
116. Merrill AE, Hebert AS, MacGilvray ME, Rose CM, Bailey DJ, et al. (2014) Neucode labels for relative protein quantification. *Molecular & Cellular Proteomics* 13: 2503–2512.
117. Overmyer KA, Tyanova S, Hebert AS, Westphall MS, Cox J, et al. (2018) Multiplexed proteome analysis with neutron-encoded stable isotope labeling in cells and mice. *Nature protocols* 13: 293–306.
118. Thompson A, Schäfer J, Kuhn K, Kienle S, Schwarz J, et al. (2003) Tandem mass tags: a novel quantification strategy for comparative analysis of complex protein mixtures by ms/ms. *Analytical chemistry* 75: 1895–1904.
119. Ross PL, Huang YN, Marchese JN, Williamson B, Parker K, et al. (2004) Multiplexed protein quantitation in *saccharomyces cerevisiae* using amine-reactive isobaric tagging reagents. *Molecular & cellular proteomics* 3: 1154–1169.
120. Choe L, D’Ascenzo M, Relkin NR, Pappin D, Ross P, et al. (2007) 8-plex quantitation of changes in cerebrospinal fluid protein expression in subjects undergoing intravenous immunoglobulin treatment for alzheimer’s disease. *Proteomics* 7: 3651–3660.
121. Huang T, Bruderer R, Muntel J, Xuan Y, Vitek O, et al. (2020) Combining precursor and fragment information for improved detection of differential abundance in data independent acquisition. *Mol Cell Proteomics* 19: 421–430.
122. Huffman G, Chen AT, Specht H, Slavov N (2019) DO-MS: Data-driven optimization of mass spectrometry methods. *Journal of Proteome Research* 18: 2493 – 2500. doi:10.1021/acs.jproteome.9b00039.
123. Ting L, Rad R, Gygi SP, Haas W (2011) Ms3 eliminates ratio distortion in isobaric multiplexed quantitative proteomics. *Nature methods* 8: 937–940.
124. Sonnett M, Yeung E, Wuhr M (2018) Accurate, sensitive, and precise multiplexed proteomics using the complement reporter ion cluster. *Analytical chemistry* 90: 5032–5039.

125. Chait BT (2006) Mass spectrometry: bottom-up or top-down? *Science* 314: 65–66.
126. van den Heuvel RH, van Duijn E, Mazon H, Synowsky SA, Lorenzen K, et al. (2006) Improving the performance of a quadrupole time-of-flight instrument for macromolecular mass spectrometry. *Analytical chemistry* 78: 7473–7483.
127. van de Waterbeemd M, Fort KL, Boll D, Reinhardt-Szyba M, Routh A, et al. (2017) High-fidelity mass analysis unveils heterogeneity in intact ribosomal particles. *Nat Methods* 14: 283–286.
128. Heck AJ (2008) Native mass spectrometry: a bridge between interactomics and structural biology. *Nature methods* 5: 927–933.
129. Belov AM, Viner R, Santos MR, Horn DM, Bern M, et al. (2017) Analysis of proteins, protein complexes, and organellar proteomes using sheathless capillary zone electrophoresis - native mass spectrometry. *J Am Soc Mass Spectrom* 28: 2614–2634.
130. Kafader JO, Melani RD, Durbin KR, Ikwuagwu B, Early BP, et al. (2020) Multiplexed mass spectrometry of individual ions improves measurement of proteoforms and their complexes. *Nature Methods* : 1–4.
131. Wörner TP, Snijder J, Bennett A, Agbandje-McKenna M, Makarov AA, et al. (2020) Resolving heterogeneous macromolecular assemblies by orbitrap-based single-particle charge detection mass spectrometry. *Nature Methods* : 1–4.
132. Donnelly DP, Rawlins CM, DeHart CJ, Fornelli L, Schachner LF, et al. (2019) Best practices and benchmarks for intact protein analysis for top-down mass spectrometry. *Nat Methods* 16: 587–594.
133. Van De Waterbeemd M, Tamara S, Fort KL, Damoc E, Franc V, et al. (2018) Dissecting ribosomal particles throughout the kingdoms of life using advanced hybrid mass spectrometry methods. *Nature communications* 9: 1–12.
134. Sinz A (2018) Cross-linking/mass spectrometry for studying protein structures and protein–protein interactions: where are we now and where should we go from here? *Angewandte Chemie International Edition* 57: 6390–6396.
135. Yu C, Huang L (2018) Cross-linking mass spectrometry: an emerging technology for interactomics and structural biology. *Analytical chemistry* 90: 144–165.
136. Bui KH, von Appen A, DiGuilio AL, Ori A, Sparks L, et al. (2013) Integrated structural analysis of the human nuclear pore complex scaffold. *Cell* 155: 1233–1243.
137. Poliakov A, Duijn Ev, Lander G, Fu Cy, Johnson JE, et al. (2007) Macromolecular mass spectrometry and electron microscopy as complementary tools for investigation of the heterogeneity of bacteriophage portal assemblies. *J Struct Biol* 157: 371–383. doi: 10.1016/j.jsb.2006.09.003.

138. Lasker K, Förster F, Bohn S, Walzthoeni T, Villa E, et al. (2012) Molecular architecture of the 26S proteasome holocomplex determined by an integrative approach. *Proc Natl Acad Sci USA* 109: 1380–1387. doi:10.1073/pnas.1120559109.
139. Noble AJ, Zhang Q, O'donnell J, Hariri H, Bhattacharya N, et al. (2013) A pseudoatomic model of the copii cage obtained from cryo-electron microscopy and mass spectrometry. *Nature structural & molecular biology* 20: 167.
140. Davis JH, Tan YZ, Carragher B, Potter CS, Lyumkis D, et al. (2016) Modular assembly of the bacterial large ribosomal subunit. *Cell* 167: 1610–1622.
141. Wu S, Tan D, Woolford Jr JL, Dong MQ, Gao N (2017) Atomic modeling of the its2 ribosome assembly subcomplex from cryo-em together with mass spectrometry-identified protein–protein crosslinks. *Protein Science* 26: 103–112.
142. Nogales E, Scheres SH (2015) Cryo-em: a unique tool for the visualization of macromolecular complexity. *Molecular cell* 58: 677–689.
143. Rozov A, Khusainov I, El Omari K, Duman R, Mykhaylyk V, et al. (2019) Importance of potassium ions for ribosome structure and function revealed by long-wavelength x-ray diffraction. *Nature communications* 10: 1–12.
144. Watson ZL, Ward FR, Méheust R, Ad O, Schepartz A, et al. (2020) Structure of the bacterial ribosome at 2 Å resolution. *bioRxiv* .
145. Mulder AM, Yoshioka C, Beck AH, Bunner AE, Milligan RA, et al. (2010) Visualizing ribosome biogenesis: parallel assembly pathways for the 30S subunit. *Science* 330: 673–677.
146. Sashital DG, Greeman CA, Lyumkis D, Potter CS, Carragher B, et al. (2014) A combined quantitative mass spectrometry and electron microscopy analysis of ribosomal 30s subunit assembly in *e. coli*. *Elife* 3: e04491.
147. Feaga HA, Kopylov M, Kim JK, Jovanovic M, Dworkin J (2020) Ribosome dimerization protects the small subunit. *Journal of bacteriology* 202.
148. Lilleorg S, Reier K, Pulk A, Liiv A, Tammsalu T, et al. (2019) Bacterial ribosome heterogeneity: Changes in ribosomal protein composition during transition into stationary growth phase. *Biochimie* 156: 169–180.