

# Starlikeness of Certain Non-Univalent Functions

Adam Lecko, V. Ravichandran, and Asha Sebastian

*Dedicated to Prof. Dato' Indera Rosihan M. Ali*

**ABSTRACT.** We consider three classes of functions defined using the class  $\mathcal{P}$  of all analytic functions  $p(z) = 1 + cz + \dots$  on the open unit disk having positive real part and study several radius problems for these classes. The first class consists of all normalized analytic functions  $f$  with  $f/g \in \mathcal{P}$  and  $g/(zp) \in \mathcal{P}$  for some normalized analytic function  $g$  and  $p \in \mathcal{P}$ . The second class is defined by replacing the condition  $f/g \in \mathcal{P}$  by  $|(f/g) - 1| < 1$  while the other class consists of normalized analytic functions  $f$  with  $f/(zp) \in \mathcal{P}$  for some  $p \in \mathcal{P}$ . We have determined radii so that the functions in these classes to belong to various subclasses of starlike functions. These subclasses includes the classes of starlike functions of order  $\alpha$ , parabolic starlike functions, as well as the classes of starlike functions associated with lemniscate of Bernoulli, reverse lemniscate, sine function, a rational function, cardioid, lune, nephroid and modified sigmoid function.

## 1. Introduction

Let  $\mathbb{D}$  denote the open unit disc in  $\mathbb{C}$ . Let  $\mathcal{A} := \{f : f \text{ is analytic on } \mathbb{D}, f(0) = 0, \text{ and } f'(0) = 1\}$ . Let  $\mathcal{S} := \{f \in \mathcal{A} : f \text{ is univalent on } \mathbb{D}\}$ . An analytic function  $f$  is subordinate to another analytic function  $g$ , written  $f \prec g$ , if  $f(z) = g(w(z))$  for some analytic function  $w : \mathbb{D} \rightarrow \mathbb{D}$  that fixes the origin. If  $g \in \mathcal{S}$ , then  $f \prec g$  if and only if the functions  $f$  and  $g$  takes the origin to the same point as well as the range of  $f$  is a subset of the range of  $g$ :  $f(\mathbb{D}) \subseteq g(\mathbb{D})$ . Several well known subclasses of starlike and convex functions were characterized by subordination of  $zf'(z)/f(z)$  or  $1 + (zf''(z)/f'(z))$  to some function in  $\mathcal{P}$ . For a univalent  $\varphi$  in unit disc  $\mathbb{D}$  with  $\operatorname{Re} \varphi(z) > 0$ , starlike with respect to  $\varphi(0) = 1$ , symmetric about the real axis and  $\varphi'(0) > 0$ , Ma and Minda [22] gave a unified treatment for functions in the class  $\mathcal{S}^*(\varphi) = \{f \in \mathcal{A} : zf'(z)/f(z) \prec \varphi(z)\}$  and  $\mathcal{K}(\varphi) = \{f \in \mathcal{A} : 1 + (zf''(z)/f'(z)) \prec \varphi(z)\}$ . Convolution theorems for these two classes in a more general setting was previously studied by Shanmugham [35] with the stronger assumption that  $\varphi$  is convex. Several authors considered these classes for various choices of the function  $\varphi$ . For  $-1 \leq B < A \leq 1$ , let  $\varphi$  be the bilinear transform that maps the unit disc  $\mathbb{D}$  onto the disc whose diametric end points are  $(1+A)/(1+B)$  and  $(1-A)/(1-B)$ ; if we impose  $\varphi(0) = 1$ , then this mapping is given by  $\varphi(z) = (1 + Az)/(1 + Bz)$ . For

---

1991 *Mathematics Subject Classification.* 30C80, 30C45.

*Key words and phrases.* Univalent functions; convex functions; starlike functions; subordination; radius of starlikeness.

Asha Sebastian is supported by an institute fellowship from NIT Tiruchirappalli.

this function  $\varphi$ , the classes  $\mathcal{S}^*(\varphi)$  and  $\mathcal{K}(\varphi)$  reduce respectively to the classes  $\mathcal{S}^*[A, B]$  and  $\mathcal{K}[A, B]$  of Janowski starlike and convex functions. Other well-known choices for  $\varphi(z)$  include  $\sqrt{1+z}$ ,  $e^z$ ,  $1 + \sin z$ ,  $z + \sqrt{1+z^2}$ . Readers may refer to [11, 39] for brief survey of these classes.

The class  $\mathcal{P}$  of analytic functions  $p(z) = 1 + cz + \dots$  on  $\mathbb{D}$  having positive real part is known as the class of Carathéodory functions or the class of functions with positive real part. For  $0 \leq \alpha < 1$ , let  $\mathcal{P}(\alpha) := \{p \in \mathcal{P} : \operatorname{Re} p(z) > \alpha\}$ . For any two subclasses  $\mathcal{F}$  and  $\mathcal{G}$  of  $\mathcal{A}$ , the  $\mathcal{G}$ -radius for the class  $\mathcal{F}$ , denoted by  $R_{\mathcal{G}}(\mathcal{F})$ , is the largest number  $R_{\mathcal{G}} \in (0, 1)$  such that  $r^{-1}f(rz) \in \mathcal{G}$ , for all  $f \in \mathcal{F}$  and for  $0 < r < R_{\mathcal{G}}$ . Among the several studies available on radius problems, a major focus has been on ratio between two classes of functions, where one of them belong to some particular subclasses of  $\mathcal{A}$  and can be seen in [23, 24, 25, 31, 30]. Let  $\Phi$  be a function defined on  $\mathbb{D}$ . For  $\Phi(z) = 1, 1/(1+z), 1/(1-z)^2, 1/(1-z^2), 1+(z/2)$ , various authors have considered radius problems for classes of functions  $f$  satisfying the following conditions: (i)  $\operatorname{Re} f(z)/g(z) > 0$  and  $\operatorname{Re} g(z)/(z\Phi(z)) > 0$ , (ii)  $|(f(z)/g(z)) - 1| < 1$  and  $\operatorname{Re} g(z)/(z\Phi(z)) > 0$ , (iii)  $\operatorname{Re} f(z)/(z\Phi(z)) > 0$ . See [2, 33, 14, 15, 7]. Also, in 2019, Lecko and Sim [20] considered two functions  $\Phi(z) = 1/(1-z)^2, 1/(1-z^2)$  where  $z\Phi(z)$  are starlike functions with integer coefficients and studied on  $\operatorname{Re} f(z)/(z\Phi(z)) > 0$ . Similar studies can be seen in [16, 17, 18].

Motivated by the aforestated works, we define three classes of functions by making use of the class  $\mathcal{P}$  as follows:

$$\mathcal{G}_1 := \left\{ f \in \mathcal{A} : \frac{f}{g} \in \mathcal{P}, \frac{g}{zp} \in \mathcal{P} \text{ for some } g \in \mathcal{A}, p \in \mathcal{P} \right\}, \quad (1.1)$$

$$\mathcal{G}_2 := \left\{ f \in \mathcal{A} : \frac{f}{g} \in \mathcal{P}, \frac{g}{zp} \in \mathcal{P}(1/2) \text{ for some } g \in \mathcal{A}, p \in \mathcal{P} \right\}, \quad (1.2)$$

and

$$\mathcal{G}_3 := \left\{ f \in \mathcal{A} : \frac{f}{zp} \in \mathcal{P} \text{ for some } p \in \mathcal{P} \right\}. \quad (1.3)$$

We determine radii for functions in the classes  $\mathcal{G}_1, \mathcal{G}_2, \mathcal{G}_3$  to belong to several subclasses of  $\mathcal{A}$  like starlike functions of order  $\alpha$ , starlike functions associated with lemniscate of Bernoulli, reverse lemniscate, sine function, exponential function, cardioid, lune, nephroid, a particular rational function, modified sigmoid function and parabolic starlike functions. The disc that contains the image of unit disc  $\mathbb{D}$  under the mapping  $zf'(z)/f(z)$  aids in determining the radius of various classes and we discuss this mapping in the following section.

## 2. Mapping of $zf'(z)/f(z)$ for $\mathcal{G}_1, \mathcal{G}_2, \mathcal{G}_3$

In this section, we show that the classes  $\mathcal{G}_1, \mathcal{G}_2, \mathcal{G}_3$  are non-empty and contains non-univalent functions. We also determine the disk containing the image of the disc  $\mathbb{D}$  under the function  $zf'/f$  when  $f$  belong to each of the classes. We shall use the function  $p_0 : \mathbb{D} \rightarrow \mathbb{C}$  defined by

$$p_0(z) = \frac{1+z}{1-z}. \quad (2.1)$$

This function  $p_0$  belongs to  $\mathcal{P}$  and maps the unit disc  $\mathbb{D}$  onto the right half-plane. For functions  $p \in \mathcal{P}(\alpha)$ , we shall use the following inequality [34, Lemma 2]:

$$\left| \frac{zp'(z)}{p(z)} \right| \leq \frac{2(1-\alpha)r}{(1-r)(1+(1-2\alpha)r)}, \quad |z| \leq r. \quad (2.2)$$

**2.1. The class  $\mathcal{G}_1$ .** Define the functions  $f_1, g_1 : \mathbb{D} \rightarrow \mathbb{C}$  by

$$f_1(z) = z \left( \frac{1+z}{1-z} \right)^3, \quad \text{and} \quad g_1(z) = z \left( \frac{1+z}{1-z} \right)^2. \quad (2.3)$$

The function  $f_1$  satisfies  $f_1/g_1 \in \mathcal{P}$  and  $g_1/zp_0 \in \mathcal{P}$  where  $p_0 \in \mathcal{P}$  is given in (2.1). Thus the function  $f_1 \in \mathcal{G}_1$  and the class  $\mathcal{G}_1$  is non-empty. It is an extremal function for the class  $\mathcal{G}_1$ . From the coefficients of Taylor series expansion of functions  $f_1$  given by

$$f_1(z) = z + 6z^2 + 18z^3 + 38z^4 + \dots,$$

it is evident that the functions  $f_1$  is not univalent. Hence the class  $\mathcal{G}_1$ , contain non-univalent functions. As

$$f_1'(z) = \frac{(1+z)^2(1+6z-z^2)}{(1-z)^4}$$

the function  $f_1'$  vanishes at  $z = -\rho = -(\sqrt{10} - 3)$ . From Theorem 3.1, it is apparent that the radii of univalence of the class  $\mathcal{G}_1$  is  $\sqrt{10} - 3$  and it coincides with its radius of starlikeness.

If the function  $f \in \mathcal{G}_1$ , then there exists an element  $g \in \mathcal{A}$  and  $p \in \mathcal{P}$  such that

$$\frac{f}{g} \in \mathcal{P} \quad \text{and} \quad \frac{g}{zp} \in \mathcal{P}. \quad (2.4)$$

Let  $p_1, p_2$  be two functions defined on unit disc  $\mathbb{D}$  by

$$p_1(z) = \frac{f(z)}{g(z)} \quad \text{and} \quad p_2(z) = \frac{g(z)}{zp(z)}. \quad (2.5)$$

From (2.4) and (2.5), we have  $f(z) = zp_1(z)p_2(z)p(z)$ . Then a calculation shows that

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| \leq \left| \frac{zp_1'(z)}{p_1(z)} \right| + \left| \frac{zp_2'(z)}{p_2(z)} \right| + \left| \frac{zp'(z)}{p(z)} \right|. \quad (2.6)$$

Since  $p_1, p_2, p \in \mathcal{P}$ , it follows from (2.2) and (2.6) that

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| \leq \frac{6r}{1-r^2}, \quad |z| \leq r. \quad (2.7)$$

**2.2. The class  $\mathcal{G}_2$ .** The functions  $f_2$  and  $g_2$  are defined on the unit disc  $\mathbb{D}$  as follows:

$$f_2(z) = z \frac{(1+z)^2}{(1-z)^3} \quad \text{and} \quad g_2(z) = z \frac{(1+z)}{(1-z)^2} \quad (2.8)$$

The function  $f_2$  satisfies  $f_2/g_2 \in \mathcal{P}$  and  $g_2/(zp_0) \in \mathcal{P}(1/2)$  where  $p_0 \in \mathcal{P}$  is given in (2.1). Thus the function  $f_2 \in \mathcal{G}_2$  and the class  $\mathcal{G}_2$  is non empty. It is an extremal function for the class  $\mathcal{G}_2$ . Note that the Taylor series expansion of

$$f_2(z) = z + 5z^2 + 13z^3 + 25z^4 + \dots$$

and it is clear that the function  $f_2$  is not univalent function, as it does not satisfy de Brange's theorem. Thus the class  $\mathcal{G}_2$  contain non univalent functions. Since

$$f_2'(z) = \frac{1 + 6z + 5z^2}{(1 - z)^4},$$

the function  $f_2'$  vanishes at  $z = -\zeta = -1/5$  and it follows that the radius of univalence of the class of functions  $\mathcal{G}_2$  is  $1/5$  by the Theorem 3.1. It also coincides with radius of starlikeness.

If the function  $f \in \mathcal{G}_2$ , then there exists a normalized analytic function  $g$  and  $p \in \mathcal{P}$  such that

$$\frac{f}{g} \in \mathcal{P} \quad \text{and} \quad \frac{g}{zp} \in \mathcal{P}(1/2). \quad (2.9)$$

Let  $h_1$  and  $h_2$  be two functions,  $h_1, h_2 : \mathbb{D} \rightarrow \mathbb{C}$  defined as

$$h_1(z) = \frac{f(z)}{g(z)} \quad \text{and} \quad h_2(z) = \frac{g(z)}{zp(z)}. \quad (2.10)$$

Using (2.9) and (2.10), we have  $f(z) = zh_1(z)h_2(z)p(z)$ , it can be shown that

$$\frac{zf'(z)}{f(z)} - 1 = \frac{zh_1'(z)}{h_1(z)} + \frac{zh_2'(z)}{h_2(z)} + \frac{zp'(z)}{p(z)}. \quad (2.11)$$

As  $h_1, p \in \mathcal{P}$  and  $h_2 \in \mathcal{P}(1/2)$ , it follows from (2.2) and (2.11) that

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| \leq \frac{r(r+5)}{1-r^2}, \quad |z| \leq r. \quad (2.12)$$

It would be interesting to find the boundary of the set  $\cup_{f \in \mathcal{G}_2} \{zf'(z)/f(z) : |z| \leq r\}$ . This would help in finding sharp radii to some of the problem where we have got only a lower bound.

**2.3. The class  $\mathcal{G}_3$ .** The function  $g_1$ , defined by (2.3), belongs to the class  $\mathcal{G}_3$  and therefore the class  $\mathcal{G}_3$  is non-empty. It is an extremal function for the class  $\mathcal{G}_3$ . From the coefficients of Taylor series expansion of functions  $g_1$  given by

$$g_1(z) = z + 4z^2 + 8z^3 + 12z^4 + \dots$$

it is evident that the functions  $g_1$  is not univalent. Hence the class  $\mathcal{G}_3$ , contain non-univalent functions. As

$$g_1'(z) = \frac{1 + 5z + 3z^2 - z^3}{(1 - z)^3},$$

the function  $g_1'$  vanishes at  $z = -\eta = -(\sqrt{5} - 2)$ . From Theorem 3.1, it is apparent that the radii of univalence of the class  $\mathcal{G}_3$  coincide with its radius of starlikeness. We now discuss the mapping of  $zf'(z)/f(z)$  when the function  $f \in \mathcal{G}_3$ . Let  $p \in \mathcal{P}$  and  $p_1$  be a function defined on unit disc  $\mathbb{D}$  such that

$$p_1(z) = \frac{f(z)}{zp(z)} \quad (2.13)$$

It is clear from (2.13) that  $f(z) = zp_1(z)p(z)$ . Then it follows that

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| \leq \left| \frac{zp_1'(z)}{p_1(z)} \right| + \left| \frac{zp'(z)}{p(z)} \right|. \quad (2.14)$$

Applying (2.2) in (2.14), we obtain

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| \leq \frac{4r}{1-r^2}, \quad |z| \leq r. \quad (2.15)$$

Using the equations (2.7), (2.12), (2.15), we investigate several radius problems associated with functions in the classes  $\mathcal{G}_1$ ,  $\mathcal{G}_2$  and  $\mathcal{G}_3$  in the next section.

### 3. Radius of starlikeness

In this section, we determine the radii of the classes  $\mathcal{G}_1$ ,  $\mathcal{G}_2$ ,  $\mathcal{G}_3$  to belong to various Ma-Minda starlike classes of functions. For  $0 \leq \alpha \leq 1$ , the class  $\mathcal{S}^*(\alpha) = \mathcal{S}^*[1 - 2\alpha, -1] = \{f \in \mathcal{A} : \operatorname{Re} zf'(z)/f(z) > \alpha\}$  is the class of starlike functions of order  $\alpha$ . These classes were studied extensively in [6, 11, 12, 13].

**THEOREM 3.1.** *The following sharp results hold for the class  $\mathcal{S}^*(\alpha)$ :*

- (i)  $R_{\mathcal{S}^*(\alpha)}(\mathcal{G}_1) = (1 - \alpha) / (3 + \sqrt{10 - 2\alpha + \alpha^2})$ .
- (ii)  $R_{\mathcal{S}^*(\alpha)}(\mathcal{G}_2) = 2(1 - \alpha) / (5 + \sqrt{25 - 4\alpha + \alpha^2})$ .
- (iii)  $R_{\mathcal{S}^*(\alpha)}(\mathcal{G}_3) = (1 - \alpha) / (2 + \sqrt{5 - 2\alpha + \alpha^2})$ .

**PROOF.** (i) The function defined by  $m(r) = (1 - 6r - r^2)(1 - r^2)^{-1}$ ,  $0 \leq r < 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}^*(\alpha)}(\mathcal{G}_1)$  is the root of the equation  $m(r) = \alpha$ . From (2.7), it follows that

$$\operatorname{Re} \frac{zf'(z)}{f(z)} \geq \frac{1 - 6r - r^2}{1 - r^2} = m(r) \geq m(\rho) = \alpha.$$

This shows that  $R_{\mathcal{S}^*(\alpha)}(\mathcal{G}_1)$  is at least  $\rho$ . At  $z = -R_{\mathcal{S}^*(\alpha)}(\mathcal{G}_1) = -\rho$ , the function  $f_1$  defined in (2.3) satisfies

$$\operatorname{Re} \frac{zf_1'(z)}{f_1(z)} = \frac{1 - 6\rho - \rho^2}{1 - \rho^2} = \alpha.$$

Thus the radius is sharp.

- (ii) The function  $n(r) : [0, 1) \rightarrow \mathbb{R}$  defined by  $n(r) = (1 - 5r)(1 - r^2)^{-1}$  is a decreasing function. Let  $\rho = R_{\mathcal{S}^*(\alpha)}(\mathcal{G}_2)$  is the root of the equation  $n(r) = \alpha$ . Let  $f \in \mathcal{G}_2$  and let  $h_1, p \in \mathcal{P}$  and  $h_2 \in \mathcal{P}(1/2)$  be the functions defined in Section 2.2. For  $h \in \mathcal{P}$ , by [4, Lemma 2.3], we have

$$\operatorname{Re} \frac{zh'(z)}{h(z)} \geq -\frac{2r}{1 - r^2}, \quad |z| \leq r. \quad (3.1)$$

For  $k \in \mathcal{P}(1/2)$ , by [4, Lemma 2.4], we have

$$\operatorname{Re} \frac{zk'(z)}{k(z)} \geq -\frac{r}{1 + r}, \quad |z| \leq r. \quad (3.2)$$

Using (3.1) and (3.2) in (2.11), it follows that

$$\operatorname{Re} \frac{zf'(z)}{f(z)} \geq 1 - \frac{4r}{1 - r^2} - \frac{r}{1 + r} = \frac{1 - 5r}{1 - r^2} = n(r) \geq n(\rho) = \alpha.$$

This shows that  $R_{\mathcal{S}^*(\alpha)}(\mathcal{G}_2)$  is at least  $\rho$ . At  $z = -R_{\mathcal{S}^*(\alpha)}(\mathcal{G}_1) = -\rho$ , the function  $f_2$  defined in (2.8) satisfies

$$\operatorname{Re} \frac{zf_2'(z)}{f_2(z)} = \frac{1 - 5\rho}{1 - \rho^2} = \alpha.$$

Thus the radius is sharp.

- (iii) The function defined by  $s(r) = (1 - 4r - r^2)(1 - r^2)^{-1}$ ,  $0 \leq r < 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}^*(\alpha)}(\mathcal{G}_3)$  is the root of the equation  $s(r) = \alpha$ . From (2.15), it follows that

$$\operatorname{Re} \frac{zf'(z)}{f(z)} \geq \frac{1 - 4r - r^2}{1 - r^2} = s(r) \geq s(\rho) = \alpha.$$

This shows that  $R_{\mathcal{S}^*(\alpha)}(\mathcal{G}_3)$  is at least  $\rho$ . For the function  $g_1$  defined in (2.3), at  $z = -R_{\mathcal{S}^*(\alpha)}(\mathcal{G}_2) = -\rho$ ,

$$\operatorname{Re} \frac{zg_1'(z)}{g_1(z)} = \frac{1 - 4\rho - \rho^2}{1 - \rho^2} = \alpha.$$

Thus the radius is sharp. ■

The class  $\mathcal{S}_L^* = \mathcal{S}^*(\sqrt{1+z})$  and it represents the collection of functions in the class  $\mathcal{A}$  whose  $zf'(z)/f(z)$  lies in the region bounded by the lemniscate of Bernoulli  $|w^2 - 1| = 1$ . Various studies on  $\mathcal{S}_L^*$  can be seen in [38, 3, 28]. Ali *et al.* [1, Lemma 2.2] had proved that for  $2\sqrt{2}/3 < a < \sqrt{2}$ ,

$$\{w : |w - a| < \sqrt{2} - a\} \subset \{w : |w^2 - 1| < 1\}. \quad (3.3)$$

Using this lemma, we obtain radii results for the classes  $\mathcal{G}_1$ ,  $\mathcal{G}_2$ ,  $\mathcal{G}_3$  to be in the class  $\mathcal{S}_L^*$  in the following theorem.

**THEOREM 3.2.** *The following results for the class  $\mathcal{S}_L^*$  are sharp.*

- (i)  $R_{\mathcal{S}_L^*}(\mathcal{G}_1) = (3 - 2\sqrt{2})/(\sqrt{2} - 1) \left(3 + \sqrt{12 - 2\sqrt{2}}\right) \approx 0.0687$ .
- (ii)  $R_{\mathcal{S}_L^*}(\mathcal{G}_2) = 2(\sqrt{2} - 1)/\left(5 + \sqrt{33 - 4\sqrt{2}}\right) \approx 0.0809$ .
- (iii)  $R_{\mathcal{S}_L^*}(\mathcal{G}_3) = (3 - 2\sqrt{2})/(\sqrt{2} - 1) \left(2 + \sqrt{7 - 2\sqrt{2}}\right) \approx 0.1025$ .

**PROOF.** (i) The function defined by  $m(r) = 6r(1 - r^2)^{-1} + 1$ ,  $0 \leq r < 1$  is an increasing function. Let  $\rho = R_{\mathcal{S}_L^*}(\mathcal{G}_1)$  is the root of the equation  $m(r) = \sqrt{2}$ . For  $0 < r \leq R_{\mathcal{S}_L^*}(\mathcal{G}_1)$ , we have  $m(r) \leq \sqrt{2}$ . That is,

$$\frac{6r}{1 - r^2} + 1 \leq \sqrt{2} = m(\rho).$$

For the class  $\mathcal{G}_1$ , the centre of the disc is 1, therefore the disc obtained in (2.7) is contained in the region bounded by lemniscate, by Lemma 3.3. For the function  $f_1$  defined in (2.3), at  $z = R_{\mathcal{S}_L^*}(\mathcal{G}_1) = \rho$ ,

$$\left| \left( \frac{zf_1'(z)}{f_1(z)} \right)^2 - 1 \right| = \left| \left( \frac{1 + 6\rho - \rho^2}{1 - \rho^2} \right)^2 - 1 \right| = 1.$$

- (ii) The function  $n(r) : [0, 1) \rightarrow \mathbb{R}$  defined by  $n(r) = (5r + r^2)(1 - r^2)^{-1} + 1$  is an increasing function. Let  $\rho = R_{\mathcal{S}_L^*}(\mathcal{G}_2)$  is the root of the equation  $n(r) = \sqrt{2}$ . For  $0 < r \leq R_{\mathcal{S}_L^*}(\mathcal{G}_2)$ , we have  $n(r) \leq \sqrt{2}$ . That is,

$$\frac{5r + r^2}{1 - r^2} + 1 \leq \sqrt{2} = n(\rho).$$

For the class  $\mathcal{G}_2$ , the centre of the disc is 1, therefore the disc obtained in (2.12) is contained in the region bounded by lemniscate, by Lemma 3.3. For the function  $f_2$  defined in (2.8), at  $z = R_{\mathcal{S}_L^*}(\mathcal{G}_2) = \rho$ ,

$$\left| \left( \frac{zf'(z)}{f(z)} \right)^2 - 1 \right| = \left| \left( \frac{1 + 5\rho}{1 - \rho^2} \right)^2 - 1 \right| = 1.$$

- (iii) The function defined by  $s(r) = 4r(1 - r^2)^{-1} + 1$ ,  $0 \leq r < 1$  is an increasing function. Let  $\rho = R_{\mathcal{S}_L^*}(\mathcal{G}_3)$  is the root of the equation  $s(r) = \sqrt{2}$ . For  $0 < r \leq R_{\mathcal{S}_L^*}(\mathcal{G}_3)$ , we have  $s(r) \leq \sqrt{2}$ . That is,

$$\frac{4r}{1 - r^2} + 1 \leq \sqrt{2} = s(\rho).$$

For the class  $\mathcal{G}_3$ , the centre of the disc is 1, therefore the disc obtained in (2.15) is contained in the region bounded by lemniscate, by Lemma 3.3. For the function  $g_1$  defined in (2.3), at  $z = R_{\mathcal{S}_L^*}(\mathcal{G}_3) = \rho$ ,

$$\left| \left( \frac{zf'(z)}{f(z)} \right)^2 - 1 \right| = \left| \left( \frac{1 + 4\rho - \rho^2}{1 - \rho^2} \right)^2 - 1 \right| = |(\sqrt{2})^2 - 1| = 1. \quad \blacksquare$$

Let  $\varphi_{PAR}(z) := 1 + \left( 2/\pi^2 (\log(1 + \sqrt{z})/(1 - \sqrt{z}))^2 \right)$ . Since  $\varphi_{PAR}(\mathbb{D}) = \{w : \operatorname{Re} w > |w - 1|\}$  is a parabolic region, the functions in the class  $\mathcal{S}_p := \mathcal{S}^*(\varphi_{PAR})$  are known as parabolic starlike functions. These functions are studied by authors in [9, 21, 32]. Shanmugam and Ravichandran [36, pp.321] had proved that for  $1/2 < a < 3/2$ , then

$$\{w : |w - a| < a - 1/2\} \subset \{w : \operatorname{Re} w > |w - 1|\}. \quad (3.4)$$

The following theorem gives the radius of parabolic starlikeness of the three classes  $\mathcal{G}_1$ ,  $\mathcal{G}_2$  and  $\mathcal{G}_3$ .

**THEOREM 3.3.** *The following results hold for the class  $\mathcal{S}_p$ :*

- (i)  $R_{\mathcal{S}_p}(\mathcal{G}_1) = \sqrt{37} - 6 \approx 0.0827$ .
- (ii)  $R_{\mathcal{S}_p}(\mathcal{G}_2) \geq (2\sqrt{7} - 5)/3 \approx 0.0972$ .
- (iii)  $R_{\mathcal{S}_p}(\mathcal{G}_3) = \sqrt{17} - 4 \approx 0.1231$ .

**PROOF.** (i) The function defined by  $m(r) = (1 - 6r - r^2)(1 - r^2)^{-1}$ ,  $0 \leq r < 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_p}(\mathcal{G}_1)$  is the root of the equation  $m(r) = 1/2$ . For  $0 < r \leq R_{\mathcal{S}_p}(\mathcal{G}_1)$ , we have  $m(r) \geq 1/2$ . That is,

$$\frac{6r}{1 - r^2} \leq \frac{1}{2} = m(\rho).$$

For the class  $\mathcal{G}_1$ , the centre of the disc is 1, therefore the disc obtained in (2.7) is contained in the region bounded by parabola, by Lemma 3.4. For the function  $f_1$  defined in (2.3), at  $z = R_{\mathcal{S}_p}(\mathcal{G}_1) = \rho$ ,

$$\operatorname{Re} \frac{zf_1'(z)}{f_1(z)} = \frac{1 + 6\rho - \rho^2}{1 - \rho^2} = \frac{1}{2} = \left| \frac{zf_1'(z)}{f_1(z)} - 1 \right|.$$

- (ii) The function  $n(r) : [0, 1) \rightarrow \mathbb{R}$  defined by  $n(r) = (1 - 5r - 2r^2)(1 - r^2)^{-1} + 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_p}(\mathcal{G}_2)$  is the root of the equation  $n(r) = 1/2$ . For  $0 < r \leq R_{\mathcal{S}_p}(\mathcal{G}_2)$ , we have  $n(r) \geq 1/2$ . That is,

$$\frac{r(r+5)}{1-r^2} \leq \frac{1}{2} = n(\rho).$$

For the class  $\mathcal{G}_2$ , the centre of the disc is 1, therefore the disc obtained in (2.12) is contained in the region bounded by parabola, by Lemma 3.4. This shows that  $R_{\mathcal{S}_p}(\mathcal{G}_2)$  is at least  $\rho$ .

- (iii) The function defined by  $s(r) = 1 - (4r(1 - r^2)^{-1})$ ,  $0 \leq r < 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_p}(\mathcal{G}_3)$  is the root of the equation  $s(r) = 1/2$ . For  $0 < r \leq R_{\mathcal{S}_p}(\mathcal{G}_3)$ , we have  $s(r) \geq 1/2$ . That is,

$$\frac{4r}{1-r^2} \leq \frac{1}{2} = s(\rho).$$

For the class  $\mathcal{G}_3$ , the centre of the disc is 1, therefore the disc obtained in (2.15) is contained in the region bounded by parabola, by Lemma 3.4. For the function  $g_1$  defined in (2.3), at  $z = -R_{\mathcal{S}_p}(\mathcal{G}_3) = -\rho$ ,

$$\operatorname{Re} \frac{zg_1'(z)}{g_1(z)} = \frac{1 - 4\rho - \rho^2}{1 - \rho^2} = \frac{1}{2} = \left| \frac{zg_1'(z)}{g_1(z)} - 1 \right|. \quad \blacksquare$$

In 2015, Mendiratta *et al.* [27] introduced the class of starlike functions associated with the exponential function as  $\mathcal{S}_e^* = \mathcal{S}^*(e^z)$  and it satisfies the condition  $|\log zf'(z)/f(z)| < 1$ . They had also proved that, for  $e^{-1} \leq a \leq (e + e^{-1})/2$ ,

$$\{w \in \mathbb{C} : |w - a| < a - e^{-1}\} \subseteq \{w \in \mathbb{C} : |\log w| < 1\}. \quad (3.5)$$

**THEOREM 3.4.** *The following results hold for the class  $\mathcal{S}_e^*$ :*

- (i)  $R_{\mathcal{S}_e^*}(\mathcal{G}_1) = (e - 1) / (3e + \sqrt{10e^2 - 2e + 1}) \approx 0.1042$ .
- (ii)  $R_{\mathcal{S}_e^*}(\mathcal{G}_2) \geq 2(e^2 + e - 2) / (2 + e) (5e + \sqrt{8 + 4e + 29e^2}) \approx 0.1213$ .
- (iii)  $R_{\mathcal{S}_e^*}(\mathcal{G}_3) = 2(e - 1)^2 / (e + 1) (4e + \sqrt{4 - 8e + 20e^2}) \approx 0.1543$ .

**PROOF.** (i) The function defined by  $m(r) = (1 - 6r - r^2)(1 - r^2)^{-1}$ ,  $0 \leq r < 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_e^*}(\mathcal{G}_1)$  is the root of the equation  $m(r) = 1/e$ . For  $0 < r \leq R_{\mathcal{S}_e^*}(\mathcal{G}_1)$ , we have  $m(r) \geq 1/e$ . That is,

$$\frac{6r}{1-r^2} \leq 1 - \frac{1}{e}.$$

For the class  $\mathcal{G}_1$ , the centre of the disc is 1, therefore the disc obtained in (2.7) is contained in the region bounded by exponential function, by Lemma 3.5. For the function  $f_1$  defined in (2.3), at  $z = R_{\mathcal{S}_e^*}(\mathcal{G}_1) = \rho$ ,

$$\left| \log \frac{zf_1'(z)}{f_1(z)} \right| = \left| \log \frac{1 + 6\rho - \rho^2}{1 - \rho^2} \right| = 1.$$

- (ii) The function  $n(r) : [0, 1) \rightarrow \mathbb{R}$  defined by  $n(r) = (1 - 5r - 2r^2)(1 - r^2)^{-1} + 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_e^*}(\mathcal{G}_2)$  is the root of the equation  $n(r) = 1/e$ . For  $0 < r \leq R_{\mathcal{S}_e^*}(\mathcal{G}_2)$ , we have  $n(r) \geq 1/e$ . That is,

$$\frac{r(r+5)}{1-r^2} \leq 1 - \frac{1}{e}.$$

For the class  $\mathcal{G}_2$ , the centre of the disc is 1, therefore the disc obtained in (2.12) is contained in the region bounded by the exponential function, by Lemma 3.5. This shows that  $R_{\mathcal{S}_e^*}(\mathcal{G}_2)$  is at least  $\rho$ .

- (iii) The function defined by  $s(r) = 1 - (4r(1 - r^2)^{-1})$ ,  $0 \leq r < 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_e^*}(\mathcal{G}_3)$  is the root of the equation  $s(r) = 1/e$ . For  $0 < r \leq R_{\mathcal{S}_e^*}(\mathcal{G}_3)$ , we have  $s(r) \geq 1/e$ . That is,

$$\frac{4r}{1-r^2} \leq \frac{e-1}{e}.$$

For the class  $\mathcal{G}_3$ , the centre of the disc is 1, therefore the disc obtained in (2.15) is contained in the region bounded by the exponential function, by Lemma 3.5. For the function  $g_1$  defined in (2.3), at  $z = R_{\mathcal{S}_e^*}(\mathcal{G}_3) = \rho$ ,

$$\left| \log \frac{zg_1'(z)}{g_1(z)} \right| = \left| \log \frac{1+4\rho-\rho^2}{1-\rho^2} \right| = 1. \quad \blacksquare$$

Theorem 3.5 provides radii results for starlike functions associated with a cardioid. Sharma *et al.* [37] studied various properties of the class  $\mathcal{S}_c^* = \mathcal{S}^*(1 + (4/3)z + (2/3)z^2)$ . Geometrically, if a function  $f \in \mathcal{S}_c^*$  then  $zf'(z)/f(z)$  lies in the region bounded by the cardioid  $\Omega_c = \{u + iv : (9u^2 + 9v^2 - 18u + 5)^2 - 16(9u^2 + 9v^2 - 6u + 1) = 0\}$ . They had also proved that, for  $1/3 < a \leq 5/3$ ,

$$\{w \in \mathbb{C} : |w - a| < (3a - 1)/3\} \subseteq \Omega_c. \quad (3.6)$$

**THEOREM 3.5.** *The following results hold for the class  $\mathcal{S}_c^*$ :*

- (i)  $R_{\mathcal{S}_c^*}(\mathcal{G}_1) = (\sqrt{85} - 9)/2 \approx 0.1097$ .
- (ii)  $R_{\mathcal{S}_c^*}(\mathcal{G}_2) \geq (\sqrt{265} - 15)/10 \approx 0.1279$ .
- (iii)  $R_{\mathcal{S}_c^*}(\mathcal{G}_3) = \sqrt{10} - 3 \approx 0.1623$ .

**PROOF.** (i) The function defined by  $m(r) = (1 - 6r - r^2)(1 - r^2)^{-1}$ ,  $0 \leq r < 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_c^*}(\mathcal{G}_1)$  is the root of the equation  $m(r) = 1/3$ . For  $0 < r \leq R_{\mathcal{S}_c^*}(\mathcal{G}_1)$ , we have  $m(r) \geq 1/3$ . That is,

$$\frac{6r}{1-r^2} \leq 1 - \frac{1}{3}.$$

For the class  $\mathcal{G}_1$ , the centre of the disc is 1, therefore the disc obtained in (2.7) is contained in the region bounded by the cardioid, by Lemma 3.6. For the function  $f_1$  defined in (2.3), at  $z = R_{\mathcal{S}_c^*}(\mathcal{G}_1) = \rho$ ,

$$\left| \frac{zf_1'(z)}{f_1(z)} \right| = \left| \frac{1+6\rho-\rho^2}{1-\rho^2} \right| = \frac{1}{3} = \Omega_c(-1).$$

- (ii) The function  $n(r) : [0, 1) \rightarrow \mathbb{R}$  defined by  $n(r) = (1 - 5r - 2r^2)(1 - r^2)^{-1} + 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_c^*}(\mathcal{G}_2)$  is the root of the equation  $n(r) = 1/3$ . For  $0 < r \leq R_{\mathcal{S}_c^*}(\mathcal{G}_2)$ , we have  $n(r) \geq 1/3$ . That is,

$$\frac{r(r+5)}{1-r^2} \leq 1 - \frac{1}{3}.$$

For the class  $\mathcal{G}_2$ , the centre of the disc is 1, therefore the disc obtained in (2.12) is contained in the region bounded by the cardioid, by Lemma 3.6. This shows that  $R_{\mathcal{S}_c^*}(\mathcal{G}_2)$  is at least  $\rho$ .

- (iii) The function defined by  $s(r) = 1 - (4r(1 - r^2)^{-1})$ ,  $0 \leq r < 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_c^*}(\mathcal{G}_3)$  is the root of the equation  $s(r) = 1/3$ . For  $0 < r \leq R_{\mathcal{S}_c^*}(\mathcal{G}_3)$ , we have  $s(r) \geq 1/3$ . That is,

$$\frac{4r}{1-r^2} \leq \frac{2}{3}.$$

For the class  $\mathcal{G}_3$ , the centre of the disc is 1, therefore the disc obtained in (2.15) is contained in the region bounded by the cardioid, by Lemma 3.6. For the function  $g_1$  defined in (2.3), at  $z = R_{\mathcal{S}_c^*}(\mathcal{G}_3) = \rho$ ,

$$\left| \frac{zf_1'(z)}{f_1(z)} \right| = \left| \frac{1 + 4\rho - \rho^2}{1 - \rho^2} \right| = \frac{1}{3} = \Omega_c(-1). \quad \blacksquare$$

In 2019, Cho *et al.* [5] considered the class of starlike functions associated with sine function where the class  $\mathcal{S}_{sin}^*$  is defined as  $\mathcal{S}_{sin}^* = \{f \in \mathcal{A} : zf'(z)/f(z) \prec 1 + \sin z := q_0(z)\}$  for  $z \in \mathbb{D}$ . For  $|a - 1| \leq \sin 1$ , they had established the following inclusion:

$$\{w \in \mathbb{C} : |w - a| < \sin 1 - |a - 1|\} \subseteq \Omega_s. \quad (3.7)$$

Here  $\Omega_s := q_0(\mathbb{D})$  is the image of the unit disk  $\mathbb{D}$  under the mappings  $q_0(z) = 1 + \sin z$ .

**THEOREM 3.6.** *The following results are sharp for the class  $\mathcal{S}_{sin}^*$ .*

- (i)  $R_{\mathcal{S}_{sin}^*}(\mathcal{G}_1) = \sin 1 / \left(3 + \sqrt{9 + \sin^2 1}\right) \approx 0.1375$ .  
(ii)  $R_{\mathcal{S}_{sin}^*}(\mathcal{G}_2) = 2 \sin 1 / \left(5 + \sqrt{25 + 4 \sin 1 + 4 \sin^2 1}\right) \approx 0.1589$ .  
(iii)  $R_{\mathcal{S}_{sin}^*}(\mathcal{G}_3) = \sin 1 / \left(2 + \sqrt{4 + \sin^2 1}\right) \approx 0.2018$ .

**PROOF.** (i) The function defined by  $m(r) = (1 - 6r - r^2)(1 - r^2)^{-1}$ ,  $0 \leq r < 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_{sin}^*}(\mathcal{G}_1)$  is the root of the equation  $m(r) = 1 - \sin 1$ . For  $0 < r \leq R_{\mathcal{S}_{sin}^*}(\mathcal{G}_1)$ , we have  $m(r) \geq 1 - \sin 1$ . That is,

$$\frac{6r}{1-r^2} \leq \sin 1.$$

For the class  $\mathcal{G}_1$ , the centre of the disc is 1, therefore the disc obtained in (2.7) is contained in the region  $\Omega_s$  bounded by the sine function, by Lemma 3.7. For the function  $f_1$  defined in (2.3), at  $z = -R_{\mathcal{S}_{sin}^*}(\mathcal{G}_1) = -\rho$ ,

$$\left| \frac{zf_1'(z)}{f_1(z)} \right| = \left| \frac{1 - 6\rho - \rho^2}{1 - \rho^2} \right| = 1 + \sin 1 = q_0(1).$$

- (ii) The function  $n(r) : [0, 1) \rightarrow \mathbb{R}$  defined by  $n(r) = (1 - 5r - 2r^2)(1 - r^2)^{-1} + 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_{sin}^*}(\mathcal{G}_2)$  is the root of the equation  $n(r) = 1 - \sin 1$ . For  $0 < r \leq R_{\mathcal{S}_{sin}^*}(\mathcal{G}_2)$ , we have  $n(r) \geq 1 - \sin 1$ . That is,

$$\frac{r(r+5)}{1-r^2} \leq \sin 1.$$

For the class  $\mathcal{G}_2$ , the centre of the disc is 1, therefore the disc obtained in (2.12) is contained in the region bounded by the sine function, by Lemma 3.7. For the function  $f_2$  defined in (2.8), at  $z = R_{\mathcal{S}_{sin}^*}(\mathcal{G}_1) = \rho$ ,

$$\left| \frac{zf_2'(z)}{f_2(z)} \right| = \left| \frac{1+5\rho}{1-\rho^2} \right| = 1 + \sin 1 = q_0(1).$$

- (iii) The function defined by  $s(r) = 1 - (4r(1 - r^2)^{-1})$ ,  $0 \leq r < 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_{sin}^*}(\mathcal{G}_3)$  is the root of the equation  $s(r) = 1 - \sin 1$ . For  $0 < r \leq R_{\mathcal{S}_{sin}^*}(\mathcal{G}_3)$ , we have  $s(r) \geq 1 - \sin 1$ . That is,

$$\frac{4r}{1-r^2} \leq \sin 1.$$

For the class  $\mathcal{G}_3$ , the centre of the disc is 1, therefore the disc obtained in (2.15) is contained in the region bounded by the sine function, by Lemma 3.7. For the function  $g_1$  defined in (2.3), at  $z = -R_{\mathcal{S}_{sin}^*}(\mathcal{G}_3) = -\rho$ ,

$$\left| \frac{zf_1'(z)}{f_1(z)} \right| = \left| \frac{1-4\rho-\rho^2}{1-\rho^2} \right| = 1 + \sin 1 = q_0(1). \quad \blacksquare$$

In 2015, Raina and Sokól[29] introduced the class  $\mathcal{S}_{\mathcal{C}}^* = \mathcal{S}^*(z + \sqrt{1+z^2})$ . They showed that a function  $f \in \mathcal{S}_{\mathcal{C}}^*$  if and only if  $zf'(z)/f(z)$  belongs to a lune shaped region  $\mathcal{L} := \{w \in \mathbb{C} : |w^2 - 1| < 2|w|\}$ . Gandhi and Ravichandran [8, Lemma 2.1] proved that

$$\{w \in \mathbb{C} : |w - a| < 1 - |\sqrt{2} - a|\} \subseteq \{w \in \mathbb{C} : |w^2 - 1| < 2|w|\}. \quad (3.8)$$

**THEOREM 3.7.** *The following results hold for the class  $\mathcal{S}_{\mathcal{C}}^*$ :*

- (i)  $R_{\mathcal{S}_{\mathcal{C}}^*}(\mathcal{G}_1) = (6 - 4\sqrt{2})/(2 - \sqrt{2}) \left(3 + \sqrt{15 - 4\sqrt{2}}\right) \approx 0.0967$ .
- (ii)  $R_{\mathcal{S}_{\mathcal{C}}^*}(\mathcal{G}_2) \geq (16 - 10\sqrt{2})/(3 - \sqrt{2}) \left(5 + \sqrt{57 - 20\sqrt{2}}\right) \approx 0.1131$ .
- (iii)  $R_{\mathcal{S}_{\mathcal{C}}^*}(\mathcal{G}_3) = (6 - 4\sqrt{2})/(2 - \sqrt{2}) \left(2 + \sqrt{10 - 4\sqrt{2}}\right) \approx 0.1434$ .

**PROOF.** (i) The function defined by  $m(r) = (1 - 6r - r^2)(1 - r^2)^{-1}$ ,  $0 \leq r < 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_{\mathcal{C}}^*}(\mathcal{G}_1)$  is the root of the equation  $m(r) = \sqrt{2} - 1$ . For  $0 < r \leq R_{\mathcal{S}_{\mathcal{C}}^*}(\mathcal{G}_1)$ , we have  $m(r) \geq \sqrt{2} - 1$ . That is,

$$\frac{6r}{1-r^2} \leq 2 - \sqrt{2}.$$

For the class  $\mathcal{G}_1$ , the centre of the disc is 1, therefore the disc obtained in (2.7) is contained in the region bounded by the intersection of disks  $\{w : |w - 1| < \sqrt{2}\}$

and  $\{w : |w + 1| < \sqrt{2}\}$ , by Lemma 3.8. For the function  $f_1$  defined in (2.3), at  $z = -R_{\mathcal{S}_{\mathbb{C}}^*}(\mathcal{G}_1) = -\rho$ ,

$$\left| \left( \frac{zf_1'(z)}{f_1(z)} \right)^2 - 1 \right| = \left| \left( \frac{1 - 6\rho - \rho^2}{1 - \rho^2} \right)^2 - 1 \right| = 2 \left| \frac{1 - 6\rho - \rho^2}{1 - \rho^2} \right|.$$

- (ii) The function  $n(r) : [0, 1) \rightarrow \mathbb{R}$  defined by  $n(r) = (1 - 5r - 2r^2)(1 - r^2)^{-1} + 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_{\mathbb{C}}^*}(\mathcal{G}_2)$  is the root of the equation  $n(r) = \sqrt{2} - 1$ . For  $0 < r \leq R_{\mathcal{S}_{\mathbb{C}}^*}(\mathcal{G}_2)$ , we have  $n(r) \geq \sqrt{2} - 1$ . That is,

$$\frac{r(r + 5)}{1 - r^2} \leq 2 - \sqrt{2}.$$

For the class  $\mathcal{G}_2$ , the centre of the disc is 1, therefore the disc obtained in (2.12) is contained in the region bounded by the lune, by Lemma 3.8. This shows that  $R_{\mathcal{S}_{\mathbb{C}}^*}(\mathcal{G}_2)$  is at least  $\rho$ .

- (iii) The function  $s(r) : [0, 1) \rightarrow \mathbb{R}$  defined by  $s(r) = 1 - (4r(1 - r^2)^{-1})$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_{\mathbb{C}}^*}(\mathcal{G}_1)$  is the root of the equation  $s(r) = \sqrt{2} - 1$ . For  $0 < r \leq R_{\mathcal{S}_{\mathbb{C}}^*}(\mathcal{G}_3)$ , we have  $s(r) \geq \sqrt{2} - 1$ . That is,

$$\frac{4r}{1 - r^2} \leq 2 - \sqrt{2}.$$

For the class  $\mathcal{G}_3$ , the centre of the disc is 1, therefore the disc obtained in (2.15) is contained in the region bounded by the lune, by Lemma 3.8. For the function  $g_1$  defined in (2.3), at  $z = -R_{\mathcal{S}_{\mathbb{C}}^*}(\mathcal{G}_3) = -\rho$ ,

$$\left| \left( \frac{zg_1'(z)}{g_1(z)} \right)^2 - 1 \right| = \left| \left( \frac{1 - 4\rho - \rho^2}{1 - \rho^2} \right)^2 - 1 \right| = 2 \left| \frac{1 - 4\rho - \rho^2}{1 - \rho^2} \right|. \quad \blacksquare$$

In the next theorem, we provide radii for starlike functions associated with a rational function. Kumar and Ravichandran [19] introduced the class of starlike functions associated with a rational function,  $\psi(z) = 1 + (z^2k + z^2/(k^2 - kz))$  where  $k = \sqrt{2} + 1$ , defined by  $\mathcal{S}_R^* = \mathcal{S}^*(\psi(z))$ . For  $2(\sqrt{2} - 1) < a \leq \sqrt{2}$ , they had proved that

$$\{w \in \mathbb{C} : |w - a| < a - 2(\sqrt{2} - 1)\} \subseteq \psi(\mathbb{D}). \quad (3.9)$$

**THEOREM 3.8.** *The following results hold for the class  $\mathcal{S}_R^*$ :*

- (i)  $R_{\mathcal{S}_R^*}(\mathcal{G}_1) = (3 - 2\sqrt{2}) / \left( 3 + \sqrt{26 - 12\sqrt{2}} \right) \approx 0.0285$ .
- (ii)  $R_{\mathcal{S}_R^*}(\mathcal{G}_2) \geq (20 - 14\sqrt{2}) / (2 - \sqrt{2}) \left( 5 + \sqrt{105 - 56\sqrt{2}} \right) \approx 0.0340$ .
- (iii)  $R_{\mathcal{S}_R^*}(\mathcal{G}_3) = (3 - 2\sqrt{2}) / \left( 2 + \sqrt{21 - 12\sqrt{2}} \right) \approx 0.0428$ .

**PROOF.** (i) The function defined by  $m(r) = (1 - 6r - r^2)(1 - r^2)^{-1}$ ,  $0 \leq r < 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_R^*}(\mathcal{G}_1)$  is the root of the equation  $m(r) = 2(\sqrt{2} - 1)$ .

For  $0 < r \leq R_{\mathcal{S}_R^*}(\mathcal{G}_1)$ , we have  $m(r) \geq 2(\sqrt{2} - 1)$ . That is,

$$\frac{6r}{1-r^2} \leq 1 - 2(\sqrt{2} - 1).$$

For the class  $\mathcal{G}_1$ , the centre of the disc is 1, therefore the disc obtained in (2.7) is contained in the region bounded by the rational function, by Lemma 3.9. For the function  $f_1$  defined in (2.3), at  $z = -R_{\mathcal{S}_R^*}(\mathcal{G}_1) = -\rho$ ,

$$\left| \frac{zf_1'(z)}{f_1(z)} \right| = \left| \frac{1 - 6\rho - \rho^2}{1 - \rho^2} \right| = 2(\sqrt{2} - 1) = \psi(1).$$

- (ii) The function  $n(r) : [0, 1) \rightarrow \mathbb{R}$  defined by  $n(r) = (1 - 5r - 2r^2)(1 - r^2)^{-1} + 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_R^*}(\mathcal{G}_2)$  is the root of the equation  $n(r) = 2(\sqrt{2} - 1)$ . The function defined by

$$n(r) = \frac{1 - 5r - 2r^2}{1 - r^2} + 1$$

is a decreasing function. For  $0 < r \leq R_{\mathcal{S}_R^*}(\mathcal{G}_2)$ , we have  $n(r) \geq 2(\sqrt{2} - 1)$ . That is,

$$\frac{r(r+5)}{1-r^2} \leq 3 - 2\sqrt{2}.$$

For the class  $\mathcal{G}_2$ , the centre of the disc is 1, therefore the disc obtained in (2.12) is contained in the region bounded by the rational function, by Lemma 3.9. This shows that  $R_{\mathcal{S}_R^*}(\mathcal{G}_2)$  is at least  $\rho$ .

- (iii) The function defined by  $s(r) = 1 - (4r(1 - r^2)^{-1})$ ,  $0 \leq r < 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_R^*}(\mathcal{G}_3)$  is the root of the equation  $s(r) = 2(\sqrt{2} - 1)$ . For  $0 < r \leq R_{\mathcal{S}_R^*}(\mathcal{G}_3)$ , we have  $s(r) \geq 2(\sqrt{2} - 1)$ . That is,

$$\frac{4r}{1-r^2} \leq 2 - \sqrt{2}.$$

For the class  $\mathcal{G}_3$ , the centre of the disc is 1, therefore the disc obtained in (2.15) is contained in the region bounded by the rational function, by Lemma 3.9. For the function  $g_1$  defined in (2.3), at  $z = -R_{\mathcal{S}_R^*}(\mathcal{G}_3) = -\rho$ ,

$$\left| \frac{zg_1'(z)}{g_1(z)} \right| = \left| \frac{1 - 4\rho - \rho^2}{1 - \rho^2} \right| = 2(\sqrt{2} - 1) = \psi(1). \quad \blacksquare$$

Mendiratta *et al.* [26] studied the subclass of starlike function associated with left half of shifted lemniscate of Bernoulli, given by  $|(w - \sqrt{2})^2 - 1| < 1$ . The class  $\mathcal{S}_{RL}^*$  is defined as

$$\mathcal{S}_{RL}^* = \mathcal{S}^* \left( \sqrt{2} - (\sqrt{2} - 1) \sqrt{\frac{1-z}{1+2(\sqrt{2}-1)z}} \right).$$

For  $\sqrt{2}/3 \leq a < \sqrt{2}$ , they had proved the following inclusion:

$$\{w \in \mathbb{C} : |w - a| < r_{RL}\} \subseteq \{w \in \mathbb{C} : |(w - \sqrt{2})^2 - 1| < 1\}, \quad (3.10)$$

where  $r_{RL} = \left( \left( 1 - (\sqrt{2} - a)^2 \right)^{1/2} - \left( 1 - (\sqrt{2} - a)^2 \right) \right)^{1/2}$ . Using this result, we obtain  $\mathcal{S}_{RL}^*$ -radii of the classes  $\mathcal{G}_1$ ,  $\mathcal{G}_2$ ,  $\mathcal{G}_3$  in the following theorem.

**THEOREM 3.9.** *Let  $\eta = \sqrt{2(\sqrt{2} - 1)} - 2(\sqrt{2} - 1)$ . Then the following sharp results hold for the class  $\mathcal{S}_{RL}^*$ .*

- (i)  $R_{\mathcal{S}_{RL}^*}(\mathcal{G}_1)$  is the smallest positive root ( $\approx 0.0475$ ) in  $(0, 1)$  of the equation  $(36 + 2\eta)r^2 - \eta r^2 - \eta = 0$ .
- (ii)  $R_{\mathcal{S}_{RL}^*}(\mathcal{G}_2)$  is the smallest positive root ( $\approx 0.0567$ ) in  $(0, 1)$  of the equation  $(1 - \eta)r^4 + 10r^3 + (25 - 2\eta)r^2 - \eta = 0$ .
- (iii)  $R_{\mathcal{S}_{RL}^*}(\mathcal{G}_3)$  is the smallest positive root ( $\approx 0.0711$ ) in  $(0, 1)$  of the equation  $\eta r^4 - (16 + 2\eta)r^2 + \eta = 0$ .

**PROOF.** (i) The function defined by  $m(r) = (6r(1 - r^2)^{-1}) + 1$ ,  $0 \leq r < 1$  is an increasing function. Let  $\rho = R_{\mathcal{S}_{RL}^*}(\mathcal{G}_3)$  is the root of the equation  $m(r) = 1 + \sqrt{\eta}$ . For  $0 < r \leq R_{\mathcal{S}_{RL}^*}(\mathcal{G}_1)$ , we have  $m(r) \leq \sqrt{2}$ . That is,

$$\left(\frac{6r}{1 - r^2}\right)^2 \leq \eta = (m(\rho) - 1)^2.$$

For the class  $\mathcal{G}_1$ , the centre of the disc is 1, therefore the disc obtained in (2.7) is contained in the region bounded by the reverse lemniscate, by Lemma 3.10. This shows that  $R_{\mathcal{S}_{RL}^*}(\mathcal{G}_1)$  is at least  $\rho$ . For the function  $f_1$  defined in (2.3), the radius is sharp.

- (ii) The function defined by  $n(r) = (5r + r^2)(1 - r^2)^{-1} + 1$ ,  $0 \leq r < 1$  is an increasing function. Let  $R_{\mathcal{S}_{RL}^*}(\mathcal{G}_2)$  is the root of the equation  $n(r) = 1 + \sqrt{\eta}$ . For  $0 < r \leq R_{\mathcal{S}_{RL}^*}(\mathcal{G}_2)$ , we have  $n(r) \leq \sqrt{2}$ . That is,

$$\left(\frac{5r + r^2}{1 - r^2}\right)^2 \leq \eta = (n(\rho) - 1)^2.$$

For the class  $\mathcal{G}_2$ , the centre of the disc is 1, therefore the disc obtained in (2.12) is contained in the region bounded by reverse lemniscate, by Lemma 3.10. This shows that  $R_{\mathcal{S}_{RL}^*}(\mathcal{G}_2)$  is at least  $\rho$ . The obtained radius is sharp for the function  $f_2$  defined in (2.8).

- (iii) The function defined by  $s(r) = 4r(1 - r^2)^{-1} + 1$ ,  $0 \leq r < 1$  is an increasing function. Let  $R_{\mathcal{S}_{RL}^*}(\mathcal{G}_3)$  is the root of the equation  $s(r) = 1 + \sqrt{\eta}$ . For  $0 < r \leq R_{\mathcal{S}_{RL}^*}(\mathcal{G}_3)$ , we have  $s(r) \leq \sqrt{2}$ . That is,

$$\left(\frac{4r}{1 - r^2}\right)^2 \leq \eta = (s(\rho) - 1)^2.$$

For the class  $\mathcal{G}_3$ , the centre of the disc is 1, therefore the disc obtained in (2.15) is contained in the region bounded by reverse lemniscate, by Lemma 3.10. This shows that  $R_{\mathcal{S}_{RL}^*}(\mathcal{G}_3)$  is at least  $\rho$ . The obtained radius is sharp for the function  $g_1$  defined in (2.3).

The sharpness of the results can be shown using the software *Wolfram Mathematica*. ■

In 2020, Wani and Swaminathan [40, Lemma 2.2] introduced the class  $\mathcal{S}_{Ne}^* = \mathcal{S}^*(1 + z - (z^3/3))$  that maps the open unit disc  $\mathbb{D}$  onto the interior of a two cusped kidney shaped curve  $\Omega_{Ne} := \{u + iv : ((u - 1)^2 + v^2 - 4/9)^3 - 4v^2/3 < 0\}$ . For  $1/3 < a \leq 1$ , they had

proved that

$$\{w \in \mathbb{C} : |w - a| < a - 1/3\} \subseteq \Omega_{Ne}. \quad (3.11)$$

Our next theorem determines the  $\mathcal{S}_{Ne}^*$ -radii results for the classes  $\mathcal{G}_1$ ,  $\mathcal{G}_2$  and  $\mathcal{G}_3$ .

**THEOREM 3.10.** *The following sharp results hold for the class  $\mathcal{S}_{Ne}^*$ .*

- (i)  $R_{\mathcal{S}_{Ne}^*}(\mathcal{G}_1) = (\sqrt{85} - 9)/2 \approx 0.1097$
- (ii)  $R_{\mathcal{S}_{Ne}^*}(\mathcal{G}_2) = (\sqrt{265} - 15)/10 \approx 0.1278$
- (iii)  $R_{\mathcal{S}_{Ne}^*}(\mathcal{G}_3) = \sqrt{10} - 3 \approx 0.1622$

**PROOF.** (i) The function defined by  $m(r) = (1 - 6r - r^2)(1 - r^2)^{-1}$ ,  $0 \leq r < 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_{Ne}^*}(\mathcal{G}_1)$  is the root of the equation  $m(r) = 1/3$ . For  $0 < r \leq R_{\mathcal{S}_{Ne}^*}(\mathcal{G}_1)$ , we have  $m(r) \geq 1/3$ . That is,

$$\frac{6r}{1 - r^2} \leq 1 - \frac{1}{3}.$$

For the class  $\mathcal{G}_1$ , the centre of the disc is 1, therefore the disc obtained in (2.7) is contained in the region bounded by the nephroid, by Lemma 3.11. For the function  $f_1$  defined in (2.3), at  $z = R_{\mathcal{S}_{Ne}^*}(\mathcal{G}_1) = \rho$ ,

$$\left| \frac{zf_1'(z)}{f_1(z)} \right| = \left| \frac{1 + 6\rho - \rho^2}{1 - \rho^2} \right| = \frac{1}{3} \in \partial\Omega_{Ne}$$

where  $\partial\Omega_{Ne}$  denotes the boundary of nephroid domain.

- (ii) The function  $n(r) : [0, 1) \rightarrow \mathbb{R}$  defined by  $n(r) = (1 - 5r - 2r^2)(1 - r^2)^{-1} + 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_{Ne}^*}(\mathcal{G}_2)$  is the root of the equation  $n(r) = 1/3$ . For  $0 < r \leq R_{\mathcal{S}_{Ne}^*}(\mathcal{G}_2)$ , we have  $n(r) \geq 1/3$ . That is,

$$\frac{r(r + 5)}{1 - r^2} \leq \frac{2}{3}.$$

For the class  $\mathcal{G}_2$ , the centre of the disc is 1, therefore the disc obtained in (2.12) is contained in the region bounded by the nephroid, by Lemma 3.11. For the function  $f_2$  defined in (2.8), at  $z = R_{\mathcal{S}_{Ne}^*}(\mathcal{G}_2) = \rho$ ,

$$\left| \frac{zf_2'(z)}{f_2(z)} \right| = \left| \frac{1 + 5\rho}{1 - \rho^2} \right| = \frac{5}{3} \in \partial\Omega_{Ne}.$$

- (iii) The function defined by  $s(r) = 1 - (4r(1 - r^2)^{-1})$ ,  $0 \leq r < 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_{Ne}^*}(\mathcal{G}_3)$  is the root of the equation  $s(r) = 1/3$ . For  $0 < r \leq R_{\mathcal{S}_{Ne}^*}(\mathcal{G}_3)$ , we have  $s(r) \geq 1/3$ . That is,

$$\frac{4r}{1 - r^2} \leq \frac{2}{3}.$$

For the class  $\mathcal{G}_3$ , the centre of the disc is 1, therefore the disc obtained in (2.15) is contained in the region bounded by the nephroid, by Lemma 3.11. For the function  $g_1$  defined in (2.3), at  $z = R_{\mathcal{S}_{Ne}^*}(\mathcal{G}_3) = \rho$ ,

$$\left| \frac{zg_1'(z)}{g_1(z)} \right| = \left| \frac{1 + 4\rho - \rho^2}{1 - \rho^2} \right| = \frac{1}{3} \in \partial\Omega_{Ne}. \quad \blacksquare$$

In 2020, Goel and Kumar [10] introduced the class  $\mathcal{S}_{SG}^*$  that maps the open unit disc  $\mathbb{D}$  onto a domain  $\Delta_{SG} := \{w \in \mathbb{C} : |\log w/(2-w)| < 1\}$  and  $\mathcal{S}_{SG}^* = \mathcal{S}^*(2/(1+e^{-z}))$ . For  $2/(1+e) < a < 2e/(1+e)$ , they had proved the following inclusion:

$$\{w \in \mathbb{C} : |w - a| < r_{SG}\} \subset \Delta_{SG}, \quad (3.12)$$

provided  $r_{SG} = ((e-1)/(e+1)) - |a-1|$ . Theorem 3.11 provides  $\mathcal{S}_{SG}^*$ -radii of the classes  $\mathcal{G}_1, \mathcal{G}_2, \mathcal{G}_3$ .

**THEOREM 3.11.** *The following sharp results hold for the class  $\mathcal{S}_{SG}^*$ .*

- (i)  $R_{\mathcal{S}_{SG}^*}(\mathcal{G}_1) = 2(e-1)/((6+6e) + \sqrt{40+64e+40e^2}) \approx 0.0766$
- (ii)  $R_{\mathcal{S}_{SG}^*}(\mathcal{G}_2) = 2(e-1)/((5+5e) + \sqrt{25+42e+33e^2}) \approx 0.0901$
- (iii)  $R_{\mathcal{S}_{SG}^*}(\mathcal{G}_3) = 2(e-1)/((4+4e) + \sqrt{20+24e+20e^2}) \approx 0.1140$

**PROOF.** (i) The function defined by  $m(r) = (1-6r-r^2)(1-r^2)^{-1}$ ,  $0 \leq r < 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_{SG}^*}(\mathcal{G}_1)$  is the root of the equation  $m(r) = 2/(1+e)$ . For  $0 < r \leq R_{\mathcal{S}_{SG}^*}(\mathcal{G}_1)$ , we have  $m(r) \geq 2/(1+e)$ . That is,

$$\frac{6r}{1-r^2} \leq \frac{e-1}{e+1}.$$

For the class  $\mathcal{G}_1$ , the centre of the disc is 1, therefore the disc obtained in (2.7) is contained in the region bounded by the modified sigmoid, by Lemma 3.12. For the function  $f_1$  defined in (2.3), at  $z = R_{\mathcal{S}_{SG}^*}(\mathcal{G}_1) = \rho$ ,

$$\left| \log \frac{zf_1'(z)/f_1(z)}{2 - (zf_1'(z)/f_1(z))} \right| = \left| \log \frac{(1+6\rho-\rho^2)/(1-\rho^2)}{2 - ((1+6\rho-\rho^2)/(1-\rho^2))} \right| = 1.$$

- (ii) The function  $n(r) : [0, 1) \rightarrow \mathbb{R}$  defined by  $n(r) = (1-5r-2r^2)(1-r^2)^{-1} + 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_{SG}^*}(\mathcal{G}_2)$  is the root of the equation  $n(r) = 2/(1+e)$ . For  $0 < r \leq R_{\mathcal{S}_{SG}^*}(\mathcal{G}_2)$ , we have  $n(r) \geq 2/(1+e)$ . That is,

$$\frac{r(r+5)}{1-r^2} \leq \frac{e-1}{e+1}.$$

For the class  $\mathcal{G}_2$ , the centre of the disc is 1, therefore the disc obtained in (2.12) is contained in the region bounded by the modified sigmoid, by Lemma 3.12. For the function  $f_2$  defined in (2.8), at  $z = R_{\mathcal{S}_{SG}^*}(\mathcal{G}_2) = \rho$ ,

$$\left| \log \frac{zf_2'(z)/f_2(z)}{2 - (zf_2'(z)/f_2(z))} \right| = \left| \log \frac{(1+5\rho)/(1-\rho^2)}{2 - ((1+5\rho)/(1-\rho^2))} \right| = 1.$$

- (iii) The function defined by  $s(r) = 1 - (4r(1-r^2)^{-1})$ ,  $0 \leq r < 1$  is a decreasing function. Let  $\rho = R_{\mathcal{S}_{SG}^*}(\mathcal{G}_3)$  is the root of the equation  $s(r) = 2/(1+e)$ . For  $0 < r \leq R_{\mathcal{S}_{SG}^*}(\mathcal{G}_3)$ , we have  $s(r) \geq 2/(1+e)$ . That is,

$$\frac{4r}{1-r^2} \leq \frac{e-1}{e+1}.$$

For the class  $\mathcal{G}_3$ , the centre of the disc is 1, therefore the disc obtained in (2.15) is contained in the region bounded by the modified sigmoid, by Lemma 3.12. For the function  $g_1$  defined in (2.3), at  $z = R_{\mathcal{S}_{SG}^*}(\mathcal{G}_3) = \rho$ ,

$$\left| \log \frac{zg_1'(z)/g_1(z)}{2 - (zg_1'(z)/g_1(z))} \right| = \left| \log \frac{(1+4\rho-\rho^2)/(1-\rho^2)}{2 - ((1+4\rho-\rho^2)/(1-\rho^2))} \right| = 1. \quad \blacksquare$$

Though we have no proof, we believe that the sharp radii for the class  $\mathcal{G}_2$  are the following:

- (1)  $R_{\mathcal{S}_p}(\mathcal{G}_2) = 5 - 2\sqrt{6} \approx 0.1010$ .
- (2)  $R_{\mathcal{S}_e^*}(\mathcal{G}_2) = 2(e^2 + e + 2)/(2 + e) (5e + \sqrt{-8 + 4e + 29e^2}) \approx 0.1276$ .
- (3)  $R_{\mathcal{S}_c^*}(\mathcal{G}_2) = (15 - \sqrt{217})/2 \approx 0.1345$ .
- (4)  $R_{\mathcal{S}_\zeta^*}(\mathcal{G}_2) = (6\sqrt{2} - 8)/(\sqrt{2} - 1) (5 + \sqrt{41 - 12\sqrt{2}}) \approx 0.1183$ .
- (5)  $R_{\mathcal{S}_R^*}(\mathcal{G}_2) = (10\sqrt{2} - 14)/(\sqrt{2} - 1) (5 + \sqrt{81 - 40\sqrt{2}}) \approx 0.0345$ .

Our estimate for these radii are respectively 0.0972, 0.1213, 0.1279, 0.1131, 0.0340 and are very much close to the above mentioned values.

## References

- [1] R. M. Ali, N. K. Jain and V. Ravichandran, *Radii of starlikeness associated with the lemniscate of Bernoulli and the left-half plane*, Appl. Math. Comput. **218** (2012), no. 11, 6557–6565.
- [2] R. M. Ali, N. K. Jain and V. Ravichandran, *On the radius constants for classes of analytic functions*, Bull. Malays. Math. Sci. Soc. (2) **36** (2013), no. 1, 23–38.
- [3] M. K. Aouf, J. Dziok and J. Sokół, *On a subclass of strongly starlike functions*, Appl. Math. Lett. **24** (2011), no. 1, 27–32.
- [4] W. M. Causey and E. P. Merkes, *Radii of starlikeness of certain classes of analytic functions*, J. Math. Anal. Appl. **31** (1970), 579–586.
- [5] N. E. Cho, V. Kumar, S. S. Kumar and V. Ravichandran, *Radius problems for starlike functions associated with the sine function*, Bull. Iranian Math. Soc. **45** (2019), no. 1, 213–232.
- [6] P. L. Duren, *Univalent Functions*, GTM, vol. 259, Springer-Verlag, New York, 1983.
- [7] A. S. A. El-Faqeer, M. H. Mohd, V. Ravichandran and S. Supramaniam, *Starlikeness of certain analytic functions*, arXiv preprint arXiv:2006.11734 (2020).
- [8] S. Gandhi and V. Ravichandran, *Starlike functions associated with a lune*, Asian-Eur. J. Math. **10** (2017), no. 4, 1750064, 12.
- [9] A. Gangadharan, V. Ravichandran and T. N. Shanmugam, *Radii of convexity and strong starlikeness for some classes of analytic functions*, J. Math. Anal. Appl. **211** (1997), no. 1, 301–313.
- [10] P. Goel and S. Sivaprasad Kumar, *Certain class of starlike functions associated with modified sigmoid function*, Bull. Malays. Math. Sci. Soc. **43** (2020), no. 1, 957–991.
- [11] A. W. Goodman, *Univalent Functions. Vol. II*, Mariner, Tampa, FL, 1983.
- [12] W. Janowski, *Extremal problems for a family of functions with positive real part and for some related families*, Ann. Polon. Math. **23** (1970/71), 159–177.
- [13] W. Janowski, *Some extremal problems for certain families of analytic functions. I*, Ann. Polon. Math. **28** (1973), 297–326.
- [14] R. Kanaga and V. Ravichandran, *Starlikeness for certain close-to-star functions*, Hacet. J. Math. Stat. to appear. (arXiv:2003.05628)
- [15] K. Khatter, S. K. Lee and V. Ravichandran, *Radius of starlikeness for classes of analytic functions*, arXiv preprint arXiv:2006.11744 (2020).
- [16] B. Kowalczyk and A. Lecko, *Radius problem in classes of polynomial close-to-convex functions I*, Bull. Soc. Sci. Lett. Łódź Sér. Rech. Déform. **63** (2013), no. 1, 65–77.
- [17] B. Kowalczyk and A. Lecko, *Radius problem in classes of polynomial close-to-convex functions II. Partial solutions*, Bull. Soc. Sci. Lett. Łódź Sér. Rech. Déform. **63** (2013), no. 2, 23–34.
- [18] B. Kowalczyk, A. Lecko and B. Śmiarowska, *On some coefficient inequality in the subclass of close-to-convex functions*, Bull. Soc. Sci. Lett. Łódź Sér. Rech. Déform. **67** (2017), no. 1, 79–90.
- [19] S. Kumar and V. Ravichandran, *A subclass of starlike functions associated with a rational function*, Southeast Asian Bull. Math. **40** (2016), no. 2, 199–212.
- [20] A. Lecko and Y. J. Sim, *Coefficient problems in the subclasses of close-to-star functions*, Results Math. **74** (2019), no. 3, Paper No. 104, 14.

- [21] W. C. Ma and D. Minda, *Uniformly convex functions*, Ann. Polon. Math. **57** (1992), no. 2, 165–175.
- [22] W. C. Ma and D. Minda, *A unified treatment of some special classes of univalent functions*, Proceedings of the Conference on Complex Analysis (Tianjin, 1992), Conf. Proc. Lecture Notes Anal., I, Int. Press, Cambridge, MA, 1994, pp. 157–169.
- [23] T. H. MacGregor, *The radius of convexity for starlike functions of order  $\frac{1}{2}$* , Proc. Amer. Math. Soc. **14** (1963), 71–76.
- [24] T. H. MacGregor, *The radius of univalence of certain analytic functions*, Proc. Amer. Math. Soc. **14** (1963), 514–520.
- [25] T. H. MacGregor, *A class of univalent functions*, Proc. Amer. Math. Soc. **15** (1964), 311–317.
- [26] R. Mendiratta, S. Nagpal and V. Ravichandran, *A subclass of starlike functions associated with left-half of the lemniscate of Bernoulli*, Internat. J. Math. **25** (2014), no. 9, 1450090, 17.
- [27] R. Mendiratta, S. Nagpal and V. Ravichandran, *On a subclass of strongly starlike functions associated with exponential function*, Bull. Malays. Math. Sci. Soc. **38** (2015), no. 1, 365–386.
- [28] E. Paprocki and J. Sokół, *The extremal problems in some subclass of strongly starlike functions*, Zeszyty Nauk. Politech. Rzeszowskiej Mat. (1996), no. 20, 89–94.
- [29] R. K. Raina and J. Sokół, *Some properties related to a certain class of starlike functions*, C. R. Math. Acad. Sci. Paris **353** (2015), no. 11, 973–978.
- [30] J. S. Ratti, *The radius of convexity of certain analytic functions*, Indian J. Pure Appl. Math. **1** (1970), no. 1, 30–36.
- [31] J. S. Ratti, *The radius of univalence of certain analytic functions*, Math. Z. **107** (1968), 241–248.
- [32] V. Ravichandran, F. Rønning and T. N. Shanmugam, *Radius of convexity and radius of starlikeness for some classes of analytic functions*, Complex Variables Theory Appl. **33** (1997), no. 1-4, 265–280.
- [33] A. Sebastian and V. Ravichandran, *Radius of starlikeness of certain analytic functions*, Math. Slovaca to appear (arXiv:2001.06999).
- [34] G. M. Shah, *On the univalence of some analytic functions*, Pacific J. Math. **43** (1972), 239–250.
- [35] T. N. Shanmugam, *Convolution and differential subordination*, Internat. J. Math. Math. Sci. **12** (1989), no. 2, 333–340.
- [36] T. N. Shanmugam and V. Ravichandran, *Certain properties of uniformly convex functions*, Computational methods and function theory 1994 (Penang), Ser. Approx. Decompos., vol. 5, World Sci. Publ., River Edge, NJ, 1995, pp. 319–324.
- [37] K. Sharma, N. K. Jain and V. Ravichandran, *Starlike functions associated with a cardioid*, Afr. Mat. **27** (2016), no. 5-6, 923–939.
- [38] J. Sokół and J. Stankiewicz, *Radius of convexity of some subclasses of strongly starlike functions*, Zeszyty Nauk. Politech. Rzeszowskiej Mat. (1996), no. 19, 101–105.
- [39] H. M. Srivastava and S. Owa (eds.), *Current topics in analytic function theory*, World Scientific Publishing Co., Inc., River Edge, NJ, 1992.
- [40] L. A. Wani and A. Swaminathan, *Starlike and convex functions associated with a nephroid domain*, Bull. Malays. Math. Sci. Soc. to appear (2020).

DEPARTMENT OF COMPLEX ANALYSIS, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF WARMIA AND MAZURY IN OLSZTYN, UL. SŁONECZNA 54, 10-710, OLSZTYN, POLAND

*E-mail address:* alecko@matman.uwm.edu.pl

DEPARTMENT OF MATHEMATICS, NATIONAL INSTITUTE OF TECHNOLOGY, TIRUCHIRAPPALLI-620015, INDIA

*E-mail address:* vravi68@gmail.com; ravic@nitt.edu

DEPARTMENT OF MATHEMATICS, NATIONAL INSTITUTE OF TECHNOLOGY, TIRUCHIRAPPALLI-620015, INDIA

*E-mail address:* ashanitt18@gmail.com