

Boundedness of Dunkl-Hausdorff operator in Lebesgue spaces

Sandhya Jain

Department of Mathematics
Vivekananda College (University of Delhi)
Vivek Vihar, Delhi - 110095, India
(email : singhal.sandhya@gmail.com)

Alberto Fiorenza

Dipartimento di Architettura
Università di Napoli Federico II, Via Monteoliveto, 3
I-80134 Napoli, Italy
and Istituto per le Applicazioni del Calcolo “Mauro Picone”, sezione di Napoli
Consiglio Nazionale delle Ricerche
via Pietro Castellino, 111
I-80131 Napoli, Italy email :fiorenza@unina.it

Pankaj Jain

Department of Mathematics
South Asian University
Akbar Bhawan, Chanakya Puri, New Delhi-110021, India
(email: pankaj.jain@sau.ac.in, pankajkrjain@hotmail.com)

Abstract

In this paper, the $L_v^p(\mathbb{R})$ -boundedness of the Dunkl-Hausdorff operator $H_{\alpha,\phi}f(x) = \int_{\mathbb{R}} \frac{|\phi(t)|}{|t|^{2\alpha+2}} f\left(\frac{x}{t}\right) dt$ has been characterized and for a certain type of weight v , the precise value of the norm $\|H_{\alpha,\phi}\|_{L_v^p(\mathbb{R}) \rightarrow L_v^p(\mathbb{R})}$ has been obtained. This covers several of the existing results. Analogous results in two dimensions have also been proved.

Keywords: Hausdorff operator, Dunkl operator, Dunkl-Hausdorff operator, weighted Lebesgue spaces.

2010 Mathematics Subject Classification. 47B38, 26D10, 26D15.

1 Introduction

Let v be a weight function, i.e., a function which is measurable, positive and finite almost everywhere on the specified domain. By $L_v^p(\mathbb{R})$, $1 \leq p < \infty$, we denote the weighted Lebesgue space and a norm of a function $f \in L_v^p(\mathbb{R})$ is given by

$$\|f\|_{L_v^p(\mathbb{R})} := \left(\int_{\mathbb{R}} |f(x)|^p v(x) dx \right)^{1/p}.$$

Occasionally, we shall be referring to the specific weight $v(x) = |x|^{2\alpha+1}$. The corresponding weighted Lebesgue space will be denoted by $L_\alpha^p(\mathbb{R})$. The non-weighted Lebesgue space, i.e., when $v \equiv 1$, will be denoted by $L^p(\mathbb{R})$.

Let $\phi \in L^1(\mathbb{R})$. In the present paper, we are concerned with the Dunkl-Hausdorff operator [4, 5, 6, 7]

$$H_{\alpha,\phi}f(x) = \int_{\mathbb{R}} \frac{|\phi(t)|}{|t|^{2\alpha+2}} f\left(\frac{x}{t}\right) dt.$$

When $\alpha = -1/2$, The operator $H_{\alpha,\phi}$ is the famous Hausdorff operator

$$H_{\phi}f(x) = \int_{\mathbb{R}} \frac{|\phi(t)|}{|t|} f\left(\frac{x}{t}\right) dt,$$

from which several well known operators can be deduced for suitable choices of ϕ , e.g., for $\phi(t) = \frac{1}{t}\chi_{(1,\infty)}(t)$, the operator H_{ϕ} reduces to the standard Hardy averaging operator

$$Hf(x) = \frac{1}{x} \int_0^x f(t) dt$$

while for $\phi(t) = \chi_{[0,1]}(t)$, it reduces to the adjoint of Hardy averaging operator

$$H^*f(x) = \int_x^{\infty} \frac{f(t)}{t} dt.$$

Similarly, other operators like Calderon operator, Cesàro operator and fractional Riemann Liouville operator can also be deduced from H_{ϕ} , see [3, 10] for details. For more updates on the Hausdorff operator, its extensions and in the framework of other function spaces one may refer to [1, 2, 11, 14, 15] and the survey [13].

By the replacement $\phi(s) = \frac{1}{s}\psi\left(\frac{1}{s}\right)$, $s > 0$, the operator H_{ϕ} (considered on \mathbb{R}^+) becomes equivalent to

$$G_{\psi}g(x) = \frac{1}{x} \int_0^{\infty} \psi\left(\frac{t}{x}\right) g(t) dt.$$

It was proved by Golberg ([8], Theorem 1) that if $\psi \geq 0$ on \mathbb{R}^+ is such that $\int_{\mathbb{R}} \frac{\psi(t)}{\sqrt{t}} dt =: K < \infty$, then the operator G_{ψ} (and consequently H_{ϕ}) is a bounded operator on $L^2(\mathbb{R}^+)$ and $\|G_{\psi}\| \leq K$. The L^p -boundedness of G_{ψ} is derived in ([9], Theorem 319) and for many other extensions with sharp constants one may refer to ([12], Theorem 6.4 and bibliographic notes to Chapter 2 therein). In [10], the authors reestablished the $L^p(\mathbb{R}^+)$ -boundedness of G_{ψ} and via a new proof of the lower bound, obtained the precise value of $\|G_{\psi}\|_{L^p(\mathbb{R}^+) \rightarrow L^p(\mathbb{R}^+)}$ as

$$\|G_{\psi}\|_{L^p(\mathbb{R}^+) \rightarrow L^p(\mathbb{R}^+)} = \int_0^{\infty} \frac{\psi(t)}{t^{1/p}} dt =: K_p, \quad 1 < p < \infty.$$

Recently, in [3], a two weight characterization of the boundedness of H_{ϕ} between $L_v^p(\mathbb{R}^+)$ and $L_w^p(\mathbb{R}^+)$ has been given. Moreover, in the same paper, the corresponding boundedness has been studied in the framework of other function spaces as well, namely, grand Lebesgue spaces and variable exponent Lebesgue spaces.

Coming back to the Dunkl-Hausdorff operator $H_{\alpha,\phi}$, its $L_{\alpha}^1(\mathbb{R})$ boundedness has been proved in [4] whereas $L_{\alpha}^p(\mathbb{R})$ boundedness is obtained in [6] and in each case, a sufficient condition has been provided. We, in this paper, generalize these results by providing a characterization for the $L_v^p(\mathbb{R})$ boundedness of $H_{\alpha,\phi}$ and for a certain type of weight v , we provide the precise value of the norm $\|H_{\alpha,\phi}\|_{L_v^p(\mathbb{R}) \rightarrow L_v^p(\mathbb{R})}$. Moreover, a sufficient condition has been proved for two weights and two indices boundedness, i.e., $H_{\alpha,\phi} : L_w^p(\mathbb{R}) \rightarrow L_v^q(\mathbb{R})$ boundedness. These results have also been proved in the two dimensional framework.

2 One dimensional case

We begin by proving the following:

Theorem 2.1. *Let $1 < p < \infty$, v be weight function and $\phi \in L^1(\mathbb{R})$ be such that*

$$\int_{\mathbb{R}} \frac{|\phi(t)|}{|t|^{2\alpha+2-\frac{1}{p}}} \left(\sup_{y \in \mathbb{R}} \frac{v(ty)}{v(y)} \right)^{1/p} dt =: A_{\text{sup}} < \infty.$$

Then the operator $H_{\alpha, \phi}$ is a bounded operator on $L_v^p(\mathbb{R})$ and

$$\|H_{\alpha, \phi} f\|_{L_v^p(\mathbb{R})} \leq A_{\text{sup}} \|f\|_{L_v^p(\mathbb{R})}.$$

Proof. If $f \in L_v^p(\mathbb{R})$ then by using generalised Minkowski inequality, change of variables and Hölder's inequality, we have

$$\begin{aligned} \|H_{\alpha, \phi} f\|_{L_v^p(\mathbb{R})} &= \left(\int_{\mathbb{R}} |H_{\alpha, \phi} f(x)|^p v(x) dx \right)^{\frac{1}{p}} \\ &= \left(\int_{\mathbb{R}} \left| \int_{\mathbb{R}} \frac{|\phi(t)|}{|t|^{2\alpha+2}} f\left(\frac{x}{t}\right) dt \right|^p v(x) dx \right)^{\frac{1}{p}} \\ &\leq \left(\int_{\mathbb{R}} \left(\int_{\mathbb{R}} \frac{|\phi(t)|}{|t|^{2\alpha+2}} \left| f\left(\frac{x}{t}\right) \right| v^{\frac{1}{p}}(x) dt \right)^p dx \right)^{\frac{1}{p}} \\ &\leq \int_{\mathbb{R}} \frac{|\phi(t)|}{|t|^{2\alpha+2}} |t|^{\frac{1}{p}} \left(\int_{\mathbb{R}} |f(y)|^p v(yt) dy \right)^{\frac{1}{p}} dt \\ &= \int_{\mathbb{R}} \frac{|\phi(t)|}{|t|^{2\alpha+2-\frac{1}{p}}} \left(\int_{\mathbb{R}} |f(y)|^p v(y) v^{-1}(y) v(yt) dy \right)^{\frac{1}{p}} dt \tag{2.1} \\ &\leq \left(\int_{\mathbb{R}} \frac{|\phi(t)|}{|t|^{2\alpha+2-\frac{1}{p}}} \left(\sup_{y \in \mathbb{R}} \frac{v(ty)}{v(y)} \right)^{\frac{1}{p}} dt \right) \left(\int_{\mathbb{R}} |f(y)|^p v(y) dy \right)^{\frac{1}{p}} \\ &= A_{\text{sup}} \|f\|_{L_v^p(\mathbb{R})} \end{aligned}$$

and the assertion follows. \square

The following theorem provides a converse of Theorem 2.1. Here and throughout p' denotes the conjugate index to p , i.e., $\frac{1}{p} + \frac{1}{p'} = 1$.

Theorem 2.2. *Let $1 < p < \infty$, v be a weight function and $\phi \in L^1(\mathbb{R})$. If the operator $H_{\alpha, \phi}$ is a bounded operator on $L_v^p(\mathbb{R})$, then*

$$\|H_{\alpha, \phi}\| \geq \int_{\mathbb{R}} \frac{|\phi(t)|}{|t|^{2\alpha+2-\frac{1}{p}}} \left(\inf_{y \in \mathbb{R}} \frac{v(ty)}{v(y)} \right)^{1/p} dt =: A_{\text{inf}}.$$

Proof. Let us consider $0 \leq f \in L_v^p(\mathbb{R})$ and $0 \leq g \in L_{v^{1-p'}}^{p'}(\mathbb{R})$. On using Fubini's Theorem and Hölder's inequality, We have

$$J := \int_{\mathbb{R}} \frac{|\phi(t)|}{|t|^{2\alpha+2}} \left(\int_{\mathbb{R}} f\left(\frac{x}{t}\right) g(x) dx \right) dt \tag{2.2}$$

$$\begin{aligned}
&= \int_{\mathbb{R}} g(x) \left(\int_{\mathbb{R}} \frac{|\phi(t)|}{|t|^{2\alpha+2}} f\left(\frac{x}{t}\right) dt \right) dx \\
&\leq \int_{\mathbb{R}} |g(x)| |(H_{\alpha,\phi}f)(x)| dx \\
&= \int_{\mathbb{R}} |g(x)| v^{\frac{1}{p}}(x) v^{-\frac{1}{p}}(x) |(H_{\alpha,\phi}f)(x)| dx \\
&\leq \left(\int_{\mathbb{R}} |(H_{\alpha,\phi}f)(x)|^p v(x) dx \right)^{1/p} \left(\int_{\mathbb{R}} |g(x)|^{p'} v^{-\frac{p'}{p}}(x) dx \right)^{\frac{1}{p'}} \\
&\leq \|H_{\alpha,\phi}\| \|f\|_{L_v^p(\mathbb{R})} \|g\|_{L_{v^{1-p'}}^{p'}(\mathbb{R})}.
\end{aligned} \tag{2.3}$$

For any interval $I = (a, b)$, I' will denote the interval $(-b, -a)$. Now, for $u \in (0, 1)$, let $I_1 = (u, 1/u)$ so that $I_1' = (-1/u, -u)$. Define the test functions

$$f_u(x) = \frac{v^{-1/p}(x)}{|x|^{1/p}} \chi_{I_1 \cup I_1'}(x), \quad g_u(x) = \frac{v^{1/p}(x)}{|x|^{1/p'}} \chi_{I_1 \cup I_1'}(x).$$

Then it can be calculated that

$$\|f_u\|_{L_v^p(\mathbb{R})}^p = \|g_u\|_{L_{v^{1-p'}}^{p'}(\mathbb{R})}^{p'} = 4 \log(1/u). \tag{2.4}$$

Also, we have

$$\begin{aligned}
h_u(t) &:= \int_{\mathbb{R}} f_u\left(\frac{x}{t}\right) g_u(x) dx \\
&= |t|^{1/p} \int_{\mathbb{R}} \frac{1}{|y|} \chi_{I_1 \cup I_1'}(y) \chi_{I_1 \cup I_1'}(ty) v^{-1/p}(y) v^{1/p}(ty) dy \\
&= |t|^{1/p} \int_{\mathbb{R}} \frac{1}{|y|} \chi_{I_{t,u}}(y) v^{-1/p}(y) v^{1/p}(ty) dy \\
&\geq \inf_{y \in \mathbb{R}} \left(\frac{v(ty)}{v(y)} \right)^{1/p} |t|^{1/p} \int_{\mathbb{R}} \frac{1}{|y|} \chi_{I_{t,u}}(y) dy
\end{aligned} \tag{2.5}$$

where

$$\begin{aligned}
I_{t,u} &= \begin{cases} (I_1 \cup I_1') \cap (I_2 \cup I_2'), & \text{if } t \geq 0 \\ (I_1 \cup I_1') \cap (I_3 \cup I_3'), & \text{if } t < 0 \end{cases} \\
&= \begin{cases} (I_1 \cap I_2) \cup (I_1' \cap I_2'), & \text{if } t \geq 0 \\ (I_1 \cap I_3) \cup (I_1' \cap I_3), & \text{if } t < 0 \end{cases}
\end{aligned}$$

with $I_2 = (\frac{u}{t}, \frac{1}{ut})$ and $I_3 = (\frac{1}{tu}, \frac{u}{t})$. We divide \mathbb{R} as

$$\mathbb{R} = \left(-\infty, -\frac{1}{u^2}, \right] \cup \left(-\frac{1}{u^2}, -1\right] \cup (-1, -u^2] \cup (-u^2, 0] \cup (0, u^2] \cup (u^2, 1] \cup \left(1, \frac{1}{u^2}\right] \cup \left(\frac{1}{u^2}, \infty\right).$$

If $t \in (-\infty, -\frac{1}{u^2},] \cup [-u^2, u^2] \cup [\frac{1}{u^2}, \infty) \cup \{-1, 1\}$, then $I_{t,u} = \emptyset$, so that in this case

$$h_u(t) = 0. \tag{2.6}$$

If $t \in (-1/u^2, -1)$, then $I_{t,u} = (u, -\frac{1}{tu}) \cup (\frac{1}{ut}, -u)$ and

$$\int_{\mathbb{R}} \frac{1}{|y|} \chi_{I_{t,u}} dy = -2 \log |t| + 4 \log \frac{1}{u}. \quad (2.7)$$

If $t \in (-1, -u^2)$, then $I_{t,u} = (-\frac{1}{u}, \frac{u}{t}) \cup (-\frac{u}{t}, \frac{1}{u})$ and we have

$$\int_{\mathbb{R}} \frac{1}{|y|} \chi_{I_{t,u}} dy = 2 \log |t| + 4 \log \frac{1}{u}. \quad (2.8)$$

If $t \in (u^2, 1)$, then $I_{t,u} = (\frac{u}{t}, \frac{1}{u}) \cup (-\frac{1}{u}, -\frac{u}{t})$ and

$$\int_{\mathbb{R}} \frac{1}{|y|} \chi_{I_{t,u}} dy = 2 \log |t| + 4 \log \frac{1}{u}. \quad (2.9)$$

If $t \in (1, \frac{1}{u^2})$, then $I_{t,u} = (u, \frac{1}{tu}) \cup (-\frac{1}{tu}, -u)$ and we have

$$\int_{\mathbb{R}} \frac{1}{|y|} \chi_{I_{t,u}} dy = -2 \log |t| + 4 \log \frac{1}{u}. \quad (2.10)$$

On taking f and g as f_u and g_u in (2.2) and then using (2.5)-(2.10), we obtain

$$\begin{aligned} J &= \left(\int_{-\infty}^{-1/u^2} + \int_{-1/u^2}^{-1} + \int_{-1}^{-u^2} + \int_{-u^2}^0 + \int_0^{u^2} + \int_{u^2}^1 + \int_1^{1/u^2} + \int_{1/u^2}^{\infty} \right) \frac{|\phi(t)|}{|t|^{2\alpha+2}} h_u(t) dt \\ &\geq 4 \log \frac{1}{u} \left[\int_{-1/u^2}^{-u^2} \inf_{y \in \mathbb{R}} \left(\frac{v(ty)}{v(y)} \right)^{\frac{1}{p}} \frac{|\phi(t)|}{|t|^{2\alpha+2-\frac{1}{p}}} \left(1 - \frac{\xi(t)}{4 \log \frac{1}{u}} \right) dt \right. \\ &\quad \left. + \int_{u^2}^{1/u^2} \inf_{y \in \mathbb{R}} \left(\frac{v(ty)}{v(y)} \right)^{\frac{1}{p}} \frac{|\phi(t)|}{|t|^{2\alpha+2-\frac{1}{p}}} \left(1 - \frac{\xi(t)}{4 \log \frac{1}{u}} \right) dt \right], \end{aligned} \quad (2.11)$$

where

$$\xi(t) = \begin{cases} 2 \log |t|, & \text{if } t \in (-1/u^2, -1) \cup (1, 1/u^2) \\ -2 \log |t|, & \text{if } t \in (-1, -u^2) \cup (u^2, 1). \end{cases}$$

Now, using (2.11) and (2.4) in (2.3) for $f = f_u$ and $g = g_u$, we get

$$\begin{aligned} &\int_{-1/u^2}^{-u^2} \inf_{y \in \mathbb{R}} \left(\frac{v(ty)}{v(y)} \right)^{\frac{1}{p}} \frac{|\phi(t)|}{|t|^{2\alpha+2-\frac{1}{p}}} \left(1 - \frac{\xi(t)}{4 \log \frac{1}{u}} \right) dt \\ &\quad + \int_{u^2}^{1/u^2} \inf_{y \in \mathbb{R}} \left(\frac{v(ty)}{v(y)} \right)^{\frac{1}{p}} \frac{|\phi(t)|}{|t|^{2\alpha+2-\frac{1}{p}}} \left(1 - \frac{\xi(t)}{4 \log \frac{1}{u}} \right) dt \leq \|H_{\alpha, \phi}\|. \end{aligned}$$

By the Monotone Convergence Theorem, the LHS $\uparrow A_{\text{inf}}$ as $u \rightarrow 0$ and we are done. \square

In view of Theorems 2.1 and 2.2, a characterization for the boundedness of $H_{\alpha, \phi} : L_v^p(\mathbb{R}) \rightarrow L_v^p(\mathbb{R})$ can be derived. In fact, the following is immediate:

Theorem 2.3. *Let $1 < p < \infty$, v be weight function and $\phi \in L^1(\mathbb{R})$. Let the following be satisfied for some constant $c > 0$:*

$$\sup_{y \in \mathbb{R}} \frac{v(ty)}{v(y)} \leq c \inf_{y \in \mathbb{R}} \frac{v(ty)}{v(y)}.$$

Then the operator $H_{\alpha,\phi} : L_v^p(\mathbb{R}) \rightarrow L_v^p(\mathbb{R})$ is bounded if and only if $A_{\text{sup}} < \infty$ and moreover the following estimates hold:

$$\frac{1}{c} A_{\text{sup}} \leq \|H_{\alpha,\phi}\|_{L_v^p(\mathbb{R}) \rightarrow L_v^p(\mathbb{R})} \leq A_{\text{sup}}.$$

Corollary 2.4. *Let $1 < p < \infty$, v be weight function and $\phi \in L^1(\mathbb{R})$. If there exists a function h such that $v(xy) = v(x)h(y)$, then $A_{\text{sup}} = A_{\text{inf}}$ and*

$$\|H_{\alpha,\phi}\|_{L_v^p(\mathbb{R}) \rightarrow L_v^p(\mathbb{R})} = \int_{\mathbb{R}} \frac{|\phi(t)|}{|t|^{2\alpha+2-\frac{1}{p}}} h^{1/p}(t) dt.$$

For $\alpha = -1/2$ and $\phi(t) = \frac{1}{t} \chi_{(1,\infty)}(t)$, as pointed out earlier, the operator $H_{\alpha,\phi}$ becomes the Hardy averaging operator

$$Hf(x) = \frac{1}{x} \int_0^x f(t) dt.$$

Further, if we take $v(t) = t^\beta$, $\beta < p - 1$, then $A_{\text{sup}} = \frac{p}{\beta - p - 1}$ and

$$\|H\|_{L_{t^\beta}^p(\mathbb{R}) \rightarrow L_{t^\beta}^p(\mathbb{R})} = \frac{p}{\beta - p + 1}.$$

The above discussion leads to the following corollary which, in fact, is the classical Hardy inequality (see e.g. [9, Theorem 330], [16, (3.6) p. 23] or [17, Theorem 6 p. 726]):

Corollary 2.5. *Let $1 < p < \infty$ and $\beta < p - 1$ be a the weight function. Then the inequality*

$$\|Hf\|_{L_{x^\beta}^p(\mathbb{R})} \leq \left(\frac{p}{\beta - p + 1} \right) \|f\|_{L_{x^\beta}^p(\mathbb{R})}$$

holds and the constant $\left(\frac{p}{\beta - p + 1} \right)$ is sharp.

Remark 2.6. (i) When $p = 1$ and $v(x) = |x|^{2\alpha+1}$, Theorem 2.1 reduces to ([4], Theorem 3.1).

(ii) When $v(x) = |x|^{2\alpha+1}$, Theorem 2.1 reduces to ([6], Theorem 1)

(iii) When $\alpha = -1/2$, Theorems 2.1, 2.2, 2.3 and Corollary 2.4 reduce to ([3], Theorems 1(i), 1(ii), Corollaries 1 and 2, respectively for $w = v$). Moreover, the functions here are defined on \mathbb{R} unlike in [3] where the domain is \mathbb{R}^+ .

3 Two dimensional case

In this section, we derive the two dimensional analogues of the results proved in Section 2.

Definition 3.1. For $\phi \in L^1(\mathbb{R}^2)$, the two dimensional Dunkl-Hausdorff operator is defined by

$$\mathcal{H}_{\alpha,\phi}(f(x_1, x_2)) = \int_{\mathbb{R}^2} \frac{|\phi(t_1, t_2)|}{|t_1 t_2|^{2\alpha+2}} f\left(\frac{x_1}{t_1}, \frac{x_2}{t_2}\right) dt_1 dt_2.$$

Now, we prove the boundedness of $\mathcal{H}_{\alpha,\phi}$. By using generalised Minkowski inequality, change of variables and Hölder's inequality in two dimensions, the following theorem can be proved along the same lines as in Theorem 2.1.

Theorem 3.2. Let $1 < p < \infty$, v be weight function and $\phi \in L^1(\mathbb{R}^2)$ be such that

$$\int_{\mathbb{R}^2} \frac{|\phi(t_1, t_2)|}{|t_1 t_2|^{2\alpha+2-\frac{1}{p}}} \left(\sup_{(y_1, y_2) \in \mathbb{R}^2} \frac{v(t_1 y_1, t_2 y_2)}{v(y_1, y_2)} \right)^{1/p} dt_1 dt_2 =: \mathcal{A}_{\text{sup}} < \infty.$$

Then the operator $\mathcal{H}_{\alpha, \phi} : L_v^p(\mathbb{R}^2) \rightarrow L_v^p(\mathbb{R}^2)$ is bounded and

$$\|\mathcal{H}_{\alpha, \phi} f\|_{L_v^p(\mathbb{R}^2)} \leq \mathcal{A}_{\text{sup}} \|f\|_{L_v^p(\mathbb{R}^2)}.$$

Towards the converse of Theorem 3.2, we prove the following:

Theorem 3.3. Let $1 < p < \infty$, v be weight function and $\phi \in L^1(\mathbb{R}^2)$. If the operator $\mathcal{H}_{\alpha, \phi} : L_v^p(\mathbb{R}^2) \rightarrow L_v^p(\mathbb{R}^2)$ is bounded, then

$$\|\mathcal{H}_{\alpha, \phi}\| \geq \int_{\mathbb{R}^2} \frac{|\phi(t_1, t_2)|}{|t_1 t_2|^{2\alpha+2-\frac{1}{p}}} \left(\inf_{(y_1, y_2) \in \mathbb{R}^2} \frac{v(t_1 y_1, t_2 y_2)}{v(y_1, y_2)} \right)^{1/p} dt_1 dt_2 =: \mathcal{A}_{\text{inf}}.$$

Proof. Let $0 \leq f \in L_v^p(\mathbb{R}^2)$ and $0 \leq g \in L_{v^{1-p'}}^{p'}(\mathbb{R}^2)$. On using Fubini's Theorem and Hölder's inequality, We have

$$\mathcal{J} := \int_{\mathbb{R}^2} \frac{|\phi(t_1, t_2)|}{|t_1 t_2|^{2\alpha+2}} \int_{\mathbb{R}^2} f\left(\frac{x_1}{t_1}, \frac{x_2}{t_2}\right) g(x_1, x_2) dx_1 dx_2 dt_1 dt_2 \quad (3.1)$$

$$= \int_{\mathbb{R}^2} g(x_1, x_2) \left(\int_{\mathbb{R}^2} \frac{|\phi(t_1, t_2)|}{|t_1 t_2|^{2\alpha+2}} f\left(\frac{x_1}{t_1}, \frac{x_2}{t_2}\right) dt_1 dt_2 \right) dx_1 dx_2$$

$$\leq \int_{\mathbb{R}^2} |g(x_1, x_2)| |(\mathcal{H}_{\alpha, \phi} f)(x_1, x_2)| dx_1 dx_2$$

$$\leq \left(\int_{\mathbb{R}^2} |(\mathcal{H}_{\alpha, \phi} f)(x_1, x_2)|^p v(x_1, x_2) dx_1 dx_2 \right)^{1/p} \left(\int_{\mathbb{R}^2} |g(x_1, x_2)|^{p'} v(x_1, x_2)^{\frac{-p'}{p}} dx_1 dx_2 \right)^{1/p'}$$

$$\leq \|\mathcal{H}_{\alpha, \phi}\| \|f\|_{L_v^p(\mathbb{R}^2)} \|g\|_{L_{v^{1-p'}}^{p'}(\mathbb{R}^2)}. \quad (3.2)$$

Now, for $u \in (0, 1)$, we define the test functions

$$f_u(x_1, x_2) = \frac{v^{-1/p}(x_1, x_2)}{|x_1 x_2|^{\frac{1}{p}}} \chi_{I_1 \cup I_1' \times I_1 \cup I_1'}(x_1, x_2),$$

$$g_u(x_1, x_2) = \frac{v^{1/p}(x_1, x_2)}{|x_1 x_2|^{\frac{1}{p'}}} \chi_{I_1 \cup I_1' \times I_1 \cup I_1'}(x_1, x_2),$$

where $I_1 = (u, 1/u)$ and as before $I_1' = (-1/u, -u)$. Then it can be calculated that

$$\|f_u\|_{L_v^p(\mathbb{R}^2)}^p = \|g_u\|_{L_{v^{1-p'}}^{p'}(\mathbb{R}^2)}^{p'} = (4 \log(1/u))^2. \quad (3.3)$$

Also, on taking $x_i/t_i = y_i$ for $i = 1, 2$, we have

$$\begin{aligned} h_u(t_1, t_2) &:= \int_{\mathbb{R}^2} f_u\left(\frac{x_1}{t_1}, \frac{x_2}{t_2}\right) g_u(x_1, x_2) dx_1 dx_2 \\ &= |t_1 t_2|^{\frac{1}{p}} \int_{\mathbb{R}^2} \frac{1}{|y_1 y_2|} \chi_{I_1 \cup I_1' \times I_1 \cup I_1'}(y_1, y_2) v^{-\frac{1}{p}}(y_1, y_2) v^{\frac{1}{p}}(t_1 y_1, t_2 y_2) \chi_{I_1 \cup I_1' \times I_1 \cup I_1'}(t_1 y_1, t_2 y_2) dy_1 dy_2 \end{aligned}$$

$$\geq \inf_{(y_1, y_2) \in \mathbb{R}^2} \frac{v^{\frac{1}{p}}(t_1 y_1, t_2 y_2)}{v^{\frac{1}{p}}(y_1, y_2)} |t_1 t_2|^{\frac{1}{p}} \int_{\mathbb{R}^2} \frac{1}{|y_1 y_2|} \chi_{I_{t_1, t_2, u}}(y_1, y_2) dy_1 dy_2,$$

where

$$I_{t_1, t_2, u} = \begin{cases} (I_1 \cup I'_1) \cap (I_2 \cup I'_2) \times (I_1 \cup I'_1) \cap (I_3 \cup I'_3), & \text{if } t_1, t_2 > 0 \\ (I_1 \cup I'_1) \cap (I_4 \cup I'_4) \times (I_1 \cup I'_1) \cap (I_5 \cup I'_5), & \text{if } t_1, t_2 < 0 \\ (I_1 \cup I'_1) \cap (I_2 \cup I'_2) \times (I_1 \cup I'_1) \cap (I_5 \cup I'_5), & \text{if } t_1 > 0, t_2 < 0 \\ (I_1 \cup I'_1) \cap (I_4 \cup I'_4) \times (I_1 \cup I'_1) \cap (I_3 \cup I'_3), & \text{if } t_1 < 0, t_2 > 0. \end{cases}$$

$$= \begin{cases} (I_1 \cap I_2) \cup (I'_1 \cap I'_2) \times (I_1 \cap I_3) \cup (I'_1 \cap I'_3), & \text{if } t_1, t_2 > 0 \\ (I_1 \cap I_4) \cup (I'_1 \cap I'_4) \times (I_1 \cap I'_5) \cup (I'_1 \cap I_5), & \text{if } t_1, t_2 < 0 \\ (I_1 \cap I_2) \cup (I'_1 \cap I'_2) \times (I_1 \cap I'_5) \cup (I'_1 \cap I_5), & \text{if } t_1 > 0, t_2 < 0 \\ (I_1 \cap I_4) \cup (I'_1 \cap I'_4) \times (I_1 \cap I_3) \cup (I'_1 \cap I'_3), & \text{if } t_1 < 0, t_2 > 0, \end{cases}$$

with

$$I_2 = \left(\frac{u}{t_1}, \frac{1}{ut_1} \right) \quad I_3 = \left(\frac{u}{t_2}, \frac{1}{ut_2} \right) \quad I_4 = \left(\frac{1}{ut_1}, \frac{u}{t_1} \right) \quad I_5 = \left(\frac{1}{ut_2}, \frac{u}{t_2} \right).$$

It is observed that if $t_1 \in (-\infty, -\frac{1}{u^2}] \cup [-u^2, u^2] \cup [\frac{1}{u^2}, \infty) \cup \{-1, 1\}$ and $t_2 \in (-\infty, \infty)$, then $I_{t_1, t_2, u} = \emptyset$ and therefore, in this case $h_u(t_1, t_2) = 0$. The same is the situation if $t_2 \in (-\infty, -\frac{1}{u^2}] \cup [-u^2, u^2] \cup [\frac{1}{u^2}, \infty) \cup \{-1, 1\}$ and $t_1 \in (-\infty, \infty)$, then $h_u(t_1, t_2) = 0$. We deal with the remaining cases as follows:

Case 1 : $t_1, t_2 \in (u^2, 1)$. In this case, it can be worked out that

$$I_{t_1, t_2, u} = I_6 \cup I'_6 \times I_7 \cup I'_7, \quad \text{where } I_6 = \left(\frac{u}{t_1}, \frac{1}{u} \right), \quad I_7 = \left(\frac{u}{t_2}, \frac{1}{u} \right)$$

and therefore,

$$\int_{\mathbb{R}^2} \frac{1}{|y_1 y_2|} \chi_{I_{t_1, t_2, u}}(y_1, y_2) dy_1 dy_2 = \left(4 \log \frac{1}{u} \right)^2 \left(1 - \frac{\log |\frac{1}{t_1}|}{2 \log \frac{1}{u}} \right) \left(1 - \frac{\log |\frac{1}{t_2}|}{2 \log \frac{1}{u}} \right).$$

Case 2 : $t_1 \in (1, \frac{1}{u^2})$, $t_2 \in (u^2, 1)$. In this case

$$I_{t_1, t_2, u} = I_8 \cup I'_8 \times I_7 \cup I'_7, \quad \text{where } I_8 = \left(u, \frac{1}{ut_1} \right)$$

so that

$$\int_{\mathbb{R}^2} \frac{1}{|y_1 y_2|} \chi_{I_{t_1, t_2, u}}(y_1, y_2) dy_1 dy_2 = \left(4 \log \frac{1}{u} \right)^2 \left(1 - \frac{\log |t_1|}{2 \log \frac{1}{u}} \right) \left(1 - \frac{\log |\frac{1}{t_2}|}{2 \log \frac{1}{u}} \right).$$

Case 3 : $t_1 \in (u^2, 1)$, $t_2 \in (1, \frac{1}{u^2})$. In this case

$$I_{t_1, t_2, u} = I_6 \cup I'_6 \times I_9 \cup I'_9, \quad \text{where } I_9 = \left(u, \frac{1}{ut_2} \right)$$

so that

$$\int_{\mathbb{R}^2} \frac{1}{|y_1 y_2|} \chi_{I_{t_1, t_2, u}}(y_1, y_2) dy_1 dy_2 = \left(4 \log \frac{1}{u} \right)^2 \left(1 - \frac{\log |\frac{1}{t_1}|}{2 \log \frac{1}{u}} \right) \left(1 - \frac{\log |t_2|}{2 \log \frac{1}{u}} \right).$$

Case 4 : $t_1, t_2 \in (1, \frac{1}{u^2})$. In this case

$$I_{t_1, t_2, u} = I_8 \cup I'_8 \times I_9 \cup I'_9$$

so that

$$\int_{\mathbb{R}^2} \frac{1}{|y_1 y_2|} \chi_{I_{t_1, t_2, u}}(y_1, y_2) dy_1 dy_2 = \left(4 \log \frac{1}{u}\right)^2 \left(1 - \frac{\log |t_1|}{2 \log \frac{1}{u}}\right) \left(1 - \frac{\log |t_2|}{2 \log \frac{1}{u}}\right).$$

Case 5 : $t_1 \in (-\frac{1}{u^2}, -1), t_2 \in (u^2, 1)$. In this case, it can be worked out that

$$I_{t_1, t_2, u} = I_{10} \cup I'_{10} \times I_7 \cup I'_7, \quad \text{where } I_{10} = \left(u, -\frac{1}{ut_1}\right)$$

and therefore,

$$\int_{\mathbb{R}^2} \frac{1}{|y_1 y_2|} \chi_{I_{t_1, t_2, u}}(y_1, y_2) dy_1 dy_2 = \left(4 \log \frac{1}{u}\right)^2 \left(1 - \frac{\log |t_1|}{2 \log \frac{1}{u}}\right) \left(1 - \frac{\log |\frac{1}{t_2}|}{2 \log \frac{1}{u}}\right).$$

Case 6 : $t_1 \in (-1, -u^2), t_2 \in (u^2, 1)$. In this case

$$I_{t_1, t_2, u} = I_{11} \cup I'_{11} \times I_7 \cup I'_7, \quad \text{where } I_{11} = \left(-\frac{u}{t_1}, \frac{1}{u}\right)$$

so that

$$\int_{\mathbb{R}^2} \frac{1}{|y_1 y_2|} \chi_{I_{t_1, t_2, u}}(y_1, y_2) dy_1 dy_2 = \left(4 \log \frac{1}{u}\right)^2 \left(1 - \frac{\log |\frac{1}{t_1}|}{2 \log \frac{1}{u}}\right) \left(1 - \frac{\log |\frac{1}{t_2}|}{2 \log \frac{1}{u}}\right).$$

Case 7 : $t_1 \in (-\frac{1}{u^2}, -1), t_2 \in (1, \frac{1}{u^2})$. In this case

$$I_{t_1, t_2, u} = I_{10} \cup I'_{10} \times I_9 \cup I'_9$$

so that

$$\int_{\mathbb{R}^2} \frac{1}{|y_1 y_2|} \chi_{I_{t_1, t_2, u}}(y_1, y_2) dy_1 dy_2 = \left(4 \log \frac{1}{u}\right)^2 \left(1 - \frac{\log |t_1|}{2 \log \frac{1}{u}}\right) \left(1 - \frac{\log |t_2|}{2 \log \frac{1}{u}}\right).$$

Case 8 : $t_1 \in (-1, -u^2), t_2 \in (1, \frac{1}{u^2})$. In this case

$$I_{t_1, t_2, u} = I_{11} \cup I'_{11} \times I_9 \cup I'_9$$

so that

$$\int_{\mathbb{R}^2} \frac{1}{|y_1 y_2|} \chi_{I_{t_1, t_2, u}}(y_1, y_2) dy_1 dy_2 = \left(4 \log \frac{1}{u}\right)^2 \left(1 - \frac{\log |\frac{1}{t_1}|}{2 \log \frac{1}{u}}\right) \left(1 - \frac{\log |t_2|}{2 \log \frac{1}{u}}\right).$$

Case 9 : $t_1 \in (-\frac{1}{u^2}, -1), t_2 \in (-1, -u^2)$. In this case

$$I_{t_1, t_2, u} = I_{10} \cup I'_{10} \times I_{12} \cup I'_{12}, \quad \text{where } I_{12} = \left(-\frac{u}{t_2}, \frac{1}{u}\right)$$

so that

$$\int_{\mathbb{R}^2} \frac{1}{|y_1 y_2|} \chi_{I_{t_1, t_2, u}}(y_1, y_2) dy_1 dy_2 = \left(4 \log \frac{1}{u}\right)^2 \left(1 - \frac{\log |t_1|}{2 \log \frac{1}{u}}\right) \left(1 - \frac{\log |\frac{1}{t_2}|}{2 \log \frac{1}{u}}\right).$$

Case 10 : $t_1, t_2 \in (-\frac{1}{u^2}, -1)$. In this case, it can be worked out that

$$I_{t_1, t_2, u} = I_{10} \cup I'_{10} \times I_{13} \cup I'_{13}, \quad \text{where } I_{13} = \left(u, -\frac{1}{ut_2}\right)$$

and therefore,

$$\int_{\mathbb{R}^2} \frac{1}{|y_1 y_2|} \chi_{I_{t_1, t_2, u}}(y_1, y_2) dy_1 dy_2 = \left(4 \log \frac{1}{u}\right)^2 \left(1 - \frac{\log |t_1|}{2 \log \frac{1}{u}}\right) \left(1 - \frac{\log |t_2|}{2 \log \frac{1}{u}}\right).$$

Case 11 : $t_1, t_2 \in (-1, -u^2)$. In this case

$$I_{t_1, t_2, u} = I_{11} \cup I'_{11} \times I_{12} \cup I'_{12}$$

so that

$$\int_{\mathbb{R}^2} \frac{1}{|y_1 y_2|} \chi_{I_{t_1, t_2, u}}(y_1, y_2) dy_1 dy_2 = \left(4 \log \frac{1}{u}\right)^2 \left(1 - \frac{\log |\frac{1}{t_1}|}{2 \log \frac{1}{u}}\right) \left(1 - \frac{\log |\frac{1}{t_2}|}{2 \log \frac{1}{u}}\right).$$

Case 12 : $t_1 \in (-1, -u^2)$, $t_2 \in (-\frac{1}{u^2}, -1)$. In this case

$$I_{t_1, t_2, u} = I_{11} \cup I'_{11} \times I_{13} \cup I'_{13}$$

so that

$$\int_{\mathbb{R}^2} \frac{1}{|y_1 y_2|} \chi_{I_{t_1, t_2, u}}(y_1, y_2) dy_1 dy_2 = \left(4 \log \frac{1}{u}\right)^2 \left(1 - \frac{\log |\frac{1}{t_1}|}{2 \log \frac{1}{u}}\right) \left(1 - \frac{\log |t_2|}{2 \log \frac{1}{u}}\right).$$

Case 13 : $t_1 \in (u^2, 1)$, $t_2 \in (-1, -u^2)$. In this case, it can be worked out that

$$I_{t_1, t_2, u} = I_6 \cup I'_6 \times I_{12} \cup I'_{12}$$

and therefore,

$$\int_{\mathbb{R}^2} \frac{1}{|y_1 y_2|} \chi_{I_{t_1, t_2, u}}(y_1, y_2) dy_1 dy_2 = \left(4 \log \frac{1}{u}\right)^2 \left(1 - \frac{\log |\frac{1}{t_1}|}{2 \log \frac{1}{u}}\right) \left(1 - \frac{\log |\frac{1}{t_2}|}{2 \log \frac{1}{u}}\right).$$

Case 14 : $t_1 \in (1, \frac{1}{u^2})$, $t_2 \in (-1, -u^2)$. In this case

$$I_{t_1, t_2, u} = I_8 \cup I'_8 \times I_{12} \cup I'_{12}$$

so that

$$\int_{\mathbb{R}^2} \frac{1}{|y_1 y_2|} \chi_{I_{t_1, t_2, u}}(y_1, y_2) dy_1 dy_2 = \left(4 \log \frac{1}{u}\right)^2 \left(1 - \frac{\log |t_1|}{2 \log \frac{1}{u}}\right) \left(1 - \frac{\log |\frac{1}{t_2}|}{2 \log \frac{1}{u}}\right).$$

Case 15 : $t_1 \in (u^2, 1)$, $t_2 \in (-\frac{1}{u^2}, -1)$. In this case

$$I_{t_1, t_2, u} = I_6 \cup I'_6 \times I_{13} \cup I'_{13}$$

so that

$$\int_{\mathbb{R}^2} \frac{1}{|y_1 y_2|} \chi_{I_{t_1, t_2, u}}(y_1, y_2) dy_1 dy_2 = \left(4 \log \frac{1}{u}\right)^2 \left(1 - \frac{\log |\frac{1}{t_1}|}{2 \log \frac{1}{u}}\right) \left(1 - \frac{\log |t_2|}{2 \log \frac{1}{u}}\right).$$

Case 16 : $t_1 \in (1, \frac{1}{u^2})$, $t_2 \in (-\frac{1}{u^2}, -1)$. In this case

$$I_{t_1, t_2, u} = I_8 \cup I'_8 \times I_{13} \cup I'_{13}$$

so that

$$\int_{\mathbb{R}^2} \frac{1}{|y_1 y_2|} \chi_{I_{t_1, t_2, u}}(y_1, y_2) dy_1 dy_2 = \left(4 \log \frac{1}{u}\right)^2 \left(1 - \frac{\log |t_1|}{2 \log \frac{1}{u}}\right) \left(1 - \frac{\log |t_2|}{2 \log \frac{1}{u}}\right).$$

Combining the above information and taking f and g as f_u and g_u respectively in (3.1), we obtain that

$$\begin{aligned} \mathcal{J} &= \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{|\phi(t_1, t_2)|}{|t_1 t_2|^{2\alpha+2}} h_u(t_1, t_2) dt_1 dt_2 \\ &\geq \int_{\mathbb{R}^2} \frac{|\phi(t_1, t_2)|}{|t_1 t_2|^{2\alpha+2-\frac{1}{p}}} \inf_{(y_1, y_2) \in \mathbb{R}^2} \left(\frac{v(t_1 y_1, t_2 y_2)}{v(y_1, y_2)} \right)^{\frac{1}{p}} \int_{\mathbb{R}^2} \frac{1}{|y_1 y_2|} \chi_{I_{t_1, t_2, u}}(y_1, y_2) dy_1 dy_2 dt_1 dt_2 \\ &= \left(4 \log \frac{1}{u}\right)^2 \left(\int_{-\frac{1}{u^2}}^{-u^2} \int_{-\frac{1}{u^2}}^{-u^2} + \int_{u^2}^{\frac{1}{u^2}} \int_{u^2}^{\frac{1}{u^2}} + \int_{u^2}^{\frac{1}{u^2}} \int_{-\frac{1}{u^2}}^{-u^2} + \int_{-\frac{1}{u^2}}^{-u^2} \int_{u^2}^{\frac{1}{u^2}} \right) \left(\frac{|\phi(t_1, t_2)|}{|t_1 t_2|^{2\alpha+2-\frac{1}{p}}} \times \right. \\ &\quad \left. \times \inf_{(y_1, y_2) \in \mathbb{R}^2} \left(\frac{v(t_1 y_1, t_2 y_2)}{v(y_1, y_2)} \right)^{\frac{1}{p}} \left(1 - \frac{\xi(t_1)}{2 \log \frac{1}{u}}\right) \left(1 - \frac{\xi(t_2)}{2 \log \frac{1}{u}}\right) dt_1 dt_2 \right), \end{aligned} \quad (3.4)$$

where for $i = 1, 2$

$$\xi(t_i) = \begin{cases} \log |\frac{1}{t_i}|, & \text{if } t_i \in (u^2, 1] \cup (-1, -u^2] \\ \log |t_i|, & \text{if } t_i \in (1, 1/u^2] \cup (-1/u^2, -1). \end{cases}$$

Now, by using the test functions f_u, g_u in (3.2) and using (3.3), (3.4), we get

$$\begin{aligned} &\left(\int_{-\frac{1}{u^2}}^{-u^2} \int_{-\frac{1}{u^2}}^{-u^2} + \int_{u^2}^{\frac{1}{u^2}} \int_{u^2}^{\frac{1}{u^2}} + \int_{u^2}^{\frac{1}{u^2}} \int_{-\frac{1}{u^2}}^{-u^2} + \int_{-\frac{1}{u^2}}^{-u^2} \int_{u^2}^{\frac{1}{u^2}} \right) \left(\frac{|\phi(t_1, t_2)|}{|t_1 t_2|^{2\alpha+2-\frac{1}{p}}} \inf_{(y_1, y_2) \in \mathbb{R}^2} \left(\frac{v(t_1 y_1, t_2 y_2)}{v(y_1, y_2)} \right)^{\frac{1}{p}} \times \right. \\ &\quad \left. \times \left(1 - \frac{\xi(t_1)}{2 \log \frac{1}{u}}\right) \left(1 - \frac{\xi(t_2)}{2 \log \frac{1}{u}}\right) dt_1 dt_2 \right) \leq \|H_{\alpha, \phi}\|. \end{aligned}$$

By the Monotone Convergence Theorem and on taking $u \rightarrow 0$ we have

$$\int_{\mathbb{R}^2} \frac{|\phi(t_1, t_2)|}{|t_1 t_2|^{2\alpha+2-\frac{1}{p}}} \inf_{(y_1, y_2) \in \mathbb{R}^2} \left(\frac{v(t_1 y_1, t_2 y_2)}{v(y_1, y_2)} \right)^{\frac{1}{p}} dt_1 dt_2 \leq \|H_{\alpha, \phi}\|$$

and we are done. \square

On the lines of Theorem 2.3, a characterization of the boundedness of $\mathcal{H}_{\alpha,\phi} : L_v^p(\mathbb{R}^2) \rightarrow L_v^p(\mathbb{R}^2)$ can be obtained. Precisely, we have the following

Theorem 3.4. *Let $1 < p < \infty$, v be weight function and $\phi \in L^1(\mathbb{R}^2)$. Let the following be satisfied for some constant $c > 0$:*

$$\sup_{(y_1,y_2) \in \mathbb{R}^2} \left(\frac{v(t_1 y_1, t_2 y_2)}{v(y_1, y_2)} \right) \leq c \inf_{(y_1,y_2) \in \mathbb{R}^2} \left(\frac{v(t_1 y_1, t_2 y_2)}{v(y_1, y_2)} \right).$$

Then the operator $\mathcal{H}_{\alpha,\phi} : L_v^p(\mathbb{R}^2) \rightarrow L_v^p(\mathbb{R}^2)$ is bounded if and only if $\mathcal{A}_{\text{sup}} < \infty$ and moreover the following estimates hold:

$$\frac{1}{c} \mathcal{A}_{\text{sup}} \leq \|\mathcal{H}_{\alpha,\phi}\|_{L_v^p(\mathbb{R}^2) \rightarrow L_v^p(\mathbb{R}^2)} \leq \mathcal{A}_{\text{sup}}.$$

Corollary 3.5. *Let $1 < p < \infty$, v be weight function and $\phi \in L^1(\mathbb{R}^2)$. If there exists a function h such that $v(x_1 y_1, x_2 y_2) = v(x_1, x_2) h(y_1, y_2)$, then $\mathcal{A}_{\text{sup}} = \mathcal{A}_{\text{inf}}$ and*

$$\|\mathcal{H}_{\alpha,\phi}\|_{L_v^p(\mathbb{R}^2) \rightarrow L_v^p(\mathbb{R}^2)} = \int_{\mathbb{R}^2} \frac{|\phi(t_1, t_2)|}{|t_1 t_2|^{2\alpha+2-\frac{1}{p}}} h^{1/p}(t_1, t_2) dt_1 dt_2.$$

Remark 3.6. For $\alpha = -1/2$, $\mathcal{H}_{\alpha,\phi}$ reduces to the two dimensional Hausdorff operator

$$\mathcal{H}_{\phi} f(x_1, x_2) = \int_{\mathbb{R}^2} \frac{|\phi(t_1, t_2)|}{|t_1 t_2|} f\left(\frac{x_1}{t_1}, \frac{x_2}{t_2}\right) dt_1 dt_2$$

which on replacement $\phi(s_1, s_2) = \frac{1}{s_1 s_2} \psi\left(\frac{1}{s_1}, \frac{1}{s_2}\right)$ becomes equivalent to

$$\mathcal{G}_{\psi} g(x_1, x_2) = \int_{\mathbb{R}^2} \psi\left(\frac{t_1}{x_1}, \frac{t_2}{x_2}\right) g(t_1, t_2) dt_1 dt_2.$$

The $L^p(\mathbb{R}^+ \times \mathbb{R}^+)$ -boundedness of \mathcal{G}_{ψ} (consequently of \mathcal{H}_{ψ}) was proved in [10]. Moreover, if we take

$$\phi(t_1, t_2) = \frac{1}{t_1 t_2} \chi_{(1,\infty)}(t_1) \chi_{(1,\infty)}(t_2)$$

then $\mathcal{H}_{\psi} f$ becomes the two-dimensional Hardy operator [18]

$$H_2 f(x_1, x_2) = \frac{1}{x_1 x_2} \int_0^{x_1} \int_0^{x_2} f(t_1, t_2) dt_1 dt_2.$$

4 Some generalizations

In this section, we shall prove generalizations of some of the theorems proved in the previous section. To begin with, the following theorem is a two-weight generalization of Theorem 2.1

Theorem 4.1. *Let $1 < p < \infty$, v, w be weight functions and $\phi \in L^1(\mathbb{R})$ be such that*

$$\int_{\mathbb{R}} \frac{|\phi(t)|}{|t|^{2\alpha+2-\frac{1}{p}}} \left(\sup_{y \in \mathbb{R}} \frac{v(ty)}{w(y)} \right)^{1/p} dt =: B_{\text{sup}} < \infty.$$

Then the operator $H_{\alpha,\phi} : L_w^p(\mathbb{R}) \rightarrow L_v^p(\mathbb{R})$ is bounded and

$$\|H_{\alpha,\phi} f\|_{L_v^p(\mathbb{R})} \leq B_{\text{sup}} \|f\|_{L_w^p(\mathbb{R})}.$$

Proof. It follows on the similar lines as that of the proof of Theorem 2.1 by replacing vv^{-1} by ww^{-1} in (2.1). \square

Theorem 4.1 has the following version for two indices p, q :

Theorem 4.2. *Let $1 < q < p < \infty$, v, w be weight functions and $\phi \in L^1(\mathbb{R})$ be such that*

$$\int_{\mathbb{R}} \frac{|\phi(t)|}{|t|^{2\alpha+2-\frac{1}{p}}} \left(\int_{\mathbb{R}} \frac{[v(ty)]^{\frac{p}{p-q}}}{[w(y)]^{\frac{q}{p-q}}} dy \right)^{\frac{p-q}{pq}} dt =: D_{\text{sup}} < \infty.$$

Then the operator $H_{\alpha, \phi} : L_w^p(\mathbb{R}) \rightarrow L_v^q(\mathbb{R})$ is bounded and

$$\|H_{\alpha, \phi} f\|_{L_v^q(\mathbb{R})} \leq D_{\text{sup}} \|f\|_{L_w^p(\mathbb{R})}.$$

Proof. Let $f \in L_w^p(\mathbb{R})$ then by using generalised Minkowski inequality, change of variables, Hölder's inequality for $p/q > 1$ and , we have

$$\begin{aligned} \|H_{\alpha, \phi} f\|_{L_v^q(\mathbb{R})} &= \left(\int_{\mathbb{R}} |H_{\alpha, \phi} f(x)|^q v(x) dx \right)^{\frac{1}{q}} \\ &= \left(\int_{\mathbb{R}} \left| \int_{\mathbb{R}} \frac{|\phi(t)|}{|t|^{2\alpha+2}} f\left(\frac{x}{t}\right) dt \right|^q v(x) dx \right)^{\frac{1}{q}} \\ &\leq \left(\int_{\mathbb{R}} \left(\int_{\mathbb{R}} \frac{|\phi(t)|}{|t|^{2\alpha+2}} \left| f\left(\frac{x}{t}\right) \right| v^{\frac{1}{q}}(x) dt \right)^q dx \right)^{\frac{1}{q}} \\ &\leq \int_{\mathbb{R}} \frac{|\phi(t)|}{|t|^{2\alpha+2}} |t|^{\frac{1}{q}} \left(\int_{\mathbb{R}} |f(y)|^q v(yt) dy \right)^{\frac{1}{q}} dt \\ &= \int_{\mathbb{R}} \frac{|\phi(t)|}{|t|^{2\alpha+2-\frac{1}{q}}} \left(\int_{\mathbb{R}} |f(y)|^q w^{\frac{q}{p}}(y) w^{-\frac{q}{p}}(y) v(yt) dy \right)^{\frac{1}{q}} dt \\ &\leq \left(\int_{\mathbb{R}} \frac{|\phi(t)|}{|t|^{2\alpha+2-\frac{1}{q}}} \left(\int_{\mathbb{R}} \frac{[v(ty)]^{\frac{p}{p-q}}}{[w(y)]^{\frac{q}{p-q}}} dt \right)^{\frac{p-q}{pq}} dt \right) \left(\int_{\mathbb{R}} |f(y)|^p w(y) dy \right)^{\frac{1}{p}} \\ &= D_{\text{sup}} \|f\|_{L_w^p(\mathbb{R})} \end{aligned}$$

and the assertion follows. \square

Two dimensional versions of Theorems 4.1 and 4.2 can also be proved. We only state the results.

Theorem 4.3. *Let $1 < p < \infty$, v, w be weight functions and $\phi \in L^1(\mathbb{R}^2)$ be such that*

$$\int_{\mathbb{R}^2} \frac{|\phi(t_1, t_2)|}{|t_1 t_2|^{2\alpha+2-\frac{1}{p}}} \left(\sup_{(y_1, y_2) \in \mathbb{R}^2} \frac{v(t_1 y_1, t_2 y_2)}{w(y_1, y_2)} \right)^{1/p} dt_1 dt_2 =: \mathcal{B}_{\text{sup}} < \infty.$$

Then the operator $\mathcal{H}_{\alpha, \phi} : L_w^p(\mathbb{R}^2) \rightarrow L_v^p(\mathbb{R}^2)$ is bounded and

$$\|\mathcal{H}_{\alpha, \phi} f\|_{L_v^p(\mathbb{R}^2)} \leq \mathcal{B}_{\text{sup}} \|f\|_{L_w^p(\mathbb{R}^2)}.$$

Theorem 4.4. Let $1 < q < p < \infty$, v, w be weight functions and $\phi \in L^1(\mathbb{R}^2)$ be such that

$$\int_{\mathbb{R}^2} \frac{|\phi(t_1, t_2)|}{|t_1 t_2|^{2\alpha+2-\frac{1}{p}}} \left(\int_{\mathbb{R}^2} \frac{[v(t_1 y_1, t_2 y_2)]^{\frac{p}{p-q}}}{[w(y_1, y_2)]^{\frac{q}{p-q}}} dt_1 dt_2 \right)^{\frac{p-q}{p}} dt_1 dt_2 =: \mathcal{D}_{\text{sup}} < \infty.$$

Then the operator $\mathcal{H}_{\alpha, \phi} : L_w^p(\mathbb{R}^2) \rightarrow L_v^q(\mathbb{R}^2)$ is bounded and

$$\|\mathcal{H}_{\alpha, \phi} f\|_{L_v^q(\mathbb{R}^2)} \leq \mathcal{D}_{\text{sup}} \|f\|_{L_w^p(\mathbb{R}^2)}.$$

Acknowledgement. The third author acknowledges the MATRICS Research Grant No. MTR/2017/000126 of SERB, Department of Science and Technology, India.

References

- [1] C. Abdelkefi, M. Rachdi, *Some results for the Dunkl transform and the generalized Cesaro operator*, 2012, Available from: <https://arxiv.org/abs/1208.5034>.
- [2] J. Chen, D. Fan, J. Li, *Hausdorff operators on function spaces*, Chin. Ann. Math. Ser. B, 33 (2013), 537–556.
- [3] R. Bandaliyev, P. Górká, *Hausdorff operator in Lebesgue spaces*, Math. Inequal. Appl., 22 (2019), 657–676.
- [4] R. Daher, F. Saadi, *The Dunkl-Hausdorff operator is bounded on the real Hardy space $H_\alpha^1(\mathbb{R})$* , Integral Transforms and Special Functions, 30 (2019), 882–892.
- [5] R. Daher, F. Saadi, *Boundedness of Multidimensional Dunkl-Hausdorff Operators*, Hindawi, J. Function Spaces, 2020, Available from: <https://doi.org/10.1155/2020/1071>.
- [6] R. Daher, F. Saadi, *The Dunkl-Hausdorff Operators and the Dunkl continuous wavelets transform*, J. Pseudo-Differ. Oper. Appl., 2020, Available from: <https://doi.org/10.1007/s11868-020-00351-1>.
- [7] C. F. Dunkl, *Differential-difference operators associated to reflection groups*, Trans. Amer. Math. Soc., 311 (1989), 167–183.
- [8] R. R. Goldberg, *Certain operators and Fourier transforms on L^2* , Proc. Amer. Math. Soc., 10 (1959), 385–390.
- [9] G. H. Hardy, J. E. Littlewood and G. Pólya, *Inequalities*, Cambridge University Press, 1934, xii+324 pp.
- [10] P. Jain, S. Jain, V. D. Stepanov, *LCT based integral transforms and Hausdorff operators*, Eurasian Math. J., 11 (2020), 57–71.
- [11] Y. Kanjin, *The Hausdorff operators on the real Hardy spaces $H_p(\mathbb{R})$* , Studia Math., 148 (2001), 37–45.
- [12] N. K. Karapetiants and S. G. Samko, *Equations with involutive operators*, Birkhäuser Boston, Inc., Boston, MA, 2001. xxiv+427 pp
- [13] E. Lifyand, *Hausdorff operators on Hardy spaces*, Eurasian Math. J., 4 (2013), 101–141.

- [14] E. Liflyand, A. Miyachi, *Boundedness of the Hausdorff operators in H_p spaces, $0 < p < 1$* , Studia Math., 194 (2009), 279–292.
- [15] E. Liflyand, F. Móricz, *The Hausdorff operator is bounded on the real Hardy spaces $H^1(\mathbb{R})$* , Proc. Amer. Math. Soc., 128 (1999), 1391–1396.
- [16] A. Kufner, L. Maligranda and L. E. Persson. *The Hardy inequality, About its history and some related results*. Pilsen 2007.
- [17] A. Kufner, L. Maligranda and L. E. Persson. *The Prehistory of the Hardy inequality*, Amer. Math. Monthly 113, 8 (2006) 715–732.
- [18] E. Sawyer, *Weighted inequalities for two-dimensional Hardy operator*, Studia Math., 82 (1985), 1–16.