On SO(N) Spin Vertex Models

VLADIMIR BELAVIN a , DORON GEPNER b , HANS WENZL c

- ^a Physics Department, Ariel University, Ariel 40700, Israel
 ^b Department of Particle Physics and Astrophysics, Weizmann Institute,
- Rehovot 76100, Israel
 ^c Department of Mathematics, University of California, San Diego, California

ABSTRACT

We describe the Boltzmann weights of the D_k algebra spin vertex models. Thus, we find the SO(N) spin vertex models, for any N, completing the B_k case found earlier. We further check that the real (self-dual) SO(N) models obey quantum algebras, which are the Birman-Murakami-Wenzl (BMW) algebra for three blocks, and certain generalizations, which include the BMW algebra as a sub-algebra, for four and five blocks. In the case of five blocks, the B_4 model is shown to satisfy additional twenty new relations, which are given. The D_6 model is shown to obey two additional relations.

1. Introduction.

Solvable lattice models in two dimensions are a fruitful ground to test phase transitions, universality, integrability [1] and conformal field theory [2]. For reviews see [3, 4].

We will concentrate here on a type of solvable lattice models which are called vertex models. Well known among these are the six, eight and nineteen vertex models [3, 4]. For recent works on vertex models see, e.g., [5, 6, 7, 8, 9]. Our purpose here is to introduce vertex models based on the algebra D_k and the spin representation. This completes the SO(N) spin vertex models for all N, where the B_k models were described before in ref. [10].

We are also interested in the algebraic structure underlying these models. We use the more general results of [11,12], which describe the three, four and five blocks algebras (where the number of blocks is the degree of polynomial equation obeyed by the Boltzmann weights), assuming only a certain ansatz for the Baxterization, described in [13], and the Yang-Baxter equation. We describe and check numerically, the algebras of B_4 , which is a five blocks theory, and the algebra of D_6 , which is a four blocks theory.

The algebras include a version of the Birman–Murakami–Wenzl algebra (BMW) [14, 15], along with two new relations for four blocks and twenty new relations for the five blocks theory, which are given here for B_4 . We check that the BMW algebra is obeyed for D_k , for any small even k, with a different skein relation.

2. D_k spin vertex models.

We wish to describe a vertex model based on the algebra $D_k = SO(2k)$ and the spin representation. This solution is an element of $\operatorname{End}(V \otimes V)$ where V is the spin representation of D_k . We denote by $\alpha_n = \epsilon_n - \epsilon_{n+1}$, for n = 1, 2, ..., k-1 and $\alpha_k = \epsilon_{k-1} + \epsilon_k$ the simple roots of D_k , where ϵ_i are orthogonal unit vectors. The spin representation, denoted by S has the weights $\sum_{i=1}^k p_i \epsilon_i/2$, where $p_i = \pm 1$ and $\prod_{i=1}^k p_i = 1$. The last product is -1 for the anti-spinor representation, denoted by \bar{S} . We find it useful to add 1/2 to these weights, and to represent weights of the spinor (anti-spinor) representation by the vector m, where $m_i = 0$ or 1.

To start constructing the vertex model, we need a solution which commutes with the co-product of $U_{q^2}(SO(2k))$. We find it convenient to first describe a solution for the larger representation $\tilde{V} = S \oplus \bar{S}$, namely the sum of the spinor and anti-spinor representations. Such a solution was described recently in [16]. It is the element C of $\operatorname{End}(\tilde{V} \otimes \tilde{V})$, given by

$$C_{m,n}^{b,c} = \sum_{j=1}^{k} \delta_{m_j,1-n_j} (-q^2)^{\{m-n\}_j} \delta_{b,\bar{m}_j} \delta_{c,\bar{n}_j}, \tag{2.1}$$

where

$$\{m\}_j = \sum_{r=1}^j m_r, \tag{2.2}$$

and \bar{n}_j is equal to n_j except at the jth coordinate where it is $1 - n_j$. Here m, n, b, c = 0 or 1 are weights of the spin or anti–spin representations shifted by 1/2. The eigenvalues of the matrix C were computed in ref. [16], and are

$$\lambda_j = \pm s(k - j), \quad \text{for } j = 0, 1, \dots, k,$$
 (2.3)

where

$$s(x) = \frac{q^{2x} - q^{-2x}}{q^2 - q^{-2}}. (2.4)$$

The solution C has the disadvantage of mapping both the spin and anti-spin

representations. We note, however, that C maps the representation $S \otimes S$ to $\bar{S} \otimes \bar{S}$, and vice versa. Thus, to get a solution in $\operatorname{End}(S \otimes S)$ all we need to do is to square the matrix C and to equate to zero all the $C_{m,n}^{b,c}$ for weights m, n, b, c which are not in S. Thus, C^2 gives the solution we want. Of course, since C commutes with the co-product, so does C^2 .

Since the matrix C^2 commutes with the co-product, it has the same eigenvectors as our desired solution which obeys the Yang-Baxter equation, but not the same eigenvalues. Thus, we define the projection operators

$$(P^{a})_{m,n}^{b,c} = \prod_{p \neq a} \left[\frac{C^{2} - \lambda_{p}^{2} I}{\lambda_{a}^{2} - \lambda_{p}^{2}} \right], \tag{2.5}$$

where the product is in $\operatorname{End}(S \otimes S)$ and I is the identity map.

We note that for even k, $P^a = 0$ for a which is odd, whereas for odd k, $P^a = 0$ for a which is even. The jth eigenvalue corresponds to the representation $V_j = \wedge^j v$, where v is the vector representation, i.e., the anti-symmetric product of j vector representations [16]. The highest weight of the representation V_j is $\epsilon_1 + \epsilon_2 + \ldots + \epsilon_j$. Thus, the non-zero P^a are in one to one correspondence with the representations that appear in the tensor product,

$$S \times S = \sum_{\substack{j=0\\j=k \bmod 2}}^{k} V_j, \tag{2.6}$$

as they should. Thus, the projection P^a projects onto the representation V_a .

We wish to make the connection between the solution C^2 and the D_k WZW conformal model. For explanation of conformal field theory see the book [2], and references therein. To do this we define,

$$q^2 = \exp[\pi i/(r+g)],$$
 (2.7)

Here r is the level of the representation and

$$g = 2k - 2, (2.8)$$

is the dual Coxeter number.

The dimension of the highest weight Λ in a WZW theory is given by

$$\Delta_{\Lambda} = \frac{\Lambda(\Lambda + 2\rho)}{2(r+g)},\tag{2.9}$$

where ρ is half the sum of positive roots and $C_{\Lambda} = \Lambda(\Lambda + 2\rho)$ is the Casimir of the representation. The Casimir of the representation V_j is given by

$$C(V_j) = C_j = j(2k - j).$$
 (2.10)

As explained in [10], the eigenvalues of the R matrix are given by

$$\beta_j = p_j e^{-i\pi\Delta_j} = p_j q^{-C(V_j)},$$
(2.11)

where $p_j = \pm 1$ is some sign which corresponds to whether the product in eq. (2.6) is symmetric or anti-symmetric. In our case, the sign is given by

$$p_j = (-1)^{(k-j)/2}. (2.12)$$

Thus, since we know the eigenvalues of the R matrix and the projection operators from eq. (2.5), we may construct the R matrix as

$$R_{m,n}^{a,b} = \sum_{j=0}^{k} \beta_j (P^j)_{m,n}^{a,b}.$$
 (2.13)

It can be verified that this R matrix satisfies the Yang–Baxter equation (YBE) which for the R matrix is the braiding relation,

$$\sum_{\alpha,\beta,\gamma} R_{j,k}^{\beta,\alpha} R_{i,\beta}^{l,\gamma} R_{\gamma,\alpha}^{m,n} = \sum_{\alpha,\beta,\gamma} R_{i,j}^{\alpha,\beta} R_{\beta,k}^{\gamma,n} R_{\alpha,\gamma}^{l,m}.$$
 (2.14)

We checked that this R matrix obeys the YBE, numerically for k = 2, 3, 4, 5, 6 and it holds, indeed, for various weights and for general q.

Now, we wish to define a trigonometric solution for the YBE. For this purpose, we use the same general ansatz for Baxterization as in [10, 13]. First, we need to decide on the order of the primary fields in eq. (2.6). The order which solves the YBE is given by

$$(h_0, h_1, \dots, h_{k/2}) = (0, 2, 4, \dots, k),$$
 (2.15)

for even k. For odd k the order is

$$(h_0, h_1, \dots, h_{(k-1)/2}) = (k, k-2, k-4, \dots, 1).$$
(2.16)

The parameters are given by [10, 13],

$$\hat{\zeta}_j = \pi (\Delta_{h_{j+1}} - \Delta_{h_j})/2, \tag{2.17}$$

for j = 0, 1, ..., m - 1, where m = k/2 for even k and m = (k - 1)/2 for odd k. Thus, the D_k theory is an m + 1 blocks theory. We thus define the parameters as

$$\zeta_j = (C_{h_{j+1}} - C_{h_j})/2. \tag{2.18}$$

We define

$$p(x) = q^x - q^{-x}. (2.19)$$

Then the trigonometric solution to the YBE assumes the form [10, 13],

$$R_{m,n}^{a,b}(u) = \sum_{j=0}^{m} f_j(u) (P^{h_j})_{m,n}^{a,b}, \qquad (2.20)$$

where

$$f_a(u) = \left[\prod_{j=1}^a p(\zeta_{j-1} - u)\right] \left[\prod_{j=a+1}^m p(\zeta_{j-1} + u)\right] / \left[\prod_{j=1}^m p(\zeta_{j-1})\right], \quad (2.21)$$

where a = 0, 1, ..., m.

For example, for k = 6, which is a four blocks theory, the parameters are $(\zeta_0, \zeta_1, \zeta_2) = (10, 6, 2)$. The crossing parameter is $\lambda = \zeta_0$. The D_k vertex models with k even are real (self-dual) as $S = S^*$. For odd k the theories are not real (not self-dual), as $S \neq S^*$.

We can check that the solution, eqs. (2.20, 2.21), obeys the Yang–Baxter equation, which is

$$\sum_{\alpha,\beta,\gamma} R_{j,k}^{\beta,\alpha}(u) R_{i,\beta}^{l,\gamma}(u+v) R_{\gamma,\alpha}^{m,n}(v) = \sum_{\alpha,\beta,\gamma} R_{i,j}^{\alpha,\beta}(v) R_{\beta,k}^{\gamma,n}(u+v) R_{\alpha,\gamma}^{l,m}(u). \tag{2.22}$$

We checked this equation, numerically, for k = 2, 3, 4, 5, 6 and various values of u, v and q and various heights. It is indeed obeyed. This gives the trigonometric D_k spin vertex model.

3. BMW' algebra and SO(N) spin vertex models.

We repeat here the definition of the BMW' algebra following [10]. We find it convenient to use an operator form for the R matrix. We define the matrix, following [4],

$$X_{i}(u) = \sum_{m,n,a,b} R_{m,n}^{a,b}(u) I^{(1)} \otimes \ldots \otimes I^{(i-1)} e_{am}^{(i)} \otimes e_{bn}^{(i+1)} \otimes I^{(i+2)} \otimes \ldots \otimes I^{(f)}, (3.1)$$

where $I^{(i)}$ is the identity matrix at position i and $(e_{rs})_{lm} = \delta_{rl}\delta_{sm}$. The YBE, eq. (2.22), then assumes a more compact form,

$$X_i(u)X_j(v) = X_j(v)X_i(u),$$
 if $|i-j| \ge 2$,

$$X_i(u)X_{i+1}(u+v)X_i(v) = X_{i+1}(v)X_i(u+v)X_{i+1}(u).$$
(3.2)

Let us denote the number of blocks by n. For the D_k models, this is n = m + 1 = k/2 + 1 (k even), or n = m + 1 = (k + 1)/2 (for odd k). In this section,

we will assume that k is even, so that the theory is real (self-dual). It is assumed that the number of blocks is greater or equal to three, $n \geq 3$. The algebras of non-real theories are also interesting, but we shall not describe it here. We define the limit of the matrix $X_i(u)$ as

$$X_i = \lim_{u \to i\infty} e^{i(n-1)u} X_i(u), \qquad X_i^t = \lim_{u \to -i\infty} e^{-i(n-1)u} X_i(u).$$
 (3.3)

We define the operators,

$$G_i = 2^{n-1} e^{-i(n-1)\zeta_0/2} \left[\prod_{r=1}^{n-1} \sin(\zeta_{r-1}) \right] X_i,$$
 (3.4)

$$G_i^{-1} = 2^{n-1} e^{i(n-1)\zeta_0/2} \left[\prod_{r=1}^{n-1} \sin(\zeta_{r-1}) \right] X_i^t, \tag{3.5}$$

and

$$E_i = X_i(\zeta_0), \qquad 1_i = X_i(0),$$
 (3.6)

where ζ_i are the parameters defined in eq. (2.17). G_i^{-1} , so defined, is the inverse of G_i , or $G_iG_i^{-1}=1_i$.

From the ansatz, eqs. (2.20, 2.21), and from the YBE, eq. (2.22), we can prove the following relations of the operators G_i , G_i^{-1} and E_i ,

$$E_i E_{i+1} E_i = b E_i, \qquad E_i^2 = b E_i, \qquad E_i E_j = E_j E_i \quad \text{if } |i-j| \ge 2,$$
 (3.7)

$$b = \prod_{r=1}^{n-1} \frac{\sin(\zeta_0 + \zeta_{r-1})}{\sin(\zeta_{r-1})},$$
(3.8)

which is the Temperley-Lieb algebra [17], and

$$G_i G_j = G_j G_i$$
 if $|i - j| \ge 2$, $G_i G_{i+1} G_i = G_{i+1} G_i G_{i+1}$, (3.9)

which is the braiding algebra. We can also prove the relations,

$$G_i E_i = E_i G_i = l^{-1} E_i, (3.10)$$

where

$$l = i^{n-1} \exp\left[i(n-1)\zeta_0/2 + i\sum_{r=0}^{n-2}\zeta_r\right].$$
 (3.11)

The following is the skein relation which stems from the definition of the projection operators along with the ansatz, eqs. (2.20, 2.21),

$$G_i^{n-2} = aE_i + \sum_{r=-1}^{n-3} b_r G_i^r, \tag{3.12}$$

where a and b_r are some coefficients, which can be expressed in terms of the parameters, ζ_r . From the skein relation we prove,

$$G_{i\pm 1}G_iE_{i\pm 1} = E_iG_{i\pm 1}G_i. (3.13)$$

The above relation, eqs. (3.7–3.13), are part of the Birman–Murakami–Wenzl algebra (BMW) [14,15]. The rest of the relations of the BMW algebra are also obeyed, except of the skein relation, eq. (3.12), which is different for more than three blocks. These are

$$G_{i\pm 1}G_iE_{i\pm 1} = E_iE_{i\pm 1}, \qquad G_{i\pm 1}E_iG_{i\pm 1} = G_i^{-1}E_{i\pm 1}G_i^{-1},$$
 (3.14)

$$G_{i\pm 1}E_iE_{i\pm 1} = G_i^{-1}E_{i\pm 1}, \qquad E_{i\pm 1}E_iG_{i\pm 1} = E_{i\pm 1}G_i^{-1},$$
 (3.15)

$$E_i G_{i\pm 1} E_i = l E_i, \qquad E_i G_{i+1}^{-1} E_i = l^{-1} E_i.$$
 (3.16)

We have verified that the full BMW' algebra (BMW with a different skein relation) is obeyed by the D_k model, with k = 4 or 6. We did this numerically,

using various heights and general q. We note that the BMW' algebra is obeyed also by the B_k spin vertex models [10], while substituting the relevant parameters ζ_r . For more than three blocks there are additional relations, except from the skein relation. These are described in the next two sections, for B_4 and D_6 .

4. n = 5 blocks case and B_4 vertex models.

It was noticed in [10] that the structure of n-CB algebra, which follows from the Baxterization of IRF models, is also applicable for vertex models. In our case, the connection is between n-CB algebra and B_{n-1} models. This algebra was checked, in particular, for B_3 models obeying 4-CB algebra [10]. In the 5-block case, the n-CB algebra reduces to a set of 20 relations in addition to BMW' sub-algebra. We get these relations by expanding the YBE, eq. (2.22) and assuming the ansatz, eqs. (2.20, 2.21). The 5-CB relations are general for all the five blocks models obeying the ansatz for Baxterization, eq. (2.20, 2.21). We specify the algebra here only for the B_4 spin vertex models for calculation reasons. In this section we summarize the 5-CB relations for B_4 spin vertex model. We give the shorter relations explicitly, here. The complete list of 5-CB relations for B_4 models can be found in the attached Mathematica file.

For the general values of the parameters, the 5-CB skein relation as well as the explicit projectors have been found in [12]. The skein relation reads

$$G_i^3 = \alpha 1_i + \beta E_i + \gamma G_i + \delta G_i^{-1} + \mu G_i^2 , \qquad (4.1)$$

where denoting $s_k = q^{\zeta_k}$ the parameters are

$$\alpha = -\frac{s_1 \left(s_1^2 s_2^2 s_3^2 - s_2^2 s_3^2 + s_3^2 - 1\right)}{s_0^3 s_2 s_3^3} ,$$

$$\beta = \frac{\left(s_1^2 - 1\right) \left(s_2^2 - 1\right) \left(s_0^2 s_1^2 s_2^2 + 1\right) \left(s_3^2 - 1\right) \left(s_0^2 s_1^2 s_2^2 s_3^2 - 1\right)}{s_0^5 s_1^3 s_2^3 \left(s_0^2 s_2^2 - 1\right) s_3^3 \left(s_0^2 s_3^2 - 1\right)} ,$$

$$\gamma = \frac{s_1^2 s_3^2 s_2^4 + s_1^2 s_2^2 - s_1^2 s_3^2 s_2^2 + s_3^2 s_2^2 - s_2^2 + 1}{s_0^2 s_2^2 s_3^2} ,$$

$$\delta = -\frac{s_1^2}{s_0^4 s_2^2} , \quad \mu = \frac{-s_2^2 s_1^2 + s_2^2 s_3^2 s_1^2 + s_1^2 - 1}{s_0 s_1 s_2 s_3} .$$

$$(4.2)$$

In the case of B_4 models the crossing parameters are

$$\zeta_0 = 7, \quad \zeta_1 = 3, \quad \zeta_2 = -1, \quad \zeta_3 = -5$$
 (4.3)

Using the results of [12] with the above explicit parameters the desired 5-CB algebra relations for B_4 models can be found. The B_4 skein relation reads explicitly

$$G_{i}^{3} = \frac{\left(q^{16} + q^{12} - q^{10} - q^{6} + q^{4} + 1\right)}{q^{14}}G_{i} - \frac{1}{q^{12}}G_{i}^{-1} + \frac{\left(q^{2} - 1\right)\left(q^{4} + 1\right)\left(q^{6} - q^{2} - 1\right)}{q^{10}}G_{i}^{2} + \frac{\left(q^{12} - q^{6} - q^{2} + 1\right)}{q^{14}}1_{i} + \frac{\left(q^{4} + 1\right)\left(q^{4} - q^{3} + q^{2} - q + 1\right)\left(q^{4} + q^{3} + q^{2} + q + 1\right)\left(q^{12} - q^{6} + 1\right)\left(q^{2} - 1\right)^{2}}{q^{38}}E_{i}.$$

To shorten notation, we denote below by $a_{l,m,n}$ the elements of the algebra $A_l[i]A_m[i+1]A_n[i]$ and by $b_{l,m,n}$ the elements of the algebra $A_l[i+1]A_m[i]A_m[i]A_n[i+1]$, where $A_l[r]$ stands for $G_r, G_r^{-1}, E_r, G_r^2$ or 1_r according to whether l = 1, 2, 3, 4, 5, respectively. A few of the relations, which are sufficiently short, are listed below:

1)
$$\frac{\left(q^{12} - q^{10} - q^6 + q^4 + 1\right) a_{5,2,3}}{q^{10}} + \frac{\left(q^{12} - q^{10} - q^6 + q^4 + 1\right) a_{5,3,1}}{q^{10}} - \frac{\left(q^{12} - q^{10} - q^6 + q^4 + 1\right) a_{5,1,3}}{q^{10}} - \frac{\left(q^{12} - q^{10} - q^6 + q^4 + 1\right) a_{5,3,2}}{q^{10}} - \frac{\left(q^{10} + q^6 - q^4 + q^2 - 1\right) \left(q^{12} - q^6 + 1\right) a_{5,5,3}}{q^{20}} - a_{4,3,3} - a_{5,3,4} + a_{5,4,3} + \frac{\left(q^{10} + q^6 - q^4 + q^2 - 1\right) \left(q^{12} - q^6 + 1\right) b_{5,5,3}}{q^{20}} + b_{3,3,4} = 0$$

2)
$$\frac{\left(q^{4}+1\right) \left(q^{12}-q^{6}+1\right) a_{5,5,3}}{q^{20}} - \frac{\left(q^{2}-1\right) \left(q^{4}+1\right) \left(q^{6}+q^{4}-1\right) b_{5,2,3}}{q^{20}} + \frac{\left(q^{2}-1\right) \left(q^{4}+1\right) \left(q^{6}+q^{4}-1\right) b_{5,3,2}}{q^{20}} + \frac{b_{5,4,3}-b_{5,3,4}}{q^{6}} - \frac{\left(q^{4}+1\right) \left(q^{12}-q^{6}+1\right) b_{5,5,3}}{q^{20}} + \frac{\left(q^{2}-1\right) \left(q^{4}+1\right) \left(q^{6}-q^{2}-1\right) b_{5,3,1}}{q^{16}} - \frac{\left(q^{2}-1\right) \left(q^{4}+1\right) \left(q^{6}-q^{2}-1\right) b_{5,1,3}}{q^{16}} + b_{1,4,3} - a_{3,4,1} = 0$$

3)
$$\frac{\left(q^{10} + q^6 - q^4 + q^2 - 1\right)\left(q^{12} - q^6 + 1\right)\left(a_{5,4,3} - a_{5,3,4} + b_{5,3,4} - b_{5,4,3}\right)}{q^{10}\left(q^{12} - q^{10} - q^6 + q^4 + 1\right)} - \frac{\left(q^{10} + q^6 - q^4 + q^2 - 1\right)^2\left(q^{12} - q^6 + 1\right)^2\left(a_{5,5,3} - b_{5,5,3}\right)}{q^{30}\left(q^{12} - q^{10} - q^6 + q^4 + 1\right)} + \frac{\left(q^{10} + q^6 - q^4 + q^2 - 1\right)\left(q^{12} - q^6 + 1\right)\left(b_{5,1,3} + b_{5,3,2} - b_{5,2,3} - b_{5,3,1}\right)}{q^{20}} + \frac{q^{10}\left(a_{4,3,4} - b_{4,3,4}\right)}{q^{12} - q^{10} - q^6 + q^4 + 1} - a_{1,3,4} + a_{2,3,4} - b_{4,3,2} + b_{4,3,1} = 0$$

4)
$$\frac{(q^{2}-1)(q^{4}+1)(q^{12}-q^{6}+1)(q^{18}-q^{16}+q^{10}-q^{8}-1)(b_{5,3,1}-b_{5,1,3})}{q^{20}} + \frac{(q^{2}-1)(q^{4}+1)(q^{6}-q^{2}-1)(b_{5,2,3}-b_{5,3,2})}{q^{4}} + b_{2,4,3} + \frac{(q^{4}+1)(q^{12}-q^{6}+1)(q^{12}-q^{10}+q^{8}-q^{2}+1)(a_{5,5,3}-b_{5,5,3})}{q^{6}} + + (q^{12}-q^{10}-q^{6}+q^{4}+1)q^{2}(b_{5,4,3}-b_{5,3,4}) - a_{3,4,2} = 0$$

5)
$$\frac{(q^{2}-1)(q^{4}+1)(q^{12}-q^{6}+1)(q^{18}-q^{16}+q^{10}-q^{8}-1)(a_{5,1,3}-a_{5,3,1})}{q^{20}} + \frac{(q^{4}+1)(q^{12}-q^{6}+1)(q^{12}-q^{10}+q^{8}-q^{2}+1)(a_{5,5,3}-b_{5,5,3})}{q^{6}} + \frac{(q^{2}-1)(q^{4}+1)(q^{6}-q^{2}-1)(a_{5,3,2}-a_{5,2,3})}{q^{4}} + (q^{12}-q^{10}-q^{6}+q^{4}+1)q^{2}(a_{5,3,4}-a_{5,4,3}) - a_{2,4,3}+b_{3,4,2} = 0$$

6)
$$\frac{\left(q^{12} - q^{10} - q^6 + q^4 + 1\right)\left(a_{1,3,1} - b_{1,3,1}\right)}{q^{10}} - a_{2,3,4} - a_{4,3,1} + b_{1,3,4} + b_{4,3,2} + \frac{\left(q^{10} + q^6 - q^4 + q^2 - 1\right)\left(q^{12} - q^6 + 1\right)\left(a_{5,3,1} - a_{5,1,3} + b_{5,2,3} - b_{5,3,2}\right)}{q^{20}} = 0$$

7)
$$\frac{\left(q^8+1\right)\left(q^8-q^4+1\right)\left(q^{16}+1\right)\left(q^4+1\right)^2\left(a_{5,5,3}-b_{5,5,3}\right)}{q^{20}}-a_{3,4,3}+b_{3,4,3}=0$$

8)
$$\frac{a_{5,3,4} - a_{5,4,3}}{q^6} + \frac{(q^2 - 1)(q^4 + 1)(q^6 + q^4 - 1)(a_{5,2,3} - a_{5,3,2})}{q^{20}} - a_{1,4,3} + \frac{(q^4 + 1)(q^{12} - q^6 + 1)(a_{5,5,3} - b_{5,5,3})}{q^{20}} + b_{3,4,1} + \frac{(q^2 - 1)(q^4 + 1)(q^6 - q^2 - 1)(a_{5,1,3} - a_{5,3,1})}{q^{16}} = 0$$

We find that the whole list of 19 5-CB relations, which can be found in the attached Mathematica file, is fulfilled for the Boltzmann weights of B_4 models. The Boltzmann weights are stated in [10]. We checked the relations numerically for a general value of the parameter q and substituting various heights.

5. 4–CB relations for D_6 .

We wish to check the 4–CB algebra for D_6 which is a four blocks model. The four blocks relations were given in [11]. The parameters for D_6 are, eq. (2.18).

$$\zeta_0 = 10, \qquad \zeta_1 = 6, \qquad \zeta_2 = 2,$$
 (5.1)

and q is given by eq. (2.7).

The skein relation is [11],

$$G_{i}^{2} = -iq^{-\frac{1}{2}\zeta_{0} - \zeta_{1} - \zeta_{2}} \left(1 - q^{2\zeta_{1}} + q^{2\zeta_{1} + 2\zeta_{2}}\right) G_{i} - iq^{-\frac{3}{2}\zeta_{0} + \zeta_{1} - \zeta_{2}} G_{i}^{-1} + \frac{q^{-2\zeta_{0} - 2\zeta_{1} - 2\zeta_{2}} \left(q^{2\zeta_{1}} - 1\right) \left(1 + q^{2\zeta_{0} + 2\zeta_{1} + 2\zeta_{2}}\right) \left(q^{2\zeta_{2}} - 1\right)}{\left(q^{2\zeta_{0} + 2\zeta_{2}} - 1\right)} E_{i}$$

$$(5.2)$$

$$-q^{-\zeta_0-2\zeta_2}\left(1-q^{2\zeta_2}+q^{2\zeta_1+2\zeta_2}\right).$$

The single additional relation is

$$g(i, i+1, i) = g(i+1, i, i+1), \tag{5.3}$$

where

$$g = a_{1,2,4} + a_{1,3,1} + a_{4,2,1} - iq^{-\zeta_0/2 + \zeta_1 - \zeta_2} (a_{1,3,4} + a_{4,2,4} + a_{4,3,1}) - (5.4)$$

$$iq^{\zeta_0/2 - \zeta_1 + \zeta_2} (a_{2,3,4} + a_{4,1,4} + a_{4,3,2}) - i\frac{q^{\zeta_1 + \zeta_2}}{(q^{2\zeta_1} - 1)(q^{2\zeta_2} - 1)} \left(q^{\zeta_0/2} a_{1,2,1} + q^{-\zeta_0/2} a_{2,1,2} \right) + z a_{4,3,4},$$

where

$$z = \frac{q^{-\zeta_0 - 2\zeta_1 - 2\zeta_2}(q^{2\zeta_1} - 1)(q^{2\zeta_2} - 1)}{q^{2\zeta_0 + 2\zeta_2} - 1} \times (5.5)$$
$$(2q^{2\zeta_0 + 2\zeta_2} + 2q^{2\zeta_0 + 2\zeta_1 + 2\zeta_2} + q^{4\zeta_0 + 2\zeta_1 + 4\zeta_2} + 1).$$

We denoted by $a_{i,j,k}(r, s, t)$ the element of the algebra $a_i[r]a_j[s]a_k[t]$ where $a_i[r]$ is G_r, G_r^{-1}, E_r or 1_r , if i = 1, 2, 3, 4, respectively.

Finally, we proceed to check these two relations, for the D_6 vertex model substituting the explicit Boltzmann weights, eqs. (2.20, 2.21). Indeed they hold for various values of the heights and for general value of q.

Acknowledgements: We thank Ida Deichaite for remarks on the manuscript.

REFERENCES

- 1. A.B. Zamolodchikov and A.B. Zamolodchikov, Ann. Phys. (NY) 120 253 (1979).
- 2. P. Francesco, P. Mathieu and D. Senechal, "Conformal field theory", Springer (1997).
- 3. R.J. Baxter, "Exactly solved models in statistical mechanics", Academic Press, London, England (1982).
- 4. M. Wadati, T. Deguchi and Y. Akutsu, Phys. Rep. 180 (4) (1989) 247.
- 5. B. Brubaker, arXiv: 1906.04140 (2019).
- 6. T.K. Kassenova, P. Yu and O.V. Razina, J. of Phys.: Conf. Series 1391 (2019).
- 7. A. Bossard and W. Galleas, J. of Math. Phys. 60, 103509 (2019).
- 8. J.H. Arbeitman, S. Mantilla and I. Sodeman, Phys. Rev. B 99 245108 (2019).
- 9. K.S. Nirov and A.V. Razumov, SIGMA 15 068 (2019).
- 10. D. Gepner, " B_k spin models and quantum algebras", arXiv: 2005.02708, Nucl. Phys. B (in press) (2020).
- 11. V. Belavin, D. Gepner, J.R. Li and R. Tessler, JHEP 11 (2019) 155.
- 12. V. Belavin and D. Gepner, arXiv: 2001.09280 (2020).
- 13. D. Gepner, "Foundations of rational quantum field theory I", arXiv: hep-th/9211100v2 (1992).
- 14. J.S. Birman and H. Wenzl, Trans. Am. Math. Soc. 313 (1) (1989) 313.
- 15. J. Murakami, Osaka J. Math. 24 (4) (1987) 745.
- 16. H. Wenzl, "Dualities for spin representation", arXiv: 2005.11299, (2020).
- 17. N. Temperley and E. Lieb, Proc. R. Soc. A 322 (1971) 251.