

On Stochastic Maximum Principle: A Backward Stochastic Partial Differential Equations Point of View

Ishak Alia

Mohamed Sofiane Alia

August 6, 2020

Abstract

In this paper, we consider a class of stochastic control problems for stochastic differential equations with random coefficients. The control domain need not to be convex but the control process is not allowed to enter in diffusion term. Moreover, the terminal cost involves a non linear term of the expected value of terminal state. Our purpose is to derive a new version of the Pontryagin's stochastic maximum principle by adopting an idea inspired from the work of Peng [S. Peng, *Maximum Principle for Stochastic Optimal Control with Nonconvex Control Domain*, Lecture Notes in Control & Information Sciences, 114, (1990), pp. 724-732]. More specifically, we show that if we combine the spike perturbation of the optimal control combined with the stochastic Feynman-Kac representation of linear backward stochastic partial differential equations (BSPDE, for short), a new version of the stochastic maximum principle can be derived. We also investigate sufficient conditions of optimality. In the last part of this paper, motivated by our version of SMP, an interesting class of forward backward stochastic partial differential equations is naturally introduced and the solvability of such kind of equations is briefly presented.

Keys words: Stochastic control, linear degenerate backward stochastic partial differential equation, optimal control, stochastic maximum principle.

MSC 2010 subject classifications, 91B51, 93E20, 60H30, 93E99.

1 Introduction

In this paper we consider a class of stochastic optimal control problems where the state of the system under consideration is governed by the controlled SDE,

$$\begin{cases} dX(t) = b(t, X(t), u(t)) dt + \sum_{j=1}^d \sigma^j(t, X(t)) dW_j(t), \\ X(0) = x_0, \end{cases} \quad (1.1)$$

and the objective of the controller is to minimize the following expected cost functional,

$$\mathbf{J}(u(\cdot)) = \mathbb{E} \left[\int_0^T f(t, X(t), u(t)) dt + h(X(T)) + G(\mathbb{E}[X(T)]) \right], \quad (1.2)$$

over the set of the admissible controls. Here $W(\cdot) = (W_1(\cdot), \dots, W_d(\cdot))^\top$ is a d-dimensional standard Brownian motion defined on some filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, T]}, \mathbb{P})$ satisfying usual conditions; the coefficients b, σ^j, f, h are sufficiently smooth $(\mathcal{F}_t)_{t \in [0, T]}$ -progressively measurable functions and G is a deterministic measurable function; $u(\cdot) : [0, T] \times \Omega \rightarrow U$ represent the control process, where (U, d) is a separable metric space; $X(\cdot)$ is the corresponding state process of $u(\cdot)$; and $x_0 \in \mathbb{R}^n$ is regarded as the initial state. A control process $\bar{u}(\cdot)$ that solves this problem is called optimal.

One of the principal methods in solving continuous-time stochastic optimal control problems is the Pontryagin's maximum principle approach, by which necessary conditions for optimality are derived by considering a *spike perturbation* of the optimal control (i.e. by perturbing an optimal control on a small time interval of length $\varepsilon > 0$). Combining a sort of Taylor expansion with respect to ε , with Euler's necessary condition of optimality one obtains a kind of variational inequality. By the duality, the stochastic maximum (or minimum) principle is obtained. It states that any optimal control along with the corresponding optimal state must solve the so called (extended) Hamiltonian system, which consists of a forward differential

equation and a linear backward differential equation called the adjoint equation, plus a maximum (or minimum) condition of a function called the Hamiltonian. The efficiency of the Pontryagin's maximum principle approach lies in the fact that maximizing (or minimizing) the Hamiltonian is much more manageable than the original control problem which is infinite-dimensional.

The Pontryagin's maximum principle approach was first performed for deterministic problems; see e.g. Pontryagin et al. [33]. Extension to stochastic diffusion control problems was first carried out by Kushner [21], followed by Haussmann ([13], [14]), Bismut [6], Elliott and Kohlmann [12] and Bensoussan [4]. However, at that time, the results were essentially obtained under the assumption that the diffusion term is independent of the control variable. Specially, in Bensoussan [4] the stochastic maximum principle (SMP, for short) is constructed for control problems, where the state is described by a stochastic differential equation (1.1) (with $n = d$) and the objective of the controller is to minimize the expected cost functional

$$\mathbf{J}_0(u(\cdot)) = \mathbb{E} \left[\int_0^T f(t, X(t), u(t)) dt + h(X(T)) \right]$$

over the set of admissible controls. The author constructs his SMP by considering a *spike perturbation* of the optimal control defined by $u^\varepsilon(t) = v\chi_{[\tau, \tau+\varepsilon)}(t) + \bar{u}(t)\chi_{[0, T] \setminus [\tau, \tau+\varepsilon)}(t)$, for any $\tau \in [0, T)$, $v \in U$ and for any $\varepsilon \in [0, T - \tau)$. Performing a Taylor expansion, the author obtained that

$$\begin{aligned} 0 &\leq \frac{d\mathbf{J}_0}{d\varepsilon}(u^\varepsilon(\cdot))|_{\varepsilon=0} \\ &= \mathbb{E} \left[h_x(\bar{X}(T))^\top z(T) + \zeta(T) \right], \end{aligned}$$

where $z(\cdot)$ is the unique solution of the so-called variational equation:

$$\begin{cases} dz(t) = b_x(t, \bar{X}(t), \bar{u}(t))^\top z(t) dt + \sum_{i=1}^d \sigma_x^i(t, \bar{X}(t))^\top z(t) dW_j(t) \\ z(\tau) = b(\tau, \bar{X}(\tau), v) - b(\tau, \bar{X}(\tau), \bar{u}(\tau)) \end{cases}$$

and $\zeta(\cdot)$ solves the following equation

$$\begin{cases} \frac{d\zeta}{dt}(t) = f_x(t, \bar{X}(t), \bar{u}(t))^\top z(t) \\ \zeta(\tau) = f(\tau, \bar{X}(\tau), v) - f(\tau, \bar{X}(\tau), \bar{u}(\tau)). \end{cases}$$

Then by introducing the first order adjoint processes $p(\cdot)$ and $q(\cdot) = (q_1(\cdot), \dots, q_d(\cdot))$ as the unique solution of the BSDE:

$$\begin{cases} dp(t) = - \left\{ \langle b_x(t, \bar{X}(t), \bar{u}(t)), p(t) \rangle + \sum_{j=1}^d \langle \sigma_x^j(t, \bar{X}(t)), q_j(t) \rangle \right. \\ \quad \left. + f_x(t, \bar{X}(t), \bar{u}(t)) \right\} dt + \sum_{j=1}^d q_j(t) dW_j(t), t \in [0, T], \\ p(T) = h_x(\bar{X}(T)), \end{cases} \quad (1.3)$$

and by applying Itô's formula to $t \rightarrow \langle p(t), z(t) \rangle$, the stochastic maximum principle can be obtained. It states that the optimal pair $(\bar{u}(\cdot), \bar{X}(\cdot))$ should satisfy the following variational inequality

$$\begin{aligned} 0 &\leq \langle b(\tau, \bar{X}(\tau), v) - b(\tau, \bar{X}(\tau), \bar{u}(\tau)), p(\tau) \rangle \\ &\quad + f(\tau, \bar{X}(\tau), v) - f(\tau, \bar{X}(\tau), \bar{u}(\tau)), \text{ a.s., } \forall v \in U, \text{ a.e. } \tau \in [0, T]. \end{aligned}$$

In the general case when the diffusion coefficient is allowed to depend on the control, the first version of the stochastic maximum principle was derived by Peng [30] in 1990. To overcome the difficulty arising from the perturbed stochastic integral term, the author considered the second-order term in the spike variation. In the form of his stochastic maximum principle the first order adjoint BSDE was accompanied by a second order BSDE involving the second order derivatives of the coefficients of the system, and the Hamiltonian was extended accordingly. In the work of Peng [31], the author introduced an alternative approach. He showed that the first and second order adjoint processes together with the variational inequality obtained in [30] can be derived in a natural way via random fields that solve a linear system of BSPDEs. With the joint effort of many researchers in the last 30 years, there is a very extensive literature on different versions of Peng's type SMP. Let us just mention a few: see Tang and Li [34] for systems with jumps, see Bahlali and

Mezerdi [3] for stochastic singular control, see Buckdan et al. [7] for stochastic systems of mean-field type, see Yong [36] and Hu [16] for general coupled forward-backward stochastic differential equations (FBSDEs). More recently, Agram and Øksendal [1] presented a new approach based on Hida-Malliavin calculus and white noise theory, which enabled them to derive the SMP without involving the second adjoint BSDE.

The purpose of this paper is to suggest a new version of Pontryagin's SMP to the class of control problems (1.1)-(1.2). More specifically, by combining the spike perturbation of the optimal control with stochastic Feynman-Kac representations of linear degenerate BSPDEs (see e.g. [Ma and Yong [22], Section 6]), we show that the optimal pair $(\bar{u}(\cdot), \bar{X}(\cdot))$ should satisfy the following variational inequality

$$\begin{aligned} 0 \leq & \langle b(\tau, \bar{X}(\tau), u) - b(\tau, \bar{X}(\tau), \bar{u}(\tau)), \bar{\theta}_x(\tau, \bar{X}(\tau)) \rangle \\ & + \sum_{i=1}^n \langle b(\tau, \bar{X}(\tau), u) - b(\tau, \bar{X}(\tau), \bar{u}(\tau)), G_{\bar{x}_i}(\mathbb{E}[\bar{X}(T)]) \bar{g}_x^i(\tau, \bar{X}(\tau)) \rangle \\ & + f(\tau, \bar{X}(\tau), u) - f(\tau, \bar{X}(\tau), \bar{u}(\tau)), \text{ for all } u \in U, \text{ a.s., a.e. } \tau \in [0, T], \end{aligned}$$

and the objective value of $\bar{u}(\cdot)$ is given by

$$\mathbf{J}(\bar{u}(\cdot)) = \bar{\theta}(0, x_0) + G(\bar{g}(0, x_0)),$$

where the pair of random fields $(\bar{\theta}(\cdot, \cdot), \bar{\psi}(\cdot, \cdot))$ is the classical solution of the following linear system of BSPDE:

$$\begin{cases} d\bar{\theta}(t, x) = - \left\{ \langle \bar{\theta}_x(t, x), b(t, x, \bar{u}(t)) \rangle + \frac{1}{2} \mathbf{tr} [\sigma \sigma^\top(t, x) \bar{\theta}_{xx}(t, x)] \right. \\ \quad \left. + \mathbf{tr} [\bar{\psi}_x(t, x) \sigma(t, x)] + f(t, x, \bar{u}(t)) \right\} dt \\ \quad + \bar{\psi}(t, x)^\top dW(t), \quad (t, x) \in [0, T] \times \mathbb{R}^n, \\ \bar{\theta}(T, x) = F(x), \text{ for } x \in \mathbb{R}^n \end{cases} \quad (1.4)$$

and

$$\bar{g}(t, x) \equiv (\bar{g}^1(t, x), \dots, \bar{g}^n(t, x))^\top$$

such that for each for $1 \leq i \leq n$, $(\bar{g}^i(\cdot, \cdot), \bar{\eta}^i(\cdot, \cdot))$ is the classical solution of the following linear BSPDE:

$$\begin{cases} d\bar{g}^i(t, x) = - \left\{ \langle \bar{g}_x^i(t, x), b(t, x, \bar{u}(t)) \rangle + \frac{1}{2} \mathbf{tr} [\sigma \sigma^\top(t, x) \bar{g}_{xx}^i(t, x)] \right. \\ \quad \left. + \mathbf{tr} [\bar{\eta}_x^i(t, x) \sigma(t, x)] \right\} dt + \bar{\eta}^i(t, x)^\top dW(t), \\ \quad \text{for } (t, x) \in [0, T] \times \mathbb{R}^n, \\ \bar{g}^i(T, x) = \bar{x}_i, \text{ for } x \in \mathbb{R}^n, \end{cases} \quad (1.5)$$

with x_i denotes the i -th coordinate of $x \in \mathbb{R}^n$.

There exists many other situations where BSPDEs are involved to construct an optimal solution of a stochastic optimal control problem. For instance, BSPDEs appear as adjoint equations in the study of stochastic maximum principle for stochastic parabolic PDEs (see e.g. [40]) and as adjoint equations of Duncan-Mortensen-Zakai filtering equations to formulate the stochastic maximum principle for stochastic control problems with partial information (see e.g. [5], [27], [39]). A class of fully nonlinear BSPDEs, the so-called backward stochastic Hamilton-Jacobi-Bellman equations, appear naturally in the dynamic programming theory of controlled non-Markovian processes [32]. More recently, Alia [2] investigated equilibrium solutions for a general class of time-inconsistent control problems by using BSPDEs. However, to our best knowledge, the above-described version of SMP for optimal controls seems to be new. Moreover, it permits us to derive the optimal solution $\bar{u}(\cdot)$ as well as its objective value $\mathbf{J}(\bar{u}(\cdot))$ in terms of the solutions of the BSPDEs (1.4)-(1.5); this is different from the traditional SMP approach which does not provide the objective value $\mathbf{J}(\bar{u}(\cdot))$ in a direct way.

In the latter part of the paper, we investigate a class of FBSPDEs which naturally arises from the above-described version of SMP. These systems differ from the classical FBSDEs (see e.g. [26]), since they are consisting of a forward SDE and a nonlinear BSPDE. Inspired by the idea of the classical four-step scheme introduced by Ma et al. [25], we show that a solution of the FBSPDEs can be constructed by solving a parabolic PDE. Under proper conditions on the involved coefficients, a well-posedness result of the PDE is established.

The plan of the paper is as follows, in the second section, we give necessary notations and some preliminaries on linear BSPDEs. In Section 3, we formulate our stochastic optimal control problem. Section 4 is devoted to the new version of the SMP. Section 5 is devoted to the proof of the first main result in this paper. In Section 6, we investigate sufficient conditions of optimality and we give a simple example to illustrate our result. Finally, in Section 7, a class of FBSPDEs is briefly presented.

2 Preliminaries

2.1 Notations

Throughout this paper $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, T]}, \mathbb{P})$ is a filtered probability space such that \mathcal{F}_0 contains all \mathbb{P} -null sets, $\mathcal{F}_T = \mathcal{F}$ for an arbitrarily fixed finite time horizon $T > 0$, and $(\mathcal{F}_t)_{t \in [0, T]}$ satisfies the usual conditions. \mathcal{F}_t stands for the information available up to time t and any decision made at time t is based on this information. We also assume that all processes and random variables are well defined and adapted in this filtered probability space. Let $W(\cdot) = (W_1(\cdot), \dots, W_d(\cdot))$ be a d -dimensional standard Brownian motion defined on $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, T]}, \mathbb{P})$. For simplicity, it is assumed that the filtration $(\mathcal{F}_t)_{t \in [0, T]}$, coincides with the one generated by the Brownian motion; that is $\mathcal{F}_t = \sigma(W(r); 0 \leq r \leq t)$.

We use C^\top to denote the transpose of any vector or matrix C , $K > 0$ is a generic constant which can be different from line to line and for a function f , we denote by f_x (resp. f_{xx}) the gradient or Jacobian (resp. the Hessian) of f with respect to the variable x . We denote by χ_A the indicator function of the set A . In addition, we use the following notations for several sets and spaces of processes on the filtered probability space, which will be used later:

- $\bar{B}_R = \{x \in \mathbb{R}^n, \text{ such that } |x| \leq R\}$, for any $R > 0$.
- (U, d) : a separable metric space.
- $\mathcal{U}[0, T]$: the set of U -valued $(\mathcal{F}_t)_{t \in [0, T]}$ -progressively measurable processes $u(\cdot)$.

For any $m \geq 1$ and $p \geq 2$ we denote by

- $\mathbb{L}^p(\Omega, \mathcal{F}_t, \mathbb{P}; \mathbb{R}^l)$: the set of \mathbb{R}^l -valued, \mathcal{F}_t -measurable random variables ζ , with

$$\|\zeta\|_{\mathbb{L}^p(\Omega, \mathcal{F}_t, \mathbb{P}; \mathbb{R}^n)} = (\mathbb{E}[|\zeta|^p])^{\frac{1}{p}} < \infty.$$

- $\mathcal{C}_{\mathcal{F}}^p(0, T; \mathbb{R}^l)$: the space of \mathbb{R}^l -valued, $(\mathcal{F}_t)_{t \in [0, T]}$ -adapted continuous processes $X(\cdot)$, with

$$\|X(\cdot)\|_{\mathcal{C}_{\mathcal{F}}^p(0, T; \mathbb{R}^l)} := \left(\mathbb{E} \left[\sup_{t \in [0, T]} |X(t)|^p \right] \right)^{\frac{1}{p}} < \infty.$$

- $\mathcal{L}_{\mathcal{F}}^p(0, T; \mathbb{R}^l)$: the space of \mathbb{R}^l -valued, $(\mathcal{F}_t)_{t \in [0, T]}$ -adapted processes $Z(\cdot)$, with

$$\|Z(\cdot)\|_{\mathcal{L}_{\mathcal{F}}^p(0, T; \mathbb{R}^l)} := \left(\mathbb{E} \left[\int_0^T |Z(t)|^p dt \right] \right)^{\frac{1}{p}} < \infty.$$

- $C^m(\mathbb{R}^n; \mathbb{R}^l)$: the set of functions from \mathbb{R}^n to \mathbb{R}^l that are continuously differentiable up to order $m \geq 1$.
- $C_b^m(\mathbb{R}^n; \mathbb{R}^l)$: the set of those functions in $C^m(\mathbb{R}^n; \mathbb{R}^l)$ whose partial derivatives up to order m are uniformly bounded.

Finally, for any infinite-dimensional Banach space E with a norm $\|\cdot\|_E$, we denote.

- $\mathcal{C}_{\mathcal{F}}^p(0, T; E)$: the set of all $(\mathcal{F}_t)_{t \in [0, T]}$ -adapted E -valued continuous processes $\theta(\cdot)$ such that:

$$\|\theta(\cdot)\|_{\mathcal{C}_{\mathcal{F}}^p(0, T; E)} := \sup_{t \in [0, T]} (\mathbb{E}[\|\theta(t)\|_E^p])^{\frac{1}{p}} < \infty.$$

- $\mathcal{L}_{\mathcal{F}}^p(0, T; E)$: the set of all $(\mathcal{F}_t)_{t \in [0, T]}$ -adapted E -valued processes $\psi(\cdot)$ such that:

$$\|\psi(\cdot)\|_{\mathcal{L}_{\mathcal{F}}^p(0, T; E)} := \left(\mathbb{E} \left[\int_0^T \|\psi(t)\|_E^p dt \right] \right)^{\frac{1}{p}} < \infty.$$

2.2 Linear Backward Stochastic Partial Differential Equations

In this paragraph, we review some well-known results on linear BSPDEs. Let $b : [0, T] \times \mathbb{R}^n \times \Omega \rightarrow \mathbb{R}^n$, $\sigma : [0, T] \times \mathbb{R}^n \times \Omega \rightarrow \mathbb{R}^{n \times d}$, $l : [0, T] \times \mathbb{R}^n \times \Omega \rightarrow \mathbb{R}$ be three $(\mathcal{F}_t)_{t \in [0, T]}$ -progressively measurable functions and $F : \mathbb{R}^n \times \Omega \rightarrow \mathbb{R}$ be an \mathcal{F}_T -measurable function. To simplify our notation, for $\varphi = b, \sigma, l$ we write $\varphi(t, x)$ for $\varphi(t, x, \omega)$ and $h(x)$ for $h(x, \omega)$.

Consider the following linear backward stochastic partial differential equation in the unknown random fields $p(t, x) \in \mathbb{R}$, $q(t, x) = (q^1(t, x), \dots, q^d(t, x))^\top \in \mathbb{R}^d$:

$$\begin{cases} dp(t, x) = - \left\{ \langle p_x(t, x), b(t, x) \rangle + \frac{1}{2} \text{tr} \left[\sigma(t, x) \sigma(t, x)^\top p_{xx}(t, x) \right] \right. \\ \quad \left. + \text{tr} [\sigma(t, x) q_x(t, x)] + l(t, x) \right\} dt \\ \quad + \langle q(t, x), dW(t) \rangle, \quad (t, x) \in [0, T] \times \mathbb{R}^n, \\ p(T, x) = h(x), \quad \text{for } x \in \mathbb{R}^n. \end{cases} \quad (2.1)$$

A pair of random fields $(p(\cdot, \cdot), q(\cdot, \cdot))$ is called a *classical solution* of (2.1) if (see e.g. [23])

$$\begin{cases} p(\cdot, \cdot) \in \mathcal{C}_{\mathcal{F}}^2(0, T; C^2(\bar{B}_R; \mathbb{R})), \\ q(\cdot, \cdot) \in \mathcal{L}_{\mathcal{F}}^2(0, T; C^1(\bar{B}_R; \mathbb{R}^d)), \end{cases} \quad \forall R > 0,$$

such that the following holds for all $(t, x) \in [0, T] \times \mathbb{R}^n$, almost surely:

$$\begin{aligned} p(t, x) &= h(x) + \int_t^T \left\{ \langle p_x(\tau, x), b(\tau, x) \rangle \right. \\ &\quad \left. + \frac{1}{2} \text{tr} \left[\sigma(\tau, x) \sigma(\tau, x)^\top p_{xx}(\tau, x) \right] + \text{tr} [\sigma(\tau, x) q_x(\tau, x)] \right. \\ &\quad \left. + l(\tau, x) \right\} d\tau - \int_t^T \langle q(\tau, x), dW(\tau) \rangle. \end{aligned}$$

The existence and uniqueness of a solution to linear or semi-linear degenerate BSPDEs driven by a Brownian motion was first proved by Ma and Yong ([22], [23]), followed by Hu et al. [15], Tang [35], Du et al. [10], Du and Zhang [11] and Ma et al. [24] in different frameworks. More recently, Chen and Tang [8] dealt with semi-linear backward stochastic integral partial differential equations with jumps. In this work, we mostly focus on the work of Tang [35], in which the author developed a probabilistic approach to construct the adapted solution of semi-linear BSPDEs in terms of those of SDEs and BSDEs.

Given an integer $m \geq 1$, we consider the following assumption.

(\mathbf{A}_m) The random fields b, σ, f and h satisfy the following conditions:

$$\begin{cases} b \in \mathcal{L}_{\mathcal{F}}^\infty(0, T; C_b^m(\mathbb{R}^n; \mathbb{R}^n)), \\ \sigma \in \mathcal{L}_{\mathcal{F}}^\infty(0, T; C_b^m(\mathbb{R}^n; \mathbb{R}^{n \times d})), \\ f \in \mathcal{L}_{\mathcal{F}}^\infty(0, T; C_b^m(\mathbb{R}^n; \mathbb{R})), \\ h \in \mathcal{L}_{\mathcal{F}_T}^\infty(\Omega; C_b^m(\mathbb{R}^n; \mathbb{R})). \end{cases}$$

The following theorem concerns the existence and uniqueness of a classical solution to the BSPDEs (2.1).

Theorem 2.1 ([35]). *Let Assumption (\mathbf{A}_m) be satisfied with $m > 2 + \frac{n}{2}$. Then the BSPDE (2.1) admits a unique adapted classical solution $(p(\cdot, \cdot), q(\cdot, \cdot))$. Moreover, the partial derivatives in x of $p(\cdot, \cdot)$ up to order m are uniformly bounded.*

The following theorem provides the stochastic Feynman–Kac representation of the linear degenerate BSPDE (2.1).

Theorem 2.2 (Feynman–Kac representation). *Suppose that Assumption (\mathbf{A}_m) is satisfied with $m > 2 + \frac{n}{2}$. Let $(p(\cdot, \cdot), q(\cdot, \cdot))$ be the classical solution of the BSPDEs (2.1), then we have the following equality: For each $(t, \zeta) \in [0, T] \times \mathbb{L}^2(\Omega, \mathcal{F}_t, \mathbb{P}; \mathbb{R}^n)$,*

$$p(t, \zeta) = \mathbb{E}_t \left[\int_t^T l(\tau, X^{t, \zeta}(\tau)) d\tau + h(X^{t, \zeta}(T)) \right], \quad (2.2)$$

where $\mathbb{E}_t[\cdot] = \mathbb{E}_t[\cdot | \mathcal{F}_t]$ and $X^{t, \zeta}(\cdot) \in \mathcal{C}_{\mathcal{F}}^2(t, T; \mathbb{R}^n)$ is the unique strong solution of the following SDE,

$$\begin{cases} dX^{t, \zeta}(\tau) = b(\tau, X^{t, \zeta}(\tau)) d\tau + \sigma(\tau, X^{t, \zeta}(\tau)) dW(\tau), \quad \tau \in [t, T], \\ X^{t, \zeta}(t) = \zeta. \end{cases} \quad (2.3)$$

Before presenting a proof of the above theorem we first recall a version of an Itô-Wentzell formula to the composition of random fields and continuous semimartingale processes; see e.g. Kunita ([18], [19], [20]), Krylov [17] and Chen and Tang [8].

Lemma 2.3 (Itô-Wentzell Formula). *Let $X(\cdot) \in \mathcal{C}_{\mathcal{F}}^2(0, T; \mathbb{R}^n)$ be a process of the form*

$$dX(t) = b(t) dt + \sigma(t) dW(t),$$

where $b(\cdot) \in \mathcal{L}_{\mathcal{F}}^{\infty}(0, T; \mathbb{R}^n)$ and $\sigma(\cdot) \in \mathcal{L}_{\mathcal{F}}^{\infty}(0, T; \mathbb{R}^{n \times d})$. Suppose that $V(\cdot, \cdot) \in \mathcal{C}_{\mathcal{F}}^2(0, T; C^2(\bar{B}_R; \mathbb{R}))$, $\forall R > 0$, is a semimartingale with spatial parameter $x \in \mathbb{R}$:

$$dV(t, x) = \Gamma(t, x) dt + \zeta(t, x)^{\top} dW(t), \quad (t, x) \in [0, T] \times \mathbb{R},$$

with $\Gamma(\cdot, \cdot) \in \mathcal{L}_{\mathcal{F}}^1(0, T; C(\bar{B}_R; \mathbb{R}))$ and $\zeta(\cdot, \cdot) \in \mathcal{L}_{\mathcal{F}}^2(0, T; C^1(\bar{B}_R; \mathbb{R}^d))$, $\forall R > 0$. Then the following holds for all $0 \leq t \leq s \leq T$, almost surely,

$$\begin{aligned} & V(s, X(s)) - V(t, X(t)) \\ &= \int_t^s \left\{ \Gamma(\tau, X(\tau)) + \langle V_x(\tau, X(\tau)), b(\tau) \rangle \right. \\ & \quad \left. + \text{tr}[\zeta_x(\tau, X(\tau)) \sigma(\tau)] + \frac{1}{2} \text{tr} \left[\sigma(\tau) \sigma(\tau)^{\top} V_{xx}(\tau, X(\tau)) \right] \right\} d\tau \\ & \quad + \int_t^s \left\{ \zeta(\tau, X(\tau))^{\top} + V_x(\tau, X(\tau))^{\top} \sigma(\tau) \right\} dW(\tau). \end{aligned}$$

Proof. A proof of this lemma can be easily obtained by adapting the proof of Lemma 3.1 in [8]. We omit it. \blacksquare

Proof of Theorem 2.2. Let $(p(\cdot, \cdot), q(\cdot, \cdot))$ be the classical solution of the BSPDEs (2.1) and $X^{t,x}(\cdot)$ is the unique strong solution of the SDE (2.3). Define

$$\mathbf{J} := \mathbb{E}_t \left[\int_t^T l(\tau, X^{t,\zeta}(\tau)) d\tau + h(X^{t,\zeta}(T)) \right].$$

By the terminal condition in (2.1) we have

$$\begin{aligned} \mathbf{J} &= \mathbb{E}_t \left[\int_t^T l(\tau, X^{t,\zeta}(\tau)) d\tau + p(T, X^{t,\zeta}(T)) \right] \\ &= \mathbb{E}_t \left[\int_t^T l(\tau, X^{t,\zeta}(\tau)) d\tau + p(t, \zeta) \right] \\ & \quad + \mathbb{E}_t [p(T, X^{t,\zeta}(T)) - p(t, \zeta)]. \end{aligned} \tag{2.4}$$

Moreover, by applying Itô-Wentzell formula to $p(\tau, X^{t,\zeta}(\tau))$ on time interval $[t, T]$, we obtain that

$$\mathbb{E}_t [p(T, X^{t,\zeta}(T)) - p(t, \zeta)] = -\mathbb{E}_t \left[\int_t^T l(\tau, X^{t,\zeta}(\tau)) d\tau \right].$$

where $\mathbb{E}_t[\cdot] = \mathbb{E}[\cdot | \mathcal{F}_t]$ is the conditional expectation with respect to \mathcal{F}_t . Invoking this into (2.4), we obtain (2.2). \blacksquare

3 Formulation of the problem

We consider a continuous-time, n-dimensional, controlled system

$$\begin{cases} dX(t) = b(t, X(t), u(t)) dt + \sigma(t, X(t)) dW(t), & t \in [0, T], \\ X(0) = x_0. \end{cases} \tag{3.1}$$

Here $b : [0, T] \times \mathbb{R}^n \times U \times \Omega \rightarrow \mathbb{R}^n$ and $\sigma : [0, T] \times \mathbb{R}^n \times \Omega \rightarrow \mathbb{R}^{n \times d}$ are $(\mathcal{F}_t)_{t \in [0, T]}$ -progressively measurable functions; $u(\cdot) \in \mathcal{U}[0, T]$ represents an admissible control process; $X(\cdot) = X^{x_0, u(\cdot)}(\cdot)$ is the controlled state process; and $x_0 \in \mathbb{R}$ is regarded as the initial state.

In order to evaluate the cost-performance of a control process $u(\cdot)$, we introduce the functional

$$\mathbf{J}(u(\cdot)) := \mathbb{E} \left[\int_0^T f(t, X(t), u(t)) dt + h(X(T)) + G(\mathbb{E}[X(T)]) \right] \quad (3.2)$$

where $f : [0, T] \times \mathbb{R}^n \times U \times \Omega \rightarrow \mathbb{R}$ is an $(\mathcal{F}_t)_{t \in [0, T]}$ -progressively measurable function, $h : \mathbb{R}^n \times \Omega \rightarrow \mathbb{R}$, is an \mathcal{F}_T -measurable function and $G : \mathbb{R}^n \rightarrow \mathbb{R}$ is a deterministic function.

Before going further, we introduce some notations. For any fixed control $u(\cdot) \in \mathcal{U}[0, T]$, we define the random fields $b^{u(\cdot)} : [0, T] \times \mathbb{R}^n \times \Omega \rightarrow \mathbb{R}^n$ and $f^{u(\cdot)} : [0, T] \times \mathbb{R}^n \times \Omega \rightarrow \mathbb{R}$ by

$$\begin{aligned} b^{u(\cdot)}(t, x) &:= b(t, x, u(t)), \\ f^{u(\cdot)}(t, x) &:= f(t, x, u(t)). \end{aligned}$$

Given an integer $m \geq 1$, we shall use the following assumption.

(H_m) For each fixed $u(\cdot) \in \mathcal{U}[0, T]$, the random fields $b^{u(\cdot)}$, σ , $f^{u(\cdot)}$, h and G satisfy the following conditions:

$$\begin{cases} b^{u(\cdot)} \in \mathcal{L}_{\mathcal{F}}^{\infty}(0, T; C_b^m(\mathbb{R}^n; \mathbb{R}^n)), \\ \sigma \in \mathcal{L}_{\mathcal{F}}^{\infty}(0, T; C_b^m(\mathbb{R}^n; \mathbb{R}^{n \times d})), \\ f^{u(\cdot)} \in \mathcal{L}_{\mathcal{F}}^{\infty}(0, T; C_b^m(\mathbb{R}^n; \mathbb{R})), \\ F \in \mathcal{L}_{\mathcal{F}_T}^{\infty}(\Omega; C_b^m(\mathbb{R}^n; \mathbb{R})), \\ G \in \mathcal{L}^{\infty}(\Omega; C_b^2(\mathbb{R}^n; \mathbb{R})) \end{cases}$$

Note that **(H_m)** implies that the derivatives in x of $b^{u(\cdot)}$ and σ up to order $m \geq 1$ are uniformly bounded by some positive constant. Under **(H_m)**, the state equation (3.1) admits a unique strong solution $X(\cdot) \in \mathcal{C}_{\mathcal{F}}^2(0, T; \mathbb{R}^n)$ (see e.g. [38]). Moreover, there exists a constant $K > 0$ such that

$$\mathbb{E} \left[\sup_{0 \leq t \leq T} |X(t)|^2 \right] \leq K \left(1 + |x_0|^2 \right).$$

Our stochastic optimal control problem can be stated as follows.

Problem (S). *Minimize (3.2) over $\mathcal{U}[0, T]$.*

Any $\bar{u}(\cdot)$ satisfying

$$\mathbf{J}(\bar{u}(\cdot)) = \inf_{u(\cdot) \in \mathcal{U}[0, T]} \mathbf{J}(u(\cdot))$$

is called an optimal control. The corresponding $\bar{X}(\cdot) = X^{x_0, \bar{u}(\cdot)}(\cdot)$ and $(\bar{u}(\cdot), \bar{X}(\cdot))$ are called an optimal state process and optimal pair, respectively.

4 Necessary Condition For Optimality

As well mentioned in Introduction Section, the first order Taylor expansion in the spike variation together with the pair of adjoint processes $(p(\cdot), q(\cdot))$ play a central role in the traditional stochastic maximum principle approach [4]. Moreover, the adjoint equation that $(p(\cdot), q(\cdot))$ satisfies is a standard BSDE having finite-dimensional state process.

In this paper, inspired by Peng [31], we follow an alternative approach which permits us to derive a stochastic maximum principle of Problem (S) without performing a Taylor expansion and without involving the adjoint processes $(p(\cdot), q(\cdot))$. More specifically, we show that if we use the spike perturbation of the optimal control combined with stochastic Feynman-Kac representations of linear degenerate BSPDEs, we can obtain a new version of the stochastic maximum principle.

In the next, we introduce the backward stochastic partial differential equations involved in the new version of the SMP.

For a given optimal control $\bar{u}(\cdot) \in \mathcal{U}[0, T]$, we define in $(t, x) \in [0, T] \times \mathbb{R}^n$, the following *linear degenerate backward stochastic partial differential equation* in the unknown random fields $\bar{\theta}(t, x) \in \mathbb{R}$, $\bar{\psi}(t, x) \in \mathbb{R}^d$:

$$\begin{cases} d\bar{\theta}(t, x) = - \left\{ \langle \bar{\theta}_x(t, x), b^{\bar{u}(\cdot)}(t, x) \rangle + \frac{1}{2} \mathbf{tr} \left[(\sigma \sigma^\top)^{u(\cdot)}(t, x) \bar{\theta}_{xx}(t, x) \right] \right. \\ \quad \left. + \mathbf{tr} [\bar{\psi}_x(t, x) \sigma(t, x)] + f^{\bar{u}(\cdot)}(t, x) \right\} dt \\ \quad + \bar{\psi}(t, x)^\top dW(t), \quad (t, x) \in [0, T] \times \mathbb{R}^n, \\ \bar{\theta}(T, x) = h(x), \quad \text{for } x \in \mathbb{R}^n. \end{cases} \quad (4.1)$$

Similarly, we define in $(t, x) \in [0, T] \times \mathbb{R}^n$ the following system of linear degenerate BSPDEs in the unknown random fields $\bar{g}(s, x) = (\bar{g}^1(s, x), \dots, \bar{g}^n(s, x))^\top \in \mathbb{R}^n$, $\bar{\eta}(s, x) = (\bar{\eta}^1(s, x), \dots, \bar{\eta}^n(s, x)) \in \mathbb{R}^{d \times n}$: For $1 \leq i \leq n$,

$$\begin{cases} d\bar{g}^i(t, x) = - \left\{ \langle \bar{g}_x^i(t, x), b^{\bar{u}(\cdot)}(t, x) \rangle + \frac{1}{2} \mathbf{tr} \left[(\sigma \sigma^\top)(t, x) \bar{g}_{xx}^i(t, x) \right] \right. \\ \quad \left. + \mathbf{tr} [\sigma(t, x) \bar{\eta}_x^i(t, x)] \right\} dt + \bar{\eta}^i(t, x)^\top dW(t), \\ \quad \text{for } (t, x) \in [0, T] \times \mathbb{R}^n, \\ \bar{g}^i(T, x) = x_i, \quad \text{for } x \in \mathbb{R}^n. \end{cases} \quad (4.2)$$

where $\bar{\eta}^i(s, x)$ is the i -th column of $\bar{\eta}(s, x)$ and x_i denotes the i -th coordinate of $x \in \mathbb{R}^n$.

According to Theorem 2.1, under Assumption (\mathbf{H}_m) , with $m > 2 + \frac{n}{2}$, the BSPDE (4.1) admits a unique adapted classical solution $(\bar{\theta}(\cdot, \cdot), \bar{\psi}(\cdot, \cdot))$. Moreover, all the partial derivatives in x of $\bar{\theta}(\cdot, \cdot)$ up to order m are uniformly bounded.

Since the terminal condition of (4.2) is unbounded, we can not directly apply Theorem 2.1 to obtain the well-posedness of (4.2). Nonetheless, we state the following result which by means of easy manipulations shows that the well-posedness of (4.2) holds indeed.

Theorem 4.1. *Let Assumption (\mathbf{H}_m) be satisfied with $m > 2 + \frac{n}{2}$. Then for each $1 \leq i \leq n$, the BSPDE (4.2) admits a unique adapted classical solution $(\bar{g}^i(\cdot, \cdot), \bar{\eta}^i(\cdot, \cdot))$. Moreover, all the partial derivatives in x of $\bar{g}^i(\cdot, \cdot)$ up to order m are uniformly bounded.*

Proof. First define for each $1 \leq i \leq n$,

$$\begin{aligned} \tilde{g}^i(t, x) &:= \bar{g}^i(t, x) - x_i, \\ \tilde{\eta}^i(t, x) &:= \bar{\eta}^i(t, x). \end{aligned}$$

Accordingly one has, for each $1 \leq i \leq n$,

$$\begin{aligned} \tilde{g}_x^i(t, x) &\equiv \bar{g}_x^i(t, x) - \mathbf{e}_i, \\ \tilde{g}_{xx}^i(t, x) &\equiv \bar{g}_{xx}^i(t, x), \\ \tilde{\eta}_x^i(t, x) &\equiv \bar{\eta}_x^i(t, x), \end{aligned}$$

where $\mathbf{e}_i = (\delta_{i,1}, \delta_{i,2}, \dots, \delta_{i,n})^\top$, with $\delta_{i,j} = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{if } i \neq j. \end{cases}$

Then it is not difficult to see that $(\bar{g}^i(\cdot, \cdot), \bar{\eta}^i(\cdot, \cdot))$ is a classical solution of (4.2), if and only if, $(\tilde{g}^i(\cdot, \cdot), \tilde{\eta}^i(\cdot, \cdot))$ is a classical solution of the following BSPDE

$$\begin{cases} d\tilde{g}^i(t, x) = - \left\{ \langle \tilde{g}_x^i(t, x), b^{\bar{u}(\cdot)}(t, x) \rangle + \frac{1}{2} \mathbf{tr} \left[(\sigma \sigma^\top)^{u(\cdot)}(t, x) \tilde{g}_{xx}^i(t, x) \right] \right. \\ \quad \left. + \mathbf{tr} [\tilde{\eta}_x^i(t, x) \sigma(t, x)] + (b^i)^{\bar{u}(\cdot)}(t, x) \right\} dt \\ \quad + \tilde{\eta}^i(t, x)^\top dW(t), \quad \text{for } (t, x) \in [0, T] \times \mathbb{R}^n, \\ \tilde{g}^i(T, x) = 0, \quad \text{for } x \in \mathbb{R}^n, \end{cases} \quad (4.3)$$

where $(b^i)^{\bar{u}(\cdot)}(t, x) = \langle \mathbf{e}_i, b^{\bar{u}(\cdot)}(t, x) \rangle$ is the i -th coordinate of $b^{\bar{u}(\cdot)}(t, x)$.

According to Theorem 2.1, under Assumption (\mathbf{H}_m) with $m > 2 + \frac{n}{2}$, the BSPDE (4.3) admits a unique adapted classical solution $(\tilde{g}^i(\cdot, \cdot), \tilde{\eta}^i(\cdot, \cdot))$ and all the partial derivatives in x of $\tilde{g}^i(\cdot, \cdot)$ up to order m are uniformly bounded. Therefor, the BSPDE (4.2) admits a unique adapted classical solution $(\bar{g}^i(\cdot, \cdot), \bar{\eta}^i(\cdot, \cdot))$. ■

In the next, we define for any $\bar{x} \in \mathbb{R}^n$,

$$G_{\bar{x}_i}(\bar{x}) := \frac{\partial}{\partial \bar{x}_i} G(\bar{x}_1, \dots, \bar{x}_n).$$

The following theorem constitutes the first main contribution of the paper.

Theorem 4.2 (Stochastic Maximum Principle). *Let Assumption (\mathbf{H}_m) be satisfied with $m > 2 + \frac{n}{2}$. If $(\bar{u}(\cdot), \bar{X}(\cdot))$ is an optimal pair of Problem (S), then the BSPDEs (4.1) and (4.2), for $1 \leq i \leq n$, admit the classical solutions $(\bar{\theta}(\cdot, \cdot), \bar{\psi}(\cdot, \cdot))$ and $(\bar{g}^i(\cdot, \cdot), \bar{\eta}^i(\cdot, \cdot))$, respectively, such that the following holds*

$$\begin{aligned} 0 \leq & \langle b(\tau, \bar{X}(\tau), u) - b(\tau, \bar{X}(\tau), \bar{u}(\tau)), \bar{\theta}_x(\tau, \bar{X}(\tau)) \rangle \\ & + \sum_{i=1}^n G_{\bar{x}_i}(\mathbb{E}[\bar{X}(T)]) \langle b(\tau, \bar{X}(\tau), u) - b(\tau, \bar{X}(\tau), \bar{u}(\tau)), \bar{g}_x^i(\tau, \bar{X}(\tau)) \rangle \\ & + f(\tau, \bar{X}(\tau), u) - f(\tau, \bar{X}(\tau), \bar{u}(\tau)), \text{ for all } u \in U, \text{ a.s., a.e. } \tau \in [0, T], \end{aligned} \quad (4.4)$$

or, equivalently,

$$\begin{aligned} & \langle b(\tau, \bar{X}(\tau), \bar{u}(\tau)), \bar{\theta}_x(\tau, \bar{X}(\tau)) + \sum_{i=1}^n G_{\bar{x}_i}(\mathbb{E}[\bar{X}(T)]) \bar{g}_x^i(\tau, \bar{X}(\tau)) \rangle \\ & + f(\tau, \bar{X}(\tau), \bar{u}(\tau)) \\ & = \min_{u \in U} \{ \langle b(\tau, \bar{X}(\tau), u), \bar{\theta}_x(\tau, \bar{X}(\tau)) + \sum_{i=1}^n G_{\bar{x}_i}(\mathbb{E}[\bar{X}(T)]) \bar{g}_x^i(\tau, \bar{X}(\tau)) \rangle \\ & + f(\tau, \bar{X}(\tau), u) \}, \text{ a.e. } \tau \in [0, T], \text{ a.s.} \end{aligned} \quad (4.5)$$

Furthermore the objective value of $\bar{u}(\cdot)$ is given by

$$\mathbf{J}(\bar{u}(\cdot)) = \bar{\theta}(0, x_0) + G(\bar{g}(0, x_0)). \quad (4.6)$$

A proof of the above theorem will be carried out in the following section. Analogous to the classical stochastic maximum principle, the optimal state equation, the corresponding BSPDEs (4.1)- (4.2), along with the minimum condition (4.5), can be written as the following system of FBSPDEs:

$$\left\{ \begin{array}{l} d\bar{X}(t) = b(t, \bar{X}(t), \bar{u}(t)) dt + \sigma(t, \bar{X}(t)) dW(t), \text{ for } t \in [0, T], \\ d\bar{\theta}(t, x) = - \{ \langle \bar{\theta}_x(t, x), b(t, x, \bar{u}(t)) \rangle + \frac{1}{2} \text{tr} [\sigma(t, x) \sigma(t, x)^\top \bar{\theta}_{xx}(t, x)] \\ \quad + \text{tr} [\bar{\psi}_x(t, x) \sigma(t, x)] + f(t, x, \bar{u}(t)) \} dt \\ \quad + \bar{\psi}(t, x)^\top dW(t), \text{ for } (t, x) \in [0, T] \times \mathbb{R}^n, \\ d\bar{g}^i(t, x) = - \{ \langle \bar{g}_x^i(t, x), b^{\bar{u}(\cdot)}(t, x) \rangle + \text{tr} [\sigma(t, x) \sigma(t, x)^\top \bar{g}_{xx}^i(t, x)] \\ \quad + \text{tr} [\bar{\eta}_x^i(t, x) \sigma(t, x)] \} dt + \bar{\eta}^i(t, x)^\top dW(t), \\ \quad \text{for } (t, x) \in [0, T] \times \mathbb{R}^n, \\ \bar{g}^i(T, x) = x_i, \text{ for } x \in \mathbb{R}^n, 1 \leq i \leq n, \\ \bar{X}(0) = x_0, \bar{\theta}(T, x) = h(x), \text{ for } x \in \mathbb{R}^n, \\ \bar{u}(t) \in \arg \min_{u \in U} \{ \langle b(t, \bar{X}(t), u), \bar{\theta}_x(t, \bar{X}(t)) \rangle + f(t, \bar{X}(t), u) \\ \quad + \sum_{i=1}^n \langle b(t, \bar{X}(t), u), G_{\bar{x}_i}(\mathbb{E}[\bar{X}(T)]) \bar{g}_x^i(t, \bar{X}(t)) \rangle \}, \\ \quad \text{a.e. } t \in [0, T], \text{ a.s.} \end{array} \right. \quad (4.7)$$

It should be noted that, the above system of FBSPDEs is quite different from the traditional stochastic (extended) Hamiltonian system (see e.g. Yong and Zhou [38]), since two backward stochastic partial differential equations are involved. We can rephrase Theorem 4.2 as the following.

Theorem 4.3. *Let Assumption (\mathbf{H}_m) be satisfied with $m > 2 + \frac{n}{2}$. If $(\bar{u}(\cdot), \bar{X}(\cdot))$ is an optimal pair of Problem (S), then the family of processes (resp. random fields) $(\bar{X}(\cdot), \bar{u}(\cdot), \{\bar{g}^i(\cdot, \cdot), \bar{\eta}^i(\cdot, \cdot)\}_{i=1, \dots, n}, \bar{\theta}(\cdot, \cdot), \bar{\psi}(\cdot, \cdot))$ satisfies the system of FBSPDEs (4.7). Furthermore the objective value of $\bar{u}(\cdot)$ is given by (4.6).*

Remark 4.4. On comparing between the SMP in Theorem 4.2 and the traditional SMP of Bensoussan [4] we find the following facts:

- (i) The advantage: The new version of the SMP enables us to derive simultaneously an optimal solution $\bar{u}(\cdot)$ as well as its objective value $\mathbf{J}(\bar{u}(\cdot))$ in terms of the solution of FBSPDEs (4.7), while the classical SMP approach permits us to derive the optimal solution $\bar{u}(\cdot)$ only.

- (ii) The disadvantage: Of course, the disadvantage of the new version of the SMP lies in the fact of assuming differentiability of higher orders in x on the coefficients, which unfortunately limits the scope of problems applicable to the theorem.

Remark 4.5. If the function $U \ni u \rightarrow \langle b(t, x, u), p \rangle + f(t, x, u) \in \mathbb{R}$ has a unique minimizer $\bar{\mathbf{u}}(t, x, p)$ for each $t \in [0, T]$ and $(x, p) \in \mathbb{R}^n \times \mathbb{R}^n$ such that $\bar{\mathbf{u}}(t, x, p)$ satisfies an appropriate regularity condition, then the optimal control is characterized by

$$\bar{u}(t) \equiv \bar{\mathbf{u}}(t, \bar{X}(t), \bar{\theta}_x(t, \bar{X}(t)) + \sum_{i=1}^n G_{\bar{x}_i}(\mathbb{E}[\bar{X}(T)]) \bar{g}_x^i(t, \bar{X}(t))),$$

where $(\bar{X}(\cdot), \{\bar{g}^i(\cdot, \cdot), \bar{\eta}^i(\cdot, \cdot)\}_{i=1, \dots, n}, \bar{\theta}(\cdot, \cdot), \bar{\psi}(\cdot, \cdot))$ is family of processes (resp. random fields) that satisfies the following system of coupled FBSPDEs:

$$\left\{ \begin{array}{l} d\bar{X}(t) = b(t, \bar{X}(t), \bar{u}(t, \bar{X}(t))) dt + \sigma(t, \bar{X}(t)) dW(t), \text{ for } t \in [0, T], \\ d\bar{\theta}(t, x) = - \left\{ \langle \bar{\theta}_x(t, x), b(t, x, \bar{u}(t, \bar{X}(t))) \rangle + \frac{1}{2} \text{tr} \left[\sigma(t, x) \sigma(t, x)^\top \bar{\theta}_{xx}(t, x) \right] \right. \\ \quad \left. + \text{tr} [\bar{\psi}_x(t, x) \sigma(t, x)] + f(t, x, \bar{u}(t, \bar{X}(t))) \right\} dt \\ \quad + \bar{\psi}(t, x)^\top dW(t), \text{ for } (t, x) \in [0, T] \times \mathbb{R}^n, \\ d\bar{g}^i(t, x) = - \left\{ \langle \bar{g}_x^i(t, x), b(t, x, \bar{u}(t, \bar{X}(t))) \rangle + \text{tr} \left[\sigma(t, x) \sigma(t, x)^\top \bar{g}_{xx}^i(t, x) \right] \right. \\ \quad \left. + \text{tr} [\bar{\eta}_x^i(t, x) \sigma(t, x)] \right\} dt + \bar{\eta}^i(t, x)^\top dW(t), \\ \quad \text{for } (t, x) \in [0, T] \times \mathbb{R}^n, \\ \bar{g}^i(T, x) = x_i, \text{ for } x \in \mathbb{R}^n, 1 \leq i \leq n, \\ \bar{X}(0) = x_0, \bar{\theta}(T, x) = h(x), \text{ for } x \in \mathbb{R}^n, \\ \bar{u}(t, \bar{X}(t)) := \bar{\mathbf{u}}(t, \bar{X}(t), \bar{\theta}_x(t, \bar{X}(t)) + \sum_{i=1}^n G_{\bar{x}_i}(\mathbb{E}[\bar{X}(T)]) \bar{g}_x^i(t, \bar{X}(t))). \end{array} \right. \quad (4.8)$$

It is well known that optimal control theory can be used to solve stochastic (extended) Hamiltonian systems, which are actually coupled forward-backward stochastic differential equation (FBSDE, for short); see e.g. Yong and Zhou [38]. However, it is seen from Theorem 4.3 that stochastic control theory can also be used to solve coupled equations of the type (4.8). In Section 7 we will study more about this type of equations.

5 A Proof of Theorem 4.2

This section is devoted to the proof of Theorem 4.2. Let $(\bar{u}(\cdot), \bar{X}(\cdot))$ be the given optimal pair. Then the following is satisfied

$$\left\{ \begin{array}{l} d\bar{X}(t) = b^{\bar{u}(\cdot)}(t, \bar{X}(t)) dt + \sigma(t, \bar{X}(t)) dW(t), t \in [0, T], \\ \bar{X}(0) = x_0. \end{array} \right.$$

For any $\tau \in [0, T)$, $u(\cdot) \in \mathcal{U}[0, T]$ and for any $\varepsilon \in [0, T - \tau)$, define

$$u^\varepsilon(t) = \begin{cases} u(t), & \text{for } t \in [\tau, \tau + \varepsilon), \\ \bar{u}(t), & \text{for } t \in [0, T] \setminus [\tau, \tau + \varepsilon). \end{cases} \quad (5.1)$$

Let $(u^\varepsilon(\cdot), X^\varepsilon(\cdot))$ satisfy the following

$$\left\{ \begin{array}{l} dX^\varepsilon(t) = b^{u^\varepsilon(\cdot)}(t, X^\varepsilon(t)) dt + \sigma(t, X^\varepsilon(t)) dW(t), t \in [0, T], \\ X^\varepsilon(0) = x_0. \end{array} \right. \quad (5.2)$$

To prove Theorem 4.2 we need to some preliminary results given in the following two lemmas.

Lemma 5.1. *Suppose that Assumption (\mathbf{H}_m) be satisfied with $m > 2 + \frac{n}{2}$. Let $(\bar{\theta}(\cdot, \cdot), \bar{\psi}(\cdot, \cdot))$ and $(\bar{g}^i(\cdot, \cdot), \bar{\eta}^i(\cdot, \cdot))$, for $1 \leq i \leq n$, be the unique classical solutions of (4.1) and (4.2), respectively. Then for each $t \in [0, T]$ and $1 \leq i \leq n$, $\bar{\theta}(t, \bar{X}(t))$ and $\bar{g}^i(t, \bar{X}(t))$ have the following probabilistic representations:*

$$\bar{\theta}(t, \bar{X}(t)) = \mathbb{E}_t \left[\int_t^T f(s, \bar{X}(s), \bar{u}(s)) ds + F(\bar{X}(T)) \right], \quad (5.3)$$

and

$$\bar{g}^i(t, \bar{X}(t)) = \mathbb{E}_t [\bar{X}_i(T)] \quad (5.4)$$

where $\bar{X}_i(T)$ denotes the i -th coordinate of $\bar{X}(T) = (\bar{X}_1(T), \dots, \bar{X}_n(T))^\top$. Furthermore, for any $\tau \in [0, T]$, $u(\cdot) \in \mathcal{U}[0, T]$ and for any $\varepsilon \in [0, T - \tau)$, the following equality holds

$$\begin{aligned} & \mathbf{J}(u^\varepsilon(\cdot)) - \mathbf{J}(\bar{u}(\cdot)) \\ &= \mathbb{E} \left[\int_\tau^{\tau+\varepsilon} \left\{ \left\langle b^{u^\varepsilon(\cdot)}(t, X^\varepsilon(t)) - b^{\bar{u}(\cdot)}(t, X^\varepsilon(t)), \bar{\theta}_x(t, X^\varepsilon(t)) \right\rangle \right. \right. \\ & \quad \left. \left. + \sum_{i=1}^n \left\langle b^{u^\varepsilon(\cdot)}(t, X^\varepsilon(t)) - b^{\bar{u}(\cdot)}(t, X^\varepsilon(t)), G_{\bar{x}_i}(\mathbb{E}[\bar{g}(t, X^\varepsilon(t))]) \bar{g}_x^i(t, X^\varepsilon(t)) \right\rangle \right. \right. \\ & \quad \left. \left. + f^{u^\varepsilon(\cdot)}(t, X^\varepsilon(t)) - f^{\bar{u}(\cdot)}(t, X^\varepsilon(t)) \right\} dt \right]. \quad (5.5) \end{aligned}$$

Proof. Equalities (5.3)-(5.4) can be easily obtained by applying Theorem 2.2. So we only need to show (5.5). Consider the difference

$$\begin{aligned} & \mathbf{J}(u^\varepsilon(\cdot)) - \mathbf{J}(\bar{u}(\cdot)) \\ &= \mathbb{E} \left[\int_0^T f^{u^\varepsilon(\cdot)}(t, X^\varepsilon(t)) dt + h(X^\varepsilon(T)) + G(\mathbb{E}[X^\varepsilon(T)]) \right] \\ & \quad - \mathbb{E} \left[\int_0^T f^{\bar{u}(\cdot)}(t, \bar{X}(t)) dt + h(\bar{X}(T)) + G(\mathbb{E}[\bar{X}(T)]) \right]. \end{aligned}$$

By the terminal conditions in BSPDEs (4.1)-(4.2) and by setting $t = 0$ in (5.3)-(5.4), we obtain that

$$\begin{aligned} \mathbf{J}(u^\varepsilon(\cdot)) &= \mathbb{E} \left[\int_0^T f^{u^\varepsilon(\cdot)}(t, X^\varepsilon(t)) dt + \bar{\theta}(T, X^\varepsilon(T)) \right] \\ & \quad + G(\mathbb{E}[\bar{g}^1(T, X^\varepsilon(T))], \dots, \mathbb{E}[\bar{g}^n(T, X^\varepsilon(T))]) \end{aligned}$$

and

$$\mathbf{J}(\bar{u}(\cdot)) = \bar{\theta}(0, x_0) + G(\mathbb{E}[\bar{g}^1(0, x_0)], \dots, \mathbb{E}[\bar{g}^n(0, x_0)]).$$

Accordingly, we have

$$\begin{aligned} & \mathbf{J}(u^\varepsilon(\cdot)) - \mathbf{J}(\bar{u}(\cdot)) \\ &= \mathbb{E} \left[\int_0^T f^{u^\varepsilon(\cdot)}(t, X^\varepsilon(t)) dt + \bar{\theta}(T, X^\varepsilon(T)) - \bar{\theta}(0, x_0) \right] \\ & \quad + G(\mathbb{E}[\bar{g}^1(T, X^\varepsilon(T))], \dots, \mathbb{E}[\bar{g}^n(T, X^\varepsilon(T))]) \\ & \quad - G(\mathbb{E}[\bar{g}^1(0, x_0)], \dots, \mathbb{E}[\bar{g}^n(0, x_0)]). \quad (5.6) \end{aligned}$$

Recall that $X^\varepsilon(0) = x_0$. Then by applying Itô-Wentzell formula to $\bar{\theta}(t, X^\varepsilon(t))$ on time interval $[0, T]$, we get

$$\begin{aligned} & \mathbb{E}[\bar{\theta}(T, X^\varepsilon(T)) - \bar{\theta}(0, x_0)] \\ &= \mathbb{E}[\bar{\theta}(T, X^\varepsilon(T)) - \bar{\theta}(0, X^\varepsilon(0))] \\ &= \mathbb{E} \left[\int_0^T \left\{ \left\langle \bar{\theta}_x(t, X^\varepsilon(t)), b^{u^\varepsilon(\cdot)}(t, X^\varepsilon(t)) - b^{\bar{u}(\cdot)}(t, X^\varepsilon(t)) \right\rangle \right. \right. \\ & \quad \left. \left. - f^{\bar{u}(\cdot)}(t, X^\varepsilon(t)) \right\} dt \right]. \quad (5.7) \end{aligned}$$

and it follows from the chain rule applied to $G(\mathbb{E}[\bar{g}^1(t, X^\varepsilon(t))], \dots, \mathbb{E}[\bar{g}^n(t, X^\varepsilon(t))])$ that

$$\begin{aligned} & G(\mathbb{E}[\bar{g}^1(T, X^\varepsilon(T))], \dots, \mathbb{E}[\bar{g}^n(T, X^\varepsilon(T))]) \\ & \quad - G(\mathbb{E}[\bar{g}^1(0, x_0)], \dots, \mathbb{E}[\bar{g}^n(0, x_0)]) \\ &= \sum_{i=1}^n \int_0^T G_{\bar{x}_i}(\mathbb{E}[\bar{g}^1(t, X^\varepsilon(t))], \dots, \mathbb{E}[\bar{g}^n(t, X^\varepsilon(t))]) d\mathbb{E}[\bar{g}^i(t, X^\varepsilon(t))] \\ &= \sum_{i=1}^n \int_0^T G_{\bar{x}_i}(\mathbb{E}[\bar{g}(t, X^\varepsilon(t))]) d\mathbb{E}[\bar{g}^i(t, X^\varepsilon(t))], \quad (5.8) \end{aligned}$$

where

$$\begin{aligned} & d\mathbb{E} [\bar{g}^i(t, X^\varepsilon(t))] \\ &= \mathbb{E} \left[\left\langle \bar{g}_x^i(t, X^\varepsilon(t)), b^{u^\varepsilon(\cdot)}(t, X^\varepsilon(t)) - b^{\bar{u}(\cdot)}(t, X^\varepsilon(t)) \right\rangle \right] dt. \end{aligned}$$

Combining (5.6) together with (5.7) and (5.8), it follows that

$$\begin{aligned} & \mathbf{J}(u^\varepsilon(\cdot)) - \mathbf{J}(\bar{u}(\cdot)) \\ &= \mathbb{E} \left[\int_0^T \left\{ \left\langle \bar{\theta}_x(t, X^\varepsilon(t)), b^{u^\varepsilon(\cdot)}(t, X^\varepsilon(t)) - b^{\bar{u}(\cdot)}(t, X^\varepsilon(t)) \right\rangle \right. \right. \\ &+ \sum_{i=1}^n G_{\bar{x}_i}(\mathbb{E}[\bar{g}(t, X^\varepsilon(t))]) \left\langle \bar{g}_x^i(t, X^\varepsilon(t)), b^{u^\varepsilon(\cdot)}(t, X^\varepsilon(t)) - b^{\bar{u}(\cdot)}(t, X^\varepsilon(t)) \right\rangle \\ &\left. \left. + f^{u^\varepsilon(\cdot)}(t, X^\varepsilon(t)) - f^{\bar{u}(\cdot)}(t, X^\varepsilon(t)) \right\} dt \right]. \end{aligned}$$

Since $u^\varepsilon(t) = u(t)\chi_{[t, t+\varepsilon]}(t) + \bar{u}(t)\chi_{[0, T]/[\tau, \tau+\varepsilon]}(t)$, we obtain that

$$\begin{aligned} & \mathbf{J}(u^\varepsilon(\cdot)) - \mathbf{J}(\bar{u}(\cdot)) \\ &= \mathbb{E} \left[\int_\tau^{\tau+\varepsilon} \left\{ \left\langle \bar{\theta}_x(t, X^\varepsilon(t)), b^{u(\cdot)}(t, X^\varepsilon(t)) - b^{\bar{u}(\cdot)}(t, X^\varepsilon(t)) \right\rangle \right. \right. \\ &+ \sum_{i=1}^n G_{\bar{x}_i}(\mathbb{E}[\bar{g}(t, X^\varepsilon(t))]) \left\langle \bar{g}_x^i(t, X^\varepsilon(t)), b^{u(\cdot)}(t, X^\varepsilon(t)) - b^{\bar{u}(\cdot)}(t, X^\varepsilon(t)) \right\rangle \\ &\left. \left. + f^{u(\cdot)}(t, X^\varepsilon(t)) - f^{\bar{u}(\cdot)}(t, X^\varepsilon(t)) \right\} dt \right]. \end{aligned}$$

This completes the proof. ■

Lemma 5.2. *Suppose that Assumption (\mathbf{H}_m) be satisfied with $m > 2 + \frac{n}{2}$. Let $(\bar{\theta}(\cdot, \cdot), \bar{\psi}(\cdot, \cdot))$ and $(\bar{g}^i(\cdot, \cdot), \bar{\eta}^i(\cdot, \cdot))$, for $1 \leq i \leq n$, be the unique classical solutions of (4.1) and (4.2), respectively. Then for any $\tau \in [0, T]$, $u(\cdot) \in \mathcal{U}[0, T]$ and for any $\varepsilon \in [0, T - \tau]$, the following equality holds*

$$\begin{aligned} & \mathbf{J}(u^\varepsilon(\cdot)) - \mathbf{J}(\bar{u}(\cdot)) \\ &= \mathbb{E} \left[\int_\tau^{\tau+\varepsilon} \left\{ \left\langle b^{u(\cdot)}(t, \bar{X}(t)) - b^{\bar{u}(\cdot)}(t, \bar{X}(t)), \bar{\theta}_x(t, \bar{X}(t)) \right\rangle \right. \right. \\ &+ \sum_{i=1}^n \left\langle b^{u(\cdot)}(t, \bar{X}(t)) - b^{\bar{u}(\cdot)}(t, \bar{X}(t)), G_{\bar{x}_i}(\mathbb{E}[\bar{g}(t, \bar{X}(t))]) \bar{g}_x^i(t, \bar{X}(t)) \right\rangle \\ &\left. \left. f^{u(\cdot)}(t, \bar{X}(t)) - f^{\bar{u}(\cdot)}(t, \bar{X}(t)) \right\} dt \right] + o(\varepsilon). \end{aligned}$$

Proof. Let $\xi^\varepsilon(\cdot) := X^\varepsilon(\cdot) - \bar{X}(\cdot)$. Then we have $\xi^\varepsilon(\cdot)$ satisfies the following SDE,

$$\begin{cases} d\xi^\varepsilon(t) = \left\{ \tilde{b}^\varepsilon(t) \xi^\varepsilon(t) + \left\{ b^{u^\varepsilon(\cdot)}(t, \bar{X}(t)) - b^{\bar{u}(\cdot)}(t, \bar{X}(t)) \right\} \chi_{[\tau, \tau+\varepsilon]}(t) \right\} dt \\ \quad + \sum_{j=1}^d \left\{ \tilde{\sigma}^j(t) \xi^\varepsilon(t) \right\} dW_j(t), \quad t \in [0, T], \\ \xi^\varepsilon(0) = 0, \end{cases}$$

where

$$\tilde{b}(t) := \int_0^1 b_x^{u^\varepsilon(\cdot)}(t, \bar{X}(t) + r\xi^\varepsilon(t)) dr$$

and

$$\tilde{\sigma}^j(t) := \int_0^1 \sigma_x^j(t, \bar{X}(t) + r\xi^\varepsilon(t)) dr.$$

By Lemma 4.2 in [[38], p. 124], we obtain for any $k \geq 1$

$$\begin{aligned} \sup_{t \in [0, T]} \mathbb{E} \left[|\xi^\varepsilon(t)|^{2k} \right] &\leq K \left(\int_\tau^{\tau+\varepsilon} \mathbb{E} \left[\left| b^{u(\cdot)}(t, \bar{X}(t)) - b^{\bar{u}(\cdot)}(t, \bar{X}(t)) \right|^{2k} \right]^{\frac{1}{2k}} dt \right)^{2k} \\ &\leq K\varepsilon^{2k}, \end{aligned}$$

for some $K > 0$.

We now consider the difference

$$\begin{aligned}
\rho(\varepsilon) &= \{\mathbf{J}(u^\varepsilon(\cdot)) - \mathbf{J}(\bar{u}(\cdot))\} \\
&\quad - \mathbb{E} \left[\int_{\tau}^{\tau+\varepsilon} \left\{ \left\langle b^{u(\cdot)}(t, \bar{X}(t)) - b^{\bar{u}(\cdot)}(t, \bar{X}(t)), \bar{\theta}_x(t, \bar{X}(t)) \right\rangle \right. \right. \\
&\quad + \left. \left\langle b(t, \bar{X}(t), u(t)) - b(t, \bar{X}(t), \bar{u}(t)), \sum_{i=1}^n G_{\bar{x}_i}(\mathbb{E}[\bar{g}(\tau, \bar{X}(\tau))]) \bar{g}_x^i(\tau, \bar{X}(\tau)) \right\rangle \right. \\
&\quad \left. \left. + f^{u(\cdot)}(t, \bar{X}(t)) - f^{\bar{u}(\cdot)}(t, \bar{X}(t)) \right\} dt \right] \\
&= \mathbb{E} \left[\int_{\tau}^{\tau+\varepsilon} \left\{ \left\langle b^{u(\cdot)}(t, X^\varepsilon(t)) - b^{\bar{u}(\cdot)}(t, X^\varepsilon(t)), \bar{\theta}_x(t, X^\varepsilon(t)) \right\rangle \right. \right. \\
&\quad - \left. \left\langle b^{u(\cdot)}(t, \bar{X}(t)) - b^{\bar{u}(\cdot)}(t, \bar{X}(t)), \bar{\theta}_x(t, \bar{X}(t)) \right\rangle \right. \\
&\quad + \left. \left\langle b^{u(\cdot)}(t, X^\varepsilon(t)) - b^{\bar{u}(\cdot)}(t, X^\varepsilon(t)), \sum_{i=1}^n G_{\bar{x}_i}(\mathbb{E}[\bar{g}(t, X^\varepsilon(t))]) \bar{g}_x^i(t, X^\varepsilon(t)) \right\rangle \right. \\
&\quad - \left. \left\langle b^{u(\cdot)}(t, \bar{X}(t)) - b^{\bar{u}(\cdot)}(t, \bar{X}(t)), \sum_{i=1}^n G_{\bar{x}_i}(\mathbb{E}[\bar{g}(t, \bar{X}(t))]) \bar{g}_x^i(t, \bar{X}(t)) \right\rangle \right. \\
&\quad \left. \left. + f^{u(\cdot)}(t, X^\varepsilon(t)) - f^{\bar{u}(\cdot)}(t, X^\varepsilon(t)) - f^{u(\cdot)}(t, \bar{X}(t)) + f^{\bar{u}(\cdot)}(t, \bar{X}(t)) \right\} dt \right].
\end{aligned}$$

Thus

$$\begin{aligned}
|\rho(\varepsilon)| &\leq \mathbb{E} \left[\int_{\tau}^{\tau+\varepsilon} \left\{ \left| b^{u(\cdot)}(t, X^\varepsilon(t)) - b^{\bar{u}(\cdot)}(t, X^\varepsilon(t)) \right| \left| \bar{\theta}_x(t, X^\varepsilon(t)) - \bar{\theta}_x(t, \bar{X}(t)) \right| \right. \right. \\
&\quad + \left| \bar{\theta}_x(t, \bar{X}(t)) \right| \left| b^{u(\cdot)}(t, X^\varepsilon(t)) - b^{u(\cdot)}(t, \bar{X}(t)) \right| \\
&\quad + \left| \bar{\theta}_x(t, \bar{X}(t)) \right| \left| b^{\bar{u}(\cdot)}(t, X^\varepsilon(t)) - b^{\bar{u}(\cdot)}(t, \bar{X}(t)) \right| \\
&\quad + \left| f^{u(\cdot)}(t, X^\varepsilon(t)) - f^{u(\cdot)}(t, \bar{X}(t)) \right| + \left| f^{\bar{u}(\cdot)}(t, X^\varepsilon(t)) - f^{\bar{u}(\cdot)}(t, \bar{X}(t)) \right| \\
&\quad + \sum_{i=1}^n |G_{\bar{x}_i}(\mathbb{E}[\bar{g}(t, X^\varepsilon(t))])| \left| g_x^i(t, X^\varepsilon(t)) - g_x^i(t, \bar{X}(t)) \right| \left| \sigma^{u(\cdot)}(t, X^\varepsilon(t)) - \sigma^{\bar{u}(\cdot)}(t, X^\varepsilon(t)) \right| \\
&\quad + \sum_{i=1}^n |\bar{g}_x^i(t, \bar{X}(t))| \left| G_{\bar{x}_i}(\mathbb{E}[\bar{g}(t, X^\varepsilon(t))]) - G_{\bar{x}_i}(\mathbb{E}[\bar{g}(t, \bar{X}(t))]) \right| \left| \sigma^{u(\cdot)}(t, X^\varepsilon(t)) - \sigma^{\bar{u}(\cdot)}(t, X^\varepsilon(t)) \right| \\
&\quad + \sum_{i=1}^n |G_{\bar{x}_i}(\mathbb{E}[\bar{g}(t, \bar{X}(t))])| \left| \bar{g}_x^i(t, \bar{X}(t)) \right| \left| \sigma^{u(\cdot)}(t, X^\varepsilon(t)) - \sigma^{u(\cdot)}(t, \bar{X}(t)) \right| \\
&\quad \left. \left. + \sum_{i=1}^n |G_{\bar{x}_i}(\mathbb{E}[\bar{g}(t, \bar{X}(t))])| \left| \bar{g}_x^i(t, \bar{X}(t)) \right| \left| \sigma^{u(\cdot)}(t, X^\varepsilon(t)) - \sigma^{u(\cdot)}(t, \bar{X}(t)) \right| \right\} dt \right].
\end{aligned}$$

Using the fact that the random fields $b_x^{u(\cdot)}$, $b_x^{\bar{u}(\cdot)}$, $f_x^{u(\cdot)}$, $f_x^{\bar{u}(\cdot)}$, $G_{\bar{x}}$, $G_{\bar{x}\bar{x}}$, $\bar{\theta}_x$, $\bar{\theta}_{xx}$, \bar{g}_x^i , \bar{g}_{xx}^i are uniformly bounded, it is not difficult to show that there exists a constant $K > 0$ such that

$$\begin{aligned}
|\rho(\varepsilon)| &\leq K \int_{\tau}^{\tau+\varepsilon} \mathbb{E} [|\bar{X}(t) - X^\varepsilon(t)|] dt \\
&= K \mathbb{E} \left[\int_{\tau}^{\tau+\varepsilon} \mathbb{E} [|\xi^\varepsilon(t)|] dt \right] \\
&\leq K \varepsilon \sup_{t \in [0, T]} \mathbb{E} [|\xi^\varepsilon(t)|^{2k}]^{\frac{1}{2k}} \\
&\leq K \varepsilon^2.
\end{aligned}$$

Hence

$$|\rho(\varepsilon)| \leq o(\varepsilon).$$

This completes the proof. ■

Now, we are ready to give a proof of Theorem 4.2.

Proof of Theorem 4.2. From Lemma 5.2, we deduce that

$$\begin{aligned}
0 &\leq \lim_{\varepsilon \downarrow 0} \frac{1}{\varepsilon} \{ \mathbf{J}(u^\varepsilon(\cdot)) - \mathbf{J}(\bar{u}(\cdot)) \} \\
&= \lim_{\varepsilon \downarrow 0} \frac{1}{\varepsilon} \int_{\tau}^{\tau+\varepsilon} \mathbb{E} [\langle b(t, \bar{X}(t), u(t)) - b(t, \bar{X}(t), \bar{u}(t)), \bar{\theta}_x(t, \bar{X}(t)) \rangle \\
&\quad + \sum_{i=1}^n \langle b(t, \bar{X}(t), u(t)) - b(t, \bar{X}(t), \bar{u}(t)), G_{\bar{x}_i}(\mathbb{E}[\bar{g}(t, \bar{X}(t))]) \bar{g}_x^i(t, \bar{X}(t)) \rangle \\
&\quad + (f(t, \bar{X}(t), u(t)) - f(t, \bar{X}(t), \bar{u}(t))) \rangle dt \\
&= \lim_{\varepsilon \downarrow 0} \frac{1}{\varepsilon} \int_{\tau}^{\tau+\varepsilon} \mathbb{E} [\langle b(t, \bar{X}(t), u(t)) - b(t, \bar{X}(t), \bar{u}(t)), \bar{\theta}_x(t, \bar{X}(t)) \rangle \\
&\quad + \sum_{i=1}^n \langle b(t, \bar{X}(t), u(t)) - b(t, \bar{X}(t), \bar{u}(t)), G_{\bar{x}_i}(\mathbb{E}[\bar{X}(T)]) \bar{g}_x^i(t, \bar{X}(t)) \rangle \\
&\quad + f(t, \bar{X}(t), u(t)) - f(t, \bar{X}(t), \bar{u}(t))] dt,
\end{aligned}$$

where we have used the following equality,

$$\begin{aligned}
\mathbb{E}[\bar{g}(t, \bar{X}(t))] &= (\mathbb{E}[\bar{g}^1(t, \bar{X}(t))], \dots, \mathbb{E}[\bar{g}^n(t, \bar{X}(t))])^\top \\
&= (\mathbb{E}[\mathbb{E}_t[\bar{X}_1(T)]], \dots, \mathbb{E}[\mathbb{E}_t[\bar{X}_n(T)]])^\top \\
&= \mathbb{E}[\bar{X}(T)].
\end{aligned}$$

Thus by Lebesgue differentiation theorem

$$\begin{aligned}
0 &\leq \mathbb{E} \left[\left\langle b(\tau, \bar{X}(\tau), u(\tau)) - b(\tau, \bar{X}(\tau), \bar{u}(\tau)), \theta_x(\tau, \hat{X}(\tau)) \right\rangle \right. \\
&\quad + \sum_{i=1}^n \langle b(\tau, \bar{X}(\tau), u(\tau)) - b(\tau, \bar{X}(\tau), \bar{u}(\tau)), G_{\bar{x}_i}(\mathbb{E}[\bar{X}(T)]) \bar{g}_x^i(\tau, \bar{X}(\tau)) \rangle \\
&\quad \left. + f(\tau, \bar{X}(\tau), u(\tau)) - f(\tau, \bar{X}(\tau), \bar{u}(\tau)) \right], \text{ a.e. } \tau \in [0, T].
\end{aligned} \tag{5.9}$$

Now let

$$u(\tau) = u\chi_A + \bar{u}(\tau)\chi_{\Omega \setminus A},$$

where A is an arbitrarily \mathcal{F}_τ -measurable set and $u \in U$.

Hence, in view of (5.9), we have

$$\begin{aligned}
0 &\leq \mathbb{E} \left[\left\langle b(\tau, \bar{X}(\tau), u) - b(\tau, \bar{X}(\tau), \bar{u}(\tau)), \theta_x(\tau, \hat{X}(\tau)) \right\rangle \right. \\
&\quad + \sum_{i=1}^n \langle b(\tau, \bar{X}(\tau), u) - b(\tau, \bar{X}(\tau), \bar{u}(\tau)), G_{\bar{x}_i}(\mathbb{E}[\bar{X}(T)]) \bar{g}_x^i(\tau, \bar{X}(\tau)) \rangle \\
&\quad \left. + f(\tau, \bar{X}(\tau), u) - f(\tau, \bar{X}(\tau), \bar{u}(\tau)) \right\} \chi_A \Big].
\end{aligned}$$

which in turn yields inequality (4.4) since $u \in U$ and the set $A \in \mathcal{F}_\tau$ are arbitrary. This completes the proof. \blacksquare

6 Sufficient Condition of Optimality

In this section, we focus on proving that provided some convexity assumptions are satisfied, the necessary condition (4.5) turns out to be sufficient.

The following fact concerning the differentiability of stochastic integrals with parameter is important for our purpose. Let $\phi(\cdot, \cdot)$ be a bounded random field such that $\phi(\cdot, \cdot) \in \mathcal{L}_{\mathcal{F}}^2(0, T; C_b^2(\mathbb{R}; \mathbb{R}))$. Then it was shown that (see, e.g. [Kunita [19], Proposition 2.3.1, Page 56] or [Kunita [18], Exercise 3.1.5, Page 78]) the stochastic integral with parameter: $\int_0^\cdot \phi(t, \cdot) dW(t)$ has a modification that belongs to $\mathcal{L}_{\mathcal{F}}^2(0, T; C^1(\mathbb{R}^n; \mathbb{R}))$ and it satisfies

$$\partial_x \int_0^t \phi(t, x) dW(t) = \int_0^t \partial_x \phi(t, x) dW(t), \tag{6.1}$$

where $\partial_x \phi(t, x)$ denote the first derivative of $\phi(t, x)$ with respect to the variable x .

Define the usual Hamiltonian as a map from $[0, T] \times \mathbb{R}^n \times U \times \mathbb{R}^n \times \mathbb{R}^{n \times d}$ into \mathbb{R} by

$$\mathbb{H}(t, x, u, p, q) := \langle b(t, x, u), p \rangle + \mathbf{tr} \left[\sigma(t, x)^\top q \right] + f(t, x, u) \tag{6.2}$$

and let us introduce two additional assumptions.

- (H1) The control domain $U \subset \mathbb{R}^l$ is a convex subset and the maps b and f are continuously differentiable with respect to u .
- (H2) The Hamiltonian $\mathbb{H}(t, x, u, p, q)$ is convex with respect to (x, u) , $h(\cdot)$ is convex with respect to x and $G(\cdot)$ is convex with respect to \bar{x} .

Theorem 6.1 (Sufficient Condition of Optimality). *Let Assumption (H1), (H2) and (H_m) be satisfied with $m > 2 + \frac{n}{2}$. Let $(\bar{u}(\cdot), \bar{X}(\cdot))$ be an admissible pair and $(\bar{\theta}(\cdot, \cdot), \bar{\psi}(\cdot, \cdot))$, $(\bar{g}^i(\cdot, \cdot), \bar{\eta}^i(\cdot, \cdot))$, for $1 \leq i \leq n$, be the solutions of the BSPDEs (4.1) and (4.2), respectively. Suppose that*

$$\bar{\psi}(\cdot, \cdot) \in \mathcal{L}_{\mathcal{F}}^2(0, T; C_b^2(\mathbb{R}^n; \mathbb{R}^d)) \quad \text{and} \quad \bar{\eta}^i(\cdot, \cdot) \in \mathcal{L}_{\mathcal{F}}^2(0, T; C_b^2(\mathbb{R}^n; \mathbb{R}^d)).$$

Suppose further, for a.e. $t \in [0, T]$,

$$\begin{aligned} \bar{u}(t) \in \arg \min_{u \in U} \{ & \langle \bar{\theta}_x(t, \bar{X}(t)), b(t, \bar{X}(t), \cdot) \rangle + f(t, \bar{X}(t), \cdot) \\ & + \sum_{i=1}^n G_{\bar{x}_i}(\mathbb{E}[\bar{X}(T)]) \langle \bar{g}_x^i(\tau, \bar{X}(\tau)), b(t, \bar{X}(t), \cdot) \rangle \}, \end{aligned} \quad (6.3)$$

a.e. $t \in [0, T]$, a.s.

Then $(\bar{u}(\cdot), \bar{X}(\cdot))$ is an optimal pair of Problem (S). Furthermore the objective value of $\bar{u}(\cdot)$ is given by

$$\mathbf{J}(\bar{u}(\cdot)) = \inf_{u(\cdot) \in \mathcal{U}[0, T]} \mathbf{J}(u(\cdot)) = \bar{\theta}(0, x_0) + G(\bar{g}(0, x_0)).$$

Proof. First, let us begin by some preparations. Let $(\bar{u}(\cdot), \bar{X}(\cdot))$ be an admissible pair and $(\bar{\theta}(\cdot, \cdot), \bar{\psi}(\cdot, \cdot))$, $(\bar{g}^i(\cdot, \cdot), \bar{\eta}^i(\cdot, \cdot))$, for $1 \leq i \leq n$, be the solutions of the following BSPDEs,

$$\begin{cases} d\bar{\theta}(t, x) = - \left\{ \langle \bar{\theta}_x(t, x), b^{\bar{u}(\cdot)}(t, x) \rangle + \frac{1}{2} \text{tr} [(\sigma \sigma^\top)(t, x) \bar{\theta}_{xx}(t, x)] \right. \\ \quad \left. + \text{tr} [\bar{\psi}_x(t, x) \sigma(t, x)] + f^{\bar{u}(\cdot)}(t, x) \right\} dt \\ \quad + \bar{\psi}(t, x)^\top dW(t), \quad (t, x) \in [0, T] \times \mathbb{R}^n, \\ \bar{\theta}(T, x) = h(x), \quad \text{for } x \in \mathbb{R}^n. \end{cases}$$

and

$$\begin{cases} d\bar{g}^i(t, x) = - \left\{ \langle \bar{g}_x^i(t, x), b^{\bar{u}(\cdot)}(t, x) \rangle + \frac{1}{2} \text{tr} [(\sigma \sigma^\top)(t, x) \bar{g}_{xx}^i(t, x)] \right. \\ \quad \left. + \text{tr} [\bar{\eta}_x^i(t, x)^\top \sigma(t, x)] \right\} dt + \bar{\eta}^i(t, x)^\top dW(t), \\ \quad \text{for } (t, x) \in [0, T] \times \mathbb{R}^n, \\ \bar{g}^i(T, x) = \bar{x}_i, \quad \text{for } x \in \mathbb{R}^n, \end{cases}$$

respectively, which also can be written as

$$\begin{cases} d\bar{\theta}(t, x) = - \left\{ \langle b^{\bar{u}(\cdot)}(t, x), \bar{\theta}_x(t, x) \rangle + \frac{1}{2} \sum_{j=1}^d \sigma^j(t, x)^\top \bar{\theta}_{xx}(t, x) \sigma^j(t, x) \right. \\ \quad \left. + \sum_{j=1}^d \langle \sigma^j(t, x), \bar{\psi}_x^j(t, x) \rangle + f^{\bar{u}(\cdot)}(t, x) \right\} dt \\ \quad + \sum_{j=1}^d \bar{\psi}^j(t, x) dW_j(t), \quad (t, x) \in [0, T] \times \mathbb{R}^n, \\ \bar{\theta}(T, x) = h(x), \quad \text{for } x \in \mathbb{R}^n \end{cases} \quad (6.4)$$

and

$$\begin{cases} d\bar{g}^i(t, x) = - \left\{ \langle \bar{g}_x^i(t, x), b^{\bar{u}(\cdot)}(t, x) \rangle + \frac{1}{2} \sum_{j=1}^d \sigma^j(t, x)^\top \bar{g}_{xx}^i(t, x) \sigma^j(t, x) \right. \\ \quad \left. + \sum_{j=1}^d \langle \sigma^j(t, x), \bar{\eta}_x^{ij}(t, x) \rangle \right\} dt + \sum_{j=1}^d \bar{\eta}^{ij}(t, x) dW_j(t), \\ \quad \text{for } (t, x) \in [0, T] \times \mathbb{R}^n, \\ \bar{g}^i(T, x) = \bar{x}_i, \quad \text{for } x \in \mathbb{R}^n. \end{cases} \quad (6.5)$$

By using the fact (6.1), we differentiate (6.4)-(6.5) in x to obtain that

$$\left\{ \begin{aligned} d\bar{\theta}_x(t, x) &= - \left\{ \left\langle b_x^{\bar{u}(\cdot)}(t, x), \bar{\theta}_x(t, x) \right\rangle + \bar{\theta}_{xx}(t, x)^\top b_x^{\bar{u}(\cdot)}(t, x) \right. \\ &\quad + \sum_{j=1}^d \sigma_x^j(t, x)^\top \bar{\theta}_{xx}(t, x) \sigma^j(t, x) + \frac{1}{2} \sum_{j=1}^d \sigma_x^j(t, x)^\top \bar{\theta}_{xxx}(t, x) \sigma^j(t, x) \\ &\quad + \sum_{j=1}^d \left[\left\langle \sigma_x^j(t, x), \bar{\psi}_x^j(t, x) \right\rangle + \bar{\psi}_{xx}^j(t, x)^\top \sigma_x^j(t, x) \right] + f_x^{\bar{u}(\cdot)}(t, x) \Big\} dt \\ &\quad + \sum_{j=1}^d \bar{\psi}_x^j(t, x) dW_j(t), \quad (t, x) \in [0, T] \times \mathbb{R}^n, \\ \bar{\theta}_x(T, x) &= h_x(x), \quad \text{for } x \in \mathbb{R}^n \end{aligned} \right.$$

and

$$\left\{ \begin{aligned} d\bar{g}_x^i(t, x) &= - \left\{ \left\langle b_x^{\bar{u}(\cdot)}(t, x), \bar{g}_x^i(t, x) \right\rangle + \bar{g}_{xx}^i(t, x)^\top b_x^{\bar{u}(\cdot)}(t, x) \right. \\ &\quad + \sum_{j=1}^d \sigma_x^j(t, x)^\top \bar{g}_{xx}^i(t, x) \sigma^j(t, x) + \frac{1}{2} \sum_{j=1}^d \sigma_x^j(t, x)^\top \bar{g}_{xxx}^i(t, x) \sigma^j(t, x) \\ &\quad + \sum_{j=1}^d \left[\left\langle \sigma_x^j, \bar{\eta}_x^{ij}(t, x) \right\rangle + \bar{\eta}_{xx}^{ij}(t, x)^\top \sigma_x^j(t, x) \right] \Big\} dt \\ &\quad + \sum_{j=1}^d \bar{\eta}_x^{ij}(t, x) dW_j(t), \quad (t, x) \in [0, T] \times \mathbb{R}^n, \\ \bar{g}_x^i(T, x) &= \mathbf{e}_i, \quad \text{for } x \in \mathbb{R}^n, \end{aligned} \right.$$

where for $\varrho = \bar{\theta}, \bar{g}^i$

$$(\sigma_x^j)^\top \varrho_{xxx}(\sigma^j) := \left((\sigma_x^j)^\top (\varrho_x)_{xx}^1(\sigma^j), \dots, (\sigma_x^j)^\top (\varrho_x)_{xx}^n(\sigma^j) \right),$$

with

$$\varrho_x := \left((\varrho_x)^1, \dots, (\varrho_x)^n \right)^\top.$$

Accordingly, applying Itô-Wentzell formula to $\bar{\theta}_x(t, \bar{X}(t))$ and $\bar{g}_x^i(t, \bar{X}(t))$, we get

$$\left\{ \begin{aligned} d\bar{\theta}_x(t, \bar{X}(t)) &= - \left\{ \left\langle b_x^{\bar{u}(\cdot)}(t, \bar{X}(t)), \bar{\theta}_x(t, \bar{X}(t)) \right\rangle + f_x(t, \bar{X}(t), \bar{u}(t)) \right. \\ &\quad + \sum_{j=1}^d \left\langle \sigma_x^j(t, \bar{X}(t)), \bar{\psi}_x^j(t, \bar{X}(t)) + \bar{\theta}_{xx}(t, \bar{X}(t)) \sigma^j(t, \bar{X}(t)) \right\rangle \Big\} dt \\ &\quad + \sum_{j=1}^d \left\{ \bar{\psi}_x^j(t, \bar{X}(t)) + \bar{\theta}_{xx}(t, \bar{X}(t)) \sigma^j(t, \bar{X}(t)) \right\} dW_j(t), \quad t \in [0, T], \\ \bar{\theta}_x(T, \bar{X}(T)) &= h_x(\bar{X}(T)), \end{aligned} \right.$$

and, for each $1 \leq i \leq n$,

$$\left\{ \begin{aligned} d\bar{g}_x^i(t, \bar{X}(t)) &= - \left\{ \left\langle b_x^{\bar{u}(\cdot)}(t, \bar{X}(t)), \bar{g}_x^i(t, \bar{X}(t)) \right\rangle \right. \\ &\quad + \sum_{j=1}^d \left\langle \sigma_x^j(t, \bar{X}(t)), \bar{\eta}_x^{ij}(t, \bar{X}(t)) + \bar{g}_{xx}^i(t, \bar{X}(t)) \sigma^j(t, \bar{X}(t)) \right\rangle \Big\} dt \\ &\quad + \sum_{j=1}^d \left\{ \bar{\eta}_x^{ij}(t, \bar{X}(t)) + \bar{g}_{xx}^i(t, \bar{X}(t)) \sigma^j(t, \bar{X}(t)) \right\} dW_j(t), \quad t \in [0, T], \\ \bar{g}_x^i(T, \bar{X}(T)) &= \mathbf{e}_i. \end{aligned} \right.$$

Define for $t \in [0, T]$,

$$\bar{p}(t) := \bar{\theta}_x(t, \bar{X}(t)) + \sum_{i=1}^n G_{\bar{x}_i}(\mathbb{E}[\bar{X}(T)]) \bar{g}_x^i(t, \bar{X}(t)) \quad (6.6)$$

and for each $j = 1, \dots, d$,

$$\begin{aligned} \bar{q}_j(t) &:= \bar{\psi}_x^j(t, \bar{X}(t)) + \bar{\theta}_{xx}(t, \bar{X}(t)) \sigma^j(t, \bar{X}(t)) \\ &\quad + \sum_{i=1}^n G_{\bar{x}_i}(\mathbb{E}[\bar{X}(T)]) \left\{ \bar{\eta}_x^{ij}(t, \bar{X}(t)) + \bar{g}_{xx}^i(t, \bar{X}(t)) \sigma^j(t, \bar{X}(t)) \right\}. \end{aligned} \quad (6.7)$$

Then it is easy to verify that $\bar{p}(\cdot)$ and $\bar{q}(\cdot) = (\bar{q}_1(\cdot), \dots, \bar{q}_d(\cdot))$ satisfy the following linear BSDEs

$$\begin{cases} d\bar{p}(t) = -\mathbb{H}_x(t, \bar{X}(t), \bar{u}(t), \bar{p}(t), \bar{q}(t)) dt + \sum_{j=1}^d \bar{q}^j(t) dW_j(t), & t \in [0, T], \\ \bar{p}(T) = h_x(\bar{X}(T)) + \bar{g}_x(\mathbb{E}[\bar{X}(T)]). \end{cases} \quad (6.8)$$

where \mathbb{H} is given by (6.2).

Recall that

$$\begin{aligned} \bar{u}(t) \in \arg \min_{u \in U} \{ & \langle \bar{\theta}_x(t, \bar{X}(t)), b(t, \bar{X}(t), \cdot) \rangle + f(t, \bar{X}(t), \cdot) \\ & + \sum_{i=1}^n G_{\bar{x}_i}(\mathbb{E}[\bar{X}(T)]) \langle \bar{g}_x^i(\tau, \bar{X}(\tau)), b(t, \bar{X}(t), \cdot) \rangle \}, \\ & \text{a.e. } t \in [0, T], \text{ a.s.} \end{aligned}$$

Accordingly, the first order optimality condition yields

$$\begin{aligned} 0 \leq & \left\langle u - \bar{u}(t), b_u(t, \bar{X}(t), \bar{u}(t))^\top \bar{\theta}_x(t, \bar{X}(t)) + f_u(t, \bar{X}(t), \bar{u}(t)) \right\rangle \\ & + \sum_{i=1}^n G_{\bar{x}_i}(\mathbb{E}[\bar{X}(T)]) \left\langle u - \bar{u}(t), b_u(t, \bar{X}(t), \bar{u}(t))^\top \bar{g}_x^i(t, \bar{X}(t)) \right\rangle \\ & \text{for all } u \in U, \text{ a.s., a.e. } t \in [0, T]. \end{aligned} \quad (6.9)$$

Combining this with Equalities (6.6)-(6.7), we get

$$0 \leq (u - \bar{u}(t))^\top \mathbb{H}_u(t, \hat{X}^{x_0}(t), \bar{u}(t), p(t), q(t), r(t, \cdot)), \mathbb{P}\text{-a.s., } \forall u \in U, \text{ a.e. } t \in [0, T]. \quad (6.10)$$

Now for an arbitrarily admissible state-control pair $(u(\cdot), X(\cdot))$, consider the difference

$$\begin{aligned} & \mathbf{J}(u(\cdot)) - \mathbf{J}(\bar{u}(\cdot)) \\ &= \mathbb{E} \left[\int_0^T \left\{ f^{u(\cdot)}(t, X(t)) - f^{\bar{u}(\cdot)}(t, \bar{X}(t)) \right\} dt + h(X(T)) - h(\bar{X}(T)) \right] \\ &+ \bar{g}(\mathbb{E}[X(T)]) - \bar{g}(\mathbb{E}[\bar{X}(T)]) \end{aligned}$$

By the convexity of $h(\cdot)$ and $\bar{g}(\cdot)$, we have

$$\mathbb{E}[h(X(T)) - h(\bar{X}(T))] \geq \mathbb{E}[(X(T) - \bar{X}(T))^\top h_x(\bar{X}(T))]$$

and

$$G(\mathbb{E}[X(T)]) - G(\mathbb{E}[\bar{X}(T)]) \geq \mathbb{E}[(X(T) - \bar{X}(T))^\top G_x(\mathbb{E}[\bar{X}(T)])].$$

Accordingly, by the terminal condition in the BSDE (6.8) we obtain that

$$\begin{aligned} & \mathbf{J}(u(\cdot)) - \mathbf{J}(\bar{u}(\cdot)) \\ & \geq \mathbb{E} \left[\int_0^T \left\{ f^{u(\cdot)}(t, X(t)) - f^{\bar{u}(\cdot)}(t, \bar{X}(t)) \right\} dt + (X(T) - \bar{X}(T))^\top \bar{p}(T) \right]. \end{aligned} \quad (6.11)$$

From Itô's lemma, applied to $t \mapsto (X(t) - \bar{X}(t))^\top \bar{p}(T)$ on time interval $[0, T]$, it follows that

$$\begin{aligned} & \mathbb{E} \left[(X(T) - \bar{X}(T))^\top \bar{p}(T) \right] \\ &= \mathbb{E} \left[\int_0^T \left\{ \left\langle b^{u(\cdot)}(t, X(t)) - b^{\bar{u}(\cdot)}(t, \bar{X}(t)), \bar{p}(t) \right\rangle \right. \right. \\ &+ \sum_{j=1}^d \left\langle \sigma^j(t, X(t)) - \sigma^j(t, \bar{X}(t)), \bar{q}_j(t) \right\rangle \\ &\left. \left. - \langle X(t) - \bar{X}(t), \mathbb{H}_x(t, \bar{X}(t), \bar{u}(t), \bar{p}(t), \bar{q}(t)) \rangle \right\} dt \right]. \end{aligned} \quad (6.12)$$

On the other hand, by the definition of the Hamiltonian \mathbb{H} , we have

$$\begin{aligned}
& \mathbb{E} \left[\int_0^T \{f(t, X(t), u(t)) - f(t, \bar{X}(t), \bar{u}(t))\} dt \right] \\
&= \mathbb{E} \left[\int_0^T \{ \mathbb{H}(t, X(t), u(t), \bar{p}(t), \bar{q}(t)) - \mathbb{H}(t, \bar{X}(t), \bar{u}(t), \bar{p}(t), \bar{q}(t)) \right. \\
&\quad - \sum_{j=1}^d \langle \sigma^j(t, X(t)) - \sigma^j(t, \bar{X}(t)), \bar{q}_j(t) \rangle \\
&\quad \left. - \langle b^{u(\cdot)}(t, X(t)) - b^{\bar{u}(\cdot)}(t, \bar{X}(t)), \bar{p}(t) \rangle \right] dt. \tag{6.13}
\end{aligned}$$

Invoking (6.12) and (6.13) into (6.11), we obtain that

$$\begin{aligned}
& \mathbf{J}(u(\cdot)) - \mathbf{J}(\bar{u}(\cdot)) \\
&\geq \mathbb{E} \left[\int_0^T \{ \mathbb{H}(t, X(t), u(t), \bar{p}(t), \bar{q}(t)) - \mathbb{H}(t, \bar{X}(t), \bar{u}(t), \bar{p}(t), \bar{q}(t)) \right. \\
&\quad \left. - \langle X(t) - \bar{X}(t), \mathbb{H}_x(t, \bar{X}(t), \bar{u}(t), \bar{p}(t), \bar{q}(t)) \rangle \right] dt. \tag{6.14}
\end{aligned}$$

By the convexity of the Hamiltonian \mathbb{H} , we have

$$\begin{aligned}
& \mathbb{H}(t, X(t), u(t), \bar{p}(t), \bar{q}(t)) - \mathbb{H}(t, \bar{X}(t), \bar{u}(t), \bar{p}(t), \bar{q}(t)) \\
&\geq \langle X(t) - \bar{X}(t), \mathbb{H}_x(t, \bar{X}(t), \bar{u}(t), \bar{p}(t), \bar{q}(t)) \rangle \\
&\quad + \langle u(t) - \bar{u}(t), \mathbb{H}_u(t, \bar{X}(t), \bar{u}(t), \bar{p}(t), \bar{q}(t)) \rangle. \tag{6.15}
\end{aligned}$$

Combining (6.14)-(6.15) together with (6.10), it follows that

$$\begin{aligned}
& \mathbf{J}(u(\cdot)) - \mathbf{J}(\bar{u}(\cdot)) \\
&\geq \mathbb{E} \left[\int_0^T \langle u(t) - \bar{u}(t), \mathbb{H}_u(t, \bar{X}(t), \bar{u}(t), \bar{p}(t), \bar{q}(t)) \rangle dt \right] \\
&\geq 0,
\end{aligned}$$

which means that $\bar{u}(\cdot)$ is an optimal control for Problem (S). ■

Remark 6.2. It should be noted that the assumptions in Theorem 6.1 can be made weaker, but we do not focus on this here.

6.1 Example

In this subsection, we consider a simple example to illustrate our results. For $n = d = 1$, consider the following controlled system:

$$\begin{cases} dX(t) = u(t) dt + dW(t), & t \in [0, 1], \\ X(0) = 0, \end{cases} \tag{6.16}$$

with the control domain being $U = [-2, 2]$ and the cost functional being

$$\mathbf{J}(u(\cdot)) := \frac{1}{2} \mathbb{E} \left[\int_0^1 (u(t) + 1)^2 dt - e^{-\mathbb{E}[X(T)]^2} \right]. \tag{6.17}$$

We want to address the following stochastic control problem.

Problem (E). Minimize (6.17) over $\mathcal{U}[0, 1]$.

Suppose that $(\bar{u}(\cdot), \bar{X}(\cdot))$ is an optimal pair (which we are going to identify), then according to Theorem 4.3 the optimal 6-tuple $(\bar{X}(\cdot), \bar{u}(\cdot), \bar{g}(\cdot, \cdot), \bar{\eta}(\cdot, \cdot), \bar{\theta}(\cdot, \cdot), \bar{\psi}(\cdot, \cdot))$ satisfies the system of FBSDEs

$$\begin{cases} d\bar{X}(t) = \bar{u}(t) dt + dW(t), & t \in [0, T], \\ d\bar{\theta}(t, x) = - \left\{ \bar{u}(t) \bar{\theta}_x(t, x) + \frac{1}{2} \bar{\theta}_{xx}(t, x) + \bar{\psi}_x(t, x) + \frac{1}{2} (\bar{u}(t) + 1)^2 \right\} dt \\ \quad + \bar{\psi}(t, x)^\top dW(t), & (t, x) \in [0, T] \times \mathbb{R}^n, \\ d\bar{g}(t, x) = - \left\{ \bar{u}(t) \bar{g}_x(t, x) + \frac{1}{2} \bar{g}_{xx}(t, x) + \bar{\eta}_x(t, x) \right\} dt \\ \quad + \bar{\eta}(t, x)^\top dW(t), & (t, x) \in [0, T] \times \mathbb{R}^n, \\ \bar{X}(0) = 0, \bar{\theta}(T, x) = 0, \bar{g}(T, x) = x, & \text{for } x \in \mathbb{R}, \end{cases} \quad (6.18)$$

with the minimum condition: For a.e. $t \in [0, 1]$,

$$\bar{u}(t) \in \arg \min_{u \in [-2, 2]} \left\{ u \left(\bar{\theta}_x(t, \bar{X}(t)) + \mathbb{E}[\bar{X}(1)] e^{-\mathbb{E}[\bar{X}(1)]^2} \bar{g}_x(t, \bar{X}(t)) \right) + \frac{1}{2} (u + 1)^2 \right\}. \quad (6.19)$$

Moreover, the objective value of $\bar{u}(\cdot)$ is given by

$$\mathbf{J}(\bar{u}(\cdot)) = \bar{\theta}(0, 0) - \frac{1}{2} e^{-g(0, 0)^2}. \quad (6.20)$$

To solve the above system, we consider the following ansatz: For all $(t, x) \in [0, T] \times \mathbb{R}$,

$$\bar{\theta}(t, x) = M(t) \quad (6.21)$$

and

$$\bar{g}(t, x) = N(t)x + L(t), \quad (6.22)$$

where $(M(\cdot), V_M(\cdot))$, $(N(\cdot), V_N(\cdot))$ and $(L(\cdot), V_L(\cdot))$ are the pairs of adapted processes which are assumed to satisfy the BSDEs,

$$\begin{cases} dM(t) = -U_M(t) dt + V_M(t) dW(t), \\ M(T) = 0, \\ dN(t) = -U_N(t) dt + V_N(t) dW(t), \\ N(T) = 1, \end{cases}$$

and

$$\begin{cases} dL(t) = -U_L(t) dt + V_L(t) dW(t), \\ L(T) = 0. \end{cases}$$

In this case, the partial derivatives of $\bar{\theta}(t, x)$ and $\bar{g}(t, x)$ are

$$\bar{\theta}_x(t, x) \equiv 0, \bar{\theta}_{xx}(t, x) \equiv 0, \bar{g}_x(t, x) \equiv N(t) \text{ and } \bar{g}_{xx}(t, x) \equiv 0. \quad (6.23)$$

We would like to determine the equation that $(M(\cdot), V_M(\cdot))$, $(N(\cdot), V_N(\cdot))$ and $(L(\cdot), V_L(\cdot))$ should satisfy. To this end, we differentiate (6.21)-(6.22) and compare them with (6.18), we obtain that

$$\begin{aligned} & -U_M(t) dt + V_M(t) dW(t) \\ & = - \left\{ \bar{\psi}_x(t, x) + \frac{1}{2} (\bar{u}(t) + 1)^2 \right\} dt + \bar{\psi}(t, x) dW(t). \end{aligned} \quad (6.24)$$

and

$$\begin{aligned} & - \{U_N(t)x + U_L(t)\} dt + \{V_N(t)x + V_L(t)\} dW(t), \\ & = - \{N(t)\bar{u}(t) + \bar{\eta}_x(t, x)\} dt + \bar{\eta}(t, x) dW(t). \end{aligned} \quad (6.25)$$

Thus for all $(t, x) \in [0, T] \times \mathbb{R}$,

$$\bar{\psi}(t, x) = V_M(t) \quad (6.26)$$

and

$$\bar{\eta}(t, x) = \{V_N(t)x + V_L(t)\}. \quad (6.27)$$

Consequently,

$$\bar{\psi}_x(t, x) \equiv 0 \quad (6.28)$$

and

$$\bar{\eta}_x(t, x) \equiv V_N(t). \quad (6.29)$$

Next, comparing the dt terms in (6.24)-(6.25) and using the equalities (6.28)- (6.29), we obtain that

$$U_M(t) = \frac{1}{2}(\bar{u}(t) + 1)^2$$

and

$$U_N(t)x + U_L(t) = \{N(t)\bar{u}(t) + V_N(t)\}$$

which leads to the following BSDEs

$$\begin{cases} dM(t) = -\frac{1}{2}(\bar{u}(t) + 1)^2 dt + V_M(t) dW(t), \\ M(T) = 0, \end{cases} \quad (6.30)$$

$$\begin{cases} dN(t) = V_N(t) dW(t), \\ N(T) = 1, \end{cases} \quad (6.31)$$

and

$$\begin{cases} dL(t) = -\{N(t)\bar{u}(t) + V_N(t)\} dt + V_L(t) dW(t), \\ L(T) = 0. \end{cases} \quad (6.32)$$

Equation (6.23) can be easily solved, whose solution is given by

$$(N(t), V_N(t)) \equiv (1, 0), \forall t \in [0, 1].$$

Moreover by taking (6.23) into (6.19) we obtain that,

$$\bar{u}(t) \in \arg \min_{u \in [-2, 2]} \left\{ u \mathbb{E}[\bar{X}(1)] e^{-\mathbb{E}[\bar{X}(1)]^2} + \frac{1}{2}(u + 1)^2 \right\}$$

which suggests that

$$\bar{u}(t) \equiv -\mathbb{E}[\bar{X}(1)] e^{-\mathbb{E}[\bar{X}(1)]^2} - 1 \in]-2, +2[, \forall t \in [0, 1]. \quad (6.33)$$

In order to see that $\bar{u}(t) \equiv -\mathbb{E}[\bar{X}(T)] e^{-\mathbb{E}[\bar{X}(T)]^2} - 1$ is indeed optimal, we note that

$$\mathbb{H}(t, x, u, p, q) = up + q + \frac{1}{2}(u + 1)^2$$

is convex with respect to (x, u) . Moreover $h(x) = 0$ and $G(\bar{x}) = -e^{-\bar{x}^2}$ are convex. So the optimality follows from the sufficient condition (Theorem 6.1).

Taking the optimal control (6.33) into the state equation (6.16), we obtain that

$$\begin{cases} d\bar{X}(t) = -\left\{ \mathbb{E}[\bar{X}(1)] e^{-\mathbb{E}[\bar{X}(1)]^2} + 1 \right\} dt + dW(t), \quad t \in [0, 1], \\ \bar{X}(0) = 0, \end{cases} \quad (6.34)$$

Accordingly we have,

$$\begin{cases} \mathbb{E}[\bar{X}(t)] = -\left\{ \mathbb{E}[\bar{X}(T)] e^{-\mathbb{E}[\bar{X}(T)]^2} + 1 \right\} t, \quad t \in [0, 1], \\ \mathbb{E}[\bar{X}(0)] = 0, \end{cases}$$

which leads to

$$\mathbb{E}[\bar{X}(t)] = -\left\{ \mathbb{E}[\bar{X}(T)] e^{-\mathbb{E}[\bar{X}(T)]^2} + 1 \right\} t, \quad \forall t \in [0, 1],$$

Setting $t = 1$ in the above, we obtain that $\mathbb{E}[\bar{X}(1)] = \mathbf{X}_*$ coincides with the unique solution of the following equation,

$$\mathbf{X}_* + \mathbf{X}_* e^{-\mathbf{X}_*^2} + 1 = 0.$$

Now invoking (6.33) into (6.30)-(6.32), the BSDEs satisfied by $(M(\cdot), V_M(\cdot))$ and $(L(\cdot), V_L(\cdot))$ reduce to

$$\begin{cases} dM(t) = -\frac{1}{2}\mathbf{X}_*^2 e^{-2\mathbf{X}_*^2} dt + V_M(t) dW(t), \\ M(1) = 0; \end{cases}$$

and

$$\begin{cases} dL(t) = \left\{ \mathbf{X}_* e^{-\mathbf{X}_*^2} + 1 \right\} dt + V_L(t) dW(t), \\ L(1) = 0. \end{cases}$$

These equations can be easily solved, whose solutions are

$$(M(t), V_M(t)) \equiv \left(\frac{1}{2}\mathbf{X}_*^2 e^{-2\mathbf{X}_*^2} (1-t), 0 \right), \forall t \in [0, 1]. \quad (6.35)$$

and

$$(L(t), V_L(t)) \equiv \left(-\left(\mathbf{X}_* e^{-\mathbf{X}_*^2} + 1 \right) (1-t), 0 \right), \forall t \in [0, 1]. \quad (6.36)$$

Taking (6.35)-(6.36) into (6.21)-(6.22) we obtain that the objective value of $\bar{u}(\cdot)$ is given by

$$\mathbf{J}(\bar{u}(\cdot)) = \frac{1}{2}\mathbf{X}_*^2 e^{-2\mathbf{X}_*^2} - \frac{1}{2}e^{-\left(\mathbf{X}_* e^{-\mathbf{X}_*^2} + 1\right)^2}.$$

Remark 6.3. Summarizing the preceding analysis, we have the following results:

(i) The optimal control of Problem (E) is given by

$$\bar{u}(t) \equiv -\mathbf{X}_* e^{-\mathbf{X}_*^2} - 1$$

with the corresponding state

$$\bar{X}(t) \equiv W(t) - t \left(\mathbf{X}_* e^{-\mathbf{X}_*^2} + 1 \right),$$

where \mathbf{X}_* satisfies the following equation

$$\mathbf{X}_* + \mathbf{X}_* e^{-\mathbf{X}_*^2} + 1 = 0.$$

(ii) The corresponding solution of the BSPDEs are given by,

$$\begin{aligned} \bar{\theta}(t, x) &\equiv \frac{1}{2}\mathbf{X}_*^2 e^{-2\mathbf{X}_*^2} (1-t), \\ \bar{\psi}(t, x) &\equiv 0. \end{aligned}$$

and

$$\begin{aligned} \bar{g}(t, x) &\equiv x - \left(\mathbf{X}_* e^{-\mathbf{X}_*^2} + 1 \right) (1-t), \\ \bar{\eta}(t, x) &\equiv 0. \end{aligned}$$

(iii) The objective value of $\bar{u}(\cdot)$ is given by

$$\begin{aligned} \mathbf{J}(\bar{u}(\cdot)) &= \inf_{u(\cdot) \in \mathcal{U}[0,1]} \mathbf{J}(u(\cdot)) \\ &= \frac{1}{2}\mathbf{X}_*^2 e^{-2\mathbf{X}_*^2} - \frac{1}{2}e^{-\left(\mathbf{X}_* e^{-\mathbf{X}_*^2} + 1\right)^2}. \end{aligned}$$

Again, we emphasize that the new version of the SMP permits us to derive the objective value $\mathbf{J}(\bar{u}(\cdot))$, which is different from the traditional SMP approach.

Remark 6.4. Approximately, we have

$$\begin{aligned} \mathbb{E}[\bar{X}(1)] &= \mathbf{X}_* \simeq -0.58462; \\ \bar{u}(t) &\simeq 0.58462, \forall t \in [0, 1]; \\ \mathbf{J}(\bar{u}(\cdot)) &\simeq -0.29. \end{aligned}$$

7 A Class of forward-backward stochastic partial differential equations

In this section, motivated by the system of FBSDEs (4.8), we are going to study the solvability of the following class of forward-backward stochastic partial differential equations:

$$\begin{cases} dX(t) = \bar{b}(t, X(t), p_x(t, X(t))) dt + \bar{\sigma}(t, X(t)) dW(t), \text{ for } t \in [0, T], \\ dp(t, x) = - \left\{ \langle p_x(t, x), \bar{b}(t, x, p_x(t, X(t))) \rangle + \frac{1}{2} \mathbf{tr} \left[\bar{\sigma}(t, x) \bar{\sigma}(t, x)^\top p_{xx}(t, X(t)) \right] \right. \\ \quad \left. + \mathbf{tr} [q_x(t, x) \bar{\sigma}(t, x)] + \bar{f}(t, x, p_x(t, X(t))) \right\} dt \\ \quad + \bar{q}(t, x)^\top dW(t), \text{ for } (t, x) \in [0, T] \times \mathbb{R}^n, \\ X(0) = x_0, p(T, x) = \bar{F}(x), \text{ for } x \in \mathbb{R}^n. \end{cases} \quad (7.1)$$

In the above, $X(\cdot)$ is the unknown process, $(p(\cdot, \cdot), q(\cdot, \cdot))$ are the unknown random fields, and they are required to be $(\mathcal{F}_t)_{t \in [0, T]}$ -adapted; $\bar{b}: [0, T] \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$, $\bar{\sigma}: [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}^{n \times d}$, $\bar{f}: [0, T] \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ and $\bar{F}: \mathbb{R}^n \rightarrow \mathbb{R}$ are given deterministic measurable functions. The main feature of the above system is that $X(\cdot)$ satisfies a forward SDE and the pair of random fields $(p(\cdot, \cdot), q(\cdot, \cdot))$ satisfies a nonlinear BSPDE.

We introduce the following definition.

Definition 7.1. A triple $(X(\cdot), p(\cdot, \cdot), q(\cdot, \cdot))$ is called a *solution* of (7.1) if the following holds

$$\begin{cases} X(\cdot) \in \mathcal{C}_{\mathcal{F}}^2(0, T; \mathbb{R}^n) \\ p(\cdot, \cdot) \in \mathcal{C}_{\mathcal{F}}^2(0, T; C^2(\bar{B}_R; \mathbb{R})), \quad \forall R > 0, \\ q(\cdot, \cdot) \in \mathcal{L}_{\mathcal{F}}^2(0, T; C^1(\bar{B}_R; \mathbb{R}^d)), \end{cases}$$

such that the following holds, for all $(t, x) \in [0, T] \times \mathbb{R}^n$, almost surely:

$$\begin{cases} X(t) = x_0 + \int_0^t \bar{b}(\tau, X(\tau), p_x(\tau, X(\tau))) d\tau + \int_0^t \bar{\sigma}(\tau, X(\tau)) dW(\tau), \\ p(t, x) = \bar{F}(x) + \int_t^T \left\{ \langle p_x(\tau, x), \bar{b}(\tau, x, p_x(\tau, X(\tau))) \rangle \right. \\ \quad \left. + \frac{1}{2} \mathbf{tr} \left[\bar{\sigma}(\tau, x) \bar{\sigma}(\tau, x)^\top p_{xx}(\tau, x) \right] \right. \\ \quad \left. + \mathbf{tr} [q_x(\tau, x) \bar{\sigma}(\tau, x)] + \bar{f}(\tau, x, p_x(\tau, X(\tau))) \right\} d\tau \\ \quad - \int_t^T \bar{q}(\tau, x)^\top dW(\tau). \end{cases}$$

7.1 A version of three-step scheme, a heuristic derivation

Now, essentially inspired by the idea of the classical four-step scheme introduced in Ma et al. [25] (see also e.g. Ma and Yong [26] or Yong and Zhou [38]), we are going to introduce a method for solving the FBSPDE (7.1) over any time duration $[0, T]$.

As in [[38], Section 7.5.2] let us give a heuristic derivation first. Suppose that $(X(\cdot), p(\cdot, \cdot), q(\cdot, \cdot))$ is an adapted solution to (7.1). We assume that $p(t, x)$ and $X(t)$ are related by

$$p(t, x) = \theta(t, x, X(t)), \text{ for any } (t, x) \in [0, T] \times \mathbb{R}^n, \quad (7.2)$$

where $\theta(t, x, y)$ is some function to be determined such that for any $(x, y) \in \mathbb{R}^n \times \mathbb{R}^n$,

$$\theta(T, x, y) = F(x).$$

We suppose that $\theta(t, x, y)$ is C^1 in t and C^2 in (x, y) . Accordingly, we have for any $(t, x) \in [0, T] \times \mathbb{R}^n$,

$$p_x(t, x) = \theta_x(t, x, X(t)),$$

and

$$p_{xx}(t, x) = \theta_{xx}(t, x, X(t)).$$

Then by Itô's formula, we differentiate (7.2) and compare it with (7.1) to obtain that

$$\begin{aligned}
dp(t, x) &= d\theta(t, x, X(t)) \\
&= \left\{ \theta_t(t, x, X(t)) + \langle \theta_y(t, x, X(t)), \bar{b}(t, X(t), \theta_x(t, X(t), X(t))) \rangle \right. \\
&\quad \left. + \frac{1}{2} \text{tr} \left[\bar{\sigma}(t, X(t)) \bar{\sigma}(t, X(t))^\top \theta_{yy}(t, x, X(t)) \right] \right\} dt \\
&\quad + \langle \theta_y(t, x, X(t)), \bar{\sigma}(t, X(t)) dW(t) \rangle \\
&= - \left\{ \langle \theta_x(t, x, X(t)), \bar{b}(t, x, \theta_x(t, X(t), X(t))) \rangle \right. \\
&\quad \left. + \frac{1}{2} \text{tr} \left[\bar{\sigma}(t, x) \bar{\sigma}(t, x)^\top \theta_{xx}(t, x, X(t)) \right] \right. \\
&\quad \left. + \text{tr} [q_x(t, x) \bar{\sigma}(t, x)] + \bar{f}(t, x, \theta_x(t, X(t), X(t))) \right\} dt \\
&\quad + q(t, x)^\top dW(t), \tag{7.3}
\end{aligned}$$

Accordingly, we have

$$q(t, x) \equiv \bar{\sigma}(t, X(t))^\top \theta_y(t, x, X(t)).$$

Thus

$$q_x(t, x) \equiv \bar{\sigma}(t, X(t))^\top \theta_{yx}(t, x, X(t)). \tag{7.4}$$

Now compare the dt terms in (7.3) and using (7.4), we obtain that

$$\begin{aligned}
0 &= \theta_t(t, x, X(t)) + \langle \theta_y(t, x, X(t)), \bar{b}(t, X(t), \theta_x(t, X(t), X(t))) \rangle \\
&\quad + \frac{1}{2} \text{tr} \left[\bar{\sigma}(t, X(t)) \bar{\sigma}(t, X(t))^\top \theta_{yy}(t, x, X(t)) \right] + \langle \theta_y(t, x, X(t)), \bar{\sigma}(t, X(t)) dW(t) \rangle \\
&\quad + \left\{ \langle \theta_x(t, x, X(t)), \bar{b}(t, x, \theta_x(t, X(t), X(t))) \rangle + \frac{1}{2} \text{tr} \left[\bar{\sigma}(t, x) \bar{\sigma}(t, x)^\top \theta_{xx}(t, x, X(t)) \right] \right. \\
&\quad \left. + \text{tr} \left[\bar{\sigma}(t, X(t))^\top \theta_{yx}(t, x, X(t)) \bar{\sigma}(t, x) \right] + \bar{f}(t, x, \theta_x(t, X(t), X(t))) \right\}.
\end{aligned}$$

The above argument suggests that we design the following three-step scheme to solve the FBSDE (7.1).

A Three-step scheme:

Step 1. Solve the following nonlinear parabolic system for $\theta(t, x, y)$:

$$\begin{cases} 0 = \theta_t(t, x, y) + \langle \theta_y(t, x, y), \bar{b}(t, y, \theta_x(t, y, y)) \rangle + \frac{1}{2} \text{tr} \left[\bar{\sigma}(t, y) \bar{\sigma}(t, y)^\top \theta_{yy}(t, x, y) \right] \\ \quad + \langle \theta_x(t, x, y), \bar{b}(t, x, \theta_x(t, y, y)) \rangle + \frac{1}{2} \text{tr} \left[\bar{\sigma}(t, x) \bar{\sigma}(t, x)^\top \theta_{xx}(t, x, y) \right] \\ \quad + \text{tr} \left[\bar{\sigma}(t, y)^\top \theta_{yx}(t, x, y) \bar{\sigma}(t, x) \right] + \bar{f}(t, x, \theta_x(t, y, y)), \text{ for } (t, x, y) \in [0, T] \times \mathbb{R}^n \times \mathbb{R}^n, \\ \theta(T, x, y) = \bar{F}(x), \text{ for } (t, x, y) \in \mathbb{R}^n \times \mathbb{R}^n. \end{cases} \tag{7.5}$$

Step 2. Use θ obtained in Steps 1 to solve the following forward SDE:

$$\begin{cases} dX(t) = \tilde{b}(t, X(t)) dt + \bar{\sigma}(t, X(t)) dW(t), \text{ for } t \in [0, T], \\ X(0) = x_0, \end{cases} \tag{7.6}$$

where

$$\tilde{b}(t, y) := \bar{b}(t, y, \theta_x(t, y, y)) \tag{7.7}$$

Step 3. Set

$$\begin{cases} p(t, x) := \theta(t, x, X(t)), \\ q(t, x) := \bar{\sigma}(t, X(t))^\top \theta_y(t, x, X(t)), \text{ for any } (t, x) \in [0, T] \times \mathbb{R}^n. \end{cases} \tag{7.8}$$

Should this scheme be realizable, $(X(\cdot), p(\cdot, \cdot), q(\cdot, \cdot))$ would give an adapted solution of (7.1). We have the following result.

Theorem 7.2. Assume that (7.5) admits a unique classical solution $\theta(t, x, y)$ with bounded $\theta_x(t, x, y)$, $\theta_y(t, x, y)$, $\theta_{xx}(t, x, y)$, $\theta_{yy}(t, x, y)$ and $\theta_{xy}(t, x, y)$. Assume further that the functions $\bar{b}(t, y, p)$ and $\bar{\sigma}(t, y)$ are uniformly Lipschitz continuous in (y, p) with $\bar{b}(t, 0, 0)$ and $\bar{\sigma}(t, 0)$ being bounded. Then the triple $(X(\cdot), p(\cdot, \cdot), q(\cdot, \cdot))$ determined by (7.6) and (7.8) is an adapted solution to (7.1).

Proof. Under our conditions both $\bar{\sigma}(t, y)$ and $\bar{b}(t, y)$ defined by (7.7) are uniformly Lipschitz continuous in y . Therefore, for any $x_0 \in \mathbb{R}^n$, (7.6) has a unique strong solution. Then, by defining $p(t, x)$ and $q(t, x)$ via (7.8) and applying Itô's formula, we can easily check that (7.1) is satisfied. Hence, $(X(\cdot), p(\cdot, \cdot), q(\cdot, \cdot))$ is a solution of (7.1). \blacksquare

We now turn our attention on the Cauchy problem (7.5). First, let us consider the following notations¹

$$\begin{aligned} \mathbf{X} &:= (x, y) \in \mathbb{R}^{2n}, \\ \varphi(\mathbf{X}) &= \varphi(x, y) := (y, y) \in \mathbb{R}^{2n}, \\ \theta_{\mathbf{X}}(t, \mathbf{X}) &:= \begin{pmatrix} \theta_x(t, x, y) \\ \theta_y(t, x, y) \end{pmatrix}, \\ \theta_{\mathbf{X}\mathbf{X}}(t, \mathbf{X}) &:= \begin{pmatrix} \theta_{xx}(t, x, y) & \theta_{xy}(t, x, y) \\ \theta_{yx}(t, x, y) & \theta_{yy}(t, x, y) \end{pmatrix}, \\ \mathbf{b}(t, \mathbf{X}, \theta_{\mathbf{X}}(t, \varphi(\mathbf{X}))) &:= \begin{pmatrix} \bar{b}(t, x, \theta_x(t, y, y)) \\ \bar{b}(t, y, \theta_x(t, y, y)) \end{pmatrix}, \\ \mathbf{f}(t, \mathbf{X}, \theta_{\mathbf{X}}(t, \varphi(\mathbf{X}))) &:= f(t, x, \theta_x(t, y, y)), \\ \mathbf{F}(\mathbf{X}) &:= \bar{F}(x), \\ \mathbf{a}(t, \mathbf{X}) &:= \frac{1}{2} \begin{pmatrix} \bar{\sigma}(t, x) \bar{\sigma}(t, x)^\top & \bar{\sigma}(t, y) \bar{\sigma}(t, x)^\top \\ \bar{\sigma}(t, x) \bar{\sigma}(t, y)^\top & \bar{\sigma}(t, y) \bar{\sigma}(t, y)^\top \end{pmatrix}. \end{aligned}$$

Then it is not difficult to see that $\theta(t, x, y)$ is a classical solution to (7.5), if and only if, $\theta(t, \mathbf{X})$ is a classical solution of the following nonlinear parabolic system

$$\begin{cases} 0 = \theta_t(t, \mathbf{X}) + \langle \theta_{\mathbf{X}}(t, \mathbf{X}), \mathbf{b}(t, \mathbf{X}, \theta_{\mathbf{X}}(t, \varphi(\mathbf{X}))) \rangle \\ \quad + \text{tr}[\mathbf{a}(t, \mathbf{X}) \theta_{\mathbf{X}\mathbf{X}}(t, \mathbf{X})] \\ \quad + \mathbf{f}(t, \mathbf{X}, \theta_{\mathbf{X}}(t, \varphi(\mathbf{X}))), \text{ for } (t, \mathbf{X}) \in [0, T] \times \mathbb{R}^{2n}, \\ \theta(T, \mathbf{X}) = \mathbf{F}(\mathbf{X}), \text{ for } \mathbf{X} \in \mathbb{R}^{2n}. \end{cases} \quad (7.9)$$

7.2 Well-posedness of the parabolic PDE

In this subsections, we discuss the well-posedness for the PDE (7.9) by adopting a fixed point method. Let us make some preparations.

Let $C^\alpha(\mathbb{R}^{2n})$ be the space of continuous functions $\varphi(\cdot)$ such that

$$\|\varphi\|_\alpha = \|\varphi\|_0 + [\varphi]_\alpha < \infty,$$

where

$$\|\varphi\|_0 = \sup_{\mathbf{X} \in \mathbb{R}^{2n}} |\varphi(\mathbf{X})|, \quad [\varphi]_\alpha = \sup_{\mathbf{X}, \mathbf{Y} \in \mathbb{R}^{2n}} \frac{|\varphi(\mathbf{X}) - \varphi(\mathbf{Y})|}{\|\mathbf{X} - \mathbf{Y}\|^\alpha} < \infty$$

Further let $C^{1+\alpha}(\mathbb{R}^{2n})$ and $C^{1+\alpha}(\mathbb{R}^{2n})$ be the space of continuous functions $\varphi(\cdot)$ such that

$$\|\varphi\|_{1+\alpha} = \|\varphi\|_0 + \|\varphi_x\|_0 + [\varphi_x]_\alpha < \infty$$

and

$$\|\varphi\|_{1+\alpha} = \|\varphi\|_0 + \|\varphi_x\|_0 + \|\varphi_{xx}\|_0 + [\varphi_{xx}]_\alpha < \infty,$$

respectively. Next, let $\mathbb{B}([0, T]; C^\alpha(\mathbb{R}^{2n}))$, the set of all measurable functions $f : [0, T] \times \mathbb{R}^{2n} \rightarrow \mathbb{R}$ such that for each $t \in [0, T]$, $f(t, \cdot) \in C^\alpha(\mathbb{R}^{2n})$ and

$$\|f\|_{\mathbb{B}([0, T]; C^\alpha(\mathbb{R}^{2n}))} = \sup_{t \in [0, T]} \|f(t, \cdot)\|_\alpha < \infty.$$

¹Here we adopt the following convention. For any $x = (x_1, \dots, x_n)^\top \in \mathbb{R}^n$ and $y = (y_1, \dots, y_n)^\top \in \mathbb{R}^n$, we consider the pair (x, y) as an element of \mathbb{R}^{2n} and write:

$$(x, y) := (x_1, \dots, x_n, y_1, \dots, y_n) \in \mathbb{R}^{2n}.$$

Also, we let $\mathbb{C}([0, T]; C^\alpha(\mathbb{R}^{2n}))$ be the set of all continuous functions that are also in $\mathbb{B}([0, T]; C^\alpha(\mathbb{R}^{2n}))$. Similarly, we define $\mathbb{B}([0, T]; C^{k+\alpha}(\mathbb{R}^{2n}))$ and $\mathbb{C}([0, T]; C^{k+\alpha}(\mathbb{R}^{2n}))$, for $k = 1, 2$.

We now introduce the following assumptions.

(H) The maps \bar{b} , $\bar{\sigma}$, \bar{f} and \bar{h} are continuous and bounded. Moreover, there exists a constant $K > 0$ such that $\forall (t, x, p) \in [0, T] \times \mathbb{R}^n \times \mathbb{R}^n$,

$$K \geq |\bar{\sigma}_x(t, x)| + |\bar{b}_x(t, x, p)| + |\bar{f}_x(t, x, p)| \\ + |\bar{b}_p(t, x, p)| + |\bar{f}_p(t, x, p)| + |\bar{h}_x(t, x)|.$$

Further $\mathbf{a}(t, \mathbf{X})^{-1}$ the inverse of $\mathbf{a}(t, \mathbf{X})$ exists for all $(t, \mathbf{X}) \in [0, T] \times \mathbb{R}^{2n}$ and there exists constants $\lambda_0, \lambda_1 > 0$ such that $\forall \mathbf{X} \in \mathbb{R}^{2n}$

$$\lambda_1 \mathbf{I}_{2n} \leq \mathbf{a}(t, \mathbf{X})^{-1} \leq \lambda_2 \mathbf{I}_{2n},$$

where \mathbf{I}_{2n} denotes the $(2n \times 2n)$ identity matrix.

The following theorem grantee the existence and uniqueness of a classical solution to the PDEs (7.9); its proof follows an argument adapted from Proof of Theorem 5.2. in Yong [37].

Theorem 7.3. *Let Assumption (H) hold. Then (7.9) admits a unique solution $\theta(\cdot, \cdot)$.*

Proof. For any fixed $v(\cdot, \cdot) \in \mathbb{C}([0, T]; C^{1+\alpha}(\mathbb{R}^{2n}))$, we consider the following linear parabolic PDE

$$\begin{cases} 0 = \theta_t(t, \mathbf{X}) + [\mathcal{L}^0 \theta(t, \cdot)](\mathbf{X}) + \langle \theta_{\mathbf{X}}(t, \mathbf{X}), \mathbf{b}(t, \mathbf{X}, v_{\mathbf{X}}(t, \varphi(\mathbf{X}))) \rangle \\ \quad + \mathbf{f}(t, \mathbf{X}, v_{\mathbf{X}}(t, \varphi(\mathbf{X}))), \text{ for } (t, \mathbf{X}) \in [0, T] \times \mathbb{R}^{2n}, \\ \theta(T, \mathbf{X}) = \mathbf{F}(\mathbf{X}), \text{ for } \mathbf{X} \in \mathbb{R}^{2n}. \end{cases} \quad (7.10)$$

where $\mathcal{L}^0(\cdot)$ is the differentiable operator defined as follows: For any $\psi(\cdot) \in C^2(\mathbb{R}^{2n})$,

$$[\mathcal{L}^0 \psi(\cdot)](\mathbf{X}) = \text{tr}[\mathbf{a}(t, \mathbf{X}) \psi_{\mathbf{X}\mathbf{X}}(\mathbf{X})]$$

Applying Proposition 5.1 in Yong [37], we obtain that

$$\begin{aligned} \theta(t, \mathbf{X}) &= \int_{\mathbb{R}^{2n}} \Gamma^0(t, \mathbf{X}; T, \mathbf{Z}) \bar{\mathbf{F}}(\mathbf{Z}) d\mathbf{Z} \\ &\quad + \int_t^T \int_{\mathbb{R}^{2n}} \Gamma^0(t, \mathbf{X}; s, \mathbf{Z}) \langle \mathbf{b}(t, \mathbf{Z}, v_{\mathbf{X}}(t, \varphi(\mathbf{Z}))), \theta_{\mathbf{X}}(t, \mathbf{Z}) \rangle d\mathbf{Z} ds \\ &\quad + \int_t^T \int_{\mathbb{R}^{2n}} \Gamma^0(t, \mathbf{X}; s, \mathbf{Z}) \mathbf{f}(t, \mathbf{Z}, v_{\mathbf{X}}(t, \varphi(\mathbf{Z}))) d\mathbf{Z} ds, \end{aligned}$$

where $\Gamma^0(t, \mathbf{X}; s, \mathbf{Z})$ is the fundamental solution of $\mathcal{L}^0(\cdot)$, given explicitly by

$$\Gamma^0(t, \mathbf{X}; s, \mathbf{Z}) = \frac{1}{(4\pi(s-t))^n (\det[\mathbf{a}(t, \mathbf{Z})])^{\frac{1}{2}}} e^{\frac{\langle \mathbf{a}(t, \mathbf{Z})^{-1}(\mathbf{X}-\mathbf{Z}), (\mathbf{X}-\mathbf{Z}) \rangle}{4(s-t)}}.$$

On the other hand, by some computations one has (see e.g. [9], Page 24.)

$$\begin{cases} |\Gamma^0(t, \mathbf{X}; s, \mathbf{Z})| \leq \frac{K}{(s-t)^n} e^{-\lambda \frac{|\mathbf{X}-\mathbf{Z}|^2}{4(s-t)}}, \\ |\Gamma_{\mathbf{X}}^0(t, \mathbf{X}; s, \mathbf{Z})| \leq \frac{K}{(s-t)^{\frac{2n+1}{2}}} e^{-\lambda \frac{|\mathbf{X}-\mathbf{Z}|^2}{4(s-t)}}. \end{cases} \quad \lambda < \lambda_0. \quad (7.11)$$

Moreover, arguing as in [37] we get,

$$\Gamma_{\mathbf{Z}}^0(t, \mathbf{X}; s, \mathbf{Z}) = -\Gamma_{\mathbf{X}}^0(t, \mathbf{X}; s, \mathbf{Z}) - \Gamma^0(t, \mathbf{X}; s, \mathbf{Z}) \rho(t, \mathbf{X}; s, \mathbf{Z}), \quad (7.12)$$

where

$$\rho(t, \mathbf{X}; s, \mathbf{Z}) := \frac{1}{2} (\det[\mathbf{a}(t, \mathbf{Z})])_{\mathbf{Z}} + \frac{\langle [a(t, \mathbf{Z})^{-1}]_{\mathbf{Z}}(\mathbf{X}-\mathbf{Z}), (\mathbf{X}-\mathbf{Z}) \rangle}{4(s-t)},$$

with

$$\langle [a(t, \mathbf{Z})^{-1}]_{\mathbf{Z}}(\mathbf{X}-\mathbf{Z}), (\mathbf{X}-\mathbf{Z}) \rangle := \begin{pmatrix} \langle [a(t, \mathbf{Z})^{-1}]_{z_1}(\mathbf{X}-\mathbf{Z}), (\mathbf{X}-\mathbf{Z}) \rangle \\ \langle [a(t, \mathbf{Z})^{-1}]_{z_2}(\mathbf{X}-\mathbf{Z}), (\mathbf{X}-\mathbf{Z}) \rangle \\ \dots \\ \langle [a(t, \mathbf{Z})^{-1}]_{z_{2n}}(\mathbf{X}-\mathbf{Z}), (\mathbf{X}-\mathbf{Z}) \rangle \end{pmatrix}.$$

Under Assumption **(H)**, we have

$$\rho(t, \mathbf{X}; s, \mathbf{Z}) \leq K \left(1 + \frac{|\mathbf{X} - \mathbf{Z}|^2}{s - t} \right). \quad (7.13)$$

Then using (7.12), we have

$$\begin{aligned} & \theta_{\mathbf{X}}(t, \mathbf{X}) \\ &= \int_{\mathbb{R}^{2n}} \Gamma_{\mathbf{X}}^0(t, \mathbf{X}; T, \mathbf{Z}) \mathbf{F}(\mathbf{Z}) d\mathbf{Z} \\ &+ \int_t^T \int_{\mathbb{R}^{2n}} \Gamma_{\mathbf{X}}^0(t, \mathbf{X}; s, \mathbf{Z}) \langle \mathbf{b}(s, \mathbf{Z}, v_{\mathbf{X}}(s, \varphi(\mathbf{Z}))), \theta_{\mathbf{X}}(s, \mathbf{Z}) \rangle d\mathbf{Z} ds \\ &+ \int_t^T \int_{\mathbb{R}^{2n}} \Gamma_{\mathbf{X}}^0(t, \mathbf{X}; s, \mathbf{Z}) \mathbf{f}(s, \mathbf{Z}, v_{\mathbf{X}}(s, \varphi(\mathbf{Z}))) d\mathbf{Z} ds \\ &= - \int_{\mathbb{R}^{2n}} \Gamma_{\mathbf{Z}}^0(t, \mathbf{X}; s, \mathbf{Z}) \mathbf{F}(\mathbf{Z}) d\mathbf{Z} - \int_{\mathbb{R}^{2n}} \Gamma^0(t, \mathbf{X}; s, \mathbf{Z}) \rho(t, \mathbf{X}; s, \mathbf{Z}) \mathbf{F}(\mathbf{Z}) d\mathbf{Z} \\ &+ \int_t^T \int_{\mathbb{R}^{2n}} \Gamma_{\mathbf{X}}^0(t, \mathbf{X}; s, \mathbf{Z}) \langle \mathbf{b}(s, \mathbf{Z}, v_{\mathbf{X}}(s, \varphi(\mathbf{Z}))), \theta_{\mathbf{X}}(s, \mathbf{Z}) \rangle d\mathbf{Z} ds \\ &+ \int_t^T \int_{\mathbb{R}^{2n}} \Gamma_{\mathbf{X}}^0(t, \mathbf{X}; s, \mathbf{Z}) \mathbf{f}(s, \mathbf{Z}, v_{\mathbf{X}}(s, \varphi(\mathbf{Z}))) d\mathbf{Z} ds \\ &= \int_{\mathbb{R}^{2n}} \Gamma^0(t, \mathbf{X}; s, \mathbf{Z}) \mathbf{F}_{\mathbf{Z}}(\mathbf{Z}) d\mathbf{Z} - \int_{\mathbb{R}^{2n}} \Gamma^0(t, \mathbf{X}; s, \mathbf{Z}) \rho(t, \mathbf{X}; s, \mathbf{Z}) \mathbf{F}(\mathbf{Z}) d\mathbf{Z} \\ &+ \int_t^T \int_{\mathbb{R}^{2n}} \Gamma_{\mathbf{X}}^0(t, \mathbf{X}; s, \mathbf{Z}) \langle \mathbf{b}(s, \mathbf{Z}, v_{\mathbf{X}}(s, \varphi(\mathbf{Z}))), \theta_{\mathbf{X}}(s, \mathbf{Z}) \rangle d\mathbf{Z} ds \\ &+ \int_t^T \int_{\mathbb{R}^{2n}} \Gamma_{\mathbf{X}}^0(t, \mathbf{X}; s, \mathbf{Z}) \mathbf{f}(s, \mathbf{Z}, v_{\mathbf{X}}(s, \varphi(\mathbf{Z}))) d\mathbf{Z} ds. \end{aligned} \quad (7.14)$$

Therefore, combining (7.11)-(7.13) together with (7.14) and using the fact that \mathbf{f} and \mathbf{b} are uniformly bounded, we obtain that

$$\begin{aligned} |\theta_{\mathbf{X}}(t, \mathbf{X})| &\leq \int_{\mathbb{R}^{2n}} \frac{K}{(T-t)^n} e^{-\lambda \frac{|\mathbf{X}-\mathbf{Z}|^2}{4(T-t)}} \left[|\mathbf{F}_{\mathbf{Z}}(\mathbf{Z})| + \left(1 + \frac{|\mathbf{X}-\mathbf{Z}|^2}{T-t} \right) |\mathbf{F}(\mathbf{Z})| \right] d\mathbf{Z} \\ &+ \int_t^T \int_{\mathbb{R}^{2n}} \frac{K}{(s-t)^{\frac{2n+1}{2}}} e^{-\lambda \frac{|\mathbf{X}-\mathbf{Z}|^2}{4(s-t)}} [|\theta_{\mathbf{X}}(t, \mathbf{Z})| + 1] d\mathbf{Z} ds \\ &\leq K \left(\|F(\cdot)\|_{C^1(\mathbb{R}^n)} + 1 \right) + \int_t^T \int_{\mathbb{R}^{2n}} \frac{K}{(s-t)^{\frac{2n+1}{2}}} e^{-\lambda \frac{|\mathbf{X}-\mathbf{Z}|^2}{4(s-t)}} |\theta_{\mathbf{X}}(s, \mathbf{Z})| d\mathbf{Z} ds \end{aligned} \quad (7.15)$$

Thus we also have

$$|\theta_{\mathbf{X}}(s, \mathbf{Z})| \leq K \left(\|F(\cdot)\|_{C^1(\mathbb{R}^n)} + 1 \right) + \int_s^T \int_{\mathbb{R}^{2n}} \frac{K}{(r-s)^{\frac{2n+1}{2}}} e^{-\lambda \frac{|\mathbf{Z}-\mathbf{Y}|^2}{4(r-s)}} |\theta_{\mathbf{X}}(r, \mathbf{Z})| d\mathbf{Z} dr.$$

Accordingly, by using Lemma 3 in ([9], Page 24), we get

$$\begin{aligned} & \int_t^T \int_{\mathbb{R}^{2n}} \frac{K}{(s-t)^{\frac{2n+1}{2}}} e^{-\lambda \frac{|\mathbf{X}-\mathbf{Z}|^2}{4(s-t)}} |\theta_{\mathbf{X}}(s, \mathbf{Z})| d\mathbf{Z} ds \\ &\leq \int_t^T \int_{\mathbb{R}^{2n}} \frac{K}{(s-t)^{\frac{2n+1}{2}}} e^{-\lambda \frac{|\mathbf{X}-\mathbf{Z}|^2}{4(s-t)}} \left(\|F(\cdot)\|_{C^1(\mathbb{R}^n)} + 1 \right) d\mathbf{Z} ds \\ &+ \int_t^T \int_{\mathbb{R}^{2n}} \frac{K}{(s-t)^{\frac{2n+1}{2}}} e^{-\lambda \frac{|\mathbf{X}-\mathbf{Z}|^2}{4(s-t)}} \int_s^T \int_{\mathbb{R}^{2n}} \frac{K}{(r-s)^{\frac{2n+1}{2}}} e^{-\lambda \frac{|\mathbf{Z}-\mathbf{Y}|^2}{4(r-s)}} |\theta_{\mathbf{X}}(r, \mathbf{Y})| d\mathbf{Y} dr d\mathbf{Z} ds \\ &\leq K \left(\|F(\cdot)\|_{C^1(\mathbb{R}^n)} + 1 \right) \\ &+ \int_t^T \int_{\mathbb{R}^{2n}} \left(\int_t^r \int_{\mathbb{R}^{2n}} \frac{K}{(s-t)^{\frac{2n+1}{2}}} e^{-\lambda \frac{|\mathbf{X}-\mathbf{Z}|^2}{4(s-t)}} \frac{K}{(r-s)^{\frac{2n+1}{2}}} e^{-\lambda \frac{|\mathbf{Z}-\mathbf{Y}|^2}{4(r-s)}} d\mathbf{Z} ds \right) |\theta_{\mathbf{X}}(r, \mathbf{Y})| d\mathbf{Y} dr \\ &\leq K \left(\|F(\cdot)\|_{C^1(\mathbb{R}^n)} + 1 \right) + \int_t^T \int_{\mathbb{R}^{2n}} \frac{K}{(r-t)^{\frac{2n}{2}}} e^{-\lambda \frac{|\mathbf{X}-\mathbf{Y}|^2}{4(r-t)}} |\theta_{\mathbf{X}}(r, \mathbf{Y})| d\mathbf{Y} dr. \end{aligned}$$

Invoking this into (7.15), we get

$$|\theta_{\mathbf{X}}(t, \mathbf{X})| \leq K \left(\|F(\cdot)\|_{C^1(\mathbb{R}^n)} + 1 \right) + \int_t^T \int_{\mathbb{R}^{2n}} \frac{K}{(r-t)^{\frac{2n}{2}}} e^{-\lambda \frac{|\mathbf{X}-\mathbf{Y}|^2}{4(r-t)}} |\theta_{\mathbf{X}}(r, \mathbf{Y})| d\mathbf{Y} dr \quad (7.16)$$

We can repeat the above procedure $2n$ times and then use Gronwall's inequality to obtain

$$\sup_{\mathbf{X} \in \mathbb{R}^{2n}} |\theta_{\mathbf{X}}(t, \mathbf{X})| \leq K \left(\|F(\cdot)\|_{C^1(\mathbb{R}^n)} + 1 \right). \quad (7.17)$$

Now let $v_1(\cdot, \cdot), v_2(\cdot, \cdot) \in \mathbb{C}([0, T]; C^1(\mathbb{R}^{2n}))$ and we consider $\theta^1(\cdot, \cdot), \theta^2(\cdot, \cdot)$ their corresponding solutions of (7.10). Then

$$\begin{aligned} & \theta^1(t, \mathbf{X}) - \theta^2(t, \mathbf{X}) \\ &= \int_t^T \int_{\mathbb{R}^{2n}} \Gamma^0(t, \mathbf{X}; s, \mathbf{Z}) \left(\langle \mathbf{b}(s, \mathbf{Z}, v_{\mathbf{X}}^1(s, \varphi(\mathbf{Z}))), \theta_{\mathbf{X}}^1(s, \mathbf{Z}) - \theta_{\mathbf{X}}^2(s, \mathbf{Z}) \rangle \right. \\ &+ \langle \mathbf{b}(s, \mathbf{Z}, v_{\mathbf{X}}^1(s, \varphi(\mathbf{Z}))) - \mathbf{b}(s, \mathbf{Z}, v_{\mathbf{X}}^2(s, \varphi(\mathbf{Z}))), \theta_{\mathbf{X}}^2(s, \mathbf{Z}) \rangle \\ &+ \mathbf{f}(s, \mathbf{Z}, v_{\mathbf{X}}^1(s, \varphi(\mathbf{Z}))) - \mathbf{f}(s, \mathbf{Z}, v_{\mathbf{X}}^2(s, \varphi(\mathbf{Z}))) \rangle d\mathbf{Z} ds \end{aligned} \quad (7.18)$$

and

$$\begin{aligned} & \theta_{\mathbf{X}}^1(t, \mathbf{X}) - \theta_{\mathbf{X}}^2(t, \mathbf{X}) \\ &= \int_t^T \int_{\mathbb{R}^{2n}} \Gamma_{\mathbf{X}}^0(t, \mathbf{X}; s, \mathbf{Z}) \left(\langle \mathbf{b}(s, \mathbf{Z}, v_{\mathbf{X}}^1(s, \varphi(\mathbf{Z}))), \theta_{\mathbf{X}}^1(s, \mathbf{Z}) - \theta_{\mathbf{X}}^2(s, \mathbf{Z}) \rangle \right. \\ &+ \langle \mathbf{b}(s, \mathbf{Z}, v_{\mathbf{X}}^1(s, \varphi(\mathbf{Z}))) - \mathbf{b}(s, \mathbf{Z}, v_{\mathbf{X}}^2(s, \varphi(\mathbf{Z}))), \theta_{\mathbf{X}}^2(s, \mathbf{Z}) \rangle \\ &+ \mathbf{f}(s, \mathbf{Z}, v_{\mathbf{X}}^1(s, \varphi(\mathbf{Z}))) - \mathbf{f}(s, \mathbf{Z}, v_{\mathbf{X}}^2(s, \varphi(\mathbf{Z}))) \rangle d\mathbf{Z} ds. \end{aligned}$$

Accordingly, by Assumption (H) we have,

$$\begin{aligned} & |\theta_{\mathbf{X}}^1(t, \mathbf{X}) - \theta_{\mathbf{X}}^2(t, \mathbf{X})| \\ &\leq \int_t^T \int_{\mathbb{R}^{2n}} \frac{K}{(s-t)^{\frac{2n+1}{2}}} e^{-\lambda \frac{|\mathbf{X}-\mathbf{Z}|^2}{4(s-t)}} \left(|\theta_{\mathbf{X}}^1(s, \mathbf{Z}) - \theta_{\mathbf{X}}^2(s, \mathbf{Z})| \right. \\ &+ (1 + |\theta_{\mathbf{X}}^2(s, \mathbf{Z})|) |v_{\mathbf{X}}^1(s, \varphi(\mathbf{Z})) - v_{\mathbf{X}}^2(s, \varphi(\mathbf{Z}))| \rangle d\mathbf{Z} ds \\ &\leq \int_t^T \int_{\mathbb{R}^{2n}} \frac{K}{(s-t)^{\frac{2n+1}{2}}} e^{-\lambda \frac{|\mathbf{X}-\mathbf{Z}|^2}{4(s-t)}} |\theta_{\mathbf{X}}^1(s, \mathbf{Z}) - \theta_{\mathbf{X}}^2(s, \mathbf{Z})| d\mathbf{Z} ds \\ &+ K(T-t)^{\frac{1}{2}} \left(\|F(\cdot)\|_{C^1(\mathbb{R}^n)} + 1 \right) \|v_{\mathbf{X}}^1(\cdot, \cdot) - v_{\mathbf{X}}^2(\cdot, \cdot)\|_{\mathbb{C}([t, T]; C(\mathbb{R}^{2n}))} \end{aligned}$$

Then, following an iterative procedure as the one used to obtain (7.17), we can obtain

$$\begin{aligned} & \sup_{\mathbf{X} \in \mathbb{R}^{2n}} |\theta_{\mathbf{X}}^1(t, \mathbf{X}) - \theta_{\mathbf{X}}^2(t, \mathbf{X})| \\ &\leq K(T-t)^{\frac{1}{2}} \left(\|F(\cdot)\|_{C^1(\mathbb{R}^n)} + 1 \right) \|v_{\mathbf{X}}^1(\cdot, \cdot) - v_{\mathbf{X}}^2(\cdot, \cdot)\|_{\mathbb{C}([t, T]; C(\mathbb{R}^{2n}))}. \end{aligned}$$

On the other hand, from (7.18), we have

$$\begin{aligned} & \sup_{\mathbf{X} \in \mathbb{R}^{2n}} |\theta^1(t, \mathbf{X}) - \theta^2(t, \mathbf{X})| \\ &\leq \int_t^T \int_{\mathbb{R}^{2n}} \left(\frac{K}{(s-t)^{\frac{2n}{2}}} e^{-\lambda \frac{|\mathbf{X}-\mathbf{Z}|^2}{4(s-t)}} |\theta_{\mathbf{X}}^1(s, \mathbf{Z}) - \theta_{\mathbf{X}}^2(s, \mathbf{Z})| \right. \\ &+ (1 + |\theta_{\mathbf{X}}^2(s, \mathbf{Z})|) |v_{\mathbf{X}}^1(s, \varphi(\mathbf{Z})) - v_{\mathbf{X}}^2(s, \varphi(\mathbf{Z}))| \rangle d\mathbf{Z} ds \\ &\leq K(T-t) \left(\|F(\cdot)\|_{C^1(\mathbb{R}^n)} + 1 \right) \|v_{\mathbf{X}}^1(\cdot, \cdot) - v_{\mathbf{X}}^2(\cdot, \cdot)\|_{\mathbb{C}([t, T]; C(\mathbb{R}^{2n}))}. \end{aligned}$$

Hence we obtain

$$\begin{aligned} & \sup_{\mathbf{X} \in \mathbb{R}^{2n}} |\theta^1(t, \mathbf{X}) - \theta^2(t, \mathbf{X})| + \sup_{\mathbf{X} \in \mathbb{R}^{2n}} |\theta_{\mathbf{X}}^1(t, \mathbf{X}) - \theta_{\mathbf{X}}^2(t, \mathbf{X})| \\ &\leq K(T-t)^{\frac{1}{2}} \left(\|F(\cdot)\|_{C^1(\mathbb{R}^n)} + 1 \right) \|v_{\mathbf{X}}^1(\cdot, \cdot) - v_{\mathbf{X}}^2(\cdot, \cdot)\|_{\mathbb{C}([t, T]; C(\mathbb{R}^{2n}))} \\ &\leq K(T-t)^{\frac{1}{2}} \left(\|F(\cdot)\|_{C^1(\mathbb{R}^n)} + 1 \right) \|v^1(\cdot, \cdot) - v^2(\cdot, \cdot)\|_{\mathbb{C}([t, T]; C^1(\mathbb{R}^{2n}))}. \end{aligned}$$

Thus, in particular, we have

$$\begin{aligned} & \|\theta^1(\cdot, \cdot) - \theta^2(\cdot, \cdot)\|_{\mathbb{C}([T-\delta, T]; C^1(\mathbb{R}^{2n}))} \\ & \leq K\delta^{\frac{1}{2}} \left(\|F(\cdot)\|_{C^1(\mathbb{R}^n)} + 1 \right) \|v^1(\cdot, \cdot) - v^2(\cdot, \cdot)\|_{\mathbb{C}([T-\delta, T]; C^1(\mathbb{R}^{2n}))}. \end{aligned}$$

Clearly, by choosing $\delta > 0$ small enough, we get a contraction mapping on $v(\cdot, \cdot) \rightarrow \theta(\cdot, \cdot)$ on $\mathbb{C}([T-\delta, T]; C^1(\mathbb{R}^{2n}))$. Therefore, this map admits a unique fixed point. Since we may obtain similar estimates on $\mathbb{C}([T-2\delta, T-\delta]; C^1(\mathbb{R}^{2n}))$, etc., one sees that the fixed point will exist on the whole space $\mathbb{C}([0, T]; C^1(\mathbb{R}^{2n}))$ for the map $v(\cdot, \cdot) \rightarrow \theta(\cdot, \cdot)$. Then we obtain the well-posedness of the following

$$\begin{aligned} & \theta(t, \mathbf{X}) \\ & = \int_{\mathbb{R}^{2n}} \Gamma^0(t, \mathbf{X}; T, \mathbf{Z}) \mathbf{F}(\mathbf{Z}) d\mathbf{Z} \\ & + \int_t^T \Gamma^0(t, \mathbf{X}; s, \mathbf{Z}) \langle \mathbf{b}(t, \mathbf{Z}, \theta_{\mathbf{X}}(t, \varphi(\mathbf{Z}))), \theta_{\mathbf{X}}(t, \mathbf{Z}) \rangle d\mathbf{Z} ds \\ & + \int_t^T \Gamma^0(t, \mathbf{X}; s, \mathbf{Z}) \mathbf{f}(t, \mathbf{Z}, \theta_{\mathbf{X}}(t, \varphi(\mathbf{Z}))) d\mathbf{Z} ds. \end{aligned}$$

Finally, by the regularity of $\Gamma^0(t, \mathbf{X}; T, \mathbf{Z})$, we know that $\theta(t, \mathbf{X})$ is $C^{2+\alpha}$ in $\mathbf{X} = (x, y)$, $C^{1+\alpha}$ in t for some $\alpha \in (0, 1)$, and the PDE (7.9) is satisfied. \blacksquare

References

- [1] N. Agram, B. Øksendal, *A Hida-Malliavin white noise calculus approach to optimal control*. Infinite Dimensional Analysis, Quantum Probability and Related Topics, 21(03), (2018), id. 1850014-496.
- [2] I. Alia, *Time-Inconsistent Stochastic Optimal Control Problems: A Backward Stochastic Partial Differential Equations Approach*. Mathematical Control & Related Fields (2020), doi:10.3934/mcrf.2020020.
- [3] S. Bahlali, B. Mezerdi, *A general stochastic maximum principle for singular control problems*, Electronic Journal of Probability. 10, (2005), 988–1004.
- [4] A. Bensoussan, *Lectures on stochastic control*. In S.K. Mitter and A. Moro, editors, Lecture Notes in Mathematics: Nonlinear Filtering and Stochastic Control, Springer-Verlag, 972, (1982), 1-62.
- [5] A. Bensoussan, *Maximum principle and dynamic programming approaches of the optimal control of partially observed diffusions*, Stochastics, 9, (1983), 169–222.
- [6] J.M. Bismut, *An Introductory Approach to Duality in Optimal Stochastic Control*. SIAM Review, 20(1), (1978), 62–78.
- [7] R. Buckdahn, B. Djehiche, J. Li, *A General Stochastic Maximum Principle for SDEs of Mean-Field Type*. Applied Math. and Optimization. 64(2), (2011), 197-216.
- [8] S. Chen, S. Tang. *Semi-linear backward stochastic integral partial differential equations driven by a Brownian motion and a Poisson point process*. Mathematical Control & Related Fields, 5(3), (2015) 401–434.
- [9] A. Friedman, *Partial Differential Equations of Parabolic Type*, Prentice Hall, Inc., Englewood Cliffs, N.J, (1964).
- [10] K. Du, S. Tang, Q. Zhang, *$W^{m,p}$ -Solution ($p \geq 2$) of Linear Degenerate Backward Stochastic Partial Differential Equations in the Whole Space*, Journal of Differential Equations, 254(7), (2013), 2877–2904.
- [11] K. Du, Q. Zhang, *Semi-linear degenerate backward stochastic partial differential equations and associated forward-backward stochastic differential equations*, Stochastic Processes and their Applications, 123, (2013), 1616–1637.
- [12] R.J. Elliott, M. Kohlmann, *The variational principle and stochastic optimal control*. Stochastics, 3, (1980), 229-241.

- [13] U.G. Haussmann, *General necessary conditions for optimal control of stochastic systems*, Mathematical Programming Studies, 6, (1986), 30-48.
- [14] U.G. Haussmann, *A Stochastic maximum principle for optimal control of diffusions*, Pitman Research Notes in Math, Series 151 (1986).
- [15] Y. Hu, J. Ma, J. Yong. *On semi-linear degenerate backward stochastic partial differential equations*, Probability Theory and Related Fields 123 (3), (2002), 381–411.
- [16] M. Hu, *Stochastic global maximum principle for optimization with recursive utilities*, Probability, Uncertainty and Quantitative Risk, 2, 1 (2017). <https://doi.org/10.1186/s41546-017-0014-7>.
- [17] N. V. Krylov, *On the Itô-Wentzell formula for distribution-valued processes and related topics*, Probability Theory and Related Fields, 150, (2011), 295-319.
- [18] H. Kunita. *Stochastic Flows and Stochastic Differential Equations*. Cambridge University Press, (1990).
- [19] H. Kunita, *Stochastic Flows and Jump-Diffusions*, Springer Singapore, (2019).
- [20] H. Kunita, *Some extensions of Ito's formula*, Séminaire de Probabilités XV 1979/80, Springer, 1981, 118–141.
- [21] H Kushner, *Necessary conditions for continuous parameter stochastic optimization problems*. SIAM Journal on Control and Optimization 10(3), (1972),550-565.
- [22] J. Ma, J. Yong, *Adapted solution of a degenerate backward SPDE, with applications*, Stochastic Processes and their Applications 70, 59–84 (1997)
- [23] J. Ma, J. Yong, *On linear degenerate backward stochastic partial differential equations*, Probability Theory and Related Fields 113, (1999), 135–170.
- [24] J. Ma, H. Yin and J. F. Zhang, *On non-Markovian forward-backward SDEs and backward stochastic PDEs*, Stochastic Processes and their Applications, 122, (2012), 3980-4004.
- [25] J. Ma, P. Protter, J. Yong. *Solving forward-backward stochastic differential equations explicitly a four step scheme*. Probability Theory and Related Fields, 98(3), (1994) ,339-359.
- [26] J. Ma, J. Yong, *Forward-Backward Stochastic Differential Equations and Their Applications*, Lecture Notes in Mathematics, 1702, Springer-Verlag, Berlin, (1999)
- [27] N. Nagasa, M. Nisio, *Optimal controls for stochastic partial differential equations*, SIAM Journal on Control and Optimization, 28, (1990), 186–213.
- [28] D. Ocone, E. Pardoux. *A generalized Itô-Wentzell formula. Application to a class of anticipating stochastic differential equations*. Annales de l'I. H. P., section B, tome 25(1), (1989), 39-71.
- [29] E. Pardoux, S. Peng, *Adapted solution of a backward stochastic differential equation*. Systems Control Letters., 14(1-2), (1990), 61-74.
- [30] S. Peng, *A general stochastic maximum principle for optimal control problems*. SIAM Journal on Control and Optimization, 28, (1990), 966-979.
- [31] S. Peng, *Maximum Principle for Stochastic Optimal Control with Nonconvex Control Domain*, in Analysis & Optimization of Systems, A. Bensoussan & J. L. Lions eds. Lecture Notes in Control & Information Sciences, 114 (1990), pp. 724-732.
- [32] S. Peng, *Stochastic Hamilton-Jacobi-Bellman equations*, SIAM Journal on Control and Optimization, 30, (1992), 284–304.
- [33] L. S. Pontryagin, V. G. Boltyanskii, R. V. Gamkrelidze and E. F. Mischenko, *The Mathematical Theory of Optimal Processes*, Wiley-Interscience, New York, 1962.
- [34] S. Tang, X. Li, *Necessary conditions for optimal control for stochastic systems with random jumps*, SIAM Journal on Control and Optimization, **32** (1994), 1447-1475.

- [35] S. Tang, *Semi-linear systems of backward stochastic partial differential equations in \mathbb{R}^n* , Chinese Annals of Mathematics, 26, (2005) 437–456.
- [36] J. Yong, , *Optimality variational principle for controlled forwardbackward stochastic differential equations with mixed initial-terminal conditions*, SIAM Journal on Control and Optimization, 48(6), (2010), 4119-4156.
- [37] J. Yong, *Time-inconsistent optimal control problems and the equilibrium HJB equation*, Mathematical Control & Related Fields, 2(3), (2012), 271-329.
- [38] J. Yong, X.Y. Zhou, *Stochastic Controls: Hamiltonian Systems and HJB Equations*. Springer-Verlag, New York. (1999).
- [39] X.Y. Zhou, *On the necessary conditions of optimal controls for stochastic partial differential equations*, SIAM Journal on Control and Optimization, 31, (1993), 1462–1478.
- [40] X. Y. Zhou, *On the necessary conditions of optimal controls for stochastic partial differential equations*, SIAM Journal on Control and Optimization, 31, (1993), 1462-1478.

Department of Mathematics University of Bordj Bou Arreridj 34000 Algeria; izacalia@yahoo.com, fermasof@yahoo.fr.