

Relative Stability in the Sup-norm and Input-to-state Stability in the Spatial Sup-norm for Parabolic PDEs

Jun Zheng^{1,2}, Guchuan Zhu², Sergey Dashkovskiy³

¹*School of Mathematics, Southwest Jiaotong University
Chengdu, Sichuan, P. R. of China 611756*

²*Department of Electrical Engineering, Polytechnique Montréal
P.O. Box 6079, Station Centre-Ville, Montreal, QC, Canada H3T 1J4*

³*Institute of Mathematics, University of Würzburg
Emil-Fischer-Str. 40, Würzburg, Germany 97074*

Abstract

In this paper, we introduce the notion of *relative \mathcal{K} -equi-stability* (RKES) to characterize the uniformly continuous dependence of (weak) solutions on external disturbances for nonlinear parabolic PDE systems. Based on the RKES, we prove the input-to-state stability (ISS) in the spatial sup-norm for a class of nonlinear parabolic PDEs with either Dirichlet or Robin boundary disturbances. Two examples, concerned respectively with a super-linear parabolic PDE with Robin boundary condition and a 1-D parabolic PDE with a destabilizing term, are provided to illustrate the obtained ISS results. Besides, as an application of the notion of RKES, we conduct stability analysis for a class of parabolic PDEs in cascade coupled over the domain or on the boundary of the domain, in the spatial and time sup-norm, and in the spatial sup-norm, respectively. The technique of De Giorgi iteration is extensively used in the proof of the results presented in this paper.

Key words: Nonlinear PDEs, relative stability, input-to-state stability, De Giorgi iteration, cascade of PDE systems.

1 Introduction

Originally introduced by Sontag in the late 1980s, the notion of *input-to-state stability* (ISS) has been proven to be a convenient tool for describing robust stability of finite dimensional systems with external inputs. The pioneering work on extending the application of ISS to infinite dimensional systems is owe to [2] and [3], where different methods were proposed for constructing ISS-Lyapunov functions for abstract equations in Banach spaces, and impulsive systems, respectively. Particularly, as an application of the proposed methods, ISS-Lyapunov functions and ISS estimates were presented for some nonlinear single and interconnected parabolic PDEs in [2]. Since then, the ISS of PDE systems has drawn much attention in the literature of PDE control. It is worth noting that applying the classical regularity theory of PDEs to ISS analysis of PDEs having only in-domain disturbances seems to be straightforward, while it is a challenge to establish the ISS for PDEs that have external disturbances distributed on the boundary of the domain.

In recent years, a great effort has been devoted to establishing the ISS for PDEs with boundary disturbances; see [9,16] for comprehensive surveys on this topic, and [23,24] for a summary of different approaches for establishing ISS of PDEs with boundary disturbances. Among the existing literature, the ISS in L^1 -norm and L^q -norm with $q \in [2, +\infty)$ has been well studied for PDEs with boundary disturbances via different methods (see, e.g., [24], and [5,6,7,15,17,18,21,20,22], etc.), while few results are concerned with ISS in L^∞ -norm except [8], where the ISS in various norms, including weighted L^∞ -norm, was considered for linear 1-D PDEs governed by Sturm-Liouville operators by exploiting the eigenfunction expansion and

Email addresses: zhengjun2014@aliyun.com (Jun Zheng^{1,2}), guchuan.zhu@polymtl.ca (Guchuan Zhu²), sergey.dashkovskiy@mathematik.uni-wuerzburg.de (Sergey Dashkovskiy³).

the finite difference scheme; [10], where ISS-style estimates in the spatial sup-norm was established for classical solutions of nonlinear 1-D parabolic PDEs by using an ISS Lyapunov functional for the sup-norm; and [21], where under an appropriate boundary feedback law and with compatibility conditions, an ISS estimate in L^∞ -norm was established for a 1-D linear parabolic equation with a destabilizing term.

The aim of this paper is to provide a new method for establishing the ISS in the spatial sup-norm for (weak) solutions of a class of higher dimensional nonlinear parabolic PDEs with boundary disturbances, which is different from those developed in [8,10], and [21] concerning classical solutions of 1-D PDEs. More precisely, in order to establish ISS estimates in the spatial sup-norm, we borrow first the notion of *relative stability* (RS) from [11], which was used to characterize a kind of relationship of stabilities for two control systems, to describe the uniformly continuous dependence on the external disturbances for weak solutions of nonlinear PDE systems. Then, based on the RS in the (spatial and time) sup-norm, we establish the ISS in the spatial sup-norm for the considered higher dimensional nonlinear PDEs with Dirichlet and Robin boundary disturbances. Moreover, we show how to apply the property of RS to characterize the stability in the sup-norm, and the spatial sup-norm, respectively, for a class of PDE systems in cascade coupled via the boundary or over the domain.

The main tool exploited in this paper for the proof of various stability properties is the De Giorgi iteration, which has been used for the first time to establish ISS estimates for classical solutions of PDEs in [21]. It should be mentioned that in [21] the De Giorgi iteration is used for addressing the ISS of 1-D parabolic PDEs with Dirichlet boundary disturbances by combining the technique of splitting, which requires certain compatibility conditions. While in this paper, the De Giorgi iteration is used for not only 1-D PDEs with Dirichlet boundary conditions, but also for higher dimensional PDEs with either Dirichlet or Robin boundary conditions. In addition, unlike [21], we consider in this paper solutions in a weak sense. Therefore, we do not use any splitting technique and impose any compatibility condition, which is an improvement of the results obtained in [21].

In summary, the main contribution of this paper includes:

- (i) introducing RKES to describe the uniformly continuous dependence of weak solutions on the external disturbances and establishing RKES estimates for a class of higher dimensional nonlinear parabolic PDEs;
- (ii) establishing the ISS in the spatial sup-norm for weak solutions of higher dimensional PDEs with Dirichlet or Robin boundary disturbances by using the property of RKES;
- (iii) establishing stability estimates in the spatial and time sup-norm and the spatial sup-norm, respectively for a class of parabolic systems in cascade, which are interconnected or coupled via the boundary of the domain;
- (iv) extending the usage of De Giorgi iteration to ISS analysis of higher dimensional PDEs with Robin boundary conditions.

In the rest of the paper, we introduce first some basic notations. Section 2 presents the problem formulation, well-posedness, notions on relative stability, and the main results on RKES in the sup-norm and ISS in the spatial sup-norm for the considered PDE systems. Section 3 provides two examples to illustrate the obtained ISS results. As an application of REKS presented in Section 2, we show in Section 4 how to apply REKS to obtain stability estimates in the spatial and time sup-norm and the spatial sup-norm, respectively, for a class of parabolic systems in cascade connected over the domain or on the boundary of the domain. Some concluding remarks are given in Section 5.

Notations. \mathbb{R}_+ denotes the set of positive real numbers and $\mathbb{R}_{\geq 0} := \{0\} \cup \mathbb{R}_+$. Ω denotes a bounded domain in $\mathbb{R}^n (n \geq 1)$ of class C^2 , that is, $\overline{\Omega}$ is an n -dimensional C^2 -submanifold of \mathbb{R}^n with boundary $\partial\Omega$. $|\Omega|$ denotes the n -dimensional Lebesgue measure of Ω . For any $T > 0$, $Q_T := \Omega \times (0, T)$, $\partial_t Q_T := \partial\Omega \times (0, T)$ and $\partial_p Q_T := \partial\Omega \times (0, T) \cup \{(x, t) | x \in \overline{\Omega}, t = 0\}$.

$\mathcal{K} := \{\gamma : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0} | \gamma(0) = 0, \gamma \text{ is continuous, strictly increasing}\}$, $\mathcal{L} := \{\gamma : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0} | \gamma \text{ is continuous, strictly decreasing, } \lim_{s \rightarrow \infty} \gamma(s) = 0\}$, $\mathcal{KL} := \{\beta : \mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0} | \beta(\cdot, t) \in \mathcal{K}, \forall t \in \mathbb{R}_{\geq 0}, \text{ and } \beta(s, \cdot) \in \mathcal{L}, \forall s \in \mathbb{R}_+\}$.

Throughout this paper, all notations on function spaces are standard, which can be found in, e.g., [4,19]. Let $\mathbb{A} := C^1(\overline{\Omega}; \mathbb{R}_+)$, $\mathbb{C} := C(\overline{\Omega}; \mathbb{R}_{\geq 0})$, $\mathbb{M} := C(\partial\Omega; \mathbb{R}_+)$, $\mathbb{H} := C^{0,1}(\overline{\Omega} \times \mathbb{R}_{\geq 0}) \times \mathbb{R}; \mathbb{R}$, $\mathbb{F} := C(\overline{\Omega} \times \mathbb{R}_{\geq 0}; \mathbb{R})$, $\mathbb{D} := C(\partial\Omega \times \mathbb{R}_{\geq 0}; \mathbb{R})$, and $\mathbb{U} := W^{1,p}(\Omega)$, where $p \geq 2$ and $p > n$. For functions $a \in \mathbb{A}$, $c \in \mathbb{C}$, $m \in \mathbb{M}$, we always denote

$$\underline{a} := \min_{x \in \overline{\Omega}} a > 0, \underline{c} := \min_{x \in \overline{\Omega}} c \geq 0, \underline{m} := \min_{x \in \partial\Omega} m > 0. \tag{1}$$

2 Problem setting and main results

2.1 Problem formulation and well-posedness

For functions $a \in \mathbb{A}, c \in \mathbb{C}, m \in \mathbb{M}, h \in \mathbb{H}, f \in \mathbb{F}, d \in \mathbb{D}, u^0 \in \mathbb{U}$, we consider the stability of the following higher dimensional nonlinear parabolic system:

$$\mathcal{L}[u] + h(x, t, u) = f \quad \text{in } \Omega \times \mathbb{R}_+, \quad (2a)$$

$$\mathcal{B}[u] = d \quad \text{on } \partial\Omega \times \mathbb{R}_+, \quad (2b)$$

$$u(\cdot, 0) = u^0(\cdot) \quad \text{in } \Omega, \quad (2c)$$

where $\mathcal{L}[u] := u_t - \operatorname{div}(a\nabla u) + cu$, and

$$\mathcal{B}[u] := a \frac{\partial u}{\partial \nu} + mu, \quad (3)$$

or

$$\mathcal{B}[u] := u, \quad (4)$$

represents the Robin boundary condition, or the Dirichlet boundary condition, respectively.

We always assume that for any $T > 0$, there exist a positive constant c_0 , an increasing function $H : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$, and a function $\Psi \in C(\mathbb{R}_{\geq 0}; \mathbb{R}_{\geq 0})$ satisfying $\Psi(0) = 0$, such that

$$|h(x, t, \xi)| \leq c_0(1 + |\xi|^\lambda), |\partial_\xi h(x, t, \xi)| \leq H(|\xi|), \quad (5a)$$

$$|h(x, s, \xi) - h(x, t, \xi) - f(x, s) + f(x, t)| \leq H(|\xi|)\Psi(|s - t|), \quad (5b)$$

holds for all $x \in \overline{\Omega}, s, t \in [0, T], \xi \in \mathbb{R}$, where $\lambda \in [1, 1 + \frac{2}{n}]$ is a constant.

We provide a definition of a (weak) solution of the system (2).

Definition 1 (i) We say that u is a weak solution of the system (2) with the Robin boundary condition (3), if for any $T > 0$:

$$u \in C([0, T]; W^{1,p}(\Omega)), u(\cdot, 0) = u^0(\cdot) \quad \text{in } \Omega,$$

and the equality

$$\begin{aligned} & - \int_0^T \int_\Omega u \eta_t dx dt + \int_0^T \int_\Omega a \nabla u \nabla \eta dx dt + \int_0^T \int_\Omega (cu + h(x, t, u)) \eta dx dt \\ & = \int_0^T \int_\Omega f \eta dx dt + \int_0^T \int_{\partial\Omega} (d - mu) \eta dx dt + \int_\Omega u^0(x) \eta(x, 0) dx \end{aligned}$$

holds true for any $\eta \in C([0, T]; (W^{1,p}(\Omega))') \cap C^1((0, T); L^{p'}(\Omega))$ with $\eta(\cdot, T) = 0$ in Ω , where $p' = \frac{p}{p-1}$, and $(W^{1,p}(\Omega))'$ is the dual space of $W^{1,p}(\Omega)$.

(ii) We say that u is a weak solution of the system (2) with the Dirichlet boundary condition (4), if for any $T > 0$:

$$u \in C([0, T]; W^{1,p}(\Omega)), u = d \text{ on } \partial_t Q_T, u(\cdot, 0) = u^0(\cdot) \quad \text{in } \Omega,$$

and the equality

$$- \int_0^T \int_\Omega u \eta_t dx dt + \int_0^T \int_\Omega a \nabla u \nabla \eta dx dt + \int_0^T \int_\Omega (cu + h(x, t, u)) \eta dx dt = \int_0^T \int_\Omega f \eta dx dt + \int_\Omega u^0(x) \eta(x, 0) dx$$

holds true for any $\eta \in C([0, T]; (W_0^{1,p}(\Omega))') \cap C^1((0, T); L^{p'}(\Omega))$ with $\eta(\cdot, T) = 0$ in Ω , where $p' = \frac{p}{p-1}$, and $(W_0^{1,p}(\Omega))'$ is the dual space of $W_0^{1,p}(\Omega)$.

For the well-posedness of the considered problem, we have the following result.

Proposition 2.1 *System (2) with either the Robin boundary condition (3), or the Dirichlet boundary condition (4), admits a unique weak solution belonging to $C([0, T]; W^{1,p}(\Omega)) \cap C^1((0, T); L^{p'}(\Omega))$ for any $T > 0$, where $p' = \frac{p}{p-1}$.*

Proof *For any $T > 0$, by Theorem 14.5 of [1], the system (2) with the Robin boundary condition (3), or the Dirichlet boundary condition (4), admits a unique maximal weak solution $u \in C([0, T_0]; W^{1,p}(\Omega)) \cap C^1((0, T_0); L^{p'}(\Omega))$ with some $T_0 \in (0, T]$. Furthermore, according to Theorem 15.2(i) of [1], if there exists a positive constant C such that the following a priori estimate holds true:*

$$\|u(\cdot, t)\|_{L^1(\Omega)} \leq C, \forall t \in (0, T_0), \quad (6)$$

then, such a maximal solution u must exist globally on $[0, T]$. Therefore, it suffices to prove that (6) holds true for the maximal solution of the system (2) with the Robin boundary condition (3), or the Dirichlet boundary condition (4), respectively.

Indeed, for system (2) with the Robin boundary condition (3), the estimate in (6) is guaranteed by [24, Theorem 3.1]. For system (2) with the Dirichlet boundary condition (4), the estimate in (6) is guaranteed by applying the estimate given in Theorem 2.2(ii) to (2) defined over Q_{T_0} . ■

Remark 1 *The growth conditions on the nonlinear term h appearing in (5) are only used for guaranteeing the existence and uniqueness of a weak solution. In particular, as indicated in [1], the exponent $1 + \frac{2}{n}$ in (5a) is optimal for the existence of a global weak solution. However, if a certain compatibility condition is imposed, and a smooth solution is considered, then the growth conditions on the nonlinear term h in (5) can be relaxed; see [24, Proposition 2.1].*

2.2 Notion on relative stability

From the point of view on PDEs, both the initial value and enforced terms (external disturbances) have a deep effect on the stability (and well-posedness) of PDE systems. In order to describe the influence induced by these data, we define some stability characteristics for PDE systems. More precisely, based on the notion of *relative stability* (RS) given by [11], which is an extension of the concept for finite dimensional systems, e.g. [13], to general control systems, we define several properties of relative stability for the considered PDE systems.

To emphasize the dependence of the solutions on the initial value and external disturbances, we denote by $\Sigma(\mathbb{U}, \mathbb{F}, \mathbb{D})$ the system (2) with data $(u^0, f, d) \in \mathbb{U} \times \mathbb{F} \times \mathbb{D}$. Note that if $v \in W^{1,p}(\Omega)$ with $p > n$, then $v \in C^{1-\frac{n}{p}}(\bar{\Omega})$; see, e.g. [19, Theorem 1.3.2]. Thus, $\sup_{x \in \Omega} |v(x)| < +\infty$ and hence, $\sup_{x \in \Omega} |v(x)|$ is well-defined.

Definition 2 *The system $\Sigma(\mathbb{U}, \mathbb{F}, \mathbb{D})$ is said to be relatively equi-stable (RES) in the sup-norm with respect to (w.r.t.) in-domain and boundary disturbances in $\mathbb{F} \times \mathbb{D}$, if for every constant $\varepsilon > 0$, there exists a positive constant δ depending only on ε , such that the following implication*

$$\sup_{(x,t) \in Q_T} |f_1(x,t) - f_2(x,t)| + \sup_{(x,t) \in \partial_t Q_T} |d_1(x,t) - d_2(x,t)| < \delta \Rightarrow \sup_{(x,t) \in Q_T} |u_1(x,t) - u_2(x,t)| < \varepsilon$$

holds true for all $(f_1, d_1), (f_2, d_2) \in \mathbb{F} \times \mathbb{D}$ and all $T > 0$, where u_i is the solution of the system $\Sigma(\mathbb{U}, \mathbb{F}, \mathbb{D})$ corresponding to the data $(u^0, f_i, d_i) \in \mathbb{U} \times \mathbb{F} \times \mathbb{D}, i = 1, 2$.

Definition 3 *The system $\Sigma(\mathbb{U}, \mathbb{F}, \mathbb{D})$ is said to be relatively \mathcal{K} -equi-stable (RKES) in the sup-norm w.r.t. in-domain and boundary disturbances in $\mathbb{F} \times \mathbb{D}$, if there exist functions $\gamma_d, \gamma_f \in \mathcal{K}$ such that*

$$\sup_{(x,t) \in Q_T} |u_1(x,t) - u_2(x,t)| \leq \gamma_f \left(\sup_{(x,t) \in Q_T} |f_1(x,t) - f_2(x,t)| \right) + \gamma_d \left(\sup_{(x,t) \in \partial_t Q_T} |d_1(x,t) - d_2(x,t)| \right), \forall T > 0,$$

where u_i is the solution of the system $\Sigma(\mathbb{U}, \mathbb{F}, \mathbb{D})$ corresponding to the data $(u^0, f_i, d_i) \in \mathbb{U} \times \mathbb{F} \times \mathbb{D}, i = 1, 2$.

Particularly, the system $\Sigma(\mathbb{U}, \mathbb{F}, \mathbb{D})$ is said to be relatively Lipschitz-equi-stable (RLES) in the sup-norm w.r.t. in-domain and boundary disturbances in $\mathbb{F} \times \mathbb{D}$, if $\gamma_f(s) = L_f s, \gamma_d(s) = L_d s$ for any $s \geq 0$, where L_f and L_d are certain positive constants.

Remark 2 *We provide some comments on RES.*

- (i) *It should be noticed that there is a slight difference for RS between the definition given in [11] and the one given in this paper. Indeed, RES defined in [11] is mainly used to describe the relationship of stabilities between two systems, namely it is in term of multiple different systems, while RES defined in this paper is mainly used to characterize the uniformly*

continuous dependence of the solution on the external disturbances for a certain PDE, namely it is in term of one system (i.e., the nominal dynamics of the PDEs are governed by the same differential operator) with different external inputs.

(ii) Note that a system with external disturbances may be RES w.r.t. external disturbances while not being asymptotically stable. For example, we consider the following systems:

$$\begin{aligned} (u_k)_t - (u_k)_{xx} &= f_k(x, t), & (x, t) &\in \left(0, \frac{\pi}{2}\right) \times \mathbb{R}_{\geq 0}, \\ u_k(0, t) &= 0, u_k\left(\frac{\pi}{2}, t\right) &= d_k(t), & t \in \mathbb{R}_+, \\ u_k(x, 0) &= u^0(x), & x &\in \left(0, \frac{\pi}{2}\right), \end{aligned}$$

where $u^0(x) := 0$, $f_k(x, t) := \sqrt{2}k \sin x \cos\left(t - \frac{\pi}{4}\right)$, $d_k(t) := k \sin t$, and $k \in \mathbb{R}_+$. It is clear that $u_k(x, t) = k \sin t \sin x$ is the unique solution, which is bounded for any fixed k . Since there exists a point (x_0, t_0) such that $u_k(x_0, t_0) > \frac{k}{2}$, u_k is unbounded as $k \rightarrow +\infty$. However, noting that

$$\sup_{(x,t) \in \left(0, \frac{\pi}{2}\right) \times (0, T)} |f_k(x, t) - f_l(x, t)| = \sqrt{2}|k - l| \sup_{(x,t) \in \left(0, \frac{\pi}{2}\right) \times (0, T)} |\sin x \cos\left(t - \frac{\pi}{4}\right)|, \forall T > 0,$$

and

$$\sup_{t \in (0, T)} |d_k(t) - d_l(t)| = |k - l| \sup_{t \in (0, T)} |\sin t|, \forall T > 0,$$

for all $k, l \in \mathbb{R}_+$, then the system is RLES, having the estimate for all $k, l \in \mathbb{R}_+$:

$$\begin{aligned} \sup_{(x,t) \in \left(0, \frac{\pi}{2}\right) \times (0, T)} |u_k(x, t) - u_l(x, t)| &= |k - l| \sup_{(x,t) \in \left(0, \frac{\pi}{2}\right) \times (0, T)} |\sin t \sin x| \\ &\leq \sup_{(x,t) \in \left(0, \frac{\pi}{2}\right) \times (0, T)} |f_k(x, t) - f_l(x, t)| + \sup_{t \in (0, T)} |d_k(t) - d_l(t)|, \forall T > 0. \end{aligned}$$

(iii) It is obvious that RLES \Rightarrow RKES \Rightarrow RES.

Definition 4 ([17]) The system $\Sigma(\mathbb{U}, \mathbb{F}, \mathbb{D})$ is said to be globally asymptotically stable at zero uniformly w.r.t. the state (0-UGAS w.r.t. the state) in the spatial sup-norm if there exists a function $\beta \in \mathcal{KL}$ such that

$$\sup_{x \in \Omega} |u(x, T)| \leq \beta \left(\sup_{x \in \Omega} |u^0(x)|, T \right), \forall u^0 \in \mathbb{U}, \forall T > 0,$$

where u is the solution of the system $\Sigma(\mathbb{U}, \mathbb{F}, \mathbb{D})$ corresponding to the data $(u^0, 0, 0) \in \mathbb{U} \times \mathbb{F} \times \mathbb{D}$.

Definition 5 The system $\Sigma(\mathbb{U}, \mathbb{F}, \mathbb{D})$ is said to be input-to-state stable (ISS) in the spatial sup-norm w.r.t. in-domain and boundary disturbances in $\mathbb{F} \times \mathbb{D}$, if there exist functions $\beta \in \mathcal{KL}$ and $\gamma_f, \gamma_d \in \mathcal{K}$ such that for all $(u^0, f, d) \in \mathbb{U} \times \mathbb{F} \times \mathbb{D}$:

$$\sup_{x \in \Omega} |u(x, T)| \leq \beta \left(\sup_{x \in \Omega} |u^0(x)|, T \right) + \gamma_f \left(\sup_{(x,t) \in Q_T} |f(x, t)| \right) + \gamma_d \left(\sup_{(x,t) \in \partial_t Q_T} |d(x, t)| \right), \forall T > 0, \quad (7)$$

where u is the solution of the system $\Sigma(\mathbb{U}, \mathbb{F}, \mathbb{D})$ corresponding to the data $(u^0, f, d) \in \mathbb{U} \times \mathbb{F} \times \mathbb{D}$.

Furthermore, the system $\Sigma(\mathbb{U}, \mathbb{F}, \mathbb{D})$ is said to be exponentially input-to-state stable (EISS) in the spatial sup-norm w.r.t. in-domain and boundary disturbances in $\mathbb{F} \times \mathbb{D}$, if there exist constants $M, \sigma > 0$ such that $\beta(r, t) = Mre^{-\sigma t}$ in (7) for all $r \geq 0$.

Proposition 2.2 If the system $\Sigma(\mathbb{U}, \mathbb{F}, \mathbb{D})$ is RKES in the sup-norm w.r.t. in-domain and boundary disturbances in $\mathbb{F} \times \mathbb{D}$ and 0-UGAS w.r.t. the state in the spatial sup-norm, then it is ISS in the spatial sup-norm w.r.t. in-domain and boundary disturbances in $\mathbb{F} \times \mathbb{D}$.

Proof For any $(u^0, f, d) \in \mathbb{U} \times \mathbb{F} \times \mathbb{D}$, let u, v be the solutions of the system $\Sigma(\mathbb{U}, \mathbb{F}, \mathbb{D})$ with data (u^0, f, d) and $(u^0, 0, 0)$, respectively. For simplicity, we write $|g|_{\infty, \omega} := \sup_{y \in \omega} |g(y)|$ for a function g defined on a domain ω of \mathbb{R}^n or \mathbb{R}^{n+1} . Since

$\Sigma(\mathbb{U}, \mathbb{F}, \mathbb{D})$ is RKES in the sup-norm w.r.t. in-domain and boundary disturbances, and u and v are continuous in T (see Proposition 2.1), there exist functions $\gamma_d, \gamma_f \in \mathcal{K}$ such that

$$\begin{aligned} |u(\cdot, T) - v(\cdot, T)|_{\infty, \Omega} &\leq |u - v|_{\infty, Q_T} \\ &\leq \gamma_f(|f - 0|_{\infty, Q_T}) + \gamma_d(|d - 0|_{\infty, \partial_t Q_T}) \\ &= \gamma_f(|f|_{\infty, Q_T}) + \gamma_d(|d|_{\infty, \partial_t Q_T}), \quad \forall T > 0. \end{aligned}$$

Since $\Sigma(\mathbb{U}, \mathbb{F}, \mathbb{D})$ is 0-UGAS w.r.t. the state in the spatial sup-norm, there exists a function $\beta \in \mathcal{KL}$ such that

$$|v(\cdot, T)|_{\infty, \Omega} \leq \beta(|u^0|_{\infty, \Omega}, T) \quad \forall T > 0.$$

It follows that

$$|u(\cdot, T)|_{\infty, \Omega} \leq |u(\cdot, T) - v(\cdot, T)|_{\infty, \Omega} + |v(\cdot, T)|_{\infty, \Omega} \leq \beta(|u^0|_{\infty, \Omega}, T) + \gamma_f(|f|_{\infty, Q_T}) + \gamma_d(|d|_{\infty, \partial_t Q_T}), \quad \forall T > 0,$$

which implies that $\Sigma(\mathbb{U}, \mathbb{F}, \mathbb{D})$ is ISS in the spatial sup-norm w.r.t. in-domain and boundary disturbances in $\mathbb{F} \times \mathbb{D}$. \blacksquare

Remark 3 In general, for a nonlinear system, it is not an easy task to establish the ISS w.r.t. boundary disturbances directly. Proposition 2.2 provides an alternative for ISS analysis of 0-UGAS systems, which amounts to only assessing the properties of RKES and may be more easily obtained.

2.3 Main stability results

Assume further that

$$(h(x, t, \xi_1) - h(x, t, \xi_2))(\xi_1 - \xi_2) \geq 0 \quad (8)$$

for all $x \in \overline{\Omega}$, $t \in \mathbb{R}_{\geq 0}$, $\xi_1, \xi_2 \in \mathbb{R}$.

The first main result is on the RKES in the sup-norm w.r.t. in-domain and boundary disturbances, whose proof is provided in Appendix.

Theorem 2.1 The following statements hold true.

- (i) Assume that $\underline{c} > 0$ in (1). System (2) with the Robin boundary condition (3) is RLES in the sup-norm w.r.t. in-domain and boundary disturbances in $\mathbb{F} \times \mathbb{D}$, having the estimate:

$$\begin{aligned} \sup_{(x,t) \in Q_T} |u_1(x, t) - u_2(x, t)| &\leq \frac{2C_S^2}{\min\{\underline{a}, \underline{c}_f\}} |\Omega|^{\frac{q-2}{q}} 2^{\frac{3q-4}{2q-4}} \sup_{(x,t) \in Q_T} |f_1(x, t) - f_2(x, t)| \\ &\quad + \frac{1}{\underline{m}} \sup_{(x,t) \in \partial_t Q_T} |d_1(x, t) - d_2(x, t)|, \quad \forall T > 0, \end{aligned} \quad (9)$$

for all $(f_i, d_i) \in \mathbb{F} \times \mathbb{D}$, $i = 1, 2$, where u_i is the solution of the system corresponding to the data $(u^0, f_i, d_i) \in \mathbb{U} \times \mathbb{F} \times \mathbb{D}$, $i = 1, 2$, and q and C_S are constants specified in Lemma A.1(i).

- (ii) Assume that $\underline{c} \geq 0$ in (1). System (2) with the Dirichlet boundary condition (4) is RLES in the sup-norm w.r.t. in-domain and boundary disturbances in $\mathbb{F} \times \mathbb{D}$, having the estimate:

$$\begin{aligned} \sup_{(x,t) \in Q_T} |u_1(x, t) - u_2(x, t)| &\leq \frac{C_P^2}{\underline{a}} |\Omega|^{\frac{q-2}{q}} 2^{\frac{3q-4}{2q-4}} \sup_{(x,t) \in Q_T} |f_1(x, t) - f_2(x, t)| \\ &\quad + \sup_{(x,t) \in \partial_t Q_T} |d_1(x, t) - d_2(x, t)|, \quad \forall T > 0, \end{aligned} \quad (10)$$

for all $(f_i, d_i) \in \mathbb{F} \times \mathbb{D}$, $i = 1, 2$, where u_i is the solution of the system corresponding to the data $(u^0, f_i, d_i) \in \mathbb{U} \times \mathbb{F} \times \mathbb{D}$, $i = 1, 2$, and q and C_P are constants specified in Lemma A.1(ii).

Furthermore, if $\underline{c} > 0$, it holds that

$$\sup_{(x,t) \in Q_T} |u_1(x, t) - u_2(x, t)| \leq C_0 |\Omega|^{\frac{q-2}{q}} 2^{\frac{3q-4}{2q-4}} \sup_{(x,t) \in Q_T} |f_1(x, t) - f_2(x, t)|$$

$$+ \sup_{(x,t) \in \partial_t Q_T} |d_1(x,t) - d_2(x,t)|, \forall T > 0, \quad (11)$$

for all $(f_i, d_i) \in \mathbb{F} \times \mathbb{D}, i = 1, 2$, where u_i is the solution of the system corresponding to the data $(u^0, f_i, d_i) \in \mathbb{U} \times \mathbb{F} \times \mathbb{D}, i = 1, 2, C_0 := \min \left\{ \frac{2C_S^2}{\min\{\underline{a}, \underline{c}\}}, \frac{C_P^2}{\underline{a}} \right\}$, and q, C_S, C_P are constants specified in Lemma A.1.

As an application of RKES, we have the second main result on the ISS in the spatial sup-norm for system (2) with in-domain and boundary disturbances, whose proof is also provided in Appendix.

Theorem 2.2 *The following statements hold true.*

(i) *Assume that $\underline{c} > 0$ in (1). System (2) with the Robin boundary condition (3) is EISS in the spatial sup-norm w.r.t. in-domain and boundary disturbances in $\mathbb{F} \times \mathbb{D}$, having the estimate:*

$$\sup_{x \in \Omega} |u(x, T)| \leq \sup_{x \in \Omega} |u^0(x)| e^{-\underline{c}T} + \frac{2C_S^2}{\min\{\underline{a}, \underline{c}\}} |\Omega|^{\frac{q-2}{q}} 2^{\frac{3q-4}{2q-4}} \sup_{(x,t) \in Q_T} |f(x, t)| + \frac{1}{\underline{m}} \sup_{(x,t) \in \partial_t Q_T} |d(x, t)|, \forall T > 0, \quad (12)$$

where q and C_S are constants specified in Lemma A.1(i).

(ii) *Assume that $\underline{c} > 0$ in (1). System (2) with the Dirichlet boundary condition (4) is EISS in the spatial sup-norm w.r.t. in-domain and boundary disturbances in $\mathbb{F} \times \mathbb{D}$, having the estimate:*

$$\sup_{x \in \Omega} |u(x, T)| \leq \sup_{x \in \Omega} |u^0(x)| e^{-\underline{c}T} + C_0 |\Omega|^{\frac{q-2}{q}} 2^{\frac{3q-4}{2q-4}} \sup_{(x,t) \in Q_T} |f(x, t)| + \sup_{(x,t) \in \partial_t Q_T} |d(x, t)|, \forall T > 0, \quad (13)$$

where the constants q, C_0 are the same as in (11).

Remark 4 *For Theorem 2.2, it is possible to weaken the condition $(u^0, f, d) \in \mathbb{U} \times \mathbb{F} \times \mathbb{D}$ to $(u^0, f, d) \in L^\infty(\Omega) \times L^\infty_{loc}(\mathbb{R}_{\geq 0}; L^\infty(\Omega)) \times L^\infty_{loc}(\mathbb{R}_{\geq 0}; L^\infty(\Omega))$, and hence obtain*

$$\|u(\cdot, T)\|_{L^\infty(\Omega)} \leq \|u^0\|_{L^\infty(\Omega)} e^{-\underline{c}T} + \frac{2C_S^2}{\min\{\underline{a}, \underline{c}\}} |\Omega|^{\frac{q-2}{q}} 2^{\frac{3q-4}{2q-4}} \|f\|_{L^\infty(Q_T)} + \frac{1}{\underline{m}} \|d\|_{L^\infty(Q_T)}, \forall T > 0,$$

and

$$\|u(\cdot, T)\|_{L^\infty(\Omega)} \leq \|u^0\|_{L^\infty(\Omega)} e^{-\underline{c}T} + C_0 |\Omega|^{\frac{q-2}{q}} 2^{\frac{3q-4}{2q-4}} \|f\|_{L^\infty(Q_T)} + \|d\|_{L^\infty(Q_T)}, \forall T > 0,$$

for a weak solution (in a certain sense differing from Definition 1) of the system (2) with the Robin boundary condition (3), and the Dirichlet boundary condition (4), respectively. Indeed, letting $\{u_n^0\}, \{f_n\}, \{d_n\}$ be sequences of sufficiently smooth functions, which satisfy $(u_n^0, f_n, d_n) \rightarrow (u^0, f, d)$ in $L^\infty(\Omega) \times L^\infty(Q_T) \times L^\infty(Q_T)$ as $n \rightarrow +\infty$, we consider the approximating equation (2) with data (u_n^0, f_n, d_n) , and establish uniform a priori estimates of strong (or smooth) solutions $\{u_n\}$ as in [25]. For the existence of a weak solution u , we may prove by using the uniform a priori estimates of $\{u_n\}$ and taking limits in appropriate functional spaces. For the ISS in the spatial L^∞ -norm of u_n , we may prove as in the proof of Theorem 2.2; see Appendix. Then, by taking limits of (u_n, u_n^0, f_n, d_n) within the ISS estimates of $\{u_n\}$, we may obtain the aforementioned ISS estimates of u .

3 Illustrative examples on ISS in the spatial sup-norm of parabolic PEDs

In this section, we illustrate the results on ISS estimates in the spatial sup-norm presented in Section 2 through two examples.

3.1 A super-linear parabolic equation with a Robin boundary condition

Consider the following super-linear parabolic equation:

$$u_t - \Delta u + cu + u \ln(1 + u^2) = f \quad \text{in } \Omega \times \mathbb{R}_+, \quad (14a)$$

$$\frac{\partial u}{\partial \nu} + mu = d \quad \text{on } \partial\Omega \times \mathbb{R}_+, \quad (14b)$$

$$u(\cdot, 0) = u^0(\cdot) \text{ in } \Omega, \quad (14c)$$

where Ω is an open bounded domain in \mathbb{R}^n ($n \geq 3$) with a smooth boundary $\partial\Omega$, Δ is the Laplace operator, c, m are positive constants with $c \geq 1$, $u^0 \in \mathbb{U}$, $d \in \mathbb{D}$, and $f \in \mathbb{F}$. We have the following result.

Proposition 3.1 *Assume that there exists a point $x^0 \in \partial\Omega$ such that $\partial\Omega$ is flat near x^0 and lies in the plane $\{x := (x_1, x_2, \dots, x_n) \in \mathbb{R}^n \mid x_i = 0\}$ for some $i \in \{1, 2, \dots, n\}$, and $f(x, t)$ is Lipschitz continuous in $t \in \mathbb{R}_{\geq 0}$ for all $x \in \bar{\Omega}$. Then, the system (14) is EISS in the spatial sup-norm, having the estimate:*

$$\sup_{x \in \Omega} |u(x, T)| \leq \sup_{x \in \Omega} |u^0(x)| e^{-cT} + \frac{9(n-1)^2}{(n-2)^2} |\Omega|^{\frac{2}{n}} 2^{\frac{n}{4}+8} \sup_{(x,t) \in Q_T} |f(x, t)| + \frac{1}{m} \sup_{(x,t) \in \partial_t Q_T} |d(x, t)|, \quad \forall T > 0.$$

Proof Setting $h(x, t, u) := u \ln(1 + u^2)$, it is easy to verify that h and f satisfy the conditions proposed in Section 2. Then according to Theorem 2.2(i), system (14) is EISS in the spatial sup-norm, having the estimate:

$$\sup_{x \in \Omega} |u(x, T)| \leq \sup_{x \in \Omega} |u^0(x)| e^{-cT} + C_S^2 |\Omega|^{\frac{q-2}{q}} 2^{\frac{5q-8}{2q-4}} \sup_{(x,t) \in Q_T} |f(x, t)| + \frac{1}{m} \sup_{(x,t) \in \partial_t Q_T} |d(x, t)|, \quad \forall T > 0, \quad (15)$$

where q and C_S are constants specified in Lemma A.1(i).

Note that $\partial\Omega$ is flat near x^0 and lies in the plane $\{x \in \mathbb{R}^n \mid x_i = 0\}$, then in the following Sobolev inequality

$$\|v\|_{L^{2^*}(\Omega)} \leq C_1 (\|v\|_{L^2(\Omega)} + \|\nabla v\|_{L^2(\Omega)}), \quad \forall v \in W^{1,2}(\Omega), \quad (16)$$

the constant C_1 can be chosen as $C_1 := \frac{2(n-1)}{n-2} \times (1 + 3 + 4 \times 2) = \frac{24(n-1)}{n-2}$; see [4, Theorem 2, §5.6.1], whose proof is based on Step 2 of the proof of [4, Theorem 1, §5.6.1], and Steps 1-4 of the proof of [4, Theorem 1, §5.4].

For any $q \in (2, 2^*)$ with $2^* := \frac{2n}{n-2}$, it follows from the Hölder's inequality that

$$\|v\|_{L^q(\Omega)}^q \leq \|v\|_{L^{2^*}(\Omega)}^q \|1\|_{L^{\frac{2^*}{2^*-q}}(\Omega)} = \|v\|_{L^{2^*}(\Omega)}^q |\Omega|^{\frac{2^*-q}{2^*}},$$

which along with (16) gives for $v \in W^{1,2}(\Omega)$:

$$\|v\|_{L^q(\Omega)} \leq \frac{24(n-1)}{n-2} |\Omega|^{\frac{2^*-q}{2^*}} (\|v\|_{L^2(\Omega)} + \|\nabla v\|_{L^2(\Omega)}).$$

Thus C_S in Lemma A.1(i) can be chosen as $C_S := \frac{24(n-1)}{n-2} |\Omega|^{\frac{2^*-q}{2^*}}$. Finally, (15) becomes

$$\begin{aligned} \sup_{x \in \Omega} |u(x, T)| &\leq \sup_{x \in \Omega} |u^0(x)| e^{-cT} + \frac{24^2(n-1)^2}{(n-2)^2} |\Omega|^{\frac{q-2}{q} + \left(\frac{2^*-q}{2^*}\right)^2} 2^{\frac{5q-8}{2q-4}} \sup_{(x,t) \in Q_T} |f(x, t)| \\ &\quad + \frac{1}{m} \sup_{(x,t) \in \partial_t Q_T} |d(x, t)|, \quad \forall T > 0. \end{aligned}$$

Letting $q \rightarrow 2^*$ and by algebraic computations, we obtain the desired result. ■

Remark 5 Note that $\frac{24(n-1)}{n-2} |\Omega|^{\frac{2^*-q}{2^*}}$ is not the best embedding constant in the Sobolev inequality given in Lemma A.1(i).

3.2 A 1-D parabolic PDE with a destabilizing term

It is well-known that the following parabolic system

$$u_t(x, t) = u_{xx}(x, t) + cu(x, t), \quad (x, t) \in (0, 1) \times \mathbb{R}_+, \quad (17a)$$

$$u(0, t) = u(1, t) = 0, \quad t \in \mathbb{R}_+, \quad (17b)$$

$$u(x, 0) = u^0(x), \quad x \in (0, 1). \quad (17c)$$

is unstable if the constant c is positive and large. While, the system (17) can be stabilized for any $c \in \mathbb{R}_+$ by a backstepping boundary control $u(1, t) = U(t)$ of the form [14,12]

$$U(t) := - \int_0^1 k(1, y)u(y, t)dy, \quad (18)$$

where k is the solution of the following equation

$$k_{xx}(x, y) - k_{yy}(x, y) = (c + \sigma)k(x, y), \quad 0 \leq y \leq x \leq 1, \quad (19a)$$

$$k(x, 0) = 0, \quad 0 \leq x \leq 1, \quad (19b)$$

$$k_x(x, x) + k_y(x, x) + \frac{d}{dx}(k(x, x)) = c + \sigma, \quad 0 \leq x \leq 1, \quad (19c)$$

with σ being an arbitrary constant. Note that the existence of k is guaranteed by [14, Lemma 2.2]. Moreover, k is twice continuously differentiable in $0 \leq y \leq x \leq 1$.

As an application of Theorem 2.2, we present below an ISS estimate in the spatial norm for the system (17) in closed loop with in-domain and boundary disturbances f, d_0, d_1 , i.e.:

$$u_t(x, t) = u_{xx}(x, t) + cu(x, t) + f(x, t), \quad (x, t) \in (0, 1) \times \mathbb{R}_+, \quad (20a)$$

$$u(0, t) = d_0(t), u(1, t) = d_1(t) + U(t), \quad t \in \mathbb{R}_+, \quad (20b)$$

$$u(x, 0) = u^0(x), \quad x \in (0, 1). \quad (20c)$$

We have the following result.

Proposition 3.2 *Assume that $c \in \mathbb{R}_+$, $f \in C([0, 1])$, $d_0, d_1 \in C(\mathbb{R}_{\geq 0})$, $u^0 \in W^{1,2}(\Omega)$. Assume further that f is Lipschitz continuous w.r.t. its second variable. Let k be the solution of (19) with some constant $\sigma \in \mathbb{R}_+$. Then, under the boundary feedback law (18), the system (20) is EISS, having the estimate:*

$$\begin{aligned} \sup_{x \in (0,1)} |u(x, T)| \leq & (1 + M) \left((1 + M) \sup_{x \in (0,1)} |u^0(x)|e^{-\sigma T} + C \sup_{(x,t) \in (0,1) \times (0,T)} |f(x, t)| \right. \\ & \left. + \sup_{t \in (0,T)} |d_0(t)| + \sup_{t \in (0,T)} |d_1(t)| \right), \forall T > 0, \end{aligned}$$

where $C := \min \left\{ \frac{8\sqrt{2}}{\pi}, \frac{2\sqrt{2}}{\min\{1, c\}} \right\}$ and $M := \sum_{i=0}^{+\infty} \frac{(c+\sigma)^{i+1} 4^i}{(i!)^2} < +\infty$.

Proof Let $w(x, t) := u(x, t) + \int_0^x k(x, y)u(y, t)dy$, which transforms (20) to (see [14, Lemma 2.4]):

$$w_t = w_{xx} - \sigma w + f, \quad (x, t) \in (0, 1) \times \mathbb{R}_+,$$

$$w(0, t) = d_0(t), w(1, t) = d_1(t), \quad t \in \mathbb{R}_+,$$

$$w(x, 0) = u^0(x) + \int_0^x k(x, y)u^0(y)dy := w^0(x), \quad x \in (0, 1).$$

The inverse of the mapping $u \mapsto w$ is given by

$$u(x, t) = w(x, t) + \int_0^x l(x, y)w(y, t)dy, \quad (21)$$

where l is twice continuously differentiable in $0 \leq y \leq x \leq 1$ and satisfies

$$l_{xx}(x, y) - l_{yy}(x, y) = (-\sigma - c)l(x, y), \quad 0 \leq y \leq x \leq 1,$$

$$l(x, 0) = 0, \quad 0 \leq x \leq 1,$$

$$l_x(x, x) + l_y(x, x) + \frac{d}{dx}(l(x, x)) = -\sigma - c, \quad 0 \leq x \leq 1.$$

Note that for $M := \sum_{i=0}^{+\infty} \frac{(c+\sigma)^{i+1} 4^i}{(i!)^2}$, which is convergent, k and l satisfy (see the proof of [14, Lemma 2.2]):

$$\max_{0 \leq y \leq x \leq 1} |k(x, y)| \leq M, \quad \max_{0 \leq y \leq x \leq 1} |l(x, y)| \leq M, \quad \forall T > 0,$$

which along with the definition of w^0 and (21) imply

$$\sup_{x \in (0,1)} |u(x, T)| \leq (1 + M) \sup_{x \in (0,1)} |w(x, T)|, \quad \forall T > 0, \quad (22a)$$

$$\sup_{x \in (0,1)} |w^0(x)| \leq (1 + M) \sup_{x \in (0,1)} |u^0(x)|. \quad (22b)$$

By Theorem 2.2(ii), it follows that

$$\sup_{x \in (0,1)} |w(x, T)| \leq \sup_{x \in (0,1)} |w^0(x)| e^{-\sigma T} + C_0 2^{\frac{3q-4}{2q-4}} \sup_{(x,t) \in (0,1) \times (0,T)} |f(x, t)| + \sup_{t \in (0,T)} |d_0(t)| + \sup_{t \in (0,T)} |d_1(t)|, \quad (23)$$

where q and C_0 are constants specified in Theorem 2.2(ii).

By (22) and (23), we obtain

$$\begin{aligned} \sup_{x \in (0,1)} |u(x, T)| \leq (1 + M) & \left((1 + M) \sup_{x \in (0,1)} |u^0(x)| e^{-\sigma T} + C_0 2^{\frac{3q-4}{2q-4}} \sup_{(x,t) \in (0,1) \times (0,T)} |f(x, t)| \right. \\ & \left. + \sup_{t \in (0,T)} |d_0(t)| + \sup_{t \in (0,T)} |d_1(t)| \right). \end{aligned} \quad (24)$$

On one hand, the Agmon's inequality [12, Lemma 2.4] and a variation of the Wirtinger's inequality [12, Remark 2.2] give

$$\begin{aligned} \|v\|_{L^q(0,1)} & \leq \sup_{x \in (0,1)} |v(x)| \\ & \leq \sqrt{2 \|v\|_{L^2(0,1)} \|v_x\|_{L^2(0,1)}} \\ & \leq \frac{2}{\sqrt{\pi}} \|v_x\|_{L^2(0,1)}, \quad \forall v \in W_0^{1,2}(0,1), \quad \forall q \in (2, +\infty). \end{aligned} \quad (25)$$

Thus, C_P in Lemma A.1(ii) can be chosen as $C_P = \frac{2}{\sqrt{\pi}}$.

On the other hand, it follows from (25) that for any $v \in W_0^{1,2}(0,1)$ and all $q \in (2, +\infty)$:

$$\|v\|_{L^q(0,1)} \leq \sqrt{2 \|v\|_{L^2(0,1)} \|v_x\|_{L^2(0,1)}} \leq \frac{1}{\sqrt{2}} (\|v\|_{L^2(0,1)} + \|v_x\|_{L^2(0,1)}).$$

Thus, C_S in Lemma A.1(ii) can be chosen as $C_S = \frac{1}{\sqrt{2}}$.

Finally, putting $C_0 := \min \left\{ \frac{2C_S^2}{\min\{1, c\}}, C_P^2 \right\} = \min \left\{ \frac{4}{\pi}, \frac{1}{\min\{1, c\}} \right\}$ into (24) and letting $q \rightarrow +\infty$, we obtain the desired result. \blacksquare

Remark 6 It is worth noting that under the boundary feedback law (18) and with additional compatibility conditions on the initial and boundary data, EISS in L^∞ -norm was established for the classical solution of system (20) by using a method combined with transforming, De Girogi iteration, splitting equations, and Lyapunov arguments in [21]. While in this paper, the EISS in the spatial sup-norm is established for the weak solution without any compatibility condition, and only the technique of transforming and De Girogi iteration are used in the proof.

Remark 7 Since it is challenge to apply the backstepping method to 1-D super-linear problems, or to PDEs defined on higher dimensional spatial domains, how to establish the ISS in L^∞ -norm w.r.t. boundary disturbances for a super-linear parabolic PDE having a destabilizing term, or a PDE defined on an arbitrary dimensional domain is still an open problem.

4 Application of RKES to nonlinear parabolic cascade systems

In this section, as an application of the main results presented in Section 2.3, we show how to apply RKES to establish stability estimates in the sup-norm, or the spatial sup-norm, for a class of nonlinear parabolic systems in cascade connected either on the boundary or in the domain. Specifically, for a fixed integer $k \geq 2$ and $j \in \{1, 2, \dots, k\}$, given functions $a_j, c_j, m_j, h_j, \phi_j, f, d$, we consider the following systems coupled on the boundary:

$$(\Sigma_j) \begin{cases} \mathcal{L}_j[u_j] + h_j(x, t, u_j) = 0 & \text{in } \Omega \times \mathbb{R}_+, \\ a_j \frac{\partial u_j}{\partial \nu} + m_j u_j = d_j & \text{on } \partial\Omega \times \mathbb{R}_+, \\ u_j(\cdot, 0) = \phi_j(\cdot) & \text{in } \Omega, \end{cases}$$

where $\mathcal{L}_j[u] := u_t - \operatorname{div}(a_j \nabla u) + c_j u$, and for $(x, t) \in \partial\Omega \times \mathbb{R}_+$:

$$d_1(x, t) := d(x, t), d_j(x, t) := u_{j-1}(x, t), \forall j \in [2, k], \quad (26)$$

or

$$d_1(x, t) := u_k(x, t), d_j(x, t) := u_{j-1}(x, t), \forall j \in [2, k]. \quad (27)$$

We also consider the following equations coupled over the domain:

$$(\Sigma'_j) \begin{cases} \mathcal{L}_j[u_j] + h_j(x, t, u_j) = f_j & \text{in } \Omega \times \mathbb{R}_+, \\ u_j = d_j & \text{on } \partial\Omega \times \mathbb{R}_+, \\ u_j(\cdot, 0) = \phi_j(\cdot) & \text{in } \Omega, \end{cases}$$

where for $(x, t) \in \Omega \times \mathbb{R}_+$:

$$f_1(x, t) := f(x, t), f_j(x, t) := u_{j-1}(x, t), \forall j \in [2, k], \quad (28)$$

or

$$f_1(x, t) := u_k(x, t), f_j(x, t) := u_{j-1}(x, t), \forall j \in [2, k]. \quad (29)$$

In this section, for any $T > 0$, we intend to establish respectively:

- (i) the estimate in the sup-norm, i.e., $\sup_{(x,t) \in Q_T} |u_j(x, t)|$; and
- (ii) the estimate in the spatial sup-norm, i.e., $\sup_{x \in \Omega} |u_j(x, T)|$,

for the considered cascade systems, where u_j is the solution of the j -th subsystem.

For $j \in [1, k]$, we always assume that $a_j \in \mathbb{A}, c_j \in \mathbb{C}, m_j \in \mathbb{M}, h_j \in \mathbb{H}, \phi_j \in \mathbb{U}$. Let $\underline{a}_j, \underline{m}_j$ and \underline{c}_j be defined by (1). Let

$$\Phi_j := \max \left\{ \sup_{x \in \Omega} |\phi_1(x)|, \dots, \sup_{x \in \Omega} |\phi_j(x)| \right\}, m_0 := \min\{\underline{m}_1, \dots, \underline{m}_k\} > 0, a_0 := \frac{1}{|\Omega|^{\frac{q-2}{q}} 2^{\frac{3q-4}{2q-4}} \min\{\tau_1, \dots, \tau_k\}} > 0,$$

where $\tau_j := \begin{cases} \frac{C_P^2}{\underline{a}_j}, & \text{if } \underline{c}_j = 0 \\ \min \left\{ \frac{2C_S^2}{\min\{\underline{a}_j, \underline{c}_j\}}, \frac{C_P^2}{\underline{a}_j} \right\}, & \text{if } \underline{c}_j > 0 \end{cases}$, and the constants q, C_S, C_P are specified in Lemma A.1. We have the

following two propositions.

Proposition 4.1 *For the system (Σ_j) , assume that $d \in \mathbb{D}$. Moreover, assume that the structural conditions (5) and (8) are satisfied with $(h, f) = (h_j, 0)$, and $h = h_j$, respectively. Then the following statements hold true:*

(i) For $j \in [1, k]$, if u_j is the solution of the system (Σ_j) with the Robin boundary condition given by (26), then

$$\sup_{(x,t) \in Q_T} |u_j(x,t)| \leq \frac{m_0}{m_0 - 1} \left(1 - \frac{1}{m_0^j}\right) \Phi_j + \frac{1}{m_0^j} \sup_{(x,t) \in \partial_t Q_T} |d(x,t)|, \forall T > 0. \quad (30)$$

Furthermore, if $\underline{c}_j > 0$, then

$$\sup_{x \in \Omega} |u_j(x,T)| \leq \frac{m_0}{m_0 - 1} \left(1 - \frac{1}{m_0^j}\right) \Phi_j e^{-\underline{c}_j T} + \frac{1}{m_0^j} \sup_{(x,t) \in \partial_t Q_T} |d(x,t)|, \forall T > 0. \quad (31)$$

(ii) Assume further that $m_0 > 1$. For $j \in [1, k]$, if u_j is the solution of the system (Σ_j) with the Robin boundary condition given by (27), then

$$\sup_{(x,t) \in Q_T} |u_j(x,t)| \leq \frac{m_0}{m_0 - 1} \Phi_k, \forall T > 0. \quad (32)$$

Furthermore, if $\underline{c}_j > 0$, then

$$\sup_{x \in \Omega} |u_j(x,T)| \leq \frac{m_0}{m_0 - 1} \Phi_k e^{-\underline{c}_j T}, \forall T > 0. \quad (33)$$

Proposition 4.2 For the system (Σ'_j) , assume that $f \in \mathbb{F}$, $d_j \in \mathbb{D}$. Moreover, assume that the structural condition (5) is satisfied with $(h, f) = (h_j, f)$ for $j = 1$, and $(h, f) = (h_j, 0)$ for $j \in [2, k]$. Meanwhile, assume that the structural condition (8) is satisfied with $h = h_j$ for all $j \in [1, k]$. Then the following statements hold true:

(i) For $j \in [1, k]$, if u_j is the solution of the system (Σ'_j) with (28), then

$$\sup_{(x,t) \in Q_T} |u_j(x,t)| \leq \frac{a_0}{a_0 - 1} \left(1 - \frac{1}{a_0^j}\right) \Phi_j + \frac{1}{a_0^j} \sup_{(x,t) \in Q_T} |f(x,t)| + \sum_{i=1}^j \frac{1}{a_0^{j-i}} \sup_{(x,t) \in \partial_t Q_T} |d_i(x,t)|, \forall T > 0. \quad (34)$$

Furthermore, if $\underline{c}_j > 0$, then

$$\sup_{x \in \Omega} |u_j(x,T)| \leq \frac{a_0}{a_0 - 1} \left(1 - \frac{1}{a_0^j}\right) \Phi_j e^{-\underline{c}_j T} + \frac{1}{a_0^j} \sup_{(x,t) \in Q_T} |f(x,t)| + \sum_{i=1}^j \frac{1}{a_0^{j-i}} \sup_{(x,t) \in \partial_t Q_T} |d_i(x,t)|, \forall T > 0. \quad (35)$$

(ii) Assume further that $a_0 > 1$. For $j \in [1, k]$, if u_j is the solution of the system (Σ'_j) with (29), then

$$\sup_{(x,t) \in Q_T} |u_j(x,t)| \leq \frac{a_0}{a_0 - 1} \Phi_k + \frac{a_0^k}{a_0^k - 1} \sum_{i=1}^k \frac{1}{a_0^{k-i}} \sup_{(x,t) \in \partial_t Q_T} |d_i(x,t)|, \forall T > 0. \quad (36)$$

Furthermore, if $\underline{c}_j > 0$, then

$$\sup_{x \in \Omega} |u_j(x,T)| \leq \frac{a_0}{a_0 - 1} \Phi_k e^{-\underline{c}_j T} + \frac{a_0^k}{a_0^k - 1} \sum_{i=1}^k \frac{1}{a_0^{k-i}} \sup_{(x,t) \in \partial_t Q_T} |d_i(x,t)|, \forall T > 0. \quad (37)$$

Remark 8 It should be mentioned that in [2], the ISS and a small-gain theorem were established for a class of interconnected systems, provided that ISS-Lyapunov functions of the subsystems are known and a small-gain condition holds. As an application of the obtained results, small-gain conditions for guaranteeing the 0-UGAS in the spatial L^2 -norm were proposed for a class of linear, and nonlinear, interconnected parabolic PDEs with homogeneous Dirichlet boundary conditions, respectively. For general interconnected reaction-diffusion systems, it is reasonable to believe that such small-gain conditions depend on the coefficients of the reaction and diffusion terms; see [2] for two special cases. For the interconnected system (Σ'_j) coupled via (27), the small-gain condition is characterized by $a_0 > 1$. While, for the system (Σ_j) coupled on the boundary given in (29),

the small-gain condition is characterized solely by $m_0 > 1$. Moreover, the small-gain conditions proposed in this paper can be used for guaranteeing not only the 0-UGAS, but also the ISS, in the spatial sup-norm, for the considered systems with either Robin or Dirichlet boundary conditions.

Proof of Proposition 4.1 The proof is based on using RKES repeatedly and composed of 4 steps.

Step 1: proof of (30). let v_j be the solution of the following system:

$$\begin{aligned} \mathcal{L}_j[v_j] + h_j(x, t, v_j) &= 0 && \text{in } \Omega \times \mathbb{R}_+, \\ a_j \frac{\partial v}{\partial \nu} + m_j v_j &= 0 && \text{on } \partial\Omega \times \mathbb{R}_+, \\ v_j(\cdot, 0) &= \phi_j(\cdot) && \text{in } \Omega. \end{aligned}$$

The maximum estimate of v_j is given by (see (A.13) in Appendix)

$$\sup_{(x,t) \in Q_T} |v_j(x, t)| \leq \sup_{x \in \Omega} |\phi_j(x)| \leq \Phi_j, \forall T > 0. \quad (38)$$

For $T > 0$, we deduce from Theorem 2.1(i) and (38) that

$$\begin{aligned} \sup_{(x,t) \in Q_T} |u_j(x, t)| &\leq \sup_{(x,t) \in Q_T} |v_j(x, t)| + \sup_{(x,t) \in Q_T} |u_j(x, t) - v_j(x, t)| \\ &\leq \Phi_j + \frac{1}{\underline{m}_j} \sup_{(x,t) \in \partial_t Q_T} |d_j(x, t) - 0| \\ &= \Phi_j + \frac{1}{\underline{m}_j} \sup_{(x,t) \in \partial_t Q_T} |u_{j-1}(x, t)| \\ &\leq \Phi_j + \frac{1}{\underline{m}_j} \sup_{(x,t) \in Q_T} |u_{j-1}(x, t)| \\ &\leq \Phi_j + \frac{1}{\underline{m}_j} \left(\sup_{(x,t) \in Q_T} |v_{j-1}(x, t)| + \sup_{(x,t) \in Q_T} |u_{j-1}(x, t) - v_{j-1}(x, t)| \right) \\ &\leq \Phi_j + \frac{1}{\underline{m}_j} \left(\Phi_j + \frac{1}{\underline{m}_{j-1}} \sup_{(x,t) \in \partial_t Q_T} |d_{j-1}(x, t)| \right) \\ &= \Phi_j \left(1 + \frac{1}{\underline{m}_j} \right) + \frac{1}{\underline{m}_j \cdot \underline{m}_{j-1}} \sup_{(x,t) \in \partial_t Q_T} |u_{j-2}(x, t)| \\ &\leq \Phi_j \left(1 + \frac{1}{\underline{m}_j} \right) + \frac{1}{\underline{m}_j \cdot \underline{m}_{j-1}} \sup_{(x,t) \in Q_T} |u_{j-2}(x, t)| \\ &\leq \dots \\ &\leq \Phi_j \left(1 + \frac{1}{\underline{m}_j} + \frac{1}{\underline{m}_j \cdot \underline{m}_{j-1}} + \dots + \frac{1}{\underline{m}_j \cdot \dots \cdot \underline{m}_{j-3}} \right) + \frac{1}{\underline{m}_j \cdot \dots \cdot \underline{m}_2} \sup_{(x,t) \in Q_T} |u_1(x, t)| \\ &\leq \Phi_j \left(1 + \frac{1}{\underline{m}_j} + \frac{1}{\underline{m}_j \cdot \underline{m}_{j-1}} + \dots + \frac{1}{\underline{m}_j \cdot \dots \cdot \underline{m}_2} \right) + \frac{1}{\underline{m}_j \cdot \dots \cdot \underline{m}_1} \sup_{(x,t) \in \partial_t Q_T} |d(x, t)| \\ &\leq \Phi_j \left(1 + \frac{1}{m_0} + \frac{1}{m_0^2} + \dots + \frac{1}{m_0^{j-1}} \right) + \frac{1}{m_0^j} \sup_{(x,t) \in \partial_t Q_T} |d(x, t)| \\ &= \frac{m_0}{m_0 - 1} \left(1 - \frac{1}{m_0^j} \right) \Phi_j + \frac{1}{m_0^j} \sup_{(x,t) \in \partial_t Q_T} |d(x, t)|, \end{aligned} \quad (39)$$

which gives (30).

Step 2: proof of (31). For any constant $\delta \in (0, \underline{c}_j)$, let $w_j := u_j e^{\delta t}$, $\tilde{c}_j := c_j - \delta$, $\tilde{h}_j(x, t, w_j) := h_j(x, t, w_j e^{\delta t}) e^{\delta t}$, $\tilde{d}_j(x, t) := e^{\delta t} d_j(x, t)$, and $\tilde{\mathcal{L}}_j[w_j] := (w_j)_t - \operatorname{div}(a_j \nabla w_j) + \tilde{c}_j w_j$. By direct computations, we have

$$(\tilde{\Sigma}_j) \begin{cases} \tilde{\mathcal{L}}_j[w_j] + \tilde{h}_j(x, t, w_j) = 0 & \text{in } \Omega \times \mathbb{R}_+, \\ a_j \frac{\partial w_j}{\partial \nu} + m_j w_j = \tilde{d}_j & \text{on } \partial\Omega \times \mathbb{R}_+, \\ w_j(\cdot, 0) = \phi_j(\cdot) & \text{in } \Omega. \end{cases}$$

Note that $\min_{x \in \bar{\Omega}} \tilde{c}_j = \underline{c}_j - \delta > 0$. Then, applying (30) to the system $(\tilde{\Sigma}_j)$, we obtain

$$\sup_{(x,t) \in Q_T} |w_j(x, t)| \leq \frac{m_0}{m_0 - 1} \left(1 - \frac{1}{m_0^j}\right) \Phi_j + \frac{1}{m_0^j} \sup_{(x,t) \in \partial_t Q_T} |\tilde{d}_1(x, t)|, \forall T > 0,$$

which implies

$$\sup_{x \in \Omega} |u_j(x, T)| \leq \frac{m_0}{m_0 - 1} \left(1 - \frac{1}{m_0^j}\right) \Phi_j e^{-\delta T} + \frac{1}{m_0^j} \sup_{(x,t) \in \partial_t Q_T} |d(x, t)|, \forall T > 0.$$

Letting $\delta \rightarrow \underline{c}_j$, we obtain (31).

Step 3: proof of (32). Without loss of generality, we consider the system (Σ_j) with (27) for $j = k$. We deduce from (39), (27), and Theorem 2.1(i) that

$$\begin{aligned} \sup_{(x,t) \in Q_T} |u_k(x, t)| &\leq \Phi_k \left(1 + \frac{1}{m_0} + \cdots + \frac{1}{m_0^{k-2}}\right) + \frac{1}{m_0^{k-1}} \sup_{(x,t) \in Q_T} |u_1(x, t)| \\ &\leq \Phi_k \left(1 + \frac{1}{m_0} + \cdots + \frac{1}{m_0^{k-2}}\right) + \frac{1}{m_0^{k-1}} \left(\Phi_k + \frac{1}{m_0} \sup_{(x,t) \in \partial_t Q_T} |u_k(x, t)|\right) \\ &\leq \frac{m_0}{m_0 - 1} \left(1 - \frac{1}{m_0^k}\right) \Phi_k + \frac{1}{m_0^k} \sup_{(x,t) \in Q_T} |u_k(x, t)|, \end{aligned} \quad (40)$$

which along with $m_0 > 1$ implies (32).

Step 4: proof of (33). Using transformation as in Step 2, considering $(\tilde{\Sigma}_j)$ with the boundary conditions given by (27), and applying (32), we get (33). \blacksquare

Proof of Proposition 4.2 Indeed, for $j \in [1, k]$, let v_j be the solution of the following system:

$$\begin{aligned} \mathcal{L}_j[v_j] + h_j(x, t, v_j) &= 0 & \text{in } \Omega \times \mathbb{R}_+, \\ v_j &= 0 & \text{on } \partial\Omega \times \mathbb{R}_+, \\ v_j(\cdot, 0) &= \phi_j(\cdot) & \text{in } \Omega, \end{aligned}$$

Analogous to (38), the maximum estimate of v_j is given by

$$\sup_{(x,t) \in Q_T} |v_j(x, t)| \leq \sup_{x \in \Omega} |\phi_j(x)|, \forall T > 0. \quad (41)$$

First, we prove (34). For $T > 0$, we deduce from Theorem 2.1(ii) and (41) that

$$\sup_{(x,t) \in Q_T} |u_j(x, t)| \leq \sup_{(x,t) \in Q_T} |v_j(x, t)| + \sup_{(x,t) \in Q_T} |u_j(x, t) - v_j(x, t)|$$

$$\begin{aligned}
&\leq \Phi_j + \frac{1}{a_0} \sup_{(x,t) \in Q_T} |f_j(x,t) - 0| + \sup_{(x,t) \in \partial_t Q_T} |d_j(x,t) - 0| \\
&\leq \Phi_j + \frac{1}{a_0} \sup_{(x,t) \in Q_T} |u_{j-1}(x,t)| + \sup_{(x,t) \in \partial_t Q_T} |d_j(x,t)| \\
&\leq \Phi_j + \frac{1}{a_0} \left(\sup_{(x,t) \in Q_T} |v_{j-1}(x,t)| + \sup_{(x,t) \in Q_T} |u_{j-1}(x,t) - v_{j-1}(x,t)| \right) + \sup_{(x,t) \in \partial_t Q_T} |d_j(x,t)| \\
&\leq \Phi_j + \frac{1}{a_0} \left(\Phi_j + \frac{1}{a_0} \sup_{(x,t) \in Q_T} |f_{j-1}(x,t)| + \sup_{(x,t) \in \partial_t Q_T} |d_{j-1}(x,t)| \right) + \sup_{(x,t) \in \partial_t Q_T} |d_j(x,t)| \\
&\leq \dots \\
&\leq \frac{a_0}{a_0 - 1} \left(1 - \frac{1}{a_0^j} \right) \Phi_j + \frac{1}{a_0^j} \sup_{(x,t) \in Q_T} |f(x,t)| + \sum_{i=1}^j \frac{1}{a_0^{j-i}} \sup_{(x,t) \in \partial_t Q_T} |d_i(x,t)|, \tag{42}
\end{aligned}$$

which gives (34).

Now we prove (36). Without loss of generality, we consider the case of $j = k$. Indeed, analogous to (42) (see also (40)), we have

$$\begin{aligned}
\sup_{(x,t) \in Q_T} |u_k(x,t)| &\leq \frac{a_0}{a_0 - 1} \left(1 - \frac{1}{a_0^k} \right) \Phi_k + \frac{1}{a_0^k} \sup_{(x,t) \in Q_T} |f_1(x,t)| + \sum_{i=1}^k \frac{1}{a_0^{k-i}} \sup_{(x,t) \in \partial_t Q_T} |d_i(x,t)|, \\
&= \frac{a_0}{a_0 - 1} \left(1 - \frac{1}{a_0^k} \right) \Phi_k + \sum_{i=1}^k \frac{1}{a_0^{k-i}} \sup_{(x,t) \in \partial_t Q_T} |d_i(x,t)| + \frac{1}{a_0^k} \sup_{(x,t) \in Q_T} |u_k(x,t)|,
\end{aligned}$$

which along with $a_0 > 1$ gives (36).

Finally, using the technique of transforming as in the proof of Proposition 4.1, we obtain (35) and (37). ■

5 Concluding remarks

This paper proposed a new method for establishing the ISS in the spatial sup-norm for nonlinear parabolic PDEs with boundary and in-domain disturbances. More precisely, we introduced the notion of *RKES* to describe the uniform dependence of solutions on the external disturbances. Based on *RKES* in the (spatial and time) sup-norm, we proved the ISS in the spatial sup-norm for a class of higher dimensional nonlinear PDEs with Dirichlet and Robin boundary disturbances, respectively. Two examples were provided to illustrate the obtained ISS results. In addition, as an application of the introduced notion of *RKES*, we also established stability estimates in the sup-norm and spatial sup-norm for a class of parabolic systems in cascade coupled over the domain and on the boundary of the domain, respectively.

It should be mentioned that the approach presented in this paper is well suited for ISS analysis of weak solutions to higher dimensional nonlinear PDEs with Dirichlet or Robin boundary conditions. However, it seems to be difficult to apply the proposed method to obtain the ISS in the spatial sup-norm for PDEs with Neumann boundary disturbances due to the usage of De Giorgi iteration. Therefore, there is a need to overcome this obstacle and establish ISS estimates in the spatial sup-norm for a wider class of PDEs with various boundary disturbances by a unified approach, which will be considered in our future work.

Besides, as indicated in Remark 7, it is also necessary to develop new methods or tools to address the ISS in the spatial sup-norm for 1-D super-linear parabolic PDEs with destabilizing terms, or parabolic PDEs defined over a higher dimensional domain and having destabilizing terms.

A Proofs of main results

We present some basic Sobolev embedding inequalities that will be used in the proofs of stabilities.

Lemma A.1 (*Theorem 1.3.2 and 1.3.4 of [19]*) *Let Ω be a bounded, open subset of \mathbb{R}^n ($n \geq 1$), and suppose that $\partial\Omega$ is C^1 . For $n = 1, 2$ and $q \in (2, +\infty)$, or $n \geq 3$ and $q \in (2, \frac{2n}{n-2})$, the following inequalities hold true:*

- (i) $\|v\|_{L^q(\Omega)} \leq C_S(\|v\|_{L^2(\Omega)} + \|\nabla v\|_{L^2(\Omega)}), \forall v \in W^{1,2}(\Omega),$
- (ii) $\|v\|_{L^q(\Omega)} \leq C_P\|\nabla v\|_{L^2(\Omega)}, \forall v \in W_0^{1,2}(\Omega),$

where C_S and C_P are positive constants depending only on q, n , and Ω .

Proof of Theorem 2.1 We prove first Theorem 2.1(i). Let u_i be the solution of the system $\Sigma(\mathbb{U}, \mathbb{F}, \mathbb{D})$ corresponding to the data $(u^0, f_i, d_i) \in \mathbb{U} \times \mathbb{F} \times \mathbb{D}, i = 1, 2$.

Consider $w = u_1 - u_2$, which satisfies:

$$\mathcal{L}[w] + h(x, t, u_1) - h(x, t, u_2) = \tilde{f} \text{ in } \Omega \times \mathbb{R}_+, \quad (\text{A.1a})$$

$$a \frac{\partial w}{\partial \nu} + mw = \tilde{d} \text{ on } \partial\Omega \times \mathbb{R}_+, \quad (\text{A.1b})$$

$$w(\cdot, 0) = 0 \text{ in } \Omega, \quad (\text{A.1c})$$

where $\tilde{d} := d_1 - d_2, \tilde{f} := f_1 - f_2$.

We proceed by De Giorgi iteration. Specifically, for any $T > 0$, let $k_0 := \max\left\{0, \frac{1}{m} \sup_{\partial_t Q_T} \tilde{d}\right\}$. For $k \geq k_0$ and $0 < t_1 < t_2 < T$, let $\eta(x, t) := (w(x, t) - k)_+ \chi_{[t_1, t_2]}(t)$, where $s_+ := \max\{s, 0\}$ for $s \in \mathbb{R}$, and $\chi_{[t_1, t_2]}(t)$ is the character function on $[t_1, t_2]$. By virtue of Proposition 2.1, and that $W^{1,p}(\Omega) \hookrightarrow W^{1,2}(\Omega) \hookrightarrow L^2(\Omega) \hookrightarrow (W^{1,2}(\Omega))' \hookrightarrow (W^{1,p}(\Omega))'$ for $p \geq 2$, we have $\eta \in L^\infty((0, T); (W^{1,p}(\Omega))')$ with $\eta_t \in L^\infty((0, T); L^{p'}(\Omega))$. Then, η can be chosen as a test function for (A.1).

By the Fubini's theorem and integrating by parts, we have

$$-\int_0^T \int_\Omega w \eta_t dx dt = -\int_\Omega w(x, T) \eta(x, T) dx + \int_\Omega w(x, 0) \eta(x, 0) dx + \int_0^T \int_\Omega w_t \eta dx dt = \int_0^T \int_\Omega w_t \eta dx dt.$$

It follows that

$$\begin{aligned} & \int_0^T \int_\Omega (w - k)_t (w - k)_+ \chi_{[t_1, t_2]}(t) dx dt - \int_0^T \int_{\partial\Omega} (\tilde{d} - mw)(w - k)_+ \chi_{[t_1, t_2]}(t) dS dt \\ & + \int_0^T \int_\Omega a |\nabla (w - k)_+|^2 \chi_{[t_1, t_2]}(t) dx dt + \int_0^T \int_\Omega cw (w - k)_+ \chi_{[t_1, t_2]}(t) dx dt \\ & + \int_0^T \int_\Omega (h(x, t, u_1) - h(x, t, u_2))(w - k)_+ \chi_{[t_1, t_2]}(t) dx dt \\ & = \int_0^T \int_\Omega \tilde{f} (w - k)_+ \chi_{[t_1, t_2]}(t) dx dt. \end{aligned} \quad (\text{A.2})$$

Note that for $w \geq k \geq k_0 \geq 0$, it follows that $-mw \leq -mk_0 \leq -\underline{m}k_0 \leq -\sup_{\partial_t Q_T} \tilde{d}$, which implies that

$$\int_0^T \int_{\partial\Omega} (\tilde{d} - mw)(w - k)_+ \chi_{[t_1, t_2]}(t) dS dt \leq \int_0^T \int_{\partial\Omega} (\tilde{d} - \sup_{\partial_t Q_T} \tilde{d})(w - k)_+ \chi_{[t_1, t_2]}(t) dS dt \leq 0. \quad (\text{A.3})$$

In addition, for $w \geq k \geq k_0 \geq 0$, it follows that $u_1 = u_2 + w \geq u_2$, which and (8) give

$$\int_0^T \int_\Omega (h(x, t, u_1) - h(x, t, u_2))(w - k)_+ \chi_{[t_1, t_2]}(t) dx dt \geq 0. \quad (\text{A.4})$$

It is obvious that

$$\int_0^T \int_\Omega cw (w - k)_+ \chi_{[t_1, t_2]}(t) dx dt = \int_0^T \int_\Omega c((w - k)_+)^2 \chi_{[t_1, t_2]}(t) dx dt + \int_0^T \int_\Omega kc (w - k)_+ \chi_{[t_1, t_2]}(t) dx ds$$

$$\geq \underline{c} \int_0^T \int_{\Omega} ((w - k)_+)^2 \chi_{[t_1, t_2]}(t) dx dt. \quad (\text{A.5})$$

Then, by (A.2), (A.3), (A.4), and (A.5), we obtain

$$\begin{aligned} & \int_0^T \int_{\Omega} (w - k)_t (w - k)_+ \chi_{[t_1, t_2]}(t) dx dt + \underline{a} \int_0^T \int_{\Omega} |\nabla (w - k)_+|^2 \chi_{[t_1, t_2]}(t) dx dt \\ & + \underline{c} \int_0^T \int_{\Omega} ((w - k)_+)^2 \chi_{[t_1, t_2]}(t) dx dt \\ & \leq \int_0^T \int_{\Omega} \tilde{f}(w - k)_+ \chi_{[t_1, t_2]}(t) dx dt. \end{aligned}$$

Hence

$$\frac{1}{2} \int_{t_1}^{t_2} \frac{d}{dt} \int_{\Omega} ((w - k)_+)^2 dx dt + \underline{a} \int_{t_1}^{t_2} \int_{\Omega} |\nabla (w - k)_+|^2 dx dt + \underline{c} \int_{t_1}^{t_2} \int_{\Omega} ((w - k)_+)^2 dx dt \leq \int_{t_1}^{t_2} \int_{\Omega} \tilde{f}(w - k)_+ dx dt,$$

i.e.,

$$\frac{1}{2} (I_k(t_2) - I_k(t_1)) + \underline{a} \int_{t_1}^{t_2} \int_{\Omega} |\nabla (w - k)_+|^2 dx dt + \underline{c} \int_{t_1}^{t_2} \int_{\Omega} ((w - k)_+)^2 dx dt \leq \int_{t_1}^{t_2} \int_{\Omega} \tilde{f}(w - k)_+ dx dt,$$

where $I_k(t) := \int_{\Omega} ((w(x, t) - k)_+)^2 dx$.

Suppose that $I_k(t_0) = \max_{t \in [0, T]} I_k(t)$ with some $t_0 \in [0, T]$. Due to $I_k(0) = 0$ and $I_k(t) \geq 0$, we can assume that $t_0 \in (0, T]$ without loss of generality.

If $t_0 = T$, then $I'_k(T) \geq 0$. Thus $I'_k(t) \geq 0$ on $(T - \delta, T]$ for some $\delta > 0$. Then, there exists a sufficiently small constant $\varepsilon > 0$ such that $I_k(T - \varepsilon) - I_k(T - 2\varepsilon) \geq 0$. Taking $t_2 = T - \varepsilon$ and $t_1 = T - 2\varepsilon > 0$, we obtain

$$\frac{\underline{a}}{\varepsilon} \int_{T-2\varepsilon}^{T-\varepsilon} \int_{\Omega} |\nabla (w - k)_+|^2 dx dt + \frac{\underline{c}}{\varepsilon} \int_{T-2\varepsilon}^{T-\varepsilon} \int_{\Omega} ((w - k)_+)^2 dx dt \leq \frac{1}{\varepsilon} \int_{T-2\varepsilon}^{T-\varepsilon} \int_{\Omega} |\tilde{f}|(w - k)_+ dx dt.$$

Letting $\varepsilon \rightarrow 0^+$, we get for such $t_0 := T$:

$$\underline{a} \int_{\Omega} |\nabla (w(x, t_0) - k)_+|^2 dx + \underline{c} \int_{\Omega} ((w(x, t_0) - k)_+)^2 dx \leq \int_{\Omega} |f(x, t_0)|(w(x, t_0) - k)_+ dx. \quad (\text{A.6})$$

If $t_0 \in (0, T)$, we can take $t_2 = t_0$ and $t_1 = t_0 - \varepsilon > 0$ for a small $\varepsilon > 0$. Analogously, we can obtain (A.6). Thus, (A.6) holds true whenever $t_0 \in (0, T]$.

Using Lemma A.1(i), we have

$$\|(w(x, t_0) - k)_+\|_{L^q(\Omega)}^2 \leq 2C_S^2 \left(\|(w(x, t_0) - k)_+\|_{L^2(\Omega)}^2 + \|\nabla (w(x, t_0) - k)_+\|_{L^2(\Omega)}^2 \right), \quad (\text{A.7})$$

where q and C_S are the same as in Lemma A.1(i).

Let $A_k(t) := \{x \in \Omega; w(x, t) > k\}$. By (A.6), (A.7), $\underline{a} > 0$, and $\underline{c} > 0$, we have

$$\frac{\min\{\underline{a}, \underline{c}\}}{2C_S^2} \left(\int_{A_k(t_0)} |w(x, t_0) - k|^q dx \right)^{\frac{2}{q}} = \frac{\min\{\underline{a}, \underline{c}\}}{2C_S^2} \|(w(x, t_0) - k)_+\|_{L^q(\Omega)}^2$$

$$\begin{aligned}
&\leq \underline{a} \|\nabla(w(x, t_0) - k)_+\|_{L^2(\Omega)}^2 + \underline{c} \|(w(x, t_0) - k)_+\|_{L^2(\Omega)}^2 \\
&\leq \int_{\Omega} |\tilde{f}(x, t_0)| (w(x, t_0) - k)_+ dx.
\end{aligned} \tag{A.8}$$

By the Hölder's inequality, we have

$$\int_{\Omega} |\tilde{f}(x, t_0)| (w(x, t_0) - k)_+ dx \leq \left(\int_{A_k(t_0)} |\tilde{f}(x, t_0)|^{q'} dx \right)^{\frac{1}{q'}} \left(\int_{A_k(t_0)} |w(x, t_0) - k|^q dx \right)^{\frac{1}{q}},$$

which along with (A.8) gives

$$\begin{aligned}
\left(\int_{A_k(t_0)} |w(x, t_0) - k|^q dx \right)^{\frac{1}{q}} &\leq \frac{2C_S^2}{\min\{\underline{a}, \underline{c}\}} \left(\int_{A_k(t_0)} |\tilde{f}(x, t_0)|^{q'} dx \right)^{\frac{1}{q'}} \\
&\leq \frac{2C_S^2}{\min\{\underline{a}, \underline{c}\}} \|\tilde{f}\|_{L^\infty(Q_T)} |A_k(t_0)|^{\frac{1}{q'}} \\
&\leq \frac{2C_S^2}{\min\{\underline{a}, \underline{c}\}} \|\tilde{f}\|_{L^\infty(Q_T)} \mu_k^{\frac{1}{q'}},
\end{aligned}$$

where $q' := \frac{q}{q-1}$, $\mu_k := \sup_{t \in (0, T)} |A_k(t)|$, and $|A_k(t_0)|$ denotes the n -dimensional Lebesgue measure of $A_k(t_0)$. Then, we may proceed exactly as in the proof of [19, Theorem 4.2.1] to obtain

$$w \leq k_0 + \frac{2C_S^2}{\min\{\underline{a}, \underline{c}\}} |\Omega|^{\frac{q-2}{q}} 2^{\frac{3q-4}{2q-4}} \|\tilde{f}\|_{L^\infty(Q_T)} \quad \text{a.e. in } Q_T,$$

which along with the continuities of w and f yields

$$w \leq \max \left\{ 0, \frac{1}{\underline{m}} \sup_{\partial_t Q_T} \tilde{d} \right\} + \frac{2C_S^2}{\min\{\underline{a}, \underline{c}\}} |\Omega|^{\frac{q-2}{q}} 2^{\frac{3q-4}{2q-4}} \sup_{Q_T} |\tilde{f}| \quad \text{in } Q_T. \tag{A.9}$$

We need to prove the lower boundedness of w . Indeed, it suffices to set $\bar{w} := -w = u_2 - u_1$, and consider the equation

$$\begin{aligned}
\mathcal{L}[\bar{w}] + h(x, t, u_2) - h(x, t, u_1) &= -\tilde{f} \quad \text{in } \Omega \times \mathbb{R}_+, \\
a \frac{\partial \bar{w}}{\partial \nu} + m\bar{w} &= -\tilde{d} \quad \text{on } \partial\Omega \times \mathbb{R}_+, \\
\bar{w}(\cdot, 0) &= 0 \quad \text{in } \Omega.
\end{aligned}$$

Let $\bar{k}_0 = \max \left\{ 0, \frac{1}{\underline{m}} \sup_{\partial_t Q_T} (-\tilde{d}) \right\}$. Proceeding as above, we obtain

$$-w = \bar{w} \leq \bar{k}_0 + \frac{2C_S^2}{\min\{\underline{a}, \underline{c}\}} |\Omega|^{\frac{q-2}{q}} 2^{\frac{3q-4}{2q-4}} \sup_{Q_T} |\tilde{f}| \quad \text{in } Q_T. \tag{A.10}$$

Finally, by (A.9) and (A.10), we have

$$\sup_{Q_T} |w| \leq \frac{1}{\underline{m}} \sup_{\partial_t Q_T} |\tilde{d}| + \frac{2C_S^2}{\min\{\underline{a}, \underline{c}\}} |\Omega|^{\frac{q-2}{q}} 2^{\frac{3q-4}{2q-4}} \sup_{Q_T} |\tilde{f}|,$$

which gives the result stated in Theorem 2.1(i).

Now we prove Theorem 2.1(ii). Consider (A.1) by replacing (A.1b) with

$$w = \tilde{d} \quad \text{on } \partial\Omega \times \mathbb{R}_+.$$

For any $T > 0$, let $k_0 := \max \left\{ 0, \sup_{\partial_t Q_T} \tilde{d} \right\}$. For $k \geq k_0$ and $0 < t_1 < t_2 < T$, let $\eta(x, t) := (w(x, t) - k)_+ \chi_{[t_1, t_2]}(t)$. It suffices to apply De Giorgi iteration as in the proof of Theorem 2.1 (i).

Indeed, if $\underline{c} \geq 0$, (A.6) can be reduced to

$$\underline{a} \int_{\Omega} |\nabla(w(x, t_0) - k)_+|^2 dx \leq \int_{\Omega} |f(x, t_0)|(w(x, t_0) - k)_+ dx.$$

By Lemma A.1(ii), (A.7) can be reduced to

$$\|(w(x, t_0) - k)_+\|_{L^q(\Omega)}^2 \leq C_P^2 \|\nabla(w(x, t_0) - k)_+\|_{L^2(\Omega)}^2,$$

where q and C_P are the same as in Lemma A.1(ii). Hence (A.8) becomes

$$\frac{\underline{a}}{C_P^2} \left(\int_{A_k(t_0)} |w(x, t_0) - k|^q dx \right)^{\frac{2}{q}} \leq \int_{\Omega} |\tilde{f}(x, t_0)|(w(x, t_0) - k)_+ dx. \quad (\text{A.11})$$

Then, analogous to (A.9), we obtain the following estimate:

$$w \leq \max \left\{ 0, \sup_{\partial_t Q_T} \tilde{d} \right\} + \frac{C_P^2}{\underline{a}} |\Omega|^{\frac{q-2}{q}} 2^{\frac{3q-4}{2q-4}} \sup_{Q_T} |\tilde{f}| \text{ in } Q_T.$$

The lower boundedness of w can be estimated in the similar way, and the boundedness of w specified in (10) is guaranteed.

Now for $\underline{c} > 0$, we shall determine an appropriate coefficient of $\left(\int_{A_k(t_0)} |w(x, t_0) - k|^q dx \right)^{\frac{2}{q}}$. Indeed, for $v \in W_0^{1,2}(\Omega)$, we also have $v \in W^{1,2}(\Omega)$. Thus (A.11) and (A.8) hold true at the same time. Then we obtain

$$\left(\int_{A_k(t_0)} |w(x, t_0) - k|^q dx \right)^{\frac{2}{q}} \leq C_0 \int_{\Omega} |\tilde{f}(x, t_0)|(w(x, t_0) - k)_+ dx,$$

where $C_0 := \min \left\{ \frac{2C_S^2}{\min\{\underline{a}, \underline{c}\}}, \frac{C_P^2}{\underline{a}} \right\}$. Finally, (11) is guaranteed. ■

Proof of Theorem 2.2 We only prove Theorem 2.2(i), since the proof of Theorem 2.2(ii) can be proceeded in the same way.

We first prove that the system (2) with the Robin boundary condition (3) is 0-UGAS w.r.t. the state in the spatial sup-norm. Indeed, let v be the solution of the following equation:

$$\mathcal{L}[v] + h(x, t, v) = 0 \quad \text{in } \Omega \times \mathbb{R}_+, \quad (\text{A.12a})$$

$$a \frac{\partial v}{\partial \nu} + mv = 0 \quad \text{on } \partial\Omega \times \mathbb{R}_+, \quad (\text{A.12b})$$

$$v(\cdot, 0) = u^0(\cdot) \text{ in } \Omega. \quad (\text{A.12c})$$

For any constant $\delta \in (0, \underline{c})$, let $w := ve^{\delta t}$, $\tilde{c} := c - \delta$, $\tilde{h}(x, t, w) := h(x, t, we^{\delta t})e^{-\delta t}$, $\tilde{f}(x, t) := \tilde{d}(x, t) := 0$. By direct computations, we have

$$\begin{aligned} w_t - \operatorname{div}(a\nabla w) + \tilde{c}w + \tilde{h}(x, t, w) &= \tilde{f} & \text{in } \Omega \times \mathbb{R}_+, \\ a \frac{\partial w}{\partial \nu} + mw &= \tilde{d} & \text{on } \partial\Omega \times \mathbb{R}_+, \end{aligned}$$

$$w(\cdot, 0) = u^0(\cdot) \text{ in } \Omega.$$

Note that $\min_{x \in \bar{\Omega}} \tilde{c} = \underline{c} - \delta > 0$. Then one may apply De Giorgi iteration as in the proof of Theorem 2.1(i), and obtain

$$\sup_{(x,t) \in Q_T} |w(x,t)| \leq k_0 := \max \left\{ \sup_{x \in \Omega} |u^0(x)|, \sup_{(x,t) \in Q_T} |\tilde{f}(x,t)|, \sup_{(x,t) \in \partial_t Q_T} |\tilde{d}(x,t)| \right\} = \sup_{x \in \Omega} |u^0(x)|, \forall T > 0, \quad (\text{A.13})$$

which along with the continuity of w in $t = T$ implies that

$$\sup_{x \in \Omega} |w(x, T)| \leq \sup_{(x,t) \in Q_T} |w(x,t)| \leq \sup_{x \in \Omega} |u^0(x)|.$$

It follows that

$$\sup_{x \in \Omega} |v(x, T)| \leq e^{-\delta T} \sup_{x \in \Omega} |u^0(x)|.$$

Letting $\delta \rightarrow \underline{c}$, we have

$$\sup_{x \in \Omega} |v(x, T)| \leq e^{-\underline{c}T} \sup_{x \in \Omega} |u^0(x)|. \quad (\text{A.14})$$

Finally, by (9), (A.14), and Proposition 2.2, we conclude that the system (2) with the Robin boundary condition (3) is EISS in the spatial sup-norm w.r.t. in-domain and boundary disturbance (f, d) in $\mathbb{F} \times \mathbb{D}$, having the estimate (12). \blacksquare

References

- [1] H. Amann, “Parabolic evolution equations and nonlinear boundary conditions,” *J. Differential Equations*, vol. 72, pp. 201–269, 1988.
- [2] S. Dashkovskiy and A. Mironchenko, “Input-to-state stability of infinite-dimensional control systems,” *Math. Control Signals Systems*, vol. 25, no. 1, pp. 1–35, 2013.
- [3] —, “Input-to-state stability of nonlinear impulsive systems,” *SIAM J. Control Optim.*, vol. 51, no. 3, pp. 1962–1987, 2013.
- [4] L. C. Evans, *Partial Differential Equations*. American Mathematical Society, Providence, Rhode Island, 2010.
- [5] B. Jacob, A. Mironchenko, J. R. Partington, and F. Wirth, “Non-coercive Lyapunov functions for input-to-state stability of infinite-dimensional systems,” 2019, arXiv:1911.01327.
- [6] B. Jacob, R. Nabiullin, J. R. Partington, and F. L. Schwenninger, “Infinite-dimensional input-to-state stability and Orlicz spaces,” *SIAM J. Control Optim.*, vol. 56, no. 2, pp. 868–889, 2018.
- [7] I. Karafyllis and M. Krstic, “ISS with respect to boundary disturbances for 1-D parabolic PDEs,” *IEEE Trans. Autom. Control*, vol. 61, no. 12, pp. 3712–3724, Dec. 2016.
- [8] —, “ISS in different norms for 1-D parabolic PDEs with boundary disturbances,” *SIAM J. Control Optim.*, vol. 55, no. 3, pp. 1716–1751, 2017.
- [9] —, *Input-to-State Stability for PDEs*. London: Springer-Verlag, 2018.
- [10] —, “ISS estimates in the spatial sup-norm for nonlinear 1-D parabolic PDEs,” 2020, arXiv:2007.15204.
- [11] P. E. Kloeden, “Some remarks on relative stability,” *J. Austral. Math. Soc. Ser. B*, vol. 19, no. 1, pp. 112–115, 1975.
- [12] M. Krstic and A. Smyshlyaev, *Boundary control of PDEs: a course on backstepping designs*. Philadelphia, PA, USA: Society for Industrial and Applied Mathematics, 2008.
- [13] V. Lakshmikantham, “Differential systems and extension of Lyapunov’s method,” *Michigan Math. J.*, vol. 9, no. 4, pp. 311–320, 1962.
- [14] W. Liu, “Boundary feedback stabilization of an unstable heat equation,” *SIAM J. Control Optim.*, vol. 42, no. 3, pp. 1033–1043, 2003.
- [15] A. Mironchenko, I. Karafyllis, and M. Krstic, “Monotonicity methods for input-to-state stability of nonlinear parabolic PDEs with boundary disturbances,” *SIAM J. Control Optim.*, vol. 57, no. 1, pp. 510–532, 2019.
- [16] A. Mironchenko and C. Prieur, “Input-to-state stability of infinite-dimensional systems: recent results and open questions,” *SIAM Rev.*, vol. 62, no. 3, pp. 529–614, 2019.
- [17] A. Mironchenko and F. Wirth, “Characterizations of input-to-state stability for infinite-dimensional systems,” *IEEE Trans. Autom. Control*, vol. 63, no. 6, pp. 1692 – 1707, 2018.
- [18] F. Schwenninger, “Input-to-state stability for parabolic boundary control: linear and semi-linear systems,” 2019, arXiv:1908.08317.
- [19] Z. Wu, J. Yin, and C. Wan, *Elliptic & Parabolic Equations*. Singapore: World Scientific Publishing, 2006.
- [20] J. Zheng and G. Zhu, “Input-to-state stability with respect to boundary disturbances for a class of semi-linear parabolic equations,” *Automatica*, vol. 97, pp. 271–277, 2018.

- [21] —, “A De Giorgi iteration-based approach for the establishment of ISS properties for Burgers’ equation with boundary and in-domain disturbances,” *IEEE Trans. Autom. Control*, vol. 64, no. 8, pp. 3476–3483, Aug. 2019.
- [22] —, “Input-to-state stability for a class of 1-D nonlinear parabolic PDEs with nonlinear boundary conditions,” *SIAM J. Control Optim.*, vol. 58, no. 4, pp. 2567–2587, 2020.
- [23] —, “ISS-like estimates for nonlinear parabolic PDEs with variable coefficients on higher dimensional domains,” *Systems & Control Letters*, vol. 146, no. 104808, 2020, DOI: <https://doi.org/10.1016/j.sysconle.2020.104808>.
- [24] —, “Approximations of Lyapunov functionals for ISS analysis of a class of higher dimensional nonlinear parabolic PDEs,” *Automatica*, vol. 125, 2021, 109414.
- [25] J. Zheng, G. Zhu, and M. Li, “A PDE-based aggregate power tracking control of large populations of heterogeneous TCLs,” 2020, arXiv:2010.10819.