Low-temperature thermodynamics of the two-leg ladder Ising model with trimer rungs: A mystery explained

Taras Hutak,¹ Taras Krokhmalskii,¹ Onofre Rojas,² Sergio Martins de Souza,² and Oleg Derzhko¹

¹Institute for Condensed Matter Physics, National Academy of Sciences of Ukraine, Svientsitskii Street 1, 79011 L'viv, Ukraine
²Departamento de Fisica, Universidade Federal de Lavras, CP 3037, 37200-000, Lavras-MG, Brazil
(Dated: August 11, 2020)

Recently, a surprising low-temperature behavior has been revealed in a two-leg ladder Ising model with trimer rungs (Weiguo Yin, arXiv:2006.08921). Motivated by these findings, we study this model from another perspective and demonstrate that the reported observations are related to a critical phenomenon in the standard Ising chain. We also discuss a related curiosity, namely, the emergence of a power-law behavior characterized by quasicritical exponents.

PACS numbers: 05.70.Fh, 75.10.-b, 75.10.Jm, 75.10.Pq Keywords: Ising model, two-leg ladder, thermodynamics, correlations

Recently, Weiguo Yin considered several two-legladder Ising models with short-range interactions which exhibit intriguing behavior at low temperatures which strongly resembles a finite-temperature phase transition [1, 2]. A simplest model is defined by the Hamiltonian

$$H = -\sum_{i=1}^{N} \left[J \left(\sigma_{i,1} \sigma_{i+1,1} + \sigma_{i,2} \sigma_{i+1,2} \right) + J_1 \left(\sigma_{i,1} \sigma_{i,3} + \sigma_{i,2} \sigma_{i,3} \right) + J_2 \sigma_{i,1} \sigma_{i,2} \right]$$
(1)

which is associated with the lattice shown in Fig. 1, top. Here $\sigma = \pm 1$, N is the number of rungs, the positive/negative sign of the Ising exchange couplings corresponds to the ferromagnetic/antiferromagnetic interaction, and the exchange-coupling scheme is illustrated in Fig. 1, top. (Our J_1 and J_2 correspond to J' and J'' of Ref. [1], respectively; we have changed the notations since in what follows we use the prime notation to denote the first and second derivatives with respect to temperature.) The model (1) is exactly solvable by the transfer-matrix method [1-3]. Weiguo Yin found that for $J_2 < 0$ and small positive values of the parameter $\alpha = (|J_1| - |J_2|)/|J|$ the exact specific heat exhibits a sharp peak at certain (low but finite) temperature, the exact entropy shows a waterfall behavior at this temperature etc [1]. He referred to such a phenomenon as marginal phase transition (MPT) [1, 2] or practically perfect phase transition (PPPT) [2] and provided an extensive mathematical analysis to explain this phenomenon. It is worthwhile noting that similar peculiar low-temperature thermodynamics was observed for other one-dimensional Ising-like models with short-range interactions [4–15].

In what follows, we will explain the astonishing low-temperature behavior of the model (1) using the approach suggested in Ref. [16]. Our present study is not restricted to model (1), but has a broader significance illustrating that the low-temperature peculiarities reported in Refs. [1, 2, 4–15] all are related to the criticality of the standard Ising chain at zero temperature [3].

The first step is to trace out in the partition function of the initial model (1) the spins $\sigma_{i,3}$, i = 1, ..., N at




FIG. 1: (Top) The initial two-leg ladder Ising model with trimer rungs, see Eq. (1). (Bottom) The effective two-leg (railroad) ladder Ising model, see Eq. (2). We denoted the bonds as follows: J (black), J_1 (blue), J_2 (red), and J_{\perp} (magenta).

the middle of the rung, see Fig. 1, top, and to arrive at the effective model – the standard two-leg rail-road Ising ladder with the interaction along the legs -J and the temperature-dependent interaction that couples two legs $-J_{\perp}$, see Fig. 1, bottom (decoration-iteration transformations [17]). The Hamiltonian of the effective model reads:

$$H = -NT \ln C$$

$$-\sum_{i=1}^{N} \left[J \left(\sigma_{i,1} \sigma_{i+1,1} + \sigma_{i,2} \sigma_{i+1,2} \right) + J_{\perp} \sigma_{i,1} \sigma_{i,2} \right],$$

$$C = 2 \sqrt{\cosh \frac{2J_1}{T}}, \ J_{\perp} = J_{\perp}(T) = J_2 + \frac{T}{2} \ln \cosh \frac{2J_1}{T}.(2)$$

This representation was noticed in Ref. [1] but not used for further analysis.

The thermodynamics of the model (2) can be found

by the transfer-matrix method [3]. The Helmholtz free energy (per rung) f reads:

$$f = -T \ln C - T \ln \lambda_1,$$

$$\lambda_1 = 2 \left(\cosh \frac{J_{\perp}}{T} \cosh \frac{2J}{T} + \sqrt{1 + \sinh^2 \frac{J_{\perp}}{T} \cosh^2 \frac{2J}{T}} \right).(3)$$

The peculiarity of the effective ladder model (2) stems from a dependence of the interleg coupling $-J_{\perp}$ on the temperature T. Therefore, the internal energy e, the entropy s, and the specific heat c are given by the formulas:

$$e = -T^{2} \frac{\partial}{\partial T} \frac{f}{T} - T \frac{\partial f}{\partial J_{\perp}} J'_{\perp} = e^{(1)} + e^{(2)},$$

$$s = -\frac{\partial f}{\partial T} - \frac{\partial f}{\partial J_{\perp}} J'_{\perp} = s^{(1)} + s^{(2)},$$

$$c = -T \frac{\partial^{2} f}{\partial T^{2}} - 2T \frac{\partial^{2} f}{\partial T \partial J_{\perp}} J'_{\perp} - T \frac{\partial^{2} f}{\partial J_{\perp}^{2}} (J'_{\perp})^{2} - T \frac{\partial f}{\partial J_{\perp}} J''_{\perp}$$

$$= c^{(1)} + c^{(2)} + c^{(3)} + c^{(4)}, (4)$$

where according to Eq. (3)

$$\frac{\partial f}{\partial J_{\perp}} = -\frac{\sinh\frac{J_{\perp}}{T}\cosh\frac{2J}{T}}{\sqrt{1 + \sinh^{2}\frac{J_{\perp}}{T}\cosh^{2}\frac{2J}{T}}},$$

$$\frac{\partial^{2} f}{\partial T \partial J_{\perp}} = \frac{J_{\perp}\cosh\frac{J_{\perp}}{T}\cosh\frac{2J}{T}\cosh\frac{2J}{T} + 2J\sinh\frac{J_{\perp}}{T}\sinh\frac{2J}{T}}{T^{2}\left(1 + \sinh^{2}\frac{J_{\perp}}{T}\cosh^{2}\frac{2J}{T}\right)^{\frac{3}{2}}},$$

$$\frac{\partial^{2} f}{\partial J_{\perp}^{2}} = -\frac{\cosh\frac{J_{\perp}}{T}\cosh^{2}\frac{2J}{T}}{T\left(1 + \sinh^{2}\frac{J_{\perp}}{T}\cosh^{2}\frac{2J}{T}\right)^{\frac{3}{2}}}, (5)$$

and $J'_{\perp} = \partial J_{\perp}/\partial T$, $J''_{\perp} = \partial^2 J_{\perp}/\partial T^2$. It is worthy noting that $\partial f/\partial J_{\perp} = -\sum_{i=1}^N \langle \sigma_{i,1}\sigma_{i,2}\rangle/N \equiv -C_{12}(0)$, where $C_{12}(0)$ is the on-rung correlation between two outer spins on the legs, see Ref. [1]. Moreover, for the correlation length ξ we have

$$\frac{1}{\xi} = \ln \frac{\cosh \frac{J_{\perp}}{T} \cosh \frac{2J}{T} + \sqrt{1 + \sinh^2 \frac{J_{\perp}}{T} \cosh^2 \frac{2J}{T}}}{\cosh \frac{J_{\perp}}{T} \cosh \frac{2J}{T} - \sqrt{1 + \sinh^2 \frac{J_{\perp}}{T} \cosh^2 \frac{2J}{T}}}.$$
 (6)

Equations (2) - (6) explain the enigmatic low-temperature behavior of the initial model (1). Assume that the effective interleg interaction in Eq. (2) is ferromagnetic for $0 \le T < T_p$, vanishes at $T = T_p$, and becomes antiferromagnetic for $T_p < T$, see Fig. 2 for examples. The equation

$$J_{\perp}(T_p) = 0 \tag{7}$$

has a simple analytical solution for $T_p > 0$ if $J_2 = -|J_2| < 0$ and $(|J_1| - |J_2|)/|J_1| \to +0$. Namely, $T_p = 2(|J_1| - |J_2|)/\ln 2 \approx 2.885(|J_1| - |J_2|)$. This is the MPT temperature reported in Ref. [1]. At $T = T_p$ the legs decouple, see Eq. (7), and the correlation length (6) becomes $1/\xi(T_p) = \ln([\cosh(2J/T_p) + 1]/[\cosh(2J/T_p) - 1])$,




FIG. 2: Dependence of J_{\perp} on T as it follows from Eq. (2) for the set of parameters $J_1=3,3.01,\ldots,3.1$ (from bottom to top in the inset) or $J_1=3,3.25,\ldots,9$ (from bottom to top in the main panel) and $J_2=-3$ in Eq. (1). Green diamonds correspond to T_p . Blue crosses correspond to the approximate analytical formula $T_p=2(|J_1|-|J_2|)/\ln 2$, which is valid if $J_2=-|J_2|<0$ and $(|J_1|-|J_2|)/|J_1|\to+0$.

cf. the correlation length for the standard Ising chain [3]. It is not surprising that $\xi(T_p)$ is extremely large if $T_p/|J|$ is small enough, i.e., the correlation length for the standard Ising chain at the temperature T_p which is low in the scale of |J| is obviously large [in this limit $\xi(T_p) \propto \exp(2|J|/T_p)$]. Hence, if

$$\frac{T_p}{|J|} \ll 1 \tag{8}$$

holds, one observes the traces of the Ising-chain criticality in the behavior of the initial model (1) which were interpreted as very unusual and puzzling phenomena. For above mentioned specific case $J_2 = -|J_2| < 0$ and $(|J_1| - |J_2|)/|J_1| \to +0$, the inequality (8) becomes $(|J_1| - |J_2|)/|J| = \alpha \ll 1$, i.e., corresponds to the strong frustration regime of Ref. [1]. Substituting $J_{\perp}(T_p) = 0$ into Eq. (5) we obtain: $\partial f/\partial J_{\perp} = \partial^2 f/\partial T \partial J_{\perp} = 0$, and $\partial^2 f/\partial J_{\perp}^2 = -[\cosh(2J/T_p)]/T_p$. If Eq. (8) holds, this results in a large value of the specific heat $c(T_p)$, see the third term $c^{(3)}$ in the formula for c in Eq. (4) which is $\propto \exp(2|J|/T_p)(J_{\perp}')^2$.

Let us consider T in the vicinity of T_p when the inequality $\sinh^2(J_\perp/T)\cosh^2(2J/T)\gg 1$ holds. Then, according to Eq. (5), $\partial f/\partial J_\perp\approx \mathrm{sgn}(J_\perp)$ whereas $\partial^2 f/\partial T\partial J_\perp\propto \mathrm{sgn}(J_\perp)/|J_\perp|^2$ and $\partial^2 f/\partial J_\perp^2\propto 1/|J_\perp|^3$. Moreover, according to Eq. (6), $1/\xi\propto |J_\perp|$. Bearing in mind that $J_\perp\propto T-T_p$ around T_p (see Fig. 2), one immediately concludes that while approaching T_p (i) e and e show a jump, see the second terms $e^{(2)}$ and $e^{(2)}$ in the formulas for e and e in Eq. (4), and (ii) e and e show power-law dependences: e constant e in the formula for e in Eq. (4) and Eq. (6). Such quasicritical exponents

 $\alpha=\alpha'=3$ and $\nu=\nu'=1$ were observed for similar one-dimensional Ising-like models in Ref. [12]. It is worthy noting that the exactly calculated specific heat c [Eq. (4)] satisfies the sum rule: $\int_0^\infty \mathrm{d}T c/T = s(\infty) - s(0)$, where $s(\infty)=3\ln 2\approx 2.079$ and s(0)=0 are the entropies at infinite and zero temperatures, respectively. Hence, the higher the peak of c around T_p is, a narrower it should be. Recall also that precisely at T_p both c and ξ are finite, see above.

To illustrate the discussion, we consider a specific example conveniently setting $J_1 = 9$, $J_2 = -3$, and J=100. In this case the frustration parameter $\alpha=0.06$ whereas $T_p \approx 24.937$ yielding $T_p/|J| \approx 0.249$ in Eq. (8) [since $(|J_1| - |J_2|)/|J_1| \approx 0.667$, the analytical formula underestimates the value of T_p]. J_{\perp} as a function of T for this set can be seen in the main panel of Fig. 2. The entropy s [Eq. (4)] exhibits a jump at T_p due to the term $s^{(2)} = -(\partial f/\partial J_{\perp})J'_{\perp}, \ s(T_p + 0) - s(T_p - 0) \approx 0.2,$ see the black curve in Fig. 3, top. The specific heat c [Eq. (4)] exhibits a peak around T_p due to the term $c^{(3)} = -T(\partial^2 f/\partial J_\perp^2)(J_\perp')^2$, $c(T_p) \approx 16.3$, see the black curves in Fig. 3, middle and bottom. The correlation length ξ [Eq. (6)] also exhibits a peak around T_p , $\xi(T_p) \approx 760$, see the blue curves in Fig. 3, middle and bottom. Since $|J'_{\perp}(T_p)| < 1$, we have $c(T_p) < \xi(T_p)$. Finally, Fig. 4 demonstrates quasicritical behavior: There is a finite range of temperatures in the vicinity of T_p on which a clear power-law behavior develops with $\alpha = \alpha' = 3$ for the specific heat (black curves) and $\nu = \nu' = 1$ for the correlation length (blue curves). Moreover, the smaller is $T_p/|J|$ in the left-hand side of the inequality in Eq. (8), the larger the region of quasicriticality is, cf. the top panel $(T_p/|J| \approx 0.249)$ and the bottom panel $(T_p/|J|\approx 0.125)$ in Fig. 4. Recall that the quasicritical behavior fails for temperatures in the immediate vicinity of T_p where both c and ξ are finite, see the values of cand ξ at the smallest values of τ in Fig. 4.

Let us summarize our findings. First of all, we have to emphasize the following. The model at hand is exactly solvable one (as well as other models discussed in Refs. [1, 2, 4–15]) and thus there is no much room for speculations about the solutions: The performed transformations are rigorous and the derived results are exact. However, the final results are somewhat astonishing: For example, the specific heat per cell c may exhibit an extra low-temperature peak of extremely large height, the correlation length ξ at this temperature is unexpectedly large etc, see Figs. 3 and 4. Naturally, these features of the rigorously found quantities call for explanations.

As a rule, the authors sought for explanations at the level of the initial models. In contrast, in Ref. [16] we suggest to examine, instead of several diverse initial models, the effective model which shows up after summing over redundant degrees of freedom. The effective model that has emerged after this rigorous decoration-iteration transformation [17] is the Ising chain with temperature-dependent parameters. In the present study, applying this way of thinking to a typical representative consid-




FIG. 3: Low-temperature properties of the two-leg ladder Ising model with trimer rungs, see Eq. (1) and Fig. 1, top, for the set of parameters $J_1 = 9$, $J_2 = -3$, and J = 100. (Top) Entropy versus temperature in the 1–1000 temperature range. (Middle) Specific heat and correlation length versus temperature in the 1–1000 temperature range. (Bottom) Specific heat and correlation length versus temperature in the vicinity of $T_p \approx 24.937$. The contribution of different terms to the entropy and the specific heat, see Eq. (4), is shown explicitly.




FIG. 4: Quasicritical exponents for the specific heat and the correlation length as evidenced by the dependences $\ln c$ (ascending and descending parts of the peak, black curves) and $\ln \xi$ (ascending and descending parts of the peak, blue curves) on $\ln |\tau|$, $\tau = (T-T^*)/T^*$, T^* stands for the temperature at which c(T) or $\xi(T)$ has the peak. The set of parameters is as follows: $J_1 = 9$, $J_2 = -3$ ($T_p \approx 24.937$) and (top) J = 100 [then $T_c \approx 24.927$, $c(T_c) \approx 16.4$ and $T_\xi \approx 24.929$, $\xi(T_\xi) \approx 7.61 \cdot 10^2$] or (bottom) J = 200 [then $T_c \approx 24.937$, $c(T_c) \approx 4.88 \cdot 10^4$ and $T_\xi \approx 24.937$, $\xi(T_\xi) \approx 2.31 \cdot 10^6$]. Dashed lines correspond to power-law dependences with the exponent 3 (gold) and the exponent 1 (green).

ered in Refs. [1, 2] - the two-leg ladder Ising model

with trimer rungs – we have arrived at the two-leg Ising ladder with temperature-dependent rung couplings $J_{\perp}=J_{\perp}(T),~{\rm Eq.}~(2).~{\rm Again},~{\rm the~analysis~of~the}$ properties of the obtained effective two-leg Ising ladder with temperature-dependent parameters explains the low-temperature thermodynamics of a whole class of the initial models which can be cast into that effective model after eliminating redundant spins.

Although we do not deny the usefulness of studies on the level of the initial model, however, we see several advantages of the study based on the effective model. First, we clearly see a universality of the phenomenon: Many initial models collapse to one effective model and, after all, the low-temperature peculiarities of two-leg-ladder Ising models [1, 2] as well as of other one-dimensional Ising-like models [16] are related to the criticality of the standard Ising chain at zero temperature [3]. From our consideration it looks that frustration is not vitally necessary (see also Ref. [10]); the only demand is to have a suitable $J_{\perp}(T)$. The characteristic temperature T_p is determined by the condition when the effective model reduces to the standard Ising-chain model, see Eq. (7). Temperature-dependent parameters of the effective model result in more complex formulas for thermodynamic quantities, see Eq. (4). If T_p is small in the scale of the effective Ising-chain model, Eq. (8), the large values of the specific heat or the correlation length are evident. Moreover, as it follows from Eqs. (4), (5), (6) and Eq. (2), the temperature dependences around T_p (but not in the immediate vicinity of T_p) follow power laws with quasicritical exponents $\alpha = \alpha' = 3$ and $\nu = \nu' = 1$.

Finally, our work paves the path for the higher-dimensional cases. For example, one can consider a two-dimensional rectangular effective Ising model with the horizontal couplings J and the vertical couplings J_{\perp} given in Eq. (2). Obviously, if condition (7) holds, the two-dimensional system becomes a set of noninteracting Ising chains and it exhibits the behavior discussed above.

The authors gratefully acknowledge helpful correspondence with Weiguo Yin. T. Hutak was supported by the fellowship of the National Academy of Sciences of Ukraine for young scientists. O. Rojas and S. M. de Souza thank CNPq (Brazil) and FAPEMIG (Brazil) for partial financial support.

^[1] Weiguo Yin, Frustration-driven unconventional phase transitions at finite temperature in a one-dimensional ladder Ising model, arXiv:2006.08921v2.

^[2] Weiguo Yin, Finding and classifying an infinite number of cases of the practically perfect phase transition in an Ising model in one dimension, arXiv:2006.15087v1.

^[3] R. J. Baxter, Exactly Solved Models in Statistical Mechanics (Academic Press, 1982).

^[4] L. Gálisová and J. Strečka, Phys. Rev. E 91, 022134 (2015).

^[5] J. Strečka, R. C. Alécio, M. L. Lyra, and O. Rojas, J. Magn. Magn. Mater. 409, 124 (2016).

^[6] J. Torrico, M. Rojas, S. M. de Souza, and O. Rojas, Phys. Lett. A 380, 3655 (2016).

^[7] O. Rojas, J. Strečka, and S. M. de Souza, Solid State Communications 246, 68 (2016).

^[8] S. M. de Souza and O. Rojas, Solid State Communications 269, 131 (2018).

^[9] I. M. Carvalho, J. Torrico, S. M. de Souza, M. Rojas, and O. Rojas, J. Magn. Magn. Mater. 465, 323 (2018).

- [10] O. Rojas, A conjecture on the relationship between critical residual entropy and finite temperature pseudo-transitions of one-dimensional models, arXiv:1810.07817v2.
- [11] I. M. Carvalho, J. Torrico, S. M. de Souza, O. Rojas, and O. Derzhko, Annals of Physics 402, 45 (2019).
- [12] O. Rojas, J. Strečka, M. L. Lyra, and S. M. de Souza, Phys. Rev. E 99, 042117 (2019).
- [13] O. Rojas, J. Strečka, O. Derzhko, and S. M. de Souza, J. Phys.: Condens. Matter 32, 035804 (2020).
- [14] J. Strečka, Acta Physica Polonica A 137, 610 (2020).
- [15] J. Strečka, Pseudo-critical behavior of spin-1/2 Ising di-

- amond and tetrahedral chains, arXiv:2002.06942v1.
- [16] T. Krokhmalskii, T. Hutak, O. Rojas, S. M. de Souza, and O. Derzhko, Towards low-temperature peculiarities of thermodynamic quantities for decorated spin chains, arXiv:1908.06419v1.
- [17] I. Syozi, Prog. Theor. Phys. 6, 341 (1951); M. Fisher, Phys. Rev. 113, 969 (1959); O. Rojas, J. S. Valverde, and S. M. de Souza, Physica A 388, 1419 (2009); J. Strečka, Phys. Lett. A 374, 3718 (2010); O. Rojas and S. M. de Souza, J. Phys. A: Math. Theor. 44, 245001 (2011).