

Darwinian evolution as Brownian motion on the simplex:

A geometric perspective on stochastic replicator dynamics

Tobias Lehmann
University of Leipzig

August 13, 2020

Abstract

We prove that stochastic replicator dynamics can be interpreted as intrinsic Brownian motion on the simplex equipped the Aitchison geometry. As an immediate consequence we derive three approximation results in the spirit of Wong-Zakai approximation, Donsker's invariance principle and a JKO-scheme. Finally, using the Fokker-Planck equation and Wasserstein-contraction estimates, we study the long time behavior of the stochastic replicator equation, as an example of a non-gradient drift diffusion on the Aitchison simplex.

MSC subject classification: 51E26, 62-07, 60J60, 60J70, 91A22, 92D25

Keywords: Brownian motion, Dirichlet distribution, Fokker-Planck equation, invariant measure, Wasserstein contraction, Aitchison geometry, replicator dynamics, random fitness landscape, evolutionary stable strategy

CONTENTS

1	Introduction and outline	2
2	A primer on the Aitchison geometry of the simplex	3
3	The Aitchison diffusion aka Brownian motion on the simplex	7
4	Three approximation results for the Aitchison diffusion	13
5	Drift diffusion on the Aitchison simplex	16
	5.1. Fokker-Planck equation and invariant measures for stochastic replicator dynamics	20
	5.2. Wasserstein contractions for stochastic replicator dynamics	27

1. INTRODUCTION AND OUTLINE

The Aitchison geometry is a Hilbert space structure on the open standard unit simplex and used prominently in compositional data analysis. The aim of this note is to present a seemingly new and interesting connection between Darwinian evolution modeled through stochastic replicator equations on one hand and Brownian motion on the Aitchison simplex on the other hand.

Let us elaborate this objective. The classical way, first introduced in [44], to reformulate Darwin's paradigm of selection in mathematical language is by means of replicator dynamics. Consider a population with n distinct types (e.g. genotypes) and denote by $p_i(t)$ the share of individuals with type i at time t . Also, given a fitness landscape $f = (f^1, \dots, f^n): \mathbb{R}^n \rightarrow \mathbb{R}^n$ we write

$$\bar{f}(p) = p \cdot f(p) = \sum_i^n p_i f^i(p), \quad i = 1, \dots, n$$

for the mean fitness. Now fix some initial datum $p \in \Delta$, where

$$\Delta = \Delta_n := \{p \in (0, 1)^n \mid p_1 + \dots + p_n = 1\}$$

is the open standard unit simplex. Then the replicator equation reads

$$\dot{p}_i(t) = p_i(t) (f^i(p(t)) - \bar{f}(p(t))), \quad p(0) = p \tag{1}$$

and models an evolution of type compositions $p(t) = (p_1(t), \dots, p_n(t)) \in \Delta$ undergoing selection through the fitness landscape f .

Especially well studied is the situation of linear fitness landscapes, i.e. when $f(p) = Ap$ for some payoff matrix $A \in \mathbb{R}^{n \times n}$, where profound connections to evolutionary game theory and Smith's concept of evolutionary stable strategies arise [c.f. 24]. But also the simple scenario in which the fitness landscape is frequency independent and thus given by a vector $f = (f^1, \dots, f^n) \in \mathbb{R}^n$ is of interest. In this situation (1) describes the prebiotic evolution of self-replicating polynucleotides (e.g RNA, DNA) without mutations [c.f. 33, 41] and is a particular example of a class of dynamical systems introduced by Eigen and Schuster in their theories of quasispecies and hypercycles [13, 14].

Often, allowing in mathematical models for uncertainty or randomness leads to a description better fitting empirical evidence. Applying this principle to a replicator equation with linear fitness landscape given by the matrix A leads to the stochastic replicator equation

$$dX_t = b(X_t)dt + \sigma(X_t)dB_t, \tag{2}$$

where B is an n -dimensional Brownian motion, the drift component is given by

$$b(x) = (\text{diag}(x_1, \dots, x_n) - x \otimes x) (A - \text{diag}(\sigma_1^2, \dots, \sigma_n^2)) x$$

and the diffusion matrix obeys

$$\sigma(x) = (\text{diag}(x_1, \dots, x_n) - x \otimes x) \text{diag}(\sigma_1, \dots, \sigma_n).$$

Initially proposed in [20], this model and in particular its long-time behavior attracted a lot of interest over the last decades. We refer exemplarily to [23] and references therein.

Although we will not take this perspective here, we want emphasize that apart from the biological application, there is also an interpretation for (2) in terms of mathematical finance. Indeed, using the language of Fernholz' stochastic portfolio theory [c.f. 18], X can be seen as the evolution of a market portfolio, for which the rates of return r_i of the underlying stock prices experience feedback through X via $r_i(t) = (AX_t)_i$.

In the second section of this note we recall basic principles of the Aitchison geometry. In short, we will see that, when equipped with appropriate vector space operations, the simplex can be given a Hilbert space structure by dint of the inner product

$$\langle p, q \rangle_A := \frac{1}{2n} \sum_{i,j=1}^n \ln \frac{p_i}{p_j} \ln \frac{q_i}{q_j}.$$

Now consider (2) with $A = 0$ and $\sigma_i = 1$ for $i = 1, \dots, n$. The corresponding diffusion process, that we refer to as the *Aitchison diffusion*, evolves according to the Stratonovich SDE

$$\begin{aligned} dX_t^i &= X_t^i \left(\circ dB_t^i - \sum_{j=1}^n X_t^j \circ dB_t^j \right), \quad i = 1, \dots, n \\ X_0 &= p \in \Delta, \end{aligned} \tag{3}$$

which can be interpreted as a replicator equation in a white noise fitness landscape. Then the main result of this note, which we present in Section 3, asserts that the Aitchison diffusion is nothing but Brownian motion on $(\Delta, \langle \cdot, \cdot \rangle_A)$.

This observation directly entails three approximation results to be presented in Section 4. Apart from a Wong-Zakai approximation and a JKO-scheme for the associated heat equation on the simplex, we show that discrete stochastic replicator dynamics are random walks on the Aitchison simplex which, in the spirit of Donsker's invariance principle, can be used to approximate (3).

In the last section we study the stochastic replicator dynamic as an example of a drift diffusion processes on Δ . Using the associated Fokker-Planck equation, we investigate the long-time behavior and recover results proven earlier in [23]. The final subsection is devoted to questions of the relaxation to equilibrium for replicator dynamics which we analyse by means of Wasserstein contraction estimates. As a major result we characterize those payoff matrices that steer trajectories of the stochastic replicator equation to synchronize with a Langevin dynamic on Δ driven by cross-entropy.

2. A PRIMER ON THE AITCHISON GEOMETRY OF THE SIMPLEX

Often, most notably in geology and chemistry but also in ecology or social sciences, one is confronted with data which represents portions of a total. We may think of the chemical composition of 100 soil samples taken at different places in Germany. Many classical statistical methods are relying on Euclidean geometry and are therefore inappropriate for analysing data constrained to a constant total sum. Traditionally, such data is called *compositional data* (CoDa) and the corresponding branch of statistics *compositional data analysis*, for which a plenty of literature is available [c.f. 1, 2, 36, 45]. In the following we will mainly rely on [36].

One of the most influential developments in the history of compositional data analysis was Aitchison's idea [3] of equipping the simplex with a Hilbert space structure, which in his honour is nowadays called *Aitchison geometry* and defined as follows:

For any $p, q \in \mathbb{R}_{>0}$ and $\alpha \in \mathbb{R}$ we define the operations:

the *closure* of p

$$\mathcal{C}(p) := \left(\sum_{j=1}^n p_j \right)^{-1} p,$$

the *perturbation* of p by q

$$p \oplus q := \mathcal{C}(p_1 q_1, \dots, p_n q_n),$$

the *powering* of p by α

$$\alpha \odot p := \mathcal{C}(p_1^\alpha, \dots, p_n^\alpha).$$

Then, one can easily check that (Δ, \oplus, \odot) is a vector space, in which the neutral element e is the barycenter of Δ , i.e.

$$e = \mathcal{C}(1, \dots, 1)$$

and where the inverse of $p \in \Delta$ is given by

$$\text{inv}(p) := p^{-1} := -1 \odot p = \mathcal{C}(p_1^{-1}, \dots, p_n^{-1}).$$

Moreover, if we introduce the the Aitchison inner product

$$\langle p, q \rangle_A := \frac{1}{2n} \sum_{i,j=1}^n \ln \frac{p_i}{p_j} \ln \frac{q_i}{q_j} \quad (4)$$

then one can show

Theorem 1 (Hilbert space structure of Δ , [c.f. 8]). *The vector space (Δ, \oplus, \odot) equipped with the inner product (4) is a Hilbert space.*

Theorem 1 is easily justify by providing a Hilbert space isomorphism. The most prominent example, which is of major importance in both, compositional data analysis as well as throughout this note, is the so called *centered log-ratio transform* which maps Δ to the Hilbert space $(H, (\cdot, \cdot)_H)$, where

$$H = H_n := \{x \in \mathbb{R}^n \mid x_1 + \dots + x_n = 0\} \quad (5)$$

and $(\cdot, \cdot)_H$ is the scalar product inherited from the standard inner product on \mathbb{R}^n . This transform is defined as

$$\text{clr}: \Delta \rightarrow H, \quad p \mapsto \text{clr}(p) := \left(\ln \frac{p_i}{g(p)} \right)_{i=1}^n,$$

where $g(p) = \left(\prod_{i=1}^n p_i \right)^{\frac{1}{n}}$ is the geometric mean of p . As a consequence we obtain the usual transformation rules

$$(i) \quad \text{clr}(\alpha \odot p \oplus q) = \alpha \text{clr}(p) + \text{clr}(q) \quad \text{and} \quad (ii) \quad \langle p, q \rangle_A = (\text{clr}(p), \text{clr}(q))_H. \quad (6)$$

For $p, q \in \Delta$ we will abbreviate $p \oplus (-1) \odot q =: p \ominus q$. Then, the Aitchison distance on Δ induced through $\langle \cdot, \cdot \rangle_A$ obeys

$$d_A^2(p, q) := \langle p \ominus q, p \ominus q \rangle = \frac{1}{2n} \sum_{i,j=1}^n \left(\ln \frac{p_i}{p_j} - \ln \frac{q_i}{q_j} \right)^2. \quad (7)$$

Because it will be of frequent use later on, we will denote the inverse log-ratio transform

$$\text{clr}^{-1}: H \rightarrow \Delta, \quad x \mapsto \text{clr}^{-1}(x) = \mathcal{C}(e^{x_1}, \dots, e^{x_n}),$$

by *sfm*. The reason for doing so, is that

$$\text{sfm}_i(x) := \text{clr}_i^{-1}(x) = \frac{e^{x_i}}{\sum_j e^{x_j}}, \quad i = 1, \dots, n$$

is an ubiquitous object in applied mathematics. Whereas it (or close variants of it) occurs as Boltzmann or Gibbs distribution in statistical mechanics, it is a well known map also in evolutionary game theory and decision theory. Over the last decades it has been used most prominently in machine learning, where it is referred to as softmax function [c.f. 21, and references therein], which is the motivation for our naming.

Remark. We defined $\text{clr}: \Delta \rightarrow H$ and $\text{sfm}: H \rightarrow \Delta$ for the sake of bijectivity, but of course a priori clr and sfm are well defined also for arguments in $\mathbb{R}_{>0}^n$ and \mathbb{R}^n , respectively.

Although the clr transform is easy to compute, it has the drawback of mapping to H , whereas often one would rather prefer an isomorphism realising $\Delta \cong \mathbb{R}^{n-1}$. This can be easily achieved by appropriately post- or pre processing clr and sfm , respectively. To this end, let $\{e_1, \dots, e_{n-1}\}$ be an orthonormal base of Δ and define the *contrast matrix* $\Psi \in \mathbb{R}^{(n-1) \times n}$ by

$$\Psi := \begin{pmatrix} \Psi_1 \\ \vdots \\ \Psi_{n-1} \end{pmatrix} := \begin{pmatrix} \text{clr } e^1 \\ \vdots \\ \text{clr } e^{n-1} \end{pmatrix}.$$

Observe that, independent of the choice of the basis, contrast matrices satisfy

$$\Psi \Psi^\top = \text{Id}_{n-1} \quad \text{and} \quad \Psi^\top \Psi = \text{Id}_n - \frac{1}{n} \mathbf{1}_n \otimes \mathbf{1}_n, \quad (8)$$

where $\mathbf{1}_n = (1, \dots, 1) \in \mathbb{R}^n$. Then the *isometric log-ratio transform* ilr is defined by

$$\text{ilr}(p) := \Psi \text{clr}(p)$$

with inverse

$$\text{ilr}^{-1}(x) = \text{sfm}(\Psi^\top x).$$

Equivalently, we can express ilr by

$$\text{ilr}(p) = (\langle p, e^1 \rangle_A, \dots, \langle p, e^{n-1} \rangle_A)^\top,$$

from which we can immediately deduce the desirable property

$$\text{ilr}(e_i) = \epsilon_i,$$

where $\{\epsilon_i\}_{i=1}^{n-1}$ is the standard basis in \mathbb{R}^{n-1} . Henceforth, we will use the symbols e_i and ϵ_i generically for basis elements in Δ and \mathbb{R}^n , respectively.

We continue with a short discussion on the relations between the Euclidean distance $d_{\mathbb{R}^n}$ and the Aitchison distance d_A on Δ and the topologies each of the two the distances induces on the simplex.

Lemma 2. *The metrics $d_{\mathbb{R}^n}$ and d_A induce the same topology on Δ .*

Proof. We need to prove that $\text{id}: \Delta \rightarrow \Delta$ is both, $(d_{\mathbb{R}^n}, d_A)$ -continuous and $(d_A, d_{\mathbb{R}^n})$ -continuous. The first statement follows immediately from the continuity of clr or ilr , respectively. As for the latter case, observe that it is well known [21, prop. 4] that $\text{sfm}: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is 1-Lipschitz continuous. Hence, for all $p, q \in \Delta$ we have

$$\|p - q\|_{\mathbb{R}^n} \leq \|\text{clr}(p) - \text{clr}(q)\|_A = d_A(p, q), \quad (9)$$

which yields the second claim. \square

Notice that, whereas d_A and $d_{\mathbb{R}^n}$ are topologically equivalent, they are not strongly equivalent. Indeed, since $d_{\mathbb{R}^n}$ is uniformly bounded on Δ by $\sqrt{2}$ we cannot find some $c > 0$ with

$$d_A(p, q) \leq c d_{\mathbb{R}^n}(p, q), \quad p, q \in \Delta.$$

Finally, let us make a few words on integration and differentiation on Δ . Since the Aitchison simplex is in particular an Abelian group it comes along with a natural reference measure λ_A , which is the Haar measure on (Δ, \oplus) . In the CoDa community it is referred to as the *Aitchison measure* [c.f. 37], which can be characterized as push-forward of the Lebesgue measure λ_{n-1} on \mathbb{R}^{n-1} under ilr^{-1} or up to multiplicative constants equivalently, as the push forward of λ_n under sfm . The joint distribution of the first $(n-1)$ marginals of λ_A (which by slight abuse of notation we call also λ_A) is absolutely continuous with respect to λ_{n-1} with Radon-Nikodym derivative

$$\frac{d\lambda_A}{d\lambda_{n-1}}(p_1, \dots, p_{n-1}) = \left(\prod_{i=1}^n p_i \right)^{-1},$$

where $p_n := 1 - \sum_{i=1}^{n-1} p_i$.

As a Hilbert space, the Aitchison simplex also exhibits a natural differential calculus, which, as we will see, is closely related to the classical Fisher information geometry of the simplex (see also [15]). Let us briefly recall the latter.

Set $|x| = \sum x_i$ and consider the positive orthant $\mathbb{R}_{>0}^n = \{x \in \mathbb{R}^n \mid x_i > 0\}$ equipped with the Riemannian metric

$$g_x(u, v) = \sum_{i=1}^n |x| \frac{u_i v_i}{x_i}.$$

As a submanifold of $(\mathbb{R}_{>0}^n, g)$, the simplex Δ inherits a Riemannian structure with inverse metric tensor

$$(g^{-1}(p))_{ij} := g^{ij}(p) := p_i(\delta_{ij} - p_j), \quad p \in \Delta \quad (10)$$

and for sufficiently smooth functions $\phi: \Delta \rightarrow \mathbb{R}$ the gradient at $p \in \Delta$ is given by

$$\nabla^g \phi(p) := g^{-1}(p) \nabla \phi(p) = \sum_{i=1}^n p_i \left(\partial_i \phi(p) - \sum_{j=1}^n p_j \partial_j \phi(p) \right) \epsilon_i$$

which is typically referred to as Fisher -or Shahshahani gradient in information geometry and mathematical biology, respectively [c.f. 5, 22, 25, 39].

Now denote by $d^A f$ the Fréchet derivative on the Aitchison simplex determined as usual via

$$\lim_{q \rightarrow e} \|q\|_A^{-1} |f(p \oplus q) - f(p) - d^A f(p)q| = 0, \quad (11)$$

provided the limit exists (recall that in the previous formula e is the neutral element of the Aitchison simplex $e = n^{-1}\mathbf{1}$). Given $f \in C^1(\Delta)$ and $p \in \Delta$, by Riesz representation theorem, we can introduce the *Aitchison gradient* $\nabla^A f(p)$ as the unique element in Δ obeying

$$d^A f(p)q = \langle \nabla^A f(p), q \rangle_A \quad \text{for all } q \in \Delta,$$

and we claim

Lemma 3. *Let $f \in C^1(\Delta)$. Then*

$$\nabla^A f = \text{sfm}(\nabla^g f).$$

Proof. We set $\bar{f} = f \circ \text{sfm}$ and rewrite $f(p \oplus q) = \bar{f}(\text{clr}(p) + \text{clr}(q))$. Now a Taylor expansion around $\text{clr}(p)$ yields

$$f(p \oplus q) = f(p) + (\nabla \bar{f}(\text{clr}(p)), \text{clr}(q)) + o(\|q\|_A^2).$$

But since $\|q\|_A^{-1} o(\|q\|_A^2) \rightarrow 0$ as $q \rightarrow e$, plugging the previous expansion into (11) necessitates

$$d^A f(p)q = (\nabla \bar{f}(\text{clr}(p)), \text{clr}(q)).$$

Next, observe that the Jacobian of sfm has the remarkable form

$$D \text{sfm} = \text{diag}(\text{sfm}) - \text{sfm} \otimes \text{sfm}, \quad (12)$$

so that by the chain rule

$$(\nabla \bar{f}) \circ \text{clr}(p) = \nabla^g f(p).$$

Hence,

$$(\nabla \bar{f}(\text{clr}(p)), \text{clr}(q)) = (\nabla^g f(p), \text{clr}(q))_H = \langle \text{sfm}(\nabla^g f(p)), q \rangle_A,$$

which yields the claim. \square

We remark that similar computations have been done before in [7]. Due to a different definition for the derivative their gradient differs from ours and simply equals ∇^g .

3. THE AITCHISON DIFFUSION AKA BROWNIAN MOTION ON THE SIMPLEX

Recall that, given a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and a topological Abelian group $(G, *)$, a G -valued stochastic process is called *Brownian motion on G* provided

A1. For $0 \leq t_1 \leq t_2 \leq \dots \leq t_m \leq T$ and every $m \in \mathbb{N}$, the increments

$X_{t_1}, X_{t_2} * X_{t_1}^{-1}, \dots, X_{t_m} * X_{t_{m-1}}^{-1}$ are mutually independent.

A2. For any $0 \leq s \leq t \leq T$ the law of the increments $X_t * X_s^{-1}$ depends only on $t - s$.

A3. $t \mapsto X_t$ is continuous a.s.

The main result of this section establishes a deep connection between stochastic replicator dynamics and Brownian motion:

Theorem 4 (Aitchison diffusion as Brownian motion on Δ). *For every $T > 0$ there exists a unique Δ -valued process X solving on $[0, T]$ the Stratonovich SDE*

$$\begin{aligned} dX_t^i &= X_t^i \left(\circ dB_t^i - \sum_{j=1}^n X_t^j \circ dB_t^j \right), \quad i = 1, \dots, n \\ X_0 &= p \in \Delta \end{aligned} \tag{3}$$

Moreover, X satisfies the properties A1.-A3. with $G = (\Delta, \oplus)$ the Aitchison simplex. Thus, X is a Brownian motion on Δ .

We approach Theorem 4 by some preliminary considerations. First observe that since (Δ, \oplus, \odot) is a finite dimensional vector space, there is a canonical way to introduce Brownian motion on the Aitchison simplex. Namely, we simply take an orthonormal basis $\{e_1, \dots, e_{n-1}\}$ of Δ and $n - 1$ independent (one-dimensional) standard Brownian motions B^i , $i = 1, \dots, n - 1$ and define for all $t \geq 0$

$$\hat{X}_t := \bigoplus_{i=1}^{n-1} B_t^i \odot e_i. \tag{13}$$

Then clearly \hat{X} satisfies A1.-A3. Next, we would like to find a characterisation of \hat{X} in conventional Euclidean coordinates. Of course, (13) means nothing but $\hat{X} = \text{ilr}^{-1}(B)$. Thus, by the Stratonovich chain rule we find that \hat{X} satisfies

$$\begin{aligned} d\hat{X}_t^i &= d \text{sfm}_i(\Psi^\top B_t) = \sum_{j=1}^{n-1} \sum_{k=1}^n \Psi_{jk} (\partial_k \text{sfm}_i)(\Psi^\top B_t) \circ dB_t^j \\ &= \sum_{j=1}^{n-1} \sum_{k=1}^n \Psi_{jk} \text{sfm}_i(\Psi^\top B_t) (\delta_{ik} - \text{sfm}_k(\Psi^\top B_t)) \circ dB_t^j \\ &= \sum_{j=1}^{n-1} \sum_{k=1}^n \Psi_{jk} g^{ik}(\hat{X}_t) \circ dB_t^j, \end{aligned}$$

with g^{-1} defined as in (10). If we introduce for $i = 1, \dots, n-1$ the maps $\hat{Z}_i: \Delta \rightarrow \mathbb{R}^n$ with

$$\hat{Z}_i(p) := \sum_{j,k=1}^n \Psi_{ik} g^{jk}(p) \epsilon_j, \quad (14)$$

then the Brownian motion on Δ defined as in (13) satisfies

$$d\hat{X}_t = \sum_{i=1}^{n-1} \hat{Z}_i(\hat{X}_t) \circ dB_t^i.$$

Therefore, the generator \hat{L} of \hat{X} is canonically given in Hörmander form as

$$\hat{L} = \frac{1}{2} \sum_{i=1}^{n-1} \hat{Z}_i^2,$$

where, by slight abuse of notation, we identified with the maps in (14) the vector fields on Δ given by

$$\hat{Z}_i(p) := \sum_{j,k=1}^n \Psi_{ik} g^{jk}(p) \partial_j.$$

Now setting

$$g_l^{ik}(p) := \partial_l g^{ik}(p) = \delta_{il} \delta_{ik} - p_k \delta_{il} - p_i \delta_{lk},$$

and

$$G^{ij}(p) := (G^{-1}(p))_{ij} := \left(g^{-1}(p)^2 \right)_{ij} = p_i \left(p_i \delta_{ij} - p_i p_j - p_j^2 + p_j \sum_{k=1}^n p_k^2 \right)$$

we can expand \hat{L} and obtain, using (8) and the the fact that the rows of g^{-1} sum to zero,

$$\begin{aligned} 2\hat{L} &= \sum_{j,k,l,m=1}^n (\Psi^\top \Psi)_{km} (g^{jk} g^{lm} \partial_{jl} + g^{jk} g_j^{lm} \partial_l) \\ &= \sum_{j,l=1}^n G^{jl} \partial_{jl} + \sum_{l=1}^n \left(\sum_{j,m=1}^n g^{jm} g_j^{lm} \right) \partial_l. \end{aligned} \quad (15)$$

From this observation we can easily infer the

Proof of Theorem 4. Although existence and uniqueness of solutions to (3) is well known, since (3) is just a special case of the general stochastic replicator equation (2), we briefly sketch the argument for the sake of completeness. Recall, that the Aitchison diffusion (3) in Itô form obeys

$$dX_t = b(X_t)dt + g^{-1}(X_t)dB_t, \quad (16)$$

with Stratonovich corrector

$$b_i(p) = \frac{1}{2} \sum_{j,k=1}^n g^{jk}(p) g_j^{ik}(p) = p_i \left(\sum_{k=1}^n p_k^2 - p_i \right) = - (g^{-1}(p)p)_i, \quad i = 1, \dots, n. \quad (17)$$

Observe that both maps, $g^{-1}: \Delta \rightarrow \mathbb{R}^{n \times n}$ and $b: \Delta \rightarrow \mathbb{R}^n$ are Lipschitz-continuous (w.r.t the standard topology on Δ). Indeed, for every $p, q \in \Delta$ we have,

$$\begin{aligned} \|g^{-1}(p) - g^{-1}(q)\|^2 &= \sum_{i,j=1}^n (g^{ij}(p) - g^{ij}(q))^2 = \sum_{i,j=1}^n ((p_i - q_i)\delta_{ij} + q_i q_j - p_i p_j)^2 \\ &\leq 2\|p - q\|^2 + 2 \sum_{i,j=1}^n (q_i q_j - q_i p_j + q_i p_j - p_i p_j)^2 \leq 10\|p - q\|^2. \end{aligned}$$

Regarding the drift component, one finds

$$\begin{aligned} \|b(p) - b(q)\|^2 &\leq 2 \sum_{i=1}^n (q_i^2 - p_i^2)^2 + 2 \sum_{i=1}^n \left(p_i \sum_{k=1}^n p_k^2 - q_i \sum_{k=1}^n q_k^2 \right)^2 \\ &\leq 8\|p - q\|^2 + 4 \sum_{i=1}^n p_i^2 \left(\sum_{k=1}^n p_k^2 - q_k^2 \right)^2 + 4 \sum_{i=1}^n (p_i - q_i)^2 \left(\sum_{k=1}^n q_k^2 \right)^2 \\ &\leq (12 + 16n)\|p - q\|^2. \end{aligned}$$

Since moreover, $\text{im } g^{-1}(p) = T_p \Delta = H$ (with H as defined in (5)) holds for every $p \in \Delta$, existence and uniqueness of a continuous and Δ -valued solution to (3) follow by standard Picard iteration. In particular such a solution satisfies A3.

We are left to check that X satisfies the properties A1. and A2. But comparing the Stratonovich corrector (17) with the first order part of \hat{L} in (15), we realise that \hat{L} and the generator L associated to the Aitchison diffusion X from (3) and (16), respectively, coincide. Consequently, the Brownian motion on the Aitchison simplex \hat{X} as introduced in (13) and the Aitchison diffusion X have the same law and thus X also satisfies A1. and A2. \square

Subsequently, we want to point out a different and instructive way to deduce Theorem 4 by rather geometric arguments. At that, we mainly follow the lines of [43, ch. 8]. First, observe that akin to \hat{X} , we can rewrite (3) as

$$dX_t = \sum_{i=1}^n Z_i(X_t) \circ dB_t^i, \quad X_0 = p \in \Delta, \quad (18)$$

where now

$$Z_i := \sum_{j=1}^n g^{ij} \epsilon_j.$$

So, the generator L of the Aitchison diffusion in Hörmander form reads

$$L = \frac{1}{2} \sum_{i=1}^n Z_i^2, \quad (19)$$

where again we identified the maps Z_i with the corresponding vector fields

$$Z_i := \sum_{j=1}^n g^{ij} \partial_j. \quad (20)$$

We omit the proof of the following result, which is tedious but straight forward.

Lemma 5. *The vector fields Z_1, \dots, Z_n as defined in (20) commute.*

Elementary facts from ODE theory ensure that we can determine uniquely a smooth map $E: \mathbb{R}^n \times \Delta \rightarrow \mathbb{R}^n$ satisfying

$$\frac{d}{dt}E(t\xi, p) = \sum_{i=1}^n \xi_i Z_i(E(t\xi, p)), \quad E(0, p) = p \in \Delta. \quad (21)$$

Moreover, Lemma 5 necessitates that

$$\partial_{\xi_k} E(\xi, p) = Z_k(E(\xi, p)) \quad (22)$$

for all $(\xi, p) \in \mathbb{R}^n \times \Delta$ and $k = 1, \dots, n$. Now take an n -dimensional standard Brownian motion B be on $(\Omega, \mathcal{F}, \mathcal{F}_t, \mathbb{P})$ and consider $X := E(B, p)$. Then, the Stratonovich chain rule and (22) entail

$$dX_t = \sum_{i=1}^n Z_i(X_t) \circ dB_t^i$$

and $X_0 = E(0, p) = p \in \Delta$. Thus, X is a solution to (3) and moreover defines a flow of diffeomorphisms on Δ . On the other hand, recall from (12) that

$$D \text{sfm} = \text{diag}(\text{sfm}) - \text{sfm} \otimes \text{sfm}.$$

Using this identity, we find that $E(\xi, p) = \text{sfm}(\xi) \oplus p$ provides a solution to (21), which in turn implies that the Aitchison diffusion starting in $p \in \Delta$ is simply given by

$$X_t = p \oplus \text{sfm}(B_t). \quad (23)$$

Of course, X inherits the properties A1.- A3. from B by the transformation rules (6). A simulation of a trajectory of the Aitchison diffusion on Δ_3 is depicted as a ternary plot in figure 1.

In the following we denote by P_t^X the Markov semigroup associated to X , i.e. for bounded and measurable functions f on Δ we set

$$P_t^X f(p) := \mathbb{E}[f(X_t) | X_0 = p]. \quad (24)$$

Also, we write P_t^B for the Brownian semigroup. The characterisations of the Aitchison diffusion which we discussed so far imply immediately the following properties of P_t^X .

Corollary 6 (invariant measure and density kernel of the Aitchison semigroup).

- (i) *The Aitchison measure λ_A is invariant and reversible for P_t^X .*
- (ii) *P_t^X admits a density kernel $(0, \infty) \times \Delta \times \Delta \ni (t, p, q) \mapsto p_t(p, q)$ with respect to λ_A , which is given by*

$$p_t(p, q) = (2\pi t)^{-\frac{n-1}{2}} e^{-\frac{\|p \ominus q\|_A^2}{2t}}.$$

In particular, d_A is the intrinsic metric for X and we have the Varadhan short time asymptotics

$$\lim_{t \downarrow 0} t \ln p_t(p, q) = -\frac{d_A^2(p, q)}{2}. \quad (25)$$




Figure 1: Simulation of a trajectory of an Aitchison diffusion starting at the barycenter

Proof. (i) Let $f: \Delta \rightarrow \mathbb{R}$ be positive, measurable and bounded. Then,

$$\begin{aligned} \int_{\Delta} P_t^X f(p) \lambda_A(dp) &= \int \mathbb{E} [\hat{X}_t | \hat{X}_0 = \text{ilr}^{-1}(x)] dx = \int P_t^B f \circ \text{ilr}^{-1}(x) dx \\ &= \int f \circ \text{ilr}^{-1}(x) dx = \int_{\Delta} f(p) \lambda_A(dp) \end{aligned}$$

proving invariance of λ_A . Reversibility follows by a similar argument.

For (ii), let $p := \text{ilr}^{-1}(x)$ for some $x \in \mathbb{R}^{n-1}$ and observe that

$$\begin{aligned} P_t^X f(p) &= \mathbb{E} [f \circ \text{ilr}^{-1}(B_t) | B_0 = x] = \int f \circ \text{ilr}^{-1}(y) (2\pi t)^{-\frac{n-1}{2}} e^{-\frac{\|x-y\|^2}{2t}} dy \\ &= \int_{\Delta} f(q) p_t(p, q) \lambda_A(dq), \end{aligned}$$

with

$$p_t(p, q) = (2\pi t)^{-\frac{n-1}{2}} e^{-\frac{\|p \ominus q\|_A^2}{2t}}.$$

The previous identity immediately gives (25). \square

Notice that λ_A is not a finite measure on Δ in accordance with the fact that X_t is transient, meaning $X_t \rightarrow \partial\Delta$ as $t \rightarrow \infty$ a.s. This again follows from the simple observation

$$\lim_{t \rightarrow \infty} \|X_t\|_A = \lim_{t \rightarrow \infty} \|B_t\| = \infty \text{ a.s.}$$

In fact, one can even identify the distribution of the limit. Namely, first observe that

$$\lim_{t \rightarrow \infty} X_t = \lim_{t \rightarrow \infty} p \oplus \text{sfm}(B_t) \stackrel{d}{=} p \oplus \lim_{t \rightarrow \infty} \text{sfm}(\sqrt{t}B_1) = p \oplus \text{argmax}(B_1).$$

Since that standard normal distribution attributes zero mass to the set of points in \mathbb{R}^n which have no distinct maximum it follow that $\lim_{t \rightarrow \infty} X_t \sim \text{Uniform}(\{\epsilon_1, \dots, \epsilon_n\})$.

4. THREE APPROXIMATION RESULTS FOR THE AITCHISON DIFFUSION

In this section we want to provide three approximation results for the Aitchison diffusion, which rely on classical theorems from stochastic analysis or optimal transport and PDE theory, namely Wong-Zakai approximation, Donsker's invariance principle and the JKO-scheme for the heat equation.

Recall, that in the introduction we alleged that the Aitchison diffusion, as described via the Stratonovich SDE

$$dX_t^i = X_t^i \left(\circ dB_t^i - \sum_{j=1}^n X_t^j \circ dB_t^j \right), \quad i = 1, \dots, n$$

$$X_0 = p \in \Delta,$$

has the natural interpretation of being a replicator equation in a white noise fitness landscape. The first statement gives a justification for this assertion. Indeed, we will see that (3) can be derived from a replicator dynamic in a coloured noise fitness landscape, when sending the correlation length to zero.

To this end, consider on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ the fitness landscape $y = (y_t)_{t \geq 0}$, which is independent of the current population state, but evolves randomly and continuously in time as a Gaussian process with correlation structure

$$\mathbb{E}[y_s^i y_t^j] = \delta_{ij} e^{\lambda^{-2}|s-t|}, \quad i, j = 1, \dots, n,$$

where $\lambda > 0$ is a correlation length parameter. In other words, we assume that the fitness landscape y is given by the n -dimensional Ornstein-Uhlenbeck process

$$\lambda^2 dy_t = -y_t dt + \lambda dB_t,$$

with B an n -dimensional Brownian motion and $y_0 \sim \mathcal{N}(0, \text{Id}_{\mathbb{R}^n})$. Then we ascertain the following

Theorem 7 (Approximation of the Aitchison diffusion by replicator dynamics in a Gaussian fitness landscape). *For every $\lambda > 0$ there exists a unique solution p^λ to*

$$\begin{aligned} \dot{p}_i(t) &= \lambda^{-1} p_i(t) (y_t^i - \bar{y}_t), \quad i = 1, \dots, n \\ p(0) &= p \in \Delta, \end{aligned} \tag{26}$$

where $\bar{y}_t = \sum_1^n p_i(t) y_t^i$ is the mean fitness. Moreover, the solutions p^λ converge weakly to the Aitchison diffusion X as λ goes to zero.

Proof. The existence of a unique solution p^λ to (26) follows again by standard Picard iteration. The convergence result is a classical incidence of the Wong-Zakai approximation [c.f. 17]. Weak convergence can be proven along the lines of [35, sec. 5.1.]. Invoking arguments from rough path theory one could obtain stronger convergence results in Hölder topologies as well [c.f. 19, 30]. \square

Our second approximation result establishes a connection between random walks on the Aitchison simplex and (3) in the spirit of Donsker's invariance principle [c.f. 32]. Therefore, first recall that the time discrete analogue of the replication dynamic (1) is given by the dynamical system

$$p_i^{(k+1)} = \frac{p_i^{(k)} f^i(p^{(k)})}{\sum_{j=1}^n p_j^{(k)} f^j(p^{(k)})}$$

$$p^{(0)} = p \in \Delta.$$

It was pointed out in [22] and [40] that we may think of this dynamic as modelling species adaptation by means of generation-wise Bayesian updating.

As previously, we replace the frequency dependent fitness f by random entities. More precisely, on some probability space $(\Omega, \mathcal{F}, \mathbb{P})$ we take iid Δ -valued random variables $f_{(1)}, f_{(2)}, \dots$ and consider

$$p_i^{(k+1)} = \frac{p_i^{(k)} f_{(k+1)}^i}{\sum_{j=1}^n p_j^{(k)} f_{(k+1)}^j}$$

$$p^{(0)} = p \in \Delta.$$

Using the Aitchison calculus we rewrite the previous updating rule simply as

$$p^{(k+1)} = p^{(k)} \oplus f_{(k+1)}$$

$$p^{(0)} = p \in \Delta. \tag{27}$$

or explicitly

$$p^{(k)} = \bigoplus_{i=1}^k f_{(i)} \oplus p.$$

That means $(p^{(k)})_{k \geq 0}$ is nothing but the random walk on the Aitchison simplex (Δ, \oplus, \odot) , induced through the iid random variables $(f_{(k)})_{k \geq 1}$. As a direct consequence of the continuous mapping theorem and Donsker's invariance principle we infer

Theorem 8 (Approximation of the Aitchison diffusion by random walks). *Let $f_{(1)}, f_{(2)}, \dots$ be iid random variables with values in Δ such that for all $k \in \mathbb{N}$*

$$\mathbb{E} \operatorname{clr}(f_{(k)}) = 0$$

and

$$\operatorname{cov}(\operatorname{ilr}_i(f_{(k)}), \operatorname{ilr}_j(f_{(k)})) = \delta_{ij}, \quad i, j = 1, \dots, n-1.$$

Let $(p^{(k)})_{k \geq 0}$ be a random walk on Δ as given in (27) and assume for simplicity $p^{(0)} = p = e$, with e being the barycenter of Δ . Define the linear interpolation of the random walk as

$$p_t := p^{(\lfloor t \rfloor)} \oplus (t - \lfloor t \rfloor) \odot p^{(\lfloor t \rfloor + 1)}.$$

Then the family of $C([0, 1], \Delta)$ -valued random elements $(n^{-\frac{1}{2}} \odot p_{nt})_{t \in [0, 1]}$, $n \in \mathbb{N}$, converges weakly to the Aitchison diffusion $(X_t)_{t \in [0, 1]}$ starting in $X_0 = e$ as $n \rightarrow \infty$.

Whereas the first two approximation results are in essence probabilistic, the last theorem in this section is based on a gradient flow interpretation of the Fokker-Planck equation associated to (3) and relies on the seminal work of Otto et al. in [28] and [34].

Let us start with a few definitions. We denote by $\mathcal{P}(\Delta)$ the set of all probability measures on Δ and by

$$\mathcal{P}_2(\Delta) := \left\{ \mu \in \mathcal{P}(\Delta) : \int_{\Delta} \|p\|_A^2 \mu(dp) < \infty \right\}.$$

Then a natural distance measure on $\mathcal{P}_2(\Delta)$ is the Wasserstein distance

$$W_A^2(\mu, \nu) := \inf_{\pi \in \Pi(\mu, \nu)} \int_{\Delta \times \Delta} d_A^2(p, q) \pi(dp dq), \quad (28)$$

where for $\mu, \nu \in \mathcal{P}_2(\Delta)$ we have

$$\Pi(\mu, \nu) := \{ \pi \in \mathcal{P}(\Delta \times \Delta) : \pi(A \times \Delta) = \mu(A) \text{ and } \pi(\Delta \times B) = \nu(B), A, B \in \mathcal{B}(\Delta) \}.$$

Notice, that the Wasserstein distance on $\mathcal{P}_2(\Delta)$ is linked to the usual Wasserstein metric on $\mathcal{P}_2(\mathbb{R}^{n-1})$ through

$$W_A^2(\mu, \nu) = \inf_{\pi \in \Pi(\text{ilr}_{\#} \mu, \text{ilr}_{\#} \nu)} \int_{\mathbb{R}^{n-1}} \|x - y\|^2 \pi(dx dy) =: W_{\mathbb{R}^{n-1}}^2(\text{ilr}_{\#} \mu, \text{ilr}_{\#} \nu). \quad (29)$$

Finally, we denote by \mathcal{S}_A the Boltzmann entropy on the Aitchison simplex:

$$\mathcal{S}_A(\mu) := \begin{cases} \int_{\Delta} \ln \left(\frac{d\mu}{d\lambda_A}(p) \right) \mu(dp), & \mu \ll \lambda_A \\ +\infty, & \text{else.} \end{cases}$$

Now let X be an Aitchison diffusion with $X_0 \sim \mu_0$ for some $\mu_0 \in \mathcal{P}_2(\Delta)$. We denote by $\mu_t(dp) = \mu_t(p) \lambda_A(dp)$ the law of X_t . Then the densities $(\mu_t)_{t \geq 0}$ provide a solution to the heat equation on the Aitchison simplex

$$\partial_t \mu = L\mu, \quad \mu|_{t=0} = \mu_0 \quad (30)$$

and satisfy for any $\nu \in \mathcal{P}_2(\Delta)$ with finite entropy

$$\frac{d}{dt} W_A^2(\mu_t, \nu) \leq \mathcal{S}_A(\nu) - \mathcal{S}_A(\mu_t). \quad (31)$$

The meaning of the previous evolution-variational inequality is that (the laws of) the Aitchison diffusion evolve as a Wasserstein gradient flow of the Boltzmann entropy. To see, why (31) is true, let ρ be a solution to the heat equation on \mathbb{R}^{n-1} . Then, denoting by

$$\mathcal{S}_{\mathbb{R}^{n-1}}(\mu) := \begin{cases} \int_{\mathbb{R}^{n-1}} \ln \left(\frac{d\mu}{d\lambda_{\mathbb{R}^{n-1}}}(x) \right) \mu(dx), & \mu \ll \lambda_{\mathbb{R}^{n-1}} \\ +\infty, & \text{else} \end{cases}$$

the usual entropy on \mathbb{R}^{n-1} , we know that

$$\frac{d}{dt} W_{\mathbb{R}^{n-1}}^2(\rho_t, \eta) \leq \mathcal{S}_{\mathbb{R}^{n-1}}(\eta) - \mathcal{S}_{\mathbb{R}^{n-1}}(\rho_t),$$

for all probability measures η on \mathbb{R}^{n-1} with finite second moment and entropy [c.f. 4, 16]. Next, notice that if $\nu \in \mathcal{P}(\Delta)$ has finite entropy, then

$$\mathcal{S}_{\mathbb{R}^{n-1}}(\text{ilr}_{\#} \nu) = \int_{\mathbb{R}^{n-1}} \ln \left(\frac{d \text{ilr}_{\#} \nu}{d \lambda_{\mathbb{R}^{n-1}}}(x) \right) \text{ilr}_{\#} \nu(dx) = \int_{\Delta} \ln \left(\frac{d \text{ilr}_{\#} \nu}{d \text{ilr}_{\#} \lambda_A}(\text{ilr}(p)) \right) \nu(dp) = \mathcal{S}_A(\nu).$$

Now we easily infer (31) from the previous identity, (29) and the fact that $\mu_t = \text{ilr}_{\#}^{-1} \rho_t$.

The observation that the Aitchison diffusion can be interpreted as a Wasserstein gradient flow now provides an immediate way to approximate the laws μ_t by means of a steepest descent algorithm, in perfect analogy to the JKO-scheme invented in [28].

Theorem 9 (Approximation of the Aitchison diffusion by JKO-scheme). *For every $t > 0$ we denote by*

$$J_t(\bar{\mu}|\mu_0) := \mathcal{S}(\bar{\mu}) - \mathcal{S}(\mu_0) - \frac{1}{2t} W_A^2(\bar{\mu}, \mu_0) \quad \text{and} \quad K_t[\rho_0] := \underset{\bar{\mu} \in \mathcal{P}_2(\Delta)}{\text{argmin}} J_t(\bar{\mu}|\mu_0)$$

Then,

$$\mu_t := \lim_{n \rightarrow \infty} (K_{t/n})^n[\mu_0], \quad t > 0$$

is the law of the Aitchison diffusion (at time t) and its λ_A -density solves the heat equation (30).

5. DRIFT DIFFUSIONS ON THE AITCHISON SIMPLEX

In this final section we are concerned with the influence of drift terms on the Aitchison diffusion. More precisely, we are interested in the long time behaviour of Markov processes which are associated to differential operators of the form $L_{Z_0} := L + Z_0$, where L is the generator of the Aitchison diffusion as in (19) and Z_0 is a vector field on Δ . Rather than aiming for maximal generality, our focus is on examples which seem interesting for applications in e.g. mathematical biology and game theory, foremost the case

$$Z_0 f(p) = Ap \cdot \nabla^g f(p),$$

which corresponds to the stochastic replication equation. Our investigations will be split into two parts. First, we derive structural properties of invariant measures for the diffusion processes in question. Afterwards, we provide quantitative statements about the relaxation to equilibrium by means of Wasserstein contraction estimates. For both concerns we will benefit from a frequent change of perspective between the usual Euclidean picture on one hand and the Aitchison geometry on the other hand.

As a first example for this approach, consider

Definition (ODE on the Aitchison simplex). Let $F: \Delta \rightarrow \Delta$. By a solution to the ordinary differential equation (ODE) on the Aitchison simplex

$$\dot{p}(t) = F(p(t)), \quad p(0) = p \in \Delta, \quad (32)$$

we mean a continuous map $[0, \infty) \ni t \mapsto p(t) \in \Delta$, obeying

$$\langle p(t), q \rangle_A = \langle p, q \rangle_A + \int_0^t \langle F(p(s)), q \rangle_A ds, \quad t \geq 0$$

for all $q \in \Delta$.

Lemma 10. *Fix $p \in \Delta$. Then $(p(t))$ is a solution to (32) if and only if it is a solution to the conventional ODE*

$$\dot{p}(t) = g^{-1}(p(t)) \operatorname{clr}(F(p(s))), \quad p(0) = p. \quad (33)$$

Proof. We set $\tilde{p}(t) := \operatorname{clr}(p(t))$. Recall that for $p, q \in \Delta$ we have $\langle p, q \rangle_A = (\operatorname{clr}(p), \operatorname{clr}(q))_H$. Thus, $(p(t))$ is a solution to (32) iff for all $h \in H$

$$(\tilde{p}(t), h)_H = (\tilde{p}, h)_H + \int_0^t (\operatorname{clr}(F(p(s))), h)_H ds, \quad (34)$$

Therefore, if $(p(t))$ solves (32), it follows

$$\dot{p}(t) = \frac{d}{dt} \operatorname{sfm}(\tilde{p}(t)) = g^{-1}(p(t)) \dot{\tilde{p}}(t) = g^{-1}(p(t)) \operatorname{clr}(F(p(s)))$$

which is (33). On the other hand, if $(p(t))$ solves (33), then since

$$\frac{d}{dt} \ln p_i(t) = \ln F_i(p(t)) - \sum_{j=1}^n p_j(t) \ln(F_j(t)),$$

we find

$$\dot{\tilde{p}}(t) = \frac{d}{dt} \operatorname{clr}(p(t)) = \operatorname{clr}(F(p(t)))$$

which, using (34), yields that $(p(t))$ solves (32). \square

Remark. *Notice that the transformation above has a nice biological interpretation. Think of F in (32) as Wrightian fitness. Then the flow $(p(t))$ on the Aitchison simplex driven by F corresponds in the usual Euclidean notation to a replicator equation with fitness landscape $(\ln F_1, \dots, \ln F_n)$, i.e. the Malthusian fitness [c.f. 48].*

In particular, given some payoff matrix $A \in \mathbb{R}^{n \times n}$, if we define

$$\theta: \Delta \rightarrow \Delta, \quad p \mapsto \theta(p) := \operatorname{sfm}(Ap), \quad (35)$$

then

$$\dot{p} = \theta(p), \quad p(0) = p \in \Delta$$

is equivalent to the replicator equation with linear fitness landscape $p \mapsto Ap$:

$$\dot{p}_i(t) = p_i(t) ((Ap(t))_i - p(t) \cdot Ap(t)), \quad i = 1, \dots, n \quad (36)$$

provided $p(0) = p \in \Delta$.

Let us try to establish a stochastic analogue of the previous observation.

Definition (SDE on the Aitchison simplex). Let $F: \Delta \rightarrow \Delta$ and $p \in \Delta$. Given a filtered probability space $\Omega = (\Omega, \mathcal{F}, (\mathcal{F})_t, \mathbb{P})$ and an Aitchison diffusion X on Ω , we call an \mathcal{F}_t -adapted, time-continuous and Δ -valued process Y a solution to the SDE on the Aitchison simplex,

$$dY_t = F(Y_t)dt \oplus dX_t, \quad Y_0 = p, \quad (37)$$

provided that \mathbb{P} -a.s.

$$\langle Y_t, q \rangle_A = \langle p, q \rangle_A + \int_0^t \langle F(Y_s), q \rangle_A ds + \langle X_t, q \rangle_A, \quad t \geq 0 \quad (38)$$

for all $q \in \Delta$.

The definition can be extended to random initial data in the usual way. Our next result is the stochastic counterpart of Lemma 10.

Theorem 11. *Let $\Omega = (\Omega, \mathcal{F}, (\mathcal{F})_t, \mathbb{P})$ be a filtered probability space. If Y is a solution to (37) on Ω , then there exists an \mathbb{R}^n -valued standard Brownian motion B on Ω , such that Y solves*

$$dY_t = g^{-1}(Y_t) \text{clr}(F(Y_t))dt + g^{-1}(Y_t) \circ dB_t. \quad (39)$$

Vice versa, if Y solves (39), then there exists an Aitchison diffusion X on Ω such that Y solves (37).

Proof. Let Y be a solution to (37). Since X is a Brownian motion on $(\Delta, \langle \cdot, \cdot \rangle_A)$, we have

$$\mathbb{E}[\langle X_t, p \rangle_A \langle X_s, q \rangle_A] = \langle p, q \rangle_{As} \wedge t, \quad (40)$$

whence, for all $g, h \in H$

$$\mathbb{E}[(\text{clr}(X_t), g)_H (\text{clr}(X_s), h)_H] = (g, h)_{Hs} \wedge t.$$

Therefore $\tilde{B} := \text{clr}(X)$ is a BM on H . Now take another one-dimensional Brownian motion W , independent from \tilde{B} and define $B := \tilde{B} + \frac{1}{\sqrt{n}}W\mathbf{1}$. Then, because

$$\begin{aligned} & \mathbb{E}[(B_t, x)(B_s, y)] \\ &= \mathbb{E} \left[(\tilde{B}_t, x)(\tilde{B}_s, y) + n^{-\frac{1}{2}}(\tilde{B}_t, x)(\mathbf{1}, y)W_s + n^{-\frac{1}{2}}(\tilde{B}_s, y)(\mathbf{1}, x)W_t + n^{-1}W_sW_t(\mathbf{1}, x)(\mathbf{1}, y) \right] \\ &= (\text{pr}_H x, \text{pr}_H y)_{Hs} \wedge t + \frac{1}{n}(\mathbf{1}, x)(\mathbf{1}, y)s \wedge t = (x, y)s \wedge t, \end{aligned}$$

we know B is a standard BM on \mathbb{R}^n . Next, set $\tilde{Y} := \text{clr}(Y)$ and $h := \text{clr}(p)$. By (38),

$$(\tilde{Y}_t, h)_H = (\tilde{Y}_0, h)_H - \int_0^t (\text{clr}(F(Y_s)), h)_H ds + (B_t - n^{-\frac{1}{2}}W_t\mathbf{1}, h)_H. \quad (41)$$

Now applying the Stratonovich chain rule to $Y = \text{sfm}(\tilde{Y})$, we recover (39).

If on the other hand Y solves (39), then

$$\begin{aligned} d \ln Y_t^i &= \frac{1}{Y_t^i} \left((g^{-1}(Y_t) \operatorname{clr}(F(Y_t)))_i dt + (g^{-1}(Y_t) \circ dB_t)_i \right) \\ &= - \left(\operatorname{clr}_i(F(Y_t)) - \sum_{k=1}^n Y_t^k \operatorname{clr}_k(F(Y_t)) \right) dt + dB_t^i - \sum_{k=1}^n Y_t^k \circ dB_t^k, \end{aligned}$$

and therefore

$$d\tilde{Y}_t^i := d \operatorname{clr}_i(Y_t) = - \operatorname{clr}_i(F(Y_t)) dt + dB_t^i - \frac{1}{n} \sum_{k=1}^n dB_t^k.$$

The noise term on the right hand side of the previous equation is a Brownian motion on H . Thus, testing \tilde{Y} with $h \in H$ and transforming back to the Aitchison simplex, we see Y obeys (37) with $X = \operatorname{sfm}(B)$. \square

Choosing again $F = \theta$ in (39), leads to

$$dY_t = g^{-1}(Y_t) A Y_t dt + g^{-1}(Y_t) \circ dB_t, \quad (42)$$

which is the stochastic replicator equation of Fudenberg and Harris as in (2) with $\sigma_i = 1$ for $i = 1, \dots, n$. We will here consider this constant coefficient case only. Yet, our methods could be easily extended to different σ_i , too. In this case, one would need to incorporate a covariance structure in the definition of X , by imposing e.g.

$$\mathbb{E}[\langle X_t, p \rangle_A \langle X_s, q \rangle_A] = (\operatorname{clr}(p), \operatorname{diag}(\sigma_1, \dots, \sigma_n) \operatorname{clr}(q)) s \wedge t. \quad (43)$$

There is another interesting and natural choice for F in (37). Namely, take a potential $V \in C^1(\Delta)$ and consider the gradient drift $F = \ominus \nabla^A V$. The associated evolution can be interpreted as a Langevin dynamic on the Aitchison simplex:

$$dY_t = \ominus \nabla^A V(Y_t) dt \oplus dX_t, \quad (44)$$

which in the standard Euclidean picture corresponds to

$$dY_t = -\nabla^G V(Y_t) dt + g^{-1}(Y_t) \circ dB_t. \quad (45)$$

where $\nabla^G := g^{-1} \nabla^g$. We will come back to those gradient drift diffusions in due course.

In preparation for the following two subsections, we need to introduce a classical notion of evolutionary game theory. As for the deterministic replicator equation (36), a key concept in the analysis of the long term behaviour of (42) are Price and Maynard Smith's *evolutionary stable strategies* (ESS) [42], which specify Nash equilibria that are non-invadable by initially rare alternative strategies.

Definition (Evolutionary stable strategy). Let $A = (a_{ij}) \in \mathbb{R}^{n \times n}$ be a payoff matrix. We call $p^* \in \bar{\Delta}$ an ESS for A provided

1. *equilibrium condition*

$$p \cdot A p^* \leq p^* \cdot A p^*, \text{ for all } p \in \bar{\Delta}$$

2. *stability condition*

$$\text{if } p \neq p^* \text{ and } p \cdot Ap^* = p^* \cdot Ap^* \text{ then } p \cdot Ap < p^* \cdot Ap. \quad (46)$$

The first condition above means that p^* is a *Nash equilibrium* (NE).

It is well-known that if $p^* \in \Delta$ is an interior ESS, then it must be unique and moreover A is conditionally negative definite [c.f. 24]:

Definition. We call $A \in \mathbb{R}^{n \times n}$ *conditionally negative semi-definite*, whenever,

$$h \cdot Ah \leq 0, \text{ for all } h \in H.$$

If the previous inequality is strict for all $h \in H \setminus \{0\}$, we say A is *conditionally negative definite*. We denote by $\Gamma^{\leq} = \Gamma_n^{\leq}$ and $\Gamma^{<} = \Gamma_n^{<}$ the sets of all $n \times n$ conditionally negative semi-definite and conditionally negative definite matrices, respectively.

If on the other hand, A is conditionally negative definite, then A has a unique ESS, possibly lying on the boundary.

Finally notice [c.f. 27], if $A \in \Gamma^{<}$ and we introduce the Rayleigh-quotient

$$\lambda := - \max_{h \in H \setminus \{0\}} \frac{h \cdot Ah}{\|h\|^2},$$

then $\lambda > 0$ and for all $h \in H$

$$h \cdot Ah \leq -\lambda \|h\|^2. \quad (47)$$

The parameter λ will play a crucial role in our final section on contraction estimates for replicator dynamics.

5.1. Fokker-Planck equation and invariant measures for stochastic replicator dynamics

In [27] Imhof studies the long-run behavior of stochastic replicator dynamics, providing in particular sufficient conditions for the existence of invariant probability measures. The authors of [23] investigate among others ergodicity properties of (42) and their consequences. Our aim for this subsection is to complement those results by a classical perspective on invariant measures, namely via the Fokker-Planck equation.

Mainly for notational convenience, and in this subsection only, X will generically be an Aitchison diffusion with covariance structure

$$\mathbb{E}[\langle X_t, p \rangle_A \langle X_s, q \rangle_A] = \sqrt{2} \langle p, q \rangle_{AS} \wedge t, \quad (48)$$

which in the Fudenberg-Harris model corresponds to $\sigma_i = \sqrt{2}$, $i = 1, \dots, n$, or equivalently to (42) if the Brownian motion obeys $\langle B^i, B^j \rangle_t = \sqrt{2} \delta_{ij} t$. We denote by (P_t) the Markov semi-group associated to (42), which we also refer to as the replicator semigroup. For the corresponding family of Markov transition kernels we write (π_t) . Of course, P_t depends

upon the choice of a payoff matrix $A \in \mathbb{R}^{n \times n}$ and so does the corresponding generator $L_A := L' + Z_A$, where $L' = 2L$ with L as in (19) and for $f: \Delta \rightarrow \mathbb{R}$ sufficiently regular,

$$Z_A f(p) := Ap \cdot \nabla^g f(p).$$

As usual, we call a σ -finite (but possibly non-finite) measure μ on Δ invariant for the replicator semigroup, if for all positive, bounded and measurable functions $f: \Delta \rightarrow \mathbb{R}$

$$\int_{\Delta} P_t f d\mu = \int_{\Delta} f d\mu, \quad t \geq 0. \quad (49)$$

Let us start with the following simple, yet important observation which is a direct consequence of the isometry between Δ and \mathbb{R}^{n-1} and Theorem 11.

Lemma 12 (Change of coordinates formula). *Denote by*

$$\hat{\theta}(x) := \text{ilr} \circ \theta \circ \text{ilr}^{-1}(x) = \Psi A \text{sfm}(\Psi^\top x) \quad (50)$$

and consider the \mathbb{R}^{n-1} -valued diffusion \hat{Y} given by

$$d\hat{Y}_t = \hat{\theta}(\hat{Y}_t) dt + \sqrt{2} dB_t, \quad (51)$$

with corresponding generator $\hat{L}_A := \Delta + \hat{\theta} \cdot \nabla$. The Markov semigroup (\hat{P}_t) given through (51) and the replicator semigroup are linked by means of the change of coordinates

$$P_t f(p) = \hat{P}_t(f \circ \text{ilr}^{-1})(\text{ilr}(p)). \quad (52)$$

Hence, the analysis of (P_t) eventually boils down to the analysis of the drifted Brownian motion \hat{Y} and its semigroup (\hat{P}_t) . Now recall that the Markov semigroup (P_t) is regular provided for all $t > 0$, the probability measures $\pi_t(p, \cdot)$ are mutually equivalent for all $p \in \Delta$ and $t > 0$. Then, as first consequence of Lemma 12, we obtain

Proposition 13. *The replicator semigroup is regular.*

Proof. Notice that, $\hat{\theta}$ is bounded and Lipschitz. Indeed, we have

$$\|\hat{\theta}(x) - \hat{\theta}(y)\| \leq \|A\| \|\Psi\|^2 \|x - y\|,$$

and

$$\|\hat{\theta}(x)\| \leq \|\Psi\| \|A\| \quad (53)$$

for all $x, y \in \mathbb{R}^{n-1}$. Therefore, (\hat{P}_t) is strongly Feller and irreducible [12, Prop. 7.20], hence regular. By Lemma 12 the replicator semigroup (P_t) inherits the regularity from (\hat{P}_t) . \square

By [11, Thm 4.2.1], regularity of a semigroup on the other hand directly entails

Corollary 14 (Uniqueness of invariant probability measures and mixing property). *Let $\mu \in \mathcal{P}(\Delta)$ be invariant for (P_t) . Then μ is the only invariant measure and moreover the replicator semigroup is strongly mixing for μ , i.e. for every $p \in \Delta$ and measurable $C \subset \Delta$*

$$\lim_{t \rightarrow \infty} \pi_t(p, C) = \mu(C). \quad (54)$$

Our next aim is a description of invariant measures by means of a stationary Fokker-Planck equation. As a preparatory result we deliver following statement on densities of invariant measures.

Proposition 15 (Existence of smooth λ_A -densities). *If μ is an invariant measure for the replicator semigroup, then μ has a C^∞ -smooth density with respect to the Aitchison measure λ_A .*

Proof. By (52) invariant measures μ for (P_t) and $\hat{\mu}$ for (\hat{P}_t) , respectively, are in one to one correspondence via

$$\mu = \text{ilr}_\#^{-1} \hat{\mu} \quad \text{and} \quad \hat{\mu} = \text{ilr}_\# \mu. \quad (55)$$

Therefore, and because $\lambda_A = \text{ilr}_\#^{-1} \lambda_{\mathbb{R}^{n-1}}$ it is enough to prove, that every invariant measure $\hat{\mu}$ of (\hat{P}_t) admits a smooth Lebesgue density. But in fact, suppose $\hat{\mu}$ is an invariant measure. Then it is a positive, weak solution to the stationary Fokker-Planck equation

$$0 = \hat{L}_A^* \hat{\mu} := \Delta \hat{\mu} - \nabla \cdot (\hat{\mu} \hat{\theta}), \quad (56)$$

\hat{L}_A^* being the formal $L^2(\lambda_{\mathbb{R}^{n-1}})$ -adjoint of \hat{L}_A . However, since \hat{L}_A^* has smooth coefficients, by Weyl's regularity theorem [c.f. 9, Thm 1.4.6] any weak solution to (56) is a smooth classical solution. In particular, $\hat{\mu}$ has a smooth density with respect to the Lebesgue measure on \mathbb{R}^{n-1} . \square

We are now ready to present the main result of this subsection. Let us define for every $p \in \Delta$ the potential

$$\Lambda(p) := \sum_{i=1}^n a_{ii} p_i - p \cdot Ap. \quad (57)$$

Occasionally, we will also write Λ_A if we want to emphasize the dependence on the matrix A . As we shall see, this potential plays a decisive role for the long time behavior of stochastic replicator dynamics.

Theorem 16 (Stationary Fokker-Planck equation). *Let μ be invariant for (P_t) . Then the λ_A density of μ is a solution to the stationary Fokker-Planck equation*

$$0 = L' \mu - Z_A \mu - \Lambda \mu. \quad (58)$$

Vice versa, assume $\mu \in C^2(\Delta)$ is a strictly positive solution to (58) whose logarithmic gradient is locally Lipschitz and has linear growth, i.e.

$$\|\nabla^g \ln \mu(p)\| \leq K(1 + \|p\|_A) \quad (59)$$

for some $K > 0$ and all $p \in \Delta$. Then $\mu d\lambda_A$ is invariant for (P_t) .

Proof. We have to determine L_A^* the formal $L^2(\lambda_A)$ -adjoint of L_A . Since, λ_A is reversible for L' , we only need to focus the vector field Z_A and claim that its formal $L^2(\lambda_A)$ -adjoint Z_A^* is given by

$$Z_A^* = -Z_A - \Lambda.$$

Indeed, consider $f, g \in C_0^\infty(\Delta)$, set $\hat{f} := f \circ \text{ilr}^{-1}$ and likewise for \hat{g} . Then, since

$$Z_A f = (\Psi A \text{ilr}^{-1}, \nabla f \circ \text{ilr}^{-1}) \circ \text{ilr},$$

it follows

$$\int_{\Delta} g Z_A f d\lambda_A = \int_{\mathbb{R}^{n-1}} \hat{g}(x) (\Psi A \text{ilr}^{-1}(x), \nabla \hat{f}(x)) dx. \quad (60)$$

But because

$$\nabla \cdot (\Psi A \text{ilr}^{-1}(x)) = \sum_{i=1}^n a_{ii} \text{ilr}_i^{-1}(x) - \text{ilr}^{-1}(x) \cdot A \text{ilr}^{-1}(x) = \Lambda \circ \text{ilr}^{-1}(x)$$

and $\hat{f}, \hat{g} \in C_0^\infty(\mathbb{R}^{n-1})$, integrating by parts in (60) yields

$$\int_{\Delta} g Z_A f d\lambda_A = \int_{\Delta} (-Z_A g - \Lambda g) f d\lambda_A. \quad (61)$$

Now if $\mu d\lambda_A$ is invariant for (P_t) , then for all smooth and compactly supported test functions f on Δ we have

$$0 = \int_{\Delta} L_A f \mu d\lambda_A = \int_{\Delta} f L_A^* \mu d\lambda_A,$$

which yields the claim by the Lemma of du Bois-Reymond.

For the converse direction assume $\mu \in C^2(\Delta)$ is a positive solution to (58). Then $\hat{\mu} := \mu \circ \text{ilr}^{-1}$ is a positive solution to

$$0 = \hat{L}_A^* \hat{\mu}, \quad (62)$$

or in other words, $\hat{\mu}$ is infinitesimally invariant for (\hat{P}_t) . Next, consider the Doob h -transform $\hat{\mathcal{L}}$ of \hat{L}_A defined by

$$\hat{\mathcal{L}} f := \frac{1}{\hat{\mu}} \hat{L}_A (\hat{\mu} f) \quad (63)$$

and set

$$\hat{\mathcal{L}}^\dagger := \hat{\mathcal{L}}^* - \hat{\mathcal{L}}^* 1. \quad (64)$$

Thus, $\hat{\mathcal{L}}^\dagger$ is the formal $L^2(\lambda_{\mathbb{R}^{n-1}})$ -adjoint of $\hat{\mathcal{L}}$ minus its zero order part and

$$\hat{\mathcal{L}}^\dagger f = \Delta f - \hat{\theta} \cdot \nabla f - \nabla \ln \hat{\mu} \cdot \nabla f \quad (65)$$

for f sufficiently smooth. Now according to [38, Thm 4.8.5], the density $\hat{\mu}$ is invariant for (\hat{P}_t) iff the diffusion given by the martingale problem for $\hat{\mathcal{L}}^\dagger$ is non-explosive. But using the growth condition (59) and identifying $x = \text{ilr}(p)$, we know

$$\|\nabla \ln \hat{\mu}(x)\| = \|\nabla^g \ln \mu(p)\| \leq K(1 + \|p\|_A) = K(1 + \|x\|) \quad (66)$$

for all $x \in \mathbb{R}^{n-1}$. Likewise, it follows that the logarithmic gradient of $\hat{\mu}$ is locally Lipschitz. Therefore, all coefficients of $\hat{\mathcal{L}}^\dagger$ are locally Lipschitz and satisfy a linear growth condition, whence classical theory [c.f. 26, ch. 6] guarantees apart from well-posedness of the martingale problem for $\hat{\mathcal{L}}^\dagger$, that the associated diffusion is conservative. \square

The rest of this subsection is devoted to the exploitation of the previous theorem. Note that some of the subsequent results were proven already in [23]. However, whereas Hofbauer and Imhof invoke elaborate Lyapunov function techniques, in our present setting they appear as direct consequences of Theorem 16.

Corollary 17. *The Aitchison measure λ_A is invariant for (P_t) if and only if the payoff matrix $\mathbb{R}^{n \times n} \ni A = (a_{ij})$ satisfies for all $i \neq j$*

$$a_{ij} + a_{ji} - a_{ii} - a_{jj} = 0. \quad (67)$$

Moreover, if $n > 3$ and A satisfies (67), then the replicator diffusion is transient in that $Y_t \rightarrow \partial\Delta$ as $t \rightarrow \infty$ almost surely.

Notice, the previous Corollary applies in particular to zero-sum games, that is, when $A = -A^\top$.

Proof. Since the Lipschitz and growth condition are trivially satisfied, by Theorem 16, λ_A is invariant iff for all $p \in \Delta$

$$0 = \Lambda(p) = \sum_{i=1}^n a_{ii} p_i - p \cdot Ap. \quad (68)$$

Observe, if A satisfies (67), then

$$\begin{aligned} p \cdot Ap &= \sum_{i \neq j} p_i a_{ij} p_j + \sum_{i=1}^n a_{ii} p_i^2 = - \sum_{i \neq j} p_i a_{ij} p_j + 2 \sum_{i \neq j} p_i a_{ii} p_j + \sum_{i=1}^n a_{ii} p_i^2 \\ &= -p \cdot Ap + 2 \sum_{i=1}^n a_{ii} p_i, \end{aligned} \quad (69)$$

whence (68) is fulfilled. On the other hand, if (68) is true for all $p \in \Delta$, then by continuity it is valid also for $p \in \bar{\Delta}$. Testing, (68) with $p = \frac{1}{2}\epsilon_i + \frac{1}{2}\epsilon_j$ for $i \neq j$ immediately yields (67).

Regarding the second statement, notice (67) entails

$$\nabla \cdot \hat{\theta}(x) = \Lambda \circ \text{ilr}^{-1}(x) = 0$$

for all $x \in \mathbb{R}^{n-1}$. Therefore, if $n \geq 4$ [38, cor. 6.3] implies \hat{Y} is transient on \mathbb{R}^{n-1} which yields the claim. \square

Occasionally, one is interested in invariant distributions of Gibbs-type

$$\mu = e^{-V} \lambda_A, \quad (70)$$

for some $V: \Delta \rightarrow \mathbb{R}$. We denote by Γ the carré du champs operator associated to L' , that is

$$\Gamma(f, g) := \langle \nabla^A f, \nabla^A g \rangle_A = (\nabla^g f, \nabla^g g) = \sum_{i=1}^n Z_i f Z_i g,$$

and $\Gamma f := \Gamma(f, f)$. Then a straight forward application of the diffusion property of L_A combined with Proposition 16 implies

Lemma 18. Let $V \in C^2(\Delta)$ and $\nabla^g V$ be locally Lipschitz and satisfy the growth condition (59). Then, $\mu = e^{-V} \lambda_A$ is an invariant measure for (P_t) if and only if V satisfies

$$0 = L'V - \Gamma V - Z_A V + \Lambda. \quad (71)$$

Remark. By classical theory [c.f. 6], we know that the Langevin dynamic on the Aitchison simplex

$$dY_t = \ominus \nabla^A V(Y_t) dt \oplus dX_t, \quad (72)$$

which is associated to the generator $L_V := L' - \Gamma(V, \cdot)$ has an invariant measure of the form (70). Since $L_A = L_V + Z_A + \Gamma(V, \cdot)$, the condition in (71) holds if and only if for all $f \in C_0^\infty(\Delta)$ we have

$$\int_{\Delta} (Z_A f + \Gamma(V, f)) d\mu = 0 \quad (73)$$

Corollary 19. Let $\alpha \in \mathbb{R}_{>0}^n$ and set $|\alpha| := \alpha_1 + \dots + \alpha_n$. The Dirichlet distribution with parameter α is the (unique) invariant measure for (P_t) if and only if

(i) the payoff matrix $A = (a_{ij})$ fulfills for $i \neq j$

$$a_{ij} + a_{ji} - a_{ii} - a_{jj} = 2|\alpha|$$

and

(ii) $\frac{\alpha}{|\alpha|}$ is a Nash equilibrium for A .

Proof. The Dirichlet distribution amounts to the choice of

$$V(p) := - \sum_{i=1}^n \alpha_i \ln p_i$$

in (70). Evidently,

$$\nabla^g V(p) = |\alpha|p - \alpha \quad (74)$$

satisfies the Lipschitz and growth conditions of Theorem 16. Plugging in the definition of V in (71) yields,

$$0 = (\alpha - |\alpha|p) \cdot Ap + \Lambda(p) - \|\alpha - |\alpha|p\|^2 + |\alpha| (1 - \|p\|^2). \quad (75)$$

Hence in order to prove Corollary 19, by Lemma 18 it is enough to show that (75) holds for all $p \in \Delta$ if and only if the conditions (i) and (ii) of Corollary 19 are fulfilled.

Let us start by assuming that A and α are such that (i) and (ii) are valid. It was shown in [23] that a matrix for which (i) holds obeys

$$h \cdot Ah = -|\alpha| \|h\|^2 \quad (76)$$

for every $h \in H$. Therefore, and because $\alpha/|\alpha|$ is an interior NE for A by (ii), we know that for any $p \in \Delta$

$$\begin{aligned} & (\alpha - |\alpha|p) \cdot Ap - \|\alpha - |\alpha|p\|^2 \\ &= -|\alpha| \left(\frac{\alpha}{|\alpha|} - p \right) \cdot A \left(\frac{\alpha}{|\alpha|} - p \right) - \|\alpha - |\alpha|p\|^2 + |\alpha| \left(\frac{\alpha}{|\alpha|} - p \right) \cdot A \left(\frac{\alpha}{|\alpha|} \right) = 0 \end{aligned}$$

Next, observe that arguing akin to (69), we see that for every $p \in \Delta$

$$p \cdot Ap = |\alpha| (1 - \|p\|^2) + \sum_{i=1}^n a_{ii} p_i, \quad (77)$$

which yields (75) for all $p \in \Delta$. Note that the previous identity can be rephrased as

$$\Lambda_A = \Lambda_{-|\alpha|\text{id}}. \quad (78)$$

Now assume on the contrary that (75) holds for all $p \in \Delta$. Testing the equation with ϵ_i , the i -th unit vector in \mathbb{R}^n (more precisely, take a sequence $(p_n)_{n \geq 0} \subset \Delta$ with $p_n \rightarrow \epsilon_i$ as $n \rightarrow \infty$, test (75) with p_n and take limits) yields

$$0 = \sum_{k=1}^n a_{ki} \alpha_k - |\alpha| a_{ii} - \|\alpha\|^2 + 2|\alpha| \alpha_i - |\alpha|^2. \quad (79)$$

Next, testing with $\frac{1}{2}\epsilon_i + \frac{1}{2}\epsilon_j$ we find

$$\begin{aligned} 0 &= \sum_{k=1}^n (a_{ki} + a_{kj}) \alpha_k - \frac{|\alpha|}{2} (a_{ii} + a_{ij} + a_{ji} + a_{jj}) + a_{ii} + a_{jj} - \frac{1}{2} (a_{ii} + a_{ij} + a_{ji} + a_{jj}) \\ &\quad - 2\|\alpha\|^2 + 2|\alpha|(\alpha_i + \alpha_j) - |\alpha|^2 + |\alpha|. \end{aligned}$$

Using (79), the previous expression simplifies to

$$0 = |\alpha|^2 + |\alpha| - \frac{|\alpha| + 1}{2} (a_{ij} + a_{ji} - a_{ii} - a_{jj}),$$

which thus gives condition (i). But then, due to (76), (77) and by assumption

$$|\alpha| \left(p - \frac{\alpha}{|\alpha|} \right) \cdot A \frac{\alpha}{|\alpha|} = \Lambda(p) + |\alpha| (1 - \|p\|^2) = 0$$

for every $p \in \Delta$. Hence $\frac{\alpha}{|\alpha|}$ is a NE for A . \square

Remark. Using (47) it is not hard to see that for every conditionally negative definite payoff matrix $A \in \mathbb{R}^{n \times n}$, there (uniquely) exists a Euclidean distance matrix D [c.f. 29] such that

$$\Lambda_A(p) = -\lambda(1 - \|p\|^2) - \frac{1}{2} p \cdot D p = \Lambda_{-\lambda\text{id} + \frac{1}{2}D}(p). \quad (80)$$

By [23, Thm 3.1] the mean of an invariant measure for the replicator semigroup constitutes a Nash equilibrium, say p^* , for A . But then in the light of Corollary 19 and in particular (78), the term $-\lambda(1 - \|p\|^2)$ in (80) corresponds to a Dirichlet distribution with parameter λp^* . This suggests that for general $A \in \Gamma^<$ invariant measures of stochastic replicator dynamics will be perturbations or generalizations of the Dirichlet family in which the matrix D enters as an additional parameter. Whether there exists an explicit expression for those measures is left as an interesting question for further investigations.

5.2. Wasserstein contractions for stochastic replicator dynamics

Corollary 19 in the previous subsection provided us with necessary and sufficient conditions for a stochastic replicator dynamic to attain the Dirichlet distribution Dir_α as invariant measure.

Another diffusion process on the Aitchison simplex for which Dir_α is invariant (in fact reversible) is the Langevin equation

$$dY_t = \ominus \nabla^A V(Y_t) dt \oplus dX_t, \quad (81)$$

with

$$V(p) = - \sum_{i=1}^n \alpha_i \ln p_i.$$

Recall that W_A is the natural Wasserstein distance on $(\Delta, \langle \cdot, \cdot \rangle_A)$ with cost d_A^2 (see (28)). Now let us also introduce

$$W_\Delta^2(\mu, \nu) := \inf_{\pi \in \Pi(\mu, \nu)} \int_{\Delta \times \Delta} \|p - q\|^2 \pi(dp dq). \quad (82)$$

The following proposition is the motivation for our subsequent investigations.

Proposition 20. *Consider the Langevin diffusion $Y = (Y_t)$ as given through (81) and denote by $\mu_t := Law(Y_t)$. Then,*

$$W_A(\mu_t, Dir_\alpha) \leq W_A(\mu_0, Dir_\alpha) \quad (83)$$

and moreover,

$$W_\Delta(\mu_t, Dir_\alpha) \leq e^{-|\alpha|t} W_A(\mu_0, Dir_\alpha). \quad (84)$$

Proof. We first show that $V(p) = - \sum \alpha_i \ln p_i$ is a convex function on the Aitchison simplex. Since $\nabla^g V(p) = |\alpha|p - \alpha$, we have

$$\langle \nabla^A V(p) \ominus \nabla^A V(q), p \ominus q \rangle_A = |\alpha|(p - q, \text{clr}(p) - \text{clr}(q)).$$

Now replacing $p = \text{sfm}(x)$ and $q = \text{sfm}(y)$ yields

$$\langle \nabla^A V(p) \ominus \nabla^A V(q), p \ominus q \rangle_A = |\alpha|(\text{sfm}(x) - \text{sfm}(y), x - y) \geq 0$$

by monotonicity of the softmax function ([c.f. 21]) and hence V is convex. Now take two solutions Y and Y' to (37), both driven by the same noise X . Then

$$\frac{d}{dt} \|Y_t \ominus Y'_t\|_A^2 = -2 \langle \nabla^A V(Y_t) \ominus \nabla^A V(Y'_t), Y_t \ominus Y'_t \rangle_A \leq 0$$

which immediately implies (83).

We move on to the ‘mixed’ Wasserstein contraction claimed in (84). First observe that sfm satisfies a stronger property than just being monotone, namely the softmax function is *co-coercive* [c.f. 21], i.e.

$$(\text{sfm}(x) - \text{sfm}(y), x - y) \geq \|\text{sfm}(x) - \text{sfm}(y)\|^2, \quad x, y \in \mathbb{R}^n. \quad (85)$$

Therefore,

$$\frac{d}{dt} \|Y_t \ominus Y'_t\|_A^2 \leq -2|\alpha| \|Y_t - Y'_t\|^2$$

and using (9) we find

$$\|Y_t - Y'_t\|_2^2 \leq \|Y_t \ominus Y'_t\|_A^2 \leq \|Y_0 \ominus Y'_0\|_A^2 - 2|\alpha| \int_0^t \|Y_s - Y'_s\|_2^2 ds.$$

Thus, by Gronwall's inequality

$$\|Y_t - Y'_t\|_2^2 \leq e^{-2|\alpha|t} \|Y_0 \ominus Y'_0\|_A^2,$$

which yields the claim after minimising on both sides over all couplings. \square

Of course, the result just proven is actually stronger than the inequalities stated in (83) and (84). Indeed, within these two estimates we may replace Dir_α by any other (law of a) solution to (81), say (μ'_t) . Then Proposition 20 asserts that with respect to W_Δ such laws attract exponentially fast, irrespective of the initial data.

The question we pose now is: can we find payoff matrices A which enforce a synchronization in the relaxation to equilibrium between replicator diffusion and the Langevin dynamic (81)? Or in other words, for which payoff matrices can we monitor for stochastic replicator dynamics the same contraction behavior as the one in Proposition 20?

In order to answer this question, let us first dwell upon the deterministic setting.

Theorem 21. *Let $A \in \mathbb{R}^{n \times n}$ be a payoff matrix. The following four statements are equivalent:*

(i) *For every $p, q \in \Delta$ let $(p(t))$ and $(q(t))$ be two solutions to the deterministic replicator equation (36) starting in p and q , respectively. Then,*

$$\|p(t) \ominus q(t)\|_A \leq \|p \ominus q\|_A, \quad t \geq 0 \tag{86}$$

(ii) *The map $p \mapsto \ominus\theta(p) = \theta(-p)$ is monotone, i.e.*

$$\langle \theta(p) \ominus \theta(q), p \ominus q \rangle_A \leq 0, \quad p, q \in \Delta. \tag{87}$$

(iii) *For every $p \in \Delta$ and $h \in H$*

$$h \cdot Ag^{-1}(p)h \leq 0. \tag{88}$$

(iv) *There exist $\lambda \geq 0$ and vectors $u, v \in \mathbb{R}^n$ such that*

$$A = -\lambda \text{id} + u \otimes \mathbf{1} + \mathbf{1} \otimes v \tag{89}$$

Proof. The chain of implications from (i) to (iii) is fairly standard and we only indicate the key ideas. (i) \implies (ii): differentiate $\|p(t) \ominus q(t)\|_A^2$ at $t = 0$. (ii) \implies (iii): observe that (87) is equivalent to

$$(A \text{sfm}(x) - A \text{sfm}(y), x - y) \leq 0, \quad x, y \in H. \tag{90}$$

Thus, for every $h \in H$ and $\tau > 0$, one has

$$(A \operatorname{sfm}(x + \tau h) - A \operatorname{sfm}(x), h) \leq 0.$$

Dividing by τ and taking $\tau \searrow 0$ yields (88).

We are now proving (iii) \implies (iv), thereby starting with the cases of dimensions $n = 2$. First observe that (88) necessitates that A is conditionally negative semi-definite, which can be seen by choosing $p = n^{-1}\mathbf{1}$. Therefore, if $n = 2$ and

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix},$$

we know $a + d \leq b + c$. Then, the claim follows by taking $2\lambda = b + c - (a + d)$, $2u = (b - d, c - a)$ and $2v = (a + c, b + d)$.

Let us now we consider the case $n \geq 3$ and write $a^i := A\epsilon_i$ for the i -th column of A . Notice that by continuity (88) holds for all $p \in \bar{\Delta}$. Testing with $2p = \epsilon_i + \epsilon_j$, for $i \neq j$, we learn that A must satisfy the peculiar monotonicity-like property

$$(h_i - h_j) \sum_{k=1}^n (a_{ki} - a_{kj}) h_k \leq 0 \quad (91)$$

for all $h \in H$ and $i \neq j$. By continuity we infer

$$h \in H \text{ with } h_i = h_j \implies (A^\top h)_i = (A^\top h)_j. \quad (92)$$

Equivalent to the previous implication is the fact that $h \in \{\mathbf{1}, \epsilon_i - \epsilon_j\}^\perp$ entails $h \in \langle a^i - a^j \rangle^\perp$. Thus, we can find scalars s_{ij}, t_{ij} such that

$$a^i - a^j = s_{ij}\mathbf{1} + t_{ij}(\epsilon_i - \epsilon_j). \quad (93)$$

Then, if $n = 3$ writing $a^i - a^j = a^i - a^k + a^k - a^j$ for distinct i, j, k and since $\mathbf{1} = \epsilon_i + \epsilon_j + \epsilon_k$ is follows

$$(\tilde{s} + t_{ij} - t_{ik})\epsilon_i + (\tilde{s} + t_{kj} - t_{ij})\epsilon_j + (\tilde{s} + t_{ik} - t_{kj})\epsilon_k = 0, \quad (94)$$

where $\tilde{s} := (s_{ij} - s_{ik} + s_{kj})$. By linear independence we deduce that $t_{ij} =: t$ must not depend on the indices and $\tilde{s} = 0$, which moreover entails $s_{ij} = v_i - v_j$ for some $v \in \mathbb{R}^3$. Likewise, for $n \geq 4$ the linear independence of $\mathbf{1}, \epsilon_i - \epsilon_k, \epsilon_i - \epsilon_j, \epsilon_j - \epsilon_k$ for pairwise distinct i, j, k , yields independence of t_{ij} on the indices as well as $s_{ij} = v_i - v_j$ for some $v \in \mathbb{R}^n$. At last, consider the vector $u := a^i - t\epsilon_i - v\mathbf{1}$ and observe that it does not depend on i . Thus, it follows $A = t \operatorname{id} + u \otimes \mathbf{1} + \mathbf{1} \otimes v$. Using (88) it is easy to see that $-\lambda := t \leq 0$, which proves the claim.

We end the proof by showing (iv) \implies (i). First notice, if A satisfies (89) then $(Ap)_i = -\lambda p_i + u_i + p \cdot v$. Hence, if $p(t)$ is a solution to the replicator equation (36)

$$\frac{d}{dt} \operatorname{clr}_i p(t) = -\lambda p_i(t) + u_i - \frac{1}{n} \sum_{k=1}^n u_k + \lambda. \quad (95)$$

But then it follows, that if $(q(t))$ is another solution to (36)

$$\frac{d}{dt} \|p(t) \ominus q(t)\|_A^2 = -2\lambda(\operatorname{clr}(p(t)) - \operatorname{clr}(q(t)), p(t) - q(t)) \leq 0$$

which yields (86). □

Note that for the case $\lambda = 0$, i.e. when $A = u \otimes \mathbf{1} + \mathbf{1} \otimes v$, we only know $A \in \Gamma^{\leq}$ from which one cannot infer the existence of ESS. However, due to the simple structure of such payoff matrices, one can easily give a full characterization of Nash equilibria and ESS. In fact, the following proposition follows straight forward from the definitions of NE and ESS.

Proposition 22. *Denote by $N(A)$ the set of all Nash equilibria of A . If $A = u \otimes \mathbf{1} + \mathbf{1} \otimes v$, then*

$$N(A) = \operatorname{argmax}_{p \in \bar{\Delta}} p \cdot u.$$

Moreover, such A has an ESS iff $N(A)$ is a singleton (i.e. when u has a distinct maximal entry).

Example. For the matrix

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix} \quad (96)$$

the pure strategy $p^* = (0, 0, 1)$ is the unique NE and ESS, because

$$A = (0, 3, 6) \otimes \mathbf{1} + \mathbf{1} \otimes (1, 2, 3).$$

If $\lambda > 0$ in (89), then $A \in \Gamma^<$ and we know A has an ESS. Clearly, if $u \in \langle \mathbf{1} \rangle$ then the barycenter e is an interior Nash equilibrium and also ESS. Otherwise, a necessary and sufficient condition on λ ensuring the existence of interior NE is given in

Proposition 23. *Let A satisfy (89) with $u \notin \langle \mathbf{1} \rangle$. Denote $|u| := u_1 + \dots + u_n$ and for $x \in \mathbb{R}$ set $x^- := -\min(x, 0)$. Then A has an interior Nash equilibrium if and only if $\lambda > |u| + n \max_i u_i^-$.*

Proof. Consider first the case when $u \in \mathbb{R}_{\geq 0}^n$. Assume $\lambda > |u|$. Then

$$\delta := \frac{1}{n} \left(1 - \frac{|u|}{\lambda} \right) > 0.$$

Define

$$p_i^* := \frac{u_i}{\lambda} + \delta.$$

Then $p^* \in \Delta$ and

$$-\lambda p_1^* + u_1 = \dots = -\lambda p_n^* + u_n, \quad (97)$$

whence p^* is an interior NE. For the converse direction, suppose p^* is an interior NE. Then p^* obeys (97). Let i^* be such that $u_{i^*} = \min_i u_i$. Using (97) it follows

$$n(-\lambda p_{i^*}^* + u_{i^*}) = -\lambda + |u|$$

and therefore

$$(1 - n p_{i^*}^*) \lambda = |u| - n u_{i^*} > 0.$$

Hence, $(1 - n p_{i^*}^*) \in (0, 1)$ and

$$\lambda \geq \frac{|u|}{1 - n p_{i^*}^*} > |u|.$$




Figure 2: Phase portraits of replicator dynamics for increasing choice of λ .

Now we drop the sign condition on u and consider some general $u \in \mathbb{R}^n$. Recall, that Nash equilibria for a payoff matrix A are invariant under the addition of a constant to any of the columns of A . Therefore,

$$N(A) = N(-\lambda \text{id} + u \otimes \mathbf{1}) = N(-\lambda \text{id} + \tilde{u} \otimes \mathbf{1}), \quad (98)$$

where $\tilde{u} \in \mathbb{R}_{\geq 0}^n$ is obtained from u by

$$\tilde{u}_i := u_i + \max_i u_i^- \quad (99)$$

and we can apply the result of the previous setting. \square

Figure 2 depicts phase portraits of replicator dynamics corresponding to A as in (96) for (a), $A - 5 \text{id}$ in (b), and $A - 10 \text{id}$ in (c). The corresponding ESS are $(0, 0, 1)$ in (a) and computed numerically using [47], $(0, \frac{1}{5}, \frac{4}{5})$ in (b) and $(\frac{1}{30}, \frac{1}{3}, \frac{19}{30})$ in (c).

Finally, observe that for such matrices one can update the estimate in (86) to an exponential contraction:

Corollary 24. *Let $A \in \mathbb{R}^{n \times n}$ be a payoff matrix with*

$$A = -\lambda \text{id} + u \otimes \mathbf{1} + \mathbf{1} \otimes v$$

for some vectors $u, v \in \mathbb{R}^n$ and $\lambda > 0$. Then, if $(p(t))$ and $(q(t))$ are solutions to the deterministic replicator equation (36), the following estimates are valid

$$\|p(t) \ominus q(t)\|_A^2 + 2\lambda \int_0^t \|p(s) - q(s)\|^2 ds \leq \|p \ominus q\|_A^2, \quad t \geq 0 \quad (100)$$

and

$$\|p(t) - q(t)\| \leq e^{-\lambda t} \|p \ominus q\|.$$

However, there does not exist any matrix $A \in \mathbb{R}^{n \times n}$ for which one can find some $\lambda > 0$ such that

$$\|p(t) \ominus q(t)\|_A \leq e^{-\lambda t} \|p \ominus q\|_A. \quad (101)$$

Proof. As in the last part of the proof of Theorem 21, we have

$$\frac{d}{dt} \|p(t) \ominus q(t)\|_A^2 = -2\lambda(\text{clr}(p(t)) - \text{clr}(q(t)), p(t) - q(t)) \leq -2\lambda \|p(t) - q(t)\|^2,$$

which immediately entails (100). On the other hand, since $\|p - q\| \leq \|p \ominus q\|_A$ we also find

$$\|p(t) - q(t)\|^2 \leq \|p \ominus q\|_A^2 - 2\lambda \int_0^t \|p(s) - q(s)\|^2 ds,$$

from which we infer (100) by Gronwall's inequality.

As for the second part of the claim let A be an arbitrary $n \times n$ payoff matrix and, aiming for a contraction, assume there is $\lambda > 0$ such that (101) holds true. Then, differentiation at $t = 0$ yields that $\ominus\theta$ ought to be λ -strongly monotone, i.e. for all $p, q \in \Delta$

$$\langle \theta(p) \ominus \theta(q), p \ominus q \rangle_A \leq -\lambda \|p \ominus q\|_A^2,$$

or equivalently

$$(A \text{sfm}(x) - A \text{sfm}(y), x - y) \leq -\lambda \|x - y\|^2, \quad x, y \in H.$$

Now rescale the previous inequality by replacing x and y by βx and βy for some $\beta \in \mathbb{R}$. Then,

$$\frac{1}{\beta} (A \text{sfm}(\beta x) - A \text{sfm}(\beta y), x - y) \leq -\lambda \|x - y\|^2 \quad (102)$$

Now choose, $x \neq y$ both such that they have one distinct maximal entry. Then [c.f. 21],

$$\lim_{\beta \rightarrow \infty} \text{sfm}(\beta x) = \text{argmax } x.$$

Therefore taking β to infinity in (102) yields

$$0 \leq -\lambda \|x - y\|^2$$

contradicting our assumption on the sign of λ . □

Whereas the previous findings might be of independent interest for evolutionary game theory, the main reason for treating in depth the deterministic dynamic is that these results have an immediate counterpart in the stochastic setting.

Theorem 25 (Wasserstein contractions for stochastic replicator dynamics). *Let $A \in \mathbb{R}^{n \times n}$ be a payoff matrix and Y and Y' be a solutions to the stochastic replicator equation (42) with $Y_0 \sim \mu_0$ and $Y'_0 \sim \mu'_0$. Denote for every $t \geq 0$ by μ_t and μ'_t the law of Y_t and Y'_t , respectively. Then,*

$$W_A(\mu_t, \mu'_t) \leq W_A(\mu_0, \mu'_0), \quad t \geq 0. \quad (103)$$

if and only if $A = -\lambda \text{id} + u \otimes \mathbf{1} + \mathbf{1} \otimes v$ for some vectors $u, v \in \mathbb{R}^n$ and $\lambda \geq 0$. If, $\lambda > 0$ then additionally

$$W_\Delta(\mu_t, \mu'_t) \leq e^{-\lambda t} W_A(\mu_0, \mu'_0), \quad t \geq 0. \quad (104)$$

However, there is no matrix A such that for some $\lambda > 0$

$$W_A(\mu_t, \mu'_t) \leq e^{-\lambda t} W_A(\mu_0, \mu'_0), \quad t \geq 0. \quad (105)$$

Proof. Using Theorem 11, we can represent a solution to the stochastic replicator equation Y by the SDE on the Aitchison simplex:

$$dY_t = \theta(Y_t)dt \oplus dX_t. \quad (106)$$

In particular, if Y and Y' are driven by the same Brownian motion B , then there is an Aitchison diffusion X , driving both Y and Y' in their Aitchison representation (106). Now if A satisfies (89), then by monotonicity of $\ominus\theta$ (c.f. 87) it follows

$$\frac{d}{dt} \|Y_t \ominus Y'_t\|_A^2 = 2\langle \theta(Y_t) \ominus \theta(Y'_t), Y_t \ominus Y'_t \rangle_A \leq 0 \quad (107)$$

and hence, $\mathbb{E}\|Y_t \ominus Y'_t\|_A^2 \leq \mathbb{E}\|Y_0 \ominus Y'_0\|_A^2$, which yields (103) after optimizing over all coupling on both sides of the inequality.

For the other direction, take (μ_t) and (μ'_t) with $\mu_0 = \delta_p$ and $\mu'_0 = \delta_q$, where $p, q \in \Delta$. By assumption, we have

$$W_{\mathbb{R}^{n-1}}(\text{ilr}_{\#}^{-1} \mu_t, \text{ilr}_{\#}^{-1} \mu'_t) \leq W_{\mathbb{R}^{n-1}}(\text{ilr}_{\#}^{-1} \delta_p, \text{ilr}_{\#}^{-1} \delta_q) \quad (108)$$

Since, the generator \hat{L} of the process $\hat{Y} = \text{ilr}(Y_t)$ is given by

$$\hat{L}f(x) = \frac{1}{2}\Delta f(x) + \nabla f(x) \cdot \hat{\theta}(x)$$

it is well-known [c.f. 10, 31, 46] that (108) implies (indeed is equivalent to)

$$(\hat{\theta}(x) - \hat{\theta}(y), x - y) \leq 0,$$

for all $x, y \in \mathbb{R}^{n-1}$. This in turn is equivalent to $\ominus\theta$ being monotone and thus, by Theorem 21 A obeys (89).

If we know that $\lambda > 0$, we can improve (109) to

$$\frac{d}{dt} \|Y_t \ominus Y'_t\|_A^2 \leq -2\lambda \|Y_t - Y'_t\|^2 \quad (109)$$

which yields (104) by the same arguments we used in the proof of Corollary 24. Finally, again by e.g. [31] the exponential contraction in (105) is equivalent to

$$(\hat{\theta}(x) - \hat{\theta}(y), x - y) \leq -\lambda \|x - y\|^2$$

and thus to $\ominus\theta$ being λ -strongly monotone, which is impossible, as we saw in Corollary 24. \square

Finally, let us relate the previous theorem to the results of Subsection 5.1. If A obeys (89) with $\lambda = 0$, then $\Lambda \equiv 0$, whence by Corollary 17 the Aitchison measure is invariant for the corresponding replicator diffusion Y . Moreover, in dimensions $n \geq 4$, Y must be transient. Indeed, transience holds also for $n = 2, 3$. To see this, note by Proposition 22, A can have interior NE only when $u \in \langle \mathbf{1} \rangle$, in which case the stochastic replicator dynamic degenerates to an undrifted Aitchison diffusion and is thus transient. Otherwise, if A has no interior NE, transience follows from [23, Cor. 4.16].

If $\lambda > 0$, then A satisfies in particular condition (i) of Corollary 19 (with $\lambda = |\alpha|$). Moreover A has an interior NE iff either $u \in \langle \mathbf{1} \rangle$ in which case $p^* = e$ or $\lambda > |u| + n \max_i u_i^-$ by Proposition 23. If $p^* \in \Delta$ is a NE for A , it follows that the Dirichlet distribution with parameter λp^* is invariant for Y . In this case we observe that the stochastic replicator dynamic obeys the same contraction behavior as the Langevin dynamic in Proposition 20.

ACKNOWLEDGEMENTS. The author is much obliged to Denis Serre, who provided the proof for the case $n \geq 4$ in (iii) \implies (iv) of Theorem 21. Much appreciated are also the critical remarks of Max von Renesse that helped to substantially improve the paper.

REFERENCES

- [1] <http://www.compositionaldata.com/>.
- [2] John Aitchison. The Statistical Analysis of Compositional Data. *Journal of the Royal Statistical Society. Series B (Methodological)*, 44(2):139–177, 1982.
- [3] John Aitchison. The statistical analysis of geochemical compositions. *Journal of the International Association for Mathematical Geology*, 16(6):531–564, Aug 1984.
- [4] Luigi Ambrosio, Nicola Gigli, and Giuseppe Savaré. *Gradient flows: in metric spaces and in the space of probability measures*. Springer Science & Business Media, 2008.
- [5] N. Ay, J. Jost, H.V. Lê, and L. Schwachhöfer. *Information Geometry*. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge / A Series of Modern Surveys in Mathematics. Springer International Publishing, 2017.
- [6] D. Bakry, I. Gentil, and M. Ledoux. *Analysis and Geometry of Markov Diffusion Operators*. Grundlehren der mathematischen Wissenschaften. Springer International Publishing, 2013.
- [7] Carles Barcel-Vidal, Josep Antoni Martí-Fernández, and Glòria Mateu-Figueras. *Compositional Differential Calculus on the Simplex*, chapter 13, pages 176–190. John Wiley & Sons, Ltd, 2011.
- [8] Dean Billheimer, Peter Guttorp, and William F. Fagan. Statistical Interpretation of Species Composition. *Journal of the American Statistical Association*, 96(456):1205–1214, 2001.
- [9] Vladimir I Bogachev, Nicolai V Krylov, Michael Röckner, and Stanislav V Shaposhnikov. *Fokker-Planck-Kolmogorov Equations*, volume 207. American Mathematical Soc., 2015.

- [10] François Bolley, Ivan Gentil, and Arnaud Guillin. Convergence to equilibrium in wasserstein distance for fokker–planck equations. *Journal of Functional Analysis*, 263(8):2430–2457, 2012.
- [11] G. Da Prato and J. Zabczyk. *Ergodicity for Infinite Dimensional Systems*. London Mathematical Society Lecture Note Series. Cambridge University Press, 1996.
- [12] Giuseppe Da Prato. *An introduction to infinite-dimensional analysis*. Springer Science & Business Media, 2006.
- [13] Manfred Eigen. Selforganization of matter and the evolution of biological macromolecules. *Naturwissenschaften*, 58(10):465–523, 1971.
- [14] Manfred Eigen and Peter Schuster. *The hypercycle: a principle of natural self-organization*. Springer Science & Business Media, 2012.
- [15] Ionas Erb and Nihat Ay. The information-geometric perspective of compositional data analysis, 2020.
- [16] Matthias Erbar. The heat equation on manifolds as a gradient flow in the Wasserstein space. *Ann. Inst. H. Poincaré Probab. Statist.*, 46(1):1–23, 02 2010.
- [17] Wong Eugene and Zakai Moshe. On the relation between ordinary and stochastic differential equations. *International Journal of Engineering Science*, 3(2):213–229, 1965.
- [18] E Robert Fernholz. Stochastic portfolio theory. In *Stochastic Portfolio Theory*, pages 1–24. Springer, 2002.
- [19] Peter Friz and Martin Hairer. A course on rough paths. *Preprint*, 2014.
- [20] Drew Fudenberg and Christopher Harris. Evolutionary dynamics with aggregate shocks. *Journal of Economic Theory*, 57(2):420–441, 1992.
- [21] Bolin Gao and Lacra Pavel. On the Properties of the Softmax Function with Application in Game Theory and Reinforcement Learning. *arXiv preprint arXiv:1704.00805*, 2017.
- [22] Marc Harper. Information geometry and evolutionary game theory. *arXiv preprint arXiv:0911.1383*, 2009.
- [23] Josef Hofbauer, Lorens A Imhof, et al. Time averages, recurrence and transience in the stochastic replicator dynamics. *The Annals of Applied Probability*, 19(4):1347–1368, 2009.
- [24] Josef Hofbauer and Karl Sigmund. *Evolutionary games and population dynamics*. Cambridge university press, 1998.
- [25] Julian Hofrichter, Jürgen Jost, and Tat Dat Tran. *Information geometry and population genetics*. Springer, 2017.

- [26] Nobuyuki Ikeda and Shinzo Watanabe. *Stochastic differential equations and diffusion processes*, volume 24. Elsevier, 2014.
- [27] Lorens A Imhof. The long-run behavior of the stochastic replicator dynamics. *The Annals of Applied Probability*, 15(1B):1019–1045, 2005.
- [28] Richard Jordan, David Kinderlehrer, and Felix Otto. The variational formulation of the Fokker–Planck equation. *SIAM journal on mathematical analysis*, 29(1):1–17, 1998.
- [29] Nathan Krislock and Henry Wolkowicz. Euclidean distance matrices and applications. In *Handbook on semidefinite, conic and polynomial optimization*, pages 879–914. Springer, 2012.
- [30] Terry J Lyons, Michael Caruana, and Thierry Lévy. *Differential equations driven by rough paths*. Springer, 2007.
- [31] Luca Natile, Mark A Peletier, and Giuseppe Savaré. Contraction of general transportation costs along solutions to Fokker–Planck equations with monotone drifts. *Journal de mathématiques pures et appliquées*, 95(1):18–35, 2011.
- [32] Charles M Newman, A Larry Wright, et al. An invariance principle for certain dependent sequences. *The Annals of Probability*, 9(4):671–675, 1981.
- [33] Martin A. Nowak. *Evolutionary dynamics*. Harvard University Press, 2006.
- [34] Felix Otto. The geometry of dissipative evolution equations: The porous medium equation. *Communications in Partial Differential Equations*, 26(1-2):101–174, 2001.
- [35] Grigorios A. Pavliotis. *Stochastic processes and applications: diffusion processes, the Fokker-Planck and Langevin equations*, volume 60. Springer, 2014.
- [36] Vera Pawlowsky-Glahn, Juan José Egozcue, and Raimon Tolosana Delgado. Lecture notes on compositional data analysis. 2007.
- [37] Vera Pawlowsky-Glahn et al. Statistical modeling on coordinates, 2003.
- [38] Ross G. Pinsky. *Positive Harmonic Functions and Diffusion*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1995.
- [39] Siavash Shahshahani. *A new mathematical framework for the study of linkage and selection*. American Mathematical Soc., 1979.
- [40] Cosma Rohilla Shalizi et al. Dynamics of bayesian updating with dependent data and misspecified models. *Electronic Journal of Statistics*, 3:1039–1074, 2009.
- [41] Karl Sigmund. A survey of replicator equations. In *Complexity, Language, and Life: Mathematical Approaches*, pages 88–104. Springer, 1986.
- [42] J Maynard Smith and George R Price. The logic of animal conflict. *Nature*, 246(5427):15, 1973.

- [43] D.W. Stroock and K. Itô. *Markov Processes from K. Itô's Perspective*. Academic Search Complete. Princeton University Press, 2003.
- [44] Peter D Taylor and Leo B Jonker. Evolutionary stable strategies and game dynamics. *Mathematical biosciences*, 40(1-2):145–156, 1978.
- [45] K.G. van den Boogaart and R. Tolosana-Delgado. *Analyzing Compositional Data with R*. Use R! Springer Berlin Heidelberg, 2013.
- [46] Max-K von Renesse and Karl-Theodor Sturm. Transport inequalities, gradient estimates, entropy and ricci curvature. *Communications on pure and applied mathematics*, 58(7):923–940, 2005.
- [47] E. Dokumaci W. H. Sandholm and F. Franchetti. *Dynamo: Diagrams for evolutionary game dynamics.*, 2012.
- [48] Bin Wu, Chaitanya S Gokhale, Matthijs van Veelen, Long Wang, and Arne Traulsen. Interpretations arising from Wrightian and Malthusian fitness under strong frequency dependent selection. *Ecology and evolution*, 3(5):1276–1280, 2013.