

# On Weighted Extropies

Narayanaswamy Balakrishnan<sup>1</sup>, Francesco Buono<sup>2</sup> and Maria Longobardi<sup>3</sup>

<sup>1</sup> Department of Mathematics and Statistics

McMaster University, Hamilton, Ontario, Canada

bala@mcmaster.ca

<sup>2</sup> Dipartimento di Matematica e Applicazioni "Renato Caccioppoli"

Universit degli Studi di Napoli Federico II, Naples, Italy

francesco.buono3@unina.it

<sup>3</sup> Dipartimento di Biologia

Universit degli Studi di Napoli Federico II, Naples, Italy

malongob@unina.it

*Keywords:* Extropy; Residual lifetime; Past lifetime; Weighted extropy; Bivariate extropy.

*AMS Subject Classification:* 62N05, 62B10.

## Abstract

The extropy is a measure of information introduced by Lad et al. (2015) as dual to entropy. As the entropy, it is a shift-independent information measure. We introduce here the notion of weighted extropy, a shift-dependent information measure which gives higher weights to large values of observed random variables. We also study the weighted residual and past extropies as weighted versions of extropy for residual and past lifetimes. Bivariate versions extropy and weighted extropy are also provided. Several examples are presented through out to illustrate the various concepts introduced here.

# 1 Introduction

## 1.1 Hazard and reversed hazard rates

Let  $X$  be a non-negative absolutely continuous random variable with probability density function (pdf)  $f$ , cumulative distribution function (cdf)  $F$  and survival function  $\bar{F}$ . In reliability theory, hazard rate function of  $X$  (also known as the force of mortality or the failure rate) where  $X$  is the lifetime of a system or a component has found many key application. In the same way the reversed hazard rate of  $X$  has also attracted much attention in the literature. In a certain sense, it is the dual function of hazard rate and it bears some interesting features useful in reliability analysis; see Block and Savits (1998) and Finkelstein (2002).

We define, for  $x$  such that  $\bar{F}(x) > 0$ , the hazard rate function of  $X$  at  $x$ ,  $r(x)$ , as

$$\begin{aligned} r(x) &= \lim_{\Delta x \rightarrow 0^+} \frac{\mathbb{P}(x < X \leq x + \Delta x | X > x)}{\Delta x} \\ &= \frac{1}{\bar{F}(x)} \lim_{\Delta x \rightarrow 0^+} \frac{\mathbb{P}(x < X \leq x + \Delta x)}{\Delta x} = \frac{f(x)}{\bar{F}(x)}. \end{aligned}$$

Analogously, we define, for  $x$  such that  $F(x) > 0$ , the reversed hazard rate function of  $X$  at  $x$ ,  $q(x)$ , as

$$\begin{aligned} q(x) &= \lim_{\Delta x \rightarrow 0^+} \frac{\mathbb{P}(x - \Delta x < X \leq x | X \leq x)}{\Delta x} \\ &= \frac{1}{F(x)} \lim_{\Delta x \rightarrow 0^+} \frac{\mathbb{P}(x - \Delta x < X \leq x)}{\Delta x} = \frac{f(x)}{F(x)}. \end{aligned}$$

The hazard rate  $r(x)$  can be interpreted as the rate of instantaneous failure occurring immediately after the time point  $x$ , given that the unit has survived up to time  $x$ . Similarly, the reversed hazard rate  $q(x)$  can be interpreted as the rate of instantaneous failure occurring immediately before the time point  $x$ , given that the unit has not survived time  $x$ . The hazard rate function and the reversed hazard rate function uniquely determine the distribution of  $X$ ; see Barlow and Proschan (1996) for details.

## 1.2 Residual and past lifetimes

In reliability theory, the residual and past lifetimes have a great importance. The residual lifetime of  $X$  at time  $t$  is defined as  $X_t = (X|X > t)$ , which is a random variable taking values in  $(t, +\infty)$  with pdf  $f_{X_t}(x) = \frac{f(x)}{F(t)}$  and survival function  $\overline{F}_{X_t}(x) = \frac{\overline{F}(x)}{F(t)}$ . The past lifetime of  $X$  at time  $t$  is defined as  ${}_tX = (X|X \leq t)$ , which is a random variable that takes values in  $(0, t]$  with pdf  $f_{{}_tX}(x) = \frac{f(x)}{F(t)}$  and distribution function  $F_{{}_tX}(x) = \frac{F(x)}{F(t)}$ .

## 1.3 Entropy, weighted entropy and extropy

The differential entropy, or Shannon entropy, of a random variable  $X$  is a basic concept as a measure of discrimination and information and is defined as

$$H(X) = -\mathbb{E}[\log f(X)] = -\int_0^{+\infty} f(x) \log f(x) dx, \quad (1)$$

where  $\log$  is the natural logarithm; see Shannon (1948).

Lad et al. (2015) introduced the concept of extropy as dual to entropy and it facilitates the comparison of uncertainties of two random variables. If the extropy of  $X$  is less than that of another variable  $Y$ , then  $X$  is said to have more uncertainty than  $Y$ . For a non-negative absolutely continuous random variable  $X$ , the extropy is defined as

$$J(X) = -\frac{1}{2}\mathbb{E}[f(X)] = -\frac{1}{2}\int_0^{+\infty} f^2(x) dx. \quad (2)$$

As a measure of information, the Shannon entropy is position-free, i.e., a random variable  $X$  has the same Shannon entropy of  $X+b$ , for any  $b \in \mathbb{R}$ . To avoid this problem, the concept of weighted entropy has been introduced (see Di Crescenzo and Longobardi (2006)) as

$$H^w(X) = -\mathbb{E}[X \log f(X)] = -\int_0^{+\infty} x f(x) \log f(x) dx. \quad (3)$$

To measure the uncertainty about the residual lifetime of  $X$  at time  $t$ , Ebrahimi (1996) introduced the residual entropy as

$$H(X_t) = -\int_t^{+\infty} \frac{f(x)}{F(t)} \log \frac{f(x)}{F(t)} dx, \quad (4)$$

and it is the differential entropy of the residual lifetime  $X_t$ . It is also possible to study the uncertainty about the past lifetime [Di Crescenzo and Longobardi (2002)] defined by

$$H({}_tX) = - \int_0^t \frac{f(x)}{F(t)} \log \frac{f(x)}{F(t)} dx, \quad (5)$$

and it is the differential entropy of the past lifetime  ${}_tX$ .

Analogous to residual entropy, Qiu (2017) defined the extropy for residual lifetime  $X_t$ , called the residual extropy at time  $t$ , as

$$J(X_t) = -\frac{1}{2} \int_0^{+\infty} f_{X_t}^2(x) dx = -\frac{1}{2\overline{F}^2(t)} \int_t^{+\infty} f^2(x) dx. \quad (6)$$

In analogous manner, Krishnan et al. (2020) and Kamari and Buono (2020) studied the past extropy defined as

$$J({}_tX) = -\frac{1}{2} \int_0^{+\infty} f_{{}_tX}^2(x) dx = -\frac{1}{2F^2(t)} \int_0^t f^2(x) dx. \quad (7)$$

Di Crescenzo and Longobardi (2006) discussed weighted versions of the residual and past entropies. The weighted residual entropy is defined as

$$H^w(X_t) = - \int_t^{+\infty} x \frac{f(x)}{\overline{F}(t)} \log \frac{f(x)}{\overline{F}(t)} dx, \quad (8)$$

while the weighted past entropy is defined as

$$H^w({}_tX) = - \int_0^t x \frac{f(x)}{F(t)} \log \frac{f(x)}{F(t)} dx. \quad (9)$$

Asadi and Zohrevand (2007) and Di Crescenzo and Longobardi (2009a) introduced the dynamic cumulative residual entropy and the dynamic cumulative entropy, respectively. Sathar and Nair (2019) introduced and studied the dynamic survival extropy  $J_s(X_t)$  defined as

$$J_s(X_t) = -\frac{1}{2\overline{F}^2(t)} \int_t^{+\infty} \overline{F}^2(x) dx. \quad (10)$$

Recently, various authors have discussed different versions of entropy and extropy and their applications, see, for example, Rao et al. (2004) for cumulative residual entropy, Di Crescenzo and Longobardi (2009b, 2013) for cumulative entropy, and Jahanshani et al. (2019) for cumulative residual extropy.

## 1.4 Scope of this paper

The rest of this paper is organized as follows. In Section 2, we introduce and study the weighted extropy analyzing some of its properties and also present some examples. In Section 3, we introduce the bivariate version of extropy and its weighted form. In section 4, we define and study the weighted residual and past extropies; we present some characterization results and also bounds for these measures of information. Finally, in Section 5, some concluding remarks are made.

## 2 Weighted extropy

Analogous to the weighted entropy, we introduce the concept of weighted extropy as

$$J^w(X) = -\frac{1}{2}\mathbb{E}[Xf(X)] = -\frac{1}{2}\int_0^{+\infty} xf^2(x)dx, \quad (11)$$

which can also be alternatively expressed as

$$J^w(X) = -\frac{1}{2}\int_0^{+\infty} f^2(x)\int_0^x dydx = -\frac{1}{2}\int_0^{+\infty} dy\int_y^{+\infty} f^2(x)dx. \quad (12)$$

We now present two examples of distributions with the same extropy, but different weighted extropy. In the first example, we note that the weighted extropy is indeed shift-dependent.

**Example 1.** Let  $X$  and  $Y$  be two random variables such that  $X \sim U(0, b)$ ,  $Y \sim U(a, a + b)$ , where  $a, b > 0$ . We have  $f_X(x) = \frac{1}{b}$ , for  $x \in (0, b)$ , and  $f_Y(x) = \frac{1}{b}$ , for  $x \in (a, a + b)$ , and then

$$J(X) = -\frac{1}{2}\int_0^b \frac{1}{b^2}dx = -\frac{1}{2b},$$

$$J(Y) = -\frac{1}{2}\int_a^{a+b} \frac{1}{b^2}dx = -\frac{1}{2b},$$

i.e.,  $X$  and  $Y$  have the same extropy, but they have different weighted extropy as seen below:

$$J^w(X) = -\frac{1}{2}\int_0^b x\frac{1}{b^2}dx = -\frac{1}{2b^2}\frac{b^2}{2} = -\frac{1}{4},$$

$$J^w(Y) = -\frac{1}{2}\int_a^{a+b} x\frac{1}{b^2}dx = -\frac{1}{2b^2}\frac{(a+b)^2 - a^2}{2} = -\frac{b^2 + 2ab}{4b^2} = -\frac{b + 2a}{4b},$$

and so if  $a \neq 0$ , i.e.,  $X$  and  $Y$  are not identically distributed, then  $J^w(X) \neq J^w(Y)$ .

**Example 2.** Let  $X$  be a random variable with piece-wise constant pdf

$$f(x) = \sum_{k=1}^n c_k \mathbf{1}_{[k-1,k)}(x),$$

where  $c_k \geq 0$ ,  $k = 1, \dots, n$ ,  $\sum_{k=1}^n c_k = 1$  and  $\mathbf{1}_{[k-1,k)}(x)$  is the indicator function of  $x$  on the interval  $[k-1, k)$ . Then, the entropy and the weighted entropy of  $X$  are

$$\begin{aligned} J(X) &= -\frac{1}{2} \int_0^n \sum_{k=1}^n c_k^2 \mathbf{1}_{[k-1,k)}(x) dx = -\frac{1}{2} \sum_{k=1}^n \int_{k-1}^k c_k^2 dx = -\frac{1}{2} \sum_{k=1}^n c_k^2, \\ J^w(X) &= -\frac{1}{2} \int_0^n x \sum_{k=1}^n c_k^2 \mathbf{1}_{[k-1,k)}(x) dx = -\frac{1}{2} \sum_{k=1}^n \int_{k-1}^k x c_k^2 dx \\ &= -\frac{1}{2} \sum_{k=1}^n c_k^2 \frac{k^2 - (k-1)^2}{2} = -\frac{1}{4} \sum_{k=1}^n c_k^2 (2k-1). \end{aligned}$$

Because we obtain different distributions through a permutation of  $c_1, \dots, c_n$ , we observe that they have the same entropy, but different weighted entropy (except in special cases).

We now present an example of distributions with the same weighted entropy, but different entropy.

**Example 3.** Let  $X$  be a random variable such that  $X \sim U(0, b)$ , with  $b > 0$ . In Example 1 earlier, we observed that  $J(X) = -\frac{1}{2b}$  and  $J^w(X) = -\frac{1}{4}$ , and so the entropy depends on  $b$  while the weighted entropy does not. Thus, if we consider  $Y \sim U(0, a)$ , with  $a > 0$  and  $a \neq b$ , we have random variables  $X$  and  $Y$  with the same weighted entropy, but different entropy.

Let us now evaluate the weighted entropy of some random variables.

**Example 4.** (a) Suppose  $X$  is exponentially distributed with parameter  $\lambda$ . Then,

$$J^w(X) = -\frac{1}{2} \int_0^{+\infty} x \lambda^2 e^{-2\lambda x} dx = \frac{1}{2} \left[ \frac{\lambda x}{2} e^{-2\lambda x} \Big|_0^{+\infty} - \int_0^{+\infty} \frac{\lambda}{2} e^{-2\lambda x} dx \right] = -\frac{1}{8}.$$

(b) Suppose  $X$  is uniformly distributed over  $(a, b)$ . Then,

$$J^w(X) = -\frac{1}{2} \int_a^b x \frac{1}{(b-a)^2} dx = -\frac{1}{2(b-a)^2} \frac{b^2 - a^2}{2} = -\frac{1}{4} \frac{b+a}{b-a}.$$

Observe that in this case the weighted extropy can be expressed as the product

$$J^w(X) = J(X)\mathbb{E}(X),$$

where  $\mathbb{E}(X) = \frac{a+b}{2}$  and  $J(X) = -\frac{1}{2(b-a)}$ . Then,  $J(X) \leq J^w(X)$  if, and only if,  $\mathbb{E}(X) \leq 1$ , due to the fact that extropy and weighted extropy are non-positive.

Let us now look for values of  $a$  and  $b$  such that the weighted extropy for  $Exp(\lambda)$  and  $U(a, b)$  are the same. We then have

$$-\frac{1}{8} = -\frac{1}{4} \frac{b+a}{b-a} \iff 2(b-a) = b+a \iff b = 3a.$$

Then,  $Exp(\lambda)$  and  $U(1, 3)$  have the same weighted extropy.

(c) Suppose  $X$  is gamma distributed with parameters  $\alpha$  and  $\beta$ , and with pdf

$$f(x) = \begin{cases} \frac{x^{\alpha-1} e^{-x/\beta}}{\beta^\alpha \Gamma(\alpha)}, & \text{if } x > 0 \\ 0, & \text{otherwise.} \end{cases}$$

Then, we have

$$\begin{aligned} J^w(X) &= -\frac{1}{2} \int_0^{+\infty} x \frac{x^{2\alpha-2} e^{-2x/\beta}}{\beta^{2\alpha} \Gamma^2(\alpha)} dx \\ &= -\frac{1}{2\beta^{2\alpha} \Gamma^2(\alpha)} \int_0^{+\infty} x^{2\alpha-1} e^{-2x/\beta} dx \\ &= -\frac{1}{2^{2\alpha+1}} \frac{\Gamma(2\alpha)}{\Gamma^2(\alpha)}, \end{aligned}$$

which is free of  $\beta$ .

(d) Suppose  $X$  is beta distributed with parameters  $\alpha$  and  $\beta$ , and with pdf

$$f(x) = \begin{cases} \frac{x^{\alpha-1} (1-x)^{\beta-1}}{B(\alpha, \beta)}, & \text{if } 0 < x < 1 \\ 0, & \text{otherwise,} \end{cases}$$

where

$$B(\alpha, \beta) = \int_0^1 x^{\alpha-1}(1-x)^{\beta-1} dx = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$$

is the complete beta function. Then, we have

$$\begin{aligned} J^w(X) &= -\frac{1}{2} \int_0^1 x \frac{x^{2\alpha-2}(1-x)^{2\beta-2}}{B^2(\alpha, \beta)} dx \\ &= -\frac{1}{2} \frac{B(2\alpha, 2\beta-1)}{B^2(\alpha, \beta)} \end{aligned}$$

if  $2\beta - 1 > 0$ , i.e.,  $\beta > \frac{1}{2}$ , but if  $0 < \beta \leq \frac{1}{2}$  we have  $J^w(X) = -\infty$ .

**Remark 1.** Let us now focus our attention on the summability of  $xf^2(x)$  on the support of  $X$ . If the support is unbounded, i.e.,  $(a, +\infty)$ , with  $a \geq 0$ , and the function is bounded, we have to investigate about the summability at infinity. Because  $\int_a^{+\infty} f(x)dx = 1$ , we have  $\lim_{x \rightarrow +\infty} f(x) = 0$  and  $f(x)$  is infinitesimal of higher order with respect to  $\frac{1}{x^{1+\varepsilon}}$ , for  $x \rightarrow +\infty$ , for some  $\varepsilon > 0$ . Then,  $xf^2(x)$  is infinitesimal of higher order with respect to  $\frac{1}{x^{1+2\varepsilon}}$ , for  $x \rightarrow +\infty$  and so it is integrable, i.e., the weighted extropy is finite. If the support and the density are unbounded, we also have to study the summability at  $a$ . Suppose  $a > 0$ . If  $\lim_{x \rightarrow a^+} f(x) = +\infty$ , by the normalization condition we know that  $f(x)$  is infinity of lower order with respect to  $\frac{1}{(x-a)^{1-\varepsilon}}$ , for  $x \rightarrow a^+$ , for some  $0 < \varepsilon < 1$ . Hence,  $xf^2(x)$  is infinity of lower order with respect to  $\frac{1}{(x-a)^{2-2\varepsilon}}$ , for  $x \rightarrow a^+$ , and so is integrable if  $\varepsilon \in (\frac{1}{2}, 1)$ . If  $a = 0$ , by the normalization condition, we know that  $f(x)$  is infinity of lower order with respect to  $\frac{1}{x^{1-\varepsilon}}$ , for  $x \rightarrow 0^+$ , for some  $0 < \varepsilon < 1$ . Hence,  $xf^2(x)$  is infinity of lower order with respect to  $\frac{1}{x^{1-2\varepsilon}}$ , for  $x \rightarrow 0^+$ , and so is integrable. If the support is bounded and  $f$  is bounded, then  $xf^2(x)$  is bounded and is integrable. If the support is bounded and  $f$  is unbounded, then we can refer to the previous cases. Observe that if the support is  $(0, +\infty)$ , the weighted extropy is always finite.

In the following theorem, we study weighted extropy under monotone transformation.

**Theorem 2.1.** *Let  $Y = \Phi(X)$ , with  $\Phi$  being strictly monotone, continuous and differentiable,*



with derivative  $\Phi'$ . Then, we have

$$J^w(Y) = \begin{cases} -\frac{1}{2} \int_0^{+\infty} \frac{\Phi(x)}{\Phi'(x)} f_X^2(x) dx, & \text{if } \Phi \text{ is strictly increasing} \\ -\frac{1}{2} \int_0^{+\infty} \frac{\Phi(x)}{|\Phi'(x)|} f_X^2(x) dx, & \text{if } \Phi \text{ is strictly decreasing.} \end{cases} \quad (13)$$

*Proof.* From (11), we have

$$J^w(Y) = -\frac{1}{2} \int_0^{+\infty} x \frac{f_X^2(\Phi^{-1}(x))}{(\Phi'(\Phi^{-1}(x)))^2} dx.$$

Let  $\Phi$  be strictly increasing. Then, with a change of variable in the above integral, we get

$$J^w(Y) = -\frac{1}{2} \int_0^{+\infty} \frac{\Phi(x)}{\Phi'(x)} f_X^2(x) dx,$$

giving the first expression in (13). If  $\Phi$  is strictly decreasing, we similarly obtain

$$J^w(Y_t) = -\frac{1}{2} \int_0^{+\infty} \frac{\Phi(x)}{|\Phi'(x)|} f_X^2(x) dx,$$

which is the second expression in (13).  $\square$

**Remark 2.** If in Theorem 2.1 we take  $\Phi(X) = F_X(X)$ , then we get

$$J^w(Y) = -\frac{1}{2} \int_0^{+\infty} F_X(x) f_X(x) dx = -\frac{1}{4},$$

which agrees with that of Uniform(0, 1) distribution, since it is known in this case that the probability integral transformation  $Y = F_X(X)$  is Uniform(0, 1).

In the following theorem, we obtain a lower bound for the weighted extropy of the sum of two independent random variables.

**Theorem 2.2.** *Let  $X$  and  $Y$  be two non-negative independent random variables with densities  $f_X$  and  $f_Y$ , respectively. Then,*

$$J^w(X + Y) \geq -2 (J(X)J^w(Y) + J^w(X)J(Y)). \quad (14)$$

*Proof.* Since  $X$  and  $Y$  are non-negative independent random variables, the density function of  $Z = X + Y$  is given, for  $z > 0$ , by

$$f_Z(z) = \int_0^z f_X(x) f_Y(z - x) dx.$$

Hence, the weighted extropy of  $Z$  is given by

$$J^w(Z) = -\frac{1}{2} \int_0^{+\infty} z \left[ \int_0^z f_X(x) f_Y(z-x) dx \right]^2 dz.$$

Using Jensen's inequality, we then have

$$\begin{aligned} J^w(Z) &\geq -\frac{1}{2} \int_0^{+\infty} z \int_0^z f_X^2(x) f_Y^2(z-x) dx dz \\ &= -\frac{1}{2} \int_0^{+\infty} f_X^2(x) \int_x^{+\infty} z f_Y^2(z-x) dz dx \\ &= -\frac{1}{2} \int_0^{+\infty} f_X^2(x) \int_0^{+\infty} (z+x) f_Y^2(z) dz dx \\ &= \int_0^{+\infty} f_X^2(x) (J^w(X) + xJ(Y)) dx \\ &= -2J(X)J^w(Y) - 2J(Y)J^w(X), \end{aligned}$$

as required. □

**Remark 3.** In particular, if  $X$  and  $Y$  are independent and identically distributed in Theorem 2.2, we simply deduce

$$J^w(X+Y) \geq -4J(X)J^w(X).$$

### 3 Bivariate version of extropy and weighted extropy

It is possible to introduce bivariate version of extropy. If  $X$  and  $Y$  are non-negative absolutely continuous random variables, the bivariate version of extropy, denoted by  $J(X, Y)$ , is defined as

$$J(X, Y) = \frac{1}{4} \mathbb{E}[f(X, Y)] = \frac{1}{4} \int_0^{+\infty} \int_0^{+\infty} f^2(x, y) dx dy, \quad (15)$$

where  $f(x, y)$  is the joint density of  $(X, Y)$ .

**Remark 4.** This measure can also be defined for a general  $k$ -dimensional vector. In that case, in the definition of  $J(X_1, X_2, \dots, X_k)$ , the multiplying factor for the integral will be  $(-\frac{1}{2})^k$ .

In the following proposition, we discuss how extropy changes under linear transformations and how bivariate version of extropy behaves in the case of independence.

**Proposition 3.1.** (i) Let  $X$  be a non-negative absolutely continuous random variable, and  $Y = aX + b$ , with  $a > 0, b \geq 0$ . Then,  $J(Y) = \frac{1}{a}J(X)$ ;

(ii) Let  $X, Y$  be non-negative absolutely continuous random variables. If  $X$  and  $Y$  are independent, then  $J(X, Y) = J(X)J(Y)$ .

*Proof.* (i) From  $Y = aX + b$ , we readily have  $f_Y(x) = \frac{1}{a}f_X\left(\frac{x-b}{a}\right)$ ,  $x > b$ . So,

$$J(Y) = -\frac{1}{2} \int_b^{+\infty} \frac{1}{a^2} f_X^2\left(\frac{x-b}{a}\right) dx = -\frac{1}{2} \int_0^{+\infty} \frac{1}{a} f_X^2(x) dx = \frac{1}{a} J(X).$$

(ii) If  $X$  and  $Y$  are independent, then  $f(x, y) = f_X(x)f_Y(y)$  and hence

$$\begin{aligned} J(X, Y) &= \frac{1}{4} \mathbb{E}[f(X, Y)] = \frac{1}{4} \mathbb{E}[f_X(X)f_Y(Y)] \\ &= \frac{1}{4} \mathbb{E}[f_X(X)] \mathbb{E}[f_Y(Y)] = J(X)J(Y). \end{aligned}$$

□

**Remark 5.** In Property (i) of Proposition 3.1, if we choose  $a = 1$ , we get the known property that extropy is invariant under translations.

In a spirit similar to that of the bivariate version of extropy, we can also introduce bivariate weighted extropy as

$$J^w(X, Y) = \frac{1}{4} \mathbb{E}[XY f(X, Y)] = \frac{1}{4} \int_0^{+\infty} \int_0^{+\infty} xy f^2(x, y) dx dy, \quad (16)$$

where  $X$  and  $Y$  are non-negative random variables with joint density function  $f(x, y)$ .

In the following proposition we discuss how weighted extropy behaves under linear transformations and the bivariate extropy in the case of independence.

**Proposition 3.2.** (i) Let  $X$  be a non-negative absolutely continuous random variable, and  $Y = aX + b$ , with  $a > 0, b \geq 0$ . Then,  $J^w(Y) = J^w(X) + \frac{b}{a}J(X)$ ;

(ii) Let  $X, Y$  be non-negative absolutely continuous random variables. If  $X$  and  $Y$  are independent, then  $J^w(X, Y) = J^w(X)J^w(Y)$ .

*Proof.* (i) From  $Y = aX + b$ , we have  $f_Y(x) = \frac{1}{a}f_X\left(\frac{x-b}{a}\right)$ ,  $x > b$ , and so

$$\begin{aligned} J^w(Y) &= -\frac{1}{2} \int_b^{+\infty} x \frac{1}{a^2} f_X^2\left(\frac{x-b}{a}\right) dx = -\frac{1}{2} \int_0^{+\infty} (ax+b) \frac{1}{a} f_X^2(x) dx \\ &= -\frac{1}{2} \int_0^{+\infty} x f_X^2(x) dx - \frac{1}{2} \frac{b}{a} \int_0^{+\infty} f_X^2(x) dx = J^w(X) + \frac{b}{a} J(X). \end{aligned}$$

(ii) If  $X$  and  $Y$  are independent, then  $f(x, y) = f_X(x)f_Y(y)$  and hence

$$\begin{aligned} J^w(X, Y) &= \frac{1}{4} \mathbb{E}[XY f(X, Y)] = \frac{1}{4} \mathbb{E}[XY f_X(X) f_Y(Y)] \\ &= \frac{1}{4} \mathbb{E}[X f_X(X)] \mathbb{E}[Y f_Y(Y)] = J^w(X) J^w(Y). \end{aligned}$$

□

**Remark 6.** In Part (i) of Proposition 3.2, if we choose  $b = 0$ , we see that the weighted extropy is invariant for proportional random variables, as we saw earlier in Example 3 in the case of uniform distribution.

**Example 5.** (a) Let us consider  $(X, Y)$  as a bivariate beta random variable with joint density function

$$f(x, y) = \frac{1}{B(\alpha, \beta, \gamma)} x^{\alpha-1} (y-x)^{\beta-1} (1-y)^{\gamma-1}, \quad 0 < x < y < 1, \quad \alpha, \beta, \gamma > 0,$$

where  $B(\alpha, \beta, \gamma) = \frac{\Gamma(\alpha)\Gamma(\beta)\Gamma(\gamma)}{\Gamma(\alpha+\beta+\gamma)}$  is the bivariate complete beta function. Then, we readily find the bivariate extropy as

$$\begin{aligned} J(X, Y) &= \frac{1}{4} \mathbb{E}[f(X, Y)] \\ &= \frac{1}{4B^2(\alpha, \beta, \gamma)} \iint_{0 < x < y < 1} x^{2\alpha-2} (y-x)^{2\beta-2} (1-y)^{2\gamma-2} dx dy \\ &= \frac{B(2\alpha-1, 2\beta-1, 2\gamma-1)}{4B^2(\alpha, \beta, \gamma)} \end{aligned}$$

provided  $\alpha, \beta, \gamma > \frac{1}{2}$ .

(b) For the above bivariate beta distribution, we readily find the bivariate weighted extropy to be

$$\begin{aligned}
J^w(X, Y) &= \frac{1}{4} \mathbb{E}[XY f(X, Y)] \\
&= \frac{1}{4B^2(\alpha, \beta, \gamma)} \iint_{0 < x < y < 1} x^{2\alpha-1} y (y-x)^{2\beta-2} (1-y)^{2\gamma-2} dx dy \\
&= \frac{1}{4B^2(\alpha, \beta, \gamma)} [B(2\alpha, 2\beta, 2\gamma-1) + B(2\alpha+1, 2\beta-1, 2\gamma-1)]
\end{aligned}$$

provided  $\alpha > 0$  and  $\beta, \gamma > \frac{1}{2}$ .

(c) In the special case of bivariate uniform distribution (i.e.,  $\alpha = \beta = \gamma = 1$ ), the above expressions readily reduce to

$$J(X, Y) = \frac{1}{2} \quad \text{and} \quad J^w(X, Y) = \frac{1}{8}.$$

## 4 Weighted residual and past extropies

In this section, we introduce and study weighted residual extropy and weighted past extropy.

**Definition 1.** Let  $X$  be a non-negative absolutely continuous random variable. For all  $t$  in the support of  $f$ , we define

(i) the weighted residual extropy of  $X$  at time  $t$  as

$$J^w(X_t) = -\frac{1}{2\overline{F}^2(t)} \int_t^{+\infty} x f^2(x) dx; \quad (17)$$

(ii) the weighted past extropy of  $X$  at time  $t$  as

$$J^w({}_tX) = -\frac{1}{2F^2(t)} \int_0^t x f^2(x) dx. \quad (18)$$

**Remark 7.** We notice that

$$\lim_{t \rightarrow 0^+} J^w(X_t) = \lim_{t \rightarrow +\infty} J^w({}_tX) = J^w(X). \quad (19)$$

In the following lemma, we evaluate the derivative of the weighted residual extropy and the derivative of the weighted past extropy.

**Lemma 4.1.** *Let  $X$  be a non-negative absolutely continuous random variable with weighted residual extropy  $J^w(X_t)$  and weighted past extropy  $J^w({}_tX)$ . Then,*

$$(i) \quad \frac{d}{dt}J^w(X_t) = \frac{r(t)}{2} [J^w(X_t) + tr(t)], \text{ where } r(t) \text{ is the hazard rate function of } X;$$

$$(ii) \quad \frac{d}{dt}J^w({}_tX) = -\frac{q(t)}{2} [J^w({}_tX) + tq(t)], \text{ where } q(t) \text{ is the reversed hazard rate function of } X.$$

*Proof.* (i) From the definitions of weighted residual extropy and hazard rate function, we have

$$\begin{aligned} \frac{d}{dt}J^w(X_t) &= -\frac{1}{4\overline{F}^3(t)}f(t) \int_t^{+\infty} xf^2(x)dx + \frac{1}{2\overline{F}^2(t)}tf^2(t) \\ &= \frac{r(t)}{2} [J^w(X_t) + tr(t)]; \end{aligned}$$

(ii) From the definition of weighted past extropy and reversed hazard rate function, we have

$$\begin{aligned} \frac{d}{dt}J^w({}_tX) &= \frac{1}{4F^3(t)}f(t) \int_0^t xf^2(x)dx - \frac{1}{2F^2(t)}tf^2(t) \\ &= -\frac{q(t)}{2} [J^w({}_tX) + tq(t)]. \end{aligned}$$

□

**Remark 8.** We may ask a natural question here whether the weighted residual extropy could be constant over the support of a non-negative absolutely continuous random variable. If  $J^w(X_t)$  is constant, then for all  $t > 0$ , we have

$$J^w(X_t) + tr(t) = 0$$

and then

$$\int_t^{+\infty} xf^2(x)dx = 2tf(t)\overline{F}(t).$$

Differentiating both sides we get

$$t f^2(t) = 2f(t)\overline{F}(t) + 2t f'(t)\overline{F}(t).$$

We know that  $r'(t) = \frac{f'(t)}{\overline{F}(t)} + r^2(t)$  and so dividing by  $\overline{F}^2(t)$  both sides of the above equality, we obtain

$$r'(t) = -\frac{r(t)}{t} + \frac{3}{2}r^2(t)$$

which is a Bernoulli differential equation with initial condition  $r(t_0) = r_0 > 0$ ,  $t_0 > 0$ . Upon solving this differential equation, we get

$$r(t) = \frac{2}{t(C - 3 \log t)},$$

where  $C = \frac{2}{t_0 r_0} + 3 \log t_0$ . We know that the hazard rate function is non-negative and this condition is satisfied if and only if  $t \leq e^C/3$ . Hence, the weighted residual extropy can not be constant over  $(0, +\infty)$ .

**Remark 9.** The weighted extropy, the weighted residual extropy and the weighted past extropy satisfy the following relationship:

$$J^w(X) = F^2(t)J^w({}_tX) + \overline{F}^2(t)J^w(X_t). \quad (20)$$

To see this, we consider

$$\begin{aligned} J^w(X) &= -\frac{1}{2} \int_0^{+\infty} x f^2(x) dx = -\frac{1}{2} \int_0^t x f^2(x) dx - \frac{1}{2} \int_t^{+\infty} x f^2(x) dx \\ &= -\frac{1}{2} F^2(t) \int_0^t x \frac{f^2(x)}{F^2(t)} dx - \frac{1}{2} \overline{F}^2(t) \int_t^{+\infty} x \frac{f^2(x)}{\overline{F}^2(t)} dx \\ &= F^2(t)J^w({}_tX) + \overline{F}^2(t)J^w(X_t). \end{aligned}$$

**Theorem 4.1.** *If  $X$  is a non-negative absolutely continuous random variable and if  $J^w(X_t)$  is increasing in  $t > 0$ , then  $J^w(X_t)$  uniquely determines the distribution of  $X$ .*

*Proof.* From Lemma 4.1, we have

$$\frac{d}{dt} J^w(X_t) = \frac{r(t)}{2} [J^w(X_t) + tr(t)].$$

Let us now introduce the following function:

$$g(x) = \frac{x}{2} [J^w(X_t) + tx] - \frac{d}{dt} J^w(X_t).$$

We know that  $g(r(t)) = 0$  and  $g(0) = -\frac{d}{dt} J^w(X_t) \leq 0$  because  $J^w(X_t)$  is increasing in  $t > 0$ . Moreover,  $\lim_{x \rightarrow +\infty} g(x) = +\infty$ . If we evaluate the derivative of  $g(x)$ , we could see that there is only one point at which it vanishes; in fact,

$$\frac{d}{dx} g(x) = \frac{1}{2} J^w(X_t) + tx$$

and so

$$\frac{d}{dx} g(x) = 0 \iff x = -\frac{1}{2t} J^w(X_t) (\geq 0).$$

Then,  $g(x) = 0$  has a unique solution and it is  $r(t)$ . From Barlow and Proschan (1996), we know that the hazard rate function uniquely determines the distribution and so  $J^w(X_t)$  also uniquely determines the distribution.  $\square$

**Theorem 4.2.** *If  $X$  is a non-negative absolutely continuous random variable and if  $J^w({}_tX)$  is decreasing in  $t > 0$ , then  $J^w({}_tX)$  uniquely determines the distribution of  $X$ .*

*Proof.* From Lemma 4.1, we have

$$\frac{d}{dt} J^w({}_tX) = -\frac{q(t)}{2} [J^w({}_tX) + tq(t)].$$

Let us now introduce the following function:

$$h(x) = \frac{x}{2} [J^w({}_tX) + tx] + \frac{d}{dt} J^w({}_tX).$$

We know that  $h(q(t)) = 0$  and  $h(0) = \frac{d}{dt} J^w({}_tX) \leq 0$  because  $J^w({}_tX)$  is decreasing in  $t > 0$ . Moreover,  $\lim_{x \rightarrow +\infty} h(x) = +\infty$ . If we evaluate the derivative of  $h(x)$ , we could see that there is only one point at which it vanishes; in fact,

$$\frac{d}{dx} h(x) = \frac{1}{2} J^w({}_tX) + tx$$

and so

$$\frac{d}{dx} h(x) = 0 \iff x = -\frac{1}{2t} J^w({}_tX) (\geq 0).$$



Then,  $h(x) = 0$  has a unique solution and it is  $q(t)$ . From Barlow and Proschan (1996), we know that the reversed hazard rate function uniquely determines the distribution and so  $J^w({}_tX)$  also uniquely determines the distribution.  $\square$

In the following two theorems, we obtain bounds for the weighted residual extropy and the weighted past extropy under the monotonicity of hazard rate and reversed hazard rate functions.

**Theorem 4.3.** *If the hazard rate function  $r$  is increasing, then*

$$J^w(X_t) \leq t r^2(t) J_s(X_t), \quad (21)$$

where  $J_s(X_t)$  is the dynamic survival extropy defined in (10).

*Proof.* From the definition of the weighted residual extropy, we have

$$J^w(X_t) = -\frac{1}{2\bar{F}^2(t)} \int_t^{+\infty} x f^2(x) dx = -\frac{1}{2\bar{F}^2(t)} \int_t^{+\infty} x r^2(x) \bar{F}^2(x) dx.$$

Because the hazard rate function is increasing, we have

$$\begin{aligned} -\frac{1}{2\bar{F}^2(t)} \int_t^{+\infty} x r^2(x) \bar{F}^2(x) dx &\leq -\frac{r^2(t)}{2\bar{F}^2(t)} \int_t^{+\infty} x \bar{F}^2(x) dx \\ &\leq -t \frac{r^2(t)}{2\bar{F}^2(t)} \int_t^{+\infty} \bar{F}^2(x) dx \\ &= t r^2(t) J_s(X_t). \end{aligned}$$

$\square$

**Theorem 4.4.** *If the reversed hazard rate function  $q$  is increasing in  $(0, T)$ , with  $T > t$ , then*

$$J^w({}_tX) \geq -t \frac{q^2(t)}{2}. \quad (22)$$

*Proof.* From the definition of the weighted past extropy, we have

$$J^w({}_tX) = -\frac{1}{2F^2(t)} \int_0^t x f^2(x) dx = -\frac{1}{2F^2(t)} \int_0^t x q^2(x) F^2(x) dx.$$

Because the reversed hazard rate function is increasing in  $(0, T)$ , we have

$$-\frac{1}{2F^2(t)} \int_0^t xq^2(x)F^2(x)dx \geq -\frac{q^2(t)}{2F^2(t)} \int_0^t xF^2(x)dx;$$

moreover, the distribution function is increasing and so

$$-\frac{q^2(t)}{2F^2(t)} \int_0^t xF^2(x)dx \geq -\frac{q^2(t)}{2} \int_0^t xdx = -t\frac{q^2(t)}{2}.$$

□

**Example 6.** For the exponential distribution with parameter  $\lambda > 0$ , the hazard function  $r(t) = \lambda$  is constant and so satisfies the conditions in Theorem 4.3. The weighted residual extropy is given by

$$J^w(X_t) = -\frac{1}{2e^{-2\lambda t}} \int_t^{+\infty} x\lambda^2 e^{-2\lambda x} dx = -\frac{\lambda t}{4} - \frac{1}{8},$$

while the upper bound given by Theorem 4.3 is

$$-t\frac{\lambda^2}{2e^{-2\lambda t}} \int_t^{+\infty} e^{-2\lambda x} dx = -\frac{\lambda t}{4},$$

and so (21) is fulfilled.

In the following theorem, we discuss weighted residual extropy and weighted past extropy under monotone transformation.

**Theorem 4.5.** *Let  $Y = \Phi(X)$ , with  $\Phi$  being strictly monotone, continuous and differentiable, with derivative  $\Phi'$ . Then, for all  $t > 0$ , we have*

$$J^w(Y_t) = \begin{cases} -\frac{1}{2F_X^2(\Phi^{-1}(t))} \int_{\Phi^{-1}(t)}^{+\infty} \frac{\Phi(x)}{\Phi'(x)} f_X^2(x) dx, & \text{if } \Phi \text{ is strictly increasing} \\ -\frac{1}{2F_X^2(\Phi^{-1}(t))} \int_0^{\Phi^{-1}(t)} \frac{\Phi(x)}{|\Phi'(x)|} f_X^2(x) dx, & \text{if } \Phi \text{ is strictly decreasing} \end{cases} \quad (23)$$

and

$$J^w({}_tY) = \begin{cases} -\frac{1}{2F_X^2(\Phi^{-1}(t))} \int_0^{\Phi^{-1}(t)} \frac{\Phi(x)}{\Phi'(x)} f_X^2(x) dx, & \text{if } \Phi \text{ is strictly increasing} \\ -\frac{1}{2F_X^2(\Phi^{-1}(t))} \int_{\Phi^{-1}(t)}^{+\infty} \frac{\Phi(x)}{|\Phi'(x)|} f_X^2(x) dx, & \text{if } \Phi \text{ is strictly decreasing.} \end{cases} \quad (24)$$

*Proof.* From (17), we have

$$J^w(Y_t) = -\frac{1}{2\overline{F}_X^2(\Phi^{-1}(t))} \int_t^{+\infty} x \frac{f_X^2(\Phi^{-1}(x))}{(\Phi'(\Phi^{-1}(x)))^2} dx.$$

Let  $\Phi$  be strictly increasing. Then, with a change of variable in the above integral, we get

$$J^w(Y_t) = -\frac{1}{2\overline{F}_X^2(\Phi^{-1}(t))} \int_{\Phi^{-1}(t)}^{+\infty} \frac{\Phi(x)}{\Phi'(x)} f_X^2(x) dx,$$

giving the first expression in (23). If  $\Phi$  is strictly decreasing, we similarly obtain

$$J^w(Y_t) = -\frac{1}{2\overline{F}_X^2(\Phi^{-1}(t))} \int_0^{\Phi^{-1}(t)} \frac{\Phi(x)}{|\Phi'(x)|} f_X^2(x) dx,$$

giving the second expression in (23). The proof of (24) is quite similar.  $\square$

## 5 Concluding remarks

In this paper, some new measures of information have been introduced and studied. We have defined the weighted extropy as well as weighted residual and past extropies. We have presented some bounds and characterization results under monotonicity of hazard and reversed hazard functions. We have also presented bivariate versions of extropy and weighted extropy. We have presented numerous examples to illustrate all the concepts introduced here.

## Acknowledgments

Francesco Buono and Maria Longobardi are partially supported by the GNAMPA research group of INdAM (Istituto Nazionale di Alta Matematica) and MIUR-PRIN 2017, Project "Stochastic Models for Complex Systems" (No. 2017 JFFHSH).

## References

- [1] Asadi, M., Zohrevand, Y. (2007). On the dynamic cumulative residual entropy. *Journal of Statistical Planning and Inference*, **137**, 1931–1941.

- [2] Barlow, R. E., Proschan, F. J. (1996). *Mathematical Theory of Reliability*. Philadelphia: Society for Industrial and Applied Mathematics.
- [3] Block, H. W., Savits, T. H. (1998). The reversed hazard rate function. *Probability in the Engineering and Informational Sciences*, **12**, 69–90.
- [4] Di Crescenzo, A., Longobardi, M. (2002). Entropy-based measure of uncertainty in past lifetime distributions. *Journal of Applied Probability*, **39**, 434–440.
- [5] Di Crescenzo, A., Longobardi, M. (2006). On weighted residual and past entropies. *Scientiae Mathematicae Japonicae*, **64** (2), 255–266.
- [6] Di Crescenzo, A., Longobardi, M. (2009a). On cumulative entropies. *Journal of Statistical Planning and Inference*, **139**, 4072–4087.
- [7] Di Crescenzo, A., Longobardi, M. (2009b). On cumulative entropies and lifetime estimations, In: J. Mira, J.M. Ferrandez, J.R. Alvarez Sanchez, F. Paz, J. Toledo (Eds.), *Methods and Models in Artificial and Natural Computation*, IWINAC 2009, Part I, in: LNCS, vol. 5601, Springer-Verlag, Berlin, Heidelberg, pp. 132–141.
- [8] Di Crescenzo, A., Longobardi, M. (2013) Stochastic comparisons of cumulative entropies, In: H. Li, X. Li (Eds.), *Stochastic Orders in Reliability and Risk, in Honor of Professor Moshe Shaked*, Lecture Notes in Statistics, vol. 208, Springer, New York, pp. 167–182.
- [9] Ebrahimi, N. (1996). How to measure uncertainty in the residual life time distribution. *Sankhya: Series A*, **58**, 48–56.
- [10] Finkelstein, M. S. (2002). On the reversed hazard rate. *Reliability Engineering & System Safety*, **78**, 71–75.
- [11] Jahanshani, S. M. A., Zarei, H., Khammar, A. H. (2019). On cumulative residual extropy. *Probability in the Engineering and Informational Sciences*, DOI:10.1017/S0269964819000196.
- [12] Kamari, O., Buono, F. (2020). On extropy of past lifetime distribution. *Ricerche di Matematica*, DOI: 10.1007/s11587-020-00488-7.
- [13] Krishnan, A. S., Sunoj S. M., Nair N. U. (2020). Some reliability properties of extropy for residual and past lifetime random variables. *Journal of the Korean Statistical Society*, <https://doi.org/10.1007/s42952-019-00023-x>.

- [14] Lad, F., Sanfilippo, G., Agr, G. (2015). Extropy: complementary dual of entropy. *Statistical Science*, **30**, 40–58.
- [15] Qiu, G. (2017). The extropy of order statistics and record values. *Statistics & Probability Letters*, **120**, 52–60.
- [16] Rao, M., Chen, Y., Vemuri, B.C., Wang, F. (2004). Cumulative residual entropy: a new measure of information, *IEEE Transactions on Information Theory*, **50**, 1220–1228.
- [17] Sathar, A. E. I., Nair, D. R. (2019). On dynamic survival extropy. *Communication in Statistics - Theory and Methods*, <https://doi.org/10.1080/03610926.2019.1649426>.
- [18] Shannon, C. E. (1948). A mathematical theory of communication. *Bell System Technical Journal*, **27**, 379–423.