

Local and nonlocal $(2 + 1)$ -dimensional Maccari systems and their soliton solutions

Aslı Pekcan *

Department of Mathematics, Faculty of Science
Hacettepe University, 06800 Ankara - Turkey

Abstract

In this work, by using the Hirota bilinear method, we obtain one- and two-soliton solutions of integrable $(2 + 1)$ -dimensional 3-component Maccari system which is used as a model describing isolated waves localized in a very small part of space and related to very well-known systems like nonlinear Schrödinger, Fokas, and long wave resonance systems. We represent all local and Ablowitz-Musslimani type nonlocal reductions of this system and obtain new integrable systems. By the help of reduction formulas and soliton solutions of the 3-component Maccari system, we obtain one- and two-soliton solutions of these new integrable local and nonlocal reduced 2-component Maccari systems. We also illustrate our solutions by plotting their graphs for particular values of the parameters.

Keywords: Maccari systems, Local and nonlocal reductions, Hirota bilinear method, Soliton solutions

1 Introduction

The generalized $(2 + 1)$ -dimensional $(N + 1)$ -component Maccari system is given by

$$iu_{k,t} + u_{k,xx} + pu_k = 0, \quad k = 1, 2, \dots, N, \quad (1)$$

$$p_y = \sum_{k=1}^N (\sigma_k u_k \bar{u}_k)_x, \quad (2)$$

where $p = p(x, y, t)$, $\sigma_k = \pm 1$, $u_k = u_k(x, y, t)$ for $k = 1, 2, \dots, N$, and the bar notation is used for complex conjugation. Here u_k denote N different complex short wave amplitudes and p denotes the real long wave amplitude. The system for $N = 2$ is

$$iu_t + u_{xx} + pu = 0, \quad (3)$$

$$iv_t + v_{xx} + pv = 0, \quad (4)$$

$$p_y = (\sigma_1 u \bar{u} + \sigma_2 v \bar{v})_x, \quad (5)$$

*aslipekcan@hacettepe.edu.tr

where $\sigma_k = \pm 1$, $k = 1, 2$. This $(2 + 1)$ -dimensional 3-component system was first derived by Maccari [1] from Nizhnik-Novikov-Veselov equation which is an S-integrable equation, with a reduction technique based on Fourier decomposition and space-time rescalings. In [2], it was noted that Maccari system can also be obtained from many nonlinear partial differential equations by following the same technique. This method preserves the integrability that is the Maccari system (3)-(5) is also integrable. Painlevé analysis was used to investigate the integrability of the Maccari system by Uthayakumar et al. in [3].

The Maccari system (3)-(5) is related to some well-known equations. If $y = x$ it reduces to coupled nonlinear Schrödinger (NLS) equation [4]. When $y = t$, we have the coupled long-wave resonance system [5]. We get $(2 + 1)$ -dimensional extension of NLS equation introduced by Fokas [6] when $u = \bar{v}$.

Maccari system has many applications in hydrodynamics, plasma physics, Bose-Einstein condensates, and nonlinear optics. It is a model describing isolated waves which are localized in a very small part of space. There are several works on obtaining solutions of the Maccari system (3)-(5). By KP hierarchy reduction method, bright-dark mixed soliton solutions [7], [8], multi-dark soliton solutions [9], semi-rational solutions [10] of the Maccari system were obtained. Solitoff and dromion solutions [11], solutions in terms of Jacobi elliptic functions [12], folded solitary waves and foldons [13] were found by using variable separation approach. Half bright, bright, dark, half dark, and combined solitons were given by extended modified auxiliary equation mapping technique in [14]. By the Hirota bilinear method, soliton solutions [3], lump and rogue wave solutions [15], and dromions [16] were derived. By means of the Bäcklund transformation, peakon and compacton solutions of the Maccari system were obtained in [17].

Recently, there is a huge interest on constructing integrable nonlocal equations and finding various types of solutions to these equations. The nonlocal reductions were first introduced by Ablowitz and Musslimani [18]-[20]. It has been indicated that systems admitting nonlocal reductions have discrete symmetry transformations that leave the systems invariant. In [24] we showed that a special case of discrete symmetry transformations gives the nonlocal reductions of the same systems. As examples of $(1 + 1)$ -dimensional nonlocal integrable equations we can mention some famous equations; nonlocal NLS [18]-[20], [25]-[38], modified Korteweg-de Vries (mKdV) [19]-[25], [29], [39]-[45], Fordy-Kulish [46], hydrodynamic type [47], KdV [48] equations, and so on. For $(2 + 1)$ -dimensional nonlocal integrable equations one can check the references [49]-[53] for Davey-Stewartson equations, [54] for Fokas equations, [55] and [56] for $(2 + 1)$ -dimensional mKdV equations, [57] for $(2 + 1)$ -dimensional NLS equations, [58] for Kadomtsev-Petviashvili equations, [59] and [60] for $(2 + 1)$ -dimensional negative AKNS equations, and so on. Possible applications of nonlocal equations have been discussed in [21]-[23].

As far as we know there is no work about the local and nonlocal reductions of the integrable $(2+1)$ -dimensional 3-component Maccari system even there are many works about different type of solutions of this system in the literature. We note that if the nonlocal (or local) reductions of integrable systems are done consistently then the reduced equations are also integrable. Finding new integrable local and nonlocal systems is very important in the integrable systems and soliton theory since they have rich physical structures and features leading to many developments in describing nature. In this work we obtain new integrable local and nonlocal reduced systems from the $(2 + 1)$ -dimensional 3-component Maccari system.

To be able to analyze all local and nonlocal reductions of the Maccari system we first let

$t \rightarrow \frac{a}{i}t$ and consider the following $(2 + 1)$ -dimensional 3-component Maccari system:

$$au_t + u_{xx} + pu = 0, \quad (6)$$

$$av_t + v_{xx} + pv = 0, \quad (7)$$

$$p_y = (\sigma_1 u \bar{u} + \sigma_2 v \bar{v})_x, \quad (8)$$

where a is an arbitrary constant. For this system we have two different local reductions,

$$i) v(x, y, t) = \rho u(x, y, t), \quad (9)$$

$$ii) v(x, y, t) = \rho \bar{u}(x, y, t), \quad (10)$$

where ρ is a real constant and \bar{u} is the complex conjugate of the function u . Both of these reductions give the same $(2 + 1)$ -dimensional 2-component Maccari system,

$$au_t + u_{xx} + pu = 0, \quad (11)$$

$$p_y = (\sigma_1 + \sigma_2 \rho^2)(u \bar{u})_x. \quad (12)$$

The Maccari system (6)-(8) has two different consistent nonlocal reductions. The first one of these reductions is

$$i) v(x, y, t) = \rho u(\varepsilon_1 x, \varepsilon_2 y, \varepsilon_3 t), \quad (13)$$

where ρ is a real constant and $\varepsilon_j^2 = 1$, $j = 1, 2, 3$ with the constraints $\varepsilon_3 = 1$ and $\varepsilon_1 \varepsilon_2 \sigma_1 = \sigma_2 \rho^2$. This reduction gives the following integrable space reversal nonlocal $(2 + 1)$ -dimensional 2-component Maccari systems:

$$au_t(x, y, t) + u_{xx}(x, y, t) + p(x, y, t)u(x, y, t) = 0, \quad (14)$$

$$p_y(x, y, t) = (\sigma_1 u(x, y, t) \bar{u}(x, y, t) + \sigma_2 u(\varepsilon_1 x, \varepsilon_2 y, t) \bar{u}(\varepsilon_1 x, \varepsilon_2 y, t))_x. \quad (15)$$

The second nonlocal reduction of the Maccari system is given by

$$ii) v(x, y, t) = \rho \bar{u}(\varepsilon_1 x, \varepsilon_2 y, \varepsilon_3 t), \quad (16)$$

yielding seven different nonlocal systems which are nonlocal time (T-), space (S-), and space-time (ST-) reversal Maccari systems represented by

$$au_t(x, y, t) + u_{xx}(x, y, t) + p(x, y, t)u(x, y, t) = 0, \quad (17)$$

$$p_y(x, y, t) = (\sigma_1 u(x, y, t) \bar{u}(x, y, t) + \sigma_2 u(\varepsilon_1 x, \varepsilon_2 y, \varepsilon_3 t) \bar{u}(\varepsilon_1 x, \varepsilon_2 y, \varepsilon_3 t))_x, \quad (18)$$

where $a = \bar{a} \varepsilon_3$ and $\varepsilon_1 \varepsilon_2 \sigma_1 = \sigma_2 \rho^2$.

In this work we study soliton solutions of local and nonlocal reduced $(2 + 1)$ -dimensional 2-component Maccari systems. We first obtain one- and two-soliton solutions of the 3-component Maccari system by Hirota bilinear method [61]-[66] which has been used as a powerful and practical tool to obtain multi-soliton solutions of integrable systems analytically. Then by using local and nonlocal Ablowitz-Musslimani type reductions, we obtain soliton solutions of the integrable local and nonlocal reduced 2-component Maccari systems. We present the graphs of some solutions for particular values of the parameters.

The lay out of the paper is as follows. In Section 2, we obtain one- and two-soliton solutions of the 3-component Maccari system by Hirota bilinear method. Then we present local and nonlocal reductions of this system in Section 3 and 4, respectively. In Section 5, we derive soliton solutions of integrable local and nonlocal reduced 2-component Maccari systems. We also give some examples of one- and two-soliton solutions of reduced Maccari systems.

2 Hirota bilinear method for 3-component Maccari system

Here by using the Hirota bilinear method [61]-[63] we will obtain one- and two-soliton solutions of 3-component Maccari system (6)-(8). Let

$$u = \frac{g}{f}, \quad v = \frac{h}{f}, \quad p = 2(\ln f)_{xx}. \quad (19)$$

Here $g(x, y, t)$, $h(x, y, t)$ are complex-valued functions and $f(x, y, t)$ is a real-valued function. We obtain the Hirota bilinear form of the system (6)-(8) as

$$(aD_t + D_x^2)\{g \cdot f\} = 0, \quad (20)$$

$$(aD_t + D_x^2)\{h \cdot f\} = 0, \quad (21)$$

$$D_x D_y \{f \cdot f\} = \sigma_1 g \bar{g} + \sigma_2 h \bar{h}. \quad (22)$$

Here D is a special differential operator called Hirota D -operator given by

$$D_t^n D_x^m D_y^r \{F \cdot G\} = \left(\frac{\partial}{\partial t} - \frac{\partial}{\partial t'}\right)^n \left(\frac{\partial}{\partial x} - \frac{\partial}{\partial x'}\right)^m \left(\frac{\partial}{\partial y} - \frac{\partial}{\partial y'}\right)^r F(x, y, t) G(x', y', t')|_{x'=x, t'=t, y'=y}. \quad (23)$$

We expand the functions g , h , and f in the form of power series as

$$\begin{aligned} g &= \epsilon g_1 + \epsilon^3 g_3 + \epsilon^5 g_5 + \dots, \\ h &= \epsilon h_1 + \epsilon^3 h_3 + \epsilon^5 h_5 + \dots, \\ f &= 1 + \epsilon^2 f_2 + \epsilon^4 f_4 + \dots. \end{aligned} \quad (24)$$

We substitute the above expansion into the Hirota bilinear form (20)-(22) and make the coefficients of ϵ^m , $m = 1, 2, \dots$ to vanish. Note that choosing the functions g_1 and h_1 as exponential functions determines the other functions g_j, h_j , and f_j as exponential functions. Due to this fact, when we consider N soliton solution we have $g_{2j+1} = h_{2j+1} = f_{2j+2} = 0$ for $j \geq N$. In the following two sections, one- and two-soliton solutions of $(2+1)$ -dimensional 3-component Maccari system will be given.

2.1 One-soliton solutions of 3-component Maccari system

In order to construct one-soliton solution, we take

$$g_1 = e^{\theta_1}, \quad h_1 = e^{\theta_2}, \quad (25)$$

in the expansion (24) and make the coefficients of ϵ^m , $m = 1, 2, \dots$ to vanish in (20)-(22). Here $\theta_j = k_j x + l_j y + \omega_j t + \delta_j$, $j = 1, 2$. By the above choice of g_1 and h_1 we have $g_{2j+1} = h_{2j+1} = f_{2j+2} = 0$ for $j \geq 1$. From the coefficients of ϵ , we get the dispersion relations,

$$\omega_j = -\frac{k_j^2}{a}, \quad j = 1, 2. \quad (26)$$

The coefficient of ϵ^2 yields the function f_2 as

$$f_2 = \frac{\sigma_1 e^{\theta_1 + \bar{\theta}_1}}{2(k_1 + \bar{k}_1)(l_1 + \bar{l}_1)} + \frac{\sigma_2 e^{\theta_2 + \bar{\theta}_2}}{2(k_2 + \bar{k}_2)(l_2 + \bar{l}_2)}. \quad (27)$$

The coefficient of ϵ^4

$$f_{2,xy}f_2 - f_{2,x}f_{2,y} = 0 \quad (28)$$

is satisfied if

$$(l_1 + \bar{l}_1 - l_2 - \bar{l}_2)(k_1 + \bar{k}_1 - k_2 - \bar{k}_2) = 0. \quad (29)$$

Therefore we have two possibilities:

$$1) l_2 + \bar{l}_2 = l_1 + \bar{l}_1, \quad (30)$$

$$2) k_2 + \bar{k}_2 = k_1 + \bar{k}_1. \quad (31)$$

The coefficients of ϵ^3 are

$$a(g_{1,t}f_2 - g_1f_{2,t}) + g_{1,xx}f_2 - 2g_{1,x}f_{2,x} + g_1f_{2,xx} = 0, \quad (32)$$

$$a(h_{1,t}f_2 - h_1f_{2,t}) + h_{1,xx}f_2 - 2h_{1,x}f_{2,x} + h_1f_{2,xx} = 0. \quad (33)$$

To satisfy the above equations we must have

$$\bar{a} = -a, \quad k_1 = k_2, \quad (34)$$

and so that (31) holds. Without loss of generality take $\epsilon = 1$. Hence one-soliton solution of the 3-component Maccari system (6)-(8) for $\bar{a} = -a$ is given by $(u(x, y, t), v(x, y, t), p(x, y, t))$ where

$$u(x, y, t) = \frac{e^{k_1x + \omega_1t + l_1y + \delta_1}}{1 + \frac{e^{(k_1 + \bar{k}_1)x + (\omega_1 + \bar{\omega}_1)t}}{2(k_1 + \bar{k}_1)} \left[\frac{\sigma_1 e^{(l_1 + \bar{l}_1)y + \delta_1 + \bar{\delta}_1}}{(l_1 + \bar{l}_1)} + \frac{\sigma_2 e^{(l_2 + \bar{l}_2)y + \delta_2 + \bar{\delta}_2}}{(l_2 + \bar{l}_2)} \right]}, \quad (35)$$

$$v(x, y, t) = \frac{e^{k_1x + \omega_1t + l_2y + \delta_2}}{1 + \frac{e^{(k_1 + \bar{k}_1)x + (\omega_1 + \bar{\omega}_1)t}}{2(k_1 + \bar{k}_1)} \left[\frac{\sigma_1 e^{(l_1 + \bar{l}_1)y + \delta_1 + \bar{\delta}_1}}{(l_1 + \bar{l}_1)} + \frac{\sigma_2 e^{(l_2 + \bar{l}_2)y + \delta_2 + \bar{\delta}_2}}{(l_2 + \bar{l}_2)} \right]}, \quad (36)$$

and

$$p(x, y, t) = \frac{(k_1 + \bar{k}_1)e^{(k_1 + \bar{k}_1)x + (\omega_1 + \bar{\omega}_1)t} \left[\frac{\sigma_1 e^{(l_1 + \bar{l}_1)y + \delta_1 + \bar{\delta}_1}}{(l_1 + \bar{l}_1)} + \frac{\sigma_2 e^{(l_2 + \bar{l}_2)y + \delta_2 + \bar{\delta}_2}}{(l_2 + \bar{l}_2)} \right]}{\left(1 + \frac{e^{(k_1 + \bar{k}_1)x + (\omega_1 + \bar{\omega}_1)t}}{2(k_1 + \bar{k}_1)} \left[\frac{\sigma_1 e^{(l_1 + \bar{l}_1)y + \delta_1 + \bar{\delta}_1}}{(l_1 + \bar{l}_1)} + \frac{\sigma_2 e^{(l_2 + \bar{l}_2)y + \delta_2 + \bar{\delta}_2}}{(l_2 + \bar{l}_2)} \right] \right)^2}, \quad (37)$$

where the dispersion relations (26) and the conditions (34) hold. Here $k_1, l_1, l_2, \delta_1, \delta_2$ are some arbitrary complex numbers.

2.2 Two-soliton solutions of 3-component Maccari system

To construct two-soliton solution, we take

$$g_1 = e^{\theta_1} + e^{\theta_2}, \quad h_1 = e^{m_1} + e^{m_2}, \quad (38)$$

in the expansion (24), where $\theta_j = k_jx + l_jy + \omega_jt + \delta_j$, $\eta_j = r_jx + m_jt + s_jy + \alpha_j$ for $j = 1, 2$. We make the coefficients of ϵ^m , $m = 1, 2, \dots$ to vanish in (20)-(22). By the choice of g_1 and h_1

we have $g_{2j+1} = h_{2j+1} = f_{2j+2} = 0$ for $j \geq 2$. From the coefficients of ϵ , we get the dispersion relations,

$$\omega_j = -\frac{k_j^2}{a}, \quad m_j = -\frac{r_j^2}{a}, \quad j = 1, 2. \quad (39)$$

The coefficient of ϵ^2 gives

$$f_2 = \sigma_1 \sum_{i \leq i, j \leq 2} e^{\theta_i + \bar{\theta}_j} \alpha_{ij} + \sigma_2 \sum_{i \leq i, j \leq 2} e^{\eta_i + \bar{\eta}_j} \beta_{ij}, \quad (40)$$

where

$$\alpha_{ij} = \frac{1}{2(k_i + \bar{k}_j)(l_i + \bar{l}_j)}, \quad \beta_{ij} = \frac{1}{2(r_i + \bar{r}_j)(s_i + \bar{s}_j)}, \quad i \leq i, j \leq 2. \quad (41)$$

From the coefficients of ϵ^3 we obtain the functions g_3 and h_3 as

$$\begin{aligned} g_3 &= A_1 e^{\theta_1 + \eta_2 + \bar{\eta}_1} + A_2 e^{\theta_2 + \eta_1 + \bar{\eta}_1} + A_3 e^{\theta_1 + \eta_2 + \bar{\eta}_2} + A_4 e^{\theta_2 + \eta_1 + \bar{\eta}_2} + A_5 e^{\theta_1 + \theta_2 + \bar{\theta}_1} + A_6 e^{\theta_1 + \theta_2 + \bar{\theta}_2}, \\ h_3 &= B_1 e^{\theta_2 + \eta_1 + \bar{\theta}_1} + B_2 e^{\theta_1 + \eta_2 + \bar{\theta}_1} + B_3 e^{\theta_2 + \eta_1 + \bar{\theta}_2} + B_4 e^{\theta_1 + \eta_2 + \bar{\theta}_2} + B_5 e^{\eta_1 + \eta_2 + \bar{\eta}_1} + B_6 e^{\eta_1 + \eta_2 + \bar{\eta}_2}, \end{aligned} \quad (42)$$

where

$$A_1 = \sigma_2 \beta_{21} \frac{(k_1 - k_2)}{(k_1 + \bar{k}_1)}, \quad A_2 = -\sigma_2 \beta_{11} \frac{(k_1 - k_2)}{(k_2 + \bar{k}_1)}, \quad A_3 = \sigma_2 \beta_{22} \frac{(k_1 - k_2)}{(k_1 + \bar{k}_2)}, \quad A_4 = -\sigma_2 \beta_{12} \frac{(k_1 - k_2)}{(k_2 + \bar{k}_2)},$$

$$A_5 = -\sigma_1 (k_1 - k_2) \frac{[\alpha_{11}(k_1 + \bar{k}_1) - \alpha_{21}(k_2 + \bar{k}_1)]}{(k_1 + \bar{k}_1)(k_2 + \bar{k}_1)}, \quad A_6 = -\sigma_1 (k_1 - k_2) \frac{[\alpha_{12}(k_1 + \bar{k}_2) - \alpha_{22}(k_2 + \bar{k}_2)]}{(k_1 + \bar{k}_2)(k_2 + \bar{k}_2)},$$

and

$$B_1 = \sigma_1 \alpha_{21} \frac{(k_1 - k_2)}{(k_1 + \bar{k}_1)}, \quad B_2 = -\sigma_1 \alpha_{11} \frac{(k_1 - k_2)}{(k_2 + \bar{k}_1)}, \quad B_3 = \sigma_1 \alpha_{22} \frac{(k_1 - k_2)}{(k_1 + \bar{k}_2)}, \quad B_4 = -\sigma_1 \alpha_{12} \frac{(k_1 - k_2)}{(k_2 + \bar{k}_2)},$$

$$B_5 = -\sigma_2 (k_1 - k_2) \frac{[\beta_{11}(k_1 + \bar{k}_1) - \beta_{21}(k_2 + \bar{k}_1)]}{(k_1 + \bar{k}_1)(k_2 + \bar{k}_1)}, \quad B_6 = -\sigma_2 (k_1 - k_2) \frac{[\beta_{12}(k_1 + \bar{k}_2) - \beta_{22}(k_2 + \bar{k}_2)]}{(k_1 + \bar{k}_2)(k_2 + \bar{k}_2)},$$

with the following conditions satisfied:

$$a = -\bar{a}, \quad r_j = k_j, \quad j = 1, 2. \quad (43)$$

The coefficient of ϵ^4 yields the function f_4 as

$$\begin{aligned} f_4 &= C_1 e^{\theta_1 + \bar{\theta}_1 + \eta_2 + \bar{\eta}_2} + C_2 e^{\theta_2 + \bar{\theta}_2 + \eta_1 + \bar{\eta}_1} + C_3 e^{\eta_1 + \bar{\eta}_1 + \eta_2 + \bar{\eta}_2} \\ &\quad + C_4 e^{\theta_1 + \bar{\theta}_1 + \theta_2 + \bar{\theta}_2} + C_5 e^{\theta_1 + \bar{\theta}_2 + \eta_2 + \bar{\eta}_1} + C_6 e^{\theta_2 + \bar{\theta}_1 + \eta_1 + \bar{\eta}_2}, \end{aligned} \quad (44)$$

where

$$C_1 = \frac{\sigma_1 \sigma_2 \kappa}{(s_2 + \bar{s}_2)(l_1 + \bar{l}_1)}, \quad C_2 = \frac{\sigma_1 \sigma_2 \kappa}{(s_1 + \bar{s}_1)(l_2 + \bar{l}_2)}, \quad C_3 = \frac{\sigma_2^2 \kappa (s_1 - s_2)(\bar{s}_1 - \bar{s}_2)}{(s_1 + \bar{s}_1)(s_1 + \bar{s}_2)(s_2 + \bar{s}_1)(s_2 + \bar{s}_2)},$$

$$C_4 = \frac{\sigma_1^2 \kappa (l_1 - l_2)(\bar{l}_1 - \bar{l}_2)}{(l_1 + \bar{l}_1)(l_1 + \bar{l}_2)(l_2 + \bar{l}_1)(l_2 + \bar{l}_2)}, \quad C_5 = -\frac{\sigma_1 \sigma_2 \kappa}{(s_2 + \bar{s}_1)(l_1 + \bar{l}_2)}, \quad C_6 = -\frac{\sigma_1 \sigma_2 \kappa}{(s_1 + \bar{s}_2)(l_2 + \bar{l}_1)},$$

for

$$\kappa = \frac{(k_1 - k_2)(\bar{k}_1 - \bar{k}_2)}{4(k_1 + \bar{k}_1)(k_1 + \bar{k}_2)(k_2 + \bar{k}_1)(k_2 + \bar{k}_2)}. \quad (45)$$

The coefficients of ϵ^j , $j = 5, 6, 7, 8$ vanish directly. Take also $\epsilon = 1$. Since two-soliton solution of the 3-component Maccari system (6)-(8) has a long expression, we will not give it here explicitly. It can be expressed as $(u(x, y, t), v(x, y, t), p(x, y, t))$ where

$$u = \frac{g_1 + g_3}{1 + f_2 + f_4}, \quad v = \frac{h_1 + h_3}{1 + f_2 + f_4}, \quad p = 2(\ln(1 + f_2 + f_4))_{xx}. \quad (46)$$

Here g_1, h_1 are given in (38), g_3, h_3 in (42), f_2 in (40), and f_4 in (44) with the dispersion relations (39) and the constraints (43) satisfied.

3 Local reductions of 3-component Maccari system

(a) $v(x, y, t) = \rho u(x, y, t)$, ρ is a real constant.

The system (6)-(8) consistently reduces to the local system

$$au_t + u_{xx} + pu = 0, \quad (47)$$

$$p_y = (\sigma_1 + \sigma_2 \rho^2)(u\bar{u})_x. \quad (48)$$

Here there is no restriction on the constant a .

(b) $v(x, y, t) = \rho \bar{u}(x, y, t)$, ρ is a real constant.

In this case the system (6)-(8) consistently reduces to the same local system (47) and (48) with the restriction $a = \bar{a}$.

4 Nonlocal reductions of 3-component Maccari system

(a) $v(x, y, t) = \rho u(\varepsilon_1 x, \varepsilon_2 y, \varepsilon_3 t) = \rho u^\varepsilon$, ρ is a real constant, and $\varepsilon_j = \pm 1$, $j = 1, 2, 3$.

We apply this reduction to 3-component Maccari system (6)-(8). We get

$$au_t + u_{xx} + uD_y^{-1}[\sigma_1 u\bar{u} + \sigma_2 \rho^2 u^\varepsilon \bar{u}^\varepsilon]_x = 0, \quad (49)$$

$$a\rho u_t^\varepsilon + \rho u_{xx}^\varepsilon + \rho u^\varepsilon D_y^{-1}[\sigma_1 u\bar{u} + \sigma_2 \rho^2 u^\varepsilon \bar{u}^\varepsilon]_x = 0. \quad (50)$$

Here D_y^{-1} is the standard anti-derivative with respect to y . Let $x \rightarrow \varepsilon_1 x$, $y \rightarrow \varepsilon_2 y$, and $t \rightarrow \varepsilon_3 t$ in (50). We have

$$a\varepsilon_3 u_t + u_{xx} + \varepsilon_1 \varepsilon_2 u D_y^{-1}[\sigma_1 u^\varepsilon \bar{u}^\varepsilon + \sigma_2 \rho^2 u\bar{u}]_x = 0. \quad (51)$$

For consistency, this equation must be same with (49) provided that

$$\varepsilon_3 = 1, \quad \varepsilon_1 \varepsilon_2 \sigma_1 = \sigma_2 \rho^2. \quad (52)$$

Since $\sigma_j = \pm 1$, $j = 1, 2$, we have $\rho = \pm 1$. Then the system (6)-(8) reduces to S-reversal nonlocal systems consistently,

$$au_t(x, y, t) + u_{xx}(x, y, t) + p(x, y, t)u(x, y, t) = 0, \quad (53)$$

$$p_y(x, y, t) = (\sigma_1 u(x, y, t)\bar{u}(x, y, t) + \sigma_2 u(\varepsilon_1 x, \varepsilon_2 y, t)\bar{u}(\varepsilon_1 x, \varepsilon_2 y, t))_x. \quad (54)$$

Explicitly, we have three different S-reversal nonlocal systems:

(i) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (1, -1, 1), \sigma_2 = -\sigma_1, \sigma_1 = \pm 1.$

$$au_t(x, y, t) + u_{xx}(x, y, t) + p(x, y, t)u(x, y, t) = 0, \quad (55)$$

$$p_y(x, y, t) = (\sigma_1 u(x, y, t)\bar{u}(x, y, t) + \sigma_2 u(x, -y, t)\bar{u}(x, -y, t))_x. \quad (56)$$

(ii) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (-1, 1, 1), \sigma_2 = -\sigma_1, \sigma_1 = \pm 1.$

$$au_t(x, y, t) + u_{xx}(x, y, t) + p(x, y, t)u(x, y, t) = 0, \quad (57)$$

$$p_y(x, y, t) = (\sigma_1 u(x, y, t)\bar{u}(x, y, t) + \sigma_2 u(-x, y, t)\bar{u}(-x, y, t))_x. \quad (58)$$

(iii) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (-1, -1, 1), \sigma_2 = \sigma_1, \sigma_1 = \pm 1.$

$$au_t(x, y, t) + u_{xx}(x, y, t) + p(x, y, t)u(x, y, t) = 0, \quad (59)$$

$$p_y(x, y, t) = (\sigma_1 u(x, y, t)\bar{u}(x, y, t) + \sigma_2 u(-x, -y, t)\bar{u}(-x, -y, t))_x. \quad (60)$$

(b) $v(x, y, t) = \rho\bar{u}(\varepsilon_1 x, \varepsilon_2 y, \varepsilon_3 t)$, ρ is a real constant, and $\varepsilon_k = \pm 1$, $k = 1, 2, 3$.

When we apply this nonlocal reduction to the system (6)-(8) and making the similar analysis as in part (a), we get the following constraints for consistency:

$$a = \bar{a}\varepsilon_3, \quad \varepsilon_1\varepsilon_2\sigma_1 = \sigma_2\rho^2, \quad (61)$$

yielding $\rho = \pm 1$. The system (6)-(8) reduces to the following nonlocal systems:

$$au_t(x, y, t) + u_{xx}(x, y, t) + p(x, y, t)u(x, y, t) = 0, \quad (62)$$

$$p_y(x, y, t) = (\sigma_1 u(x, y, t)\bar{u}(x, y, t) + \sigma_2 u(\varepsilon_1 x, \varepsilon_2 y, \varepsilon_3 t)\bar{u}(\varepsilon_1 x, \varepsilon_2 y, \varepsilon_3 t))_x. \quad (63)$$

Explicitly, here we have seven different nonlocal T-, S-, and ST-reversal 2-component Maccari systems corresponding to

(i) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (1, 1, -1), \sigma_2 = \sigma_1, a = -\bar{a},$

(ii) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (1, -1, -1), \sigma_2 = -\sigma_1, a = -\bar{a},$

(iii) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (-1, -1, -1), \sigma_2 = \sigma_1, a = -\bar{a},$

(iv) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (-1, 1, -1), \sigma_2 = -\sigma_1, a = -\bar{a},$

(v) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (1, -1, 1), \sigma_2 = -\sigma_1, a = \bar{a},$

(vi) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (-1, -1, 1), \sigma_2 = \sigma_1, a = \bar{a},$

(vii) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (-1, 1, 1), \sigma_2 = -\sigma_1, a = \bar{a}.$

Note that the systems obtained for the cases (v)-(vii) have already been given in the part (a) with no restriction on the constant a .

In [54], the $(2 + 1)$ -dimensional nonlocal Fokas system given by

$$\begin{aligned} iu_t(x, y, t) + u_{xx}(x, y, t) + u(x, y, t)p(x, y, t) &= 0, \\ p_y &= (u(x, y, t)\bar{u}(-x, -y, t))_x, \end{aligned} \quad (64)$$

was studied and semi-rational solutions of this system were obtained. The above system is similar but different than the system (62) and (63) corresponding to (vi).

5 Soliton solutions of the local and nonlocal reduced systems

In Sections 3 and 4 we have obtained consistent local and nonlocal reductions of 3-component Maccari system (6)-(8) and the constraints to be satisfied. By using the reduction formulas with one- and two-soliton solutions of (6)-(8) and following Type 1 or Type 2 approaches [28], [29], [39] we can obtain one- and two-soliton solutions of the local and nonlocal reduced 2-component Maccari systems.

5.1 One-soliton solutions of the local reduced systems

Both of the local reductions; $v(x, y, t) = \rho u(x, y, t)$ and $v(x, y, t) = \rho \bar{u}(x, y, t)$, yield the same reduced local system (47) and (48). The latter one puts a condition on the constant a as $a = \bar{a}$. Note that we have given one-soliton solution of 3-component Maccari system (6)-(8) under the constraints (34). Therefore we only consider the first local reduction $v(x, y, t) = \rho u(x, y, t)$ giving

$$\frac{h}{f} = \rho \frac{g}{f}. \quad (65)$$

When we use Type 1 that is the approach based on equating numerators and denominators separately, we get the following conditions:

$$l_2 = l_1, \quad e^{\delta_2} = \rho e^{\delta_1}. \quad (66)$$

Hence one-soliton solution of the local reduced system (47) and (48) is given by the pair $(u(x, y, t), p(x, y, t))$ where

$$u(x, y, t) = \frac{e^{k_1 x + \omega_1 t + l_1 y + \delta_1}}{\left(1 + \frac{(\sigma_1 + \rho^2 \sigma_2)}{2(k_1 + \bar{k}_1)(l_1 + \bar{l}_1)} e^{(k_1 + \bar{k}_1)x + (\omega_1 + \bar{\omega}_1)t + (l_1 + \bar{l}_1)y + \delta_1 + \bar{\delta}_1}\right)} \quad (67)$$

and

$$p(x, y, t) = \frac{(k_1 + \bar{k}_1)(\sigma_1 + \rho^2 \sigma_2) e^{(k_1 + \bar{k}_1)x + (\omega_1 + \bar{\omega}_1)t + (l_1 + \bar{l}_1)y + \delta_1 + \bar{\delta}_1}}{(l_1 + \bar{l}_1) \left(1 + \frac{(\sigma_1 + \rho^2 \sigma_2)}{2(k_1 + \bar{k}_1)(l_1 + \bar{l}_1)} e^{(k_1 + \bar{k}_1)x + (\omega_1 + \bar{\omega}_1)t + (l_1 + \bar{l}_1)y + \delta_1 + \bar{\delta}_1}\right)^2}. \quad (68)$$

Here $\omega_1 = -\frac{k_1^2}{a}$, $\sigma_j = \pm 1$ for $j = 1, 2$, and $a = -\bar{a}$. Let $k_1 = \alpha_1 + i\beta_1$, $\omega_1 = \alpha_2 + i\beta_2$, $a = i\beta_3$,

$l_1 = \alpha_4 + i\beta_4$, and $e^{\delta_1} = \alpha_5 + i\beta_5$ where $\alpha_1, \alpha_2, \alpha_4, \alpha_5 \in \mathbb{R}$, $\beta_j \in \mathbb{R}$ for $1 \leq j \leq 5$. Indeed, $\alpha_2 = -\frac{2\alpha_1\beta_1}{\beta_3}$ and $\beta_2 = \frac{(\alpha_1^2 - \beta_1^2)}{\beta_3}$. Then from the solution (67) we have

$$|u(x, y, t)|^2 = \frac{2\alpha_1\alpha_4}{(\sigma_1 + \rho^2\sigma_2)} \operatorname{sech}^2(\alpha_1x + \alpha_2t + \alpha_4y + \phi) \quad (69)$$

and from (68) we get

$$p(x, y, t) = 2\alpha_1^2 \operatorname{sech}^2(\alpha_1x + \alpha_2t + \alpha_4y + \phi) \quad (70)$$

with $\phi = \frac{1}{2} \ln \left| \frac{(\alpha_5^2 + \beta_5^2)(\sigma_1 + \rho^2\sigma_2)}{8\alpha_1\alpha_4} \right|$. These solutions are nonsingular and bounded if $\frac{(\sigma_1 + \rho^2\sigma_2)}{\alpha_1\alpha_4} > 0$. They are bell-shaped soliton solutions.

Example 1. Take the parameters of one-soliton solution of the local system (47) and (48) as $(k_1, l_1, a, \rho) = (1 + \frac{i}{2}, \frac{1}{3}, 2i, 4)$ with $\sigma_1 = \sigma_2 = e^{\delta_1} = 1$. Then the pair $(|u(x, y, t)|^2, p(x, y, t))$ becomes

$$|u(x, y, t)|^2 = \frac{2}{51} \operatorname{sech}^2(x + \frac{1}{3}y - \frac{1}{2}t + \delta), \quad p(x, y, t) = 2 \operatorname{sech}^2(x + \frac{1}{3}y - \frac{1}{2}t + \frac{1}{2} + \delta), \quad (71)$$

where $\delta = \frac{1}{2} \ln(\frac{51}{8})$. The graphs of the above solutions at $t = 0$ are given in Figure 1.




Figure 1: One-soliton solutions of the local system (47) and (48) at $t = 0$ for the parameters $(k_1, l_1, a, \rho) = (1 + \frac{i}{2}, \frac{1}{3}, 2i, 4)$ with $\sigma_1 = \sigma_2 = e^{\delta_1} = 1$ (a) $|u(x, y, t)|^2$, (b) $p(x, y, t)$.

Figures 2(a) and 2(b) depict the movements of the solitons from $|u(x, y, t)|^2$ and $p(x, y, t)$ for different times in different positions at the y -axis, respectively.




Figure 2: The movement of the waves (71) as time increases.

5.2 One-soliton solutions of the nonlocal reduced systems

In this section we present one-soliton solutions of the nonlocal systems obtained from 3-component Maccari system (6)-(8) by two types of nonlocal reductions.

(a) $v(x, y, t) = \rho u(\varepsilon_1 x, \varepsilon_2 y, \varepsilon_3 t)$, $\rho = \pm 1$, and $\varepsilon_j = \pm 1$, $j = 1, 2, 3$.

By this nonlocal reduction we obtain three types of nonlocal systems corresponding to $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (1, -1, 1)$ with $\sigma_2 = -\sigma_1$; $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (-1, 1, 1)$ with $\sigma_2 = -\sigma_1$; and $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (-1, -1, 1)$ with $\sigma_2 = \sigma_1$. We analyze these cases separately.

(a).(i) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (1, -1, 1)$, $\sigma_2 = -\sigma_1$, $\sigma_1 = \pm 1$.

When we use Type 1, the reduction formula yields the conditions,

$$l_2 = -l_1, \quad e^{\delta_2} = \rho e^{\delta_1}. \quad (72)$$

Therefore one-soliton solution of the nonlocal system (55) and (56) is the pair $(u(x, y, t), p(x, y, t))$ where

$$u(x, y, t) = \frac{e^{k_1 x + \omega_1 t + l_1 y + \delta_1}}{1 + \sigma_1 \frac{e^{(k_1 + \bar{k}_1)x + (\omega_1 + \bar{\omega}_1)t + \delta_1 + \bar{\delta}_1}}{(k_1 + \bar{k}_1)(l_1 + \bar{l}_1)} \cosh((l_1 + \bar{l}_1)y)} \quad (73)$$

and $p(x, y, t)$ is

$$p(x, y, t) = \frac{\frac{2\sigma_1(k_1 + \bar{k}_1)}{(l_1 + \bar{l}_1)} e^{(k_1 + \bar{k}_1)x + (\omega_1 + \bar{\omega}_1)t + \delta_1 + \bar{\delta}_1} \cosh((l_1 + \bar{l}_1)y)}{\left[1 + \frac{\sigma_1}{(k_1 + \bar{k}_1)(l_1 + \bar{l}_1)} e^{(k_1 + \bar{k}_1)x + (\omega_1 + \bar{\omega}_1)t + \delta_1 + \bar{\delta}_1} \cosh((l_1 + \bar{l}_1)y)\right]^2}. \quad (74)$$

Let $k_1 = \alpha_1 + i\beta_1$, $\omega_1 = \alpha_2 + i\beta_2$, $a = i\beta_3$, $l_1 = \alpha_4 + i\beta_4$, and $e^{\delta_1} = \alpha_5 + i\beta_5$ where $\alpha_1, \alpha_2, \alpha_4, \alpha_5 \in \mathbb{R}$, $\beta_j \in \mathbb{R}$ for $1 \leq j \leq 5$. Here $\alpha_2 = -\frac{2\alpha_1\beta_1}{\beta_3}$ and $\beta_2 = \frac{\alpha_1^2 - \beta_1^2}{\beta_3}$. Then solutions become

$$|u(x, y, t)|^2 = \frac{e^{2\alpha_1 x + 2\alpha_2 t + 2\alpha_4 y} (\alpha_5^2 + \beta_5^2)}{\left(1 + \frac{\sigma_1(\alpha_5^2 + \beta_5^2)}{4\alpha_1\alpha_4} e^{2\alpha_1 x + 2\alpha_2 t} \cosh(2\alpha_4 y)\right)^2}, \quad (75)$$

and

$$p(x, y, t) = \frac{2\alpha_1(\alpha_5^2 + \beta_5^2)\sigma_1 e^{2\alpha_1 x + 2\alpha_2 t} \cosh(2\alpha_4 y)}{\alpha_4 \left[1 + \frac{\sigma_1}{4\alpha_1\alpha_4} (\alpha_5^2 + \beta_5^2) e^{2\alpha_1 x + 2\alpha_2 t} \cosh(2\alpha_4 y)\right]^2}. \quad (76)$$

The above solutions are nonsingular and bounded if $\frac{\sigma_1}{\alpha_1\alpha_4} > 0$. Consider the following particular example.

Example 2. Choose the parameters of the one-soliton solution (73) as $(k_1, l_1, a, \sigma_1, e^{\delta_1}, \rho) = (1 + \frac{i}{4}, \frac{1}{2}, i, 1, 1, 1)$. Then the solutions of the nonlocal system (55) and (56) become

$$|u(x, y, t)|^2 = \frac{e^{2x-t+y}}{(1 + \frac{1}{2}e^{2x-t} \cosh(y))^2}, \quad p(x, y, t) = \frac{4e^{2x-t} \cosh(y)}{(1 + \frac{1}{2}e^{2x-t} \cosh(y))^2}. \quad (77)$$

Both of the above solutions are nonsingular and bounded. The graphs of (77) at $t = 0$ are given in Figure 3.




Figure 3: One-soliton solutions of the nonlocal system (55) and (56) at $t = 0$ for the parameters $k_1 = 1 + \frac{i}{4}, l_1 = \frac{1}{2}, a = i, \sigma_1 = e^{\delta_1} = \rho = 1$. (a) Solitoff solution $|u(x, y, t)|^2$, (b) V-type solitary wave solution $p(x, y, t)$.

Figures 4(a) and 4(b) show the movements of the waves from $|u(x, y, t)|^2$ and $p(x, y, t)$ for different times in different positions at the y -axis, respectively.




Figure 4: The movement of the waves (77) as time increases.

(a).(ii) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (-1, 1, 1)$, $\sigma_2 = -\sigma_1$, $\sigma_1 = \pm 1$.

For this case if we use Type 1 we get $k_1 = 0$ giving trivial solutions $u(x, y, t) = 0$ and $p(x, y, t) = 0$ for the nonlocal system (57) and (58). Therefore we apply Type 2 approach based on the cross multiplication of the reduction formula. But in this case we obtain $l_1 = -\bar{l}_1$ yielding again a trivial solution.

(a).(iii) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (-1, -1, 1)$, $\sigma_2 = \sigma_1$, $\sigma_1 = \pm 1$.

Here similar to the previous case, Type 1 approach gives a trivial solution. Using Type 2 yields the following constraints:

$$k_1 = \bar{k}_1, \quad l_2 = \bar{l}_1, \quad e^{\delta_2} = \sigma_3 e^{\delta_1}, \quad e^{\delta_1 + \bar{\delta}_1} = \frac{2k_1 \sigma_3 (l_1 + \bar{l}_1)}{\rho \sigma_1}, \quad (78)$$

where $\sigma_3 = \pm 1$. Since $k_1 = \bar{k}_1$ and $a = -\bar{a}$ by (34), we have $\omega_1 = -\bar{\omega}_1$. Hence one-soliton solution of the nonlocal system (59) and (60) is given by the pair $(u(x, y, t), p(x, y, t))$ where

$$u(x, y, t) = \frac{e^{k_1 x + \omega_1 t + l_1 y + \delta_1}}{1 + \rho \sigma_3 e^{2k_1 x + (l_1 + \bar{l}_1)y}} \quad (79)$$

and

$$p(x, y, t) = \frac{8k_1^2 \sigma_3 e^{2k_1 x + (l_1 + \bar{l}_1)y}}{(1 + \rho \sigma_3 e^{2k_1 x + (l_1 + \bar{l}_1)y})^2}. \quad (80)$$

The above solutions are nonsingular and bounded if $\rho \sigma_3 > 0$. Let $a = i\beta_1$, $\omega_1 = i\frac{k_1^2}{\beta_1} = i\beta_2$, $l_1 = \alpha_3 + i\beta_3$, where $\alpha_3 \in \mathbb{R}$, $\beta_j \in \mathbb{R}$ for $j = 1, 2, 3$. Let also $\rho \sigma_3 = e^{2\delta}$. Then from (79) and (80) we have

$$|u(x, y, t)|^2 = k_1 \sigma_1 \alpha_3 \operatorname{sech}^2(k_1 x + \alpha_3 y + \delta) \quad \text{and} \quad p(x, y, t) = 2k_1^2 \operatorname{sech}^2(k_1 x + \alpha_3 y + \delta). \quad (81)$$

They are bell-shaped soliton solutions.

(b) $v(x, y, t) = \rho \bar{u}(\varepsilon_1 x, \varepsilon_2 y, \varepsilon_3 t)$, $\rho = \pm -1$, and $\varepsilon_k = \pm 1$, $k = 1, 2, 3$.

By this nonlocal reduction we obtain seven different nonlocal reduced systems. As we noted previously the systems corresponding to (v)-(vii) are same with the systems obtained by the reduction $v(x, y, t) = \rho u(\varepsilon_1 x, \varepsilon_2 y, \varepsilon_3 t)$ and we have already presented one-soliton solutions of these systems in the previous part (a). Therefore here we will only consider one-soliton solutions of the nonlocal reduced systems corresponding to (i)-(iv).

(b).(i) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (1, 1, -1)$, $\sigma_2 = \sigma_1$, $\sigma_1 = \pm 1$.

When we use Type 1 with the reduction formula we obtain the following constraints:

$$k_1 = \bar{k}_1, \quad l_2 = \bar{l}_1, \quad e^{\delta_2} = \rho e^{\bar{\delta}_1}. \quad (82)$$

In this case we have $\omega_1 = -\bar{\omega}_1$. Hence one-soliton solution of the nonlocal reduced system (62) and (63) with $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (1, 1, -1)$ is given by the pair $(u(x, y, t), p(x, y, t))$ where

$$u(x, y, t) = \frac{e^{k_1 x + \omega_1 t + l_1 y + \delta_1}}{1 + \frac{\sigma_1}{2k_1(l_1 + \bar{l}_1)} e^{2k_1 x + (l_1 + \bar{l}_1)y + \delta_1 + \bar{\delta}_1}} \quad (83)$$

and

$$p(x, y, t) = \frac{4k_1\sigma_1 e^{2k_1x+(l_1+\bar{l}_1)y+\delta_1+\bar{\delta}_1}}{(l_1 + \bar{l}_1)[1 + \frac{\sigma_1}{2k_1(l_1+l_1)} e^{2k_1x+(l_1+\bar{l}_1)y+\delta_1+\bar{\delta}_1}]^2}. \quad (84)$$

These solutions are nonsingular and bounded for $\frac{\sigma_1}{k_1\alpha_3} > 0$. Let $a = i\beta_1$, $\omega_1 = i\frac{k_1^2}{\beta_1} = i\beta_2$, $l_1 = \alpha_3 + i\beta_3$, $e^{\delta_1} = \alpha_4 + i\beta_4$, where $\alpha_3, \alpha_4 \in \mathbb{R}$, $\beta_j \in \mathbb{R}$ for $1 \leq j \leq 4$. Let also $\frac{\sigma_1(\alpha_4^2 + \beta_4^2)}{4k_1\alpha_3} = e^{2\delta}$. Then from (83) and (84) we get

$$|u(x, y, t)|^2 = \frac{k_1\alpha_3}{\sigma_1} \text{sech}^2(k_1x + \alpha_3y + \delta) \quad \text{and} \quad p(x, y, t) = 2k_1 \text{sech}^2(k_1x + \alpha_3y + \delta). \quad (85)$$

These are bell-shaped soliton solutions.

(b).(ii) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (1, -1, -1)$, $\sigma_2 = -\sigma_1$, $\sigma_1 = \pm 1$.

Using Type 1 yields the following constraints:

$$k_1 = \bar{k}_1, \quad l_2 = -\bar{l}_1, \quad e^{\delta_2} = \rho e^{\bar{\delta}_1}. \quad (86)$$

Here we also get $\omega_1 = -\bar{\omega}_1$ directly. Then we obtain one-soliton solution $(u(x, y, t), p(x, y, t))$ of the nonlocal reduced system (62) and (63) with $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (1, -1, -1)$ as

$$u(x, y, t) = \frac{e^{k_1x+\omega_1t+l_1y+\delta_1}}{1 + \frac{\sigma_1}{2k_1(l_1+l_1)} e^{2k_1x+\delta_1+\bar{\delta}_1} \cosh((l_1 + \bar{l}_1)y)} \quad (87)$$

and

$$p(x, y, t) = \frac{4k_1\sigma_1 e^{2k_1x+\delta_1+\bar{\delta}_1} \cosh((l_1 + \bar{l}_1)y)}{(l_1 + \bar{l}_1)[1 + \frac{\sigma_1}{2k_1(l_1+l_1)} e^{2k_1x+\delta_1+\bar{\delta}_1} \cosh((l_1 + \bar{l}_1)y)]^2}. \quad (88)$$

Let $a = i\beta_1$, $\omega_1 = i\frac{k_1^2}{\beta_1} = i\beta_2$, $l_1 = \alpha_3 + i\beta_3$, $e^{\delta_1} = \alpha_4 + i\beta_4$, where $\alpha_3, \alpha_4 \in \mathbb{R}$, $\beta_j \in \mathbb{R}$ for $1 \leq j \leq 4$. Then from (87) and (88) we obtain

$$|u(x, y, t)|^2 = \frac{e^{2k_1x+2\alpha_3y}(\alpha_4^2 + \beta_4^2)}{[1 + \frac{\sigma_1(\alpha_4^2 + \beta_4^2)}{4k_1\alpha_3} e^{2k_1x} \cosh(2\alpha_3y)]^2}, \quad p(x, y, t) = \frac{k_1\sigma_1(\alpha_4^2 + \beta_4^2)e^{2k_1x} \cosh(2\alpha_3y)}{\alpha_3[1 + \frac{\sigma_1(\alpha_4^2 + \beta_4^2)}{4k_1\alpha_3} e^{2k_1x} \cosh(2\alpha_3y)]^2}. \quad (89)$$

The above solutions are nonsingular and bounded for $\frac{\sigma_1}{k_1\alpha_3} > 0$. As in the nonlocal case (a).(i), the function $|u(x, y, t)|^2$ defines a solitoff and $p(x, y, t)$ a V-type solitary wave solution.

(b).(iii) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (-1, -1, -1)$, $\sigma_2 = \sigma_1$, $\sigma_1 = \pm 1$.

If we use Type 1, we get $k_1 = -\bar{k}_1$ which gives trivial solutions $u(x, y, t) = 0$ and $p(x, y, t) = 0$. Therefore we apply Type 2 approach. Here we obtain the following constraints:

$$l_2 = l_1, \quad e^{\delta_2} = \sigma_3 e^{\bar{\delta}_1}, \quad e^{\delta_1+\bar{\delta}_1} = \frac{(k_1 + \bar{k}_1)(l_1 + \bar{l}_1)}{\rho\sigma_1}, \quad (90)$$

where $\sigma_3 = \pm 1$. Hence one-soliton solution of the nonlocal reduced system (62) and (63) with $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (-1, -1, -1)$ is obtained as the pair $(u(x, y, t), p(x, y, t))$ where

$$u(x, y, t) = \frac{e^{k_1x+\omega_1t+l_1y+\delta_1}}{1 + \rho e^{(k_1+\bar{k}_1)x+(\omega_1+\bar{\omega}_1)t+(l_1+\bar{l}_1)y}} \quad (91)$$

and

$$p(x, y, t) = \frac{2\rho(k_1 + \bar{k}_1)^2 e^{(k_1 + \bar{k}_1)x + (\omega_1 + \bar{\omega}_1)t + (l_1 + \bar{l}_1)y}}{[1 + \rho e^{(k_1 + \bar{k}_1)x + (\omega_1 + \bar{\omega}_1)t + (l_1 + \bar{l}_1)y}]^2}. \quad (92)$$

The above solutions are nonsingular and bounded if $\rho = 1$. For $\rho = 1$, let $k_1 = \alpha_1 + i\beta_1$, $\omega_1 = \alpha_2 + i\beta_2$, $a = i\beta_3$, $l_1 = \alpha_4 + i\beta_4$, where $\alpha_1, \alpha_2, \alpha_4 \in \mathbb{R}$, $\beta_j \in \mathbb{R}$ for $1 \leq j \leq 4$. Here $\alpha_2 = -\frac{2\alpha_1\beta_1}{\beta_3}$ and $\beta_2 = \frac{(\alpha_1^2 - \beta_1^2)}{\beta_3}$. Then from (91) and (92) we get

$$|u(x, y, t)|^2 = \alpha_1\alpha_4\sigma_1 \operatorname{sech}^2(\alpha_1x + \alpha_2t + \alpha_4y + \delta), \quad p(x, y, t) = 2\alpha_1^2 \operatorname{sech}^2(\alpha_1x + \alpha_2t + \alpha_4y + \delta). \quad (93)$$

The solutions are nonsingular and bounded bell-shaped soliton solutions.

(b).(iv) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (-1, 1, -1)$, $\sigma_2 = -\sigma_1$, $\sigma_1 = \pm 1$.

For this case if we use Type 1 we get $k_1 = -\bar{k}_1$ which gives trivial solutions $u(x, y, t) = 0$ and $p(x, y, t) = 0$. We then apply Type 2 and get $l_1 = -\bar{l}_1$ which also yields trivial solutions.

5.3 Two-soliton solutions of the local reduced systems

Similar to one-soliton solution, we only consider the local reduction $v(x, y, t) = \rho u(x, y, t)$ since the second local reduction $v(x, y, t) = \rho \bar{u}(x, y, t)$ is consistent if $a = \bar{a}$ different than the constraint $a = -\bar{a}$ that we obtain for two-soliton solution of Maccari system.

(a) $v(x, y, t) = \rho u(x, y, t)$, ρ is a real constant.

Under this reduction, when we use Type 1 approach with two-soliton solution (46), in addition to the conditions (43) we get the following constraints:

$$s_j = l_j, \quad e^{\alpha_j} = \rho e^{\delta_j}, \quad j = 1, 2. \quad (94)$$

Example 3. Let us take the parameters of two-soliton solution as $(k_1, k_2, l_1, l_2, a) = (2, \frac{1}{2}, \frac{1}{3}, 1, 2i)$ with $\sigma_1 = \sigma_2 = \rho = e^{\delta_1} = e^{\delta_2} = 1$. Then the graphs of the two-soliton solutions of the local reduced 2-component Maccari system (47) and (48) at $t = 0$ are given in Figure 5.

Figures 6(a) and 6(b) show the interactions of the waves from $|u(x, y, t)|^2$ and $p(x, y, t)$ for different times in different positions at the y -axis, respectively.

5.4 Two-soliton solutions of the nonlocal reduced systems

In this section we obtain two-soliton solutions of nonlocal systems obtained from 3-component Maccari systems by two nonlocal reductions.

(a) $v(x, y, t) = \rho u(\varepsilon_1x, \varepsilon_2y, \varepsilon_3t)$, $\rho = \pm 1$, and $\varepsilon_j = \pm 1$, $j = 1, 2, 3$.

Under this reduction we will consider soliton solutions of the three different S-reversal nonlocal reduced 2-component Maccari systems.




Figure 5: Two-soliton solutions of the local system (47) and (48) at $t = 0$ for the parameters $(k_1, k_2, l_1, l_2, a) = (2, \frac{1}{2}, \frac{1}{3}, 1, 2i)$ with $\sigma_1 = \sigma_2 = \rho = e^{\delta_1} = e^{\delta_2} = 1$. (a) Two V-type solitary waves from $|u(x, y, t)|^2$, (b) V-type interaction of localized solitons from $p(x, y, t)$.




Figure 6: The movement of the waves as time increases.

(a).(i) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (1, -1, 1), \sigma_2 = -\sigma_1, \sigma_1 = \pm 1$.

If we use this nonlocal reduction with two-soliton solution (46) and apply Type 1, besides (43) we get the following constraints:

$$s_j = -l_j, \quad e^{\alpha_j} = \rho e^{\delta_j}, \quad j = 1, 2. \quad (95)$$

Example 4. Let us choose the two-soliton solution parameters as $(k_1, k_2, l_1, l_2, a) = (1 + \frac{i}{2}, \frac{1}{4}, \frac{1}{3}, 1, i)$ with $\sigma_1 = e^{\delta_1} = e^{\delta_2} = \rho = 1$. Then the graphs of the two-soliton solutions of (55) and (56) are given in Figure 7.




Figure 7: Two-soliton solutions of the nonlocal system (55) and (56) at $t = 0$ for the parameters $(k_1, k_2, l_1, l_2, a) = (1 + \frac{i}{2}, \frac{1}{4}, \frac{1}{3}, 1, i)$ with $\sigma_1 = e^{\delta_1} = e^{\delta_2} = \rho = 1$. (a) V-type interaction of two solitons from $|u(x, y, t)|^2$, (b) Interaction of V-type solitary waves from $p(x, y, t)$.

Figures 8(a) and 8(b) depict the interactions of the waves from $|u(x, y, t)|^2$ and $p(x, y, t)$ for different times in different positions at the y -axis, respectively.




Figure 8: The movement of the waves as time increases.

(a).(ii) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (-1, 1, 1), \sigma_2 = -\sigma_1, \sigma_1 = \pm 1$.

When we use this nonlocal reduction with (46) and apply Type 1, in addition to (43) we obtain the following constraints:

$$k_2 = -k_1, \quad s_1 = l_2, \quad s_2 = l_1, \quad e^{\alpha_1} = \rho e^{\delta_2}, \quad e^{\alpha_2} = \rho e^{\delta_1}. \quad (96)$$

Example 5. Consider the two-soliton solution parameters as $(k_1, l_1, l_2, a) = (1 + \frac{i}{2}, \frac{1}{2}, -1, 2i)$ with $\sigma_1 = e^{\delta_1} = e^{\delta_2} = \rho = 1$. Then the graphs of the two-soliton solutions of the nonlocal system (57) and (58) are given in Figure 9.




Figure 9: Two-soliton solutions of the nonlocal system (57) and (58) at $t = 0$ for the parameters $(k_1, l_1, l_2, a) = (1 + \frac{i}{2}, \frac{1}{2}, -1, 2i)$ with $\sigma_1 = e^{\delta_1} = e^{\delta_2} = \rho = 1$. (a) Interaction of two solitons from $|u(x, y, t)|^2$, (b) Interaction of two V-type solitary waves from $p(x, y, t)$.

Figures 10(a) and 10(b) depict the interactions of the waves from $|u(x, y, t)|^2$ and $p(x, y, t)$ for different times in different positions at the y -axis, respectively.




Figure 10: The movement of the waves as time increases.

(a).(iii) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (-1, -1, 1)$, $\sigma_2 = \sigma_1$, $\sigma_1 = \pm 1$.

For this case, applying Type 1 approach yields the conditions,

$$k_2 = -k_1, \quad s_1 = -l_2, \quad s_2 = -l_1, \quad e^{\alpha_1} = \rho e^{\delta_2}, \quad e^{\alpha_2} = \rho e^{\delta_1}, \quad (97)$$

besides the constraints (43). Here we obtain nonsingular solutions very similar to the cases (a).(i) and (a).(ii), hence we will not give a particular example for this case.

(b) $v(x, y, t) = \rho \bar{u}(\varepsilon_1 x, \varepsilon_2 y, \varepsilon_3 t)$, $\rho = \pm -1$, and $\varepsilon_j = \pm 1$, $j = 1, 2, 3$.

In this part we will consider soliton solutions of four different nonlocal reduced 2-component Maccari systems.

(b).(i) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (1, 1, -1)$, $\sigma_2 = \sigma_1$, $\sigma_1 = \pm 1$.

Here using two-soliton solution (46) with this reductions and applying Type 1 yield the following constraints:

$$k_j = \bar{k}_j, \quad s_j = \bar{l}_j, \quad e^{\alpha_j} = \rho e^{\bar{\delta}_j}, \quad j = 1, 2. \quad (98)$$

Example 6. Consider the two-soliton solution parameters as $(k_1, k_2, l_1, l_2, a) = (1, 2, \frac{1}{2}, 2, i)$ with $\sigma_1 = e^{\delta_1} = e^{\delta_2} = \rho = 1$. Then the graphs of the two-soliton solutions of the nonlocal system (57) and (58) are given in Figure 11.




Figure 11: Two-soliton solutions of the nonlocal system (57) and (58) at $t = 0$ for the parameters $(k_1, k_2, l_1, l_2, a) = (1, 2, \frac{1}{2}, 2, i)$ with $\sigma_1 = e^{\delta_1} = e^{\delta_2} = \rho = 1$. (a) Interaction of localized two-solitons from $|u(x, y, t)|^2$, (b) Interaction of localized two-solitons from $p(x, y, t)$.

Here the form of the waves changes as time changes but the their locations does not change. Figures 12(a) and 12(b) show the interactions of the waves from $|u(x, y, t)|^2$ and $p(x, y, t)$ for different times in different positions at the y -axis, respectively.




Figure 12: The movement of the waves as time increases.

(b).(ii) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (1, -1, -1)$, $\sigma_2 = -\sigma_1$, $\sigma_1 = \pm 1$.

Applying Type 1 approach, here we obtain the constraints,

$$k_j = \bar{k}_j, \quad s_j = -\bar{l}_j, \quad e^{\alpha_j} = \rho e^{\bar{\delta}_j}, \quad j = 1, 2. \quad (99)$$

Example 7. Take the two-soliton solution parameters as $(k_1, k_2, l_1, l_2, a) = (1, 2, \frac{1}{2}, 1, i)$ with $\sigma_1 = e^{\delta_1} = e^{\delta_2} = \rho = 1$. Then the graphs of the two-soliton solutions of the nonlocal system (57) and (58) are given in Figure 13.




Figure 13: Two-soliton solutions of the nonlocal system (57) and (58) at $t = 0$ for the parameters $(k_1, k_2, l_1, l_2, a) = (1, 2, \frac{1}{2}, 1, i)$ with $\sigma_1 = e^{\delta_1} = e^{\delta_2} = \rho = 1$. (a) Localized combined two solitoffs from $|u(x, y, t)|^2$, (b) Localized combined two V-type solitary waves from $p(x, y, t)$.

In Figure 13(a) there occurs a bump near the combined solitoffs. The amplitudes of the solitoffs change as time changes but they are localized. This property is also same for combined V-type waves in Figure 13(b). Figures 14(a) and 14(b) show the interactions of the waves from $|u(x, y, t)|^2$ and $p(x, y, t)$ for different times in different positions at the y -axis, respectively.




Figure 14: The movement of the waves as time increases.

(b).(iii) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (-1, -1, -1)$, $\sigma_2 = \sigma_1$, $\sigma_1 = \pm 1$.

In this case Type 1 gives trivial solution. Therefore we apply Type 2 approach and obtain the constraints,

$$s_j = l_j, \quad e^{\alpha_j} = \sigma_3 e^{\delta_j}, \quad j = 1, 2, \quad (100)$$

and

$$e^{2\delta_1} = \frac{\rho\sigma_1(k_1 + \bar{k}_2)(k_1 + \bar{k}_1)(l_1 + \bar{l}_2)(l_1 + \bar{l}_1)}{\sigma_3(l_1 - l_2)(k_1 - k_2)}, \quad e^{2\delta_2} = \frac{\rho\sigma_1(k_2 + \bar{k}_1)(k_2 + \bar{k}_2)(l_2 + \bar{l}_1)(l_2 + \bar{l}_2)}{\sigma_3(l_1 - l_2)(k_1 - k_2)}, \quad (101)$$

where $\sigma_3 = \pm 1$. Nonsingular two-soliton solutions obtained here are very similar to the ones given in (b).(i) case. Therefore we will not give a particular example corresponding to this case.

(b).(iv) $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (-1, 1, -1)$, $\sigma_2 = -\sigma_1$, $\sigma_1 = \pm 1$.

In this case using both Type 1 and Type 2 approaches gives trivial solutions $u(x, y, t) = 0$ and $p(x, y, t) = 0$.

6 Conclusion

In this work we considered integrable $(2 + 1)$ -dimensional 3-component Maccari system which has many applications in several areas such as hydrodynamics, plasma physics, Bose-Einstein condensates, and nonlinear optics. Under some special transformations, this system can be reduced to well-known systems like NLS, Fokas, and long wave resonance systems. We obtained one- and two-soliton solutions of the 3-component Maccari system by using the Hirota bilinear method. Then we presented all possible local and nonlocal reductions of the system and obtained several new integrable systems. By using one- and two-soliton solutions of the 3-component Maccari system with reduction formulas, we obtained soliton solutions of the reduced integrable 2-component Maccari systems. Types of the one-soliton solutions obtained for these reduced local and nonlocal systems are solitons, solitoffs, and V-type solitary waves. Two-soliton solutions of reduced 2-component systems define interactions of mentioned type of waves.

As a future work we plan to obtain different types of exact solutions to local and nonlocal reductions of 3-component Maccari system. In recent years, rogue wave (also known as freak wave, killer wave) phenomenon has become a hot topic in the nonlinear science. Rogue waves are large and spontaneous ocean surface waves. They also appear in optics, superfluids, Bose-Einstein condensates, cold matter systems, and so on. By finding higher-order soliton solutions via the Hirota bilinear method and applying long wave limit we can obtain rogue wave solutions [67]-[71] to the 3-component Maccari system and its local and nonlocal reductions. To study integrability properties of the Maccari system we will consider Lie symmetry analysis of this system [72]-[76]. Under this analysis we can also obtain new solutions of the 3-component Maccari system and its reductions.

References

- [1] A. Maccari, Universal and integrable nonlinear evolution systems of equations in (2+1) dimensions, *J. Math. Phys.* **38**, 4151, 1997.
- [2] A. Maccari, The Maccari system as model system for rogue waves, *Phys. Lett. A* **384**, 126740, 2020.
- [3] A. Uthayakumar, K. Nakkeeran, and K. Porsezian, Soliton solutions of new (2+1) dimensional nonlinear partial differential equations, *Chaos Solitons Fractals* **10** (9), 1513–1518, 1999.
- [4] V.F. Zakharov, A.B. Shabat, Exact theory of two-dimensional self-focusing and one-dimensional self-modulation of wave in nonlinear media, *Soviet Phys. JETP* **34** (1), 62–69, 1972.
- [5] A.D.D. Craik, *Wave interactions and fluid flows*, Cambridge University Press, Cambridge, 1985.
- [6] A.S. Fokas, On the simplest integrable equation in 2+1, *Inverse Probl.* **10**, L19–L22, 1994.
- [7] Z. Han, Y. Chen, Bright-dark mixed N-soliton solution of the two-dimensional Maccari system, *Chin Phys. Lett.* **34** (7), 070202, 2017.
- [8] Z. Han, Y. Chen, Bright-dark mixed N-soliton solution of the two-dimensional multicomponent Maccari system, *Z. Naturforsch. A* **72** (8), 2017.
- [9] T. Xu, Y. Chen, and Z. Qiao, Multi-dark soliton solutions for the (2+1)-dimensional multicomponent Maccari system, *Modern Phys. Lett. B* **33** (31), 1950390, 2019.
- [10] W. Liu, A.-M. Wazwaz, Dynamics of fusion and fission collisions between lumps and line solitons in the Maccari’s system, *Rom. J. Phys.* **64**, 111, 2019.
- [11] W.-H. Huang, C.-L. Zheng, and J.-F. Zhang, Coherent soliton structures of a new (2+1) dimensional evolution equation, *Acta Phys. Sin.* (in Chinese) **51**, 2676, 2002.
- [12] W.-H. Huang, Y.-L. Liu, and Z.-Y. Ma, Doubly periodic propagating wave patterns of (2+1)-dimensional Maccari system, *Commun. Theor. Phys.* **47**, 397–402, 2007.
- [13] W.-H. Huang, J.-F. Zhang, Folded localized excitations of the Maccari system, *Acta Phys. Polon. B* **35** (8), 2004.
- [14] N. Cheemaa, S. Chen, and A.R. Seadawy, Propagation of isolated waves of coupled nonlinear (2+1)-dimensional Maccari system in plasma physics, *Results Phys.* **17**, 102987, 2020.
- [15] F. Yuan, J. Rao, K. Porsezian, D. Mihalache, and J. He, Various exact rational solutions of the two-dimensional Maccari’s system, *Rom. J. Phys.* **61** (3-4), 378–399, 2016.
- [16] J.-F. Zhang, Generalized dromion structures of new (2+1)-dimensional nonlinear evolution equation, *Commun. Theor. Phys.* **35**, 267–270, 2001.

- [17] J.-F. Zhang, J.-P. Meng, and W.-H. Huang, Novel class of coherent localized structures for the Maccari system, *Commun. Theor. Phys.* **40**, 443–446, 2003.
- [18] M.J. Ablowitz and Z.H. Musslimani, Integrable nonlocal nonlinear Schrödinger equation, *Phys. Rev. Lett.* **110**, 064105, 2013.
- [19] M.J. Ablowitz and Z.H. Musslimani, Inverse scattering transform for the integrable nonlocal nonlinear Schrödinger equation, *Nonlinearity* **29**, 915–946, 2016.
- [20] M.J. Ablowitz and Z.H. Musslimani, Integrable nonlocal nonlinear equations, *Stud. App. Math.* **139** (1), 7–59, 2016.
- [21] S.Y. Lou, F. Huang, Alice-Bob physics: Coherent solutions of nonlocal KdV systems, *Sci. Rep.* **7**, 869, 2017.
- [22] S.Y. Lou, Alice-Bob systems, \hat{P} - \hat{T} - \hat{C} symmetry invariant and symmetry breaking soliton solutions, *J. Math. Phys.* **59**, 083507, 2018.
- [23] T.A. Gadzhimuradov, A.M. Agalarov, Towards a gauge equivalent magnetic structure of the nonlocal nonlinear Schrödinger equation, *Phys. Rev. A* **93**, 062124, 2016.
- [24] M. Gürses, A. Pekcan, and K. Zheltukhin, Discrete symmetries and nonlocal reductions, *Phys. Lett. A* **384**, 120065, 2020.
- [25] K. Chen, X. Deng, S. Lou, and D. Zhang, Solutions of local and nonlocal equations reduced from the AKNS hierarchy, *Stud. Appl. Math.* **141** (1), 113–141, 2018.
- [26] B.F. Feng, X.D. Luo, M.J. Ablowitz, and Z.H. Musslimani, General soliton solution to a nonlocal nonlinear Schrödinger equation with zero and nonzero boundary conditions, *Nonlinearity* **31** (12), 5385–5409, 2018.
- [27] V.S. Gerdjikov and A. Saxena, Complete integrability of nonlocal nonlinear Schrödinger equation, *J. Math. Phys.* **58** (1), 013502, 2017.
- [28] M. Gürses and A. Pekcan, Nonlocal nonlinear Schrödinger equations and their soliton solutions, *J. Math. Phys.* **59**, 051501, 2018.
- [29] M. Gürses and A. Pekcan, Integrable nonlocal reductions, *Symmetries, Differential Equations and Applications SDEA-III*, Istanbul, Turkey, August 2017, in: V.G. Kac, P.J. Olver, P. Winternitz, T. Ozer (Eds), *Springer Proceedings in Mathematics and Statistics*, **266**, 27–52, 2018.
- [30] X. Huang, L. Ling, Soliton solutions for the nonlocal nonlinear Schrödinger equation, *Eur. Phys. J. Plus* **131**, 148, 2016.
- [31] M. Li, T. Xu, Dark and antidark soliton interactions in the nonlocal nonlinear Schrödinger equation with the self-induced parity-time-symmetric potential, *Phys. Rev. E* **91**, 033202, 2015.
- [32] M.J. Ablowitz, B.F. Feng, X.D. Luo, and Z.H. Musslimani, Inverse scattering transform for the nonlocal reverse space-time nonlinear Schrödinger equation, *Theor. Math. Phys.* **196** (3), 1241–1267, 2018.

- [33] X.Y. Wen, Z. Yan, and Y. Yang, Dynamics of higher-order rational solitons for the nonlocal nonlinear Schrödinger equation with the self-induced parity-time-symmetric potential, *Chaos* **26**, 063123, 2015.
- [34] A. Khare, A. Saxena, Periodic and hyperbolic soliton solutions of a number of nonlocal nonlinear equations, *J. Math. Phys.* **56**, 032104, 2015.
- [35] J. Yang, General N-solitons and their dynamics in several nonlocal nonlinear Schrödinger equations, *Phys. Lett A* **383** (4), 328–337, 2019.
- [36] T. Xu, S. Lan, M. Li, L.-L. Li, and G.-W. Zhang, Mixed soliton solutions of the defocusing nonlocal nonlinear Schrödinger equation, *Phys. D* **390**, 47–61, 2019.
- [37] T. Xu, Y. Chen, M. Li, and D.-X. Meng, General stationary solutions of the nonlocal nonlinear Schrödinger equation and their relevance to the \mathcal{PT} -symmetric system, *Chaos* **29**, 123124, 2019.
- [38] W.-X. Ma, Inverse scattering for nonlocal reverse-time nonlinear Schrödinger equations, *Appl. Math. Lett.* **102**, 106161, 2020.
- [39] M. Gürses, A. Pekcan, Nonlocal nonlinear modified KdV equations and their soliton solutions, *Commun. Nonlinear Sci. Numer. Simul.* **67**, 427, 2019.
- [40] J.L. Ji, Z.N. Zhu, On a nonlocal modified Korteweg-de Vries equation: Integrability, Darboux transformation and soliton solutions, *Commun. Non. Sci. Numer. Simul.* **42**, 699, 2017.
- [41] J.L. Ji, Z.N. Zhu, Soliton solutions of an integrable nonlocal modified Korteweg-de Vries equation through inverse scattering transform, *J. Math. Anal. Appl.* **453**, 973, 2017.
- [42] L.Y. Ma, S.F. Shen, and Z.N. Zhu, Soliton solution and gauge equivalence for an integrable nonlocal complex modified Korteweg-de Vries equation, *J. Math. Phys.* **58**, 103501, 2017.
- [43] X. Shi, P. Lv, and C. Qi, Explicit solutions to a nonlocal 2-component complex modified Korteweg-de Vries equation, *Appl. Math. Lett.* **100**, 106043, 2020.
- [44] X.-D. Luo, Inverse scattering transform for the complex reverse space-time nonlocal modified Korteweg-de Vries equation with nonzero boundary conditions and constant phase shift, *Chaos* **29**, 073118, 2019.
- [45] G. Zhang, Z. Yan, Inverse scattering transforms and soliton solutions of focusing and defocusing nonlocal mKdV equations with nonzero boundary conditions, *Phys. D* **402**, 132170, 2020.
- [46] M. Gürses, Nonlocal Fordy-Kulish equations on symmetric spaces, *Phys. Lett. A* **381**, 1791–1794, 2017.
- [47] M. Gürses, A. Pekcan, and K. Zheltukhin, Nonlocal hydrodynamic type of equations, *Commun. Nonlinear Sci. Numer. Simul.* **85**, 105242, 2020.
- [48] M. Gürses, A. Pekcan, Nonlocal KdV equations, *Phys. Lett. A* **384** (35), 126894, 2020.

- [49] J. Rao, Y. Zhang, A.S. Fokas, and J. He, Rogue waves of the nonlocal Davey-Stewartson I equation, *Nonlinearity* **31**, 4090–4107, 2018.
- [50] T. Xu, M. Li, Y. Huang, Y. Chen, and C. Yu, Nonsingular localized wave solutions for the nonlocal Davey-Stewartson I equation with zero background, *Modern Phys. Lett. B* **31** (35), 1750338, 2017.
- [51] Z.-X. Zhou, Darboux transformations global explicit solutions for nonlocal Davey-Stewartson I equation, *Stud. Appl. Math.* **141** (2) 186–204, 2018.
- [52] Y. Zhang, Y. Liu, Breather and lump solutions for nonlocal Davey-Stewartson II equation, *Nonlinear Dyn.* **96**, 107–113, 2019.
- [53] J. Rao, Y. Cheng, and J. He, Rational and semirational solutions of the nonlocal Davey-Stewartson equations, *Stud. Appl. Math.* **139** (4), 2017.
- [54] Y. Cao, J. Rao, D. Mihalache, and J. He, Semi-rational solutions for the (2+1)-dimensional nonlocal Fokas system, *Appl. Math. Lett.* **80**, 27–34, 2018.
- [55] X. Zhu, A coupled (2+1)-dimensional mKdV system and its nonlocal reductions, *Commun. Nonlinear Sci. Numer. Simul.* **91**, 105438, 2020.
- [56] N.-N. Li, H.-Q. Hao, and R. Guo, Darboux transformation and the exact solutions of the (2+1)-dimensional nonlocal complex modified Korteweg-de Vries and Maxwell-Bloch equations, *Modern Phys. Lett. B* **34** (22), 2050230, 2020.
- [57] Y.-K. Liu, B. Li, Rogue waves in the (2+1)-dimensional nonlinear Schrödinger equation with a parity-time symmetric potential, *Chin. Phys. Lett.* **34** (1), 010202, 2017.
- [58] X. Zhang, Y. Chen, and Y. Zhang, Breather, lump and X soliton solutions to nonlocal KP equation, *Comput. Math. Appl.* **74**, 2341–2347, 2017.
- [59] M. Gürses, A. Pekcan, (2+1)-dimensional local and nonlocal reductions of the negative AKNS system: Soliton solutions, *Commun. Nonlinear Sci. Numer. Simul.* **71**, 161–173, 2019.
- [60] M. Gürses, A. Pekcan, (2+1)-Dimensional AKNS($-N$) Systems: $N = 3, 4$, arXiv:1910.11298 [nlin.SI].
- [61] R. Hirota, Exact solution of the Korteweg-de Vries equation for multiple collisions of solitons, *Phys. Rev. Lett.* **27**, 1192, 1971.
- [62] R. Hirota, *The Direct Method in Soliton Theory*, Cambridge University Press, Cambridge, 2004.
- [63] J. Hietarinta, Introduction to the Bilinear Method, in: *Integrability of Nonlinear Systems*, eds. Y. Kosman-Schwarzbach, B. Grammaticos, and K.M. Tamizhmani, Springer Lecture Notes in Physics **495**, 95–103, 1997.
- [64] J.-P. Yu, Y.-L. Sun, Study of lump solutions to dimensionally reduced generalized KP equations, *Nonlinear Dyn.* **87**, 2755–2763, 2017.

- [65] J.-P. Yu, Y.-L. Sun, A direct Bäcklund transformation for a (3+1)-dimensional Kadomtsev-Petviashvili-Boussinesq-like equation, *Nonlinear Dyn.* **90**, 2263–2768, 2017.
- [66] Y.-L. Sun, W.-X. Ma, J.-P. Yu, and C.M. Khalique, Dynamics of lump solitary wave of Kadomtsev-Petviashvili-Boussinesq-like equation, *Comput. Math. Appl.* **78**, 840–847, 2019.
- [67] W. Gai-Hua, W. Li-Hong, R. Ji-Guang, and H. Jing-Song, New patterns of the two-dimensional rogue waves: (2 + 1)-dimensional Maccari system, *Commun. Theor. Phys.* **67**, 601, 2017.
- [68] H.M. Baskonus, T.A. Sulaiman, and H. Bulut, On the novel wave behaviors to the coupled nonlinear Maccari’s system with complex structure, *Optik* **131**, 1036, 2017.
- [69] M. Shakeel, S.T. Mohyud-Din, and M.A. Iqbal, Closed form solutions for coupled nonlinear Maccari system, *Comput. Math. Appl.* **76** (4) 799, 2018.
- [70] N. Chema, S. Chen, and A.R. Seadawy, Propagation of isolated waves of coupled nonlinear (2 + 1)-dimensional Maccari system in plasma physics, *Results Phys.* **17**, 102987, 2020.
- [71] Y. Jiang, D.-Q. Xian, and X.-R. Kang, Homoclinic breather and rogue wave solutions to Maccari equation, *Comput. Math. Appl.* **79** (7), 1890–1894, 2020.
- [72] S. Kumar, M. Niwas, and A.-M. Wazwaz, Lie symmetry analysis, exact analytical solutions, and dynamics of solitons for (2+1)-dimensional NNV equations, *Phys. Scr.* **95** (9), 095204, 2020.
- [73] S. Kumar, S. Rani, Lie symmetry reductions and dynamics of soliton solutions of (2+1)-dimensional Pavlov equation, *Pramana* **94**, Art. No. 116, 2020.
- [74] S. Kumar, A. Kumar, and H. Kharbanda, Lie symmetry analysis and generalized invariant solutions of (2+1)-dimensional dispersive long wave (DLW) equations, *Phys. Scr.* **95** (6), 065207, 2020.
- [75] S. Kumar, A. Kumar, Lie symmetry reductions and group invariant solutions of (2+1)-dimensional modified Veronese web equation, *Nonlinear Dyn.* **98** (3), 1891–1903, 2019.
- [76] S. Kumar, A. Kumar, Dynamical structures of solitons and some new types of exact solutions for the (2+1)-dimensional DJKM equation using Lie symmetry analysis, *Modern Phys. Lett. B* **34**, 2150015, 2020.