On the Truth of Gödelian and Rosserian Sentences

ZIBA ASSADI

Independent Scholar, Tabriz, IRAN. E-mail:assadi.golzar@gmail.com SAEED SALEHI Research Institute for Fundamental Sciences (RIFS), University of Tabriz, P.O.Box 51666–16471, Tabriz, IRAN. E-mail:salehipour@tabrizu.ac.ir

Abstract

There is a longstanding debate in the logico-philosophical community as to why the Gödelian sentences of a consistent and sufficiently strong theory are true. The prevalent argument seems to be something like this: since every one of the Gödelian sentences of such a theory is equivalent to the theory's consistency statement, even provably so inside the theory, the truth of those sentences follows from the consistency of the theory in question. So, Gödelian sentences of consistent theories should be true. In this paper, we show that Gödelian sentences of only sound theories are true; and there is a long road from consistency to soundness, indeed a hierarchy of conditions which are satisfied by some theories and falsified by others. We also study the truth of Rosserian sentences and provide necessary and sufficient conditions for the truth of Rosserian (and also Gödelian) sentences of theories.

Keywords: The Incompleteness Theorem; Gödelian Sentences, Rosser's Trick, Rosserian Sentences, Soundness, Consistency, Σ_n -Soundness.

2020 AMS MSC: 03F40.

1 Introduction

By the first incompleteness theorem of Gödel (1931), for every consistent and sufficiently strong arithmetical theory there are sentences which are undecidable in the theory ([5]). Examples of such undecidable sentences are actually constructed in Gödel's original proof (hence *Gödelian sentences*) each of which is equivalent to its own unprovability in the theory; see Definition 2.1 below. A natural question here is that while the theory in question cannot decide the truth of its Gödelian sentences, what about us (humans)? Can we "see" (or demonstrate) their truth? This question has attracted the attention of many philosophers, physicists, computer scientists, as well as mathematical logicians. As there are numerous papers and books on this subject, it is not possible to cite them all here; see the Conclusions for a few. In this paper, we present necessary and sufficient conditions for the truth of Gödelian sentences (and Rosserian sentences) of consistent and sufficiently strong arithmetical theories (see the diagram in the Conclusions section).

We assume familiarity with the notions of Π_n and Σ_n formulae, Robinson's Arithmetic Q, and the fact that Q is a sound and Σ_1 -complete theory (i.e., every Q-provable sentence is true and every true Σ_1 -sentence is Q-provable); see e.g. [9]. By the Diagonal Lemma of Gödel and Carnap, see e.g. [24], for every formula $\Psi(x)$ with the only free variable x, there exists a sentence θ such that $\theta \leftrightarrow \Psi(\#\theta)$ is true (in the standard model of natural numbers \mathbb{N}) and also provable in Q; here #A denotes the numeral of the Gödel code of A, relative to

 \diamond

a fixed Gödel numbering (arithmetization) of the syntax. Moreover, if $\Psi(x)$ is a Π_n -formula, then θ can be taken to be a Π_n -sentence; and if $\Psi(x)$ is Σ_n , then θ can be taken to be Σ_n too. We provide more details in the following:

LEMMA 1.1 (The Diagonal Lemma): Let $n \ge 1$.

For every Π_n -formula $\Psi(x)$ there exists a Π_n -sentence θ such that $\boldsymbol{Q} \vdash \theta \leftrightarrow \Psi(\#\theta)$.

And for every Σ_n -formula $\Psi(x)$ there exists a Σ_n -sentence θ with the above property.

PROOF: There exists a Σ_1 -formula $\delta(x, y)$, in the language of arithmetic, that strongly represents the diagonal (primitive recursive) function **d** which assigns to a given m the Gödel code of the expression that results from substituting \overline{m} (the numeral of m, a term in the language of arithmetic representing m) for all the free variables (if any) of the expression coded by m. So, for every $m \in \mathbb{N}$ we have $\mathbf{Q} \vdash \forall y[\delta(\overline{m}, y) \leftrightarrow y = \overline{\mathbf{d}(m)}]$.

If $\Psi(x)$ is a Π_n -formula, then put $\alpha(x) = \forall y[\delta(x, y) \to \Psi(y)]$, and let a be its Gödel code. Now, let $\theta = \alpha(\overline{a})$; then θ is a Π_n -sentence and we have provably in \boldsymbol{Q} that

$$\theta \!=\! \forall y [\delta(\overline{\mathbf{a}}, y) \!\rightarrow\! \Psi(y)] \!\leftrightarrow\! \forall y [y \!=\! \overline{\mathbf{d}(\mathbf{a})} \!\rightarrow\! \Psi(y)] \!\leftrightarrow\! \forall y [y \!=\! \# \theta \!\rightarrow\! \Psi(y)] \!\leftrightarrow\! \Psi(\# \theta).$$

If $\Psi(x)$ is a Σ_n -formula, then put $\eta(x) = \exists y [\delta(x, y) \land \Psi(y)]$, and let e be its Gödel code. Now, let $\theta = \alpha(\overline{e})$; then θ is a Σ_n -sentence and we have provably in Q that

$$\theta = \exists y [\delta(\overline{\mathbf{e}}, y) \land \Psi(y)] \leftrightarrow \exists y [y = \overline{\mathbf{d}(\mathbf{e})} \land \Psi(y)] \leftrightarrow \exists y [y = \#\theta \land \Psi(y)] \leftrightarrow \Psi(\#\theta).$$

This lemma is one of the breakthroughs of Gödel's theorem (and of modern logic). The incompleteness theorem is usually stated for recursively enumerable (RE) theories that extend Q; though it also holds for more general theories, see e.g. [23]. If a theory is RE, then it can be defined by a Σ_1 -formula, see [9, Theorem 3.3.], and so its theorems (i.e., provable sentences) can also be defined by a Σ_1 -formula.

CONVENTION 1.2: Throughout, let T be an RE extension of Q. Let $\Pr_T(x)$ be a provability predicate for T, which is a Σ_1 -formula, relative to a fixed Gödel numbering. A basic property of $\Pr_T(x)$ is that for every sentence φ we have

$$T \vdash \varphi \iff \mathbb{N} \vDash \Pr_T(\# \varphi) \iff \boldsymbol{Q} \vdash \Pr_T(\# \varphi)$$

Let $\operatorname{Con}_T = \neg \operatorname{Pr}_T(\#[0 \neq 0])$ be the $(\Pi_1$ -)sentence that expresses T's consistency.

2 Gödel Sentences and their Truth.

Gödel's proof of his incompleteness theorem uses the diagonal lemma for the negation of the provability predicate of T.

DEFINITION 2.1 (Gödelian Sentences): A sentence γ is called a Gödelian sentence of T when we have $T \vdash \gamma \leftrightarrow \neg \Pr_T(\#\gamma)$.

Since for sufficiently strong theories T any two Gödelian sentences are T-provably equivalent, see e.g. [28, Remark 2.2.5.], many authors talk of *the* Gödel sentence of T. However, we will show that every unsound theory has both true and false Gödel sentences; so, even though the theory proves (unsoundly) that those sentences are equivalent, in reality they are not. Gödel [5], and several authors after him, e.g. [28, p. 825], [29, p. 7], or [17, p. 6], argue that Gödelian sentences of a sufficiently strong theory are true, since

(1) they are provably equivalent with their unprovability in the theory, and

(2) they are indeed unprovable in the theory; and so

(3) they must be true.

It is argued in [12] that this line of reasoning does not demonstrate the truth of Gödelian sentences, and indeed some (Σ_1 -)unsound theories may have false Gödelian sentences. In fact, the assumption (2) in the above argument (of Gödel) is redundant:

LEMMA 2.2: If T is consistent, then for every sentence φ , $T \vdash \varphi \rightarrow \neg \Pr_T(\#\varphi)$ implies $T \nvDash \varphi$.

PROOF: Since $T \vdash \varphi$ would imply on the one hand $T \vdash \neg \Pr_T(\#\varphi)$ by the assumption, and on the other hand $T \vdash \Pr_T(\#\varphi)$ by Convention 1.2.

So, the question of the validity of the above reasoning for the truth of Gödelian sentences boils down to the following question:

Does $T \vdash \gamma \leftrightarrow \neg \Pr_T(\#\gamma)$, for a consistent T, imply that $\mathbb{N} \models \gamma$?

Or put in another way, under which conditions all the Gödelian sentences of T are true?

We answer this question in the present section, and in the next section we answer a similar question for the Rosserian sentences (of arithmetical theories). Let us start with a characterization of the unprovable sentences:

PROPOSITION 2.3 (Characterizing Unprovable Sentences): Suppose that Löb's Rule holds for T. The following are equivalent for every sentence φ :

(1) φ is unprovable in *T*, i.e., $T \nvDash \varphi$;

(2) φ is a Gödelian sentence of some consistent extension U of T;

(3) $T + [\varphi \leftrightarrow \neg \Pr_T(\#\varphi)]$ is consistent.

PROOF:

 $(1 \Rightarrow 2)$: There exists, by Lemma 1.1, a sentence ξ such that

 $T \vdash \xi \leftrightarrow [\varphi \leftrightarrow \neg \Pr_{T+\xi}(\#\varphi)].$

Let $U = T + \xi$; then $U \vdash \varphi \leftrightarrow \neg \Pr_U(\#\varphi)$ and it remains to show that U is consistent. If not, then $T \vdash \neg \xi$. So, on the one hand we have (i) $T \vdash \neg [\varphi \leftrightarrow \neg \Pr_U(\#\varphi)]$, and on the other hand $U \vdash \varphi$ which implies (ii) $T \vdash \Pr_U(\#\varphi)$ by Convention 1.2. Now, (i) and (ii) imply that $T \vdash \varphi$, contradicting the assumption.

 $(2 \Rightarrow 3)$: If $T + [\varphi \leftrightarrow \neg \Pr_T(\#\varphi)]$ is not consistent, then $T \vdash \neg [\varphi \leftrightarrow \neg \Pr_T(\#\varphi)]$, and so $T \vdash \Pr_T(\#\varphi) \rightarrow \varphi$, which implies $T \vdash \varphi$ by Löb's Rule. So, for every extension U of T we have $U \vdash \varphi$, and so, by Convention 1.2, $U \vdash \Pr_U(\#\varphi)$. Therefore, for every such U we have $U \vdash \neg [\varphi \leftrightarrow \neg \Pr_U(\#\varphi)]$, which contradicts the assumption.

 $(3 \Rightarrow 1)$: If $T \vdash \varphi$, then, by Convention 1.2, we have that $T \vdash \Pr_T(\#\varphi)$, and so we should have also $T \vdash \neg[\varphi \leftrightarrow \neg \Pr_T(\#\varphi)]$.

It should be noted that the assumption of holding Löb's Rule for T was used only in the implication $(2 \Rightarrow 3)$. So, (1) and (2) are equivalent with each other, and are implied by (3), even when this rule does not hold. We now provide a necessary and sufficient condition, on a sufficiently strong T, for the truth of all the Gödelian Π_1 -sentences of T: **THEOREM** 2.4 (On the Truth and Independence of Gödelian Π_1 -Sentences): Suppose that *T* satisfies the following two conditions:

- (I) $T \vdash \neg \operatorname{Con}_T \rightarrow \operatorname{Pr}_T(\#\varphi)$ for every sentence φ , and
- (II) $T \vdash \operatorname{Con}_T \to \neg \operatorname{Pr}_T(\#\gamma)$ for every Gödelian sentence γ of T.

If $T \vdash \neg \text{Con}_T$, then every false Π_1 -sentence is a Gödelian sentence of T, and no Gödelian sentence of T is independent from T.

If $T \nvDash \neg Con_T$, then all the Gödelian Π_1 -sentences of T are true, and all the Gödelian sentences of T are independent from T.

PROOF: If $T \vdash \neg \text{Con}_T$, then by (I) we have $T \vdash \Pr_T(\#\varphi)$ for every sentence φ . So, for every Gödelian sentence γ of T we have $T \vdash \neg \gamma$; thus no Gödelian sentence of T can be independent from T. Now, let ϕ be an arbitrary false Π_1 -sentence; then $\neg \phi$ is a true Σ_1 -sentence, and so provable in Q. Thus, $T \vdash \neg \phi$; and so from $T \vdash \Pr_T(\#\phi)$ we have $T \vdash \phi \leftrightarrow \neg \Pr_T(\#\phi)$, which means that ϕ is a (false) Gödelian Π_1 -sentence of T.

If $T \nvDash \neg \operatorname{Con}_T$, then by (II) for every Gödelian sentence γ of T we have $T \nvDash \operatorname{Pr}_T(\#\gamma)$, and so $T \nvDash \neg \gamma$; thus γ is independent from T (noting that T is consistent and so we also have $T \nvDash \gamma$ by Lemma 2.2). If a Gödelian Π_1 -sentence τ of T is not true, then $\neg \tau$ is a true Σ_1 -sentence, and so should be Q-provable; a contradiction with the T-independence of τ proved above.

Every extension of Peano's Arithmetic **PA** satisfies (I) and (II) in Theorem 2.4; as a matter of fact (II) is a formalization of Gödel's first incompleteness theorem (in T). If T is $(\Sigma_1$ -)sound, then $T \nvDash \neg \operatorname{Con}_T$. If T is inconsistent or $T = S + \neg \operatorname{Con}_S$, where S is a consistent extension of **PA**, then $T \vdash \neg \operatorname{Con}_T$; in the latter case T is consistent by Gödel's second incompleteness theorem. Thus, by Theorem 2.4, a necessary and sufficient condition for the truth of all the Gödelian Π_1 -sentences of T is the consistency of T with Con_T , a condition obviously implied by ω -consistency; though, this condition is stronger than the mere consistency of T, see [8, Theorem 36]. For investigating the truth of Gödelian Π_{n+1} -sentences (and Σ_{n+1} -sentences) we make a definition and an observation. Before that let us note that no Gödelian Σ_1 -sentence of a consistent extension of **Q** can be true:

PROPOSITION 2.5 (On the Truth of Gödelian Σ_1 -Sentences): No Gödelian Σ_1 -sentence of T can be true if T is consistent.

PROOF: If a Gödelian Σ_1 -sentence of T were true, then it would have been provable in Q, and this would have contradicted Lemma 2.2 for consistent T.

DEFINITION 2.6 (Γ -Soundness): Let Γ be a class of sentences. A theory *S* is called Γ -sound when every *S*-provable Γ -sentence is true.

The following lemma has been proved for $\Gamma = \Sigma_1, \Sigma_2$ in [8, Theorems 25, 27, 30,32]:

LEMMA 2.7 (On Extensions of Γ -Sound Theories): Let Γ be a class of sentences that is closed under disjunction. If T is a Γ -sound theory, then for every sentence φ , either $T + \varphi$ or $T + \neg \varphi$ is Γ -sound.

PROOF: If neither $T + \varphi$ nor $T + \neg \varphi$ is Γ -sound, then for some false Γ -sentences ς and ς' we have $T + \varphi \vdash \varsigma$ and $T + \neg \varphi \vdash \varsigma'$. Thus, $T \vdash \varsigma \lor \varsigma'$, and $\varsigma \lor \varsigma'$ is a false Γ -sentence; a contradiction.

One of our main results is the following necessary and sufficient condition for the truth of Gödelian (Π_{n+1} and Σ_{n+1}) sentences:

THEOREM 2.8 (The Truth of Gödel Sentences): Let $n \ge 1$.

All the Gödelian Π_{n+1} -sentences of T are true if and only if T is Π_{n+1} -sound. All the Gödelian Σ_{n+1} -sentences of T are true if and only if T is Σ_{n+1} -sound.

PROOF: Let Γ be any of Π_{n+1} or Σ_{n+1} .

First, suppose that T is Γ -sound, and let γ be a Gödelian Γ -sentence of the theory T. By Lemma 2.2 and Convention 1.2 we have $\mathbb{N} \models \neg \Pr_T(\#\gamma)$, and so $\Pr_T(\#\gamma)$ is a false Σ_1 -sentence. Now, $T + \neg \gamma \vdash \Pr_T(\#\gamma)$, and so $T + \neg \gamma$ is not Σ_1 -sound; whence, it is not Γ -sound either. Thus, by Lemma 2.7, the theory $T + \gamma$ should be Γ -sound. Therefore, γ must be true.

Now, suppose that all the Gödelian Γ -sentences of T are true. We show that the theory T is Γ -sound. Assume that $T \vdash \varsigma$ for a Γ -sentence ς . We prove that ς is true. By Lemma 1.1 there exists a Γ -sentence ζ such that $\mathbf{Q} \vdash \zeta \leftrightarrow [\varsigma \land \neg \Pr_T(\#\zeta)]$. Thus, from $T \vdash \varsigma$ we have $T \vdash \zeta \leftrightarrow \neg \Pr_T(\#\zeta)$, and so ζ is a Gödelian Γ -sentence of T. Whence, ζ is true, and so, by the soundness of \mathbf{Q} , we have $\mathbb{N} \models \varsigma$.

Whence, all the Gödelian sentences of a theory are true if and only if the theory is sound; cf. also [27, Theorem 24.7.].

REMARK 2.9 (On the Hierarchy of Π_n , Σ_n -Soundness): Let us note that an extension of Q is consistent if and only if it is Π_1 -sound: indeed, no consistent extension of Q can prove a false Π_1 -sentence, since the negation of such a sentence would be a true Σ_1 -sentence and so would be provable in Q.

One can also show that a theory is Σ_n -sound if and only if it is Π_{n+1} -sound: if the theory S is Σ_n -sound and $S \vdash \pi$, where π is a Π_{n+1} -sentence, then write $\pi = \forall x \sigma(x)$ for a Σ_n -formula σ ; since for every $k \in \mathbb{N}$ we have $S \vdash \sigma(\overline{k})$, and $\sigma(\overline{k})$ is a Σ_n -sentence, then $\mathbb{N} \models \sigma(\overline{k})$ for every $k \in \mathbb{N}$, so $\mathbb{N} \models \forall x \sigma(x) = \pi$.

The hierarchy of Σ_n -sound theories is strict, since there are some Σ_n -sound theories which are not Σ_{n+1} sound; this is proved in e.g. [23, Theorem 2.5.] and also [10, Theorem 4.8.]. Therefore, the truth of (even all) the Gödelian Π_{n+1} -sentences (respectively, Σ_{n+1} -sentences) of a theory does not necessarily imply the truth of its Gödelian Π_{n+2} -sentences (respectively, Σ_{n+2} -sentences).

3 Rosserian Sentences and their Truth.

In Theorem 2.4 we saw that Gödelian sentences of some theories could be refutable in them (though, they are always unprovable in consistent theories, see Lemma 2.2). Rosser's trick ([22]) constructs an independent sentence for a given theory (which is an RE extension of Q), when it is consistent. Before going into Rosser's construction, let us note that no construction similar to Gödel's can result in an independent sentence.

DEFINITION 3.1 (Pseudo-Gödelian Sentences): Let us call ψ a pseudo-Gödelian sentence of a theory T when there are some propositional formulas $C_1(p), \dots, C_n(p)$, over the one propositional variable p, and there is one propositional formula $B(p_1, \dots, p_n)$, over the propositional variables p_1, \dots, p_n , such that we have $T \vdash \psi \leftrightarrow B(\Pr_T[\#C_1(\psi)], \dots, \Pr_T[\#C_n(\psi)])$.

For example, the sentences \mathfrak{P}_T and \mathfrak{R}_T for which we have

$$T \vdash \mathfrak{Pr}_T \leftrightarrow \left[\neg \mathtt{Pr}_T(\#\mathfrak{P}_T) \land \neg \mathtt{Pr}_T(\#[\neg \mathfrak{P}_T])\right] \text{ and } T \vdash \mathfrak{R}_T \leftrightarrow \left[\mathtt{Pr}_T(\#\mathfrak{R}_T) \rightarrow \neg \mathtt{Pr}_T(\#[\neg \mathfrak{R}_T])\right],$$

are both some peudo-Gödelian sentences of T. For an alternative formulation of the following result see [29, Exercise 1, p. 149].

 \diamond

PROPOSITION 3.2 (Decidability of Pseudo-Gödelian Sentences): No pseudo-Gödelian sentence of $U = T + \neg \text{Con}_T$ can be independent from U.

PROOF: Let ψ be a pseudo-Gödelian sentence of the theory $U = T + \neg \text{Con}_T$. For every sentence ζ we have $U \vdash \Pr_U(\#\zeta)$. Now, $B(\top, \dots, \top)$, where \top denotes the propositional truth, is equivalent to either \top or \bot , where \bot denotes the propositional falsum. If $B(\top, \dots, \top) \equiv \top$, then $U \vdash B(\Pr_U[C_1(\#\psi)], \dots, \Pr_U[C_n(\#\psi)])$; and if $B(\top, \dots, \top) \equiv \bot$, then $U \vdash \neg B(\Pr_U[C_1(\#\psi)], \dots, \Pr_U[C_n(\#\psi)])$. As a result, we have either $U \vdash \psi$ or $U \vdash \neg \psi$; thus ψ is not independent from U.

In both of the above examples, it can be seen that $U \vdash \neg \mathfrak{P}_U$ and $U \vdash \neg \mathfrak{R}_U$ hold for the theory $U = T + \neg \operatorname{Con}_T$. Thus, for getting independent sentences (of consistent theories) one should go beyond the (pesudo-)Gödelian sentences.

The *T*-provability predicate $\Pr_T(x)$ in Convention 1.2 is usually constructed from a *T*-proof predicate $\Pr_T(y, x)$, as $\Pr_T(x) = \exists y \Pr_T(y, x)$; where $\Pr_T(y, x)$ is a Δ_1 -formula stating that "y is (the Gödel code of) a proof in *T* of the formula (coded by) x".

CONVENTION 3.3: Let us fix a proof predicate of T as the Δ_1 -formula $prf_T(y, x)$ that satisfies the following for every sentence φ :

$$\begin{array}{ll} T \vdash \varphi & \Longleftrightarrow & \boldsymbol{\mathcal{Q}} \vdash \operatorname{prf}_{T}(\overline{m}, \#\varphi) \text{ for some } m \in \mathbb{N}. \\ T \nvDash \varphi & \Longleftrightarrow & \boldsymbol{\mathcal{Q}} \vdash \neg \operatorname{prf}_{T}(\overline{n}, \#\varphi) \text{ for every } n \in \mathbb{N}. \end{array}$$

DEFINITION 3.4 (Rosserian Provability and Rosserian Sentences): The following Σ_1 -formula, with the free variable *x*, is the Rosserian Provability predicate of *T*:

$$\texttt{R.Pr}_T(x) = \exists y \, [\texttt{prf}_T(y, x) \land \forall z < y \, \neg \texttt{prf}_T(z, \neg x)].$$

A sentence ρ is called a Rosserian sentence of T when we have $T \vdash \rho \leftrightarrow \neg R.Pr_T(\#\rho)$.

The independence of the Rosserian sentences (from the theory in question) follows form the following basic properties of the Rosserian provability:

LEMMA 3.5: If T is consistent, then for every sentence φ we have

- (1) $T \vdash \varphi \iff \boldsymbol{Q} \vdash \operatorname{R.Pr}_T(\#\varphi).$
- (2) $T \vdash \neg \varphi \implies \boldsymbol{Q} \vdash \neg \mathtt{R}.\mathtt{Pr}_T(\#\varphi).$

PROOF: For (1) it suffices to note that for a consistent theory T we have: $T \vdash \varphi$ if and only if the Σ_1 sentence $\mathbb{R}.\Pr_T(\#\varphi)$ is true. For (2) suppose that $T \vdash \neg \varphi$; then by Convention 3.3 for some m we have $\mathbf{Q} \vdash \Pr_T(\overline{m}, \#[\neg \varphi])$. Now, reason inside \mathbf{Q} : for any y with $\Pr_T(y, \#\varphi)$ we should have $y > \overline{m}$, since no $i \leq \overline{m}$ (which are $i = 0, \dots, \overline{m}$) could satisfy $\Pr_T(\overline{i}, \#\varphi)$ by Convention 3.3, and so for some z < y, which is $z = \overline{m}$, we have $\Pr_T(z, \#[\neg \varphi])$. Thus, we have $\forall y[\Pr_T(y, \#\varphi) \to \exists z < y \Pr_T(z, \#[\neg \varphi])]$ or, equivalently, $\neg \mathbb{R}.\Pr_T(\#\varphi)$.

Now, we can characterize the independent sentences of T similarly to Proposition 2.3:

PROPOSITION 3.6 (Characterizing Independent Sentences): Let φ be a sentence.

The following are equivalent:

(1) φ is independent from T, i.e., $T \nvDash \varphi$ and $T \nvDash \neg \varphi$;

(2) φ is a Rosserian sentence of some consistent extension U of T;

and are implied by the following:

(3) $T + [\varphi \leftrightarrow \neg \mathbb{R}.\Pr_T(\#\varphi)]$ is consistent.

PROOF: First we show the equivalence of (1) and (2).

 $(1 \Rightarrow 2)$: By Lemma 1.1 for some sentence ξ we have $T \vdash \xi \leftrightarrow [\varphi \leftrightarrow \neg \mathbb{R}.\Pr_{T+\xi}(\#\varphi)]$. Let $U = T + \xi$; then $U \vdash \varphi \leftrightarrow \neg \mathbb{R}.\Pr_U(\#\varphi)$ which shows that φ is a Rosserian sentence of U. We show that U is consistent. If not, then $T \vdash \neg \xi$. Thus, (*) $T \vdash \neg [\varphi \leftrightarrow \neg \mathbb{R}.\Pr_U(\#\varphi)]$. Also, $U \vdash \varphi$ and $U \vdash \neg \varphi$, and so by Convention 3.3 there are $m, n \in \mathbb{N}$ such that $\mathbf{Q} \vdash \operatorname{prf}_U(\overline{m}, \#\varphi)$ and $\mathbf{Q} \vdash \operatorname{prf}_U(\overline{n}, \#[\neg \varphi])$; we can assume that m and n are the least such numbers.

(i) If $m \leq n$, then $\mathbf{Q} \vdash \operatorname{prf}_U(\overline{m}, \#\varphi) \land \forall z < \overline{m} \neg \operatorname{prf}_U(z, \#[\neg \varphi])$ and so $\mathbf{Q} \vdash \operatorname{R.Pr}_U(\#\varphi)$, which implies by (*) that $T \vdash \varphi$; contradicting (1).

(ii) If n < m, then $\mathbf{Q} \vdash \forall y [\operatorname{prf}_U(y, \#\varphi) \rightarrow y \ge \overline{m} \rightarrow \exists z < y \operatorname{prf}_U(z, \#[\neg \varphi])]$, since one can take z = n, and so $\mathbf{Q} \vdash \neg \mathbb{R}.\operatorname{Pr}_U(\#\varphi)$, which implies by (*) that $T \vdash \neg \varphi$; contradicting (1) again.

So, U must be consistent.

 $(2 \Rightarrow 1)$: It suffices to show that φ is independent from U. If $U \vdash \rho$, then we should have on the one hand $U \vdash \mathbb{R}.\Pr_U(\#\rho)$ by Lemma 3.5(1), and on the other hand $U \vdash \neg \mathbb{R}.\Pr_U(\#\rho)$ by Definition 3.4; thus, U could not be consistent. Also $U \vdash \neg \rho$ would imply on the one hand $U \vdash \neg \mathbb{R}.\Pr_U(\#\rho)$ by Lemma 3.5(2), and on the other hand $U \vdash \mathbb{R}.\Pr_U(\#\rho)$ by Lemma 3.5(2), and on the other hand $U \vdash \mathbb{R}.\Pr_U(\#\rho)$ by Definition 3.4; contradicting U's consistency again.

Now, we show that (3) implies (1); and so (2) too.

 $(3 \Rightarrow 1)$: Note that the theory T is consistent by the assumption. If $T \vdash \varphi$, then by Lemma 3.5(1) we would have $T \vdash \mathbb{R}.\Pr_T(\#\varphi)$, and so $T \vdash \neg[\varphi \leftrightarrow \neg \mathbb{R}.\Pr_T(\#\varphi)]$. If $T \vdash \neg\varphi$, then Lemma 3.5(2) would imply that $T \vdash \neg \mathbb{R}.\Pr_T(\#\varphi)$, and so we would have $T \vdash \neg[\varphi \leftrightarrow \neg \mathbb{R}.\Pr_T(\#\varphi)]$ again.

REMARK 3.7 (Löb-Like Rule for Rosserian Provability): Let us note that the contraposition of the implication $(1 \Rightarrow 3)$ in Proposition 3.6 says that if $T \vdash \varphi \leftrightarrow \mathbb{R}.\Pr_T(\#\varphi)$, i.e., if φ is a Rosser-type Henkin sentence so called by Kurahashi (in [11]), then φ is not independent from T. Actually, it is shown in [11] that there are *standard* proof predicates $\operatorname{prf'}_T(y, x)$ which have independent Rosser-type Henkin sentences, and there are standard proof predicates $\operatorname{prf'}_T(y, x)$ none of whose Rosser-type Henkin sentences are independent. The latter proof predicates $\operatorname{satisfy}(1 \Rightarrow 3)$ in Proposition 3.6 and satisfy a Löb-like rule for Rosserian provability; while the former ones do not satisfy $(1 \Rightarrow 3)$ in Proposition 3.6, and do not satisfy any Löb-like rule for Rosserian provability. So, the implication $(1 \Rightarrow 3)$ in Proposition 3.6 depends on the proof predicate $\operatorname{prf}_T(y, x)$, and is not robust.

Unlike Gödelian Π_1 -sentences, all the Rosserian Π_1 -sentences of consistent theories are true, and like Gödelian Σ_1 -sentence, all of their Rosserian Σ_1 -sentences are false:

THEOREM 3.8 (On the Truth of the Rosserian Π_1, Σ_1 -Sentences): Every Rosserian Π_1 -sentence of T is true, and every Rosserian Σ_1 -sentence of T is false, if T is consistent.

PROOF: If a Rosserian Π_1 -sentence were false, then its negation would be a true Σ_1 -sentence, and so would be provable in the theory Q; contradicting Rosser's theorem on the independence of Rosserian sentences

(Proposition 3.6). If a Rosserian Σ_1 -sentence were true, then it would be provable in the theory Q; contradicting the unprovability of Rosserian sentences.

However, for $n \ge 1$, the truth of all the Gödelian \prod_{n+1} -sentences is equivalent to the truth of all the Rosserian \prod_{n+1} -sentences; and the truth of all the Gödelian \sum_{n+1} -sentences is equivalent to the truth of all the Rosserian \sum_{n+1} -sentences:

THEOREM 3.9 (On the Truth of the Rosserian Π_{n+1} , Σ_{n+1} -Sentences): Fix $n \ge 1$.

All the Rosserian Π_{n+1} -sentences of T are true if and only if T is Π_{n+1} -sound.

All the Rosserian Σ_{n+1} -sentences of T are true if and only if T is Σ_{n+1} -sound.

PROOF: This can be proved in the same lines of the proof of Theorem 2.8 by using Lemma 3.5 (instead of Lemma 2.2) which implies that $\mathbb{R}.\Pr_T(\#\rho)$ is a false Σ_1 -sentence when ρ is a Rosserian sentence of consistent theory T.

Whence, all the Rosserian sentences of T are true if and only if T is sound; cf. [27, Theorem 24.7.].

4 Conclusions.

The first one who talked about the truth of Gödelian sentences was Gödel himself [5]. This turned into a serious debate with [6] in which (what we call now) the *Gödel Disjunction* was announced; see [4] and [7] and the references therein. The so called *Anti-Mechanism Thesis*, or the *Lucas-Penrose Argument*, started with [13] and popularized by [16]; see also [15] and [20]. After that, there has been a large discussion on the truth of Gödelian sentences; see e.g. [3], [28], [1], [26], [30], [21], [17], [14], [25], [8], [2], [18] and [19]. As shown above, the consistency of a theory need not imply the truth of (all of) its Gödelian (Π_1 -)sentences; but does imply the truth of its all Rosserian Π_1 -sentences. One wonders why the proponents of the anti-mechanism thesis have not used the Rosserian (Π_1 -)sentences for their reasoning; since the truth of those sentences are straightforward (and immediately follows from the consistency of the theory). Though, the opponents have argued that actually for "seeing" the truth of Gödelian (Π_1 -)sentences one should "see" (at least) the consistency of the theory (and indeed, more than that). Our old and new results are summarized in the following diagram; note that the conditions get (strictly) stronger from bottom to top.

Soundness \equiv Truth of Gödelian and Rosserian Sentences \vdots $\Sigma_{n+1} (\Pi_{n+2})$ Soundness \equiv Truth of Gödelian, Rosserian Σ_{n+1}, Π_{n+2} Sentences \vdots $\Sigma_2 (\Pi_3)$ Soundness \equiv Truth of Gödelian, Rosserian Σ_2, Π_3 Sentences $\Sigma_1 (\Pi_2)$ Soundness \equiv Truth of Gödelian, Rosserian Π_2 Sentences Consistency of $T + \text{Con}_T \equiv$ Truth of Gödelian Π_1 Sentences Consistency (Π_1 Soundness) \equiv Falsity of Gödelian, Rosserian Σ_1 Sentences

Acknowledgements. This research is supported by the grant $\mathcal{N}^{\underline{O}}$ 98013437 of the Iran National Science Foundation (INSF). The authors warmly thank KAAVE LAJEVARDI for the most helpful discussions and comments; this is a continuation of a project that he started a while ago and which have resulted in [12] and some other works to appear in the near future.

References

- BOOLOS, GEORGE (1990); On "Seeing" the Truth of the Gödel Sentence, *Behavioral and Brain Sciences* 13(4):655–656. Reprinted in: R. Jeffrey (ed.), G. Boolos, *Logic, Logic and Logic*, Harvard University Press (1999) pp. 389–391.
- [2] BOYER, JULIEN & SANDU, GABRIEL (2012); Between Proof and Truth, *Synthese* 187(3):821–832. Erratum: *ibid* 973–974.
- [3] DUMMETT, MICHAEL (1963); The Philosophical Significance of Gödel's Theorem, *Ratio* 5:140–155. Reprinted in: M. Dummett, *Truth and Other Enigmas*, Harvard University Press (1978, 6th print. 1996) pp. 186–201.
- [4] FEFERMAN, SOLOMON (2006); Are There Absolutely Unsolvable Problems? Gödel's Dichotomy, *Philosophia Mathematica* 14(2):134–152.
- [5] GÖDEL, KURT (1931); Über Formal Unentscheidbare Sätze der Principia Mathematica und Verwandter Systeme I, *Monatshefte für Mathematik und Physik* 38(1):173–198 (in German). English Translation: "On Formally Undecidable Propositions of *Principia mathematica* and Related Systems I", in: S. Feferman et al. (eds.), *Kurt Gödel Collected Works, Volume I: Publications 1929–1936*, Oxford University Press (1986) pp. 135–152.
- [6] GÖDEL, KURT (1951); "Some Basic Theorems on the Foundations of Mathematics and their Implications", in: S. Feferman et al. (eds.), *Kurt Gödel Collected Works, Volume III: Unpublished Essays and Lectures*, Oxford University Press (1995) pp. 304–323.
- [7] HORSTEN, LEON & WELCH, PHILIP (eds.) (2016); Gödel's Disjunction: The Scope and Limits of Mathematical Knowledge, Oxford University Press.
- [8] ISAACSON, DANIEL (2011); "Necessary and Sufficient Conditions for Undecidability of the Gödel Sentence and its Truth", in: D. DeVidi & M. Hallett & P. Clarke (eds.), *Logic, Mathematics, Philosophy: Vintage Enthusiasms—Essays in Honour of John L. Bell*, Springer, pp. 135–152.
- [9] KAYE, RICHARD (1991); Models of Peano Arithmetic, Oxford University Press.
- [10] KIKUCHI, MAKOTO & KURAHASHI, TAISHI (2017); Generalizations of Gödel's Incompleteness Theorems for Σ_n -Definable Theories of Arithmetic, *The Review of Symbolic Logic* 10(4):603–616.
- [11] KURAHASHI, TAISHI (2016); Henkin Sentences and Local Reflection Principles for Rosser Provability, *Annals of Pure and Applied Logic* 167(2):73–94.
- [12] LAJEVARDI, KAAVE & SALEHI, SAEED (2019); On the Arithmetical Truth of Self-Referential Sentences, *Theoria*—A Swedish Journal of Philosophy 85(1):8–17.
- [13] LUCAS, JOHN R. (1961); Minds, Machines and Gödel, *Philosophy* 36(137):112–127. Reprinted in: K.M. Sayre & F.J. Crosson (eds.), *The Modeling of Mind*, University of Notre Dame Press (1963) pp. 255–271; and in: A.R. Anderson (ed.), *Minds and Machines*, Prentic-Hall (1964) pp. 43–59.
- [14] MILNE, PETER (2007); On Gödel Sentences and What They Say, *Philosophia Mathematica* 15(2):193–226.
- [15] NAGEL, ERNEST & NEWMAN, JAMES R. (1958); Gödel's Proof, New York University Press (revised: 3rd ed. Routledge 2005).

- [16] PENROSE, ROGER (1989); *The Emperor's New Mind*, Oxford University Press (republished with a new preface 1999).
- [17] PEREGRIN, JAROSLAV (2007); Gödel, Truth & Proof, Journal of Physics: Conference Series 82:012005.
- [18] PIAZZA, MARIO & PULCINI, GABRIELE (2015); "A Deflationary Account of the Truth of the Gödel Sentence G", in: G. Lolli & M. Panza & G. Venturi (eds.), From Logic to Practice, Springer, pp. 71–90.
- [19] PIAZZA, MARIO & PULCINI, GABRIELE (2016); "What's so Special About the Gödel Sentence G", in:
 F. Boccuni & A. Sereni (eds.), *Objectivity, Realism, and Proof*, Springer, pp. 245–263.
- [20] PUTNAM, HILLARY (1960); "Minds and Machines", in: S. Hood (ed.), Dimensions of Mind: A Symposium, New York University Press, pp. 138–164. Reprinted in: H. Putnam, Philosophical Papers, Volume 2: Mind, Language and Reality, Harvard University Press (1975) pp. 362–385.
- [21] RAATIKAINEN, PANU (2005); On the Philosophical Relevance of Gödel's Incompleteness Theorems, *Revue Internationale de Philosophie* 59(4):513–534.
- [22] ROSSER, BARKLEY (1936); Extensions of Some Theorems of Gödel and Church, *The Journal of Symbolic Logic* 1(3):87–91.
- [23] SALEHI, SAEED & SERAJI, PAYAM (2017); Gödel-Rosser's Incompleteness Theorem, Generalized and Optimized for Definable Theories, *Journal of Logic and Computation* 27(5):13911397.
- [24] SALEHI, SAEED (2020); On the Diagonal Lemma of Gödel and Carnap, *The Bulletin of Symbolic Logic*, to appear. DOI: 10.1017/bsl.2019.58
- [25] SERÉNY, GYÖRGY (2011); How do We Know that the Gödel Sentence of a Consistent Theory Is True?, *Philosophia Mathematica* 19(1):47–73.
- [26] SHAPIRO, STEWART (1998); Induction and Indefinite Extensibility: The Gödel Sentence Is True, But Did Someone Change the Subject?, *Mind* 107(427):597–624.
- [27] SMITH, PETER (2013); An Introduction to Gödel's Theorems, Cambridge University Press (2nd ed.).
- [28] SMORYŃSKI, CRAIG (1977); "The Incompleteness Theorems", in: J. Barwise (ed.), *Handbook of Mathematical Logic*, North-Holland, pp. 821–865.
- [29] SMORYŃSKI, CRAIG (1985); Self-Reference and Modal Logic, Springer.
- [30] TENNANT, NEIL (2001); On Turing Machines Knowing Their Own Gödel-Sentences, *Philosophia Mathematica* 9(1):72–79.