arXiv:2009.04610v2 [quant-ph] 13 Sep 2020

Sample efficient tomography via Pauli Measurements

Nengkun Yu^1

¹Centre for Quantum Software and Information, Faculty of Engineering and Information Technology, University of Technology, Sydney, NSW 2007, Australia (Dated: September 15, 2020)

Pauli Measurements are the most important measurements in both theoretical and experimental aspects of quantum information science. In this paper, we explore the power of Pauli measurements in the state tomography related problems. Firstly, we show that the quantum state tomography problem of *n*-qubit system can be accomplished with $\mathcal{O}(\frac{10^n}{\epsilon^2})$ copies of the unknown state using Pauli measurements. As a direct application, we studied the quantum overlapping tomography problem introduced by Cotler and Wilczek in Ref. [1]. We show that the sample complexity is $\mathcal{O}(\frac{10^k \cdot \log(\binom{n}{k}/\delta)}{\epsilon^2})$ for quantum overlapping tomography of *k*-qubit reduced density matrices among *n* is quantum system, where $1 - \delta$ is the confidential level, and ϵ is the trace distance error. This can be achieved using Pauli measurements. Moreover, we prove that $\Omega(\frac{\log(n\delta)}{\epsilon^2})$ copies are needed. In other words, for constant *k*, joint, highly entangled, measurements are not asymptotically more efficient than Pauli measurements.

INTRODUCTION

It is increasingly important to understand how the cost scales of learning useful information as the experiments can control larger and larger quantum systems. Quantum state tomography refers to a procedure of reconstructing the density matrix of a quantum state from various measurements on multiple copies of the state. This fundamental task is crucial for quantum information experiments and theoretically goes back at least to the work of Helstrom, Holevo, and others from around 1970.

In quantum state tomography, one is given k copies of an n-qubit quantum state ρ and is required to output a classical description of density matrix $\hat{\rho}$ close to ρ by performing quantum measurements on $\rho^{\otimes k}$. Broadly speaking, there are three categories of measurements: one consists of joint (entangled) measurements on $\rho^{\otimes k}$, as in [2–4]. In [5–7], the authors showed that the optimal scaling of the sample number k as a function of trace distance goal ϵ is $n \propto \frac{4^n}{\epsilon^2}$. The second category consists of measurements on each copy of the state ρ , whose results are to be combined to reconstruct the density matrix, as in [8]. Ref. [9] showed that if one can perform many-outcome measurements, tomography is possible using $n \propto 8^n/\delta^2$ copies, and this is optimal if the measurements on every copy are independent as showed in [5]. The third category local measurements consists of measurements on each qubit of each copy of the state ρ , whose results are to be combined to reconstruct the density matrix [10]. Generally, joint measurements over several copies of the state can usually achieve lower sampling rates [5–7], but are much more challenging to implement. On the other hand, Pauli measurements are experimental friendly, therefore, extremely important both theoretically and experimentally. If one is restricted to use Pauli measurements, Flammia, Gross, Liu and Eisert observed that the quantum state tomography can be accomplished using $\mathcal{O}(\frac{n \cdot 16^n}{\epsilon^2})$ copies in [11]. This was improved the bound to $\mathcal{O}(\frac{n \cdot 12^n}{\epsilon^2})$ in [10].

In [1], Cotler and Wilczek introduced the problem called quantum overlapping tomography problem. The goal is to output the classical description of all k-qubit reduced density matrix of an n-qubit system. This problem is also of great importance in quantum information science, because many important physical quantities, for instance, energy and entropy, depend on very small parts of the whole system only, in other words, depends on the set of local reduced density matrices. By using the perfect hash families, they showed that one only needs to use $e^{\mathcal{O}(k)} \log^2 n$ rounds of parallel measurements to achieve this goal. Ref. [12] proposed a measurement scheme to perform two-qubit tomography of all pairs. Later Ref. [13] provided an algorithm to estimate the expectation value of m Pauli operators with weight $\leq k$ using $\mathcal{O}(3^k \log m)$ measurements for small k. All of these bounds can be achieved using Pauli measurements.

Our results

In this paper, we study the power of Pauli measurements, the most important class of measurements, in two tomography related problems. In the first part of this paper, we revisit the sample complexity of quantum state tomography problem,

Problem 1. Given an unknown n-qubit quantum mixed state ρ , the goal is to output density matrices σ such that

$$\|\rho - \sigma\|_1 < \epsilon \tag{1}$$

for a given $\epsilon > 0$. How many copies of ρ are needed to achieve this goal, with high probability?

The sample complexity of this problem is nearly resolved in the general joint measurement setting and independent measurement setting [5]. However, it is still unclear in the local measurement, in particular, the Pauli measurement setting. We show that

Theorem 1. The quantum state tomography problem can be solved using $\mathcal{O}(\frac{10^n \log \frac{1}{\delta}}{\epsilon^2})$ copies of ρ via Pauli measurement on each qubit to success with probability at least $1 - \delta$.

Secondly, we study the sample complexity of the quantum overlapping tomography problem: "How many copies of states are necessary and sufficient to reconstruct all the k-reduced density matrices of unknown n-qubit state ρ , to within additive error ϵ , for constant k?" More precisely, the formula of this problem is

Problem 2. Given an unknown n-qubit quantum mixed state $\rho_{1,2,...,n}$, and $1 \le k \le n$, the goal of quantum overlapping tomography is to output density matrices σ_S for all $S \subseteq \{1, 2, ..., n\}$ with $|S| \le k$ such that

$$||\rho_S - \sigma_S||_1 < \epsilon \tag{2}$$

for a given $\epsilon > 0$, where ρ_S denotes the reduced density matrix of $\rho_{1,2,\dots,n}$ on S.

If one only cares about M density matrices $\sigma_{S_1}, \sigma_{S_2}, \cdots, \sigma_{S_M}$ with $|S_i| \leq k$, this problem is called partial quantum overlapping tomography.

How many copies of ρ are needed to achieve this goal, with probability at least $1 - \delta$ for $\delta > 0$?

We show that,

Theorem 2. The sample complexity of quantum overlapping tomography of n-qubit system is $\Theta(\epsilon^{-2} \cdot \log(n/\delta))$ for constant k to succeed with probability at least $1 - \delta$, even using general joint measurement schemes. General quantum overlapping tomography problem for $1 \le k \le n$ can be accomplished by performing Pauli measurements on $\mathcal{O}(\epsilon^{-2} \cdot 10^k \cdot \log(\binom{n}{k}/\delta))$ copies. Moreover, for partial quantum overlapping tomography with M outcomes, the sample complexity is $\mathcal{O}(\epsilon^{-2} \cdot 10^k \cdot \log(2M/\delta))$ using Pauli measurements.

For the lower bound part, we show that to succeed with probability at least $1-\delta$, it is necessary to have $C_{\ell} \frac{\log(n/\delta)}{\epsilon^2}$ copies even if joint measurement on many copies of ρ are allowed, where $C_{\ell} > 0$ is a constant. For example, if m copies of the *n*-qubit system are prepared and measured. In the joint measure setting, the mn qubits may be accessed collectively.

The upper bound can be achieved by the following algorithm,

Algorithm 1: Quantum Overlapping Tomography
1 Repeat the following measurement $\mathcal{O}(g(k) \cdot \epsilon^{-2} \cdot \log(\binom{n}{k}/\delta))$ times;
2 Measuring each qubit using some informationally complete measurement for any copy of ρ ;

Here g(k) > 0 is a function depends on the informationally complete measurements and k only. Therefore, for constant k and fixed informationally complete measurements, the used number of copies becomes $\mathcal{O}(\epsilon^{-2} \cdot \log(n/\delta))$.

As an example, we choose informationally complete measurement $\mathcal{M} = \frac{1}{6} \{ \sigma_I + \sigma_X, \sigma_I - \sigma_X, \sigma_I + \sigma_Y, \sigma_I - \sigma_Y, \sigma_I + \sigma_Z, \sigma_I - \sigma_Z \}$, which can be regarded as random Pauli measurement, on each qubit. Then we can obtain the measurement scheme using $\mathcal{O}(\epsilon^{-2} \cdot 10^k \cdot \log(M/\delta))$ copies of ρ .

This is the first nontrivial example that Pauli measurements are as powerful as general joint measurements asymptotically. For this example, the number of unknown parameters of the output is exponentially larger than the number of copies.

This implies that for all observables $O_S \otimes I_{\bar{S}}$ with S being a set consisting of at most k-qubit, \bar{S} being the complementary set of S and $-I \leq O_S \leq I$, we can output estimation o_S such that

$$|\operatorname{tr}[\rho(O_S \otimes I_{\bar{S}})] - o_S| \le \epsilon$$

PRELIMINARIES

A positive-operator valued measure (POVM) on a finite dimensional Hilbert space \mathcal{H} is a set of positive semi-definite matrices $\mathcal{M} = \{M_i\}$ such that

$$\sum M_i = I_{\mathcal{H}}$$

We need the concept of the informationally complete POVM as follows,

Definition 3. An informationally complete POVM is a POVM whose outcome probabilities are sufficient to determine any state.

Equaivalently, a POVM $\mathcal{M} = \{M_i\}$ on *d*-dimensional \mathcal{H} is informationally complete if the linear span of $\{M_i\}$ equals to the whole $d \times d$ matrix space. In qubit system, it means $\sigma_I, \sigma_X, \sigma_Y$ and σ_Z all live in the linear span of $\{M_i\}$, where $\sigma_I, \sigma_X, \sigma_Y$ and σ_Z are Pauli matrices,

$$\sigma_I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \sigma_X = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \sigma_Z = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \sigma_Y = \begin{bmatrix} 0 & i \\ -i & 0 \end{bmatrix}.$$

It is direct to observe that $\mathcal{M} = \frac{1}{6} \{ \sigma_I + \sigma_X, \sigma_I - \sigma_X, \sigma_I + \sigma_Y, \sigma_I - \sigma_Y, \sigma_I + \sigma_Z, \sigma_I - \sigma_Z \}$ is an informationally complete measurement.

Observation 1. Given sufficient measurement outcomes of an informationally complete POVM, one can determine the state with high accuracy and confidence.

Observation 2. For information complete POVMs, \mathcal{M}_1 on \mathcal{H}_1 and \mathcal{M}_2 on \mathcal{H}_2 , $\mathcal{M}_1 \otimes \mathcal{M}_2$ is an informationally complete POVM on $\mathcal{H}_1 \otimes \mathcal{H}_2$.

Directly, $\mathcal{M}^{\otimes n} = \mathcal{M} = \frac{1}{6^n} \{ \sigma_I + \sigma_X, \sigma_I - \sigma_X, \sigma_I + \sigma_Y, \sigma_I - \sigma_Y, \sigma_I + \sigma_Z, \sigma_I - \sigma_Z \}^{\otimes n}$ is an informationally complete POVM of *n*-qubit system.

Definition 4. Let X_1, X_2, \dots, X_n be *n* samples of a distribution on $\{1, 2, \dots, n\}$. Then the empirical distribution is defined as

$$p(i) = \frac{\text{number of occurrences of } i}{n}$$

The following McDiarmids inequality [14] will be used in this paper.

Lemma 5. Consider independent random variables X_1, X_2, \ldots, X_n on probability space (Ω, \mathcal{F}, P) where $X_i \in \mathcal{X}_i$ for all *i* and a mapping $f : \mathcal{X}_1 \times \mathcal{X}_2 \times \cdots \times \mathcal{X}_n \to \mathbb{R}$. Assume there exist constant c_1, c_2, \ldots, c_n such that for all *i*,

$$\sup_{x_1,\cdots,x_{i-1},x_i,x'_i,x_{i+1},\cdots,x_n} |f(x_1,\dots,x_{i-1},x_i,x_{i+1},\cdots,x_n) - f(x_1,\dots,x_{i-1},x'_i,x_{i+1},\cdots,x_n)| \le c_i.$$
(3)

(In other words, changing the value of the *i*-th coordinate x_i changes the value of f by at most c_i .) Then, for any $\epsilon > 0$,

$$\Pr(f(X_1, X_2, \cdots, X_n) - \mathbb{E}[f(X_1, X_2, \cdots, X_n)] \ge \epsilon) \le \exp\left(-\frac{2\epsilon^2}{\sum_{i=1}^n c_i^2}\right).$$
(4)

QUANTUM TOMOGRAPHY USING PAULI MEASUREMENT

We start from the following observation:

When we measure an element of Pauli group, for instance $\sigma_X \sigma_Y$, on a two-qubit state ρ , the outcome is a sample from a 4-dimensional probability distribution, says $(p_{00}, p_{01}, p_{10}, p_{11})$, such that

$$tr(\rho(\sigma_X \otimes \sigma_Y)) = p_{00} - p_{01} - p_{10} + p_{11}.$$

One can easily observe that

$$tr(\rho(\sigma_X \otimes \sigma_I)) = p_{00} + p_{01} - p_{10} - p_{11}, tr(\rho(\sigma_I \otimes \sigma_Y)) = p_{00} - p_{01} + p_{10} - p_{11}, tr(\rho(\sigma_I \otimes \sigma_I)) = p_{00} + p_{01} + p_{10} + p_{11}.$$

In other words, by measuring XY, we actually obtained a sample of $\sigma_X \sigma_I$, a sample of $\sigma_Y \sigma_I$, and a sample of $\sigma_I \sigma_I$. This is also true for general *n*-qubit system as the following observation shows,

Observation 3. For any $P = P_1 \otimes P_2 \otimes \cdots \otimes P_n \in {\sigma_X, \sigma_Y, \sigma_Z}^{\otimes n}$, the measurement result of performing measurement P_i on the *i*-th qubit is an *n*-bit string *s*. One can interpret s the measurement outcome of performing $Q_i \in \{\sigma_I, \sigma_X, \sigma_Y, \sigma_Z\}$ on the *i*-th qubit if $Q_i = P_i$ or $Q_i = \sigma_I$. We call these $Q = Q_1 \otimes Q_2 \otimes \cdots \otimes Q_n$ corresponds to P.

Our measurement scheme is as follows: For any $\epsilon > 0$, fix an integer $m = 16 \cdot \frac{10^n \log \frac{1}{\delta}}{3^n \cdot \epsilon^2}$.

For any $P \in \{\sigma_X, \sigma_Y, \sigma_Z\}^{\otimes n}$, one performs m times P on ρ , and records the m samples of the 2^n dimensional outcome distribution.

According to the key observation, this measurement scheme provides $m \cdot 3^{n-w}$ samples of the expectation $tr(\rho P)$, says $\frac{\mu_P}{m \cdot 3^{n-w}}$ for each Pauli operator $P \in \{\sigma_I, \sigma_X, \sigma_Y, \sigma_Z\}^{\otimes n}$ with weight w, where $-m \cdot 3^{n-w} \leq \mu_P \leq m \cdot 3^{n-w}$. Output

$$\sigma = \sum_{P} \frac{\mu_P}{m \cdot 3^{n-w} \cdot 2^n} P.$$

Using this scheme, we obtained $m \cdot 3^n$ independent samples,

$$X_1, X_2, \cdots, X_{m \cdot 3^r}$$

where each $0 \leq X_i \leq 2^n - 1$.

Fruthermore, X_1, X_2, \dots, X_m corresponds to the measurement $\sigma_X^{\otimes n}$; $X_{m+1}, X_{m+2}, \dots, X_{2m}$ corresponds to the measurement $\sigma_X^{\otimes n-1} \otimes \sigma_Y$; \dots

It is direct to observe that, for any P of weight w, $\mu_P = \sum_{j=0}^{m \cdot 3^{n-w}-1} Z_j$, where Z_j are independent samples from distribution Z

$$\Pr(Z = 1) = \frac{1 + tr(\rho P)}{2}$$
$$\Pr(Z = -1) = \frac{1 - tr(\rho P)}{2}.$$

Furthermore, Z_j can be obtained from samples

$$X_1, X_2, \cdots, X_{m \cdot 3^n}.$$

Therefore,

$$\sigma = \sum_{P} \frac{\mu_P}{m \cdot 3^{n - w_P} \cdot 2^n} P.$$

is defined according to samples

$$X_1, X_2, \cdots, X_{m \cdot 3^n}.$$

It is direct to verify that

$$\mathbb{E}\sigma = \rho$$

where the expectation is taken over the probabilistic distribution according to the measurements. For any

$$\rho = \sum_{P} \frac{\alpha_P}{2^n} P$$

we can define the function $f: X_1 \times X_2 \times \cdots \times X_{m \cdot 3^n} \mapsto \mathbb{R}$

$$f = ||\rho - \sigma||_2$$

According to Cauchy inequality, we have

$$\begin{split} \mathbb{E}f \\ \leq &\sqrt{\mathbb{E}f^2} \\ = &\sqrt{\mathbb{E}(\operatorname{tr}\rho^2 - 2\operatorname{tr}\rho\sigma + \operatorname{tr}\sigma^2)} \\ = &\sqrt{\mathbb{E}(\operatorname{tr}\sigma^2 - \operatorname{tr}\rho^2)} \\ = &\sqrt{\frac{1}{2^n}\sum_P (\mathbb{E}\frac{\mu_P^2}{m^2 \cdot 9^{n-w_P}} - \alpha_P^2)} \\ = &\sqrt{\frac{1}{m \cdot 2^n} \cdot \sum_P \frac{1 - \alpha_P^2}{3^{n-w_P}}} \\ < &\sqrt{\frac{1}{m \cdot 2^n} \cdot \sum_P \frac{1}{3^{n-w_P}}} \\ = &\sqrt{\frac{1}{m \cdot 2^n} \cdot \sum_{w_P=0}^n \frac{1}{3^{n-w_P}} \binom{n}{w_P}} 3^{w_P} \\ = &\sqrt{\frac{1}{m \cdot 6^n} \cdot \sum_{w_P=0}^n (1+9)^n} \\ = &\sqrt{\frac{5^n}{m \cdot 3^n}} \\ < &\frac{\epsilon}{2 \cdot \sqrt{2^n}}. \end{split}$$

For any sample X_i corresponding to $P \in \{\sigma_X, \sigma_Y, \sigma_Z\}^{\otimes n}$. If only X_i is changed, it would only change μ_Q for those $Q \in \{\sigma_I, \sigma_X, \sigma_Y, \sigma_Z\}^{\otimes n}$ where Q is obtained by replacing some $\{\sigma_X, \sigma_Y, \sigma_Z\}$ s of P by σ_I . Moreover, those μ_Q would change at most 2. According to triangle inequality, f would change at most

$$\begin{split} ||\sum_{Q} \frac{\Delta \mu_Q}{m \cdot 3^{n-w_Q} \cdot 2^n} Q||_2 \\ = & \sqrt{\sum_{Q} \frac{\Delta \mu_Q^2}{m^2 \cdot 9^{n-w_Q} \cdot 2^n}} \\ \leq & \sqrt{\sum_{Q} \frac{2^2}{m^2 \cdot 9^{n-w_Q} \cdot 2^n}} \\ = & \sqrt{\sum_{w_Q=0}^n \frac{2^2}{m^2 \cdot 9^{n-w_Q} \cdot 2^n}} \begin{pmatrix} n \\ w_Q \end{pmatrix}} \\ = & \frac{2 \cdot \sqrt{5}^n}{m \cdot 3^n}, \end{split}$$

where Q ranges over all Paulis which corresponds to P, and $\delta \mu_Q$ denotes the difference of μ_Q when X_i is changed.

For $\epsilon > 0$, let $\epsilon' = \frac{\epsilon}{\sqrt{2^n}}$. We have

$$\begin{aligned} &\Pr(||\rho - \sigma||_1 > \epsilon) \\ \leq &\Pr(||\rho - \sigma||_2 > \epsilon') \\ = &\Pr(f > \epsilon') \\ \leq &\Pr(f - \mathbb{E}f > \frac{\epsilon'}{2}) \\ < &\exp(-\frac{\epsilon'^2}{4 \cdot \frac{4\cdot 5^n}{m^2 \cdot 9^n} \cdot m \cdot 3^n}) \\ = &\exp(-\frac{m \cdot 3^n \cdot \epsilon'^2}{16 \cdot 5^n}) \\ < &\delta, \end{aligned}$$

where the first step is by the relation between the trace norm and 2 norm; the third step is by $\mathbb{E}f \leq \frac{\epsilon'}{2}$; the fourth step is by Lemma 5 (McDiarmids inequality).

The total number of used copies is

$$m \cdot 3^n = 16 \cdot \frac{10^n \log \frac{1}{\delta}}{\epsilon^2}.$$

QUANTUM OVERLAPPING TOMOGRAPHY

In this section, we study the quantum overlapping tomography.

We first analyze Algorithm 1.

Fixing *n* informationally complete POVM of one-qubit system, says $\mathcal{M}_1, \mathcal{M}_2, \dots, \mathcal{M}_n$. For any given unknown *n*-qubit state ρ , we perform $\mathcal{M}_1 \otimes \mathcal{M}_2 \otimes \dots \otimes \mathcal{M}_n$ on ρ and obtain a string $s_1 s_2 \cdots s_n$ with s_i denoting the measurement outcome of \mathcal{M}_i . Assume the output probability distribution is $p_{1,2,\dots,n}$. we know that $s_1 s_2 \cdots s_n$ obeys the distribution of $p_{1,2,\dots,n}$.

The key observation is that measuring each qubit independently preserves the local structure. More precisely,

Observation 4. For any $S = \{i_1, i_2, \dots, i_r\} \subseteq \{1, 2, \dots, n\}$, the distribution of $s_{i_1}s_{i_2}\cdots s_{i_r}$ obeys the outcome distribution of performing $\mathcal{M}_{i_1} \otimes \mathcal{M}_{i_2} \otimes \cdots \otimes \mathcal{M}_{i_r}$ on ρ_S . Moreover, $s_{i_1}s_{i_2}\cdots s_{i_r}$ obeys the marginal distribution p_S .

By Algorithm 1, we repeat the measurement $\mathcal{M}_1 \otimes \mathcal{M}_2 \otimes \cdots \otimes \mathcal{M}_n$ m times to obtain m samples of $p_{1,2,\dots,n}$. For any sets $S_1, S_2, \dots, S_M \subseteq \{1, 2, \dots, n\}$, Obervation 3 implies that we have m samples of the marginal distributions $p_{S_1}, p_{S_2} \cdots$, and p_{S_M} , not independent. Although, m samples maybe not enough to recover $p_{1,2,\dots,n}$, but enough to recover p_{S_i} for each S_i with high accuracy and high confidence using Chernoff bound. According to Observation 1 and Observation 2, this implies one can recover ρ_{S_i} with high accuracy, trace distance error ϵ , and successful probability at least $1 - \delta'$ for each S_i using $m = \mathcal{O}(g(k) \cdot \frac{\log \delta'}{\epsilon^2})$.

To successfully recover ρ_{S_i} simultaneously, we only need the recovery of each ρ_{S_i} with high accuracy and probability at least $1 - \frac{\delta}{\binom{n}{2}}$ according to union bound.

Therefore, we only need $m = \mathcal{O}(g(k) \cdot \frac{\log \binom{n}{k}}{\epsilon^2})$ copies.

Joint Measurement Lower bound

In this subsection, we show that $\Omega(\frac{\log(n/\delta)}{\epsilon^2})$ copies are necessary to quantum overlapping tomography by proving that: $\Omega(\frac{\log(n/\delta)}{\epsilon^2})$ copies are necessary for quantum overlapping tomography with k = 1, even if general *joint* measurements are used.

Because trace distance is non-increasing according to partial trace, we can conclude that any measurement scheme can solve the quantum overlapping tomography problem for $k \ge 1$, automatically, it solves the case that k = 1. Moreover, to deal with general joint measurement schemes, we focus on classical distributions.

We first consider a simple question: Given a binary random variable X obeys distribution $q_0 = (1/2 - \epsilon, 1/2 + \epsilon)$ or $q_1 = (1/2 + \epsilon, 1/2 - \epsilon)$. The goal is to find out which distribution is the true distribution. It is widely known that: For any fixed m as the number of tossing this coin, the best strategy is to toss the coin m times and declare the index (0 or 1) that appears less.

Let the X_1, X_2, \ldots, X_m be the *m* samples of q_1 and any $0 \le t \le 2m\epsilon$, Ref. [15] proved the following:

$$\Pr(\sum_{i=1}^{m} X_i > t + m(1/2 - \epsilon)) \ge \frac{1}{4} \cdot \exp(-\frac{2t^2}{m(1/2 - \epsilon)})$$
(5)

By choosing $t = m\epsilon$, $\Pr(\sum_{i=1}^{m} X_i > t + m(1/2 - \epsilon)) = \Pr(\sum_{i=1}^{m} X_i > m/2)$ is the probability of answering q_0 , a lower bound of the failure probability.

To success with probability at least $1 - \delta'$, we must have

$$\delta' \ge \frac{1}{4} \cdot \exp(-\frac{2m\epsilon^2}{1/2 - \epsilon}) \tag{6}$$

That is, $\frac{1-2\epsilon}{4\epsilon^2}\log(\frac{1}{4\delta'})$ samples are needed to distinguish q_0 and q_1 with confidence at least $1-\delta'$.

Back to our problem of showing $\Omega(\frac{\log(n/\delta)}{\epsilon^2})$ samples of *n*-qubit state ρ are necessary to solve the quantum overlapping tomography for $k \geq 1$, to within additive error ϵ and confidence at least $1 - \delta$, we consider the classical distributions $p_{i_1,i_2,\dots,i_n} = q_{i_1} \otimes q_{i_2} \otimes \dots \otimes q_{i_n}$ where $q_0 = (1/2 - \epsilon, 1/2 + \epsilon)$ and $q_1 = (1/2 + \epsilon, 1/2 - \epsilon)$. In total, there are 2^n different distributions.

Suppose there is a quantum procedure \mathcal{A} which uses *m* copies of ρ to accomplish the quantum overlapping tomography with probability at least $1 - \delta$.

Let Z_1, Z_2, \dots, Z_n be random variables which obey uniform binary distribution. Choose each p_{Z_1, Z_2, \dots, Z_n} with probability $1/2^n$, and apply \mathcal{A} on m copies of p_{Z_1, Z_2, \dots, Z_n} . Because the ℓ_1 norm is non-increasing under partial trace, we know that according to the output of \mathcal{A} , we can successfully recover the indices Z_1, Z_2, \dots, Z_n with probability at least $1 - \delta$.

In the following, we first observe that quantum procedure does not help in recovering Z_1, Z_2, \dots, Z_n from samples of p_{Z_1, Z_2, \dots, Z_n} . We assume the joint measurement $(M_{0,0,\dots,0}, \dots, M_{1,1,\dots,1})$ applied on m copies (samples) of p such that the measurement outcome M_{i_1,i_2,\dots,i_n} allows us to answer $Z_1 = i_1, Z_2 = i_2, \dots, Z_n = i_n$. Here M_{i_1,i_2,\dots,i_n} s are $2^{mn} \times 2^{mn}$ matrices.

We first observe that p_{Z_1,Z_2,\dots,Z_n} s are all diagonal, so are $p_{Z_1,Z_2,\dots,Z_n}^{\otimes m}$. Then, the off diagonal elements of M_{i_1,i_2,\dots,i_n} has no effect for this task. Therefore, we only need to consider the procedure in the following two steps: At the first step, measure m copies of p_{Z_1,Z_2,\dots,Z_n} s in the diagonal basis; at the second step, output according to some probability distributions.

The first step ensures that we only need to measure each copy of p_{Z_1,Z_2,\dots,Z_n} s in the diagonal basis, as there is no difference. By the convexity of the successful probability, we know that deterministic function works best in the second step, that is declare the index (0 or 1) that appears less for each $1 \leq j \leq n$.

Now, we assume the output random virable is Y_1, Y_2, \dots, Y_n , our goal is

$$\Pr(Y_1 = Z_1, Y_2 = Z_2, \cdots, Y_n = Z_n) \ge 1 - \delta.$$
(7)

By Bayes' theorem, we know that

$$Pr(Y_1 = Z_1, Y_2 = Z_2, \cdots, Y_n = Z_n)$$

=Pr(Y_1 = Z_1) × Pr(Y_2 = Z_2 | Y_1 = Z_1) × \cdots × Pr(Y_n = Z_n | Y_1 = Z_1, \cdots, Y_{n-1} = Z_{n-1})
 $\leq (1 - \delta') \cdot (1 - \delta') \cdots (1 - \delta')$
= $(1 - \delta')^m$,

where we use the fact that p_{Z_1,Z_2,\dots,Z_n} s are all in tensor product form, therefore, $\Pr(Y_2 = Z_2 | Y_1 = Z_1) = \Pr(Y_2 = Z_2)$, and so on. $(1 - \delta')$ denotes the successful probability of discriminate q_0 and q_1 with m copies.

Therefore, we require that

$$(1-\delta')^n > 1-\delta.$$

That is $\delta' = \Theta(\frac{\delta}{n})$, this implies the bound $m = \Omega(\frac{\log(n/\delta)}{\epsilon^2})$.

Pauli Measurement Upper bound

In this subsection, we analyze Algorithm 1 using the measurement scheme $\{M_0, M_1, M_2, M_3, M_4, M_5\}^{\otimes n}$ as a special case, where

$$M_0 = \frac{\sigma_I + \sigma_X}{6}, \qquad M_1 = \frac{\sigma_I - \sigma_X}{6},$$
$$M_2 = \frac{\sigma_I + \sigma_Y}{6}, \qquad M_3 = \frac{\sigma_I - \sigma_Y}{6},$$
$$M_4 = \frac{\sigma_I + \sigma_Z}{6}, \qquad M_5 = \frac{\sigma_I - \sigma_Z}{6}.$$

This can be regarded as a random chosen of $P \in \{\sigma_X, \sigma_Y, \sigma_Z\}^{\otimes}$ and measure ρ in the basis of P, and repeat it m times. That is,

Algorithm 2: Quantum Overlapping Tomography by Pauli Measurements	
1Repeat the following measurement $32 \cdot 10^k \cdot \epsilon^{-2} \cdot \log(2\binom{n}{k}/\delta)$ times; 2For $i = 1$ to n : measure the <i>i</i> -th qubit in a random chosen basis from $\{\sigma_X, \sigma_Y, \sigma_Z\}$;	

We observe the following,

Observation 5. For any $S = \{i_1, i_2, \dots, i_k\} \subseteq \{1, 2, \dots, n\}$, with probability at least $1 - \frac{2^m}{e^m} \cdot 3^k$, each $P \in \{\sigma_X, \sigma_Y, \sigma_Z\}^{\otimes k}$ was measured for ρ_S at least $m = 16 \cdot \frac{10^k \cdot \log(2\binom{n}{k})/\delta}{3^k \cdot \epsilon^2}$ times.

Proof. It is direct to observe that for any $P \in {\{\sigma_X, \sigma_Y, \sigma_Z\}}^{\otimes k}$, it was chosen as a basis to measure is no more than m times with probability at most

$$\sum_{i=0}^{m} \binom{2m \cdot 3^{k}}{i} (1 - \frac{1}{3^{k}})^{2m \cdot 3^{k} - i} \frac{1}{3^{ki}}$$
$$= (1 - \frac{1}{3^{k}})^{2m \cdot 3^{k}} \cdot \sum_{i=0}^{m} \binom{2m \cdot 3^{k}}{i} (1 - \frac{1}{3^{k}})^{-i} \frac{1}{3^{ki}}$$
$$= (1 - \frac{1}{3^{k}})^{2m \cdot 3^{k}} \cdot \sum_{i=0}^{m} \binom{2m \cdot 3^{k}}{i} \frac{1}{(3^{k} - 1)^{i}}$$

The ratio of the i + 1-th term and the *i*-th term is

$$\frac{1}{3^k} \cdot \frac{2m \cdot 3^k - i}{i+1} \ge 2$$

Therefore, the serie grows faster than a geometric progression with common ratio 2. Then,

$$\begin{aligned} &(1-\frac{1}{3^{k}})^{2m\cdot 3^{k}}\cdot \sum_{i=0}^{m}\binom{2m\cdot 3^{k}}{i}\frac{1}{(3^{k}-1)^{i}}\\ &\leq 2\cdot (1-\frac{1}{3^{k}})^{2m\cdot 3^{k}}\binom{2m\cdot 3^{k}}{m}\frac{1}{(3^{k}-1)^{m}}\\ &\leq 3\sqrt{\frac{2\cdot 3^{k}}{2\pi\cdot m\cdot (2\cdot 3^{k}-1)}}[2(1-\frac{1}{2\cdot 3^{k}-1})^{2\cdot 3^{k}-1}]^{m}\\ &\leq 3\sqrt{\frac{2\cdot 3^{k}}{2\pi\cdot m\cdot (2\cdot 3^{k}-1)}}\frac{2^{m}}{e^{m}}\\ &\leq \frac{2^{m}}{e^{m}}.\end{aligned}$$

where we use the bound that

$$\sqrt{2\pi n} \cdot n^n \cdot e^{-n} \cdot e^{\frac{1}{12n}} < n! < \sqrt{2\pi n} \cdot n^n \cdot e^{-n} \cdot e^{\frac{1}{12n+1}}$$

and the number series

$$(1-\frac{1}{d})^d$$

is increasing and goes to $\frac{1}{e}$.

Then, by union bound, we know that for any S of size k, with probability at least $1 - \frac{2^m}{e^m} \cdot 3^k$, each $P \in \{\sigma_X, \sigma_Y, \sigma_Z\}^{\otimes k}$ was measured for ρ_S at least m times.

According to Theorem 1, the tomography of ρ_S with trace distance error ϵ was successful with probability at least

$$1 - [3^k \cdot \frac{2^m}{e^m} + \frac{\delta}{2\binom{n}{k}}] > 1 - \frac{\delta}{\binom{n}{k}},$$

the last inequality follows from

$$3^{k} \cdot \frac{2^{m}}{e^{m}}$$

$$<3^{k} \left(\frac{2}{e}\right)^{12 \cdot 3^{k}} \cdot \left(\frac{2}{e}\right)^{4 \cdot 3^{k} \cdot \log\left(2\binom{n}{k}\right)/\delta}$$

$$<3^{k} \left(\frac{2}{e}\right)^{12 \cdot k} \cdot \left(\frac{1}{e}\right)^{\log\left(2\binom{n}{k}\right)/\delta}$$

$$<\frac{\delta}{2\binom{n}{k}}.$$

By union bound, the quantum overlapping tomography was with trace distance error ϵ was successful with probability at least

$$1 - \binom{n}{k} \cdot \frac{\delta}{\binom{n}{k}} = 1 - \delta$$

DISCUSSION

For the general quantum state tomography problem, our measurement scheme is much more efficient than previous schemes in the Pauli measurements setting. Our result raises an important question on the performance of *local* measurements, in particular Pauli measurements, in which each qubit is measured at a time.

The sample complexity of the quantum overlapping tomography problem is nearly resolved here up to a constant factor.

An independent work [16] analyzes random Clifford measurement and random Pauli measurement and proves that it only requires $\mathcal{O}(\frac{\log(M) \cdot s}{\epsilon^2})$ copies to achieve ϵ accuracy estimation for $\operatorname{tr}(O_1\rho), \cdots, \operatorname{tr}(O_M\rho)$, a more general question, where s depends on the structure of O_i s. They also show that this is tight if only single-copy measurements are allowed. If O_i s are k-qubit Paulis, $\mathcal{O}(\frac{\log(M/\delta) \cdot 3^k}{\epsilon^2})$ which coinsides with the bound provided in Ref. [13]. These upper bound does not cover ours, because a 4^k coefficient would be added when we move from Pauli estimation to state tomography, thus, these results would lead to a $\mathcal{O}(\frac{\log(M/\delta) \cdot 12^k}{\epsilon^2})$ measurement scheme. Also, this lower bound does not imply our lower bound, for the specific quantum overlapping tomography problem, because this bound is not for joint measurement and does not consider the successful probability parameter.

ACKNOWLEDGMENTS

We thank Steve Flammia for the discussion on quantum overlapping tomography. We thank Richard Keung for clasifying their bound in [10]. This work was supported by DE180100156.

Jordan Cotler and Frank Wilczek, "Quantum overlapping tomography," Physical Review Letters 124 (2020), 10.1103/physrevlett.124.100401.

- [2] E. Bagan, M. Baig, R. Muñoz Tapia, and A. Rodriguez, "Collective versus local measurements in a qubit mixed-state estimation," Phys. Rev. A 69, 010304 (2004).
- [3] M. Keyl, "Quantum state estimation and large deviations," Reveiws in Mathematical Physics 18, 19–60 (2006).
- [4] Mădălin Guţă and Jonas Kahn, "Optimal estimation of qubit states with continuous time measurements," Communications in Mathematical Physics 277, 127–160 (2008), quant-ph/0608074.
- [5] J. Haah, A. W. Harrow, Z. Ji, X. Wu, and N. Yu, "Sample-optimal tomography of quantum states," in Proceedings of the Forty-Seventh Annual ACM on Symposium on Theory of Computing, STOC '16 (2016) pp. 913–925.
- [6] R. O'Donnell and J. Wright, "Efficient quantum tomography," in Proceedings of the Forty-Seventh Annual ACM on Symposium on Theory of Computing, STOC '16 (2016) pp. 899–912.
- [7] R. O'Donnell and J. Wright, "Efficient quantum tomography ii," in Proceedings of the Forty-Seventh Annual ACM on Symposium on Theory of Computing, STOC '17 (2017) pp. 962–974.
- [8] S. T. Flammia, D. Gross, Y. Liu, and J. Eisert, "Quantum tomography via compressed sensing: Error bounds, sample complexity, and efficient estimators," New J. Phys. 14, 095022 (2012).
- [9] R. Kueng, H. Rauhut, and U. Terstiege, "Low rank matrix recovery from rank one measurements," Applied and Computational Harmonic Analysis 42, 88–116 (2017).
- [10] M Guţă, J Kahn, R Kueng, and J A Tropp, "Fast state tomography with optimal error bounds," Journal of Physics A: Mathematical and Theoretical 53, 204001 (2020).
- [11] Steven T Flammia, David Gross, Yi-Kai Liu, and Jens Eisert, "Quantum tomography via compressed sensing: error bounds, sample complexity and efficient estimators," New Journal of Physics 14, 095022 (2012).
- [12] Guillermo Garca-Prez, Matteo A. C. Rossi, Boris Sokolov, Elsi-Mari Borrelli, and Sabrina Maniscalco, "Pairwise tomography networks for many-body quantum systems," Physical Review Research 2 (2020), 10.1103/physrevresearch.2.023393.
- [13] Tim J. Evans, Robin Harper, and Steven T. Flammia, "Scalable bayesian hamiltonian learning," (2019), arXiv:1912.07636 [quant-ph].
- [14] Colin McDiarmid, "On the method of bounded differences," in Surveys in Combinatorics, 1989: Invited Papers at the Twelfth British Combinatorial Conference, London Mathematical Society Lecture Note Series, edited by J.Editor Siemons (Cambridge University Press, 1989) p. 148188.
- [15] Nima Mousavi, "How tight is chernoff bound?" (2016), https://ece.uwaterloo.ca/ nmousavi/Papers/Chernoff-Tightness.pdf.
- [16] Hsin-Yuan Huang, Richard Kueng, and John Preskill, "Predicting many properties of a quantum system from very few measurements," Nature Physics (2020), 10.1038/s41567-020-0932-7.