

Unlocking of time reversal, space-time inversion and rotation invariants in magnetic materials

Jian Yang,¹ Zheng-Xin Liu,^{2,*} and Chen Fang^{1,†}

¹*Beijing National Laboratory for Condensed Matter Physics and Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China*

²*Department of Physics, Renmin University, Beijing, China*

(Dated: September 18, 2020)

Time reversal (T) and space inversion are symmetries of our universe in the low-energy limit. Fundamental theorems relate their corresponding quantum numbers to the spin for elementary particles: $\hat{T}^2 = (\hat{P}\hat{T})^2 = -1$ for half-odd-integer spins and $\hat{T}^2 = (\hat{P}\hat{T})^2 = +1$ for integer spins. Here we show that for elementary excitations in magnetic materials, this “locking” between quantum numbers is lifted: \hat{T}^2 and $(\hat{P}\hat{T})^2$ take all four combinations of $+1$ and -1 regardless of the value of the spin, where T now represents the composite symmetry of time reversal and lattice translation. Unlocked quantum numbers lead to new forms of minimal coupling between these excitations and external fields, enabling novel physical phenomena such as the “cross-Larmor precession”, indirectly observable in a proposed light-absorption experiment. We list the magnetic space groups with certain high-symmetry momenta where such excitations may be found.

I. INTRODUCTION

Spatial inversion (P) and time reversal (T) are considered as fundamental symmetries of our universe at low energy (but are broken in weak forces)[1]. As operators, inversion acts on the spatial degrees of freedom of particles, and time reversal the internal ones: they naturally commute $[\hat{P}, \hat{T}] = 0$. (We use hatted letters for operators in Hilbert space, and unhatted ones for symmetries themselves.) Restricting the discussion to the single-particle sector of the Hilbert space, we understand that inversion squares to unity, $\hat{P}^2 = 1$, because (i) two consecutive inversions equal identity and (ii) inversion does not act on spin. Time reversal, on the other hand, an anti-unitary symmetry reversing time and inverting spin, is represented by $\hat{T} = e^{i\hat{S}_y\pi}K$, where \hat{S}_y is the y -component of the spin operator. Therefore we have $\hat{T}^2 = (-1)^{2s}$, where $s(s+1)\hbar^2 = \hat{\mathbf{S}}^2$.

Symbolically, we have

$$[\hat{P}, \hat{T}] = 0, \hat{P}^2 = 1, \hat{T}^2 = (-1)^{2s}. \quad (1)$$

Eq.(1) relates the space-inversion, the time-reversal and the rotation invariants for particles in vacuum. These relations can be more concisely represented by three *invariants* $\chi_T := \hat{T}^2$, $\chi_{PT} := (\hat{P}\hat{T})^2$ and $\chi_S := (-1)^{2s}$:

$$\chi_T = \chi_{PT} = \chi_S. \quad (2)$$

For a given type of particle with spin- s , χ_T and χ_{PT} are completely fixed to be $+1$ if s is an integer (particle being boson) and -1 if s is a half-odd-integer (particle being fermion). Eq.(2) applies for any Lorentz-invariant theory

given that \hat{P} and \hat{T} do not act on the species degrees of freedom [2, 3].

In this paper, we show that for elementary excitations in certain lattices with magnetic ordering, the three invariants, χ_S (the definition of which is modified due to the absence of continuous rotation), χ_T and χ_{PT} , are independent from each other, and can take all eight combinations, in contrast to being locked with each other as in Eq.(2) in real vacuum. Here, what T represents is not time reversal, broken by the magnetic order, but a composite symmetry of time reversal and some lattice translation, which is present in many antiferromagnets. The properties of elementary excitations having unlocked invariants are demonstrated through the example of $\mathbf{R} = (\pi/a, \pi/a, \pi/a)$ in magnetic space group 222.103, where a is the lattice constant of the magnetic unit cell. The unlocked invariants enable unconventional linear responses to electromagnetic field. We show that polarization operators and magnetization operators together furnish an $SO(4)$ algebra unseen in previous studies to our best knowledge. The $SO(4)$ algebra leads to a new type of Larmor precession, where a precession of polarization is driven by a magnetic field, and a precession of magnetization by an electric field. We propose a light-absorption experiment to observe this “cross-Larmor precession”.

II. INVARIANTS ON A NON-MAGNETIC LATTICE

When we put any theory on a lattice, the three-dimensional-rotation symmetry reduces to a point group symmetry, so the rotation invariant χ_S is no longer well-defined. For a modified version, we require the point group in question to contain at least three orthogonal twofold rotation axes. For example, the group D_{2h} having three such twofold axes $C_{2x,2y,2z}$, satisfies our constraint, while the group C_{2h} , having only one, does not.

*Electronic address: liuzxphys@ruc.edu.cn

†Electronic address: cfang@iphy.ac.cn




FIG. 1: Electron excitations centering at BZ corners. (a) In SnTe, the conduction bottom (and valence top) is located at L instead of Γ . (b) In BaPpT, the conduction and the valence bands touch each other at R. In both cases, there are bands of elementary excitations centering at some nonzero and high-symmetry momentum. These band structures are downloaded from <http://materiae.iphy.ac.cn>

The redefined rotation invariant χ_S is

$$\hat{C}_{2m}\hat{C}_{2n} = \chi_S\hat{C}_{2n}\hat{C}_{2m}, \quad (3)$$

where $m \neq n$ take values x, y, z .

In real vacuum, for any stable theory, the lowest particle excitations are near zero momentum, because this is the only special point having higher symmetry than its neighborhood. However, on a lattice, the elementary excitation may also appear at the corners of the Brillouin zone (BZ), which are nonzero crystal momenta with highest symmetry in the neighborhood[4]. This is the case when, for example, the band bottom of the conduction band appears at a BZ corner [Fig.1(a)], or when the conduction and the valence bands touch each other at some BZ corner [Fig.1(b)].

Yet another complication in lattices is that some spatial symmetries are neither point-group operations nor lattice translations, but are composite operations of a point group symmetry and a fractional lattice translation. Lattice symmetry groups containing these spatial symmetries are called nonsymmorphic space groups[5]. Consider for example space group $Pbca$, generated by three orthogonal lattice translations $t_{\mathbf{x},\mathbf{y},\mathbf{z}}$ along x, y, z -directions with lattice constants a, b, c , respectively, space inversion about the origin, P , and three twofold screw axes parallel to x, y, z -axes, $C_{2x} \rightarrow 2_{100}t_{\mathbf{x}/2+\mathbf{y}/2}$, $C_{2y} \rightarrow 2_{010}t_{\mathbf{y}/2+\mathbf{z}/2}$, $C_{2z} \rightarrow 2_{001}t_{\mathbf{x}/2+\mathbf{z}/2}$; 2_{mnl} represents a π -rotation on both spatial coordinates and spin components about the $[mnl]$ -direction, where $[mnl]$ are Miller indices. $t_{d_1\mathbf{x}+d_2\mathbf{y}+d_3\mathbf{z}}$ represents translation by $d_1\mathbf{x}+d_2\mathbf{y}+d_3\mathbf{z}$ where $\mathbf{x}, \mathbf{y}, \mathbf{z}$ are unit vectors and a, b, c lattice constants. It is straightforward to check that $\hat{C}_{2x}\hat{C}_{2y} = (-1)^{2s_0}\hat{C}_{2y}\hat{C}_{2x}\hat{t}_{-\mathbf{x}+\mathbf{y}+\mathbf{z}}$. Here s_0 is the physical spin of the particle, $s_0 \in \text{even}$ for bosons and $s_0 \in \text{odd}$ for fermions. Consider excitations at $\mathbf{R} = (\pi/a, \pi/b, \pi/c)$, where $\hat{t}_{-\mathbf{x}+\mathbf{y}+\mathbf{z}} = -1$, then we have $\chi_S = -(-1)^{2s_0}$. The additional minus sign means that the half-lattice translations give a ‘‘twist’’ to the rotation invariants, making integer spins look like half-odd-integer spins, and vice versa[6].

However, on a nonmagnetic, centrosymmetric lattice, similar twist does not occur in the invariants (χ_T, χ_{PT}). For excitations on any nonmagnetic lattice, we always have $\chi_T = \chi_{PT} = (-1)^{2s_0}$. Therefore, for excitations at $\mathbf{R} = (\pi/a, \pi/b, \pi/c)$ in space group $Pbca$, we have

$$\chi_T = \chi_{PT} = -\chi_S. \quad (4)$$

III. FURTHER UNLOCKING OF INVARIANTS ON A MAGNETIC LATTICE

Is it possible to further unlock the values of χ_T and χ_{PT} from the constraint $\chi_T = \chi_{PT}$? We seek this possibility in elementary excitations in magnetically ordered lattices, the band topology of which has recently become a research focus[7–14]. While the physical time reversal is broken by magnetism, many antiferromagnetic materials preserve a symmetry in the form $T \rightarrow T_{phy}t_{\mathbf{L}/2}$, where T_{phy} is pure time reversal operator acting only on internal degrees of freedom, and $\mathbf{L}/2$ is a lattice vector (which becomes a half-lattice vector in the magnetic unit cell)[7, 8]. Symmetries of magnetic systems are classified by magnetic space groups, and when the above T is a symmetry, the magnetic space groups are called type-IV[15], which are of our interest. When this is the case, excitations at any time-reversal invariant momentum in the magnetic Brillouin zone (mBZ) have this T symmetry. We then naturally redefine χ_T as $\chi_T \equiv \hat{T}^2 = (\hat{T}_{phy}\hat{t}_{\mathbf{L}/2})^2$. As expected, the translation part in T in general changes the value of χ_T , and could also make P and T not commute[7, 12]. One can use the theory of projective representation(Rep)s to show that (χ_T, χ_{PT}) can take all possible values of $(+1, +1)$, $(+1, -1)$, $(-1, +1)$ and $(-1, -1)$, irrespective of the rotation invariant χ_S . Let us use an example to illustrate one of these possibilities here, and defer the general proof in Table IV of appendix.

In Fig.2, a conjectured magnetic structure of NdZn[13, 16] (magnetic space group 222.103) is shown. Looking from O or O' , the magnetic structure is invariant under all proper rotations of a cube (point group O); the structure is also centro-symmetric about I , and has a composite symmetry $T \rightarrow T_{phy}t_{(\mathbf{x}+\mathbf{y}+\mathbf{z})/2}$, where $(\mathbf{x}+\mathbf{y}+\mathbf{z})/2$ is a lattice vector of the nonmagnetic lattice. It is straightforward to check $\hat{T}^2 = (-1)^{2s_0}\hat{t}_{\mathbf{x}+\mathbf{y}+\mathbf{z}}$, $\hat{P}\hat{T} = \hat{t}_{\mathbf{x}+\mathbf{y}+\mathbf{z}}\hat{T}\hat{P}$, and $\chi_S = (-1)^{2s_0}$. Therefore, at mBZ corner $\mathbf{R} = (\pi/a, \pi/a, \pi/a)$, we have

$$\chi_T = -\chi_{PT} = -\chi_S. \quad (5)$$

For magnons on this lattice, we have $(\chi_S, \chi_T, \chi_{PT}) = (+1, -1, +1)$, and for electrons $(-1, +1, -1)$, neither of which is possible on nonmagnetic lattices.




FIG. 2: The magnetic unit cell with lattice constant a of a magnetic structure having magnetic space group 222.103, conjectured as the symmetry for the magnetic ground state of NdZn. Yellow circles O, O' are body centers of point group symmetry O , and orange circle I is an inversion center. T is time reversal combined with a lattice translation that relates the two cubes outlined in red. The local moments are polarized so that they are all-out with respect to O and all-in with respect to O' .

IV. EFFECTIVE THEORY WITH MINIMAL COUPLING

In quantum field theory, particles couple to an external source (field) if there exists a bilinear of particle field operators that carries the same symmetry Rep as the one carried by the source. The form of coupling hence depends on the projective Rep carried by particle field, as well as the linear Rep carried by the external field. The symmetry invariants $\chi_{S,T,PT}$ do not denote specific Reps, but classes of Reps. For example, for group $SO(3)$, all integer spins belong to one class ($\chi_S = 1$) while all half-odd-integer spins to another class ($\chi_S = -1$).

We postpone the former definition of “classes of Reps” and invariants to a later part. For now, we only need that fact that new invariants unlocked in magnetic systems allow new classes of Reps for field operators, which in turn bring about new forms of linear coupling to external sources such as applied electromagnetic field and strain tensor field. In this section we restrict the discussion to electron excitations, $s_0 = 1/2$, linearly coupled to an electromagnetic field. Physically, since electrons have charge, the electromagnetic field has two effects: the orbital effect that changes the crystal momentum of the electron in a fixed band structure, and the Zeeman effect which modifies the periodic part of Bloch wavefunctions at each momentum, thereby modifying the band struc-

ture itself[17]. The invariants, $\chi_{S,T,PT}$, have little to do with the first effect, so we focus on the second, altogether ignoring the change of crystal momentum by the external fields.

Again let us use the magnetic lattice in Fig.2 as the example, and focus on mBZ corner $\mathbf{R} = (\pi/a, \pi/a, \pi/a)$ having little co-group $O_h \times Z_2^T$. There are three different irreducible Reps, $R_{1/2}$ and $R_{3/2}^\pm$, of $G_{\mathbf{R}}$, all satisfying $(\chi_S, \chi_T, \chi_{PT}) = (-1, +1, -1)$. Here we pick one Rep $R_{1/2}$ for our analysis, where the generators are given by

$$\begin{aligned} \hat{T} &= \tau_y \sigma_y K, \\ \hat{P} &= i\tau_y, \\ \hat{C}_{2x} &= i\sigma_x, \\ \hat{C}_{31} &\equiv \hat{C}_{3,111} = -\exp(-i\sigma_{111}\pi/3), \\ \hat{C}_{4z} &= i\tau_z \exp(-i\sigma_z\pi/4), \end{aligned} \quad (6)$$

where $C_{3,111}$ is the threefold rotation about the $[111]$ -direction, and $\sigma_{111} \equiv (\sigma_x + \sigma_y + \sigma_z)/\sqrt{3}$. Group theory allows us to classify all hermitian bilinears of field operators in the form $\psi^\dagger \tau_\mu \sigma_\nu \psi$ into the Reps of the symmetry group as in Table I. On the other hand, all monomials of crystal momentum relative to \mathbf{R} , $\mathbf{q} = \mathbf{k} - \mathbf{R}$, and the components of electric, magnetic and strain fields can also be put into these Reps, also summarized in Table I.

Matching the Reps between fermion bilinear operators and monomials of momenta and of fields, we find the effective theory with minimal couplings. The free part of the theory is obtained by matching the monomials of momenta to the bilinears:

$$h_0[\psi] = -\frac{1}{2m} \int dr^3 \psi^\dagger \tau_y (\sigma_x \partial_{yz}^2 + \sigma_y \partial_{zx}^2 + \sigma_z \partial_{xy}^2) \psi. \quad (7)$$

The minimal coupling terms to the electric field are

$$h_E[\psi, \mathbf{E}] = -\lambda_E \mathbf{E} \cdot \int dr^3 \psi^\dagger \tau_z \boldsymbol{\sigma} \psi; \quad (8)$$

the minimal coupling terms to the magnetic field

$$h_B[\psi, \mathbf{B}] = -\lambda_B \mathbf{B} \cdot \int dr^3 \psi^\dagger \boldsymbol{\sigma} \psi, \quad (9)$$

where $\lambda_{E,B}$ are coupling constants. From Eq.(8,9), we have the effective polarization and magnetization operators:

$$\begin{aligned} \hat{P}_i &= -\frac{\delta(h_0 + h_E + h_B)}{\delta E_i} = \lambda_E \tau_z \sigma_i, \\ \hat{M}_i &= -\frac{\delta(h_0 + h_E + h_B)}{\delta B_i} = \lambda_B \sigma_i. \end{aligned} \quad (10)$$

We note that Eq. (10) implies the quantum nature of polarization \hat{P}_i in this minimal effective theory, as different components do not commute

$$[\hat{p}_i, \hat{p}_j] = i\epsilon_{ijk} \hat{m}_k, \quad [\hat{p}_i, \hat{m}_j] = i\epsilon_{ijk} \hat{p}_k, \quad (11)$$

where we have defined the dimensionless reduced dipole operator $\hat{p}_i \equiv \hat{P}_i/(2\lambda_E)$ and magnetization operator $\hat{m}_i \equiv \hat{M}_i/(2\lambda_B)$.

TABLE I: Classification of hermitian operators, monomials of relative momentum and external electric, magnetic and strain tensor fields into their respective irreducible Reps of $O_h \times Z_2^T$, for a minimal effective theory having symmetries given in Eq.(6). The symbols of the irreducible Reps follow the conventional definition of Ref.[5], and the \pm in superscript denotes whether this Rep changes sign under \hat{T} . $\mathbf{B} \star \mathbf{q}$ stands for $(B_y q_z + B_z q_y, B_z q_x + B_x q_z, B_x q_y + B_y q_x)$.

Irreducible Rep	A_{2u}^-	A_{2g}^-	A_{1u}^-	T_{2u}^+	T_{2g}^+	T_{1u}^+	T_{1g}^-	A_{1g}^+
bilinear Operators	$\psi^\dagger \tau_x \psi$	$\psi^\dagger \tau_y \psi$	$\psi^\dagger \tau_z \psi$	$\psi^\dagger \tau_x \sigma_i \psi$	$\psi^\dagger \tau_y \sigma_i \psi$	$\psi^\dagger \tau_z \sigma_i \psi$	$\psi^\dagger \sigma_i \psi$	$\psi^\dagger \psi$
Applied fields		$B_x B_y B_z$	$\mathbf{E} \cdot \mathbf{B}$		$(E_y E_z, E_x E_z, E_x E_y)$ $(B_y B_z, B_x B_z, B_x B_y)$ $(\epsilon_{yz}, \epsilon_{xz}, \epsilon_{xy})$	E_i	B_i	E^2 B^2
Monomials of momentum	$q_x q_y q_z$				$(q_y q_z, q_x q_z, q_x q_y)$			q^2
cross coupling	$\epsilon_{yz} q_x + \epsilon_{zx} q_y + \epsilon_{xy} q_z$			$(\mathbf{B} \star \mathbf{q})_i$		$(\mathbf{B} \times \mathbf{q})_i$	$(\mathbf{E} \times \mathbf{q})_i$	$\mathbf{E} \times \mathbf{B} \cdot \mathbf{q}$

V. HIDDEN $SO(4)$ ALGEBRA, CROSS-LARMOR PRECESSION AND ABSORPTION

The commutation relations in Eq.(11) imply that \hat{p}_i and \hat{m}_i together form an $SO(4)$ algebra. Consider one quasiparticle magnetically polarized in the z -direction, that is, $\hat{m}_z |\psi(q=0)\rangle = |\psi(q=0)\rangle$. At $t=0$, an electric field was applied along x -direction. According to Eq.(11), at $t > 0$, we have

$$[\cos(2\lambda_E t) \hat{m}_z + \sin(2\lambda_E t) \hat{p}_y] |\psi(t)\rangle = |\psi(t)\rangle. \quad (12)$$

Specially, at $t = (n-1/4)\pi/\lambda_E$, the quasiparticle is completely electrically polarized along y -direction, and is spin unpolarized $\langle \psi(t) | \hat{m}_i | \psi(t) \rangle = 0$. Therefore an electric field rotates a magnetic dipole into an electric dipole, hence the name ‘‘cross-Larmor precession’’.

How could cross-Larmor precession be observed in experiments? The original Larmor precession is observed in the absorption in ferromagnets[18], where two elements are essential: a polarized spin configuration and an oscillating magnetic field component perpendicular to the magnetization. In our system, we first use a magnetic field \mathbf{B}_0 to spin-polarize the electrons at $\mathbf{q} = 0$ along z -direction. Then we use a polarized light propagating along y -direction, the \mathbf{E} -vector of which is polarized in the x -direction. With this geometry the Larmor precession is minimized, because the \mathbf{B} -vector of the light is parallel to the spin polarization. The cross-Larmor precession, on the other hand, is maximized because the electric field is perpendicular to the spin polarization. We therefore predict a resonant absorption edge at a frequency $|2\lambda_B \mathbf{B}_0|$, assuming that the system be exactly at half-filling. This is what we call a cross-resonant absorption, because the level splitting is obtained using a magnetic field, while the resonance is realized using an electric field. While $SO(4)$ algebra in Eq.(11) is a sufficient, but not necessary, condition for the presence of cross-resonant absorption. An example illustrating the non-necessity is shown in sec.D of appendix, where the other two Reps at \mathbf{R} in 222.103 are analyzed. There we show, though the $SO(4)$ algebra no longer holds, the cross-resonant absorption is still present.

We emphasize that neither the $SO(4)$ algebra nor the cross-resonant absorption is restricted to a specific

momentum $[\mathbf{R} = (\pi/a, \pi/a, \pi/a)]$ or a specific magnetic space group (222.103). The little group at \mathbf{R} in 222.103 is $G_{\mathbf{R}} = O_h \times Z_2^T$. As one breaks O_h down to T_h , D_{4h} and D_{2h} , one finds that $R_{1/2}$ remains an irreducible Rep. When $G_{\mathbf{R}} = T_h \times Z_2^T$, both the $SO(4)$ algebra and the cross-resonant absorption hold; and as $G_{\mathbf{R}}$ is reduced to $D_{4h,2h} \times Z_2^T$, the polarization operators $\hat{p}_{x,y,z}$ and magnetization operators $\hat{m}_{x,y,z}$ no longer form the six generators of $SO(4)$, but the cross-resonant absorption still appears (See sec.C of appendix for the derivation of the above statements). Due to the preservation of cross-resonant absorption in symmetry reduction, we predict that at least four magnetic space groups (222.103, 223.109, 200.17 and 201.21) in which certain irreducible Reps at \mathbf{R} shows $SO(4)$, and 12 groups (222.103, 223.109, 200.17, 201.21, 126.386, 130.432, 131.446, 135.492, 47.256, 48.264, 55.361 and 56.373) that potentially host cross-resonant absorption.

VI. GUIDELINES FOR FINDING MATERIALS

The four combinations $(\chi_S, \chi_T, \chi_{PT}) = (-1, +1, -1), (+1, -1, +1), (+1, +1, -1), (-1, -1, +1)$ are restricted to magnetic materials, realized as elementary excitations with momentum at corners of the mBZ, and they bring about novel physical effects as are discussed above. Hence it is vital that we provide clues as to where these excitations may be found in real materials. In mathematics, invariants such as χ_T , χ_{PT} and χ_S exactly correspond to the invariants of the second group cohomology[19], $\mathcal{H}^2[G_{\mathbf{Q}}, U(1)]$, of the little co-group $G_{\mathbf{Q}}$, which classify the projective Reps of $G_{\mathbf{Q}}$. Here we insert a few technical comments on $G_{\mathbf{Q}}$. As we focus on the elementary excitations centered at high-symmetry momentum \mathbf{Q} , the full magnetic space group M is reduced to the little group $M(\mathbf{Q}) \subset M$ that leaves \mathbf{Q} invariant, up to a reciprocal lattice vector. This little group has lattice translation Tr as its normal subgroup, so that a quotient group can be defined $M(\mathbf{Q})/Tr$, which is always isomorphic to a magnetic point group $G_{\mathbf{Q}}$, called the little co-group of M at \mathbf{Q} . As far as type-IV magnetic space groups are concerned, $G_{\mathbf{Q}}$ has the simple structure of a point group direct-product

time reversal. The invariants of each of the 32 $G_{\mathbf{Q}}$'s can be calculated, and are listed in Table II.

However, it is not obvious that all the possible values of these invariants may be taken in elementary excitations of real materials. While we are unable to address this general question, a full answer can be obtained, as discussion is restricted to excitations that form bands (magnon or electron). Throughout this article, symbols such as C_{2x} or T all refer to elements of some $G_{\mathbf{Q}}$. Yet, one should keep in mind that physically, $g \in G_{\mathbf{Q}}$ represent a coset in $M(\mathbf{Q})/Tr$, out of which a representative element $\tilde{g} \in M(\mathbf{Q})$ may be chosen. A degenerate-multiplet of Bloch eigenstates at \mathbf{Q} can either be considered as a *projective* Rep of $G_{\mathbf{Q}}$, or a linear Rep of $M(\mathbf{Q})$. Generically, \tilde{g} contains a fractional lattice translation, and the fractional translations and the particle statistics in \tilde{g} uniquely determine the invariants of $G_{\mathbf{Q}}$ [20]. In sec.B of appendix, we calculate the invariants for each high-symmetry momentum in every type-IV magnetic space group. In Table IV, we list all magnetic space groups with respective high-symmetry momenta where magnons and electrons carry $(\chi_S, \chi_T, \chi_{PT}) = (-1, +1, -1), (+1, -1, +1), (+1, +1, -1), (-1, -1, +1)$, values restricted to magnetic systems.

A final technical remark on terminologies. We use ‘‘quantum numbers’’ and ‘‘invariants’’ interchangeably for readability, as the former concept is familiar to non-experts. Rigorously speaking they are certainly not the same: different quantum numbers correspond to different irreducible Reps, while different invariants refer to different classes of Reps. The invariants of $G_{\mathbf{Q}}$ solely depends on M and \mathbf{Q} , and all the Reps at \mathbf{Q} share the same set of invariants.

Appendix A: Classification of Projective representations of anti-unitary groups $H \times Z_2^T$

1. Factor Systems and the Invariants

Projective representations(Reps) and the factor systems. As discussed in the main text, the little co-groups of type-IV magnetic space groups have the structure $G = H \times Z_2^T$ with $Z_2^T = \{E, \mathbb{T}\}$, $\mathbb{T}^2 = E$, where the physical meaning of \mathbb{T} will be specified later. Supposing g is a group element of G , then it is represented by $M(g)$ if g is a unitary element and represented by $M(g)K$ if g is anti-unitary, where K is the complex-conjugate operator satisfying $Ku = u^*K$ with u an arbitrary matrix and u^* its complex conjugation.

The multiplication of (projective) Reps of g_1, g_2 depends on if they are unitary or anti-unitary. If we define

$$s(g) = \begin{cases} 1, & \text{if } g \text{ is unitary,} \\ -1, & \text{if } g \text{ is antiunitary,} \end{cases}$$

and define the corresponding operator $K_{s(g)}$ as

$$K_{s(g)} = \begin{cases} I, & \text{if } s(g) = 1, \\ K, & \text{if } s(g) = -1, \end{cases}$$

then we have the multiplication rule of a projective Rep,

$$M(g_1)K_{s(g_1)}M(g_2)K_{s(g_2)} = M(g_1g_2)e^{i\theta_2(g_1, g_2)}K_{s(g_1g_2)},$$

where the $U(1)$ phase factor $\omega_2(g_1, g_2) \equiv e^{i\theta_2(g_1, g_2)}$ is a function of two group variables and is called the *factor system*. If $\omega_2(g_1, g_2) = 1$ for any $g_1, g_2 \in G$, then above projective Rep is trivial, namely, it is a linear Rep.

Substituting above results into the associativity relation of the sequence of operations $g_1 \times g_2 \times g_3$, we can obtain

$$\begin{aligned} & M(g_1)K_{s(g_1)}M(g_2)K_{s(g_2)}M(g_3)K_{s(g_3)} \\ &= M(g_1g_2g_3)\omega_2(g_1, g_2)\omega_2(g_1g_2, g_3)K_{s(g_1g_2g_3)} \\ &= M(g_1g_2g_3)\omega_2(g_1, g_2g_3)\omega_2^{s(g_1)}(g_2, g_3)K_{s(g_1g_2g_3)}, \end{aligned}$$

namely,

$$\omega_2(g_1, g_2)\omega_2(g_1g_2, g_3) = \omega_2^{s(g_1)}(g_2, g_3)\omega_2(g_1, g_2g_3). \quad (\text{A1})$$

Eq.(A1) is the general relation that the factor systems of any finite group (no matter unitary or anti-unitary) should satisfy. If we introduce a gauge transformation $M'(g)K_{s(g)} = M(g)\Omega_1(g)K_{s(g)}$, where the phase factor $\Omega_1(g) = e^{i\theta_1(g)}$ depends on a single group variable, then the factor system changes into

$$\omega_2'(g_1, g_2) = \omega_2(g_1, g_2)\Omega_2(g_1, g_2), \quad (\text{A2})$$

with

$$\Omega_2(g_1, g_2) = \frac{\Omega_1(g_1)\Omega_1^{s(g_1)}(g_2)}{\Omega_1(g_1g_2)}. \quad (\text{A3})$$

The equivalent relations (A2) and (A3) define the equivalent classes of the solutions of (A1). The number of equivalent classes for a finite group is usually finite.

The 2-cocycles and the 2nd group cohomology.

The factor systems (A1) of projective Reps of group G are also called cocycles. The equivalent classes of the cocycles form a group, *i.e.* the group-cohomology group. The group cohomology ${}^1\{\text{Kernel } d/\text{Image } d\}$ is defined by the coboundary operator d

$$\begin{aligned} & (d\omega_n)(g_1, \dots, g_{n+1}) \\ &= [g_1 \cdot \omega_n(g_2, \dots, g_{n+1})]\omega_n^{(-1)^{n+1}}(g_1, \dots, g_n) \times \\ & \prod_{i=1}^n \omega_n^{(-1)^i}(g_1, \dots, g_{i-1}, g_i g_{i+1}, g_{i+2}, \dots, g_{n+1}). \quad (\text{A4}) \end{aligned}$$

¹ For an introduction to group cohomology, see wiki http://en.wikipedia.org/wiki/Group_cohomology and Romyar Sharifi, [AN INTRODUCTION TO GROUP COHOMOLOGY](#)

where $g_1, \dots, g_{n+1} \in G$ and the variables $\omega_n(g_1, \dots, g_n)$ take value in an Abelian coefficient group \mathcal{A} [usually \mathcal{A} is a subgroup of $U(1)$, in the present work $\mathcal{A} = U(1)$]. The set of variables $\omega_n(g_1, \dots, g_n)$ is called a n -cochain. For anti-unitary groups the module $g \cdot$ is defined by

$$g \cdot \omega_n(g_1, \dots, g_n) = \omega_n^{s(g)}(g_1, \dots, g_n).$$

With this notation, Eq. (A1) can be rewritten as

$$(d\omega_2)(g_1, g_2, g_3) = 1,$$

the solutions of above equations are called 2-cocycles with $U(1)$ -coefficient. Similarly, Eq. (A3) can be rewritten as

$$\Omega_2(g_1, g_2) = (d\Omega_1)(g_1, g_2),$$

where $\Omega_1(g_1), \Omega_1(g_2) \in U(1)$ and $\Omega_2(g_1, g_2)$ are called 2-coboundaries. Two 2-cocycles $\omega_2'(g_1, g_2)$ and $\omega_2(g_1, g_2)$ are equivalent if they differ by a 2-coboundary, see Eq. (A2). The equivalent classes of the 2-cocycles $\omega_2(g_1, g_2)$ form the second group cohomology $\mathcal{H}^2(G, U(1))$.

Writing $\omega_2(g_1, g_2) = e^{i\theta_2(g_1, g_2)}$, where $\theta_2(g_1, g_2) \in [0, 2\pi)$, then the cocycle equations $(d\omega_2)(g_1, g_2, g_3) = 1$ can be written in terms of linear equations,

$$\begin{aligned} s(g_1)\theta_2(g_2, g_3) - \theta_2(g_1g_2, g_3) + \theta_2(g_1, g_2g_3) \\ - \theta_2(g_1, g_2) = 0. \end{aligned} \quad (\text{A5})$$

Similarly, if we write $\Omega_1(g_1) = e^{i\theta_1(g_1)}$ and $\Omega_2(g_1, g_2) = e^{i\Theta_2(g_1, g_2)}$, then the 2-coboundary (A3) can be written as

$$\Theta_2(g_1, g_2) = s(g_1)\theta_1(g_2) - \theta_1(g_1g_2) + \theta_1(g_1). \quad (\text{A6})$$

The equal sign in (A5) and (A6) means equal mod 2π . From these linear equations, we can obtain the solution space of the cocycle equations, as well as the classes that the solutions belong to. The set of classes forms a finite Abelian group, which labels the classification of the projective Reps.

Invariants of projective Reps. The second group-cohomology group is generated by a certain number of *invariants*. The invariants are, by definition, invariant under the gauge transformation (A2) and (A3). They are formed by independent functions of the cocycles $\omega_2(g_1, g_2)$. For instance, for the unitary group $D_2 = Z_2 \times Z_2 = \{E, P\} \times \{E, Q\}$ with $P^2 = E, Q^2 = E$, the classification of 2-cocycles is

$$\mathcal{H}^2(D_2, U(1)) = \mathbb{Z}_2,$$

there is only one independent invariant which is given by $\chi = \frac{\omega_2(P, Q)}{\omega_2(Q, P)}$. Another example is the simplest anti-unitary group $Z_2^T = \{E, \mathbb{T}\}$, which has

$$\mathcal{H}^2(Z_2^T, U(1)) = \mathbb{Z}_2,$$

with the invariant $\chi_T = \omega_2(\mathbb{T}, \mathbb{T})$.

Generally, for any group G , if $\mathcal{H}^2(G, U(1)) = \mathbb{Z}_2^n$, then the 2-cocycles of G have n invariants, each taking value

+1 or -1. The invariants of some anti-unitary groups will be given later.

Irreducible projective Reps. For the cocycle solution of the each class, we can construct the regular projective Rep, from which all of the inequivalent irreducible projective Reps can be obtained. The method is provided in Ref. [21] and we will not go to details here.

2. Relation Between the Classification of projective Reps of $H \times Z_2^T$ and that of H

The group cohomology of $H \times Z_2^T$ is closely related to that of H . Before identifying this relation, we first introduce Z_2 -coefficient group cohomology and clarify its relation to the $U(1)$ -coefficient group cohomology.

$U(1)$ -coefficient and Z_2 -coefficient It can be shown that the $U(1)$ -coefficient and Z_2 -coefficient second cohomology has the following relation,

$$\begin{aligned} \mathcal{H}^2(G, Z_2) &= [\mathcal{H}^2(G, U(1))]_{\mathbb{Z}_2} \times [\mathcal{H}^2(G, Z)]_{\mathbb{Z}_2} \\ &= [\mathcal{H}^2(G, U(1))]_{\mathbb{Z}_2} \times \mathcal{H}^1(G, Z_2), \end{aligned} \quad (\text{A7})$$

where $[\mathcal{H}^2(G, U(1))]_{\mathbb{Z}_2} = \mathcal{H}^1(\mathcal{H}^2(G, U(1)), Z_2)$. The first part on the right-hand side indicates that the $U(1)$ -coefficient cohomology group $\mathcal{H}^2(G, U(1))$ contributes to the Z_2 -coefficient classification only if it has a Z_2 representations; the second part $[\mathcal{H}^2(G, Z)]_{\mathbb{Z}_2} = \mathcal{H}^1(G, Z_2)$ originates from the collapsing of certain $U(1)$ -coboundaries of G when their values are constrained to $Z_2 = \{1, -1\}$.

For example, $\mathcal{H}^2(Z_4 \times Z_4, U(1)) = \mathbb{Z}_4$ and $\mathcal{H}^1(Z_4 \times Z_4, Z_2) = \mathbb{Z}_2 \times \mathbb{Z}_2$, therefore

$$\mathcal{H}^2(Z_4 \times Z_4, Z_2) = \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2,$$

where one of the \mathbb{Z}_2 comes from $[\mathcal{H}^2(Z_4 \times Z_4, U(1))]_{\mathbb{Z}_2}$ and the other two \mathbb{Z}_2 come from $\mathcal{H}^1(Z_4 \times Z_4, Z_2)$.

Classification of $H \times Z_2^T$. Firstly, for 2-cocycles of any group H , we adopt the gauge such that $\omega_2(E, h) = \omega_2(h, E) = 1$ for any $h \in H$. Furthermore, for anti-unitary groups $H \times Z_2^T$, one can further fix the gauge such that the following relations are satisfied (see Ref.[21] for details)

$$\begin{aligned} \omega_2(\mathbb{T}, h) &= \omega_2(h, \mathbb{T}) = 1, \\ \omega_2(\mathbb{T}, \mathbb{T}h) &= \omega_2(\mathbb{T}h, \mathbb{T}) = \omega_2(\mathbb{T}, \mathbb{T}), \\ \omega_2(\mathbb{T}h_1, \mathbb{T}h_2) &= \omega_2(\bar{h}_1, h_2)\omega_2(\mathbb{T}, \mathbb{T}), \\ \omega_2(\mathbb{T}h_1, h_2) &= \omega_2^{-1}(h_1, h_2), \\ \omega_2(h_1, \mathbb{T}h_2) &= \omega_2(h_1, \bar{h}_2), \end{aligned}$$

with the constraint

$$\omega_2(h_1, h_2)\omega_2(\bar{h}_1, \bar{h}_2) = 1,$$

where $\bar{h} = \mathbb{T}h\mathbb{T}$, $h, h_1, h_2 \in H$. Therefore, the classification of the second group cohomology is dependent on

$\omega_2(\mathbb{T}, \mathbb{T})$ and $\omega_2(h_1, h_2)$. For the direct product group $H \times Z_2^T$, we have $\tilde{h} = h, \omega_2(h_1, h_2) = \pm 1$ and consequently

$$\begin{aligned} \mathcal{H}^2(H \times Z_2^T, U(1)) &= \mathcal{H}^2(Z_2^T, U(1)) \times \mathcal{H}^2(H, Z_2) \quad (\text{A8}) \\ &= \mathbb{Z}_2 \times [\mathcal{H}^2(H, U(1))]_{\mathbb{Z}_2} \times \mathcal{H}^1(H, Z_2), \end{aligned}$$

where $\mathcal{H}^1(H, Z_2)$ stands for the twisting between H and \mathbb{T} , which corresponds to the general Kramers classes $(g_0\mathbb{T})^2 = -1$ for $g_0^2 = E$. In the second line of the equation we have used the result (A7).

For instance, for the group $G = Z_2 \times Z_2^T$, since $\mathcal{H}^2(Z_2, U(1)) = \mathbb{Z}_1, \mathcal{H}^1(Z_2, Z_2) = \mathbb{Z}_2$, we have

$$\begin{aligned} \mathcal{H}^2(Z_2 \times Z_2^T, U(1)) \\ = \mathcal{H}^2(Z_2^T, U(1)) \times [\mathcal{H}^2(Z_2, U(1))]_{\mathbb{Z}_2} \times \mathcal{H}^1(Z_2, Z_2) = \mathbb{Z}_2^2. \end{aligned}$$

3. Results for abstract and concrete groups

In Table II, we give the results of the second group cohomology of several direct product groups $G = H \times Z_2^T$. All the groups appeared in Table II are abstract groups. The physical meaning of these groups are specified by mapping these abstract groups into concrete magnetic point groups.

In Table III, we list the correspondence between the abstract group $H \times Z_2^T$ and the isomorphic magnetic point groups. As before, the abstract group Z_2^T stands for $Z_2^T = \{E, \mathbb{T}\}$, where the anti-unitary generator \mathbb{T} can be mapped to either T or a product of T with other unitary element. Here we have exclusively denoted T as pure time reversal operation. The following notations have been used: $C_{2m}, C_{4m}^\pm (m = x, y, z), C_{3j}^\pm (j = 1, 2, 3, 4), C_2, C_{2i}', C_{2i}'' (i = 1, 2, 3), C_{2p} (p = a, b, c, d, e, f)$, see Fig 1.1, 1.2 and 1.3 of Ref.[5], $P =$ space inversion, $\mathcal{M} =$ mirror reflection plane, $PC_2 = \mathcal{M}_h, PC_{2m} = \mathcal{M}_m, PC_{2i}' = \mathcal{M}_{di}, PC_{2i}'' = \mathcal{M}_{vi}, PC_{2p} = \mathcal{M}_{dp}, PC_3^\pm = S_6^\mp, PC_{4m}^\pm = S_{4m}^\mp, PC_6^\pm = S_3^\mp, PC_{3j}^\pm = S_{6j}^\mp$.

The definitions of invariants of the second group cohomology for the concrete groups are also given in Table III. Notice that for $G_{\mathbf{Q}} = D_{2h} \times Z_2^T, D_{4h} \times Z_2^T, D_{6h} \times Z_2^T, T_h \times Z_2^T, O_h \times Z_2^T$, the first three invariants correspond to the quantum numbers $(\chi_S, \chi_T, \chi_{PT})$ that were discussed in main text.

Appendix B: Invariants for the little co-groups in type-IV magnetic space groups

For a Shubnikov magnetic space group M , the magnetic little group $M(\mathbf{Q})$ is the subgroup of M which transforms the wave vector \mathbf{Q} in the first Brillouin zone (BZ) into its equivalent wave vector $\mathbf{Q} + \mathbf{K}$, here \mathbf{K} is reciprocal lattice vector. $M(\mathbf{Q})$ is also a Shubnikov space group which has the translation group as its normal subgroup. The point group of $M(\mathbf{Q})$ is called magnetic little

co-group $G_{\mathbf{Q}}$, which is the quotient group of $M(\mathbf{Q})$ with respect to the translation group as discussed in the main text.

A type IV Shubnikov space group [5] has the form $M = H + T\{E|\mathbf{t}_0\}H$, where H belongs to the 230 (Fedorov) space groups and \mathbf{t}_0 is the extra fractional translation about the time-reversal operation T . The magnetic little co-group of M at point \mathbf{Q} is isomorphic to the direct product of a point group H and Z_2^T , namely, $G_{\mathbf{Q}} = H \times Z_2^T$, where $Z_2^T = \{E, T\}$ [if T is an element of the magnetic point group of $M(\mathbf{Q})$, or equivalently $\tilde{T} \in M(\mathbf{Q})$, where \tilde{T} is defined in (B1) below] or $Z_2^T = \{E, PT\}$ (if $\tilde{T} \notin M(\mathbf{Q})$) depending on the symmetry of the high symmetry point (also see Table III). For all the type IV magnetic space groups of our interest whose point group contains P , we always have $PT \in G_{\mathbf{Q}}$.

As illustrated in the main text, the linear Rep of the Shubnikov space group M at some high symmetry point \mathbf{Q} (or on a high symmetry line) defines a projective Rep of the little co-group $G_{\mathbf{Q}}$. Now we derive the details here. For any group element $g \in G_{\mathbf{Q}}$, we denote

$$\tilde{g} = \{g|\mathbf{t}_g\} \in M, \quad (\text{B1})$$

where $\mathbf{t}_g = x_g\mathbf{a} + y_g\mathbf{b} + z_g\mathbf{c}$ is the unique fractional translation associated with g , with $\mathbf{a}, \mathbf{b}, \mathbf{c}$ the basis vectors of the lattice and $0 \leq x_g, y_g, z_g < 1$. Then we further denote \hat{g} as the linear Rep matrix of $\tilde{g} \in M$. Then for $\tilde{g}_1, \tilde{g}_2 \in M$, we have the group multiplication rule (for simplicity, we temporarily assume that both of \tilde{g}_1, \tilde{g}_2 are unitary operations)

$$\begin{aligned} \tilde{g}_1\tilde{g}_2 &= \{g_1|\mathbf{t}_{g_1}\}\{g_2|\mathbf{t}_{g_2}\} = \{g_1g_2|g_1\mathbf{t}_{g_2} + \mathbf{t}_{g_1}\} \\ &= \{g_1g_2|\mathbf{t}_{g_1g_2}\}\omega(g_1, g_2), \end{aligned}$$

where $\omega(g_1, g_2) = \{E|g_1\mathbf{t}_{g_2} + \mathbf{t}_{g_1} - \mathbf{t}_{g_1g_2}\}$ is an element of the translation group. Accordingly, we have

$$\hat{g}_1\hat{g}_2 = \widehat{g_1g_2}\hat{\omega}(g_1, g_2)$$

where $\hat{\omega}(g_1, g_2) = e^{-i\mathbf{Q}\cdot(g_1\mathbf{t}_{g_2} + \mathbf{t}_{g_1} - \mathbf{t}_{g_1g_2})}$. Therefore, if we treat \hat{g} as a projective Rep of $g \in G_{\mathbf{Q}}$, then $\hat{\omega}(g_1, g_2)$ defines the factor system. It can be verified that thus defined factor system satisfy the 2-cocycle equation. On the other hand, if we know the projective Rep of $G_{\mathbf{Q}}$ with the factor system $\hat{\omega}(g_1, g_2)$ defined above, then we can easily obtain the linear Rep of the total magnetic space group M .

Actually, up to a gauge transformation, the factor system mentioned above can be obtained in a systematic way[5, 20, 22]. In the following we generalize the rule introduced in Ref.[5, 20, 22] to the anti-unitary groups (*i.e.* magnetic space groups). For arbitrary elements $g_1, g_2 \in G_{\mathbf{Q}}$ (and accordingly $\tilde{g}_1, \tilde{g}_2 \in M$), we have the factor system

$$\omega_2(g_1, g_2) = e^{-i\mathbf{K}_{g_1}\cdot\mathbf{t}_{g_2}}, \quad (\text{B2})$$

where $\mathbf{K}_{g_1} = s(g_1)(g_1^{-1}\mathbf{Q} - \mathbf{Q})$ is a reciprocal lattice vector and \mathbf{t}_{g_2} is the fractional translation associated with

g_2 . For any $g_1, g_2, g_3 \in G_{\mathbf{Q}}$, one can verify that the cycle equation (A1) is satisfied,

$$\begin{aligned}
& \omega_2(g_1, g_2 g_3) \omega_2^{s(g_1)}(g_2, g_3) \\
&= e^{-i\mathbf{K}_{g_1} \cdot \mathbf{t}_{g_2 g_3}} e^{-is(g_1)\mathbf{K}_{g_2} \cdot \mathbf{t}_{g_3}} \\
&= e^{-i\mathbf{K}_{g_1} \cdot \mathbf{t}_{g_2}} e^{-i\mathbf{K}_{g_1 \cdot g_2} \mathbf{t}_{g_3}} e^{-is(g_1)\mathbf{K}_{g_2} \cdot \mathbf{t}_{g_3}} \\
&= e^{-i\mathbf{K}_{g_1} \cdot \mathbf{t}_{g_2}} e^{-is(g_2)g_2^{-1}\mathbf{K}_{g_1} \cdot \mathbf{t}_{g_3}} e^{-is(g_1)\mathbf{K}_{g_2} \cdot \mathbf{t}_{g_3}} \\
&= e^{-i\mathbf{K}_{g_1} \cdot \mathbf{t}_{g_2}} e^{-is(g_2)s(g_1)[(g_1 g_2)^{-1}\mathbf{Q} - g_2^{-1}\mathbf{Q}] \cdot \mathbf{t}_{g_3}} \\
&\quad \cdot e^{-is(g_1)s(g_2)(g_2^{-1}\mathbf{Q} - \mathbf{Q}) \cdot \mathbf{t}_{g_3}} \\
&= e^{-i\mathbf{K}_{g_1} \cdot \mathbf{t}_{g_2}} e^{-is(g_1 g_2)[(g_1 g_2)^{-1}\mathbf{Q} - \mathbf{Q}] \cdot \mathbf{t}_{g_3}} \\
&= e^{-i\mathbf{K}_{g_1} \cdot \mathbf{t}_{g_2}} e^{-i\mathbf{K}_{g_1 g_2} \cdot \mathbf{t}_{g_3}} \\
&= \omega_2(g_1, g_2) \omega_2(g_1 g_2, g_3), \tag{B3}
\end{aligned}$$

here $\mathbf{t}_{g_2 g_3} = \mathbf{t}_{g_2} + g_2 \mathbf{t}_{g_3} + \mathbf{R}_{2g_3}$, $\mathbf{K}_{g_1 \cdot g_2} \mathbf{t}_{g_3} = s(g_2)g_2^{-1}\mathbf{K}_{g_1} \cdot \mathbf{t}_{g_3}$ and \mathbf{R}_{2g_3} is a lattice vector.

For bosonic particles whose bare spin s_0 is an integer, the band structure is characterized by projective Rep of $G_{\mathbf{Q}}$ with the factor system (B2), now we note it as $\omega_{2b}(g_1, g_2)$. The bosonic particles are characterized by the invariant $\chi_{PT} = \omega_2(PT, PT) = (-1)^{2s_0} = 1$ (this is no surprise because $(PT)^{-1}\mathbf{Q} = \mathbf{Q}$ such that $\mathbf{K}_{PT}=0$). The magnon excitations in magnetic ordered ground states belong to this case.

For fermionic particles whose bare spin s_0 is a half-odd-integer, an extra factor system $\omega_2^{(\frac{1}{2})}(g_1, g_2)$ is contributed from the spin rotation (owing to spin-orbit coupling), which corresponds to the double valued Reps of the little co-group. The non-trivial invariants of factor system $\omega_2^{(\frac{1}{2})}(g_1, g_2)$ are $\chi_S = \frac{\omega_2(C_{2m}, C_{2n})}{\omega_2(C_{2n}, C_{2m})} = -1, \chi_T = \omega_2(T, T) = -1, \chi_{PT} = \omega_2(PT, PT) = -1$, where $m \neq n$ take values x, y, z . Combining with the factor system from fractional translation, the total factor system for fermions is given by

$$\omega_{2f}(g_1, g_2) = \omega_2^{(\frac{1}{2})}(g_1, g_2) \omega_{2b}(g_1, g_2).$$

The fermionic particles (such as electrons) are characterized by the invariant $\chi_{PT} = -1$.

From the given factor systems and the definition of the invariants in Table III, we can easily obtain the values of the invariants. In Table IV, we list the little co-groups of the high symmetry points of type IV Shubnikov space groups, and also provide the values of the invariants for the two different classes of factors system discussed above. The values $(-1, +1, -1), (+1, -1, +1), (+1, +1, -1), (-1, -1, +1)$ of $(\chi_S, \chi_T, \chi_{PT})$ are only accessible in magnetic materials, which are marked with red for electrons, and blue for magnons.

Appendix C: $SO(4)$ algebra and cross-resonant absorption under symmetry reduction

In this section we derive a few statements made in the main article, with respect to whether the $SO(4)$ al-

gebra, among polarization and magnetization operators for the $R_{1/2}$ -Rep, as well as the cross-resonant absorption phenomenon, in 222.103 is preserved or broken, if the little co-group is broken from $O_h \times Z_2^T$ to $T_h \times Z_2^T$, $D_{4h} \times Z_2^T$ and $D_{2h} \times Z_2^T$. These symmetry reduction naturally comes into consideration as one (i) breaks the fourfold axis into a twofold axis (T_h), (ii) breaks the threefold axis (D_{4h}), and (iii) both the fourfold and the threefold axis (D_{2h}), leaving the parity, time-reversal, and three twofold axes intact.

Our first statement is that, after the reduction, $R_{1/2}$ remains an irreducible Rep for the little co-group. In all three scenarios, $D_{2h} \times Z_2^T$ is a subgroup of the little co-group, therefore, the three invariants $\chi_{S,T,PT}$ simply ‘‘inherit’’ their respective values in 222.103, $(\chi_S, \chi_T, \chi_{PT}) = (-1, +1, -1)$, because these three invariants only involve parity, time-reversal, and twofold axes, all inside $D_{2h} \times Z_2^T$. From Eq.(6) in main text, we have

$$\hat{T} = \tau_y \sigma_y K, \tag{C1}$$

$$\hat{P}T = i\sigma_y K, \tag{C2}$$

$$\hat{C}_{2x,2y,2z} = i\sigma_{x,y,z}. \tag{C3}$$

From Eq.(C3), we see that mass terms (matrices that commute with all symmetris) must take the form $\tau_{\mu=0,1,2,3}\sigma_0$, yet Eq.(C1) narrows these possibilities down to $\tau_0\sigma_0$. A constant mass term cannot open a gap, so $R_{1/2}$ remains an irreducible Rep under all three types of symmetry reduction.

The next statement is that for $O_h \rightarrow T_h$, the $SO(4)$ algebra among polarization and magnetization operators still holds. We make use of Table I to simplify our derivation. We observe that as $O_h \rightarrow T_h$, the Reps such as A_{1u}^+ and A_{2u}^+ become identical, because their difference solely appears in how they transform under C_4 , broken in T_h . Similarly, T_{1u}^+ and T_{2u}^+ become the same T_u^+ . The electric-field components (E_x, E_y, E_z) form a T_u^+ -Rep, so there are two linearly independent sets of Γ -matrices that couple to the electric field:

$$\hat{d}_{i=x,y,z}^{(1)} = \tau_z \sigma_i / 2, \tag{C4}$$

$$\hat{d}_{i=x,y,z}^{(2)} = \tau_x \sigma_i / 2.$$

The physical (reduced) polarization \hat{p} is a linear superposition of $\hat{d}^{(1)}$ and $\hat{d}^{(2)}$:

$$\hat{p}_i = \hat{d}_i^{(1)} \cos \theta + \hat{d}_i^{(2)} \sin \theta. \tag{C5}$$

On the other hand, while $T_{1g,2g}^- \rightarrow T_g^-$ as $O_h \rightarrow T_h$, according to Table I there is not a T_{2g}^- for all 16 Γ -matrices, so T_g^- -matrices are just the T_{1g}^- -matrices in Table I. Physically, this means that there is not any other choice for the (reduced) magnetization:

$$\hat{m}_i = \tau_0 \sigma_i / 2. \tag{C6}$$

It is obvious from Eq.(C5,C6) that \hat{p}_i, \hat{m}_i still satisfy Eq.(11) in the main text, i.e., the $SO(4)$ algebra is preserved.

Our third statement is that as symmetry is reduced to D_{4h} , the $SO(4)$ algebra is broken. We show this, again, by using Table I and the reduction of irreducible Rep from O_h to D_{4h} : $T_{1u}^+ \rightarrow A_{2u} \oplus E_u, T_{2u}^+ \rightarrow B_{2u} \oplus E_u$. The electric-field component E_z belongs to A_{2u} and (E_x, E_y) belong to E_u . Therefore there is only one choice for \hat{p}_z , yet two choices for $\hat{p}_{x,y}$:

$$\begin{aligned}\hat{p}_z &= \tau_z \sigma_z / 2, \\ \hat{p}_{x,y} &= \hat{d}_{x,y}^{(1)} \cos \theta + \hat{d}_{x,y}^{(2)} \sin \theta.\end{aligned}\quad (\text{C7})$$

Since there is not a T_{2g}^- -Rep in Table I, the choice for \hat{m}_i remains the same. Obviously, the \hat{p}_i operators in Eqs.(C7) do not form an $SU(2)$ algebra, so the $SO(4)$ algebra is also broken. For the D_{2h} scenario, the derivation is similar to the T_h case, with the only difference that, in D_{2h} , the θ in Eq.(C5) becomes three different values $\theta_{x,y,z}$ for three components.

$$\hat{p}_i = \hat{d}_i^{(1)} \cos \theta_i + \hat{d}_i^{(2)} \sin \theta_i. \quad (\text{C8})$$

Due to the component-dependent θ_i , \hat{p}_i do not form an $SU(2)$ algebra, and the $SO(4)$ algebra is again broken.

Our final statement is that even in the absence of $SO(4)$ algebra, the cross-resonant absorption still holds under symmetry reduction. For this purpose, we only need to show the absorption in D_{2h} -scenario, because it has the lowest symmetry. Consider an \hat{m}_z -polarized state ψ_0 : $\hat{m}_i \psi_0 = \delta_{iz} \psi_0$. Using the explicit expression of \hat{m}_z , we have

$$\psi = u|11\rangle + v|\bar{1}\bar{1}\rangle, \quad (\text{C9})$$

where $|mn\rangle$ denotes an eigenstate of τ_z, σ_z of eigenvalues m and n , respectively. Now consider an electric field along x -axis, so that the Hamiltonian is

$$\hat{h} = \lambda_{E,x} (\tau_z \sigma_x \cos \theta_x + \tau_x \sigma_x \sin \theta_x). \quad (\text{C10})$$

At $t > 0$, we have

$$\begin{aligned}\psi(t) &= e^{-i\hat{h}t} \psi_0 \\ &= \cos(\lambda_{E,x} t) \psi_0 - i \sin(\lambda_{E,x} t) (\tau_z \sigma_x \cos \theta_x + \tau_x \sigma_x \sin \theta_x) \psi_0 \\ &= \cos(\lambda_{E,x} t) (u|11\rangle + v|\bar{1}\bar{1}\rangle) - i \sin(\lambda_{E,x} t) \cos \theta_x (u|1\bar{1}\rangle - v|\bar{1}\bar{1}\rangle) - i \sin(\lambda_{E,x} t) \sin \theta_x (u|\bar{1}\bar{1}\rangle + v|1\bar{1}\rangle) \\ &= \cos(\lambda_{E,x} t) (u|11\rangle + v|\bar{1}\bar{1}\rangle) - i \sin(\lambda_{E,x} t) (u \cos \theta_x + v \sin \theta_x) |1\bar{1}\rangle - i \sin(\lambda_{E,x} t) (u \sin \theta_x - v \cos \theta_x) |\bar{1}\bar{1}\rangle.\end{aligned}\quad (\text{C11})$$

Eq.(C11) shows a clear oscillation between ψ_0 and an orthogonal $\psi'_0 = (u \cos \theta_x + v \sin \theta_x) |1\bar{1}\rangle + (u \sin \theta_x - v \cos \theta_x) |\bar{1}\bar{1}\rangle$, with period $T = \pi / \lambda_{E,x}$. Therefore, if an incident light is propagating along y -axis and polarized along x -axis, it has an absorption peak at $\omega_0 = 2\pi / T = 2\lambda_{E,x}$.

Appendix D: The other two Reps of $O_h \times Z_2^T$ at $(\pi/a, \pi/a, \pi/a)$ in 222.103

In addition to Eq.(6), there exist another two inequivalent 4-dim irreducible Reps $R_{3/2}^\pm$. By using the three spin- $\frac{3}{2}$ angular momentum operators

$$J_x = \frac{1}{2} \begin{pmatrix} 0 & \sqrt{3} & 0 & 0 \\ \sqrt{3} & 0 & 2 & 0 \\ 0 & 2 & 0 & \sqrt{3} \\ 0 & 0 & \sqrt{3} & 0 \end{pmatrix}, \quad (\text{D1})$$

they are written as

$$J_y = -\frac{i}{2} \begin{pmatrix} 0 & \sqrt{3} & 0 & 0 \\ -\sqrt{3} & 0 & 2 & 0 \\ 0 & -2 & 0 & \sqrt{3} \\ 0 & 0 & -\sqrt{3} & 0 \end{pmatrix}, \quad (\text{D2})$$

$$J_z = \frac{1}{2} \begin{pmatrix} 3 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -3 \end{pmatrix}, \quad (\text{D3})$$

$$\begin{aligned}\hat{T} &= \tau_z \sigma_x K, \\ \hat{P} &= i\tau_y \sigma_z, \\ \hat{C}_{2x} &= \exp(iJ_x \pi) = -i\tau_x \sigma_x, \\ \hat{C}_{31} &\equiv \hat{C}_{3,111} = -\exp\left(-\frac{2\pi i J_{111}}{3}\right), \\ \hat{C}_{4z} &= -i \exp\left(-\frac{iJ_z \pi}{2}\right) = -\frac{1}{\sqrt{2}} (\tau_z \sigma_0 - i\tau_z \sigma_z); (\text{D4})\end{aligned}$$

and

$$\begin{aligned}
\hat{T} &= \tau_z \sigma_x K, \\
\hat{P} &= -i\tau_y \sigma_z, \\
\hat{C}_{2x} &= \exp(iJ_x \pi) = -i\tau_x \sigma_x, \\
\hat{C}_{31} &\equiv \hat{C}_{3,111} = -\exp\left(-\frac{2\pi i J_{111}}{3}\right), \\
\hat{C}_{4z} &= i \exp\left(-\frac{iJ_z \pi}{2}\right) = \frac{1}{\sqrt{2}}(\tau_z \sigma_0 - i\tau_z \sigma_z), \quad (\text{D5})
\end{aligned}$$

here $J_{111} \equiv \frac{J_x + J_y + J_z}{\sqrt{3}}$. The two Reps (D4) and (D5) have quadratic (T_{2g}^+) and quintic (E_u^-) dispersion terms. They can couple electric and magnetic fields, but the minimal coupling terms about electric field do not have the $SO(4)$ algebra as Eq.(11). Even so, all the three inequivalent Reps (6), (D4) and (D5) can lead to cross-resonant absorption. Furthermore, when the above two Reps are reduced from $O_h \times Z_2^T$ to $T_h \times Z_2^T$, $D_{4h} \times Z_2^T$ and $D_{2h} \times Z_2^T$, the cross-resonant absorption also appears, which will be discussed in the future.

TABLE II: Some direct product groups $G = H \times Z_2^T$ and their 2nd group cohomology which classify their projective Reps.

Abstract group G	$\mathcal{H}^2(G, U(1))$	Abstract group G	$\mathcal{H}^2(G, U(1))$	Abstract group G	$\mathcal{H}^2(G, U(1))$
Z_2^T	\mathbb{Z}_2	$(Z_4 \times Z_2) \times Z_2^T$	\mathbb{Z}_2^4	$(Z_6 \times Z_2) \times Z_2^T$	\mathbb{Z}_2^4
$Z_2 \times Z_2^T$	\mathbb{Z}_2^2	$[(Z_4 \rtimes Z_2) \times Z_2] \times Z_2^T$	\mathbb{Z}_2^7	$[(Z_6 \rtimes Z_2) \times Z_2] \times Z_2^T$	\mathbb{Z}_2^7
$(Z_2 \times Z_2) \times Z_2^T$	\mathbb{Z}_2^4	$Z_3 \times Z_2^T$	\mathbb{Z}_2	$A_4 \times Z_2^T$	\mathbb{Z}_2^2
$[(Z_2 \times Z_2) \times Z_2] \times Z_2^T$	\mathbb{Z}_2^7	$Z_6 \times Z_2^T$	\mathbb{Z}_2^2	$(A_4 \times Z_2) \times Z_2^T$	\mathbb{Z}_2^3
$Z_4 \times Z_2^T$	\mathbb{Z}_2^2	$(Z_3 \rtimes Z_2) \times Z_2^T$	\mathbb{Z}_2^2	$(A_4 \rtimes Z_2) \times Z_2^T$	\mathbb{Z}_2^3
$(Z_4 \rtimes Z_2) \times Z_2^T$	\mathbb{Z}_2^4	$(Z_6 \rtimes Z_2) \times Z_2^T$	\mathbb{Z}_2^4	$[(A_4 \rtimes Z_2) \times Z_2] \times Z_2^T$	\mathbb{Z}_2^5

TABLE III: Correspondence between abstract groups and concrete magnetic point groups, where the physical meaning of T (the generator of Z_2^T) is specified as the last generator as listed in the third column. In the fourth column, we list all the invariants of the second group cohomology. Here, $\chi_T \equiv \omega_2(T, T)$, $\chi_{PT} \equiv \omega_2(PT, PT)$, $\chi_{TC_2} \equiv \omega_2(TC_2, TC_2)$, $\chi_{TM} \equiv \omega_2(TM, TM)$.

Abstract group	Concrete group	Generators	Invariants
$Z_2 \times Z_2^T$	$C_i \times Z_2^T$	P, T	(χ_T, χ_{PT})
$(Z_2 \times Z_2) \times Z_2^T$	$C_{2v} \times Z_2^T$ $D_2 \times Z_2^T$ $C_{2h}^1 \times Z_2^T$ $C_{2h}^2 \times Z_2^T$ $C_{2h}^3 \times Z_2^T$ $C_{2h}^4 \times Z_2^T$ $C_{2h}^5 \times Z_2^T$ $C_{2h}^6 \times Z_2^T$ $C_{2h}^7 \times Z_2^T$ $C_{2h}^8 \times Z_2^T$	$C_{2x}, \mathcal{M}_y, PT$ C_{2x}, C_{2y}, PT C_{2z}, P, T C_{2x}, P, T C_{2y}, P, T C'_{21}, P, T C'_{22}, P, T C'_{23}, P, T C''_{21}, P, T C_2, P, T	$(\frac{\omega_2(\mathcal{M}_y, C_{2x})}{\omega_2(C_{2x}, \mathcal{M}_y)}, \chi_{PT}, \chi_{TM_x}, \chi_{TC_{2y}})$ $(\frac{\omega_2(C_{2y}, C_{2x})}{\omega_2(C_{2x}, C_{2y})}, \chi_{PT}, \chi_{TM_x}, \chi_{TM_y})$ $(\frac{\omega_2(P, C_{2z})}{\omega_2(C_{2z}, P)}, \chi_T, \chi_{TC_{2z}}, \chi_{PT})$ $(\frac{\omega_2(P, C_{2x})}{\omega_2(C_{2x}, P)}, \chi_T, \chi_{TC_{2x}}, \chi_{PT})$ $(\frac{\omega_2(P, C_{2y})}{\omega_2(C_{2y}, P)}, \chi_T, \chi_{TC_{2y}}, \chi_{PT})$ $(\frac{\omega_2(P, C'_{21})}{\omega_2(C'_{21}, P)}, \chi_T, \chi_{TC'_{21}}, \chi_{PT})$ $(\frac{\omega_2(P, C'_{22})}{\omega_2(C'_{22}, P)}, \chi_T, \chi_{TC'_{22}}, \chi_{PT})$ $(\frac{\omega_2(P, C'_{23})}{\omega_2(C'_{23}, P)}, \chi_T, \chi_{TC'_{23}}, \chi_{PT})$ $(\frac{\omega_2(P, C''_{21})}{\omega_2(C''_{21}, P)}, \chi_T, \chi_{TC''_{21}}, \chi_{PT})$ $(\frac{\omega_2(P, C_2)}{\omega_2(C_2, P)}, \chi_T, \chi_{TC_2}, \chi_{PT})$
$(Z_2 \times Z_2 \times Z_2) \times Z_2^T$	$D_{2h}^1 \times Z_2^T$ $D_{2h}^2 \times Z_2^T$ $D_{2h}^3 \times Z_2^T$	C_{2x}, C_{2y}, P, T C_{2b}, C_{2a}, P, T C'_{21}, C''_{21}, P, T	$(\frac{\omega_2(C_{2x}, C_{2y})}{\omega_2(C_{2y}, C_{2x})}, \chi_T, \chi_{PT}, \chi_{TC_{2x}}, \chi_{TC_{2y}}, \frac{\omega_2(C_{2x}, P)}{\omega_2(P, C_{2x})}, \frac{\omega_2(C_{2y}, P)}{\omega_2(P, C_{2y})})$ $(\frac{\omega_2(C_{2b}, C_{2a})}{\omega_2(C_{2a}, C_{2b})}, \chi_T, \chi_{PT}, \chi_{TC_{2b}}, \chi_{TC_{2a}}, \frac{\omega_2(C_{2b}, P)}{\omega_2(P, C_{2b})}, \frac{\omega_2(C_{2a}, P)}{\omega_2(P, C_{2a})})$ $(\frac{\omega_2(C'_{21}, C''_{21})}{\omega_2(C''_{21}, C'_{21})}, \chi_T, \chi_{PT}, \chi_{TC'_{21}}, \chi_{TC''_{21}}, \frac{\omega_2(C'_{21}, P)}{\omega_2(P, C'_{21})}, \frac{\omega_2(C''_{21}, P)}{\omega_2(P, C''_{21})})$
$Z_4 \times Z_2^T$	$S_4 \times Z_2^T$	S_{4z}^-, PT	(χ_{PT}, χ_{TM_z})
$(Z_4 \rtimes Z_2) \times Z_2^T$	$D_{2d}^1 \times Z_2^T$ $D_{2d}^2 \times Z_2^T$	S_{4z}^+, C_{2x}, PT $S_{4x}^+, \mathcal{M}_y, PT$	$(\frac{\omega_2(C_{2y}, C_{2x})}{\omega_2(C_{2x}, C_{2y})}, \chi_{PT}, \chi_{TM_x}, \chi_{TC_{2a}})$ $(\frac{\omega_2(\mathcal{M}_y, \mathcal{M}_z)}{\omega_2(\mathcal{M}_z, \mathcal{M}_y)}, \chi_{PT}, \chi_{TC_{2y}}, \chi_{TM_{df}})$
$(Z_4 \times Z_2) \times Z_2^T$	$C_{4h} \times Z_2^T$	C_{4z}^+, P, T	$(\frac{\omega_2(C_{4z}^+, P)}{\omega_2(P, C_{4z}^+)}, \chi_T, \chi_{PT}, \chi_{TM_z})$
$[(Z_4 \rtimes Z_2) \times Z_2] \times Z_2^T$	$D_{4h}^1 \times Z_2^T$ $D_{4h}^2 \times Z_2^T$	C_{4z}^+, C_{2x}, P, T C_{4y}^+, C_{2z}, P, T	$(\frac{\omega_2(C_{2x}, C_{2y})}{\omega_2(C_{2y}, C_{2x})}, \chi_T, \chi_{PT}, \chi_{TC_{2x}}, \chi_{TC_{2a}}, \frac{\omega_2(C_{2x}, P)}{\omega_2(P, C_{2x})}, \frac{\omega_2(C_{2a}, P)}{\omega_2(P, C_{2a})})$ $(\frac{\omega_2(C_{2z}, C_{2x})}{\omega_2(C_{2x}, C_{2z})}, \chi_T, \chi_{PT}, \chi_{TC_{2z}}, \chi_{TC_{2c}}, \frac{\omega_2(C_{2z}, P)}{\omega_2(P, C_{2z})}, \frac{\omega_2(C_{2c}, P)}{\omega_2(P, C_{2c})})$
$Z_3 \times Z_2^T$	$C_3 \times Z_2^T$	C_3^+, PT	χ_{PT}
$Z_6 \times Z_2^T$	$C_{3i}^1 \times Z_2^T$ $C_{3i}^2 \times Z_2^T$ $C_{3h} \times Z_2^T$	S_6^+, T S_{61}^+, T S_3^-, PT	(χ_T, χ_{PT}) (χ_T, χ_{PT}) (χ_{PT}, χ_{TC_2})
$(Z_3 \rtimes Z_2) \times Z_2^T$	$C_{3v} \times Z_2^T$ $D_3 \times Z_2^T$	$C_3^+, \mathcal{M}_{d1}, PT$ C_3^+, C''_{21}, PT	$(\chi_{PT}, \chi_{TC'_{21}})$ $(\chi_{PT}, \chi_{TM_{v1}})$
$(Z_6 \rtimes Z_2) \times Z_2^T$	$D_{3h} \times Z_2^T$ $D_{3d}^1 \times Z_2^T$ $D_{3d}^2 \times Z_2^T$ $D_{3d}^3 \times Z_2^T$	$S_3^-, \mathcal{M}_{d1}, PT$ S_6^+, C'_{21}, T S_6^+, C''_{21}, T S_{61}^+, C_{2b}, T	$(\frac{\omega_2(\mathcal{M}_h, \mathcal{M}_{d1})}{\omega_2(\mathcal{M}_{d1}, \mathcal{M}_h)}, \chi_{PT}, \chi_{TC_2}, \chi_{TC'_{21}})$ $(\frac{\omega_2(P, C'_{21})}{\omega_2(C'_{21}, P)}, \chi_T, \chi_{PT}, \chi_{TC'_{21}})$ $(\frac{\omega_2(P, C''_{21})}{\omega_2(C''_{21}, P)}, \chi_T, \chi_{PT}, \chi_{TC''_{21}})$ $(\frac{\omega_2(P, C_{2b})}{\omega_2(C_{2b}, P)}, \chi_T, \chi_{PT}, \chi_{TC_{2b}})$
$(Z_6 \times Z_2) \times Z_2^T$	$C_{6h} \times Z_2^T$	C_6^+, P, T	$(\frac{\omega_2(C_6^+, P)}{\omega_2(P, C_6^+)}, \chi_T, \chi_{PT}, \chi_{TM_h})$
$[(Z_6 \rtimes Z_2) \times Z_2] \times Z_2^T$	$D_{6h} \times Z_2^T$	C_6^+, C'_{21}, P, T	$(\frac{\omega_2(C'_{21}, C''_{21})}{\omega_2(C''_{21}, C'_{21})}, \chi_T, \chi_{PT}, \chi_{TC'_{21}}, \chi_{TC''_{21}}, \frac{\omega_2(C'_{21}, P)}{\omega_2(P, C'_{21})}, \frac{\omega_2(C''_{21}, P)}{\omega_2(P, C''_{21})})$
$(A_4 \times Z_2) \times Z_2^T$	$T_h \times Z_2^T$	$C_{31}^+, C_{2x}, C_{2y}, P, T$	$(\frac{\omega_2(C_{2x}, C_{2y})}{\omega_2(C_{2y}, C_{2x})}, \chi_T, \chi_{PT})$
$[(A_4 \rtimes Z_2) \times Z_2] \times Z_2^T$	$O_h \times Z_2^T$	$C_{31}^+, C_{2x}, C_{2y}, C_{2a}, P, T$	$(\frac{\omega_2(C_{2x}, C_{2y})}{\omega_2(C_{2y}, C_{2x})}, \chi_T, \chi_{PT}, \chi_{TC_{2a}}, \frac{\omega_2(C_{2a}, P)}{\omega_2(P, C_{2a})})$

-
- [1] T. D. Lee, R. Oehme, and C. N. Yang, Phys. Rev. **106**, 340 (1957), URL <https://link.aps.org/doi/10.1103/PhysRev.106.340>.
- [2] R. Streater and A. Wightman, *PCT, spin and statistics, and all that* (Princeton University Press, 1989), ISBN 978-0-691-07062-9.
- [3] S. Weinberg, *The Quantum theory of fields. Vol. 1: Foundations* (Cambridge University Press, 2005), ISBN 978-0-521-67053-1, 978-0-511-25204-4.
- [4] M. G. Vergniory, L. Elcoro, Z. Wang, J. Cano, C. Felser, M. I. Aroyo, B. A. Bernevig, and B. Bradlyn, Phys. Rev. E **96**, 023310 (2017), URL <https://link.aps.org/doi/10.1103/PhysRevE.96.023310>.
- [5] C. Bradley and A. Cracknell, *The Mathematical Theory of Symmetry in Solids: Representation Theory for Point Groups and Space Groups*, EBSCO ebook academic collection (OUP Oxford, 2010), ISBN 9780199582587, URL <https://books.google.com/books?id=mNQDAAAQBAJ>.
- [6] B. Bradlyn, J. Cano, Z. Wang, M. G. Vergniory, C. Felser, R. J. Cava, and B. A. Bernevig, Science **353** (2016), ISSN 0036-8075, <https://science.sciencemag.org/content/353/6299/aaf5037.full.pdf>, URL <https://science.sciencemag.org/content/353/6299/aaf5037>.
- [7] R. S. K. Mong, A. M. Essin, and J. E. Moore, Phys. Rev. B **81**, 245209 (2010), URL <https://link.aps.org/doi/10.1103/PhysRevB.81.245209>.
- [8] C. Fang, M. J. Gilbert, and B. A. Bernevig, Phys. Rev. B **88**, 085406 (2013), URL <https://link.aps.org/doi/10.1103/PhysRevB.88.085406>.
- [9] C. Fang, M. J. Gilbert, and B. A. Bernevig, Phys. Rev. Lett. **112**, 106401 (2014), URL <https://link.aps.org/doi/10.1103/PhysRevLett.112.106401>.
- [10] H. Watanabe, H. C. Po, and A. Vishwanath, Science Advances **4** (2018), <https://advances.sciencemag.org/content/4/8/eaat8685.full.pdf>, URL <https://advances.sciencemag.org/content/4/8/eaat8685>.
- [11] G. Hua, S. Nie, Z. Song, R. Yu, G. Xu, and K. Yao, Physical Review B **98**, 201116 (2018).
- [12] J. Cano, B. Bradlyn, and M. G. Vergniory, APL Materials **7**, 101125 (2019), <https://doi.org/10.1063/1.5124314>, URL <https://doi.org/10.1063/1.5124314>.
- [13] Y. Xu, L. Elcoro, Z. Song, B. J. Wieder, M. G. Vergniory, N. Regnault, Y. Chen, C. Felser, and B. A. Bernevig, *High-throughput calculations of antiferromagnetic topological materials from magnetic topological quantum chemistry* (2020), 2003.00012.
- [14] J. Zou, Q. Xie, G. Xu, and Z. Song, National Science Review (2020), ISSN 2095-5138, nwaal69, <https://academic.oup.com/nsr/advance-article-pdf/doi/10.1093/nsr/nwaa169/33528562/nwaa169.pdf>, URL <https://doi.org/10.1093/nsr/nwaa169>.
- [15] C. J. BRADLEY and B. L. DAVIES, Rev. Mod. Phys. **40**, 359 (1968), URL <https://link.aps.org/doi/10.1103/RevModPhys.40.359>.
- [16] S. V. Gallego, J. M. Perez-Mato, L. Elcoro, E. S. Tasci, R. M. Hanson, K. Momma, M. I. Aroyo, and G. Madariaga, Journal of Applied Crystallography **49**, 1750 (2016), URL <https://doi.org/10.1107/S1600576716012863>.
- [17] P. Anderson, *Concepts in Solids: Lectures on the Theory of Solids*, Advanced book classics series (World Scientific, 1997), ISBN 9789810232313, URL <https://books.google.com/books?id=2NEKzzbMbRgC>.
- [18] J. H. E. GRIFFITHS, Nature **158**, 670 (1946), URL <https://doi.org/10.1038/158670a0>.
- [19] A. Hatcher, *Algebraic Topology* (Cambridge University Press, 2005), ISBN 9780521541862, URL <https://books.google.com/books?id=cFgAPAAACAAJ>.
- [20] J.-Q. Chen, M.-J. Gao, and G.-Q. Ma, Rev. Mod. Phys. **57**, 211 (1985), URL <https://link.aps.org/doi/10.1103/RevModPhys.57.211>.
- [21] J. Yang and Z.-X. Liu, J. Phys. A: Math. Theor. **51**, 025207 (2018), URL <https://iopscience.iop.org/article/10.1088/1751-8121/aa971a>.
- [22] J. Chen, J. Ping, and F. Wang, *Group Representation Theory for Physicists* (World Scientific, 2002), ISBN 9789812380654, URL <https://books.google.com/books?id=yWhIDQAAQBAJ>.

TABLE IV: All invariants of magnetic point groups $H \times Z_2^T$ and symmetry points in Brillouin zone to realize them in type IV Shubnikov space groups with $P =$ space inversion, $T =$ time reversal symmetries.

Shubnikov space group	symmetry point	G_Q	invariants (electron)	invariants (Magnon)
2.7 IV	Γ	$C_i \times Z_2^T$	(-1, -1)	(+1, +1)
	B	$C_i \times Z_2^T$	(+1, -1)	(-1, +1)
	F	$C_i \times Z_2^T$	(-1, -1)	(+1, +1)
	G	$C_i \times Z_2^T$	(-1, -1)	(+1, +1)
10.47 IV	Γ	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	B	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	Y	$C_{2h}^1 \times Z_2^T$	(+1, +1, +1, -1)	(+1, -1, +1, +1)
	Z	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	C	$C_{2h}^1 \times Z_2^T$	(+1, +1, +1, -1)	(+1, -1, +1, +1)
	D	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	A	$C_{2h}^1 \times Z_2^T$	(+1, +1, +1, -1)	(+1, -1, +1, +1)
10.48 IV	Γ	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	B	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	Y	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	Z	$C_{2h}^1 \times Z_2^T$	(+1, +1, -1, -1)	(+1, -1, -1, +1)
	C	$C_{2h}^1 \times Z_2^T$	(+1, +1, -1, -1)	(+1, -1, -1, +1)
	D	$C_{2h}^1 \times Z_2^T$	(+1, +1, -1, -1)	(+1, -1, -1, +1)
	A	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
10.49 IV	Γ	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	B	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	Y	$C_{2h}^1 \times Z_2^T$	(+1, +1, +1, -1)	(+1, -1, +1, +1)
	Z	$C_{2h}^1 \times Z_2^T$	(+1, +1, -1, -1)	(+1, -1, -1, +1)
	C	$C_{2h}^1 \times Z_2^T$	(+1, -1, -1, -1)	(+1, +1, -1, +1)
	D	$C_{2h}^1 \times Z_2^T$	(+1, +1, -1, -1)	(+1, -1, -1, +1)
	A	$C_{2h}^1 \times Z_2^T$	(+1, +1, +1, -1)	(+1, -1, +1, +1)
11.55 IV	Γ	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	B	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	Y	$C_{2h}^1 \times Z_2^T$	(+1, +1, +1, -1)	(+1, -1, +1, +1)
	Z	$C_{2h}^1 \times Z_2^T$	(-1, -1, -1, -1)	(-1, +1, -1, +1)
	C	$C_{2h}^1 \times Z_2^T$	(-1, +1, -1, -1)	(-1, -1, -1, +1)
	D	$C_{2h}^1 \times Z_2^T$	(-1, -1, -1, -1)	(-1, +1, -1, +1)
	A	$C_{2h}^1 \times Z_2^T$	(+1, +1, +1, -1)	(+1, -1, +1, +1)
11.56 IV	Γ	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	B	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	Y	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	Z	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	C	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	D	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	A	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
11.57 IV	Γ	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	B	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	Y	$C_{2h}^1 \times Z_2^T$	(+1, +1, +1, -1)	(+1, -1, +1, +1)
	Z	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	C	$C_{2h}^1 \times Z_2^T$	(-1, -1, +1, -1)	(-1, +1, +1, +1)
	D	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	A	$C_{2h}^1 \times Z_2^T$	(+1, +1, +1, -1)	(+1, -1, +1, +1)
12.63 IV	Γ	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	A	$C_{2h}^1 \times Z_2^T$	(+1, +1, +1, -1)	(+1, -1, +1, +1)
	Z	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	M	$C_{2h}^1 \times Z_2^T$	(+1, +1, +1, -1)	(+1, -1, +1, +1)
	V	$C_i \times Z_2^T$	(+1, -1)	(-1, +1)

(Continued from the previous page)

Shubnikov space group	symmetry point	G_Q	invariants (electron)	invariants (Magnon)
12.64 IV	Γ	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	A	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	Z	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	M	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	L	$C_i \times Z_2^T$	(+1, -1)	(-1, +1)
	V	$C_i \times Z_2^T$	(+1, -1)	(-1, +1)
13.70 IV	Γ	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	B	$C_{2h}^1 \times Z_2^T$	(-1, -1, +1, -1)	(-1, +1, +1, +1)
	Y	$C_{2h}^1 \times Z_2^T$	(+1, +1, +1, -1)	(+1, -1, +1, +1)
	Z	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	C	$C_{2h}^1 \times Z_2^T$	(+1, +1, +1, -1)	(+1, -1, +1, +1)
	D	$C_{2h}^1 \times Z_2^T$	(-1, -1, +1, -1)	(-1, +1, +1, +1)
	A	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
E	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)	
13.71 IV	Γ	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	B	$C_{2h}^1 \times Z_2^T$	(-1, -1, +1, -1)	(-1, +1, +1, +1)
	Y	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	Z	$C_{2h}^1 \times Z_2^T$	(+1, +1, -1, -1)	(+1, -1, -1, +1)
	C	$C_{2h}^1 \times Z_2^T$	(+1, +1, -1, -1)	(+1, -1, -1, +1)
	D	$C_{2h}^1 \times Z_2^T$	(-1, +1, -1, -1)	(-1, -1, -1, +1)
	A	$C_{2h}^1 \times Z_2^T$	(-1, -1, +1, -1)	(-1, +1, +1, +1)
E	$C_{2h}^1 \times Z_2^T$	(-1, +1, -1, -1)	(-1, -1, -1, +1)	
13.72 IV	Γ	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	B	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	Y	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	Z	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	C	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	D	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	A	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
E	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)	
13.73 IV	Γ	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	B	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	Y	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	Z	$C_{2h}^1 \times Z_2^T$	(+1, +1, -1, -1)	(+1, -1, -1, +1)
	C	$C_{2h}^1 \times Z_2^T$	(+1, +1, -1, -1)	(+1, -1, -1, +1)
	D	$C_{2h}^1 \times Z_2^T$	(-1, -1, -1, -1)	(-1, +1, -1, +1)
	A	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
E	$C_{2h}^1 \times Z_2^T$	(-1, -1, -1, -1)	(-1, +1, -1, +1)	
13.74 IV	Γ	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	B	$C_{2h}^1 \times Z_2^T$	(-1, -1, +1, -1)	(-1, +1, +1, +1)
	Y	$C_{2h}^1 \times Z_2^T$	(+1, +1, +1, -1)	(+1, -1, +1, +1)
	Z	$C_{2h}^1 \times Z_2^T$	(+1, +1, -1, -1)	(+1, -1, -1, +1)
	C	$C_{2h}^1 \times Z_2^T$	(+1, -1, -1, -1)	(+1, +1, -1, +1)
	D	$C_{2h}^1 \times Z_2^T$	(-1, +1, -1, -1)	(-1, -1, -1, +1)
	A	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
E	$C_{2h}^1 \times Z_2^T$	(-1, -1, -1, -1)	(-1, +1, -1, +1)	
14.80 IV	Γ	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	B	$C_{2h}^1 \times Z_2^T$	(-1, -1, +1, -1)	(-1, +1, +1, +1)
	Y	$C_{2h}^1 \times Z_2^T$	(+1, +1, +1, -1)	(+1, -1, +1, +1)
	Z	$C_{2h}^1 \times Z_2^T$	(-1, -1, -1, -1)	(-1, +1, -1, +1)
	C	$C_{2h}^1 \times Z_2^T$	(-1, +1, -1, -1)	(-1, -1, -1, +1)
	D	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	A	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
E	$C_{2h}^1 \times Z_2^T$	(+1, +1, -1, -1)	(+1, -1, -1, +1)	
14.81 IV	Γ	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	B	$C_{2h}^1 \times Z_2^T$	(-1, -1, +1, -1)	(-1, +1, +1, +1)
	Y	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	Z	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	C	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	D	$C_{2h}^1 \times Z_2^T$	(+1, +1, +1, -1)	(+1, -1, +1, +1)
	A	$C_{2h}^1 \times Z_2^T$	(-1, -1, +1, -1)	(-1, +1, +1, +1)
E	$C_{2h}^1 \times Z_2^T$	(+1, +1, +1, -1)	(+1, -1, +1, +1)	

(Continued from the previous page)

Shubnikov space group	symmetry point	$G_{\mathbf{Q}}$	invariants (electron)	invariants (Magnon)
14.82 IV	Γ	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	B	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	Y	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	Z	$C_{2h}^1 \times Z_2^T$	(-1, -1, -1, -1)	(-1, +1, -1, +1)
	C	$C_{2h}^1 \times Z_2^T$	(-1, -1, -1, -1)	(-1, +1, -1, +1)
	D	$C_{2h}^1 \times Z_2^T$	(+1, +1, -1, -1)	(+1, -1, -1, +1)
	A	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	E	$C_{2h}^1 \times Z_2^T$	(+1, +1, -1, -1)	(+1, -1, -1, +1)
14.83 IV	Γ	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	B	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	Y	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	Z	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	C	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	D	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	A	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	E	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
14.84 IV	Γ	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	B	$C_{2h}^1 \times Z_2^T$	(-1, -1, +1, -1)	(-1, +1, +1, +1)
	Y	$C_{2h}^1 \times Z_2^T$	(+1, +1, +1, -1)	(+1, -1, +1, +1)
	Z	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	C	$C_{2h}^1 \times Z_2^T$	(-1, -1, +1, -1)	(-1, +1, +1, +1)
	D	$C_{2h}^1 \times Z_2^T$	(+1, +1, +1, -1)	(+1, -1, +1, +1)
	A	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	E	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
15.90 IV	Γ	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	A	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	Z	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	M	$C_{2h}^1 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	L	$C_i \times Z_2^T$	(+1, -1)	(-1, +1)
	V	$C_i \times Z_2^T$	(-1, -1)	(+1, +1)
15.91 IV	Γ	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	A	$C_{2h}^1 \times Z_2^T$	(-1, -1, +1, -1)	(-1, +1, +1, +1)
	Z	$C_{2h}^1 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	M	$C_{2h}^1 \times Z_2^T$	(-1, -1, +1, -1)	(-1, +1, +1, +1)
	L	$C_i \times Z_2^T$	(+1, -1)	(-1, +1)
	V	$C_i \times Z_2^T$	(+1, -1)	(-1, +1)
47.254 IV	Γ	$D_{2h}^1 \times Z_2^T$	(-1, -1, -1, +1, +1, +1, +1)	(+1, +1, +1, +1, +1, +1, +1)
	Y	$D_{2h}^1 \times Z_2^T$	(-1, -1, -1, +1, +1, +1, +1)	(+1, +1, +1, +1, +1, +1, +1)
	X	$D_{2h}^1 \times Z_2^T$	(-1, +1, -1, -1, +1, +1, +1)	(+1, -1, +1, -1, +1, +1, +1)
	Z	$D_{2h}^1 \times Z_2^T$	(-1, -1, -1, +1, +1, +1, +1)	(+1, +1, +1, +1, +1, +1, +1)
	U	$D_{2h}^1 \times Z_2^T$	(-1, +1, -1, -1, +1, +1, +1)	(+1, -1, +1, -1, +1, +1, +1)
	T	$D_{2h}^1 \times Z_2^T$	(-1, -1, -1, +1, +1, +1, +1)	(+1, +1, +1, +1, +1, +1, +1)
	S	$D_{2h}^1 \times Z_2^T$	(-1, +1, -1, -1, +1, +1, +1)	(+1, -1, +1, -1, +1, +1, +1)
	R	$D_{2h}^1 \times Z_2^T$	(-1, +1, -1, -1, +1, +1, +1)	(+1, -1, +1, -1, +1, +1, +1)
47.255 IV	Γ	$D_{2h}^1 \times Z_2^T$	(-1, -1, -1, +1, +1, +1, +1)	(+1, +1, +1, +1, +1, +1, +1)
	Y	$D_{2h}^1 \times Z_2^T$	(-1, +1, -1, +1, -1, +1, +1)	(+1, -1, +1, +1, -1, +1, +1)
	X	$D_{2h}^1 \times Z_2^T$	(-1, +1, -1, -1, +1, +1, +1)	(+1, -1, +1, -1, +1, +1, +1)
	Z	$D_{2h}^1 \times Z_2^T$	(-1, -1, -1, +1, +1, +1, +1)	(+1, +1, +1, +1, +1, +1, +1)
	U	$D_{2h}^1 \times Z_2^T$	(-1, +1, -1, -1, +1, +1, +1)	(+1, -1, +1, -1, +1, +1, +1)
	T	$D_{2h}^1 \times Z_2^T$	(-1, +1, -1, +1, -1, +1, +1)	(+1, -1, +1, +1, -1, +1, +1)
	S	$D_{2h}^1 \times Z_2^T$	(-1, -1, -1, -1, -1, +1, +1)	(+1, +1, +1, -1, -1, +1, +1)
	R	$D_{2h}^1 \times Z_2^T$	(-1, -1, -1, -1, -1, +1, +1)	(+1, +1, +1, -1, -1, +1, +1)
47.256 IV	Γ	$D_{2h}^1 \times Z_2^T$	(-1, -1, -1, +1, +1, +1, +1)	(+1, +1, +1, +1, +1, +1, +1)
	Y	$D_{2h}^1 \times Z_2^T$	(-1, +1, -1, +1, -1, +1, +1)	(+1, -1, +1, +1, -1, +1, +1)
	X	$D_{2h}^1 \times Z_2^T$	(-1, +1, -1, -1, +1, +1, +1)	(+1, -1, +1, -1, +1, +1, +1)
	Z	$D_{2h}^1 \times Z_2^T$	(-1, +1, -1, +1, +1, +1, +1)	(+1, -1, +1, +1, +1, +1, +1)
	U	$D_{2h}^1 \times Z_2^T$	(-1, -1, -1, -1, +1, +1, +1)	(+1, +1, +1, -1, +1, +1, +1)
	T	$D_{2h}^1 \times Z_2^T$	(-1, -1, -1, +1, -1, +1, +1)	(+1, +1, +1, +1, -1, +1, +1)
	S	$D_{2h}^1 \times Z_2^T$	(-1, -1, -1, -1, -1, +1, +1)	(+1, +1, +1, -1, -1, +1, +1)
	R	$D_{2h}^1 \times Z_2^T$	(-1, +1, -1, -1, -1, +1, +1)	(+1, -1, +1, -1, -1, +1, +1)

(Continued from the previous page)

Shubnikov space group	symmetry point	$G_{\mathbf{Q}}$	invariants (electron)	invariants (Magnon)
74.562 IV	Γ	$D_{2h}^1 \times Z_2^T$	$(-1, -1, -1, +1, +1, +1, +1)$	$(+1, +1, +1, +1, +1, +1, +1)$
	X	$D_{2h}^1 \times Z_2^T$	$(-1, -1, -1, +1, +1, +1, +1)$	$(+1, +1, +1, +1, +1, +1, +1)$
	R	$C_{2h}^3 \times Z_2^T$	$(+1, +1, +1, -1)$	$(+1, -1, +1, +1)$
	S	$C_{2h}^2 \times Z_2^T$	$(+1, -1, +1, -1)$	$(+1, +1, +1, +1)$
	T	$C_{2h}^1 \times Z_2^T$	$(-1, +1, +1, -1)$	$(-1, -1, +1, +1)$
	W	$D_2 \times Z_2^T$	$(+1, -1, +1, -1)$	$(-1, +1, +1, -1)$
83.48 IV	Γ	$C_{4h} \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	M	$C_{4h} \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	Z	$C_{4h} \times Z_2^T$	$(+1, +1, -1, +1)$	$(+1, -1, +1, +1)$
	A	$C_{4h} \times Z_2^T$	$(+1, +1, -1, +1)$	$(+1, -1, +1, +1)$
	R	$C_{2h}^1 \times Z_2^T$	$(+1, +1, -1, -1)$	$(+1, -1, -1, +1)$
	X	$C_{2h}^1 \times Z_2^T$	$(+1, -1, +1, -1)$	$(+1, +1, +1, +1)$
83.49 IV	Γ	$C_{4h} \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	M	$C_{4h} \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	Z	$C_{4h} \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	A	$C_{4h} \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	R	$C_{2h}^1 \times Z_2^T$	$(+1, +1, +1, -1)$	$(+1, -1, +1, +1)$
	X	$C_{2h}^1 \times Z_2^T$	$(+1, +1, +1, -1)$	$(+1, -1, +1, +1)$
83.50 IV	Γ	$C_{4h} \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	M	$C_{4h} \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	Z	$C_{4h} \times Z_2^T$	$(+1, +1, -1, +1)$	$(+1, -1, +1, +1)$
	A	$C_{4h} \times Z_2^T$	$(+1, +1, -1, +1)$	$(+1, -1, +1, +1)$
	R	$C_{2h}^1 \times Z_2^T$	$(+1, -1, -1, -1)$	$(+1, +1, -1, +1)$
	X	$C_{2h}^1 \times Z_2^T$	$(+1, +1, +1, -1)$	$(+1, -1, +1, +1)$
84.56 IV	Γ	$C_{4h} \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	M	$C_{4h} \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	Z	$C_{4h} \times Z_2^T$	$(-1, +1, -1, +1)$	$(-1, -1, +1, +1)$
	A	$C_{4h} \times Z_2^T$	$(-1, +1, -1, +1)$	$(-1, -1, +1, +1)$
	R	$C_{2h}^1 \times Z_2^T$	$(+1, +1, -1, -1)$	$(+1, -1, -1, +1)$
	X	$C_{2h}^1 \times Z_2^T$	$(+1, -1, +1, -1)$	$(+1, +1, +1, +1)$
84.57 IV	Γ	$C_{4h} \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	M	$C_{4h} \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	Z	$C_{4h} \times Z_2^T$	$(-1, -1, -1, +1)$	$(-1, +1, +1, +1)$
	A	$C_{4h} \times Z_2^T$	$(-1, -1, -1, +1)$	$(-1, +1, +1, +1)$
	R	$C_{2h}^1 \times Z_2^T$	$(+1, +1, +1, -1)$	$(+1, -1, +1, +1)$
	X	$C_{2h}^1 \times Z_2^T$	$(+1, +1, +1, -1)$	$(+1, -1, +1, +1)$
84.58 IV	Γ	$C_{4h} \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	M	$C_{4h} \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	Z	$C_{4h} \times Z_2^T$	$(-1, +1, -1, +1)$	$(-1, -1, +1, +1)$
	A	$C_{4h} \times Z_2^T$	$(-1, +1, -1, +1)$	$(-1, -1, +1, +1)$
	R	$C_{2h}^1 \times Z_2^T$	$(+1, -1, -1, -1)$	$(+1, +1, -1, +1)$
	X	$C_{2h}^1 \times Z_2^T$	$(+1, +1, +1, -1)$	$(+1, -1, +1, +1)$
85.64 IV	Γ	$C_{4h} \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	M	$C_{4h} \times Z_2^T$	$(-1, -1, -1, +1)$	$(-1, +1, +1, +1)$
	Z	$C_{4h} \times Z_2^T$	$(+1, +1, -1, +1)$	$(+1, -1, +1, +1)$
	A	$C_{4h} \times Z_2^T$	$(-1, +1, -1, +1)$	$(-1, -1, +1, +1)$
	R	$C_{2h}^1 \times Z_2^T$	$(-1, +1, -1, -1)$	$(-1, -1, -1, +1)$
	X	$C_{2h}^1 \times Z_2^T$	$(-1, -1, +1, -1)$	$(-1, +1, +1, +1)$
85.65 IV	Γ	$C_{4h} \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	M	$C_{4h} \times Z_2^T$	$(-1, -1, -1, +1)$	$(-1, +1, +1, +1)$
	Z	$C_{4h} \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	A	$C_{4h} \times Z_2^T$	$(-1, -1, -1, +1)$	$(-1, +1, +1, +1)$
	R	$C_{2h}^1 \times Z_2^T$	$(-1, +1, +1, -1)$	$(-1, -1, +1, +1)$
	X	$C_{2h}^1 \times Z_2^T$	$(-1, +1, +1, -1)$	$(-1, -1, +1, +1)$
85.66 IV	Γ	$C_{4h} \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	M	$C_{4h} \times Z_2^T$	$(-1, -1, -1, +1)$	$(-1, +1, +1, +1)$
	Z	$C_{4h} \times Z_2^T$	$(+1, +1, -1, +1)$	$(+1, -1, +1, +1)$
	A	$C_{4h} \times Z_2^T$	$(-1, +1, -1, +1)$	$(-1, -1, +1, +1)$
	R	$C_{2h}^1 \times Z_2^T$	$(-1, -1, -1, -1)$	$(-1, +1, -1, +1)$
	X	$C_{2h}^1 \times Z_2^T$	$(-1, +1, +1, -1)$	$(-1, -1, +1, +1)$

(Continued from the previous page)

Shubnikov space group	symmetry point	$G_{\mathbf{Q}}$	invariants (electron)	invariants (Magnon)
138.529 IV	Γ	$D_{4h}^1 \times Z_2^T$	$(-1, -1, -1, +1, +1, +1, +1)$	$(+1, +1, +1, +1, +1, +1, +1)$
	M	$D_{4h}^1 \times Z_2^T$	$(-1, -1, -1, +1, +1, -1, +1)$	$(+1, +1, +1, +1, +1, -1, +1)$
	Z	$D_{4h}^1 \times Z_2^T$	$(-1, -1, -1, +1, +1, -1, +1)$	$(+1, +1, +1, +1, +1, -1, +1)$
	A	$D_{4h}^1 \times Z_2^T$	$(-1, -1, -1, +1, +1, +1, +1)$	$(+1, +1, +1, +1, +1, +1, +1)$
	R	$D_{2h}^1 \times Z_2^T$	$(+1, +1, -1, +1, +1, -1, +1)$	$(-1, -1, +1, +1, +1, -1, +1)$
	X	$D_{2h}^1 \times Z_2^T$	$(+1, +1, -1, +1, +1, +1, -1)$	$(-1, -1, +1, +1, +1, +1, -1)$
138.530 IV	Γ	$D_{4h}^1 \times Z_2^T$	$(-1, -1, -1, +1, +1, +1, +1)$	$(+1, +1, +1, +1, +1, +1, +1)$
	M	$D_{4h}^1 \times Z_2^T$	$(-1, -1, -1, +1, +1, -1, +1)$	$(+1, +1, +1, +1, +1, -1, +1)$
	Z	$D_{4h}^1 \times Z_2^T$	$(-1, +1, -1, +1, +1, -1, +1)$	$(+1, -1, +1, +1, +1, -1, +1)$
	A	$D_{4h}^1 \times Z_2^T$	$(-1, +1, -1, +1, +1, +1, +1)$	$(+1, -1, +1, +1, +1, +1, +1)$
	R	$D_{2h}^1 \times Z_2^T$	$(+1, -1, -1, +1, +1, -1, +1)$	$(-1, +1, +1, +1, +1, -1, +1)$
	X	$D_{2h}^1 \times Z_2^T$	$(+1, +1, -1, +1, +1, +1, -1)$	$(-1, -1, +1, +1, +1, +1, -1)$
139.540 IV	Γ	$D_{4h}^1 \times Z_2^T$	$(-1, -1, -1, +1, +1, +1, +1)$	$(+1, +1, +1, +1, +1, +1, +1)$
	N	$C_{2h}^3 \times Z_2^T$	$(+1, +1, +1, -1)$	$(+1, -1, +1, +1)$
	X	$D_{2h}^2 \times Z_2^T$	$(-1, -1, -1, +1, +1, +1, +1)$	$(+1, +1, +1, +1, +1, +1, +1)$
	Z	$D_{4h}^1 \times Z_2^T$	$(-1, -1, -1, +1, +1, +1, +1)$	$(+1, +1, +1, +1, +1, +1, +1)$
	P	$D_{2d}^1 \times Z_2^T$	$(-1, -1, -1, +1)$	$(+1, +1, -1, +1)$
	140.550 IV	Γ	$D_{4h}^1 \times Z_2^T$	$(-1, -1, -1, +1, +1, +1, +1)$
N		$C_{2h}^3 \times Z_2^T$	$(-1, +1, +1, -1)$	$(-1, -1, +1, +1)$
X		$D_{2h}^2 \times Z_2^T$	$(-1, -1, -1, +1, +1, +1, +1)$	$(+1, +1, +1, +1, +1, +1, +1)$
Z		$D_{4h}^1 \times Z_2^T$	$(-1, -1, -1, +1, +1, +1, +1)$	$(+1, +1, +1, +1, +1, +1, +1)$
P		$D_{2d}^1 \times Z_2^T$	$(-1, -1, +1, +1)$	$(+1, +1, +1, +1)$
141.560 IV		Γ	$D_{4h}^1 \times Z_2^T$	$(-1, -1, -1, +1, +1, +1, +1)$
	N	$C_{2h}^3 \times Z_2^T$	$(+1, +1, +1, -1)$	$(+1, -1, +1, +1)$
	X	$D_{2h}^2 \times Z_2^T$	$(-1, -1, -1, +1, +1, -1, +1)$	$(+1, +1, +1, +1, +1, -1, +1)$
	Z	$D_{4h}^1 \times Z_2^T$	$(-1, -1, -1, +1, +1, +1, -1)$	$(+1, +1, +1, +1, +1, +1, -1)$
	P	$D_{2d}^1 \times Z_2^T$	$(+1, -1, -1, +1)$	$(-1, +1, -1, +1)$
	142.570 IV	Γ	$D_{4h}^1 \times Z_2^T$	$(-1, -1, -1, +1, +1, +1, +1)$
N		$C_{2h}^3 \times Z_2^T$	$(-1, +1, +1, -1)$	$(-1, -1, +1, +1)$
X		$D_{2h}^2 \times Z_2^T$	$(-1, -1, -1, +1, +1, -1, +1)$	$(+1, +1, +1, +1, +1, -1, +1)$
Z		$D_{4h}^1 \times Z_2^T$	$(-1, -1, -1, +1, +1, +1, -1)$	$(+1, +1, +1, +1, +1, +1, -1)$
P		$D_{2d}^1 \times Z_2^T$	$(+1, -1, +1, +1)$	$(-1, +1, +1, +1)$
147.16 IV		Γ	$C_{3i}^1 \times Z_2^T$	$(-1, -1)$
	M	$C_i \times Z_2^T$	$(-1, -1)$	$(+1, +1)$
	A	$C_{3i}^1 \times Z_2^T$	$(+1, -1)$	$(-1, +1)$
	L	$C_i \times Z_2^T$	$(+1, -1)$	$(-1, +1)$
	K	$C_3 \times Z_2^T$	-1	$+1$
	H	$C_3 \times Z_2^T$	-1	$+1$
148.20 IV	Γ	$C_{3i}^1 \times Z_2^T$	$(-1, -1)$	$(+1, +1)$
	Z	$C_{3i}^1 \times Z_2^T$	$(+1, -1)$	$(-1, +1)$
	L	$C_i \times Z_2^T$	$(+1, -1)$	$(-1, +1)$
	F	$C_i \times Z_2^T$	$(-1, -1)$	$(+1, +1)$
162.78 IV	Γ	$D_{3d}^1 \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	M	$C_{2h}^4 \times Z_2^T$	$(+1, -1, +1, -1)$	$(+1, +1, +1, +1)$
	A	$D_{3d}^1 \times Z_2^T$	$(+1, +1, -1, +1)$	$(+1, -1, +1, +1)$
	L	$C_{2h}^4 \times Z_2^T$	$(+1, +1, +1, -1)$	$(+1, -1, +1, +1)$
	K	$C_{3v} \times Z_2^T$	$(-1, +1)$	$(+1, +1)$
	H	$C_{3v} \times Z_2^T$	$(-1, +1)$	$(+1, +1)$
163.84 IV	Γ	$D_{3d}^1 \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	M	$C_{2h}^4 \times Z_2^T$	$(+1, -1, +1, -1)$	$(+1, +1, +1, +1)$
	A	$D_{3d}^1 \times Z_2^T$	$(-1, +1, -1, +1)$	$(-1, -1, +1, +1)$
	L	$C_{2h}^4 \times Z_2^T$	$(-1, +1, +1, -1)$	$(-1, -1, +1, +1)$
	K	$C_{3v} \times Z_2^T$	$(-1, +1)$	$(+1, +1)$
	H	$C_{3v} \times Z_2^T$	$(-1, +1)$	$(+1, +1)$
164.90 IV	Γ	$D_{3d}^2 \times Z_2^T$	$(+1, -1, -1, +1)$	$(+1, +1, +1, +1)$
	M	$C_{2h}^7 \times Z_2^T$	$(+1, -1, +1, -1)$	$(+1, +1, +1, +1)$
	A	$D_{3d}^2 \times Z_2^T$	$(+1, +1, -1, +1)$	$(+1, -1, +1, +1)$
	L	$C_{2h}^7 \times Z_2^T$	$(+1, +1, +1, -1)$	$(+1, -1, +1, +1)$
	K	$D_3 \times Z_2^T$	$(-1, +1)$	$(+1, +1)$
	H	$D_3 \times Z_2^T$	$(-1, -1)$	$(+1, -1)$

(Continued from the previous page)

Shubnikov space group	symmetry point	$G_{\mathbf{Q}}$	invariants (electron)	invariants (Magnon)
165.96 IV	Γ	$D_{3d}^2 \times Z_2^T$	(+1, -1, -1, +1)	(+1, +1, +1, +1)
	M	$C_{2h}^7 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	A	$D_{3d}^2 \times Z_2^T$	(-1, +1, -1, +1)	(-1, -1, +1, +1)
	L	$C_{2h}^7 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	K	$D_3 \times Z_2^T$	(-1, +1)	(+1, +1)
	H	$D_3 \times Z_2^T$	(-1, +1)	(+1, +1)
166.102 IV	Γ	$D_{3d}^1 \times Z_2^T$	(+1, -1, -1, +1)	(+1, +1, +1, +1)
	Z	$D_{3d}^1 \times Z_2^T$	(+1, +1, -1, +1)	(+1, -1, +1, +1)
	L	$C_{2h}^5 \times Z_2^T$	(+1, +1, +1, -1)	(+1, -1, +1, +1)
	F	$C_{2h}^4(C_{2h}^6) \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
167.108 IV	Γ	$D_{3d}^1 \times Z_2^T$	(+1, -1, -1, +1)	(+1, +1, +1, +1)
	Z	$D_{3d}^1 \times Z_2^T$	(-1, +1, -1, +1)	(-1, -1, +1, +1)
	L	$C_{2h}^5 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	F	$C_{2h}^4(C_{2h}^6) \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
175.142 IV	Γ	$C_{6h} \times Z_2^T$	(+1, -1, -1, +1)	(+1, +1, +1, +1)
	M	$C_{2h}^8 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	A	$C_{6h} \times Z_2^T$	(+1, +1, -1, +1)	(+1, -1, +1, +1)
	L	$C_{2h}^8 \times Z_2^T$	(+1, +1, -1, -1)	(+1, -1, -1, +1)
	K	$C_{3h} \times Z_2^T$	(-1, +1)	(+1, +1)
	H	$C_{3h} \times Z_2^T$	(-1, -1)	(+1, -1)
176.148 IV	Γ	$C_{6h} \times Z_2^T$	(+1, -1, -1, +1)	(+1, +1, +1, +1)
	M	$C_{2h}^8 \times Z_2^T$	(+1, -1, +1, -1)	(+1, +1, +1, +1)
	A	$C_{6h} \times Z_2^T$	(-1, +1, -1, +1)	(-1, -1, +1, +1)
	L	$C_{2h}^8 \times Z_2^T$	(-1, +1, +1, -1)	(-1, -1, +1, +1)
	K	$C_{3h} \times Z_2^T$	(-1, +1)	(+1, +1)
	H	$C_{3h} \times Z_2^T$	(-1, +1)	(+1, +1)
191.242 IV	Γ	$D_{6h} \times Z_2^T$	(-1, -1, -1, +1, +1, +1)	(+1, +1, +1, +1, +1, +1)
	M	$D_{2h}^3 \times Z_2^T$	(-1, -1, -1, +1, +1, +1)	(+1, +1, +1, +1, +1, +1)
	A	$D_{6h} \times Z_2^T$	(-1, +1, -1, +1, +1, +1)	(+1, -1, +1, +1, +1, +1)
	L	$D_{2h}^3 \times Z_2^T$	(-1, +1, -1, +1, +1, +1)	(+1, -1, +1, +1, +1, +1)
	K	$D_{3h} \times Z_2^T$	(-1, -1, +1, +1)	(+1, +1, +1, +1)
	H	$D_{3h} \times Z_2^T$	(-1, -1, -1, +1)	(+1, +1, -1, +1)
192.252 IV	Γ	$D_{6h} \times Z_2^T$	(-1, -1, -1, +1, +1, +1)	(+1, +1, +1, +1, +1, +1)
	M	$D_{2h}^3 \times Z_2^T$	(-1, -1, -1, +1, +1, +1)	(+1, +1, +1, +1, +1, +1)
	A	$D_{6h} \times Z_2^T$	(-1, +1, -1, +1, +1, -1)	(+1, -1, +1, +1, +1, -1)
	L	$D_{2h}^3 \times Z_2^T$	(-1, +1, -1, +1, +1, -1)	(+1, -1, +1, +1, +1, -1)
	K	$D_{3h} \times Z_2^T$	(-1, -1, +1, +1)	(+1, +1, +1, +1)
	H	$D_{3h} \times Z_2^T$	(+1, -1, -1, +1)	(-1, +1, -1, +1)
193.262 IV	Γ	$D_{6h} \times Z_2^T$	(-1, -1, -1, +1, +1, +1)	(+1, +1, +1, +1, +1, +1)
	M	$D_{2h}^3 \times Z_2^T$	(-1, -1, -1, +1, +1, +1)	(+1, +1, +1, +1, +1, +1)
	A	$D_{6h} \times Z_2^T$	(+1, +1, -1, +1, +1, -1)	(-1, -1, +1, +1, +1, -1)
	L	$D_{2h}^3 \times Z_2^T$	(+1, +1, -1, +1, +1, -1)	(-1, -1, +1, +1, +1, -1)
	K	$D_{3h} \times Z_2^T$	(-1, -1, +1, +1)	(+1, +1, +1, +1)
	H	$D_{3h} \times Z_2^T$	(-1, -1, +1, +1)	(+1, +1, +1, +1)
194.272 IV	Γ	$D_{6h} \times Z_2^T$	(-1, -1, -1, +1, +1, +1)	(+1, +1, +1, +1, +1, +1)
	M	$D_{2h}^3 \times Z_2^T$	(-1, -1, -1, +1, +1, +1)	(+1, +1, +1, +1, +1, +1)
	A	$D_{6h} \times Z_2^T$	(+1, +1, -1, +1, +1, -1)	(-1, -1, +1, +1, +1, -1)
	L	$D_{2h}^3 \times Z_2^T$	(+1, +1, -1, +1, +1, -1)	(-1, -1, +1, +1, +1, -1)
	K	$D_{3h} \times Z_2^T$	(-1, -1, +1, +1)	(+1, +1, +1, +1)
	H	$D_{3h} \times Z_2^T$	(+1, -1, +1, +1)	(-1, +1, +1, +1)
200.17 IV	Γ	$T_h \times Z_2^T$	(-1, -1, -1)	(+1, +1, +1)
	X	$D_{2h}^1 \times Z_2^T$	(-1, +1, -1, +1, -1, +1)	(+1, -1, +1, +1, -1, +1)
	M	$D_{2h}^1 \times Z_2^T$	(-1, -1, -1, -1, -1, +1)	(+1, +1, +1, -1, -1, +1)
	R	$T_h \times Z_2^T$	(-1, +1, -1)	(+1, -1, +1)
201.21 IV	Γ	$T_h \times Z_2^T$	(-1, -1, -1)	(+1, +1, +1)
	X	$D_{2h}^1 \times Z_2^T$	(-1, +1, -1, +1, -1, -1)	(+1, -1, +1, +1, -1, -1)
	M	$D_{2h}^1 \times Z_2^T$	(-1, -1, -1, -1, -1, -1)	(+1, +1, +1, -1, -1, -1)
	R	$T_h \times Z_2^T$	(-1, +1, -1)	(+1, -1, +1)
202.25 IV	Γ	$T_h \times Z_2^T$	(-1, -1, -1)	(+1, +1, +1)
	X	$D_{2h}^1 \times Z_2^T$	(-1, -1, -1, +1, +1, +1)	(+1, +1, +1, +1, +1, +1)
	L	$C_{3i}^2 \times Z_2^T$	(+1, -1)	(-1, +1)
	W	$C_{2v} \times Z_2^T$	(-1, -1, +1, +1)	(+1, +1, +1, +1)

(Continued from the previous page)

Shubnikov space group	symmetry point	$G_{\mathbf{Q}}$	invariants (electron)	invariants (Magnon)
203.29 IV	Γ	$T_h \times Z_2^T$	$(-1, -1, -1)$	$(+1, +1, +1)$
	X	$D_{2h}^1 \times Z_2^T$	$(-1, -1, -1, +1, +1, -1, +1)$	$(+1, +1, +1, +1, +1, -1, +1)$
	L	$C_{3i}^2 \times Z_2^T$	$(+1, -1)$	$(-1, +1)$
	W	$C_{2v} \times Z_2^T$	$(+1, -1, -1, +1)$	$(-1, +1, -1, +1)$
205.36 IV	Γ	$T_h \times Z_2^T$	$(-1, -1, -1)$	$(+1, +1, +1)$
	X	$D_{2h}^1 \times Z_2^T$	$(+1, +1, -1, +1, +1, -1, -1)$	$(-1, -1, +1, +1, +1, -1, -1)$
	M	$D_{2h}^1 \times Z_2^T$	$(-1, -1, -1, +1, +1, +1, -1)$	$(+1, +1, +1, +1, +1, +1, -1)$
	R	$T_h \times Z_2^T$	$(+1, +1, -1)$	$(-1, -1, +1)$
221.97 IV	Γ	$O_h \times Z_2^T$	$(-1, -1, -1, +1, +1)$	$(+1, +1, +1, +1, +1)$
	X	$D_{4h}^2 \times Z_2^T$	$(-1, +1, -1, +1, +1, +1, +1)$	$(+1, -1, +1, +1, +1, +1, +1)$
	M	$D_{4h}^1 \times Z_2^T$	$(-1, -1, -1, -1, +1, +1, +1)$	$(+1, +1, +1, -1, +1, +1, +1)$
	R	$O_h \times Z_2^T$	$(-1, +1, -1, +1, +1)$	$(+1, -1, +1, +1, +1)$
222.103 IV	Γ	$O_h \times Z_2^T$	$(-1, -1, -1, +1, +1)$	$(+1, +1, +1, +1, +1)$
	X	$D_{4h}^2 \times Z_2^T$	$(-1, +1, -1, +1, +1, -1, -1)$	$(+1, -1, +1, +1, +1, -1, -1)$
	M	$D_{4h}^1 \times Z_2^T$	$(-1, -1, -1, -1, +1, -1, +1)$	$(+1, +1, +1, -1, +1, -1, +1)$
	R	$O_h \times Z_2^T$	$(-1, +1, -1, +1, -1)$	$(+1, -1, +1, +1, -1)$
223.109 IV	Γ	$O_h \times Z_2^T$	$(-1, -1, -1, +1, +1)$	$(+1, +1, +1, +1, +1)$
	X	$D_{4h}^2 \times Z_2^T$	$(-1, +1, -1, +1, +1, +1, -1)$	$(+1, -1, +1, +1, +1, +1, -1)$
	M	$D_{4h}^1 \times Z_2^T$	$(-1, -1, -1, -1, +1, +1, +1)$	$(+1, +1, +1, -1, +1, +1, +1)$
	R	$O_h \times Z_2^T$	$(-1, +1, -1, +1, -1)$	$(+1, -1, +1, +1, -1)$
224.115 IV	Γ	$O_h \times Z_2^T$	$(-1, -1, -1, +1, +1)$	$(+1, +1, +1, +1, +1)$
	X	$D_{4h}^2 \times Z_2^T$	$(-1, +1, -1, +1, +1, -1, +1)$	$(+1, -1, +1, +1, +1, -1, +1)$
	M	$D_{4h}^1 \times Z_2^T$	$(-1, -1, -1, -1, +1, -1, +1)$	$(+1, +1, +1, -1, +1, -1, +1)$
	R	$O_h \times Z_2^T$	$(-1, +1, -1, +1, +1)$	$(+1, -1, +1, +1, +1)$
225.121 IV	Γ	$O_h \times Z_2^T$	$(-1, -1, -1, +1, +1)$	$(+1, +1, +1, +1, +1)$
	X	$D_{4h}^2 \times Z_2^T$	$(-1, -1, -1, +1, +1, +1, +1)$	$(+1, +1, +1, +1, +1, +1, +1)$
	L	$D_{3d}^3 \times Z_2^T$	$(+1, +1, -1, +1)$	$(+1, -1, +1, +1)$
	W	$D_{2d}^2 \times Z_2^T$	$(-1, -1, +1, -1)$	$(+1, +1, +1, -1)$
226.127 IV	Γ	$O_h \times Z_2^T$	$(-1, -1, -1, +1, +1)$	$(+1, +1, +1, +1, +1)$
	X	$D_{4h}^2 \times Z_2^T$	$(-1, -1, -1, +1, +1, +1, +1)$	$(+1, +1, +1, +1, +1, +1, +1)$
	L	$D_{3d}^3 \times Z_2^T$	$(-1, +1, -1, +1)$	$(-1, -1, +1, +1)$
	W	$D_{2d}^2 \times Z_2^T$	$(-1, -1, +1, +1)$	$(+1, +1, +1, +1)$
227.133 IV	Γ	$O_h \times Z_2^T$	$(-1, -1, -1, +1, +1)$	$(+1, +1, +1, +1, +1)$
	X	$D_{4h}^2 \times Z_2^T$	$(-1, -1, -1, +1, +1, -1, +1)$	$(+1, +1, +1, +1, +1, -1, +1)$
	L	$D_{3d}^3 \times Z_2^T$	$(+1, +1, -1, +1)$	$(+1, -1, +1, +1)$
	W	$D_{2d}^2 \times Z_2^T$	$(+1, -1, +1, -1)$	$(-1, +1, +1, -1)$
228.139 IV	Γ	$O_h \times Z_2^T$	$(-1, -1, -1, +1, +1)$	$(+1, +1, +1, +1, +1)$
	X	$D_{4h}^2 \times Z_2^T$	$(-1, -1, -1, +1, +1, -1, +1)$	$(+1, +1, +1, +1, +1, -1, +1)$
	L	$D_{3d}^3 \times Z_2^T$	$(-1, +1, -1, +1)$	$(-1, -1, +1, +1)$
	W	$D_{2d}^2 \times Z_2^T$	$(+1, -1, +1, +1)$	$(-1, +1, +1, +1)$