Upper bounds for the maximum deviation of the Pearcey process

Christophe Charlier

January 6, 2023

Abstract

The Pearcey process is a universal point process in random matrix theory and depends on a parameter $\rho \in \mathbb{R}$. Let N(x) be the random variable that counts the number of points in this process that fall in the interval [-x, x]. In this note, we establish the following global rigidity upper bound:

$$\lim_{s \to \infty} \mathbb{P}\left(\sup_{x>s} \left| \frac{N(x) - \left(\frac{3\sqrt{3}}{4\pi}x^{\frac{4}{3}} - \frac{\sqrt{3}\rho}{2\pi}x^{\frac{2}{3}}\right)}{\log x} \right| \le \frac{4\sqrt{2}}{3\pi} + \epsilon\right) = 1,$$

where $\epsilon > 0$ is arbitrary. We also obtain a similar upper bound for the maximum deviation of the points, and a central limit theorem for the individual fluctuations. The proof is short and combines a recent result of Dai, Xu and Zhang with another result of Charlier and Claeys.

AMS SUBJECT CLASSIFICATION (2010): 41A60, 60B20, 33B15, 33E20, 35Q15. Keywords: Pearcey process, Rigidity, Random matrix theory.

Introduction and statement of results

The Pearcey process describes the local eigenvalue statistics of large random matrices near the points of the spectrum where the limiting mean eigenvalue density vanishes as a cubic root. This has been established rigorously for random matrix ensembles with an external source [9, 10, 34, 6], a two-matrix model [21], large complex correlated Wishart matrices [23, 24], and for general complex Hermitian Wigner-type matrices [17]. Beyond matrix ensembles, the Pearcey process also appears in Brownian motion models [1, 2, 6] and in random plane partition models [30].

The Pearcey process is a determinantal point process on \mathbb{R} whose kernel is given by

$$K_{\rho}^{\text{Pe}}(x,y) = \frac{1}{(2\pi)^2} \int_0^\infty \left(\int_{-\infty}^\infty e^{-\frac{1}{4}t^4 - \frac{\rho}{2}t^2 + it(x+z)} dt \right) \left(\int_{\Sigma} e^{\frac{1}{4}t^4 + \frac{\rho}{2}t^2 + it(y+z)} dt \right) dz$$

where $\rho \in \mathbb{R}$ is a parameter of the model and Σ consists of four rays: $\Sigma = (e^{\frac{\pi i}{4}} \infty, 0) \cup (0, e^{\frac{3\pi i}{4}} \infty) \cup (e^{-\frac{3\pi i}{4}} \infty, 0) \cup (0, e^{-\frac{\pi i}{4}} \infty)$. If $\rho \to +\infty$, the Pearcey process favors the point configurations with fewer points near 0, and in this case the large gap asymptotics are closely related to the Airy gap probabilities [5, Theorem 5.1]. Large gap asymptotics and first exponential moment asymptotics for any fixed ρ have only recently been obtained by Dai, Xu and Zhang [15, 16].

In this note, we obtain a central limit theorem (CLT) and establish two global rigidity upper bounds for the Pearcey process. For $x \ge 0$, let N(x) denote the random variable that counts the number of points in the Pearcey process that fall in the interval [-x, x], and let $x_k \ge 0$ denote the smallest number such that $N(x_k) = k$. All the points $\{x_k\}_{k\ge 1}$ almost surely exist, and by definition they satisfy $0 < x_1 < x_2 < x_3 < \dots$ **Theorem 1.** As $k \to +\infty$, we have

$$\pi \frac{\frac{3\sqrt{3}}{4\pi} x_k^{\frac{4}{3}} - \frac{\sqrt{3}\rho}{2\pi} x_k^{\frac{2}{3}} - k}{\sqrt{\log k}} \quad \stackrel{d}{\longrightarrow} \quad \mathcal{N}(0,1) \tag{1}$$

where $\stackrel{d}{\longrightarrow}$ means convergence in distribution and $\mathcal{N}(0,1)$ is a zero-mean normal random variable with variance 1. Furthermore, for any $\epsilon > 0$, we have

$$\lim_{s \to \infty} \mathbb{P}\left(\sup_{x > s} \left| \frac{N(x) - \left(\frac{3\sqrt{3}}{4\pi}x^{\frac{4}{3}} - \frac{\sqrt{3}\rho}{2\pi}x^{\frac{2}{3}}\right)}{\log x} \right| \le \frac{4\sqrt{2}}{3\pi} + \epsilon\right) = 1,\tag{2}$$

$$\lim_{k_0 \to \infty} \mathbb{P}\left(\sup_{k \ge k_0} \frac{|\frac{3\sqrt{3}}{4\pi} x_k^{\frac{4}{3}} - \frac{\sqrt{3}\rho}{2\pi} x_k^{\frac{2}{3}} - k|}{\log k} \le \frac{\sqrt{2}}{\pi} + \epsilon\right) = 1.$$
(3)

Remark 2. We emphasize that the x_k 's are not exactly the points of the Pearcey process, but rather they are the points of the "absolute value" of the Pearcey process. More precisely, if ξ_1, ξ_2, \ldots are the points of the Pearcey process ordered such that $|\xi_1| < |\xi_2| < \ldots$, then $x_k = |\xi_k|$.

The CLT (1) is a rather straightforward consequence of the following CLT obtained by Dai, Xu and Zhang [16, Corollary 2.4]:

$$\frac{N(s) - \left(\frac{3\sqrt{3}}{4\pi}s^{\frac{4}{3}} - \frac{\sqrt{3}\rho}{2\pi}s^{\frac{2}{3}}\right)}{\frac{2\sqrt{\log s}}{\sqrt{3\pi}}} \xrightarrow{d} \mathcal{N}(0,1) \quad \text{as} \quad s \to +\infty.$$
(4)

The CLTs (1) and (4) give information about the fluctuations of a single point around its classical location. Similar CLTs exist for various other point processes, see e.g. [26, 22]. On the other hand, (2)–(3) are upper bounds for the maximum (properly rescaled) fluctuations of the points; they give information about the global rigidity of the Pearcey process. Over the past few years, we have witnessed significant progress in understanding the global rigidity of various point processes, see e.g. [18, 3, 12, 25, 31, 28, 13, 11, 14], and this note can be viewed as a modest contribution to this ongoing effort. The global rigidity upper bounds (2)–(3) follow (almost) directly from two recent results: a global rigidity theorem from [11] and the first exponential moment asymptotics for the Pearcey process from [16]. This fact has remained unnoticed until now probably because the general result of [11] applies to point processes with almost surely a smallest particle, and that the Pearcey process does not meet this criteria. Our simple idea in this note is to apply [11] to the "absolute value" of the Pearcey process.

On the supposition that upper bounds obtained via [11] are sharp, see in particular [11, Remark 1.3] and Figures 1 and 2 below, we also conjecture the following global rigidity lower bounds.

Conjecture 3. For any $\epsilon > 0$, we have

$$\lim_{s \to \infty} \mathbb{P}\left(\sup_{x > s} \left| \frac{N(x) - \left(\frac{3\sqrt{3}}{4\pi}x^{\frac{4}{3}} - \frac{\sqrt{3}\rho}{2\pi}x^{\frac{2}{3}}\right)}{\log x} \right| \ge \frac{4\sqrt{2}}{3\pi} - \epsilon\right) = 1,\tag{5}$$

$$\lim_{k_0 \to \infty} \mathbb{P}\left(\sup_{k \ge k_0} \frac{|\frac{3\sqrt{3}}{4\pi} x_k^{\frac{4}{3}} - \frac{\sqrt{3}\rho}{2\pi} x_k^{\frac{2}{3}} - k|}{\log k} \ge \frac{\sqrt{2}}{\pi} - \epsilon\right) = 1.$$
 (6)

Remark 4. For several point processes in random matrix theory, the so-called second exponential moment asymptotics have turned out to be important in obtaining optimal rigidity lower bounds, see e.g. [3, 12, 31, 13]. Therefore, the second exponential moment asymptotics of the Pearcey process are expected to be relevant in proving (or disproving) Conjecture 3. These asymptotics are currently not available in the literature.

Proofs of the main results

Proof of (1). The proof mainly follows Gustavsson [22, Theorem 1.2]. For $s > \frac{|\rho|^{\frac{3}{2}}}{3\sqrt{3}}$, define

$$\mu(s) = \frac{3\sqrt{3}}{4\pi}s^{\frac{4}{3}} - \frac{\sqrt{3}\rho}{2\pi}s^{\frac{2}{3}}, \qquad \sigma^2(s) = \frac{4}{3\pi^2}\log s.$$
(7)

It is easy to verify that the function μ is strictly increasing on its domain of definition, and hence is invertible. Since

$$\sqrt{\sigma^2 \circ \mu^{-1}(k)} = \frac{\sqrt{\log k}}{\pi} (1 + o(1)), \quad \text{as } k \to +\infty,$$

(1) follows if we prove that

$$\frac{\mu(x_k) - k}{\sqrt{\sigma^2 \circ \mu^{-1}(k)}} \quad \xrightarrow{d} \quad \mathcal{N}(0, 1), \qquad \text{as } k \to +\infty.$$
(8)

Let $y \in \mathbb{R}$ be an arbitrary constant. For all sufficiently large k, we define $s_k = \mu^{-1} \left(k + y \sqrt{\sigma^2 \circ \mu^{-1}(k)} \right)$. We have

$$\mathbb{P}\Big[\frac{\mu(x_k)-k}{\sqrt{\sigma^2 \circ \mu^{-1}(k)}} \le y\Big] = \mathbb{P}\Big[N(s_k) \ge k\Big] = \mathbb{P}\Big[\frac{N(s_k)-\mu(s_k)}{\sqrt{\sigma^2(s_k)}} \ge \frac{k-\mu(s_k)}{\sqrt{\sigma^2(s_k)}}\Big]$$
$$= \mathbb{P}\Big[\frac{\mu(s_k)-N(s_k)}{\sqrt{\sigma^2(s_k)}} \le y\frac{\sqrt{\sigma^2 \circ \mu^{-1}(k)}}{\sqrt{\sigma^2(s_k)}}\Big] = \mathbb{P}\Big[\frac{\mu(s_k)-N(s_k)}{\sqrt{\sigma^2(s_k)}} \le y(1+o(1))\Big], \quad \text{as } k \to +\infty.$$

The CLT (8), and hence (1), now follows directly from (4).

As already mentioned, the upper bounds (2) and (3) are rather direct consequences of two recent results from [16] and [11]. Let us briefly recall these results.

Theorem 5. (First exponential moment asymptotics from [16, Theorem 2.3]). We have

$$\mathbb{E}\left[e^{\gamma N(s)}\right] = C(\gamma)e^{\gamma \mu(s) + \frac{\gamma^2}{2}\sigma^2(s)}(1 + \mathcal{O}(s^{-2/3})), \qquad as \ s \to +\infty$$

uniformly for γ in compact subsets of \mathbb{R} , where μ, σ^2 are given by (7) and $C(\gamma)$ is independent of s and continuous as a function of γ .

Remark 6. $C(\gamma)$ has in fact also been obtained in [16], but this is not needed for us.

Theorem 7. ([11, Lemma 2.1 and Theorem 1.2]) Let X be a locally finite random point configuration on \mathbb{R} distributed according to a given point process. Assume that X has almost surely a smallest particle, let x_k denote the k-th smallest point of X, and let $\tilde{N}(x)$ be the random variable that counts the number of points in X that are $\leq x$. Assume that there exist constants C, a > 0, $s_0 \in \mathbb{R}$, $M > \sqrt{2/a}$ and functions $\tilde{\mu}, \tilde{\sigma} : [s_0, +\infty) \to [0, +\infty)$ such that the following holds:

(1) We have

$$\mathbb{E}\left[e^{\gamma \widetilde{N}(s)}\right] \le C e^{\gamma \widetilde{\mu}(s) + \frac{\gamma^2}{2}\widetilde{\sigma}^2(s)},\tag{9}$$

for all $\gamma \in [-M, M]$ and for all $s > s_0$.

(2) The functions $\tilde{\mu}$ and $\tilde{\sigma}$ are strictly increasing and differentiable, and they satisfy

$$\lim_{s \to +\infty} \tilde{\mu}(s) = +\infty, \qquad \lim_{s \to +\infty} \tilde{\sigma}(s) = +\infty, \qquad \lim_{s \to +\infty} \frac{s\tilde{\mu}'(s)}{\tilde{\sigma}^2(s)} = +\infty$$

Moreover, $s \mapsto s\tilde{\mu}'(s)$ is weakly increasing, $\tilde{\sigma}^2 \circ \tilde{\mu}^{-1} : [\tilde{\mu}(s_0), +\infty) \to [0, +\infty)$ is strictly concave and $(\tilde{\sigma}^2 \circ \tilde{\mu}^{-1})(s) \sim a \log s$ as $s \to +\infty$.

Then, for any $\epsilon > 0$, it holds that

$$\lim_{s \to +\infty} \mathbb{P}\left(\sup_{x>s} \left| \frac{\widetilde{N}(x) - \widetilde{\mu}(x)}{\widetilde{\sigma}^2(x)} \right| \le \sqrt{\frac{2}{a}} + \epsilon \right) = 1, \tag{10}$$

$$\lim_{k_0 \to \infty} \mathbb{P}\left(\sup_{k \ge k_0} \frac{|\tilde{\mu}(x_k) - k|}{\tilde{\sigma}^2(\tilde{\mu}^{-1}(k))} \le \sqrt{\frac{2}{a}} + \epsilon\right) = 1.$$
(11)

Proof of (2) and (3). Let $Y = \{\xi_k\}_{k \ge 1}$ be a random point configuration distributed according to the Pearcey process. Since Y does not have almost surely a smallest particle, we cannot apply Theorem 7 to the Pearcey process. Instead we define

$$X = \{ |\xi| : \xi \in Y \}$$

Since Y is locally finite, X has a smallest particle. Note that X is distributed according to a point process which is not determinantal, but that does not matter for Theorem 7. Let $\tilde{N}(x)$ denote the number of points in X that are in the interval [0, x]. Using Theorem 5, it is directly seen that X verifies the assumptions of Theorem 7 with

$$\tilde{\mu} = \mu$$
, $\tilde{\sigma} = \sigma$, $M = 10$, $C = 2 \max_{\gamma \in [-M,M]} C(\gamma)$, $a = \frac{1}{\pi^2}$, s_0 sufficiently large.

By definition of \widetilde{N} , we have

$$\widetilde{N}(x) = \#(X \cap [0, x]) = \#(Y \cap [-x, x]) = N(x).$$

Now (2)–(3) follow straightforwardly from (10)–(11) after substituting $\tilde{\mu}, \tilde{\sigma}, a, \tilde{N}(x)$ by $\mu, \sigma, \frac{1}{\pi^2}, N(x)$, respectively.

Numerical support for Theorem 1 and Conjecture 3

We provide here some numerical data to support the validity of Theorem 1 and Conjecture 3. For this, we use the result [6, Theorem 1.1] of Bleher and Kuijlaars which states that the (properly rescaled) eigenvalues around 0 of a large Gaussian random matrix with an external source approximate the random points of the Pearcey process. More precisely, consider the random matrix ensemble

$$\frac{1}{Z_n}e^{-n\operatorname{Tr}(\frac{1}{2}M^2-AM)}dM, \qquad n \in \mathbb{N}_{>0}, \ n \text{ even},$$

defined on the set $\{M\}$ of all $n \times n$ Hermitian matrices, where A is a diagonal matrix with two eigenvalues $\pm (1 + \frac{\rho}{2\sqrt{n}})$ of equal multiplicities. The eigenvalues $\lambda_1^{(n)}, \ldots, \lambda_n^{(n)}$ of M are distributed according to a determinantal point process on \mathbb{R} whose kernel $K_{n,\rho}$ satisfies (see [6, Theorem 1.1]):

$$\lim_{n \to +\infty} \frac{1}{n^{3/4}} K_{n,\rho}\left(\frac{x}{n^{3/4}}, \frac{y}{n^{3/4}}\right) = K_{\rho}^{\text{Pe}}(x, y), \qquad x, y \in \mathbb{R}.$$
 (12)

Let $N_n(x) := \#\{\lambda_j^{(n)} : \lambda_j^{(n)} \in (-\frac{x}{n^{3/4}}, \frac{x}{n^{3/4}})\}$. One expects from (12) that $N_n(x)$ converges in distribution to N(x) as $n \to +\infty$. Let us choose the numbering of the $\lambda_j^{(n)}$'s such that $|\lambda_1^{(n)}| \leq \ldots \leq |\lambda_n^{(n)}|$. From (12), one also expects that for any fixed $k \in \mathbb{N}_{>0}$, the random variables $n^{3/4}\lambda_1^{(n)}, \ldots, n^{3/4}\lambda_k^{(n)}$ converge in distribution to x_1, \ldots, x_k as $n \to +\infty$.



Figure 1: Numerical support for (2) and (5).

The probabilistic bound (2) implies that for any $\epsilon > 0$, the probability that

$$\mu(x) - \left(\frac{4\sqrt{2}}{3\pi} + \epsilon\right)\log x \le N(x) \le \mu(x) + \left(\frac{4\sqrt{2}}{3\pi} + \epsilon\right)\log x \quad \text{for all } x > s \tag{13}$$

tends to 1 as $s \to +\infty$. In Figure 1 (left), the blue step-like curves represent some graphs of the random function $x \mapsto N_n(x)$ for n = 400 and two values of ρ , the blue smooth curves are the upper and lower bounds in (13) with $\epsilon = 0.05$, and the orange curves represent $x \mapsto \mu(x)$. In Figure 1 (right), the orange lines indicate the heights $\pm \frac{4\sqrt{2}}{3\pi} \pm \epsilon$ with $\epsilon = 0.05$, and the blue curve is a graph of the random function

$$x \mapsto \frac{N_n(x) - \mu(x)}{\log x},$$

for n = 400 and $\rho = -1.31$. We see that some of the local minimums and maximums of this function are in the bands between the orange lines, which supports the validity of both (2) and (5).



Figure 2: Numerical support for (3) and (6).

The result (3) implies that for any $\epsilon > 0$, the probability that

$$\mu^{-1}\left(k - \left(\frac{\sqrt{2}}{\pi} + \epsilon\right)\log k\right) \le x_k \le \mu^{-1}\left(k + \left(\frac{\sqrt{2}}{\pi} + \epsilon\right)\log k\right) \quad \text{for all } k \ge k_0 \tag{14}$$

tends to 1 as $k_0 \to +\infty$. In Figure 2 (left), the blue dots represent the random points $(k, n^{3/4} \lambda_k^{(n)})$ for n = 400 and two values of ρ , the blue curves are the upper and lower bounds in (14) with $\epsilon = 0.05$, and the orange curves represent

$$k \mapsto \mu^{-1}(k) = \left(\frac{1}{3}\left(\rho + \sqrt{4\sqrt{3}\pi k + \rho^2}\right)\right)^{3/2}.$$

In Figure 2 (right), the orange lines indicate the heights $\pm \frac{\sqrt{2}}{\pi} \pm \epsilon$ with $\epsilon = 0.05$, and the blue dots have coordinates

$$\left(k, \frac{\mu(n^{3/4}\lambda_k^{(n)}) - k}{\log k}\right),$$

for n = 400 and $\rho = 2.19$. We see that some of these points are in the bands between the orange lines, which supports the validity of both (3) and (6).

Acknowledgements. The author was supported by the European Research Council, Grant Agreement No. 682537.

References

- M. Adler, N. Orantin and P. van Moerbeke, Universality for the Pearcey process, *Phys. D* 239 (2010), 924–941.
- [2] M. Adler and P. van Moerbeke, PDEs for the Gaussian ensemble with external source and the Pearcey distribution, Comm. Pure Appl. Math. 60 (2007), 1261–1292.
- [3] L.-P. Arguin, D. Belius, and P. Bourgade, Maximum of the characteristic polynomial of random unitary matrices. Comm. Math. Phys. 349 (2017), 703–751.
- [4] N. Berestycki, C. Webb, and M. D. Wong, Random Hermitian matrices and Gaussian multiplicative chaos, Probab. Theory Relat. Fields 172 (2018), no. 172, 103–189.
- [5] M. Bertola and M. Cafasso, The transition between the gap probabilities from the Pearcey to the Airy process–a Riemann–Hilbert approach, Int. Math. Res. Not. 2012 (2012), 1519–1568.
- [6] P.M. Bleher and A.B.J. Kuijlaars, Large n limit of Gaussian random matrices with external source, part III: double scaling limit, Comm. Math. Phys. 270 (2007), 481–517.
- [7] A. Borodin, *Determinantal point processes*, The Oxford handbook of random matrix theory, edited by G. Akemann, J. Baik, and P. Di Francesco, 231–249, Oxford Univ. Press, Oxford, 2011.
- [8] P. Bourgade, L. Erdős, and H.-T. Yau, Edge universality of beta ensembles, Comm. Math. Phys. 332 (2014), 261–353.
- [9] E. Brézin and S. Hikami, Level spacing of random matrices in an external source, *Phys. Rev. E.* 58 (1998), 7176–7185.
- [10] E. Brézin and S. Hikami, Universal singularity at the closure of a gap in a random matrix theory, *Phys. Rev. E.* 57 (1998), 4140–4149.
- [11] C. Charlier and T. Claeys, Global rigidity and exponential moments for soft and hard edge point processes, arXiv:2002.03833.
- [12] R. Chhaibi, T. Madaule, and J. Najnudel, On the maximum of the CβE field. Duke Math. J. 167 (2018), 2243–2345.
- [13] T. Claeys, B. Fahs, G. Lambert, and C. Webb, How much can the eigenvalues of a random Hermitian matrix fluctuate?, arxiv:1906.01561.

- [14] T. Claeys, G. Glesner, A. Minakov and M. Yang, Asymptotics for averages over classical orthogonal ensembles, arXiv:2008.07785.
- [15] D. Dai, S-X Xu and L. Zhang, Asymptotics of Fredholm determinant associated with the Pearcey kernel, arXiv:2002.06370.
- [16] D. Dai, S-X Xu and L. Zhang, On the deformed Pearcey determinant, arXiv:2007.12691.
- [17] L. Erdős, T. Krüger and D. Schröder, Cusp universality for random matrices I: local Law and the complex Hermitian case, Comm. Math. Phys. 378 (2020), 1203–1278.
- [18] L. Erdős, B. Schlein, and H.-T. Yau, Semicircle law on short scales and delocalization of eigenvectors for Wigner random matrices, Ann. Probab. 37 (2009), no. 3, 815–852.
- [19] L. Erdős and H.T. Yau, A dynamical approach to random matrix theory, Courant Lecture Notes in Mathematics 28, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 2017.
- [20] L. Erdős, H.-T. Yau, and J. Yin, Rigidity of eigenvalues of generalized Wigner matrices, Adv. Math. 229 (2012), no. 3, 1435–1515.
- [21] D. Geudens and L. Zhang, Transitions between critical kernels: from the tacnode kernel and critical kernel in the two-matrix model to the Pearcey kernel, Int. Math. Res. Not. IMRN 2015 (2015), 5733– 5782.
- [22] J. Gustavsson, Gaussian fluctuations of eigenvalues in the GUE, Ann. Inst. H. Poincare Probab. Statist. 41 (2005), 151–178.
- [23] W. Hachem, A. Hardy and J. Najim, Large complex correlated Wishart matrices: fluctuations and asymptotic independence at the edges, Ann. Probab. 44 (2016), 2264–2348.
- [24] W. Hachem, A. Hardy and J. Najim, Large complex correlated Wishart matrices: the Pearcey kernel and expansion at the hard edge, *Electron. J. Probab.* **21** (2016), 36 pp.
- [25] D. Holcomb and E. Paquette, The maximum deviation of the Sine-β counting process, *Electron. Com*mun. Probab. 23 (2018), paper no. 58, 13 pp.
- [26] K. Johansson, On fluctuations of eigenvalues of random Hermitian matrices, Duke Math. J. 91 (1998), no. 1, 151–204.
- [27] K. Johansson. Random matrices and determinantal processes, Mathematical statistical physics, 1–55, Elsevier B.V., Amsterdam, 2006.
- [28] G. Lambert, Mesoscopic central limit theorem for the circular beta-ensembles and applications, arxiv:1902.06611.
- [29] M.Y. Mo, Universality in complex Wishart ensembles for general covariance matrices with 2 distinct eigenvalues, *Journal of Multivariate Analysis* 101 (2009), 1203–1225.
- [30] A. Okounkov and N. Reshetikhin, Random skew plane partitions and the Pearcey process, Comm. Math. Phys. 269 (2007), 571–609.
- [31] E. Paquette and O. Zeitouni, The maximum of the CUE field, Int. Math. Res. Not. 2018 (2018), no. 16, 5028–5119.
- [32] L.A. Pastur, The spectrum of random matrices, Teoret. Mat. Fiz. 10 (1972), 102–112.
- [33] A. Soshnikov, Determinantal random point fields, Russian Math. Surveys 55 (2000), no. 5, 923–975.
- [34] C. Tracy and H. Widom, The Pearcey process, Comm. Math. Phys. 263 (2006), 381-400.