

# Sharp Hardy inequalities in an exterior of a ball

Nikolai Kutev\*      Tsviatko Rangelov \*

## Abstract

New Hardy type inequalities in sectorial area and as a limit in an exterior of a ball are proved. Sharpness of the inequalities is shown as well.

**Keywords:** Hardy inequalities, sharp estimates.

**2010 Mathematics Subject Classification:** 26D10, 35P15

## 1 Introduction

The classical Hardy inequality, proved in Hardy [7, 8] states

$$\int_0^\infty |u'(x)|^p x^\alpha dx \geq \left( \frac{p-1-\alpha}{p} \right)^p \int_0^\infty x^{-p+\alpha} |u(x)|^p dx \quad (1)$$

where  $1 < p < \infty$ ,  $\alpha < p-1$  and  $u(x)$  is absolutely continuous on  $[0, \infty)$ ,  $u(0) = 0$ .

There are several generalizations of (1) in multidimensional case mostly in bounded domains, see Davies [4], Ghoussoub and Moradifam [6], Balinsky et al. [3], Kutev and Rangelov [9] and the literature therein. For unbounded domains, in the exterior of a ball, only few generalizations of (1) are reported in Wang and Zhu [10], Adimurthi et al. [1], Avkhadiiev and Laptev [2].

For example, in Theorem 1.1 in Wang and Zhu [10] for all function  $u \in D_a^{1,2}(R^n)$ , where  $D_a^{1,2}(R^n)$  is the weighted Sobolev space - the completion of  $C_0^\infty(R^n)$  with the norm  $\int_{R^n} |x|^{-2a} |u|^2 dx$ , the following inequality for  $a < \frac{n-2}{2}$  is proved

$$\int_{B_1^c} |x|^{-2a} |\nabla u|^2 dx \geq \left( \frac{n-2-2a}{2} \right)^2 \int_{B_1^c} \frac{|u|^2}{|x|^2(a+1)} dx + \frac{n-2-2a}{2} \int_{\partial B_1^c} u^2, \quad (2)$$

where  $B_1^c = \{|x| > 1\}$ .

Let us mention that in Wang and Zhu [10] Hardy inequalities with weights in unbounded domains  $\Omega \subset R^n, 0 \notin \partial\Omega$  are also considered, see Theorem 1.3 and Remark 1.5.

In Adimurthi et al. [1], Corollary, for  $n \geq 3$  and  $u \in C_0^\infty(B_1^c)$  the limiting case of the Caffarelli–Kohn–Nirenberg inequality

$$\int_{B_1^c} |\nabla u|^2 dx \geq \left( \frac{n-2}{2} \right)^2 \int_{B_1^c} \frac{u^2}{|x|^2} dx + C_n(a) \left[ \int_{B_1^c} X_1^{\frac{2(n-1)}{n-2}} \left( a, \frac{1}{|x|} \right) u^{\frac{2n}{n-2}} dx \right]^{\frac{n-2}{n}}, \quad (3)$$

---

\*Institute of Mathematics and Informatics, Bulgarian Academe of Sciences, 1113, Sofia, Bulgaria  
Corresponding author: T. Rangelov, rangelov@math.bas.bg

is proved, where  $X_1(a, s) = a - \ln s)^{-1}$ ,  $a > 0, 0 < s \leq 1$  and the constant  $C_n(a)$  is given explicitly.

Finally, in Corollary 1 and Remark 1 in Avkhadiev and Laptev [2] the following Hardy inequality is proved for  $n \geq 3$  and  $u \in W_0^{1,2}(B_r^c)$

$$\int_{B_r^c} |\nabla u|^2 dx \geq \left(\frac{n-2}{2}\right)^2 \int_{B_r^c} \frac{u^2}{|x|^2} dx + \frac{1}{4} \int_{B_r^c} \left(\frac{1}{|x-r|^2} - \frac{1}{|x|^2}\right) u^2 dx. \quad (4)$$

At the end of the paper we compare the inequalities (2), (3), (4) with our results.

The aim of the present work is to derive new Hardy inequalities in the exterior of a ball. There are listed also functions for which these inequalities are sharp, i.e., inequalities with an optimal constant of the leading term become equations.

## 2 Inequalities in sectorial area

We start with Hardy inequalities in sectorial area  $B_R \setminus B_r$  where  $B_R, B_r$  are balls centered at zero,  $0 < r < R$ . Let  $1 < p, p' = \frac{p}{p-1}$ ,  $2 \leq n$  and denote  $m = \frac{p-n}{p-1} = \frac{p-n}{p} p'$ .

For functions  $u$  such that  $\int_{B_R \setminus B_r} \left| \frac{\langle x, \nabla u \rangle}{|x|} \right|^p dx < \infty$  let us define two sets

$$M_1(r, R) = \begin{cases} \left| \frac{R^m - \hat{R}^m}{m} \right|^{1-p} \int_{\partial B_{\hat{R}}} |u|^p d\sigma \rightarrow 0, & \hat{R} \rightarrow R - 0, \quad m \neq 0, \\ \left| \ln \frac{R}{\hat{R}} \right|^{1-n} \int_{\partial B_{\hat{R}}} |u|^n d\sigma \rightarrow 0, & \hat{R} \rightarrow R - 0, \quad m = 0 \end{cases}$$

$$M_2(r, R) = \begin{cases} \left| \frac{\hat{r}^m - r^m}{m} \right|^{1-p} \int_{\partial B_{\hat{r}}} |u|^p d\sigma \rightarrow 0, & \hat{r} \rightarrow r + 0, \quad m \neq 0, \\ \left| \ln \frac{\hat{r}}{r} \right|^{1-n} \int_{\partial B_{\hat{r}}} |u|^n d\sigma \rightarrow 0, & \hat{r} \rightarrow r + 0, \quad m = 0 \end{cases}$$

where  $\langle \cdot, \cdot \rangle$  is a scalar product in  $R^n$ .

Let functions  $\psi_j$  be solutions of the problems:

$$-\Delta_p \psi_1 = 0, \quad \text{in } B_R \setminus \bar{B}_r, \quad \psi_1|_{\partial B_R} = 0, \quad \psi_1|_{\partial B_r} = 1,$$

$$-\Delta_p \psi_2 = 0, \quad \text{in } B_R \setminus \bar{B}_r, \quad \psi_2|_{\partial B_R} = 1, \quad \psi_2|_{\partial B_r} = 0.$$

Their explicit form is:

$$\psi_1(x) = \begin{cases} \frac{R^m - |x|^m}{R^m - r^m}, & m \neq 0, \\ \frac{\ln \frac{R}{|x|}}{\ln \frac{R}{r}}, & m = 0 \end{cases}, \quad \psi_2(x) = \begin{cases} \frac{|x|^m - r^m}{R^m - r^m}, & m \neq 0, \\ \frac{\ln \frac{|x|}{r}}{\ln \frac{R}{r}}, & m = 0 \end{cases}.$$

We can define vector functions  $f_i$  as  $f_i = \left| \frac{\nabla \psi_i}{\psi_i} \right|^{p-2} \frac{\nabla \psi_i}{\psi_i}$ . in  $B_R \setminus \bar{B}_r$  and

$$\begin{aligned} L^i(u) &= \int_{B_R \setminus B_r} \left| \frac{\langle \nabla \psi_i, \nabla u \rangle}{|\nabla \psi_i|} \right|^p dx = \int_{B_R \setminus B_r} \left| \frac{\langle x, \nabla u \rangle}{|x|} \right|^p dx, \\ K^i(u) &= \int_{B_R \setminus B_r} \left| \frac{\nabla \psi_i}{\psi_i} \right|^p |u|^p dx = \int_{B_R \setminus B_r} |f_i|^{p'} |u|^p dx, \\ K_0^i(u) &= \int_{\partial(B_R \setminus B_r)} \left| \frac{\nabla \psi_i}{\psi_i} \right|^{p-2} \left\langle \frac{\nabla \psi_i}{\psi_i}, \nu \right\rangle |u|^p ds = \int_{\partial(B_R \setminus B_r)} \langle f_i, \nu \rangle |u|^p dx. \end{aligned} \quad (5)$$

Here  $\nu$  is the outward normal to  $B_R \setminus B_r$ .

The following Theorem takes place.

**Theorem 2.1.** *For every  $u \in M_i(r, R)$  we have*

$$L^i(u) \geq \left( \frac{1}{p} \right)^p \frac{[K_0^i(u) + (p-1)K_1^i(u)]^p}{(K_1^i(u))^{p-1}} = K^i(u). \quad (6)$$

where  $\nu$  is the outward normal to  $B_R \setminus B_r$ .

*Proof.* We follow the proof of Proposition 1 in Fabricant et al. [5]. Since

$$\int_{B_{\hat{R}} \setminus B_{\hat{r}}} \langle f_i, \nabla |u|^p \rangle dx = p \int_{B_{\hat{R}} \setminus B_{\hat{r}}} |u|^{p-2} u \langle f_i, \nabla u \rangle dx, \quad (7)$$

where  $r < \hat{r} < \hat{R} < R$ . Then applying Hölder inequality on the *rhs* of (7) with  $\frac{\langle x, \nabla u \rangle}{|x|}$  and  $|f_i| |u|^{p-2} u$  as factors of the integrand we get

$$\begin{aligned} \int_{B_{\hat{R}} \setminus B_{\hat{r}}} \langle f_i, \nabla |u|^p \rangle dx &\leq p \left( \int_{B_{\hat{R}} \setminus B_{\hat{r}}} \left| \frac{\langle x, \nabla u \rangle}{|x|} \right|^p dx \right)^{1/p} \\ &\quad \times \left( \int_{B_{\hat{R}} \setminus B_{\hat{r}}} |f_i|^{p'} |u|^p dx \right)^{1/p'}. \end{aligned} \quad (8)$$

Rising to  $p$  power both sides of (8) it follows that

$$\int_{B_{\hat{R}} \setminus B_{\hat{r}}} \left| \frac{\langle x, \nabla u \rangle}{|x|} \right|^p dx \geq \frac{\left| \frac{1}{p} \int_{B_{\hat{R}} \setminus B_{\hat{r}}} \langle f_i, \nabla |u|^p \rangle dx \right|^p}{\left( \int_{B_{\hat{R}} \setminus B_{\hat{r}}} |f_i|^{p'} |u|^p dx \right)^{p-1}}. \quad (9)$$

Integrating by parts the numerator of the right hand side of (9) we get

$$\begin{aligned} \frac{1}{p} \int_{B_{\hat{R}} \setminus B_{\hat{r}}} \langle f_i, \nabla |u|^p \rangle dx &= \frac{1}{p} \int_{\partial B_{\hat{R}} \cup \partial B_{\hat{r}}} \langle f_i, \nu \rangle |u|^p dS - \frac{1}{p} \int_{B_{\hat{R}} \setminus B_{\hat{r}}} \operatorname{div} f_i |u|^p dx \\ &= \frac{1}{p} \int_{\partial B_{\hat{R}} \cup \partial B_{\hat{r}}} \langle f_i, \nu \rangle |u|^p dS + \left( \frac{p-1}{p} \right) \int_{B_{\hat{R}} \setminus B_{\hat{r}}} |f_i|^{p'} |u|^p dx \\ &\rightarrow \frac{1}{p} ((p-1)K^i + K_0^i), \text{ when } \hat{R} \rightarrow R-0, \hat{r} \rightarrow r+0. \end{aligned} \quad (10)$$

Note that  $\int_{\partial B_R \cup \partial B_r} \langle f_i, \nu \rangle |u|^p dS \geq 0$  for  $u \in M_i(r, R)$ , since  $\nu|_{\partial B_R} = \frac{x}{|x|}|_{\partial B_R}$ ,  $\nu|_{\partial B_r} = -\frac{x}{|x|}|_{\partial B_r}$ . From (9) and (10) we obtain (6) since

$$-\operatorname{div} f_i = (p-1)|f_i|^{p'}.$$

□

### 3 Inequalities in an exterior of a ball $B_r^c = R^n \setminus \bar{B}_r$

Let us introduce functions  $u$  such that

$$\int_{B_r^c} \left| \frac{\langle x, \nabla u \rangle}{|x|} \right|^p dx < \infty, \quad \int_{B_r^c} \frac{|u|^p}{|x|^{(n-1)p'}} dx < \infty$$

and define two sets

$$M_1(r, \infty) = \begin{cases} R^{1-n} \int_{\partial B_R} |u|^p d\sigma \rightarrow 0, & R \rightarrow \infty, \quad m \geq 0, \\ R^{1-p} \int_{\partial B_R} |u|^p d\sigma \rightarrow 0, & R \rightarrow \infty, \quad m < 0. \end{cases}$$

$$M_2(r, \infty) = \begin{cases} \left| \frac{\hat{r}^m - r^m}{m} \right|^{1-p} \int_{\partial B_{\hat{r}}} |u|^p d\sigma \rightarrow 0, & \hat{r} \rightarrow r+0, \quad m \neq 0, \\ \left( \ln \frac{\hat{r}}{r} \right)^{1-p} \int_{\partial B_{\hat{r}}} |u|^p d\sigma \rightarrow 0, & \hat{r} \rightarrow r+0, \quad m = 0. \end{cases}$$

In a similar way we can prove Theorem 2.1, replacing  $B_R \setminus \bar{B}_r$  with  $B_r^c = R^n \setminus \bar{B}_r$  and  $\partial(B_R \setminus \bar{B}_r)$  with  $\partial B_r^c = \partial B_r$  and  $L^i(u), K_0^i(u), K_1^i(u)$  define in (5) for  $R \rightarrow \infty$ . The inequalities below for functions of  $M_i(r, \infty), i = 1, 2$  can be obtained with a limit  $R \rightarrow \infty$ .

**Proposition 3.1.** *For every  $u \in M_1(r, \infty)$  the following inequalities hold:*

(i)

$$\left( \int_{B_r^c} \frac{|u|^p}{|x|^{(n-1)p'}} dx \right)^{\frac{1}{p'}} \left( \int_{B_r^c} \left| \frac{\langle x, \nabla u \rangle}{|x|} \right|^p dx \right)^{\frac{1}{p}} \geq \frac{1}{p} r^{1-n} \int_{\partial B_r} |u|^p dS, \quad \text{for } m > 0. \quad (11)$$

With function  $u_\alpha(x) = e^{-\alpha|x|^m}$ ,  $\alpha > 0$  inequality (11) becomes equality.

(ii) For  $m < 0$  we get:

$$\left( \int_{B_r^c} \left| \frac{\langle x, \nabla u \rangle}{|x|} \right|^p dx \right)^{\frac{1}{p}} \geq \frac{|m|}{p'} \left( \int_{B_r^c} \frac{|u|^p}{|x|^p} dx \right)^{\frac{1}{p}} + \frac{1}{p} r^{1-p} \int_{\partial B_r} |u|^p dS \left( \int_{B_r^c} \frac{|u|^p}{|x|^p} dx \right)^{-\frac{1}{p'}}, \quad \text{for } m < 0. \quad (12)$$

With function  $u_k(x) = |x|^{km}$ ,  $k > p'$  inequality (12) becomes an equality.

(iii)

$$\begin{aligned} & \left( \int_{B_r^c} \frac{|u|^n}{|x|^n} dx \right)^{\frac{n-1}{n}} \left( \int_{B_r^c} \left| \frac{\langle x, \nabla u \rangle}{|x|} \right|^n dx \right)^{\frac{1}{n}} \\ & \geq \frac{1}{n} r^{1-n} \int_{\partial B_r} |u|^n dS, \quad \text{for } m = 0. \end{aligned} \quad (13)$$

With function  $u_q(x) = |x|^q$ ,  $q < 0$  inequality (13) becomes equality.

*Proof.* For  $m \neq 0$  the inequality (6) has the form

$$\begin{aligned} & \left( \int_{B_R \setminus B_r} \left| \frac{\langle x, \nabla u \rangle}{|x|} \right|^p \right)^{\frac{1}{p}} \geq \frac{1}{p'} \left( \int_{B_R \setminus B_r} \frac{|u|^p}{|x|^{(n-1)p'} \left| \frac{R^m - |x|^m}{m} \right|^p} dx \right)^{\frac{1}{p}} \\ & + \frac{1}{p} r^{1-n} \left| \frac{R^m - r^m}{m} \right|^{1-p} \int_{\partial B_r} |u|^p dS \left( \int_{B_R \setminus B_r} \frac{|u|^p}{|x|^{(n-1)p'} \left| \frac{R^m - |x|^m}{m} \right|^p} dx \right)^{-\frac{1}{p'}}. \end{aligned} \quad (14)$$

Analogously, for  $m = 0$ , i. e.  $p = n$  the inequality (6) becomes

$$\begin{aligned} & \left( \int_{B_R \setminus B_r} \left| \frac{\langle x, \nabla u \rangle}{|x|} \right|^n dx \right)^{\frac{1}{n}} \geq \frac{n-1}{n} \left( \int_{B_R \setminus B_r} \frac{|u|^n}{|x|^n \left| \ln \frac{R}{|x|} \right|^n} dx \right)^{\frac{1}{n}} \\ & + \frac{1}{n} \left( r \ln \frac{R}{r} \right)^{1-n} \int_{\partial B_r} |u|^n dS \left( \int_{B_R \setminus B_r} \frac{|u|^n}{|x|^n \left| \ln \frac{R}{|x|} \right|^n} dx \right)^{\frac{1-n}{n}}. \end{aligned} \quad (15)$$

(i) For  $m > 0$ , after the limit  $R \rightarrow \infty$  in (14) we obtain

$$\begin{aligned} & \frac{1}{p'} \left( \int_{B_R \setminus B_r} \frac{|u|^p}{|x|^{(n-1)p'} \left| \frac{R^m - |x|^m}{m} \right|^p} dx \right)^{\frac{1}{p}} \rightarrow_{R \rightarrow \infty} 0 \quad \text{and} \\ & \frac{1}{p} r^{1-n} \left| \frac{R^m - r^m}{m} \right|^{1-p} \int_{\partial B_r} |u|^p dS \left( \int_{B_R \setminus B_r} \frac{|u|^p}{|x|^{(n-1)p'} \left| \frac{R^m - |x|^m}{m} \right|^p} dx \right)^{-\frac{1}{p'}} \\ & \rightarrow_{R \rightarrow \infty} \frac{1}{p} r^{1-n} \int_{\partial B_r} |u|^p dS \left( \int_{B_r^c} \frac{|u|^p}{|x|^{(n-1)p'}} dx \right)^{-\frac{1}{p'}}. \end{aligned}$$

and hence (11) holds.

Let us check that with the function  $u_\alpha(x) = e^{-\alpha|x|^m}$ ,  $\alpha > 0$  inequality (11) becomes an

equality. Simply computation gives us

$$\begin{aligned}
I_{1\alpha} &= \left( \int_{B_r^c} \frac{e^{-\alpha p|x|^m}}{|x|^{(n-1)p'}} dx \right)^{\frac{1}{p'}} = \left( \int_{S_1} \int_r^\infty e^{-\alpha p \rho^m} \rho^{-(n-1)p'} \rho^{n-1} d\rho dS \right)^{\frac{1}{p'}} \\
&= |S_1|^{\frac{1}{p'}} \left( \frac{1}{m} \int_r^\infty e^{-\alpha p \rho^m} d\rho^m \right)^{\frac{1}{p'}} = |S_1|^{\frac{1}{p'}} \frac{1}{(\alpha p m)^{\frac{1}{p'}}} e^{-\alpha(p-1)r^m}, \\
I_{2\alpha} &= \left( \int_{B_r^c} \left| \frac{\langle x, \nabla u \rangle}{|x|} \right|^p dx \right)^{\frac{1}{p}} = \left( \int_{S_1} \int_r^\infty e^{-\alpha p \rho^m} (\alpha m)^p \rho^{-(n-1)p'} \rho^{n-1} d\rho dS \right)^{\frac{1}{p}} \\
&= |S_1|^{\frac{1}{p}} \left( \frac{(\alpha m)^p}{m \alpha p} \int_r^\infty e^{-\alpha p \rho^m} d\rho^m \right)^{\frac{1}{p}} = |S_1|^{\frac{1}{p}} \frac{\alpha m}{(\alpha p m)^{\frac{1}{p}}} e^{-\alpha r^m}, \\
I_{3\alpha} &= \frac{1}{p} r^{1-n} \int_{\partial B_r} |u_\alpha|^p dS = \frac{1}{p} r^{1-n} \int_{S_1} e^{-\alpha p r^m} r^{n-1} dS = |S_1| \frac{1}{p} e^{-\alpha p r^m},
\end{aligned}$$

and we get the equality  $I_{1\alpha} I_{2\alpha} = I_{3\alpha}$ .

(ii) For  $m < 0$ , after the limit  $R \rightarrow \infty$  in (14) we obtain (12). Since  $k > \frac{1}{p'}$  it follows that  $u_k(x) \in M_1(r, \infty)$ . Moreover, inequality (12) becomes equality. Indeed,

$$\begin{aligned}
I_{1k} &= \left( \int_{B_r^c} \left| \frac{\langle x, \nabla u_k \rangle}{|x|} \right|^p dx \right)^{\frac{1}{p}} = \left( \int_{B_r^c} |km|x|^{km-1}|^p dx \right)^{\frac{1}{p}} \\
&= |km| \left( |S_1| \int_r^\infty \rho^{(km-1)p} \rho^{n-1} d\rho \right)^{\frac{1}{p}} = |km| \left( |S_1| \frac{r^{(km-1)p+n}}{|(km-1)p+n|} \right)^{\frac{1}{p}}, \\
I_{2k} &= \frac{|m|}{p'} \left( \int_{B_r^c} \frac{|u_k|^p}{|x|^p} dx \right)^{\frac{1}{p}} = \frac{|m|}{p'} \left( |S_1| \int_r^\infty \rho^{(km-1)p+n-1} d\rho \right)^{\frac{1}{p}} \\
&= \frac{|m|}{p'} \left( |S_1| \frac{r^{(km-1)p+n}}{|(km-1)p+n|} \right)^{\frac{1}{p}}, \\
I_{3k} &= \frac{1}{p} r^{1-p} \int_{\partial B_r} |u_k|^p dS \left( \int_{B_r^c} \frac{|u_k|^p}{|x|^p} dx \right)^{-\frac{1}{p'}} \\
&= \frac{1}{p} r^{1-p} |S_1| r^{kmp+n-1} \left( |S_1| \frac{r^{(km-1)p+n}}{|(km-1)p+n|} \right)^{-\frac{1}{p'}} \\
&= \frac{1}{p} \left( |S_1| r^{|(km-1)p+n|} \right)^{\frac{1}{p}} \left( |(km-1)p+n| \right)^{\frac{1}{p'}}.
\end{aligned}$$

Since

$$\frac{|km|}{|(km-1)p+n|} = \frac{|m|}{p'} \frac{1}{|(km-1)p+n|} + \frac{1}{p}$$

we get the equality  $I_{1k} = I_{2k} + I_{3k}$ .

(iii) For  $m = 0$ , after the limit  $R \rightarrow \infty$  in (15), since

$$\begin{aligned} & \frac{n-1}{n} \left( \int_{B_R \setminus B_r} \frac{|u|^n}{|x|^n \left| \ln \frac{R}{|x|} \right|^n} dx \right)^{\frac{1}{n}} \rightarrow_{R \rightarrow \infty} 0 \quad \text{and} \\ & \frac{1}{n} \left( r \ln \frac{R}{r} \right)^{1-n} \int_{\partial B_r} |u|^n dS \left( \int_{B_R \setminus B_r} \frac{|u|^n}{|x|^n \left| \ln \frac{R}{|x|} \right|^n} dx \right)^{\frac{1-n}{n}} \\ &= \frac{1}{n} r^{1-n} \ln^{1-n} R \left( 1 - \frac{\ln r}{\ln R} \right)^{1-n} \int_{\partial B_r} |u|^n dS \frac{1}{\ln^{1-n} R} \left( \int_{B_R \setminus B_r} \frac{|u|^n}{|x|^n \left| 1 - \frac{\ln |x|}{\ln R} \right|^n} dx \right)^{\frac{1-n}{n}} \\ &\rightarrow_{R \rightarrow \infty} \frac{1}{n} r^{1-n} \int_{\partial B_r} |u|^n dS \left( \int_{B_r^c} \frac{|u|^n}{|x|^n} dx \right)^{\frac{1-n}{n}} \end{aligned}$$

we obtain (13). Let us check that with function  $u_q(x) = |x|^q$ ,  $q < 0$  inequality (13) becomes equality. Indeed,

$$\begin{aligned} I_{1q} &= \left( \int_{B_r^c} \frac{|u_q|^n}{|x|^n} dx \right)^{\frac{n-1}{n}} = \left( |S_1| \int_r^\infty \rho^{(q-1)n} \rho^{n-1} d\rho \right)^{\frac{n-1}{n}} = |S_1|^{\frac{n-1}{n}} (n|q|)^{-\frac{n-1}{n}} r^{(n-1)q}, \\ I_{2q} &= \left( \int_{B_r^c} \left| \frac{\langle x, \nabla u_q \rangle}{|x|} \right|^n dx \right)^{\frac{1}{n}} = \left( |S_1| \int_r^\infty |q|^n \rho^{(q-1)n} \rho^{n-1} d\rho \right)^{\frac{1}{n}} = |S_1|^{\frac{1}{n}} |q|^{\frac{n-1}{n}} n^{-\frac{1}{n}} r^q, \\ I_{3q} &= \frac{1}{n} r^{1-n} \int_{\partial B_r} |u_q|^n dS = \frac{1}{n} r^{1-n} |S_1| r^{nq} r^{n-1} = |S_1| \frac{1}{n} r^{qn}, \end{aligned}$$

and we get the equality  $I_{1q} I_{2q} = I_{3q}$ . □

**Proposition 3.2.** *For every  $u \in M_2(r, \infty)$  the following inequalities hold:*

(i)

$$\begin{aligned} & \left( \int_{B_r^c} \left| \frac{\langle x, \nabla u \rangle}{|x|} \right|^p dx \right)^{\frac{1}{p}} \geq \frac{m}{p'} \left( \int_{B_r^c} \frac{|u|^p}{|x|^{(n-1)p'} |r^m - |x|^m|^p} dx \right)^{\frac{1}{p}} \\ & + \frac{1}{p} \limsup_{R \rightarrow \infty} \left[ R^{1-p} \int_{\partial B_R} |u|^p dS \right] \left( \int_{B_r^c} \frac{|u|^p}{|x|^{(n-1)p'} ||x|^m - r^m|^p} dx \right)^{-\frac{1}{p'}}, \end{aligned}$$

for  $m > 0$ .

(16)

With function  $u_\varepsilon(x) = |x|^{-\frac{m(1-\varepsilon)}{p'}} (|x|^m - r^m)^{\frac{(1+\varepsilon)}{p'}}$ ,  $0 < \varepsilon < 1$  inequality (16) is  $\varepsilon$ -sharp, i.e.

$$\left(\frac{m}{p'}\right)^p \leq \frac{L^1(u_\varepsilon)}{\left[(K^1(u_\varepsilon))^{\frac{1}{p}} + K_0^1(u_\varepsilon) (K^1(u_\varepsilon))^{-\frac{1}{p'}}\right]^p} \leq \frac{L^1(u_\varepsilon)}{K^1(u_\varepsilon)} \leq \left(\frac{m}{p'}\right)^p (1 + \varepsilon)^p. \quad (17)$$

(ii)

$$\begin{aligned} & \left( \int_{B_r^c} \left| \frac{\langle x, \nabla u \rangle}{|x|} \right|^p dx \right)^{\frac{1}{p}} \geq \frac{|m|}{p'} \left( \int_{B_r^c} \frac{|u|^p}{|x|^{(n-1)p'} |r^m - |x|^m|^p} dx \right)^{\frac{1}{p}} \\ & + \frac{r^{n-p}}{p} \limsup_{R \rightarrow \infty} \left[ R^{1-n} \int_{\partial B_R} |u|^p dS \right] \left( \int_{B_r^c} \frac{|u|^p}{|x|^{(n-1)p'} ||x|^m - r^m|^p} dx \right)^{-\frac{1}{p'}}, \\ & \text{for } m < 0. \end{aligned} \quad (18)$$

With function  $u_s(x) = (r^m - |x|^m)^s$ ,  $s > \frac{1}{p'}$  inequality (18) becomes equality.

(iii)

$$\begin{aligned} & \left( \int_{B_r^c} \left| \frac{\langle x, \nabla u \rangle}{|x|} \right|^n dx \right)^{\frac{1}{n}} \geq \frac{n-1}{n} \left( \int_{B_r^c} \frac{|u|^n}{|x|^n \left| \ln \frac{|x|}{r} \right|^n} dx \right)^{\frac{1}{n}} \\ & + \frac{1}{n} \limsup_{R \rightarrow \infty} \left[ \left( R \ln \frac{R}{r} \right)^{1-n} \int_{\partial B_R} |u|^n dS \right] \left( \int_{B_r^c} \frac{|u|^n}{|x|^n \left| \ln \frac{|x|}{r} \right|^n} dx \right)^{\frac{1-n}{n}}, \\ & \text{for } m = 0. \end{aligned} \quad (19)$$

For function

$$u_\eta(x) = \begin{cases} \left( \ln \frac{|x|}{r} \right)^{\frac{n-1}{n}(1+\frac{\eta}{2})} M^{\frac{n-1}{n}(1-\frac{\eta}{2})}, & r < |x| < M \\ \left( \ln \frac{|x|}{r} \right)^{\frac{n-1}{n}(1+\frac{\eta}{2})} |x|^{\frac{n-1}{n}(1-\frac{\eta}{2})}, & M < |x| \end{cases},$$

where  $0 < \eta < 1$ , inequality (19) is  $\eta$ -sharp, i.e.

$$\left(\frac{n-1}{n}\right)^n \leq \frac{L^1(u_\eta)}{K^1(u_\eta)} \leq \left(\frac{n-1}{n}\right)^n (1 + \eta)^n. \quad (20)$$



*Proof.* For  $m \neq 0$  inequality (6) has the form

$$\begin{aligned} & \left( \int_{B_R \setminus B_r} \left| \frac{\langle x, \nabla u \rangle}{|x|} \right|^p \right)^{\frac{1}{p}} \geq \frac{|m|}{p'} \left( \int_{B_R \setminus B_r} \frac{|u|^p}{|x|^{(n-1)p'} |r^m - |x|^m|^p} dx \right)^{\frac{1}{p}} \\ & + \frac{1}{p} R^{1-n} |R^m - r^m|^{1-p} \int_{\partial B_R} |u|^p dS \left( \int_{B_R \setminus B_r} \frac{|u|^p}{|x|^{(n-1)p'} |r^m - |x|^m|^p} dx \right)^{-\frac{1}{p'}}. \end{aligned} \quad (21)$$

while for  $m = 0$  (6) becomes

$$\begin{aligned} & \left( \int_{B_R \setminus B_r} \left| \frac{\langle x, \nabla u \rangle}{|x|} \right|^n dx \right)^{\frac{1}{n}} \geq \frac{n-1}{n} \left( \int_{B_R \setminus B_r} \frac{|u|^n}{|x|^n \left| \ln \frac{|x|}{r} \right|^n} dx \right)^{\frac{1}{n}} \\ & + \frac{1}{n} \left( R \ln \frac{R}{r} \right)^{1-n} \int_{\partial B_r} |u|^n dS \left( \int_{B_R \setminus B_r} \frac{|u|^n}{|x|^n \left| \ln \frac{|x|}{r} \right|^n} dx \right)^{-\frac{1}{n'}}. \end{aligned} \quad (22)$$

(i) If  $m > 0$  after the limit  $R \rightarrow \infty$  in (21) we obtain

$$\begin{aligned} & \frac{1}{p} \limsup_{R \rightarrow \infty} \left[ R^{1-n} |R^m - r^m|^{1-p} \int_{\partial B_R} |u|^p dS \right] \\ & = \frac{1}{p} \limsup_{R \rightarrow \infty} \left[ R^{1-n} R^{n-p} \left| 1 - \frac{r^m}{R^m} \right|^{1-p} \int_{\partial B_R} |u|^p dS \right] \\ & = \frac{1}{p} \limsup_{R \rightarrow \infty} R^{1-p} \int_{\partial B_R} |u|^p dS \end{aligned}$$

which proves (16).

For the function  $u_\varepsilon(x) = |x|^{-\frac{m(1-\varepsilon)}{p'}} (|x|^m - r^m)^{\frac{(1+\varepsilon)}{p'}}$ ,  $0 < \varepsilon < 1$ ,  $u_\varepsilon(x) \in M_2(r, \infty)$

simple computation give us

$$\begin{aligned}
I_{1\varepsilon} &= L^1(u_\varepsilon) = \int_{B_r^\varepsilon} \left| \frac{\langle x, \nabla u_\varepsilon \rangle}{|x|} \right|^p dx \\
&= \left( \frac{m}{p'} \right)^p \int_{B_r^\varepsilon} |x|^{-m(1-\varepsilon)(p-1)-p} (|x|^m - r^m)^{(1+\varepsilon)(p-1)-p} \\
&\quad \times [(1+\varepsilon)|x|^m - (1-\varepsilon)(|x|^m - r^m)]^p dx \\
&= \left( \frac{m}{p'} \right)^p |S_1| \int_r^\infty (\rho^m - r^m)^{(1+\varepsilon)(p-1)-p} \\
&\quad \times [(1+\varepsilon)\rho^m - (1-\varepsilon)(\rho^m - r^m)]^p \rho^{-m(1-\varepsilon)(p-1)-p+n-1} d\rho \\
&\leq \left( \frac{m}{p'} \right)^p |S_1| (1+\varepsilon)^p \int_r^\infty (\rho^m - r^m)^{(1+\varepsilon)(p-1)-p} \rho^{-m(1-\varepsilon)(p-1)-p+n-1+mp} d\rho \\
&= \left( \frac{m}{p'} \right)^p |S_1| (1-\varepsilon)^p \int_r^\infty (\rho^m - r^m)^{(1+\varepsilon)(p-1)-p} \rho^{-m(1+\varepsilon)(p-1)-\frac{n-1}{p-1}} d\rho,
\end{aligned}$$

because  $n - 1 - p + mp = -\frac{n-1}{p-1}$ .

$$\begin{aligned}
I_{2\varepsilon} &= K^1(u_\varepsilon) = \int_{B_r^\varepsilon} |x|^{-m(1-\varepsilon)(p-1)-(n-1)p'} (|x|^m - r^m)^{(1+\varepsilon)(p-1)-p} dx \\
&= |S_1| \int_r^\infty (\rho^m - r^m)^{(1+\varepsilon)(p-1)-p} \rho^{-m(1-\varepsilon)(p-1)-\frac{n-1}{p-1}} d\rho.
\end{aligned}$$

Thus (17) follows immediately from expressions of  $I_{1\varepsilon}$  and  $I_{2\varepsilon}$ . (ii) When  $m < 0$ , after the limit  $R \rightarrow \infty$  in (21) we get

$$\begin{aligned}
&\frac{1}{p} \limsup_{R \rightarrow \infty} \left[ R^{1-n} |R^m - r^m|^{1-p} \int_{\partial B_R} |u|^p dS \right] \\
&= \frac{1}{p} \limsup_{R \rightarrow \infty} \left[ r^{n-p} R^{1-n} \left| \frac{R^m}{r^m} - 1 \right|^{1-p} \int_{\partial B_R} |u|^p dS \right] \\
&= \frac{r^{n-p}}{p} \limsup_{R \rightarrow \infty} R^{1-n} \int_{\partial B_R} |u|^p dS
\end{aligned}$$

which proves (18).

For the function  $u_s(x) = (r^m - |x|^m)^s$ ,  $s > \frac{1}{p'}$  we have the identities

$$\begin{aligned} I_{1s} &= \left( \int_{B_r^c} \left| \frac{\langle x, \nabla u_s \rangle}{|x|} \right|^p dx \right)^{\frac{1}{p}} = \left( s^p |m|^p |S_1| \int_r^\infty \rho^{p(m-1)+n-1} (r^m - \rho^m)^{(s-1)p} d\rho \right)^{\frac{1}{p}} \\ &= s|m| |S_1|^{\frac{1}{p}} \left( -\frac{1}{m} \int_r^\infty (r^m - \rho^m)^{(s-1)p} d(r^m - \rho^m) \right)^{\frac{1}{p}}, \\ &= s|m|^{\frac{1}{p'}} |S_1|^{\frac{1}{p}} r^{\frac{m}{p}[(s-1)p+1]} [(s-1)p+1]^{-\frac{1}{p}} \end{aligned}$$

because  $p(m-1) + n - 1 = m - 1$ .

$$\begin{aligned} I_{2s} &= \int_{B_r^c} \frac{|u_s|^p}{|x|^{(n-1)p'} |r^m - |x|^m|^p} dx = |S_1| \int_r^\infty \rho^{(n-1)(1-p')} |r^m - \rho^m|^{(s-1)p} d\rho \\ &= \frac{|S_1| r^{m[(s-1)p+1]}}{|m| [(s-1)p+1]}, \end{aligned}$$

$$\begin{aligned} I_{3s} &= \frac{r^{n-p}}{p} \limsup_{R \rightarrow \infty} \left[ R^{1-n} \int_{\partial B_R} |r^m - R^m|^{sp} dS \right] \\ &= \frac{r^{n-p}}{p} \limsup_{R \rightarrow \infty} |R^m - r^m|^{sp} |S_1| = \frac{|S_1|}{p} r^{n-p+mps}. \end{aligned}$$

Simple computation gives us the equality

$$\begin{aligned} &\frac{|m|}{p'} (I_{2s})^{\frac{1}{p}} + I_{3s} (I_{2s})^{-\frac{1}{p'}} \\ &= \frac{|S_1| |m|^{\frac{1}{p'}} r^{\frac{m}{p}[(s-1)p+1]}}{p' [(s-1)p+1]^{\frac{1}{p}}} + \frac{|S_1| r^{n-p+mps}}{p} \left( \frac{|S_1|}{|m|} \right)^{-\frac{1}{p'}} \left[ \frac{r^{m[(s+1)p+1]}}{[(s-1)p+1]} \right]^{-\frac{1}{p'}} \\ &= \frac{1}{p} |m|^{\frac{1}{p'}} |S_1|^{\frac{1}{p}} r^{\frac{m}{p}[(s-1)p+1]} \left( \frac{p-1}{[(s-1)p+1]^p} + [(s-1)p+1]^{\frac{1}{p'}} \right) \\ &= s|m|^{\frac{1}{p'}} |S_1|^{\frac{1}{p}} r^{\frac{m}{p}[(s-1)p+1]} [(s-1)p+1]^{-\frac{1}{p}} = I_{1s} \end{aligned}$$

because

$$n - p + mps - \frac{m}{p'} [(s-1)p+1] = \frac{m}{p} [(s-1)p+1]$$

(iii) After the limit  $R \rightarrow \infty$  in (22) we get (19).

The function  $u_\eta(x)$  belongs to  $M_2(r, \infty)$  for  $m = 0$ . Moreover, for  $r < |x| < M$  we have the equalities

$$I_{1\eta} = L^1(u_\eta) = M^{(n-1)(1-\eta)} \left( 1 + \frac{\eta}{2} \right)^n \left( \frac{n-1}{n} \right)^n |S_1| \int_r^M \left( \ln \frac{\rho}{r} \right)^{(n-1)(1+\frac{\eta}{2})-n} \rho^{-1} d\rho,$$

$$I_{2\eta} = K^1(u_\eta) = M^{(n-1)(1+\frac{\eta}{2})} |S_1| \int_r^M \left( \ln \frac{\rho}{r} \right)^{(n-1)(1+\frac{\eta}{2})-n} \rho^{-1} d\rho,$$

and hance

$$\frac{L^1(u_\eta)}{K^1(u_\eta)} = \left(\frac{n-1}{n}\right)^n \left(1 + \frac{\eta}{2}\right)^n \leq \left(\frac{n-1}{n}\right)^n (1 + \eta)^n.$$

Tedious calculations give us for  $|x| \geq M$  the identities

$$\begin{aligned} I_{1\eta} &= L^1(u_\eta) = \left(\frac{n-1}{n}\right)^n |S_1| \int_M^\infty \rho^{(n-1)(1-\frac{\eta}{2})-1} \left(\ln \frac{\rho}{r}\right)^{(n-1)(1+\frac{\eta}{2})-n} \\ &\times \left[ \left(1 - \frac{\eta}{2}\right) \frac{1}{\rho} \ln \frac{\rho}{r} + \left(1 + \frac{\eta}{2}\right) \right]^n d\rho \\ I_{2\eta} &= K^1(u_\eta) = |S_1| \int_r^M \rho^{(n-1)(1-\frac{\eta}{2})-1} \left(\ln \frac{\rho}{r}\right)^{(n-1)(1+\frac{\eta}{2})-n}, \end{aligned}$$

If  $M$  is sufficiently large, i.e.,

$$\frac{1}{M} \ln \frac{M}{r} < \frac{\eta}{2-\eta} \quad \text{and} \quad M > er,$$

then

$$\left(1 - \frac{\eta}{2}\right) \frac{1}{\rho} \ln \frac{\rho}{r} + 1 + \frac{\eta}{2} \leq \left(1 - \frac{\eta}{2}\right) \frac{\eta}{2-\eta} + 1 + \frac{\eta}{2} = 1 + \eta,$$

because the function  $h(\rho) = \frac{1}{\rho} \ln \frac{\rho}{r}$  is monotone decreasing for  $\rho > er$ . Thus (20) follows from expressions of  $I_{1\eta}$  and  $I_{2\eta}$  above for  $|x| \geq r$ .  $\square$

#### 4 Sharp inequalities in an exterior of a ball $B_r^c$ for $u \in W_0^{1,p}(B_r^c)$

For  $m < 0$ , i.e.  $p < n$ , we can combine inequalities (14) and (12) for functions  $u \in M(r, \infty)$ , where

$$M(r, \infty) = \begin{cases} u \in W_0^{1,p}(B_r^c), \\ \left(\frac{\hat{r}^m - r^m}{m}\right)^{1-p} \int_{\partial B_{\hat{r}}} |u|^p dx \rightarrow 0, \quad \hat{r} \rightarrow r + 0, \\ R^{1-p} \int_{\partial B_R} |u|^p dx \rightarrow 0, \quad \text{for } R \rightarrow \infty. \end{cases}$$

For  $r < \gamma < \infty$ ,  $\gamma = 2^{\frac{1}{|m|}} r$ , we define

$$L_1(u) = \int_{B_\gamma \setminus B_r} \left| \frac{\langle x, \nabla u \rangle}{|x|} \right|^p dx, \quad L_2(u) = \int_{B_\gamma^c} \left| \frac{\langle x, \nabla u \rangle}{|x|} \right|^p dx,$$

$$K_{11}(u) = |m|^p \int_{B_\gamma \setminus B_r} \frac{|u|^p}{|x|^{(n-1)p'} (|x|^m - r^m)^p} dx, \quad K_{12}(u) = |m|^p \int_{B_\gamma^c} \frac{|u|^p}{|x|^p} dx,$$

$$K_{01}(u) = |m|^{p-1} \gamma^{1-p} \int_{\partial B_\gamma} |u|^p d\sigma = K_{02}(u) = |m|^{p-1} \gamma^{1-p} \int_{\partial B_\gamma} |u|^p d\sigma = K_0.$$

**Proposition 4.1.** *If  $\gamma = 2^{1/|m|}r$  then for every  $u \in M(r, \infty)$  the inequality*

$$L(u) = \sum_1^2 L_j(u) = \int_{B_\gamma^c} \left| \frac{\langle x, \nabla u \rangle}{|x|} \right|^p dx \geq \left( \frac{1}{p} \right)^p \sum_1^2 \frac{[K_{0j}(u) + (p-1)K_{1j}(u)]^p}{(K_{1j}(u))^{p-1}} = K(u), \quad (23)$$

holds in  $B_r^c$ .

The inequality (23) becomes an equality for functions  $u_\beta(x) \in M(r, \infty)$ ,  $\beta > 1/p'$  where

$$u_\beta(x) = \begin{cases} (r^m - |x|^m)^\beta & x \in B_\gamma \setminus B_r, \\ |x|^{m\beta}, & x \in B_\gamma^c. \end{cases}$$

*Proof.* As in Theorem 2.1 and Proposition 3.1 ii) we get

$$\begin{aligned} L_1(u) &\geq K_1(u) && \text{in } B_\gamma \setminus B_r, \\ L_2(u) &\geq K_2(u) && \text{in } B_\gamma^c \end{aligned} \quad (24)$$

where

$$K_j(u) = \frac{[K_{0j}(u) + (p-1)K_{1j}(u)]^p}{(K_{1j}(u))^{p-1}}.$$

With the choice of  $\gamma = 2^{1/|m|}r$ , so  $\gamma^p = \gamma^{(n-1)p'}(r^m - \gamma^m)^p$ , we have continuous kernel for  $K_{j1}$  on  $\partial B_\gamma$ .

$$\text{Adding inequalities in (24) we get for } u \in M(r, \infty) \quad L(u) = \sum_1^2 L_j(u) \geq \sum_1^2 K_j(u).$$

Hence from the Joun inequality we obtain (23)

Using (12) in  $B_\gamma^c$  and (15) in  $B_\gamma \setminus B_r$  it follows that the inequality (23) becomes equality for functions  $u_\beta(x)$ ,  $\beta > \frac{1}{p'}$ .  $\square$

Proposition 4.1 gives sharp Hardy inequality in an exterior of a ball  $B_r^c$  for functions  $u \in W_0^{1,p}(B_r^c)$ ,  $p < n$ .

We will illustrate Proposition 3.2 ii) and Proposition 4.1 in the following examples.

**Example 4.1.** For  $p = 2$ ,  $n \geq 3$ ,  $r = 1$ ,  $a = 0$  and  $m = 2 - n < 0$  from Proposition 3.1 ii) it follows that the Hardy inequality

$$\int_{B_1^c} |\nabla u|^2 dx \geq \left( \frac{n-2}{2} \right)^2 \int_{B_1^c} \frac{|u|^2}{|x|^2} dx + \frac{n-2}{2} \int_{\partial B_1^c} u^2 + \frac{1}{4} \int_{\partial B_1^c} u^2 \left( \int_{B_1^c} \frac{|u|^2}{|x|^2} \right)^{-1}, \quad (25)$$

holds. Note that (25) has an additional term in the right hand side in comparison with (2). Moreover, for the functions  $u_k(x) = |x|^{k(2-n)}$ ,  $k > 2$  inequality (25) becomes an equality, i.e., inequality (25) is sharp.

**Example 4.2.** For  $p = 2, n \geq 3, m = 2 - n < 0, \gamma = 2^{\frac{1}{n-2}}r$ , and every function  $u \in W_0^{1,2}(B_r^c)$  the Hardy inequality (24) becomes

$$\begin{aligned}
& \int_{B_r^c} \left| \frac{\langle x, \nabla u \rangle}{|x|} \right|^p dx \geq \left( \frac{n-2}{2} \right)^2 \int_{B_r^c} \frac{u^2}{|x|^2} dx \\
& + 2 \left( \frac{n-2}{2} \right)^2 \int_{B_\gamma \setminus B_r} \frac{\left| \frac{x}{r} \right|^{n-2} \left( 1 - \left| \frac{x}{\gamma} \right|^{n-2} \right)}{\left( \left| \frac{x}{r} \right|^{n-2} - 1 \right)^2} u^2 dx + \frac{2^{-\frac{1}{n-2}}}{r} (n-2) \int_{\partial B_\gamma} u^2 dS \\
& + \frac{2^{-\frac{2}{n-2}}}{r^2} \left( \frac{4}{n-2} \right)^2 \left( \int_{\partial B_\gamma} u^2 dS \right)^2 \\
& \times \left[ \left( \int_{B_\gamma \setminus B_r} \frac{u^2}{|x|^2 \left( \left| \frac{x}{r} \right|^{n-2} - 1 \right)^2} dx \right)^{-2} + \left( \int_{B_r^c} \frac{u^2}{|x|^2} dx \right)^{-2} \right]
\end{aligned} \tag{26}$$

Moreover, for function  $u_\beta(x)$  defined in Proposition 4.1 for  $m = 2 - n$ , inequality (26) becomes equality.

Let us mention that Hardy inequality (26) has the same leading term in the right hand side as in inequality (3), but inequality (26) is sharp one.

Finally, it is difficult to compare inequality (4) with (26), but (26) is sharp one, i.e., for the functions  $u_\beta(x)$  defined in Proposition 4.1, inequality (26) becomes an equality.

**Acknowledgement** This paper is partially supported by the National Scientific Program "Information and Communication Technologies for a Single Digital Market in Science, Education and Security (ICTinSES)", contract No DOI205/23.11.2018, financed by the Ministry of Education and Science in Bulgaria and also by the Grant No BG05M2OP001-1.001-0003, financed by the Science and Education for Smart Growth Operational Program (2014-2020) in Bulgaria and co-financed by the European Union through the European Structural and Investment Funds.

## References

- [1] Adimurthi, S. Filippas, and A. Tertikas. On the best constant of Hardy–Sobolev inequalities. *Nonlinear Anal. T.M.A.*, 70:2826–2833, 2009.
- [2] F. Avkhadiev and A. Laptev. Hardy inequalities for nonconvex domains. In *International Mathematical Series, vol 11, Around Research of Vladimir Maz'ya, I*, pages 1–12, Springer, New York, 2010.
- [3] A. A. Balinsky, W. D. Evans, and R. T. Lewis. *The analysis and geometry of Hardy's Inequality*. Springer Int. Publisher, Cham, 2015.
- [4] E. Davies. A review of Hardy inequalities. *Oper. Theory Adv. Appl.*, 110:57–67, 1998.
- [5] A. Fabricant, N. Kutev, and T. Rangelov. Sharp hardy inequalities in a ball. *Pliska Stud. Math. Bulgar.*, 23:67–80, 2014.
- [6] N. Ghoussoub and A. Moradifam. *Functional inequalities: New perspectives and new applications*. AMS, Math. Surveys and Monographs, v. 187, Rhode Island, 2013.

- [7] G. Hardy. Note on a theorem of Hilbert. *Math. Zeit.*, 6:314–317, 1920.
- [8] G. Hardy. Notes on some points in the integral calculus, LX. an inequality between integrals. *Messenger of Math.*, 54:150–156, 1925.
- [9] N. Kutev and T. Rangelov. Hardy inequalities with double singular weights. 2020. arXiv.2001.07368v2.
- [10] Z-Q. Wang and M. Zhu. Hardy inequalities with boundary terms. *Electron. J. Diff. Eq.*, 43: 1–8, 2003.