

# Balancing costs and benefits of pandemic control in an outbreak phase

Tsuyoshi Hondou

Graduate School of Science, Tohoku University

Sendai 980-8578, Japan

After the first lockdowns in response to the COVID-19 outbreak, many countries faced difficulties in balancing infection control with economics. Because there was scant prior knowledge<sup>1)</sup>, economists began researching this issue<sup>2, 3, 4, 5, 6)</sup> using cost-benefit analysis<sup>7, 8)</sup> and found that infection control processes seriously affect economic efficiency. Rowthorn and Maciejowski<sup>2, 4)</sup>, using economic parameters in the U.K., demonstrated numerically that an optimal balance was found in the process, including keeping infected population stationary. However, because these analyses assume a specific disease and regional parameters, universally applicable knowledge, indispensable for guiding principles of infection control, has not yet been developed. Here we prove a universal result of economic irreversibility<sup>24, 25, 26)</sup> that delaying infection control measures is more expensive than implementing infection control measures early while keeping infected populations stationary. That is, once the infected population increases, society cannot return to its previous state without extra expenditures. This universal result is analytically obtained by focusing on the infection-spreading phase of pandemics, which is applicable not only to COVID-19,

**and whether or not ‘herd immunity’ exists<sup>11, 12)</sup>. It also confirms the numerical observation of stationary infected population in its optimally efficient process<sup>2, 4)</sup>. Our findings suggest that economic irreversibility is a guiding principle for balancing infection control with economic effects.**

Governments in several countries fear adverse economic effects and have hesitated to take measures to control COVID-19 infection, as the economic effects may result in illness and death in the non-infected population<sup>2)</sup>. For example, Japan hesitated to respond to the pandemic. The Japanese government requested that governors increase their medical capacities<sup>10)</sup> as they determined the upper limit for the infected population. This social turbulence is attributed to insufficient knowledge about the relationship between infection control and the economy.

Several economists, perceiving a serious lack of knowledge<sup>1)</sup>, started studying this issue this past spring<sup>3, 4, 5, 6)</sup>. Rowthorn<sup>2)</sup>, along with his colleague Maciejowski<sup>4)</sup>, utilized the cost-benefit analysis (CBA)<sup>7, 8)</sup> to determine how infection control intervention costs could efficiently be utilized (inhibition of infection). Using the susceptible-infected-recovered (SIR) model to simulate the epidemic<sup>9)</sup>, they discussed several infection control processes to determine the optimal process. The optimal process includes the stationary state of the constant infection population in its principal part. While the methodology and results of Rowthorn et al. are pioneering and suggestive, they are not straightforward enough to generalize their results because the study investigated specific situations with given parameter sets. Therefore, explicit solutions independent of specific parameters are needed to reveal the universal property, while Rowthorn stated that one must explore using numerical simulation as an explicit solution is absent<sup>2)</sup>. Explicit solutions could be applicable in the United Kingdom and other countries during different situations, including COVID-19 and other pandemics.

From the physicists’ perspective, the optimization in CBA is similar to finding the minimum state of free energy. In addition, the finding<sup>2, 4)</sup> that the most efficient process includes the

stationary state suggests an analogous structure with thermodynamic irreversibility.

In this paper, we analytically show the basic property of economic cost in the infection control process by analysing the cyclic processes of the system's state variable. For this purpose, we restrict ourselves to the infection-spreading phase in the pandemic model, in which the infected population grows exponentially in the absence of infection control. In several pandemics, including COVID-19, the society may not arrive at a traditional immune state called 'herd immunity' as indicated in several studies<sup>11, 12)</sup>. However, the infection-spreading phase is universal and principal, irrespective of whether or not herd immunity exists. By comparing the stationary state of a constant infected population, we will derive several explicit solutions and inequalities of costs in infection control processes and show economic irreversibility in infection control. With these explicit results, we prove that delaying infection control measures is always more expensive than implementing early measures while keeping the infected population stationary.

## Formulation with Cost-Benefit Analysis

Infection control comprises measures taken to decrease the number of patients infected by an individual. The average number within society is called the "effective reproduction number",  $R_t$ . When  $R_t$  drops below 1, epidemics subside. Several measures, including handwashing, wearing masks, suspension of business activities, and lockdowns can be taken to reduce  $R_t$  from its uncontrolled (natural) value,  $R_N > 1$ .  $R_N$  equals the basic reproduction number  $R_0$  for the initial phase of infection. These measures have a negative influence on the economy and society<sup>2)</sup>. This social cost,  $\hat{C}$ , is positively correlated to the strength of the measure. Rowthorn assumed<sup>2)</sup> that the infection control measure is taken through the valuable  $q$  as  $R_t = R_N(1 - q(t))$ , where  $q$  represents the intensity of social intervention against pandemics. Then, he defined the social cost per unit of time as a function of  $q$ :  $\hat{C} = \hat{C}(q)^{2, 4)$ . He assumed  $\hat{C}(0) = 0$  because there is

no infection control at  $q = 0$ .

Here, we consider the social cost induced by infection measure as a function of the effective reproduction number,  $R_t$ , instead of  $q$ . While Rowthorn<sup>2)</sup> assumes maximum strength,  $q_{\max}$ , which corresponds to the minimum effective reproduction number,  $R_t$ , we do not adopt this inessential assumption. Our functional form of the function  $C(R_t)$  itself is different from  $\hat{C}(q)$ , while the following basic assumptions, Eqs.(1,2,3,4), are essentially the same as Rowthorn. Hereafter, we refer to the social cost per unit time as “intervention cost” in the form of  $C(R_t)$ . The following are assumed in the function  $C(R_t)$ .

The condition without intervention measures corresponds to  $R_t = R_N$ , in which  $C(R_N) = 0$ . The cost should increase, as the effective reproduction number decreases. The rate of increase of  $C(R_t)$  should also increase as the effective reproduction number decreases. This is because the society can take cost-effective measures such as handwashing to achieve a small decrease in  $R_t$ . If the society must decrease  $R_t$  more, it must take costlier measures. Thus, we can set the following conditions on the intervention cost function  $C(R_t)$  ( $0 < R_t \leq R_N$ ), where an example is shown in Figure 1.

$$C(R_t) \text{ is twice continuously differentiable,} \tag{1}$$

$$C(R_N) = 0, \tag{2}$$

$$\frac{dC(R_t)}{dR_t} \leq 0, \tag{3}$$

$$\frac{d^2C(R_t)}{dR_t^2} \geq 0. \tag{4}$$

The measure taken by spending the intervention cost,  $C(R)$ , is to decrease infected population (the number of infected persons who are capable of transmitting infections),  $I$ . The more infected the population decreases for fixed intervention costs, the more the society benefits from the measure. The benefit of a decrease in the infected population is evaluated as the decrease in the cost of the infected population. We set this “infection cost”  $M$  to be proportional to the infected population,  $I$ , which includes medical cost and patients’ losses incurred by being infected. This yields

$$M(t) = c_1 I(t), \tag{5}$$

where  $c_1$  is a constant. This assumption is also the same as Rowthorn<sup>2)</sup>. The total cost per unit time is the sum of the intervention cost and the infection cost, namely  $C(t) + M(t)$ . The optimization issue is to find  $R(t)$  that minimizes the integrated total cost over a certain period:

$$\int [C(t) + M(t)] dt. \tag{6}$$

To find the optimized intervention process specified by a protocol of  $R(t)$  for a targeted period, we must take into account the dynamics of the infected population. Here, we utilize the SIR model proposed by Kermack and McKendrick<sup>9)</sup>. It is the simplest fundamental model that describes the basic dynamics of epidemics. It models the exponential growth of the infected population in the outbreak stage, the peak of the infected population, and transitioning to the end stage<sup>13)</sup>.

## **SIR Model**

The SIR model is a set of differential equations that describes epidemic disease propagation, in which the population is divided into three states:  $S(t)$ , the population ratio of susceptible

persons,  $I(t)$ , the ratio of infected persons, and  $\hat{R}_{\text{rec}}(t)$ , the ratio of those who have recovered (or died). This formulation considers a closed population that is conserved. Note that we use the notation  $\hat{R}_{\text{rec}}$  for recovered persons, instead of the conventional notation,  $R$ , because we use  $R_t$  for the average reproduction number.

$$\frac{dS(t)}{dt} = -\beta S(t)I(t), \quad (7)$$

$$\frac{dI(t)}{dt} = \beta S(t)I(t) - \gamma I(t), \quad (8)$$

$$\frac{d\hat{R}_{\text{rec}}(t)}{dt} = \gamma I(t), \quad (9)$$

where  $\beta$  and  $\gamma$  are the infection and recovery rates, respectively. The sum of the three population ratios remains constant:

$$S(t) + I(t) + \hat{R}_{\text{rec}}(t) = 1. \quad (10)$$

In the following, we evaluate the infected population,  $I(t)$ . Equation (8) leads to

$$\frac{dI(t)}{dt} = \gamma \left[ \frac{\beta S(t)}{\gamma} - 1 \right] I(t). \quad (11)$$

We restrict ourselves to the period before the vicinity of the infection peak, in which  $S(t)$  is well replaced by  $S(0)$ . This approximation is accurate in major parts of the first outbreak and its recurrent phases<sup>14)</sup>, as shown in the figure in the Extended Data. Then, Eq. (11) leads,

$$\frac{dI(t)}{dt} = \gamma \left[ \frac{\beta S(0)}{\gamma} - 1 \right] I(t). \quad (12)$$

We restrict ourselves to  $\gamma$  being fixed, like Rowthorn<sup>2)</sup>. If the set of parameters  $\frac{\beta S(0)}{\gamma} > 1$ , the

infections start spreading in Eq. (12)<sup>15</sup>). The change in  $\beta$  in  $\frac{\beta S(0)}{\gamma}$  changes the dynamics of the pandemic. The set of parameters is the effective reproduction number:

$$R_t = \frac{\beta S(0)}{\gamma}, \quad (13)$$

where  $R_t$  corresponds to the basic reproduction number,  $R_0$ , if the following two assumptions are satisfied: 1)  $\beta$  has an uncontrolled value and 2)  $S(0) = 1$ . The infected population increases when  $R_t > 1$  and decreases for  $R_t < 1$ . Equation 12 becomes, with  $\Delta_R = R_t - 1 (> 0)$ ,

$$\frac{dI(t)}{dt} = \gamma \Delta_R I(t). \quad (14)$$

At  $R_t = 1$ , the infected population is stationary, as  $\Delta_R = 0$ . In this formulation, the infected population,  $I(t)$ , is the only variable that describes the state of the system. In the following sections, we will show the universal properties of the system by analysing the cyclic process of the state variable  $I(t)$ .

## Irreversible cost in on/off-type intervention process

We next evaluate the costs of on/off-type infection control (see Figure 2) and compare it with the costs of keeping the infected population stationary, where we assume that both processes have the same average effective reproduction number,  $\langle R_t \rangle = 1$ . The present on/off-type intervention forms a cycle of both  $R_t$  and  $I(t)$ , as shown in the following, in which a set of lockdown and recurrence is the extreme example. We set the amplitude of the cycle in the effective reproduction number around  $R_t = 1$  as “ $\Delta$ ”, where  $\Delta = |R_t - 1|$ . The cyclic process (with time interval  $T$ ) is as follows:

Stage 1)  $0 < t < T$ :  $I_0 \rightarrow I_1 (> I_0)$  with  $R_t = 1 + \Delta$ ,

Stage 2)  $T < t < 2T$ :  $I_1 \rightarrow I_0$  with  $R_t = 1 - \Delta$ ,

Stage 3)  $2T < t < 3T$ :  $I_0 \rightarrow I_3 (< I_0)$  with  $R_t = 1 - \Delta$ ,

Stage 4)  $3T < t < 4T$ :  $I_3 \rightarrow I_0$  with  $R_t = 1 + \Delta$ .

In the second and fourth steps, the infected population returns to the original state,  $I_0$ , as shown in *Methods*.

From the straightforward analytical calculation in *Methods* (Analytical calculation I), we obtain the average infected population during the cycle (Figure 2):

$$\frac{1}{4T} \int_0^{4T} I(t) dt = \frac{I_0}{\gamma \Delta T} \sinh(\gamma \Delta T) = I_0 + \frac{I_0 (\gamma \Delta T)^2}{3!} + O((\gamma \Delta T)^4). \quad (15)$$

The stationary infected population at  $R_t = 1$  during the same period,  $4T$ , is  $I_0$ . This proves that the average infected population in this cycle is always higher than that of the stationary state. This results directly yields through Eq.(5) :

$$\langle M \rangle_{\text{cycle}} > \langle M \rangle_{R_t=1}, \quad (16)$$

where  $\langle M \rangle$  denotes the time-average of  $M$ . Thus, the average infection cost for this cycle is higher than that of the stationary state. Figure 3 shows how the average infection cost depends on the amplitude of the cycle  $\Delta$ .

Next, we calculate the average intervention cost over the cycle. The average intervention cost, weighting the two effective reproduction numbers,  $R_t = 1 + \Delta$  and  $R_t = 1 - \Delta$  equally ( $\Delta > 0$ ) for the same period is

$$\langle C(R_t) \rangle_{\text{cycle}} = \frac{C(1 + \Delta) + C(1 - \Delta)}{2}. \quad (17)$$



From the straightforward calculation in *Methods* (Analytical calculation II), we obtain

$$\langle C(R_t) \rangle_{\text{cycle}} = \frac{C(1 + \Delta) + C(1 - \Delta)}{2} > C(1). \quad (18)$$

Thus, we find that the average intervention cost,  $\langle C(R_t) \rangle$ , is also higher in this cycle than keeping a stationary state with  $R_t = 1$ . Figure 4 illustrates how the intervention cost depends on the amplitude of the cycle  $\Delta$ , where we use the model in Fig. 1.

The results show that the cycle of infection control around the stationary state provokes an excess average infected population,  $\langle I(t) \rangle$ , and also, excess intervention cost compared to that at stationary state. Because the variable of the state,  $I(t)$ , finally returns to the initial state in the cycle, the cycle above results in waste of social resource compared to a stationary state. The economic irreversibility that society cannot retrieve the dissipated social resource is similar to entropy production (or free energy decreases) in thermodynamics<sup>16</sup>).

The cost,  $C(R_t) + M(t)$ , for the cycle thus satisfies the inequality

$$\text{Average cost of the cyclic process} > \text{Average cost of the stationary process} \quad (19)$$

even if the two processes have the same average effective reproduction number  $\langle R_t \rangle = 1$ .

We have learned that society cannot produce extra benefits (decrease of infected population) in the cyclic process compared to keeping the infected population constant, while it pays extra intervention cost in the cycle. Not only that, but society also incurs the demerit (increase of infected population) in the cycle. Note that this inequality holds irrespective of specific parameters, which contrasts with previous studies on the economic efficiency of infection control. This inequality clearly illustrates how on/off-type infection control against pandemic costs to society.

## Irreversible Cost for Delaying Measures

Now, we will show the implication of economic irreversibility by the effect of delaying measures against pandemics. We compare the two processes having the same initial and final states,  $I_0$ . Only the intermediate states are different between the two processes.

Process 1) Do not perform infection control initially or perform small intervention at  $t = 0$  with  $R_t = R_a$ , in which  $1 < R_a \leq R_N$ , until some critical time ( $t = t_a$ ) just before serious problems such as the crash of medical capacity arise. Then, infection control is performed at  $t = t_a$  to achieve a constant  $R_t < 1$  to decrease  $I(t)$  back to  $I_0$ . This process is similar to the combined process of Stages 1 and 2 in Figure 2. However, the choice of  $R(t)$  before and after  $t = t_a$  is arbitrary.

Process 2) Perform infection control to achieve  $R_t = 1$  immediately at  $t = 0$ .

Here, we assume  $R_N > 1$  for both processes.

Process 1 has the “benefit” that there is no or small intervention cost,  $C(R_a) < C(1)$ , between  $t = 0$  and  $t = t_a$ . Compared with the decision to initially take measure  $R_t = 1$  (Process 2), this saves intervention cost between  $t = 0$  and  $t_a$ :

$$\int_0^{t_a} [C(1) - C(R_a)] dt. \quad (20)$$

Thus, it is the matter whether the saving of the intervention cost (Eq. (20) at  $t = t_a$ ) remains positive even at the final stage,  $t = t_b$ , when the state returns to its initial state,  $I_0$ . Thus, we calculate the average intervention cost of Process 1,  $\langle C(R_t) \rangle_{\text{delay}}$ , during the period from  $t = 0$  to  $t = b$ .

From straightforward calculation in *Methods* (Calculation III), we obtain

$$\langle C(R_t) \rangle_{\text{delay}} > C(1). \quad (21)$$

The right-hand side is the average intervention cost of Process 2. The average intervention cost,  $\langle C(R_t) \rangle_{\text{delay}}$  in the delaying measure is found to be higher than that for a stationary infection state. The inequality has universality because Eq. (21) holds for any process with linear functions with parameters  $\Delta_a$  and  $\Delta_b$ . Furthermore, because any integrable function can be decomposed into a set of linear functions with arbitrary precision, Eq. (21) holds even for any process of integrable  $R(t)$  on the condition that the variable of state  $I(t)$  returns to its initial state.

Apparently, infection cost satisfies the similar inequality as above,

$$\langle M(I(R_t)) \rangle_{\text{delay}} > M(I(1)), \quad (22)$$

as the average infected population is higher in the delaying measure than in a stationary infection population with  $R_t = 1$ . The results show that society with a delaying measure must pay more intervention and infection costs in the process until the state  $I(t)$  returns to its original state even if it temporarily saves intervention cost. In other words, once the infected population increases, the society cannot return to the previous lower infection state without paying extra cost than keeping a stationary state, as in Process 2. An increase of infected population always results in this economic irreversibility in pandemics, except for the vicinity of the infection peak. The universal result of the model is again independent of the details of the system.

## Discussion and Conclusion

This study theoretically analysed in infection-spreading phase the fundamental structure of economic irreversibility in infection control process. Delaying measures against the spread of infection results in increases in costs. Once the state variable  $I(t)$  is increased, the system is irreversible in the sense that it cannot return to the previous low-infection state without spending extra cost, compared to keeping the stationary state of the low-infection. These general results contradict the naive idea that infection control always results in economic damage.

The merit of keeping the infection population constant has been previously discussed by Rowthorn<sup>2)</sup>, saying, “The most robust conclusion is that, if a relatively inexpensive way can be found to reduce the net reproduction ratio to  $r = 1$ , that is, the policy to aim for in the medium term”. This numerical finding is consistent with our analytical result. It should be noted that the present result cannot show the level to which the society should decrease the infected population. Additionally, our analysis is restricted to a principal part of the pandemic, namely, the infection-spreading phase. These are limitations of our study.

The validity of the present study is subject to assumptions of the methodology. In addition to the conventional methodological assumptions of mean field theory<sup>17)</sup> and of constant rates<sup>9)</sup>, as in the SIR model, we used the following assumptions:

1. The intervention cost depends on the effective reproduction number,  $R_t$ , and its cost function  $C(R_t)$  is concave up as in Eq.(4).
2. Infection cost is proportional to the infected population,  $I(t)$ .
3. Without infection control, the epidemic is in the infection-spreading phase, and increases and decreases in the infected population obey exponential dynamics as in Eq. (14).

The first assumption is the same as the assumption of previous research<sup>2, 4)</sup>, through the relation

$R_t = R_N(1 - q(t))$ . The assumptions set out in Eqs. (1), (2), (3), and (4) are mathematically simple, as Assumption 2. The third assumption is justified when the effective reproduction number,  $R_t$ , describes the system well, and when the infection-spreading phase is expected to last for more period than the period of variation of the infected population.

Our study itself does not offer concrete cost values as conventional cost-benefit analyses do. However, the present result reveals the universal structure of the costs, which is independent of the coefficients of the cost, for example,  $c_1$ , of Eq. (5). The universality found in this study is similar to thermodynamics in physics<sup>18)</sup>. The theory of thermodynamics itself does not give the physical quantity of a system by itself, as in the present study. However, it provides a quantitative relationship among physical variables and shows physical irreversibility, the latter of which is similar to the present result that an increase in the infected population is economically irreversible.

Irreversibility of thermodynamics is caused by the deviation from thermal equilibrium. Carnot's cycle is known as a reversible thermodynamics process, which converts thermal energy into mechanical energy at maximum efficiency<sup>18)</sup>, which is called Carnot's efficiency and analogous to CBA in the sense that CBA evaluates the efficiency of conversion from social intervention cost into benefit (decrease in infected population, in the present case). Optimal energy conversion is available in the Carnot's cycle, as the cycle is at equilibrium and thus there is no entropy production. In non-equilibrium stationary state, it requires a finite cost to keep the system stationary<sup>19, 20)</sup>, in which the efficiency of energy conversion is different from that at equilibrium. However, even if the system is out of equilibrium, the efficiency of energy conversion<sup>21)</sup> and an equality on irreversibly work<sup>22)</sup> can be analytically discussed with thermodynamics and statistical mechanics. The present system corresponds to non-equilibrium even in the stationary state of a constant infected population, because the stationarity is maintained by spending the infection control cost with  $C(R_t = 1) > 0$  to inhibit an increase in the infected

population. Therefore, the application of concepts and methodology of non-equilibrium thermodynamics into CBA would be interesting<sup>23)</sup>, as economic irreversibility<sup>24, 25, 26)</sup> exists and has universality, as shown here.

Our analysis of the conventional pandemic model in the infection-spreading phase explicitly showed that once the infected population increases, the increased state is economically irreversible. To the best of our knowledge, this is the first analytical study of economic efficiency in pandemic control. The result may provide guiding principles for infection control in pandemics as thermodynamics gives several guiding principles for nature and industries. The following question has not yet been clarified by our model, ‘To which level we should decrease infected population?’ The question includes whether we should aim at the eradication of infection. Analytical studies to find conditions that determine the most effective infection control are important and challenging for the future.


## References

- [1] Bloom, D.E., & Canning, D. Epidemics and Economics, *PGDA Working Paper No. 9* <http://www.hsph.harvard.edu/pgda/working.htm>, Sch. of Pub. Health, Harvard University (2006).
- [2] Rowthorn, R. A cost-benefit analysis of the COVID-19 disease *Covid Economics*, Issue 9, 97-106 (2020).
- [3] For example, articles at *Covid Economics*, *Centre for Economic Policy Research* (Vetted and Real-Time Papers): <https://cepr.org/content/covid-economics-vetted-and-real-time-papers-0> (2020).


- [4] Rowthorn, R. & Maciejowski, J.A. Cost-benefit analysis of the Covid-19 disease, *Oxford Rev. Econ. Policy*, <https://doi.org/10.17863/CAM.54369> (2020).
- [5] Eichenbaum, M.S., Rebelo, S. & Trabandt, M. The Macroeconomics of Epidemics, Preprint at <https://www.kellogg.northwestern.edu/faculty/rebelo/htm/epidemics.pdf> (2020).
- [6] Acemoglu, D., Chernozhokov, V., Werning, I., Whinston, M.D. *Working Paper 27102*, (National Bureau of Economic Research, 2020).
- [7] Boardman, A.E., Greenberg, D.H., Vining, A.R., & Weimer, D.L. *Cost-benefit analysis: concept and practice (2nd Ed.)* (Prentice Hall, 2001).
- [8] Nas, T. F. *Cost-Benefit Analysis: Theory and Application*, (Sage Publication, 1996).
- [9] Kermack, W.O. & McKendrick, A.G. A contribution to the mathematical theory of epidemics, *Proc. Royal Soc. A*, **115**, 700-721 (1927).
- [10] NHK News, Chief Cabinet Secretary asked governors to keep hospitalization facilities for infected citizens with mild symptoms, *Japan Broadcasting Corporation*, July 28 (in Japanese), <https://www.nhk.or.jp/politics/articles/statement/42463.html> (2020).
- [11] To, K. K-W., Hung, I.F-N. *et al.* COVID-19 re-infection by a phylogenetically distinct SARS-coronavirus-2 strain confirmed by whole genome sequencing, *Clin. Infect. Dis.*, <https://doi.org/10.1093/cid/ciaa1275> (2020).
- [12] Tillett, R., Sevinsky, J. *et al.* Genomic evidence for a case of reinfection with SARS-CoV-2, *SSRN*, <https://dx.doi.org/10.2139/ssrn.3680955> (2020).
- [13] Martcheva, M. *An Introduction to Mathematical Epidemiology* (Springer-Verlag 2010).

- [14] Vynnycky, E. & White, R.G. *An Introduction to Infectious Disease Modeling* (Oxford University Press, 2010).
- [15] Earn, D.J.D. A light introduction to modelling recurrent epidemics, in *Mathematical Epidemiology* (eds F. Brauer, P. Driessche, J. Wu) 3-17 (Springer-Verlag, 2008).
- [16] Nicolis, G. & Prigogine, I. *Exploring Complexity* (R. Piper GmbH & Co., 1989).
- [17] Giesecke, J. *Modern Infection Disease Epidemiology (3rd Ed.)*, (Taylor & Francis, 2017)
- [18] Callen, H. *Thermodynamics* (John Wiley & Sons, 1960).
- [19] Hondou, T. & Sekimoto, K. Unattainability of Carnot efficiency in Brownian heat engine, *Phys. Rev. E*, **62**, 6021-6025 (2000).
- [20] Hondou, T. & Takagi, F. Irreversible operation in a Stalled state of Feynman's ratchet, *J. Phys. Soc. Jpn.*, **67**, 2974-2976 (1998).
- [21] Kamegawa, H. & Hondou, T. & Takagi, F. Energetics of a forced thermal ratchet, *Phys. Rev. Lett.*, **80**, 5251-5254 (1998).
- [22] Jarzynski, C. Nonequilibrium equality for free energy differences, *Phys. Rev. Lett.*, **78**, 2690-2693 (1997).
- [23] Tsirlin, A.M. Irreversible microeconomics: Optimal processes and control, *Autom. Remote. Control.*, **62**, 820-830 (2001).
- [24] Martinas K. About Irreversibility in Economics, *Open Sys. Inf. Dyn.*, **7**, 329-364 (2000).
- [25] Amir, S. The role of thermodynamics in the study of economic and ecological systems, *Ecol. Econ.*, **10**, 125-142 (1994).
- [26] Ayres, R.U. & Nair I. Thermodynamics and economics, *Phys. Today*, **37**, 62-71 (1984).






**Fig. 1.** An example of intervention cost,  $C$ .  
Here  $C(R_t) = (2 - R_t)/R_t$ , where  $R_N = 2$ .




**Fig. 2.** Trace of infected population during the cyclic process of infection control.

It is shown that the infected population is also cyclic, and returns to the initial state at the end of the cycle. The average infected population  $\langle I(t) \rangle$  over the cycle is larger than that for keeping the infected population stationary.



**Fig. 3.** Large oscillation of intervention results in large infections cost.

The average infection cost,  $\langle M(I(t)) \rangle$ , increases monotonically and exponentially as the amplitude of  $R_t$  in the cycle,  $\Delta$ , increases. The vertical axis is normalized by the average infection cost for the stationary state with  $R_t = 1$ , having an average effective reproduction number equal to that of the cycle. As the state variable  $I(t)$  returns to its initial state in the cycle, the increase in average infection cost is irreversible.



**Fig. 4.** Large oscillation of intervention also results in large intervention cost.

The average intervention cost,  $\langle C(R_t) \rangle$ , increases exponentially as the amplitude of  $R_t$  in the cycle,  $\Delta$ , increases. The vertical axis is normalized by the average intervention cost for the stationary state with  $R_t = 1$ , having an average effective reproduction number equal to that of the cycle. We use  $R_N = 2$  and  $C(R_t)$  of Figure 1. The increase in average intervention cost in the cycle does not contribute to the benefit (decrease in average infection cost) at all, as Figure 3 shows.

# Methods

## Analytical calculation I

By integrating Eq. (14) from  $t = 0$  to  $T$  with  $R_t = 1 + \Delta$ , we obtain the infected population  $I$  at the end of Stage 1,

$$I(T) = I_0 e^{\gamma T \Delta}. \quad (23)$$

Similarly, replacing  $\Delta_R$  in Eq. (14) by “ $-\Delta$ ” and using Eq. (23), we obtain  $I(2T)$  at the end of Stage 2:

$$I(2T) = I_0 \quad (24)$$

Stages 3 and 4 also yield

$$I(4T) = I_0. \quad (25)$$

We have confirmed that Stages 1 through 4 form a typical cyclic process of state variable,  $I(t)$ , around a stationary state kept by  $R_t = 1$ , where the infected population returns to its original value.

We calculated the average infected population to evaluate the infection cost in the cycle. Using Eqs. (14) and (23), we have, for Stages 1 and 2,

$$\int_0^T I_{\text{Stage1}}(t) dt + \int_T^{2T} I_{\text{Stage2}}(t) dt = I_0 \left[ \int_0^T e^{\gamma \Delta t} dt + \int_T^{2T} e^{\gamma \Delta T} e^{-\gamma \Delta (t-T)} dt \right] = I_0 \int_0^T [e^{\gamma \Delta t} + e^{\gamma \Delta (T-t)}] dt. \quad (26)$$

Similarly, for Stages 3 and 4, we have

$$\int_{2T}^{3T} I_{\text{Stage3}}(t) dt + \int_{3T}^{4T} I_{\text{Stage4}}(t) dt = I_0 \int_0^T [e^{-\gamma \Delta t} + e^{\gamma \Delta (t-T)}] dt. \quad (27)$$

Thus, we obtain,

$$\frac{1}{4T} \int_0^{4T} I(t) dt = \frac{I_0}{\gamma \Delta T} \sinh(\gamma \Delta T) = I_0 + \frac{I_0(\gamma \Delta T)^2}{3!} + O((\gamma \Delta T)^4). \quad (28)$$

## Analytical calculation II

The cost  $C(1 + \Delta)$  is evaluated as follows:

$$C(1 + \Delta) = C(1) + \int_1^{1+\Delta} \frac{dC(R_t)}{dR_t} dR_t, \quad (29)$$

From Eq. (4) we find

$$\frac{dC(R_t)}{dR_t} > \left. \frac{dC(R_t)}{dR_t} \right|_{R_t=1} \quad (\text{for } 1 < R_t \leq R_N). \quad (30)$$

Then, we have

$$C(1 + \Delta) > C(1) + \left. \frac{dC(R_t)}{dR_t} \right|_{R_t=1} \Delta. \quad (31)$$

Since  $\frac{dC(R_t)}{dR_t} < \left. \frac{dC(R_t)}{dR_t} \right|_{R_t=1}$  for  $0 < R_t < 1$ ,

$$C(1 - \Delta) > C(1) - \left. \frac{dC(R_t)}{dR_t} \right|_{R_t=1} \Delta. \quad (32)$$

We obtain through Eqs. (31) and (32)

$$\langle C(R_t) \rangle_{\text{cycle}} = \frac{C(1 + \Delta) + C(1 - \Delta)}{2} > C(1). \quad (33)$$

### Analytical calculation III

From Eq. (14), the state of  $I(t)$  at  $t = t_a$  is  $I(t_a) = I_0 e^{\gamma t_a \Delta_a}$ , where  $\Delta_a = R_a - 1$ . We assume that  $I(t)$  returns to  $I_0$  at  $t = t_a + t_b$ , and  $R_t = R_b = 1 - \Delta_b$  ( $0 < \Delta_b < 1$ ) for  $t_a < t \leq t_b$ . Then, we have  $I(t_a + t_b) = I(t_a) e^{-\gamma t_b \Delta_b}$ . As  $I(t_a + t_b) = I_0$ , we obtained the equality

$$t_a \Delta_a = t_b \Delta_b. \quad (34)$$

Then, the average intervention cost between  $t = 0$  and  $t = t_b$  is written as

$$\langle C(R_t) \rangle_{\text{delay}} = \frac{t_a}{t_a + t_b} C(1 + \Delta_a) + \frac{t_b}{t_a + t_b} C(1 - \Delta_b). \quad (35)$$

From Eqs. (31) and (32), Eq. (35) satisfies the following condition:

$$\langle C(R_t) \rangle_{\text{delay}} > \frac{t_a}{t_a + t_b} \left[ C(1) + \left. \frac{dC}{dR_t} \right|_{R_t=1} \Delta_a \right] + \frac{t_b}{t_a + t_b} \left[ C(1) - \left. \frac{dC}{dR_t} \right|_{R_t=1} \Delta_b \right]. \quad (36)$$

Using Eq. (34), the right-hand side of Eq. (36) equals  $C(1)$ .

### Data Availability

Data sharing is not applicable to this article as no datasets were generated or analysed during the current study. All figures except for the lower curve in Extended Figure 1 are generated by the explicit solutions shown in the text. The lower curve in Extended Figure 1 is generated by numerical simulation (Euler method). As the initial condition is fully described in Extended figure legend, the figure is also reproducible.

## **End notes**

### **Acknowledgments**

The author acknowledges T. Onai, S. Yonemura, K. Hirata, and S. Munakata for their fruitful discussions, S. Takagi, R. Seto, M. Arikawa, M. Sano, H. Chaté and M. Wallis for critical reading of the manuscript and helpful comments. We would like to thank Editage for English language editing. This work was supported by the Japanese Grants-in-Aid JSPS Nos. 16H01820 and 20H00002.

### **Authors contributions**

T.H. solely contributed to this study.

### **Competing interests**


The author has no completing interest.

### **Corresponding author**

Tsuyoshi Hondou, [hondou@mail.sci.tohoku.ac.jp](mailto:hondou@mail.sci.tohoku.ac.jp)



## Extended Data



### Extended Figure 1:

The validity of theoretical approximation leading to Equation 12 is shown.

We replaced the variable of susceptible persons,  $S(t)$ , in Eq. (11) with respect to the initial value  $S(0)$  (Eq.(12)) for the evaluation of the infected population in the infection-spreading phase because  $S(t)$  is a slow variable. The upper line is a solution of Equation 12. The lower line is a solution of Equation 11, which is a simple solution of the SIR model without approximation. The figure illustrates that the approximation over  $S(t)$  is precise in the infection-spreading phase, except for the vicinity of the infection peak. Here, we used  $\beta = 0.51$  and  $\gamma = 0.204$ , which corresponds to the basic reproduction number  $R_0 = 2.5$  for demonstration. Numerical calculation is performed using the Euler method, in which the initial values are as follows: Total population  $N = 1.2 \times 10^8 + 1$ ,  $S(0) = 1.2 \times 10^8 / N$ ,  $I(0) = 1/N$ ,  $\hat{R}_{\text{rec}}(0) = 0/N$ .