ISOMORPHIC INVERSE PROBLEMS

EVGENY L. KOROTYAEV

ABSTRACT. Consider two inverse problems for Sturm-Liouville problems on the unit interval. It means that there are two corresponding mappings F, f from a Hilbert space of potentials H into their spectral data. They are called isomorphic if F is a composition of f and some isomorphism U of H onto itself. A isomorphic class is a collection of inverse problems isomorphic to each other. We consider basic Sturm-Liouville problems on the unit interval and on the circle and describe their isomorphic classes of inverse problems. For example, we prove that the inverse problems for the case of Dirichlet and Neumann boundary conditions are isomorphic. The proof is based on the non-linear analysis.

1. Introduction and main results

- 1.1. **Introduction.** We consider the Sturm-Liouville problems $-y'' + qy = \lambda y$ on the unit interval [0,1] under basic boundary conditions or on the unit circle \mathbb{S}^1 . Here the potential q is real and belongs to the space $L^2(0,1)$. There are a lot of results about the inverse problems for the Sturm-Liouville operators on [0,1] and on \mathbb{S}^1 . These inverse problems were investigated by many authors (G. Borg, I. M. Gel'fand, B. M. Levitan, V. A. Marchenko, E. Trubowitz, ..), see the monographs [29], [33], [34] and references therein. In general, the study of an inverse spectral problem consists of the following parts:
- (i) Uniqueness: prove that the spectral data (eigenvalues plus some additional parameters) determine the potential uniquely);
- (ii) Reconstruction: reconstruct the potential from spectral data;
- (iii) Characterization: describe all spectral data corresponding to fixed classes of potentials.
- (iv) Stability estimates: obtain a priori two sided estimates of the potential and spectral data. We will discuss their additional *isomorphic* properties.

Definition.Let f and f_o be mappings from a Hilbert space K to a set X. They are called isomorphic if $f_o = f \circ U$ for some isomorphism (in general, non-linear) U of K onto itself.

Note that if some of them is a bijection, then U is a unique canonical automorphism of \mathcal{K} .

Consider two inverse problems for Sturm-Liouville problems, when potentials q belong to a corresponding Hilbert space \mathcal{K} . Thus there are two mappings $f: \mathcal{K} \to X$ and $f_o: \mathcal{K} \to X$, where X is their set of spectral data. They are called isomorphic if $f_o = f \circ U$ for some isomorphism (in general, it is non-linear) U of \mathcal{K} onto itself. If U is an unitary linear operator, then these two inverse problems are called unitarily equivalent.

Note that if two Sturm-Liouville inverse problems are isomorphic, then we have

1) If the first has some property from (i)-(iv), then the second also has it. For example, the first has uniqueness iff the second has uniqueness.

Date: January 13, 2023.

²⁰²⁰ Mathematics Subject Classification. 34A55 (34B24 47E05 47N50 81Q10).

Key words and phrases. inverse problem, eigenvalues, Sturm-Liouville problem.

- 2) Eigenvalues of the first problem have some asymptotics for each potential iff eigenvalues of the second problem have similar asymptotics.
- 3) The first problem has some trace formula iff the second problem has a similar trace formula.

We write our main results:

- A) We describe all isomorphic Sturm-Liouville inverse problems on the unit interval under the basic boundary conditions and on the circle. The corresponding automorphism U is obtained in explicit form.
- B) The same is made for potentials from Sobolev spaces. The needed new sharp asymptotics of norming constants are determined.

To the best of our knowledge the obtained results have no analogies in existing literature. Our proof uses observations 1)-3) and also the following results and methods:

- two spectra mapping (Marchenko-Ostrovski [32]),
- the Dirichlet eigenvalues and norming constants mapping (Pöschel-Trubowitz [34]),
- the four spectra mapping (Korotyaev [18]),
- results of Marchenko-Ostrovski [32] and Korotyaev [22] about inverse periodic problems.

We consider four Sturm-Liouville problems on the interval [0,1] with the Dirichlet and Neumann boundary conditions:

$$-f'' + qf = \lambda f, \qquad f(0) = f(1) = 0, f'(0) = f'(1) = 0,$$
(1.1)

and with the so-called mixed boundary conditions:

$$-f'' + qf = \lambda f, \qquad f(0) = f'(1) = 0, f'(0) = f(1) = 0,$$
 (1.2)

where $\lambda \in \mathbb{C}$. Here the potential q belongs to the real Hilbert space \mathcal{L} defined by

$$\mathcal{L} = \left\{ q \in L^2([0,1], \mathbb{R}) : \int_0^1 q dx = 0 \right\}$$

equipped with the norm $||q||^2 = \int_0^1 q^2(x) dx$. There are a lot of results about these problems, see, e.g., the books [29], [33], [34]. Let μ_n and $\nu_0, \nu_n, n \ge 1$ be eigenvalues of the Dirichlet and Neumann problems respectively. Let τ_n , and $\varrho_n, n \ge 1$ be eigenvalues of the first and the second problem respectively with mixed boundary conditions (1.2), and we say shortly mixed eigenvalues. All these eigenvalues are simple and satisfy

$$\nu_0 < \overline{\tau_1, \varrho_1} < \overline{\mu_1, \nu_1} < \overline{\tau_2, \varrho_2} < \overline{\mu_2, \nu_2} < \dots,
\nu_n, \mu_n = \mu_n^o + o(1), \quad \tau_n, \varrho_n = \tau_n^o + o(1) \quad \text{as } n \to \infty,$$
(1.3)

where $\overline{u,v}$ denotes $\min\{u,v\} \leqslant \max\{u,v\}$ for shortness, and $\nu_0^o = 0, \nu_n^o = \mu_n^o = (\pi n)^2$ and $\tau_n^o = \varrho_n^o = \pi^2 (n - \frac{1}{2})^2, n \geqslant 1$ are the corresponding unperturbed eigenvalues.

We introduce the fundamental solutions $\varphi(x,\lambda)$, $\vartheta(x,\lambda)$ of the equation

$$-f'' + q(x)f = \lambda f, \qquad \lambda \in \mathbb{C},$$

under conditions: $\varphi'(0,\lambda) = \vartheta(0,\lambda) = 1$ and $\varphi(0,\lambda) = \vartheta'(0,\lambda) = 0$. Here and below $(') = \frac{\partial}{\partial x}$ and $(\dot{}) = \frac{\partial}{\partial \lambda}$. Note that $\{\mu_n\}, \{\nu_n\}, \{\tau_n\}$ and $\{\varrho_n\}$ are zeros of $\varphi(1,\lambda), \vartheta'(1,\lambda), \varphi'(1,\lambda)$ and

 $\vartheta(1,\lambda)$ respectively. Introduce the real Hilbert spaces $\ell_k^2 = \ell_k^2(\mathbb{N}), k \in \mathbb{R}$ of real sequences $v = (v_n)_1^{\infty}$ equipped with the norm

$$||v||_k^2 = \sum_{n\geq 1} (2\pi n)^{2k} v_n^2$$
, and let $\ell^2 = \ell_0^2$, $||\cdot|| = ||\cdot||_0$.

Following the book of Pöschel and Trubowitz [34] we define a set \mathfrak{J}^o of all real, strictly increasing sequences by

$$\mathfrak{J}^{o} = \left\{ s = (s_{n})_{1}^{\infty} : s_{1} < s_{2} < \dots, \quad s_{n} = \mu_{n}^{o} + \widetilde{s}_{n}, \quad \widetilde{s} = (\widetilde{s}_{n})_{1}^{\infty} \in \ell^{2} \right\}.$$

Note that the mapping $s \leftrightarrow \widetilde{s}$ is a natural coordinate map between \mathfrak{J}^o and some open convex subset $\widetilde{\mathfrak{J}}^o = \left\{ \widetilde{s} = (\widetilde{s}_n)_1^\infty \in \ell^2 : \mu_1^o + \widetilde{s}_1 < \mu_2^o + \widetilde{s}_2 < \right\}$ of ℓ^2 . Following [34] we identify \mathfrak{J}^o and $\widetilde{\mathfrak{J}}^o$ using this mapping. As in [34] this identification allows to do analysis on \mathfrak{J}^o as if it was an open convex subset of ℓ^2 . We also define similar sets \mathfrak{J}^1 and \mathfrak{J} of all real, strictly increasing sequences by

$$\mathfrak{J}^{1} = \left\{ t = (t_{n})_{1}^{\infty} : t_{1} < t_{2} < \dots, \quad t_{n} = \tau_{n}^{o} + \widetilde{t}_{n}, \qquad \widetilde{t} = (\widetilde{t}_{n})_{1}^{\infty} \in \ell^{2} \right\},$$

$$\mathfrak{J} = \left\{ t = (t_{n})_{1}^{\infty} : t_{1} < t_{2} < \dots, \quad t_{n} = (\frac{\pi n}{2})^{2} + \widetilde{t}_{n}, \qquad \widetilde{t} = (\widetilde{t}_{n})_{1}^{\infty} \in \ell^{2} \right\}.$$

Introduce 1-spectra mappings μ and ν from \mathcal{L} into \mathfrak{J}^o and τ and ϱ from \mathcal{L} into \mathfrak{J}^1 by

$$q \to \mu = (\mu_n)_1^{\infty}, \quad q \to \nu = (\nu_n)_1^{\infty},$$

$$q \to \tau = (\tau_n)_1^{\infty}, \quad q \to \varrho = (\varrho_n)_1^{\infty}.$$
(1.4)

For two 1-spectra mappings (only for strongly increasing and alternate) we construct standard 2-spectra mappings of strongly increasing sequences. For example, for $\tau = (\tau_n)_1^{\infty} \in \mathfrak{J}^1$ and $\mu = (\mu_n)_1^{\infty} \in \mathfrak{J}^o$ such that $\tau_1 < \mu_1 < \tau_2 < \mu_2 < \dots$ we define a 2-spectra mapping $\tau \star \mu$ from \mathcal{L} into \mathfrak{J} as

$$q \to \tau \star \mu = (\tau_1, \mu_1, \tau_2, \mu, \dots). \tag{1.5}$$

The mapping $\tau \star \mu$ is a bijection between \mathcal{L} and \mathfrak{J} . It is a classical result of Marchenko and Ostrovski [32].

Following Trubowitz and co-authors [12], [6] we introduce norming constants $h_{s,n}$, $\mathfrak{h}_{s,n}$ (associated with the Dirichlet and Neumann eigenvalues) and the corresponding mappings by

$$h_{s,n} = \ln |\varphi'(1,\mu_n)|, \qquad \mathfrak{h}_{s,n} = \ln |\vartheta(1,\nu_n)|, \qquad n \geqslant 1, \quad \mathfrak{h}_{s,0} = \ln |\vartheta(1,\nu_0)|,$$

$$q \to h_{(s)} = (h_{s,n})_{n=1}^{\infty}, \qquad q \to \mathfrak{h}_{(s)} = (\mathfrak{h}_{s,n})_{n=1}^{\infty}.$$

$$(1.6)$$

It is known that the mappings $\mu \times h_{(s)}$ is a bijections between \mathcal{L} and $\mathfrak{J}^o \times \ell_1^2$. This is a classical result of Pöschel and Trubowitz [34].

We sometimes write $\mu_n(q), \nu_n(q), ...$ instead of $\mu_n, \nu_n, ...$, when several potentials are being dealt with. We discuss a mapping $q \to \mu \times (D_n)_1^\infty$ introduce by Marchenko [31], where D_n is the normalizing constant associated with Dirichlet eigenvalue μ_n and defined by

$$D_n(q) = \int_0^1 \varphi^2(x, \mu_n(q), q) dx, \quad n \in \mathbb{N}, \quad \text{where } D_n(0) = \frac{1}{2\mu^2}.$$
 (1.7)

Marchenko [31] proved that the spectral data μ_n , D_n , $n \in \mathbb{N}$ determines the potential uniquely. Gel'fand and Levitan [9] created a basic method to reconstruct the potential q from μ_n , D_n , $n \in \mathbb{N}$: they determined an integral equation and expressed q in terms of its solution. It is

convenient to modify constants D_n and define another mapping $q \to \alpha = (\alpha_n)_1^{\infty}$, where the components α_n are given by

$$\alpha_n = \log \frac{D_n(q)}{D_n(0)} = \log \left[2\mu_n^o D_n(q) \right], \qquad n \in \mathbb{N}.$$
(1.8)

We discuss similar mapping $q \to \nu \times (N_n)_1^{\infty}$, where N_n is a normalizing constant, associated with Neumann eigenvalue ν_n and defined by

$$N_n(q) = \int_0^1 \vartheta^2(x, \nu_n, q) dx, \quad n \in \mathbb{N}, \quad \text{where } N_n(0) = \frac{1}{2}.$$
 (1.9)

It is more convenient to modify constants N_n and to define another mapping $q \to \beta = (\beta_n)_1^{\infty}$, where the components β_n are given by

$$\beta_n = \ln[2N_n(q)]. \tag{1.10}$$

Recall some definitions. We write $\mathcal{K}_{\mathbb{C}}$ for the complexification of the real Hilbert space \mathcal{K} . Suppose that \mathcal{K}, \mathcal{S} are real separable Hilbert spaces. The mapping $f: \mathcal{K} \to \mathcal{S}$ is a local real analytic isomorphism iff for any $y \in \mathcal{K}$ it has an analytic continuation \widetilde{f} into some complex neighborhood $V \subset \mathcal{K}_{\mathbb{C}}$ of y, which is a bijection between V and some open set $\widetilde{f}(V) \subset \mathcal{S}_{\mathbb{C}}$ and if \widetilde{f} , \widetilde{f}^{-1} are analytic mappings on V, $\widetilde{f}(V)$ respectively. The mapping f is a real-analytic bijection (shortly a RAB) between \mathcal{K} and \mathcal{S} if it is both a bijection and a local real analytic isomorphism.

1.2. Short review. We shortly discuss well-known results about inverse problems for Sturm-Liouville operators on the unit interval under different boundary conditions, which are used in our paper. We recall only some important steps mainly on the characterization problem. Marchenko and Ostrovski in [32] solved the inverse problem for the 2-spectra mapping $\tau \star \mu$. Their proof is based on the inverse scattering on the half line (with decreasing potentials) and sharp asymptotics for eigenvalues μ_n, τ_n .

Trubowitz and co-authors ([6], [12], [13], [34]) suggested an analytic approach. It is based on analyticity of the mapping {potentials} \mapsto {spectral data} and an explicit reconstruction procedure for the special case when only one spectral parameter has been changed. The excellent book [34] is devoted to the mapping $q \to \mu \star h_{(s)}$, where the inverse problem is solved, including the characterization of the spectral data. Also, this approach was applied to other inverse problems with purely discrete spectrum: for impedance [4], [5], [21]; singular Sturm-Liouville operators on a finite interval [11]; periodic potentials [8], [16], [22], perturbed harmonic oscillators [1], vector-valued operators [2], and Birkhoff coordinates for the KdV equation on the circle [15], and see references therein.

Now we discuss the periodic case. Note that only the periodic eigenvalues do not determine a potential uniquely and we need to add auxiliary spectral data: Dirichlet (or Neumann) eigenvalues plus sequence of signs \pm . There are only two mappings related to the characterization for inverse periodic problems:

- 1) in terms of local maxima and minima of the Lyapunov functions on the real line,
- 2) in terms of gaps.

Marchenko and Ostrovski [32] solved the inverse problems (including characterization and stability estimates) in terms of the local maxima and minima of Lyapunov functions on the real line. The proof is based on the inverse scattering on the half line (with decreasing potentials), sharp asymptotics of periodic eigenvalues and new results about conformal mappings for

the quasimomentum. A shorter proof was given in [24]. Korotyaev [22], [23] solved the inverse problems in terms of gap lengths. The proof (including characterization) was based on analyticity of the mapping and a priori estimate of potentials in terms of gap lengths from [23].

1.3. Main results on the unit interval. In order to discuss main results we introduce our basic transformations. We define the 4-spectra mapping $\mathfrak{f}:\mathcal{L}\to\ell^2$ from a recent paper [18] by

$$q \to \mathfrak{f}(q) = (\mathfrak{f}_n(q))_1^{\infty}, \quad \mathfrak{f}_{2n-1} = \varrho_n - \tau_n, \qquad \mathfrak{f}_{2n} = \nu_n - \mu_n, \qquad n \geqslant 1,$$
 (1.11)

which will be basic for us. Recall a result from [18]: the 4-spectra mapping $\mathfrak{f}: \mathcal{L} \to \ell^2$ defined by (1.11) is a RAB between \mathcal{L} and ℓ^2 , see more in Theorem 2.1. Let \mathfrak{S} be a set of all diagonal operators $\sigma = \operatorname{diag}(\sigma_1, \sigma_2, ...)$ on ℓ^2 , or shortly $\sigma = (\sigma_j)_1^{\infty}$, where $\sigma_j \in \{\pm 1\}, j \in \mathbb{N}$. This set \mathfrak{S} defines so-called the lamplighter group, see [3]. For each $\sigma \in \mathfrak{S}$ we define a mapping \mathcal{U}_{σ} by

$$\mathcal{U}_{\sigma} = \mathfrak{f}^{-1}\sigma\mathfrak{f}: \mathcal{L} \to \mathcal{L},\tag{1.12}$$

where \mathfrak{f} is given by (1.11). These mappings have the following properties:

Proposition 1.1. Let $\mathcal{U}_{\sigma}, \sigma \in \mathfrak{S}$ be defined by (1.12). Then each \mathcal{U}_{σ} is a RAB of \mathcal{L} onto itself and satisfies

$$\mathcal{U}_{\sigma} = \mathcal{U}_{\sigma}^{-1}, \qquad \mathcal{U}_{\sigma} \ \mathcal{U}_{\sigma'} = \mathcal{U}_{\sigma\sigma'} = \mathcal{U}_{\sigma'} \ \mathcal{U}_{\sigma}, \qquad \forall \quad \sigma, \sigma' \in \mathfrak{S},$$
 (1.13)

$$\|\mathcal{U}_{\sigma}(q)\| = \|q\| \qquad \forall \ q \in \mathcal{L}. \tag{1.14}$$

Remark. 1) The mapping \mathfrak{f} is non-linear, but \mathcal{U}_{σ} keeps the norm on \mathcal{L} , see (1.14).

2) To prove Proposition 1.1 we use the bijection of the mapping \mathfrak{f} from [18]. Its proof is based on the bijections of the mapping $\tau \star \mu$ and on inverse problems for periodic potentials [22].

Define two specific mappings, often used in our paper:

$$U_1 = \mathcal{U}_{\sigma}, \quad \text{where} \quad \sigma = (\sigma_j)_1^{\infty}, \ \sigma_j = -1 \quad \forall \ j \in \mathbb{N},$$

 $U_o = \mathcal{U}_{\sigma}, \quad \text{where} \quad \sigma = (\sigma_j)_1^{\infty}, \ \sigma_j = (-1)^j \ \forall \ j \in \mathbb{N}.$ (1.15)

Below we show that $U_o = \mathcal{R}$, where \mathcal{R} is a reflection (unitary) operator on \mathcal{L} given by $(\mathcal{R}y)(x) = y(1-x), x \in (0,1)$. For each $\mathfrak{n} = (\mathfrak{n}_n)_1^{\infty} \in \mathfrak{J}$ we define a potential \mathfrak{q} by

$$\mathfrak{q}(x) = -2\frac{d^2}{dx^2}\log\left(\Gamma_{\mathfrak{n}}\det\Omega(x,\mathfrak{n})\right), \quad x \in (0,1), \tag{1.16}$$

where $\Omega(x, \mathbf{n}), x \in (0, 1)$ is the infinite matrix whose elements $\Omega_{n,j}(x, \mathbf{n})$ are given by

$$\Omega_{n,j}(x,\mathfrak{n}) = \frac{\mathfrak{n}_n - \mathfrak{n}_n^o}{\mathfrak{n}_n - \mathfrak{n}_j^o} \left\{ \cos\sqrt{\mathfrak{n}_n} x + \frac{(-1)^n - \cos 2\sqrt{\mathfrak{n}_n}}{\sin 2\sqrt{\mathfrak{n}_n}} \sin\sqrt{\mathfrak{n}_n} x, \frac{\sin\frac{\pi j}{2}x}{\frac{\pi j}{2}} \right\}_w, \tag{1.17}$$

where $\{u, v\}_w = uv' - u'v$ and

$$\Gamma_{\mathfrak{n}} = \prod_{j>n\geqslant 1} \left(\frac{\mathfrak{n}_n - \mathfrak{n}_n^o}{\mathfrak{n}_n - \mathfrak{n}_j} \cdot \frac{\mathfrak{n}_n^o - \mathfrak{n}_j^o}{\mathfrak{n}_n^o - \mathfrak{n}_j^o} \right), \qquad \mathfrak{n}_n^o = \left(\frac{\pi n}{2} \right)^2.$$
 (1.18)

Here $\Omega - I$ is a trace class operator, so det $\Omega(x, p)$ is well defined. A recovering of a potential q by (1.16) via the 2-spectra mapping $\mathfrak{n} = \tau \star \mu$, $\mathfrak{n}_{2j-1} = \tau_j$, $\mathfrak{n}_{2j} = \mu_j, j \in \mathbb{N}$, was performed in [18] (see also [34] for even potentials). The first main result describes equivalent inverse problems on the unit interval.

Theorem 1.2. Let mappings U_0, U_1 be defined by (1.15). Then 1-spectra mappings $\mu = (\mu_n)_1^{\infty}, \nu = (\nu_n)_1^{\infty}, \text{ and } \tau = (\tau_n)_1^{\infty}, \varrho = (\varrho_n)_1^{\infty} \text{ defined by (1.4) satisfies}$

i) All 2-spectra mappings $\tau \star \mu$, $\varrho \star \nu$, $\varrho \star \mu$ and $\tau \star \nu$ acting from $\mathcal L$ into $\mathfrak J$ are isomorphic. Moreover, each of them is a RAB between $\mathcal L$ and $\mathfrak J$ and satisfies

$$\tau \star \mu = (\varrho \star \nu) \circ U_1 = (\varrho \star \mu) \circ U_o = (\tau \star \nu) \circ U_o \circ U_1. \tag{1.19}$$

- ii) Let $q \in \mathcal{L}$. Then a potential $U_1(q)$ is given by (1.16), where $\mathfrak{n} = \varrho \star \nu$.
- iii) Let $q \in \mathcal{L}$. Then a potential $(U_0U_1)(q)$ is given by (1.16), where $\mathfrak{n} = \tau \star \nu$.

Remark. 1) In the proof we show (1.19), then we obtain i), since the mapping $\tau \star \mu$ is a bijection between \mathcal{L} and \mathfrak{J} [32] and U_0, U_1 are bijections of \mathcal{L} onto itself.

2) We connect all 2-spectra mappings by the bijections U_o, U_1 and U_oU_1 .

We discuss the mapping $q \to \text{eigenvalues plus norming (or normalizing)}$ constants.

Theorem 1.3. i) Mappings $\mu \times h_{(s)}, \nu \times \mathfrak{h}_{(s)}, \mu \times \mathfrak{h}_{(s)}$ and $\nu \times h_{(s)}$ (defined by (1.4), (1.6)) are isomorphic as mappings from \mathcal{L} into $\mathfrak{J}^o \times \ell_1^2$ and satisfy

$$\mu \times h_{(s)} = (\nu \times \mathfrak{h}_{(s)}) \circ U_1 = (\mu \times \mathfrak{h}_{(s)}) \circ U_o = (\nu \times h_{(s)}) \circ U_o \circ U_1. \tag{1.20}$$

Moreover, each of them is a RAB between \mathcal{L} and $\mathfrak{J}^o \times \ell_1^2$.

ii) Each of the mappings $\mu \times \alpha$ and $\nu \times \beta$ (defined by (1.8), (1.10)) acting from \mathcal{L} into $\mathfrak{J}^o \times \ell_1^2$ is a bijection between \mathcal{L} and $\mathfrak{J}^o \times \ell_1^2$ and the following trace formula holds true

$$\frac{1}{N_0} - 1 = \sum_{n \ge 1} \left(2 - \frac{1}{N_n} \right),\tag{1.21}$$

where the series converges absolutely and uniformly on bounded subsets of \mathcal{L} .

iii) Define another normalizing mapping $q \to \hat{\beta} = (\hat{\beta}_n)_1^{\infty}$, where $\hat{\beta}_n = \beta_n - \ln \frac{\nu_n - \nu_0}{\nu_n^{\circ}}$. Then the mapping $\nu \times \hat{\beta}$ is a bijection between \mathcal{L} and $\mathfrak{J}^{\circ} \times \ell_1^2$ and satisfies

$$\mu \times \alpha = (\nu \times \hat{\beta}) \circ U_1. \tag{1.22}$$

Remark. 1) The mapping $\nu \times \mathfrak{h}_{(s)}$ is a bijection between \mathcal{L} and \mathfrak{J} , see [26]. The mapping $\nu \times \mathfrak{h}_{(s)}$ does not use the eigenvalue ν_0 and the norming constant $\mathfrak{h}_{s,0}$ since $\nu_n, \mathfrak{h}_{s,n}, n \geq 0$ are dependent, see trace formulas (2.3), (1.21).

2) In order to prove that the mappings in ii) are bijections we show, that they are isomorphic to the mapping $\mu \times h_{(s)}$, which is well studied in [34]. Thus the proof is short.

Final Remarks. Below we discuss also following isomorphic inverse problems:

- The case of mixed eigenvalues is considered in Sect. 4.
- The case of smooth potentials is considered in Sect. 5. The needed new sharp asymptotics for the norming constants for potentials from Sobolev space are obtained in Sect. 7.
- The case of periodic potentials is discussed in Sect. 6 (including stability eliminates).

In Sect. 2 we prove preliminary results about the mappings \mathcal{U}_{σ} , U_1 , U_o . In Sect. 3 the main results for Dirichlet and Neumann b.c. are proved. In Sect. 7 we determined some specific asymptotics for the fundamental solutions, and trace formulas.

2. Properties of $\mathcal{U}_{\sigma}, U_1, U_{\sigma}$

2.1. **Preliminaries.** We introduce mappings $q \to \mathfrak{t} = (\mathfrak{t}_n)_1^{\infty}$ and $q \to \mathfrak{r} = (\mathfrak{r}_n)_1^{\infty}$, where the components are norming constants $\mathfrak{t}_n, \mathfrak{r}_n$ (associated with the mixed eigenvalues τ_n, ϱ_n , respectively) and are given by

$$\mathfrak{t}_n = \ln|\varphi(1, \tau_n)\sqrt{\tau_n^o}|, \qquad \mathfrak{r}_n(q) = -\ln|\vartheta'(1, \varrho_n)/\sqrt{\tau_n^o}| = \ln|\varphi(1, \varrho_n)\sqrt{\varrho_n^o}|, \tag{2.1}$$

where $\vartheta'(1,\varrho_n)\varphi(1,\varrho_n)=-1$ has been used. Note that $\mathfrak{t}(q)\in\ell_1^2$, see [26]. In the case of mix boundary conditions y(0) = 0, y'(1) = 0 the spectral data $(\tau_n)_1^{\infty}$, $(\mathfrak{t}_n)_1^{\infty}$ are not independent since they satisfy the trace formula (4.4). It turns out that the first eigenvalue τ_1 can be uniquely reconstructed from the other spectral data $(\tau_n)_2^{\infty}$ and $(\mathfrak{t}_n)_1^{\infty}$. It is possible to "exclude" from the spectral data not the first eigenvalue τ_1 but an arbitrary norming constant \mathfrak{t}_m , see [26]. Thus we define the spectral data

$$\mathfrak{J}^{1}(2) = \left\{ t = (t_{n})_{2}^{\infty} : t_{2} < t_{3} < \dots, \quad t_{n} = \tau_{n}^{o} + \varepsilon_{n}, \qquad \varepsilon = (\varepsilon_{n-1})_{1}^{\infty} \in \ell^{2} \right\}.$$
 (2.2)

We recall known results about a RAB (i.e., a real-analytic bijection) for inverse problems on a unit interval. We formulate only results needed below.

Theorem 2.1. i) The mapping $q \mapsto (\tau \star \mu)(q)$ defined by (1.4), (1.5) is a RAB between \mathcal{L} and \mathfrak{J} .

- ii) The mapping $q \mapsto (\mu \times h_{(s)})(q)$ defined by (1.4), (1.6) is a RAB between \mathcal{L} and $\mathfrak{J}^o \times \ell_1^2$.
- iii) The mapping $(\tau_n)_2^{\infty} \times \mathfrak{t}$ from \mathcal{L} into $\mathfrak{J}^1(2) \times \ell_1^2$ is a RAB between \mathcal{L} and $\mathfrak{J}^1(2) \times \ell_1^2$.
- iv) The mappings $q \to (\nu \times \mathfrak{h}_{(s)})(q)$ from \mathcal{L} into $\mathfrak{J}^1 \times \ell_1^2$ is a RAB between \mathcal{L} and $\mathfrak{J}^1 \times \ell_1^2$. Moreover, all eigenvalues $(\nu_n)_0^{\infty}$ and norming constants $(\mathfrak{h}_{s,n})_0^{\infty}$ satisfy

$$\frac{e^{\pm \mathfrak{h}_{s,0}}}{|\dot{\vartheta}'(1,\nu_0)|} - 1 = \sum_{n=1}^{+\infty} \left(2 - \frac{e^{\pm \mathfrak{h}_{s,n}}}{|\dot{\vartheta}'(1,\nu_n)|}\right),\tag{2.3}$$

where the series converges absolutely and uniformly on bounded subsets of \mathcal{L} .

Remark. A bijection i) was proved in [32], see [34], [18] about a RAB. The results of ii) were proved in [34]. The results of iii) were proved in [26]. The results of iv) were proved in [26], the proof is based on [12] and the identities (2.3).

In order to study \mathcal{U}_{σ} , we describe basic properties of the 4-spectra mapping $\mathfrak{f}: \mathcal{L} \to \ell^2$.

Theorem 2.2. i) The 4-spectra mapping $\mathfrak{f}: \mathcal{L} \to \ell^2$ defined by

$$q \to \mathfrak{f}(q) = (\mathfrak{f}_n(q))_1^{\infty}, \quad \mathfrak{f}_{2n-1} = \varrho_n - \tau_n, \qquad \mathfrak{f}_{2n} = \nu_n - \mu_n, \qquad n \geqslant 1,$$
 (2.4)

is a RAB between \mathcal{L} and ℓ^2 . Furthermore, the following estimates hold true:

$$||q|| \le 2||\mathbf{f}||(1+2||\mathbf{f}||^{\frac{1}{3}}), \qquad ||\mathbf{f}|| \le 2||q||(1+2||q||^{\frac{1}{3}}),$$
 (2.5)

where $\|\mathfrak{f}\|^2 = \sum_{n\geqslant 1} (|\nu_n - \mu_n|^2 + |\tau_n - \varrho_n|^2)$. ii) Let $q, q' \in \mathcal{L}$. Then the following identities hold true:

$$\frac{q(0)-q(1)}{2} = \sum_{n\geq 1} \mathfrak{f}_{2n-1}, \qquad \frac{q(0)+q(1)}{2} = \nu_0 + \sum_{n\geq 1} \mathfrak{f}_{2n},
q(0) = \nu_0 + \sum_{n\geq 1} \mathfrak{f}_n, \qquad q(1) = \nu_0 + \sum_{n\geq 1} (-1)^n \mathfrak{f}_{2n}.$$
(2.6)

Proof. The results of i) were proved in [18]. Identities (2.6) follow from (7.51), (7.50) and (7.47).

We consider the operator Ty = -y'' + qy on $L^2(0,2)$ with 2-periodic conditions $y(x+2) = y(x), x \in \mathbb{R}$, where the potential q is 1-periodic and belongs to the real space \mathcal{H} defined by

$$\mathcal{H} = \Big\{ f \in L^2(\mathbb{T}, \mathbb{R}) : \int_0^1 f dx = 0 \Big\}, \qquad \mathbb{T} = \mathbb{R}/\mathbb{Z}.$$

The spectrum of T is eigenvalues $\lambda_0^+, \lambda_n^{\pm}, n \ge 1$ which satisfy

$$\lambda_0^+ < \lambda_1^- \leqslant \lambda_1^+ < \dots \leqslant \lambda_{n-1}^+ < \lambda_n^- \leqslant \lambda_n^+ < \dots,$$

$$\lambda_n^{\pm} = (\pi n)^2 + o(1) \quad \text{as} \quad n \to \infty.$$

These eigenvalues have the known relations (see Fig. 1)

$$\nu_0 \leqslant \lambda_0^+, \quad \tau_n, \varrho_n \in (\lambda_{n-1}^+, \lambda_n^-), \quad \text{and} \quad \mu_n, \nu_n \in [\lambda_n^-, \lambda_n^+], \quad \forall n \geqslant 1.$$
 (2.7)

Here the equality $\lambda_n^- = \lambda_n^+$ means that λ_n^- is a double eigenvalue. The lowest eigenvalue λ_0^+ is simple, and the corresponding eigenfunction has period 1. The eigenfunctions corresponding to λ_n^{\pm} have period 1 when n is even and they are antiperiodic, $y(x+1) = -y(x), \ x \in \mathbb{R}$, when n is odd.

FIGURE 1. Periodic λ_n^{\pm} , Dirichlet μ_n , Neumann ν_n and mixed τ_n , ϱ_n eigenvalues.

For each $q \in \mathcal{L}_0$ we consider the auxiliary Sturm-Liouville problems on the interval [0, 2] with the Dirichlet and Neumann boundary conditions with even potentials \tilde{q} on [0, 2]:

$$-f'' + \widetilde{q}f = \lambda f, \qquad \begin{array}{c} f(0) = f(2) = 0, \\ f'(0) = f'(2) = 0, \end{array} \qquad \widetilde{q}(x) = \begin{cases} q(x), & 0 < x < 1 \\ q(2 - x), & 1 < x < 2 \end{cases}$$
 (2.8)

Let $\widetilde{\mu}_n$ and $\widetilde{\nu}_0, \widetilde{\nu}_n, n \geqslant 1$ be eigenvalues of the Dirichlet and Neumann problems respectively. For a potential $\widetilde{q} \in L^2(0,2)$ given by (2.8) we denote by the same letter \widetilde{q} its 2-periodic extension to the real line. Introduce a operator $\widetilde{T}y = -y'' + \widetilde{q}y$ on $L^2(0,4)$ with 4-periodic conditions, that is $y(x+4) = y(x), x \in \mathbb{R}$. The spectrum of \widetilde{T} is a union of the eigenvalues $\widetilde{\lambda}_0^+, \widetilde{\lambda}_n^\pm, n \geqslant 1$ which satisfy

$$\begin{split} \widetilde{\lambda}_0^+ < \widetilde{\lambda}_1^+ \leqslant \widetilde{\lambda}_1^+ < \ldots \leqslant \widetilde{\lambda}_{n-1}^+ < \widetilde{\lambda}_n^- \leqslant \widetilde{\lambda}_n^+ < \ldots \\ \widetilde{\lambda}_n^\pm = (\frac{\pi n}{2})^2 + o(1) \quad \text{as} \quad n \to \infty. \end{split}$$

Here the equality $\widetilde{\lambda}_n^- = \widetilde{\lambda}_n^+$ means that $\widetilde{\lambda}_n^-$ is a double eigenvalue. The lowest eigenvalue $\widetilde{\lambda}_0^+$ is simple, and the corresponding eigenfunction has period 2. The eigenfunctions corresponding to $\widetilde{\lambda}_n^{\pm}$ have period 2 when n is even and they are antiperiodic, $y(x+2) = -y(x), \ x \in \mathbb{R}$, when n is odd. It is well known the Dirichlet and Neumann eigenvalues for even potentials \widetilde{q} satisfy

$$\tilde{\nu}_0 = \tilde{\lambda}_0^+, \qquad \{\tilde{\lambda}_n^-, \tilde{\lambda}_n^+\} = \{\tilde{\mu}_n, \tilde{\nu}_n\} \quad \text{for all } n \geqslant 1,$$
 (2.9)

see e.g., [8], [16]. Recall standards results, see e.g., [18].

Lemma 2.3. Let \widetilde{q} be given by (2.8) for some $q \in \mathcal{L}_0$. Then Dirichlet $\widetilde{\mu}_n$ and Neumann eigenvalues $\widetilde{\nu}_{n-1}$ and periodic eigenvalues $\lambda_n^{\pm}, n \geqslant 1$ satisfy

$$\widetilde{\mu}_{2n-1} = \tau_n, \qquad \widetilde{\mu}_{2n} = \mu_n, \tag{2.10}$$

$$\widetilde{\nu}_{2n-1} = \varrho_n, \qquad \widetilde{\nu}_{2n} = \nu_n, \tag{2.11}$$

$$\widetilde{\nu}_{2n-1} = \varrho_n, \qquad \widetilde{\nu}_{2n} = \nu_n,
\{\widetilde{\lambda}_{2n-1}^-, \widetilde{\lambda}_{2n-1}^+\} = \{\varrho_n, \tau_n\}, \qquad \{\widetilde{\lambda}_{2n}^-, \widetilde{\lambda}_{2n}^+\} = \{\mu_n, \nu_n\},$$
(2.11)

where $\{A, B\}$ is a set of two elements A, B

The functions $\varphi(1,\lambda), \varphi'(1,\lambda)$ and $\vartheta(1,\lambda)$ are entire and have the Hadamard factorizations

$$\varphi(1,\lambda) = \prod_{1}^{\infty} \frac{\mu_{n} - \lambda}{\mu_{n}^{o}}, \quad \varphi'(1,\lambda) = \prod_{1}^{\infty} \frac{\tau_{n} - \lambda}{\tau_{n}^{o}},
\vartheta(1,\lambda) = \prod_{1}^{\infty} \frac{\varrho_{n} - \lambda}{\varrho_{n}^{o}}, \quad \vartheta'(1,\lambda) = (\nu_{0} - \lambda)\vartheta_{*}(\lambda), \qquad \vartheta_{*}(\lambda) = \prod_{1}^{\infty} \frac{\nu_{n} - \lambda}{\nu_{n}^{o}},$$
(2.13)

uniformly in every bounded disk, see e.g., [34], [33].

2.2. Mappings \mathcal{U}_{σ} . Now we describe properties of the mapping \mathcal{U}_{σ} .

Proposition 2.4. Let $\mathcal{U}_{\sigma} = \mathfrak{f}^{-1}\sigma\mathfrak{f}: \mathcal{L} \to \mathcal{L}$ for some operator $\sigma = (\sigma_i)_1^{\infty} \in \mathfrak{S}$. Then

i) The 4-periodic eigenvalues $\{\widetilde{\lambda}_0^+, \widetilde{\lambda}_n^\pm, n \geqslant 1\}$ are invariant under \mathcal{U}_{σ} and

$$(\widetilde{\lambda}_0^+, (\widetilde{\lambda}_n^\pm)_1^\infty) = (\widetilde{\lambda}_0^+, (\widetilde{\lambda}_n^\pm)_1^\infty) \circ \mathcal{U}_\sigma. \tag{2.14}$$

ii) If $q^{\bullet} = \mathcal{U}_{\sigma}(q)$ for some $q \in \mathcal{L}$, then for each $j \in \mathbb{N}$ we have

if
$$n = 2j - 1 \Rightarrow \begin{cases} \tau_j(q) = \varrho_j(q^{\bullet}), & \varrho_j(q) = \tau_j(q^{\bullet}) & \text{if } \sigma_n = -1 \\ \tau_j(q) = \tau_j(q^{\bullet}), & \varrho_j(q) = \varrho_j(q^{\bullet}) & \text{if } \sigma_n = 1 \end{cases}$$
, (2.15)

if
$$n = 2j \Rightarrow \begin{cases} \mu_j(q) = \nu_j(q^{\bullet}), & \nu_j(q) = \mu_j(q^{\bullet}) & \text{if } \sigma_n = -1\\ \mu_j(q) = \mu_j(q^{\bullet}), & \nu_j(q) = \nu_j(q^{\bullet}) & \text{if } \sigma_n = 1 \end{cases}$$
 (2.16)

iii) Let $\sigma_n = -1$ (or $\sigma_n = 1$) for all odd $n \ge 1$. Let $\sigma_n \in \{\pm 1\}$ for all even $n \ge 1$. Then 2-periodic eigenvalues $\{\lambda_0^+, \lambda_n^{\pm}, n \geq 1\}$ are invariant under \mathcal{U}_{σ} and

$$(\lambda_0^+, (\lambda_n^\pm)_1^\infty) = (\lambda_0^+, (\lambda_n^\pm)_1^\infty) \circ \mathcal{U}_{\sigma}.$$
 (2.17)

iv) Neumann ν_0 is invariant under the mapping \mathcal{U}_{σ} , i.e., $\nu_0 = \nu_0 \circ \mathcal{U}_{\sigma}$ for all $\sigma \in \mathfrak{S}$.

Proof. The statements i-ii) follow from Lemma 2.3.

iii) Consider the case when $\sigma_n = -1$ for all odd $n \ge 1$, the proof for $\sigma_n = 1$ is similar. Let $\sigma_n \in \{\pm 1\}$ for all even $n \ge 1$. Then from (2.15) we have

if
$$n = 2j - 1 \Rightarrow \tau_j(q) = \varrho_j(q^{\bullet}), \quad \varrho_j(q) = \tau_j(q^{\bullet}), \quad q^{\bullet} = \mathcal{U}_{\sigma}.$$

These identities and (2.13) imply

$$2\Delta(\cdot,q) = \varphi'(1,\cdot,q) + \vartheta(1,\cdot,q) = \vartheta(1,\cdot,q^{\bullet}) + \varphi'(1,\cdot,q^{\bullet}) = 2\Delta(\cdot,q^{\bullet}),$$

which yields $\lambda_0^+(q) = \lambda_0^+(q^{\bullet}), \lambda_n^{\pm}(q) = \lambda_n^{\pm}(q^{\bullet})$ for all $n \ge 1$. iv) Let $q^{\bullet} = \mathcal{U}_{\sigma}(q)$ for some $q \in \mathcal{L}$. Rewrite a Wronskian in the form

$$f(\lambda,q)-g(\lambda,q)=1, \quad \text{where} \quad f(\lambda,q)=\vartheta(1,\lambda,q)\varphi'(1,\lambda,q), \quad g(\lambda,q)=\vartheta'(1,\lambda,q)\varphi(1,\lambda,q).$$

The function $f(\cdot,q)$ is entire and has zeros ϱ_n, τ_n and due to (2.15) we obtain $f(\cdot,q) = f(\cdot,q^{\bullet})$ and then $g(\cdot,q) = g(\cdot,q^{\bullet})$. The function $g(\cdot,q)$ is entire and has zeros $\nu_0(q)$ and a collection $A(q) = \{\nu_n(q), \lambda_n(q), n \ge 1\}$. Note that due to (2.16) we obtain $A(q) = A(q^{\bullet})$, which yields $\nu_0(q^{\bullet}) = \nu_0(q)$.

We discuss properties of a mapping $U_1 = \mathcal{U}_{\sigma} = \mathfrak{f}^{-1}\sigma\mathfrak{f}$, where $\sigma_n = -1$ for all $n \in \mathbb{N}$.

Lemma 2.5. Let $q^{(1)} = U_1(q)$ for some $q \in \mathcal{L}$ and \widetilde{q} is given by (2.8). Then the eigenvalues for $q, q^{(1)}, \widetilde{q}$ satisfy

$$\begin{cases}
\widetilde{\mu}_{2n} = \mu_n = \nu_n(q^{(1)}), \\
\widetilde{\mu}_{2n-1} = \tau_n = \varrho_n^{(1)}
\end{cases}, \qquad
\begin{cases}
\widetilde{\nu}_{2n} = \nu_n = \mu_n^{(1)} \\
\widetilde{\nu}_{2n-1} = \varrho_n = \tau_n^{(1)}
\end{cases}, \qquad (2.18)$$

$$\begin{cases}
\{\widetilde{\lambda}_{2n-1}^{-}, \widetilde{\lambda}_{2n-1}^{+}\} = \{\varrho_n, \tau_n\} = \{\varrho_n^{(1)}, \tau_n^{(1)}\}, \\
\{\widetilde{\lambda}_{2n}^{-}, \widetilde{\lambda}_{2n}^{+}\} = \{\mu_n, \nu_n\} = \{\mu_n^{(1)}, \nu_n^{(1)}\}
\end{cases},$$
(2.19)

$$(\mu, \nu, \tau, \varrho)(q) = (\nu, \mu, \varrho, \tau)(q^{(1)}),$$
 (2.20)

for all $n \ge 1$, where $\mu_n = \mu_n(q), \mu_n^{(1)} = \mu_n(q^{(1)}), \dots$ and $\widetilde{\mu}_n = \mu_n(\widetilde{q}), \dots$ for shortness, and

$$\varphi'(1,\cdot,q) = \vartheta(1,\cdot,q^{(1)}), \quad \vartheta(1,\cdot,q) = \varphi'(1,\cdot,q^{(1)}),$$
 (2.21)

$$\mathfrak{h}_{s,n}(q) = \ln|\vartheta(1,\nu_n,q)| = \ln|\varphi'(1,\mu_n^{(1)},q^{(1)})| = h_{s,n}(q^{(1)}). \tag{2.22}$$

Proof. Fix $q \in \mathcal{L}$ and the corresponding $\mathfrak{f}(q) = (\mathfrak{f}_n(q))_1^{\infty}$. Define a mapping $q \to \mathfrak{f}^{\bullet}(q) = -\mathfrak{f}(q) \in \ell^2$. For this case there exists $q^{(1)} \in \mathcal{L}$ such that $\mathfrak{f}(q^{(1)}) = \mathfrak{f}^{\bullet} = -\mathfrak{f}(q)$, since \mathfrak{f} is a homeomorphism. Then we obtain

$$\begin{cases}
-\mathfrak{f}_{2n-1}(q) = -\tau_n(q) + \varrho_n(q) = \mathfrak{f}_{2n-1}^{\bullet}(q^{(1)}) = \tau_n(q^{(1)}) - \varrho_n(q^{(1)}), \\
-\mathfrak{f}_{2n}(q) = -\mu_n(q) + \nu_n(q) = \mathfrak{f}_{2n}^{\bullet}(q^{(1)}) = \mu_n(q^{(1)}) - \nu_n(q^{(1)}),
\end{cases} \qquad n \geqslant 1. \tag{2.23}$$

Due to Lemma 2.3 we have

$$\begin{cases}
\{\widetilde{\lambda}_{2n-1}^{-}, \widetilde{\lambda}_{2n-1}^{+}\} = \{\varrho_n, \tau_n\} = \{\varrho_n^{(1)}, \tau_n^{(1)}\} \\
\{\widetilde{\lambda}_{2n}^{-}, \widetilde{\lambda}_{2n}^{+}\} = \{\mu_n, \nu_n\} = \{\mu_n^{(1)}, \nu_n^{(1)}\}
\end{cases}$$
(2.24)

and jointly with (2.10), (2.11), (2.23) we obtain

$$\begin{cases} \widetilde{\mu}_{2n} = \mu_n = \nu_n^{(1)}, \\ \widetilde{\mu}_{2n-1} = \tau_n = \varrho_n^{(1)} \end{cases}, \qquad \begin{cases} \widetilde{\nu}_{2n} = \nu_n = \mu_n^{(1)} \\ \widetilde{\nu}_{2n-1} = \varrho_n = \tau_n^{(1)} \end{cases}, \qquad \forall \ n \geqslant 1.$$
 (2.25)

These identities give (2.18)–(2.20). Due to (2.9) the eigenvalues $\widetilde{\nu}_n, \widetilde{\mu}_n$ belong to the ends of "gaps" $[\widetilde{\lambda}_n^-, \widetilde{\lambda}_n^+]$ for the even potential \widetilde{q} . From (2.13), (2.20) we have (2.21), which yields (2.22).

We discuss properties of a mapping $U_o = \mathcal{U}_\sigma$, where $\sigma_j = (-1)^j$ for all $j \in \mathbb{N}$.

Lemma 2.6. Let $q \in \mathcal{L}$ and \widetilde{q} be given by (2.8). Then eigenvalues for $q, q^{(o)} = U_o(q)$ and \widetilde{q} satisfy

$$\begin{cases}
\mu_n = \mu_n^{(o)} = \widetilde{\mu}_{2n}, \\
\tau_n = \varrho_n^{(o)} = \widetilde{\mu}_{2n-1}
\end{cases}, \qquad
\begin{cases}
\nu_n = \nu_n^{(o)} = \widetilde{\nu}_{2n} \\
\varrho_n = \tau_n^{(o)} = \widetilde{\nu}_{2n-1}
\end{cases}, \qquad (2.26)$$

$$\begin{cases}
\{\widetilde{\lambda}_{2n-1}^{-}, \widetilde{\lambda}_{2n-1}^{+}\} = \{\varrho_n, \tau_n\} = \{\varrho_n^{(o)}, \tau_n^{(o)}\}, \\
\{\widetilde{\lambda}_{2n}^{-}, \widetilde{\lambda}_{2n}^{+}\} = \{\mu_n, \nu_n\} = \{\mu_n^{(o)}, \nu_n^{(o)}\}
\end{cases},$$
(2.27)

$$(\mu, \nu, \tau, \varrho)(q) = (\mu, \nu, \varrho, \tau)(q^{(o)}), \tag{2.28}$$

for all $n \ge 1$, where $\mu_n = \mu_n(q), \mu_n^{(o)} = \mu_n(q^{(o)}), \dots$ and $\widetilde{\mu}_n = \mu_n(\widetilde{q}), \dots$ for shortness, and

$$\varphi'(1,\cdot,q) = \vartheta(1,\cdot,q^{(o)}), \quad \vartheta(1,\cdot,q) = \varphi'(1,\cdot,q^{(o)}), \tag{2.29}$$

$$\mathfrak{h}_{s,n}(q) = \ln|\vartheta(1,\nu_n,q)| = \ln|\varphi'(1,\mu_n^{(o)},q^{(o)})| = h_{s,n}(q^{(o)}). \tag{2.30}$$

Proof. Fix $q \in \mathcal{L}$ and the corresponding $\mathfrak{f}(q) = (\mathfrak{f}_n(q))_1^{\infty}$. Define a mapping

$$q \to \mathfrak{f}^{(o)}(q) = ((-1)^n \mathfrak{f}_n(q))_1^{\infty} \in \ell^2.$$

For this case there exists $q^{(o)} \in \mathcal{L}$ such that $\mathfrak{f}(q^{(o)}) = \mathfrak{f}^{\bullet}(q^{(o)})$, since \mathfrak{f} is a homeomorphism. Then we obtain

$$\begin{cases} -\mathfrak{f}_{2n-1}(q) = -\tau_n(q) + \varrho_n(q) = \mathfrak{f}_{2n-1}^{\bullet}(q^{(o)}) = \tau_n(q^{(o)}) - \varrho_n(q^{(o)}), \\ \mathfrak{f}_{2n}(q) = \mu_n(q) - \nu_n(q) = \mathfrak{f}_{2n}^{\bullet}(q^{(o)}) = \mu_n(q^{(o)}) - \nu_n(q^{(o)}) \end{cases} \qquad n \geqslant 1.$$
 (2.31)

Due to Lemma 2.3 we have (2.27)

$$\begin{cases}
\{\widetilde{\lambda}_{2n-1}^{-}, \widetilde{\lambda}_{2n-1}^{+}\} = \{\varrho_n, \tau_n\} = \{\varrho_n^{(o)}, \tau_n^{(o)}\}, \\
\{\widetilde{\lambda}_{2n}^{-}, \widetilde{\lambda}_{2n}^{+}\} = \{\mu_n, \nu_n\} = \{\mu_n^{(o)}, \nu_n^{(o)}\}
\end{cases},$$
(2.32)

and jointly with (2.10), (2.11), (2.23) we obtain (2.26). These identities give (2.27)–(2.28). Due to (2.9) the eigenvalues $\widetilde{\nu}_n$, $\widetilde{\mu}_n$ belong to the ends of gaps $(\widetilde{\lambda}_n^-, \widetilde{\lambda}_n^+)$ for the even potential \widetilde{q} . From (2.13), (2.20) we have (2.29) and then (2.30).

Proof of Proposition 1.1 Due to the estimate (2.5) the mapping $\mathcal{U}_{\sigma} = \mathfrak{f}^{-1}\sigma\mathfrak{f}: \mathcal{L} \to \mathcal{L}$ is bounded in any ball $\{\|q\| \leqslant R\}$. The definition $\mathcal{U}_{\sigma} = \mathfrak{f}^{-1}\sigma\mathfrak{f}: \mathcal{L} \to \mathcal{L}$ implies that $\mathcal{U}_{\sigma} = \mathcal{U}_{\sigma}^{-1}$. The definition $\mathcal{U}_{\sigma} = \mathfrak{f}^{-1}\sigma\mathfrak{f}$ implies $\mathcal{U}_{\sigma}\mathcal{U}_{\sigma'} = \mathcal{U}_{\sigma\sigma'} = \mathcal{U}_{\sigma'}\mathcal{U}_{\sigma}$ for all $\sigma, \sigma' \in \mathfrak{S}$.

The definition $\mathcal{U}_{\sigma} = \mathfrak{f}^{-1}\sigma\mathfrak{f}$ implies $\mathcal{U}_{\sigma} \mathcal{U}_{\sigma'} = \mathcal{U}_{\sigma\sigma'} = \mathcal{U}_{\sigma'} \mathcal{U}_{\sigma}$ for all $\sigma, \sigma' \in \mathfrak{S}$. Due to (2.14) the 4-periodic eigenvalues $\{\widetilde{\lambda}_0^+, \widetilde{\lambda}_n^\pm, n \geq 1\}$ are invariant under \mathcal{U}_{σ} and then the Lyapunov function for the potential $\widetilde{q}(x)$ given by (2.8) is also invariant under \mathcal{U}_{σ} . Thus a functional $\int_0^2 |\widetilde{q}(x)|^2 dx$ is invariant under \mathcal{U}_{σ} (see e.g., [19]) and we obtain for $r = \mathcal{U}_{\sigma}(q)$:

$$2\int_0^1 |r(x)|^2 dx = \int_0^2 |(\widetilde{r})(x)|^2 dx = \int_0^2 |\widetilde{q}(x)|^2 dx = 2\int_0^1 |q(x)|^2 dx,$$

which yields $\|\mathcal{U}_{\sigma}(q)\| = \|q\|$.

3. Proof of main results for the unit interval

We discuss main results about 2-spectra mappings.

Proof of Theorem 1.2. i) From (2.20) we obtain $(\tau \star \mu)(q) = (\varrho \star \nu)(q^{(1)})$, where $q^{(1)} = U_1 q$, which yields $\tau \star \mu = (\varrho \star \nu) \circ U_1$. By Theorem 2.1, the mapping $q \mapsto (\tau \star \mu)(q)$ from \mathcal{L} to \mathfrak{J} is a RAB between \mathcal{L} and \mathfrak{J} . Then the mapping $q \mapsto (\varrho \star \nu)(q)$ from \mathcal{L} to \mathfrak{J} is a RAB between \mathcal{L} and \mathfrak{J} , since due to Proposition 1.1 U_1 is a RAB of \mathcal{L} onto itself.

From (2.28) we obtain $(\tau \star \mu)(q) = (\varrho \star \mu)(q^{(o)})$, where $q^{(o)} = U_o q$. Then due to Theorem 2.1 the mapping $q \mapsto (\varrho \star \mu)(q)$ from \mathcal{L} to \mathfrak{J} is a RAB between \mathcal{L} and \mathfrak{J} , since due to Proposition 1.1 U_o is an unitary operator on \mathcal{L} .

Above we have obtained $\tau \star \mu = (\varrho \star \nu) \circ U_1$. From (2.28) we have $\varrho \star \nu = (\tau \star \nu) \circ U_o$. A combination of these identities gives $\tau \star \mu = (\tau \star \nu) \circ U_o \circ U_1$. Then due to Theorem 2.1 the mapping $q \mapsto (\varrho \star \nu)(q)$ from \mathcal{L} to \mathfrak{J} is a RAB between \mathcal{L} and \mathfrak{J} , since due to Proposition 1.1 U_o, U_1 are RABs of \mathcal{L} onto itself. Collecting all identities we obtain (1.19).

ii) Let $q \in \mathcal{L}$ and let $\mathfrak{n} = \varrho \star \nu$, $\mathfrak{n}_{2j-1} = \varrho_j$, $\mathfrak{n}_{2j} = \nu_j$ be given. Above we have obtained $(\tau \star \mu)(q) = (\varrho \star \nu)(q^{(1)})$, where $q^{(1)} = U_1q$. This yields $\tau(q) = \varrho(q^{(1)})$ and $\mu(q) = \nu(q^{(1)})$, which implies $\tau(q^{(1)}) = \varrho(q)$, $\mu(q^{(1)}) = \nu(q)$, since $U_1^2 = I$. Recall that the recovering of a potential q by (1.16) via the 2-spectra mapping $\mathfrak{n} = \tau \star \mu$, $\mathfrak{n}_{2j-1} = \tau_j$, $\mathfrak{n}_{2j} = \mu_j$, $j \in \mathbb{N}$, was obtained in [18]. Then we obtain the recovering a potential $q^{(1)}$ by (1.16) via the 2-spectra mapping $\mathfrak{n} = \varrho \star \nu$, $\mathfrak{n}_{2j-1} = \varrho_j$, $\mathfrak{n}_{2j} = \nu_j$. The proof of iii) is similar.

Below we need identities, see e.g. [34]: let y be a solution to the equation $-y'' + qy = \lambda y$, then

$$\int_{0}^{1} y^{2}(x,\lambda)dx = \{\dot{y}, y\}_{w}\Big|_{0}^{1}, \quad \text{where} \quad \{f, y\}_{w} = fy' - f'y, \quad \dot{y} = \frac{\partial y}{\partial \lambda}.$$
 (3.1)

We discuss results about mappings with eigenvalues plus norming (or normolizing) constants. **Proof of Theorem 1.3.** i) From (2.20), (2.22) we obtain $\mu \times h_{(s)} = (\nu \times \mathfrak{h}_{(s)}) \circ U_1$. Then due to Theorem 2.1 the mapping $q \mapsto \nu \times \mathfrak{h}_{(s)}$ from \mathcal{L} to $\mathfrak{J}^o \times \ell_1^2$ is a RAB between \mathcal{L} and $\mathfrak{J}^o \times \ell_1^2$, since by Proposition 1.1, U_1 is a RAB of \mathcal{L} onto itself and $\mu \times h_{(s)} : \mathcal{L} \to \mathfrak{J}^o \times \ell_1^2$ is a RAB.

From (2.28), (2.30) we obtain $\mu \times h_{(s)} = (\mu \times \mathfrak{h}_{(s)})U_o$. Then due to Theorem 2.1 the mapping $q \mapsto (\mu \times \mathfrak{h}_{(s)})(q)$ from \mathcal{L} to $\mathfrak{J}^o \times \ell_1^2$ is a RAB between \mathcal{L} and $\mathfrak{J}^o \times \ell_1^2$, since due to Proposition 1.1 U_o is an unitary operator on \mathcal{L} .

Above we have obtained $\mu \times h_{(s)} = (\nu \times \mathfrak{h}_{(s)}) \circ U_1$. From (2.28) we have $\nu \times \mathfrak{h}_{(s)} = (\nu \times h_{(s)}) \circ U_o$. A combination of these identities gives

$$\mu \times h_{(s)} = (\nu \times \mathfrak{h}_{(s)}) \circ U_1 = (\nu \times h_{(s)}) \circ U_o \circ U_1.$$

Then due to Theorem 2.1 the mapping $q \mapsto (\nu \times \mathfrak{h}_{(s)})(q)$ from \mathcal{L} to $\mathfrak{J}^o \times \ell_1^2$ is a RAB between \mathcal{L} and $\mathfrak{J}^o \times \ell_1^2$, since due to Proposition 1.1 U_o, U_1 are RABs of \mathcal{L} onto itself. Collecting all identities we obtain (1.20).

ii) Consider the mapping $\mu \times \alpha$. The identity (3.1) implies

$$D_n = \dot{\varphi}(1, \mu_n) \varphi'(1, \mu_n), \qquad \forall \ n \in \mathbb{N}. \tag{3.2}$$

Thus using (3.2), $D_n(0) = \frac{1}{2\mu_n^o}$ and $(-1)^n \varphi'(1,\mu_n) = e^{h_{s,n}}$ we rewrite α_n in the form

$$e^{\alpha_n} = \frac{D_n(q)}{D_n(0)} = [(-1)^n 2\mu_n^o \dot{\varphi}(1, \mu_n)] [(-1)^n \varphi'(1, \mu_n)] = e^{M_n + h_{s,n}}, \tag{3.3}$$

where $M_n \in \mathbb{R}$ is defined by $0 < (-1)^n 2\mu_n^o \dot{\varphi}(1,\mu_n) = e^{M_n}$ and due to (7.1) the sequence $(M_n)_1^\infty \in \ell_1^2$. Thus we obtain

$$\alpha_n = h_{s,n} + M_n, \quad \forall \quad n \in \mathbb{N}.$$
 (3.4)

Due to Theorem 2.1 the mapping $q \mapsto \mu \times h_{(s)}$ is a bijection between \mathcal{L} and $\mathfrak{J}^o \times \ell_1^2$ and the mapping $\mu \times h_{(s)} \to \mu \times \alpha$ is a bijection from $\mathfrak{J}^o \times \ell_1^2$ onto itself, since $\alpha_n, h_{s,n}$ satisfy (3.4). This gives that the mapping $q \mapsto (\mu \times \alpha)(q)$ is a bijection between \mathcal{L} and $\mathfrak{J}^o \times \ell_1^2$.

Consider the mapping $\nu \times \beta$. The identity (3.1) implies

$$N_n(q) = -\dot{\vartheta}'(1, \nu_n, q)\vartheta(1, \nu_n, q). \tag{3.5}$$

At q = 0 we have

$$N_n(0) = \frac{1}{2}, \qquad \dot{\vartheta}'(1, \nu_n^o, 0) = -\frac{(-1)^n}{2}, \qquad \vartheta(1, \nu_n^o, 0) = (-1)^n.$$
 (3.6)

Then due to the definition (1.10) of β_n , and (3.5),(3.6) and $(-1)^n \vartheta(1,\nu_n,q) = e^{\mathfrak{h}_{s,n}}$ we get

$$e^{\beta_n} = 2N_n = -2\dot{\vartheta}'(1,\nu_n)(-1)^n e^{\mathfrak{h}_{s,n}} = b_n[2(-1)^n \nu_n^o \vartheta_*(\nu_n)] e^{\mathfrak{h}_{s,n}} = e^{\mathfrak{h}_{s,n} + K_n + K_n^o}, \tag{3.7}$$

where $K_n^o, K_n \in \mathbb{R}$ are defined by $b_n = \frac{\nu_n - \nu_0}{\nu_n^o} = e^{K_n^o}$ and $0 < (-1)^{n+1} 2\nu_n^o \dot{\vartheta}_*(\nu_n) = e^{K_n}$ and we have used the Hadamard factorization of $\vartheta'(1,\lambda) = -(\lambda - \nu_0)\vartheta_*(\lambda)$ from (2.13). Thus we obtain

$$\beta_n = \mathfrak{h}_{s,n} + K_n + K_n^o, \quad K_n^o = \ln \frac{\nu_n - \nu_0}{\nu_n^o} = \frac{O(1)}{n^2},$$
 (3.8)

and (7.3) gives $(K_n)_1^{\infty} \in \ell_1^2$. Repeating arguments from the proof for α we obtain that the mapping $q \mapsto (\mu \times \beta)(q)$ is a bijection between \mathcal{L} and $\mathfrak{J}^o \times \ell_1^2$.

We show the trace formulas (1.21). The identity (3.7) yields $e^{-\mathfrak{h}_{s,n}} = |\dot{\vartheta}'(1,\nu_n)|/N_n$ and substituting one into (2.3) we obtain $\frac{1}{N_0} - 1 = \sum_{1}^{\infty} \left(2 - \frac{1}{N_n}\right)$, i.e., we have (1.21), where the series converges absolutely and uniformly on bounded subsets of \mathcal{L} .

iii) Let $q^{\bullet} = U_1(q)$ and let $\mu_n = \mu_n(q), \alpha_n = \alpha_n(q), \dots$ and $\mu_n^{\bullet} = \mu_n(q^{\bullet}), \alpha_n^{\bullet} = \alpha_n(q^{\bullet}), \dots$ From (2.18), (3.3), (3.7) we obtain for all $n \in \mathbb{N}$:

$$\mu_n^{\bullet} = \nu_n(q), \qquad e^{M_n^{\bullet}} = (-1)^n 2 \mu_n^o \dot{\varphi}(1, \mu_n^{\bullet}, q^{\bullet}) = (-1)^n 2 \nu_n^o \dot{\vartheta}_*(\nu_n, q) = e^{K_n}.$$

From (2.22) we have $h_{s,n}^{\bullet} = \mathfrak{h}_{s,n}$. Then these identities and (3.8) imply

$$\alpha_n^{\bullet} = h_{s,n}^{\bullet} + M_n^{\bullet} = \mathfrak{h}_{s,n} + K_n = \beta_n - K_n^o = \hat{\beta}_n,$$

which yields $(\mu \times \alpha) \circ U_1 = (\nu \times \hat{\beta})$. This gives that the mapping $q \mapsto (\nu \times \hat{\beta})(q)$ is a bijection between \mathcal{L} and $\mathfrak{J}^o \times \ell_1^2$, since all other mappings $\mu \times \alpha$ and U_1 are bijections.

Remark. In [36] the inverse problem for the mapping $q \to \mu \times (\widetilde{D}_n)_1^{\infty}$ is discussed, where \widetilde{D}_n is the normalizing constant given by $\widetilde{D}_n = \mu_n D_n(q)$. But the presentation and the proof is not entirely clear.

We discuss a new type of inverse problems. Let the Dirichlet mapping $q \to \mu = (\mu_n)_1^{\infty}$ be given and replace some μ_n by the Neumann eigenvalues ν_n . Then we obtain a replacing mapping \mathfrak{c} . For example, we have $\mathfrak{c} = (\mu_1, \nu_2, \nu_3, \mu_4, \mu_5, ...)$. There is a question: it is a good 1-spectra mapping? We discuss *replacing* mappings.

Corollary 3.1. (Replacing mappings.) Let $\sigma = (\sigma_n)_1^{\infty}$, where $\sigma_n \in \{\pm 1\}$. Define replacing mappings $q \to \mathfrak{a} = (\mathfrak{a}_n)_1^{\infty}$, $q \to \mathfrak{b} = (\mathfrak{b}_n)_1^{\infty}$ and $q \to \mathfrak{c} = (\mathfrak{c}_n)_1^{\infty}$ and their components by

$$\mathfrak{a}_{n} = \begin{cases} \tau_{n}, & \text{if } \sigma_{2n-1} = 1\\ \varrho_{n}, & \text{if } \sigma_{2n-1} = -1 \end{cases}, \quad \mathfrak{b}_{n} = \begin{cases} \mu_{n}, & \text{if } \sigma_{2n} = 1\\ \nu_{n}, & \text{if } \sigma_{2n} = -1 \end{cases}, \quad \mathfrak{c}_{n} = \begin{cases} h_{s,n}, & \text{if } \sigma_{2n-1} = 1\\ \mathfrak{h}_{s,n}, & \text{if } \sigma_{2n-1} = -1 \end{cases}.$$
(3.9)

i) Then the mapping $\mathfrak{a} \star \mathfrak{b} : \mathcal{L} \to \mathfrak{J}$ is a RAB between \mathcal{L} and \mathfrak{J} and satisfies

$$(\mathfrak{a} \star \mathfrak{b}) \circ \mathcal{U}_{\sigma} = \tau \star \mu = (\varrho \star \nu) \circ U_{1} = (\varrho \star \mu) \circ U_{o} = (\tau \star \nu) \circ U_{o} \circ U_{1}, \tag{3.10}$$

and all 2-spectra mappings $\mathfrak{a}\star\mathfrak{b}$, $\varrho\star\nu$, $\tau\star\nu$, $\varrho\star\mu$ and $\tau\star\mu$ acting from $\mathcal L$ into $\mathfrak J$ are isomorphic. ii) The mapping $\mathfrak{b}\times\mathfrak{c}:\mathcal L\mapsto\mathfrak J^o\times\ell_1^2$ is a RAB between $\mathcal L$ and $\mathfrak J^o\times\ell_1^2$ and satisfies

$$(\mathfrak{b} \times \mathfrak{c}) \circ \mathcal{U}_{\sigma} = \mu \times h_{(s)} = (\nu \times \mathfrak{h}_{(s)}) \circ U_1 = (\mu \times \mathfrak{h}_{(s)}) \circ U_o = (\nu \times h_{(s)}) \circ U_o \circ U_1,$$
 (3.11) and all these mappings are isomorphic.

Proof. i) Let $q^{\sigma} = \mathcal{U}_{\sigma}(q)$, where $q \in \mathcal{L}$. From (2.15) we deduce that

if $\sigma_{2n-1} = 1$ then the eigenvalues $\mathfrak{a}_n(q^{\sigma}) = \tau_n(q^{\sigma}) = \tau_n(q)$,

if $\sigma_{2n-1} = -1$ then the eigenvalues $\mathfrak{a}_n(q^{\sigma}) = \varrho_n(q^{\sigma}) = \tau_n(q)$.

From (2.16) we deduce that

if $\sigma_{2n} = 1$ then the eigenvalues $\mathfrak{b}_n(q^{\sigma}) = \mu_n(q^{\sigma}) = \mu_n(q)$,

if $\sigma_{2n} = -1$ then the eigenvalues $\mathfrak{b}_n(q^{\sigma}) = \nu_n(q^{\sigma}) = \mu_n(q)$.

Thus all these identities give $\mathfrak{a} \star \mathfrak{b} = (\tau \star \mu) \circ \mathcal{U}_{\sigma}$. Then due to Theorem 2.1 the mapping $q \mapsto (\mathfrak{a} \star \mathfrak{b})(q)$ from \mathcal{L} to \mathfrak{J} is a RAB between \mathcal{L} and \mathfrak{J} , since the mapping $U_{\sigma} : \mathcal{L} \to \mathcal{L}$ is a RAB from \mathcal{L} onto itself. Moreover, we have (3.10). The proof of ii) is similar.

4. MIXED BOUNDARY CONDITIONS

We discuss isomorphic inverse problems for mixed eigenvalues. Recall that we have introduced norming constants \mathfrak{t}_n , \mathfrak{r}_n (associated with mixed eigenvalues τ_n , ϱ_n respectively) and the corresponding mappings by

$$\mathfrak{t}_n = \ln |\varphi(1, \tau_n) \sqrt{\tau_n^o}|, \qquad \mathfrak{r}_n(q) = -\ln |\vartheta'(1, \varrho_n) / \sqrt{\tau_n^o}| = \ln |\varphi(1, \varrho_n) \sqrt{\varrho_n^o}|,
q \to \mathfrak{t} = (\mathfrak{t}_n)_1^{\infty}, \qquad q \to \mathfrak{r} = (\mathfrak{r}_n)_1^{\infty},$$
(4.1)

where $\mathfrak{t}(q) \in \ell_1^2$ if $q \in \mathcal{L}$, see [26]. We consider the known facts about properties of Sturm-Liouville problems under the reflection (unitary) operator $\mathcal{R}: \mathcal{L} \to \mathcal{L}$ defined by $(\mathcal{R}y)(x) = y(1-x), x \in (0,1)$. Let $y_n(x,q), n \geqslant 1$ be the eigenfunction corresponding be the Dirichlet eigenvalue $\mu_n(q)$:

$$-y_n'' + qy_n = \mu_n y_n, \quad q \in \mathcal{L}.$$

Then the function $u_n = \mathcal{R}y_n$ satisfies $-u_n'' + q^{\bullet}u_n = \mu_n u_n$, where $q^{\bullet} = \mathcal{R}q$ and then $\mu_n(q) = \mu_n(q^{\bullet})$. We apply similar arguments for the Neumann and mixed eigenvalues and we obtain

$$(\mu_n, \nu_n, \tau_n, \rho_n) = (\mu_n, \nu_n, \rho_n, \tau_n) \circ \mathcal{R} \quad \forall \ n \geqslant 1.$$
 (4.2)

In particular, it gives that the two mappings $\tau \times \mathfrak{t}$ and $\tau \times \mathfrak{r}$ acting from \mathcal{L} into $\mathfrak{J}^1(2) \times \ell_1^2$ are unitarily equivalent. We discuss the mappings $q \to \tau^{(2)} = (\tau_n)_2^{\infty}$ and $q \to \varrho^{(2)} = (\varrho_n)_2^{\infty}$ and formulate results based on [26] about inverse problems for mixed eigenvalues. In this case we obtain unitarily equivalent mappings.

Proposition 4.1. i) The operator $U_o = \mathcal{R}$ and satisfies

$$\varrho \times \mathfrak{r} = (\tau \times \mathfrak{t}) \circ U_o. \tag{4.3}$$

The two mappings $\tau \times \mathfrak{t}$ and $\tau \times \mathfrak{r}$ acting from \mathcal{L} into $\mathfrak{J}^1(2) \times \ell_1^2$ are unitarily equivalent. ii) Let $\mathfrak{J}^1(2)$ be defined by (2.2). Define a mapping $q \to \varrho^{(2)} = (\varrho_n)_2^{\infty}$. Then a mapping $\varrho^{(2)} \times \mathfrak{r}$ is a RAB between \mathcal{L} and $\mathfrak{J}^1(2) \times \ell_1^2$. Moreover, the following trace formulas hold true

$$\sum_{n=1}^{\infty} \left(2 - \frac{e^{\mathfrak{t}_n}}{\sqrt{\tau_n^o} \left| \frac{\varphi'}{\partial \lambda}(1, \tau_n) \right|} \right) = 0, \qquad \sum_{n=1}^{+\infty} \left(2 - \frac{e^{\mathfrak{r}_n(q)}}{\sqrt{\tau_n^o} \left| \frac{\partial \vartheta}{\partial \lambda}(1, \rho_n, q) \right|} \right) = 0. \tag{4.4}$$

Proof. i) Due to (4.2) the mappings ϱ and τ from \mathcal{L} into \mathfrak{J}^1 are unitarily equivalent and satisfy $\varrho = \tau \circ \mathcal{R}$. From (4.2), (1.19) we obtain $\tau \star \mu = (\varrho \star \mu) \circ U_o = (\varrho \star \mu) \circ \mathcal{R}$, which yields that $\mathcal{R} = U_0$, since due to Theorem 1.2 the mappings $\tau \star \mu$ and $\varrho \star \mu$ are bijections.

From $\mathcal{R} = U_o$ and (4.2) we have $\mu_n(q) = \mu_n(q^{\bullet}), q^{\bullet} = U_o q$, which jointly with (2.13) yield $\varphi(1, \lambda, q) = \varphi(1, \lambda, q^{\bullet})$. Then the definitions (4.1) and (4.2) imply

$$\mathfrak{t}_n(q) = \ln|\varphi(1, \tau_n(q), q)\sqrt{\tau_n^o}| = \ln|\varphi(1, \varrho_n(q^{\bullet}), q^{\bullet})\sqrt{\tau_n^o}| = \mathfrak{r}_n(q^{\bullet}), \quad \forall \ n \geqslant 1,$$

$$(4.5)$$

which jointly with $\mathcal{R} = U_0$, and (4.2) yield (4.3).

ii) Due to Theorem 2.1 and (4.3) the mapping $\varrho^{(2)} \times \mathfrak{r}$ is a RAB between \mathcal{L} and $\mathfrak{J}^1(2) \times \ell_1^2$, since $U_o = \mathcal{R}$ is an unitary operator on \mathcal{L} .

The first trace formula in (4.4) was proved in [26]. We show the second one. Identities (2.29) give $\varphi'(1,\lambda,q) = \vartheta(1,\lambda,q^{\bullet})$. Then substituting this identity, $\tau_n(q) = \varrho_n(q^{\bullet})$ from (4.2) and (4.5) into the first trace formula in (4.4) we obtain the second one in (4.4).

Consider inverse problems for a mapping $q \to \tau \times (\mathcal{D}_n)_1^{\infty}$, where \mathcal{D}_n is a normalizing constant defined by

$$\mathcal{D}_n(q) = \int_0^1 \varphi^2(x, \tau_n, q) dx, \quad n \in \mathbb{N}, \quad \text{where} \quad \mathcal{D}_n(0) = \frac{1}{2\tau^2}. \tag{4.6}$$

It is more convenient to modify constants \mathcal{D}_n and define another mapping $q \to \eta = (\eta_n)_1^{\infty}$, where the components η_n are given by

$$\eta_n = \log \frac{\mathcal{D}_n(q)}{\mathcal{D}_n(0)} = \log \left[2\tau_n^o \mathcal{D}_n(q) \right], \qquad n \in \mathbb{N}.$$
(4.7)

Consider inverse problems for a mapping $q \to \varrho \times (\mathcal{N}_n)_1^{\infty}$, where \mathcal{N}_n is a normalizing constant defined by

$$\mathcal{N}_n(q) = \int_0^1 \vartheta^2(x, \varrho_n, q) dx, \quad n \in \mathbb{N}, \quad \text{where} \quad \mathcal{N}_n(0) = \frac{1}{2}.$$
 (4.8)

It is more convenient to define modified normalizing constant χ_n given

$$\chi_n = -\log \frac{\mathcal{N}_n(q)}{\mathcal{N}_n(0)} = -\log \left[2\mathcal{N}_n(q)\right], \qquad n \in \mathbb{N}.$$
(4.9)

and introduce a mapping $q \to \chi = (\chi_n)_1^{\infty}$. We discuss results about inverse problems for mixed boundary conditions.

Theorem 4.2. i) Let $\sigma_{2n} = 1$ and $\sigma_{2n-1} \in \{\pm 1\}$ for all $n \in \mathbb{N}$. Define replacing mappings $q \to \mathfrak{a} = (\mathfrak{a}_n)_2^{\infty}$ and $q \to \mathfrak{c} = (\mathfrak{c}_n)_1^{\infty}$, where the components are given by

$$\mathfrak{a}_{n} = \begin{cases} \tau_{n}, & \text{if } \sigma_{2n-1} = 1\\ \varrho_{n}, & \text{if } \sigma_{2n-1} = -1 \end{cases}, \quad \mathfrak{c}_{n} = \begin{cases} \mathfrak{t}_{n}, & \text{if } \sigma_{2n-1} = 1\\ \mathfrak{r}_{n}, & \text{if } \sigma_{2n-1} = -1 \end{cases}. \tag{4.10}$$

Then the two mappings $\mathfrak{a} \times \mathfrak{c}$ and $\tau^{(2)} \times \mathfrak{t}$ from $\mathcal{L} \to \mathfrak{J}^1(2) \times \ell_1^2$ are isomorphic. Moreover, each of them is a RAB between \mathcal{L} and $\mathfrak{J}^1(2) \times \ell_1^2$ and satisfies

$$\mathfrak{a} \times \mathfrak{c} = (\tau^{(2)} \times \mathfrak{t}) \circ \mathcal{U}_{\sigma}. \tag{4.11}$$

ii) Each of two mappings $\tau^{(2)} \times \eta$ and $\varrho^{(2)} \times \chi$ acting from \mathcal{L} into $\mathfrak{J}^1(2) \times \ell_1^2$ is a bijection between \mathcal{L} and $\mathfrak{J}^1(2) \times \ell_1^2$.

Remark. 1) The mapping $q \to \tau \times (\mathcal{D}_n)_1^{\infty}$ is considered in [35], but the authors do not see that spectral data $\tau, (\mathcal{D}_n)_1^{\infty}$ are dependent due to (4.4).

Proof. i) Let $q^{\sigma} = \mathcal{U}_{\sigma}(q)$. Consider \mathfrak{a} . From (2.15) we deduce that the eigenvalues

$$\mathfrak{a}_n(q^{\sigma}) = \begin{cases} \tau_n(q^{\sigma}) = \tau_n(q) & \text{if } \sigma_{2n-1} = 1\\ \varrho_n(q^{\sigma}) = \tau_n(q) & \text{if } \sigma_{2n-1} = -1 \end{cases}$$
(4.12)

This yields $\mathfrak{a}_n(q^{\sigma}) = \tau_n(q)$ for all $n \ge 2$.

Consider the mapping \mathfrak{c} : if $\sigma_{2n-1}=1$, then due to (2.13), (4.12) the norming constant

$$\mathfrak{c}_n(q^{\sigma}) = \mathfrak{t}_n(q^{\sigma}) = \ln|\varphi(1, \tau_n(q^{\sigma}), q^{\sigma})\sqrt{\tau_n^{\sigma}}| = \ln|\varphi(1, \tau_n(q^{\sigma}), q)\sqrt{\tau_n^{\sigma}}|$$
$$= \ln|\varphi(1, \tau_n(q), q)\sqrt{\tau_n^{\sigma}}| = \mathfrak{t}_n(q),$$

if $\sigma_{2n-1} = -1$ then due to (2.13), (4.12) the norming constant

$$\mathfrak{c}_n(q^{\sigma}) = \mathfrak{r}_n(q^{\sigma}) = \ln |\varphi(1, \tau_n(q^{\sigma}), q^{\sigma}) \sqrt{\tau_n^{o}}| = \ln |\varphi(1, \varrho_n(q), q) \sqrt{\tau_n^{o}}| = \mathfrak{t}_n(q).$$

This yields $\mathfrak{c}_n(q^{\sigma}) = \mathfrak{t}_n(q)$ for all $n \in \mathbb{N}$. Collecting the identities we obtain (4.11). Then due to Theorem 2.1 the mapping $q \mapsto (\mathfrak{a} \times \mathfrak{c})(q)$ from \mathcal{L} to $\mathfrak{J}^1(2) \times \ell_1^2$ is a RAB between \mathcal{L} and $\mathfrak{J}^1(2) \times \ell_1^2$, since due to Proposition 1.1 U_{σ} is a RAB of \mathcal{L} onto itself.

ii) Consider the mapping $\tau^{(2)} \times \eta$. The identity (3.1) implies

$$\mathcal{D}_n = -\dot{\varphi}'(1, \tau_n)\varphi(1, \tau_n), \qquad \forall \ n \in \mathbb{N}. \tag{4.13}$$

Then using $(-1)^{n+1}\varphi(1,\tau_n)\sqrt{\tau_n^o}=e^{t_n}$ we rewrite η_n in the form

$$e^{\eta_n} = \frac{\mathcal{D}_n(q)}{\mathcal{D}_n(0)} = [(-1)^n 2\sqrt{\tau_n^{\bar{o}}} \ \dot{\varphi}'(1,\tau_n)] \ [(-1)^{n+1}\sqrt{\tau_n^{\bar{o}}}\varphi(1,\tau_n)] = e^{\mathcal{M}_n + \mathfrak{t}_n}, \tag{4.14}$$

where $\mathcal{M}_n \in \mathbb{R}$ is defined by $0 < (-1)^n 2\sqrt{\tau_n^o} \ \dot{\varphi}'(1,\tau_n) = e^{\mathcal{M}_n}$ and due to (7.2) the sequence $(\mathcal{M}_n)_1^{\infty} \in \ell_1^2$. Thus we obtain

$$\eta_n = \mathfrak{t}_n + \mathcal{M}_n, \quad \forall \quad n \in \mathbb{N}.$$
(4.15)

The mapping $q \mapsto \tau^{(2)} \times \mathfrak{t}$ is a bijection between \mathcal{L} and $\mathfrak{J}^1(2) \times \ell_1^2$ and the mapping $\tau^{(2)} \times \mathfrak{t} \to \tau^{(2)} \times \eta$ is a bijection from $\mathfrak{J}^1(2) \times \ell_1^2$ onto itself, since η_n, \mathfrak{t}_n satisfy (4.15). This gives that the mapping $q \mapsto (\tau^{(2)} \times \eta)(q)$ is a bijection between \mathcal{L} and $\mathfrak{J}^1(2) \times \ell_1^2$.

• Consider the mapping $\tau^{(2)} \times \chi$. The identity (3.1) implies

$$\mathcal{N}_n = \dot{\vartheta}(1, \varrho_n)\vartheta'(1, \varrho_n), \qquad \forall \ n \in \mathbb{N}. \tag{4.16}$$

Then using $e^{-\mathfrak{r}_n} = (-1)^{n+1} \vartheta'(1,\varrho_n)/\sqrt{\varrho_n^o}$ we rewrite χ_n in the form

$$e^{-\chi_n} = \frac{\mathcal{N}_n(q)}{\mathcal{N}_n(0)} = \left[(-1)^n 2\sqrt{\varrho_n^o} \ \dot{\vartheta}(1,\varrho_n) \right] \left[(-1)^n \vartheta'(1,\varrho_n) / \sqrt{\varrho_n^o} \right] = e^{-\mathcal{K}_n - \mathfrak{r}_n}, \tag{4.17}$$

where $\mathcal{K}_n \in \mathbb{R}$ is defined by $e^{-\mathcal{K}_n} = (-1)^n 2\sqrt{\varrho_n^o} \ \dot{\vartheta}(1,\varrho_n) > 0$ and due to (7.3) the sequence $(\mathcal{K}_n)_1^{\infty} \in \ell_1^2$. Thus we obtain

$$\chi_n = \mathfrak{r}_n + \mathcal{K}_n, \quad \forall \quad n \in \mathbb{N}.$$
 (4.18)

The mapping $q \mapsto \varrho^{(2)} \times \mathfrak{r}$ is a bijection between \mathcal{L} and $\mathfrak{J}^1 \times \ell_1^2$ and the mapping $\varrho^{(2)} \times \mathfrak{r} \to \varrho^{(2)} \times \eta$ is a bijection from $\mathfrak{J}^1(2) \times \ell_1^2$ onto itself, since η_n, \mathfrak{r}_n satisfy (4.15). This gives that the mapping $q \mapsto (\varrho^{(2)} \times \chi)(q)$ is a bijection between \mathcal{L} and $\mathfrak{J}^1(2) \times \ell_1^2$.

5. Smooth potentials

5.1. Smooth potentials. Consider the case of potentials from Sobolev spaces \mathcal{L}_k given by

$$\mathcal{L}_k = \{q, q^{(k)} \in \mathcal{L}\}, \quad k \geqslant 0, \quad \mathcal{L} = \mathcal{L}_0.$$

Following the book of Pöschel and Trubowitz [34] in analogy to the notation O(1/n), we use the notation $\ell_k^2(n)$ for an arbitrary sequence of numbers which is an element of ℓ_k^2 :

$$y_n = y_n^o + \ell_k^2(n) \Longleftrightarrow \sum_{n \ge 1} n^{2k} |y_n - y_n^o|^2 < \infty.$$

We define the spectral data \mathfrak{J}_k for potentials from $\mathcal{L}_k, k \in \mathbb{N}$ by

$$\mathfrak{J}_k = \left\{ (s_n)_1^{\infty} \in \mathfrak{J} : \sqrt{s_n} = \frac{\pi n}{2} + \sum_{1 \le j \le d} \frac{a_j}{(\pi n)^{2j+1}} + \ell_{k+1}^2(n), \quad (a_j)_1^d \in \mathbb{R}^d \right\}, \quad d = \left[\frac{k+1}{2}\right],$$

here [r] is the integer part of $r \in \mathbb{R}$ and the coefficients a_j depend on a sequence $s = (s_n)_1^{\infty}$. We define two sets \mathfrak{J}_k^o and \mathfrak{J}_k^1 of all real, strictly increasing sequences by

$$\mathfrak{J}_{k}^{o} = \left\{ s = (s_{n})_{1}^{\infty} \in \mathfrak{J}^{o} : \sqrt{s_{n}} = \pi n + \sum_{1 \leq j \leq d} a_{j} \varepsilon^{2j+1} + \ell_{k+1}^{2}(n), \quad (a_{j})_{1}^{d} \in \mathbb{R}^{d} \right\}, \quad \varepsilon = \frac{1}{2\pi n},$$

$$\mathfrak{J}_k^1 = \left\{ (s_n)_1^\infty \in \mathfrak{J}^1 : \sqrt{s_n} = \pi(n - \frac{1}{2}) + \sum_{1 \le j \le d} b_j \delta_n^{2j+1} + \ell_{k+1}^2(n), \quad (b_j)_1^d \in \mathbb{R}^d \right\}, \ \delta_n = \frac{1}{2\pi(n - \frac{1}{2})},$$

Note that the coefficients a_j, b_j depend on a sequence $s = (s_n)_1^{\infty}$. Recall results from [32]: if $q \in \mathcal{L}_k$, then we have asymptotics of the Dirichlet eigenvalues μ_n :

$$\sqrt{\mu_n} = \pi n + \sum_{1 \le i \le d} a_i \varepsilon^{2j+1} + \varepsilon^{k+1} \widetilde{r}_n, \quad \widetilde{r}_n = q_{\bullet}^{(k)}(n) + \ell_1^2(n), \quad (\widetilde{r}_n)_1^{\infty} \in \ell^2, \tag{5.1}$$

where the coefficients $(a_j)_1^d \in \mathbb{R}^d$ depend on $q, d = \left[\frac{k+1}{2}\right]$, and \widetilde{r}_n has the form

$$q_{\bullet}^{(k)}(n) = (-1)^{1 + \left[\frac{k}{2}\right]} \int_{0}^{1} q^{(k)}(x) \mathcal{F}_{k}(nx) dx, \quad \mathcal{F}_{k}(nx) = \begin{cases} \sin[2\pi nx], & k+1 \in 2\mathbb{N} \\ \cos[2\pi nx], & k \in 2\mathbb{N} \end{cases} , \quad (5.2)$$

and asymptotics of the mixed eigenvalues τ_n :

$$\sqrt{\tau_n} = \pi n' + \sum_{1 \le j \le d} \widetilde{\tau}_j \delta_n^{2j+1} + \widetilde{q}_{\bullet}^{(k)}(n) \delta_n^{k+1} + \ell_{k+2}^2(n) \quad \text{as} \quad n \to \infty, \quad n' = n - \frac{1}{2},
\widetilde{q}_{\bullet}^{(k)}(n) = (-1)^{1+\left[\frac{k}{2}\right]} \int_0^1 q^{(k)}(x) \mathcal{F}_k(n'x) dx, \quad \mathcal{F}_k(nx) = \begin{cases} \sin[2\pi n'x], & k+1 \in 2\mathbb{N} \\ \cos[2\pi n'x], & k \in 2\mathbb{N} \end{cases},$$
(5.3)

where the coefficients $\widetilde{\tau}_j \in \mathbb{R}$, $j \in \mathbb{N}_d$ depend on q. We recall the famous results of Marchenko and Ostrovski (Corollary 5.1 from [32]) about the mapping $\tau \star \mu$ from \mathcal{L}_k to \mathfrak{J}_k .

Theorem 5.1. Each 2-spectra mapping $q \mapsto (\tau \star \mu)(q)$ is a bijection between \mathcal{L}_k and $\mathfrak{J}_k, k \in \mathbb{N}$.

Asymptotics (5.1), (5.3) give a 2-spectra mapping $q \to \tau \star \mu$ from \mathcal{L}_k to \mathfrak{J}_k for any $k \in \mathbb{N}$. We show that the mappings $\tau \star \mu$, $\varrho \star \nu$, $\tau \star \nu$ and $\varrho \star \mu$ are isomorphic bijections between \mathcal{L}_k and \mathfrak{J}_k . Moreover, we show that the mappings $\mu \times h_{(s)}$, $\nu \times \mathfrak{h}_{(s)}$, $\mu \times \mathfrak{h}_{(s)}$, $\nu \times h_{(s)}$ are isomorphic bijections between \mathcal{L}_k and $\mathfrak{J}_k^o \times \mathfrak{J}_k^e$.

Theorem 5.2. i) The mappings U_1, U_o are bijections of \mathcal{L}_k onto itself for any $k \in \mathbb{N}$.

- ii) All 2-spectra mappings $\varrho \star \nu$, $\tau \star \nu$, $\varrho \star \mu$ and $\tau \star \mu$ acting from \mathcal{L}_k into \mathfrak{J}_k are isomorphic for any $k \in \mathbb{N}$, and each of them is a bijection between \mathcal{L}_k and \mathfrak{J}_k and satisfy (1.19).
- iii) Mappings $\mu \times h_{(s)}$, $\nu \times \mathfrak{h}_{(s)}$, $\mu \times \mathfrak{h}_{(s)}$ and $\nu \times h_{(s)}$ (defined by (1.4), (1.6)) acting from \mathcal{L}_k into $\mathfrak{J}_k^o \times \mathfrak{J}_k^e$ are isomorphic for any $k \in \mathbb{N}$, and each of them is a bijection between \mathcal{L}_k and $\mathfrak{J}_k^o \times \mathfrak{J}_k^e$ and satisfy

$$\mu \times h_{(s)} = (\nu \times \mathfrak{h}_{(s)}) \circ U_1 = (\mu \times \mathfrak{h}_{(s)}) \circ U_o = (\nu \times h_{(s)}) \circ U_o \circ U_1. \tag{5.4}$$

Proof. i) Let $q \in \mathcal{L}_k$ for some $k \ge 1$. Repeating arguments from [32] for the case (5.1), (5.3) we determine asymptotics the eigenvalues $(\nu_n)_1^{\infty}$ and $(\varrho_n)_1^{\infty}$ of the boundary value problems y'(0) = y'(1) = 0 and y'(0) = y(1) = 0:

$$\sqrt{\nu_n} = \pi n + \sum_{1 \le j \le d} \widetilde{a}_j \varepsilon^{2j+1} + \ell_{k+1}^2(n), \quad \varepsilon = \frac{1}{2\pi n}, \tag{5.5}$$

$$\sqrt{\varrho_n} = \pi n' + \sum_{1 \le i \le d} \widetilde{b}_j \delta_n^{2j+1} + \ell_{k+1}^2(n), \quad \delta_n = \frac{1}{2\pi n'}, \quad n' = n - \frac{1}{2}$$
 (5.6)

for some constants $(\widetilde{a}_j)_1^d, (\widetilde{b}_j)_1^d \in \mathbb{R}^d$ depending on q.

Due to (1.19) the Dirichlet μ_n , Neumann ν_n and mixed τ_n , ϱ_n eigenvalues satisfy

$$(\mu_n, \nu_n, \tau_n, \varrho_n)(q) = (\nu_n, \mu_n, \varrho_n, \tau_n)(q^{\bullet}), \quad \forall n \geqslant 1, \text{ where } q^{\bullet} = U_1(q).$$
 (5.7)

Then the Dirichlet $\mu_n(q^{\bullet})$, and mixed $\tau_n(q^{\bullet})$ eigenvalues have the corresponding asymptotics (5.1), (5.3). Thus from Theorem 5.1 we deduce that $q^{\bullet} = U_1(q) \in \mathcal{L}_k$, i.e., $U_1\mathcal{L}_k \subset \mathcal{L}_k$, and $U_1^2 = I$ gives $U_1\mathcal{L}_k = \mathcal{L}_k$.

ii) We will show that 2-spectra mappings $\varrho \star \nu$, $\tau \star \nu$, $\varrho \star \mu$ and $\tau \star \mu$ acting from \mathcal{L}_k into \mathfrak{J}_k are isomorphic. Consider the 2-spectra mapping $\varrho \star \nu$ acting from \mathcal{L}_k into \mathfrak{J}_k . The proof for other mappings is similar. In i) we have obtained that if $q \in \mathcal{L}_k$, then the eigenvalues $(\nu_n)_1^{\infty}$ and $(\varrho_n)_1^{\infty}$ alternate and the asymptotics (5.5), (5.6) hold true, this yields $q \to \varrho \star \nu$ is a mapping from \mathcal{L}_k into \mathfrak{J}_k^{\bullet} .

In (5.7) we have an identity $\tau \star \mu = (\varrho \star \nu) \circ U_1$, which yields $(\tau \star \mu) \circ U_1 = \varrho \star \nu$. Then the mapping $\varrho \star \nu : \mathcal{L}_k \to \mathfrak{J}_k$ is a bijection between \mathcal{L}_k and \mathfrak{J}_k , since due to Theorem 5.1 the mapping $\tau \star \mu : \mathcal{L}_k \to \mathfrak{J}_k$ is a bijection between \mathcal{L}_k and \mathcal{J}_k and $U_1 : \mathcal{L}_k \to \mathcal{L}_k$ is a bijection from \mathcal{L}_k onto itself. Moreover, due to the identity $\tau \star \mu = (\varrho \star \nu) \circ U_1$ the mappings $\tau \star \mu$ and $\varrho \star \nu$ are isomorphic.

iii) Due to (5.7), (7.13) we have a mapping $q \mapsto (\mu \times h_{(s)})(q)$ from \mathcal{L}_k into $\mathfrak{J}_k^o \times \mathfrak{J}_k^e$ for any integer $k \in \mathbb{N}$. Theorem 1.3 gives an injection of this mapping.

In order to show a surjection of this mapping we use arguments from the proof of Theorem 5.1 from [32]. Let $\mu \times h_{(s)}^{\cdot} \in \mathfrak{J}_{k}^{o} \times \mathfrak{J}_{k}^{e}$ be given for some $k \geq 1$. Then due to Theorem 2.1, ii) there exist unique $q \in \mathcal{L}_{0}$ such that $\mu(q) \times h_{(s)}(q) = \mu \times h_{(s)}^{\cdot}$. Assume that $q \in \mathcal{L}_{m}$ but $q \notin \mathcal{L}_{m+1}$, where m < k. Further we actually repeat the proof from [32] verbatim and using sharp asymptotics (7.13) of $h_{s,n}$ and (5.1) of μ_{n} show that $q \in \mathcal{L}_{m+1}$, which gives a contradiction. Here it is important to determine the new sharp asymptotics of $h_{s,n}$ from (7.13) with the Fourier coefficients (7.14).

Using the identities (1.20), and the bijection of the mapping $q \to \mu(q) \times h_{(s)}(q)$ acting from \mathcal{L}_k into $\mathfrak{J}_k^o \times$ we have the proof of iii), since by i), the mappings U_1, U_o are bijections of \mathcal{L}_k onto itself for any $k \in \mathbb{N}$.

We discuss inverse problems for mixed eigenvalue mapping $\tau^{(2)} = (\tau_n)_2^{\infty}$ for potentials $q \in \mathcal{L}_k, k \in \mathbb{N}$. In this case we define the corresponding spectral data $\mathfrak{J}_k^1(2)$ by

$$\mathfrak{J}_k^1(2) = \left\{ s = (s_n)_2^{\infty} \in \mathfrak{J}^1 : s_n = \sum_{1 \le j \le d} \frac{a_j}{(2\pi n)^{2j+1}} + \ell_{k+1}^2(n), \quad (a_j)_1^d \in \mathbb{R}^d \right\}, \quad k \in \mathbb{N}, \quad (5.8)$$

and a set of all possible norming constants $\mathfrak{t} = (\mathfrak{t}_n)_1^{\infty}$ by

$$\mathfrak{J}_{k}^{e} = \left\{ s = (s_{n})_{1}^{\infty} \in \ell_{1}^{2} : s_{n} = \sum_{1 \leq j \leq d} \frac{\widetilde{a}_{j}}{(2\pi(n - \frac{1}{2})^{2j})} + \ell_{k+1}^{2}(n), \quad (\widetilde{a}_{j})_{1}^{d} \in \mathbb{R}^{d} \right\}, \quad k \in \mathbb{N},$$

Note that $(a_i)_1^d \in \mathbb{R}^d$ and $(\widetilde{a}_i)_1^d \in \mathbb{R}^d$ depends on q.

Theorem 5.3. Let $\tau^{(2)} = (\tau_n)_2^{\infty}$. The mapping $\tau^{(2)} \times \mathfrak{t}$ from \mathcal{L}_k into $\mathfrak{J}_k^1(2) \times \mathfrak{J}_k^e$ is a bijection between \mathcal{L}_k and $\mathfrak{J}_k^1(2) \times \mathfrak{J}_k^e$.

Proof. Due to (5.3), (7.25) we have a mapping $q \mapsto (\tau^{(2)} \times \mathfrak{t})(q)$ from \mathcal{L}_k into $\mathfrak{J}_k^1(2) \times \mathfrak{J}_k^e$ for any integer $k \in \mathbb{N}$. Theorem 2.1 iii) gives an injection of this mapping.

In order to show a surjection of this mapping we use arguments from the proof of Theorem 5.1 from [32]. Let $\tau_{\cdot} \times \mathfrak{t}_{\cdot} \in \mathfrak{J}_{k}^{1}(2) \times \mathfrak{J}_{k}^{e}$ be given for some $k \geqslant 1$. Then due to Theorem 2.1, ii) there exist unique $q \in \mathcal{L}_0$ such that $(\tau^{(2)} \times \mathfrak{t})(q) = \tau \times \mathfrak{t}$. Assume that $q \in \mathcal{L}_m$ but $q \notin \mathcal{L}_{m+1}$, where m < k. Further we actually repeat the proof from [32] verbatim and using sharp asymptotics (5.3) of τ_n and (7.25) of \mathfrak{t}_n show that $q \in \mathcal{L}_{m+1}$, which gives a contradiction. Here it is important to have the sharp asymptotics of τ_n , \mathfrak{t}_n from (5.3), (7.25) with the Fourier coefficients.

The identities (4.3), i.e., $\varrho^{(2)} \times \mathfrak{r} = (\tau^{(2)} \times \mathfrak{t}) \circ U_o$, and the bijection of the mapping $q \to 0$ $(\tau^{(2)} \times \mathfrak{t})(q)$ between \mathcal{L}_k and $\mathfrak{J}_k^1(2) \times \mathfrak{J}_k^e$ imply that the mapping $q \to (\varrho^{(2)} \times \mathfrak{r})(q)$ is a bijection between \mathcal{L}_k and $\mathfrak{J}_k^1(2) \times \mathfrak{J}_k^e$, since $U_o = \mathcal{R}$ is an unitary operator on \mathcal{L}_k for any $k \in \mathbb{N}$.

6. Periodic problems

6.1. **Periodic potentials.** We consider inverse problems on the circle. Firstly we define the gap mapping $q \to \psi = (\psi_n)^{\infty}$ acting from \mathcal{H} into $\ell^2 \oplus \ell^2$ from [22]. The components $\psi_n \in \mathbb{R}^2$ are constructed via the periodic plus Dirichlet eigenvalues plus signs by

$$\psi_{n} = (p_{c,n}, \psi_{s,n}) \in \mathbb{R}^{2}, \qquad |\psi_{n}|^{2} = \psi_{c,n}^{2} + \psi_{s,n}^{2} = \frac{1}{4}(\lambda_{n}^{+} - \lambda_{n}^{-})^{2},
\psi_{c,n} = \frac{1}{2}(\lambda_{n}^{+} + \lambda_{n}^{-}) - \mu_{n}, \quad \psi_{s,n} = \left||\psi_{n}|^{2} - \psi_{c,n}^{2}|^{\frac{1}{2}} \operatorname{sign} h_{s,n}, \quad h_{s,n} = \log|\varphi'(1, \mu_{n})|.$$
(6.1)

The mapping ψ is a RAB between \mathcal{H} and $\ell^2 \oplus \ell^2$, see Theorem 6.1 below.

We define another gap mapping $\mathfrak{p}:\mathcal{H}\to\ell^2\oplus\ell^2$ by $q\to\mathfrak{p}=(\mathfrak{p}_n)_1^\infty$. The components $\mathfrak{p}_n \in \mathbb{R}^2$ are constructed via the periodic plus Neumann eigenvalues plus signs by

$$\mathfrak{p}_{n} = (\mathfrak{p}_{c,n}, \mathfrak{p}_{s,n}) \in \mathbb{R}^{2}, \quad |\mathfrak{p}_{n}|^{2} = \mathfrak{p}_{c,n}^{2} + \mathfrak{p}_{s,n}^{2} = \frac{1}{4}(\lambda_{n}^{+} - \lambda_{n}^{-})^{2},
\mathfrak{p}_{c,n} = \frac{1}{2}(\lambda_{n}^{+} + \lambda_{n}^{-}) - \nu_{n}, \qquad \mathfrak{p}_{s,n} = \left| |\mathfrak{p}_{n}|^{2} - (\mathfrak{p}_{c,n})^{2} \right|^{\frac{1}{2}} \operatorname{sign} \mathfrak{h}_{s,n}, \quad \mathfrak{h}_{s,0} = \ln |\vartheta(1,\nu_{0})|.$$
(6.2)

Secondly we consider inverse problems in terms of local maxima and minima of the Lyapunov function, given by $\Delta(\lambda) = \frac{1}{2}(\varphi'(1,\lambda) + \vartheta(1,\lambda))$. The Lyapunov function on the real line has local maxima and minima at points $\lambda_n \in [\lambda_n^-, \lambda_n^+]$ for all $n \in \mathbb{N}$, where $(-1)^n \Delta(\lambda_n^{\pm}) = 1$ and $(-1)^n \Delta(\lambda_n) \ge 1$. Define the corresponding mapping $h : \mathcal{H} \to \ell^2 \oplus \ell^2$ as $h : q \to h = ([2\pi n]h_n)_1^{\infty}$ from [32]. The components $h_n \in \mathbb{R}^2$ are constructed via maxima and minima of the Lyapunov function plus Dirichlet eigenvalues plus signs by

$$h_{n} = (h_{c,n}, h_{s,n}) \in \mathbb{R}^{2}, \qquad |h_{n}|^{2} = h_{c,n}^{2} + h_{s,n}^{2},$$

$$h_{c,n} = \left| |h_{n}|^{2} - h_{s,n}^{2} \right|^{\frac{1}{2}} \operatorname{sign}(\lambda_{n} - \mu_{n}), \qquad h_{s,n} = \log |\varphi'(1, \mu_{n})|,$$

$$(6.3)$$

here the value $|h_n| \ge 0$ is uniquely defined by the equation $\cosh |h_n| = |\Delta(\lambda_n)| \ge 1$. Recall that $(-1)^n \Delta(\mu_n) = \cosh h_{s,n}$ for all $n \ge 1$ and $|h_n| \ge |h_{s,n}|$, since $(-1)^n \Delta$ has the maximum at λ_n on the segment $[\lambda_n^-, \lambda_n^+]$. The mapping h is a RAB between \mathcal{H} and $\ell^2 \oplus \ell^2$, see below. We introduce similar mapping $\mathfrak{h} : \mathcal{H} \to \ell^2 \oplus \ell^2$ as $\mathfrak{h} : q \to \mathfrak{h}(q) = ([2\pi n]\mathfrak{h}_n(q))_1^{\infty}$. The

components $\mathfrak{h}_n \in \mathbb{R}^2$ are constructed via maxima and minima of the Lyapunov function plus

Neumann eigenvalues plus signs by

$$\mathfrak{h}_{n} = (\mathfrak{h}_{c,n}, \mathfrak{h}_{s,n}) \in \mathbb{R}^{2}, \quad |\mathfrak{h}_{n}| = |h_{n}|,$$

$$\mathfrak{h}_{c,n} = \left| |\mathfrak{h}_{n}|^{2} - \mathfrak{h}_{s,n}^{2} \right|^{\frac{1}{2}} \operatorname{sign}(\lambda_{n} - \nu_{n}), \qquad \mathfrak{h}_{s,n} = \log |\vartheta(1, \nu_{n})|.$$
(6.4)

Recall that $(-1)^n \Delta(\nu_n) = \cosh \mathfrak{h}_{s,n}$ for all $n \ge 1$ and $|\mathfrak{h}_n| \ge |\mathfrak{h}_{s,n}|$, since $(-1)^n \Delta$ has the local maximum at λ_n on the segment $[\lambda_n^-, \lambda_n^+]$. We describe well-known results about inverse problems on the circle.

Theorem 6.1. i) The mapping $h = ([2\pi n]h_n)_1^{\infty} : \mathcal{H} \to \ell^2 \oplus \ell^2$ given by (6.3) is a RAB between \mathcal{H} and $\ell^2 \oplus \ell^2$. Furthermore, the following estimates hold true:

$$||q|| \le 3||h||(6+h_+)^{\frac{1}{2}}, \qquad ||h|| \le 2||q||(1+||q||^{\frac{1}{3}}),$$

$$(6.5)$$

where $||q||^2 = \int_0^1 q^2(x) dx$ and $||h||^2 = \sum_{n \ge 1} |2\pi n h_n|^2$ and $h_+ = \sup_{n \ge 1} |h_n|$. ii) The mapping $\psi : \mathcal{H} \to \ell^2 \oplus \ell^2$ given by (6.1) is a RAB between \mathcal{H} and $\ell^2 \oplus \ell^2$. Furthermore, the following estimates hold true:

$$||q|| \le 2||\psi||(1+||\psi||^{\frac{1}{3}}), \qquad ||\psi|| \le ||q||(1+||q||^{\frac{1}{3}}),$$

$$(6.6)$$

where $\|\psi\|^2 = \sum_{n\geq 1} (\psi_{c,n}^2 + \psi_{s,n}^2) = \frac{1}{4} \sum_{n\geq 1} |\lambda_n^+ - \lambda_n^-|^2$.

Remark. 1) A bijection in i) was proved in [32]. It was reproved in [24], including the RAB. The proof in [24] is simpler and is based on analytic approach from [16], [34].

- 2) A bijection of ψ was proved in [22]. The proof is sufficiently short, since the estimates (6.6) from [23] were used. Note that there is a unique way of placing the sequence of open tiles of lengths $\lambda_n^+ - \lambda_n^-, n \ge 1$, in order on the half line $[\lambda_0^+, \infty)$ so that the compliment is the set of bands for a function $q \in \mathcal{H}$, so that they are genuine gaps, see [22]. It does not depend on the positions of the Dirichlet spectrum $\{\mu_n\}$ and $\{\operatorname{sign} h_{s,n}\}$.
- 3) The estimates (6.5), (6.6) were obtained in [23], [20] (some estimates of h were determined in [32]). Their proof is based on the conformal mapping theory and trace formulas [24]. We describe inverse problems on the circle.

Theorem 6.2. i) The mappings $\mathfrak{p}: \mathcal{H} \to \ell^2 \oplus \ell^2$ and $\psi: \mathcal{H} \to \ell^2 \oplus \ell^2$ given by (6.2), (6.1) are isomorphic. Moreover, $\mathfrak p$ is a RAB between $\mathcal H$ and $\ell^2 \oplus \ell^2$ and satisfies

$$\mathfrak{p} = \psi \circ U_1. \tag{6.7}$$

ii) The mappings $\mathfrak{h}:\mathcal{H}\to\ell^2\oplus\ell^2$ and $h:\mathcal{H}\to\ell^2\oplus\ell^2$ given by (6.4), (6.3) are isomorphic. Moreover, $\hat{\mathfrak{h}}$ is a RAB between \mathcal{H} and $\ell_1^2 \oplus \ell_1^2$ and satisfies

$$\mathfrak{h} = h \circ U_1. \tag{6.8}$$

iii) Let $\sigma = (\sigma_n)_1^{\infty}$, where $\sigma_{2n} \in \{\pm 1\}$ and $\sigma_{2n-1} = -1$ for all $n \in \mathbb{N}$. Define replacing mappings $q \to \phi = (\phi_n)_1^{\infty}$ and $q \to \omega = ([2\pi n]\omega_n)_1^{\infty}$, where

$$\phi_n = \begin{cases} \psi_n \\ \mathfrak{p}_n \end{cases}, \qquad \omega_n = \begin{cases} h_n, & \text{if } \sigma_{2n} = 1 \\ \mathfrak{h}_n, & \text{if } \sigma_{2n} = -1 \end{cases}. \tag{6.9}$$

Then ϕ is a RAB between \mathcal{H} and $\ell^2 \oplus \ell^2$ and ω is a RAB between \mathcal{H} and $\ell^2 \oplus \ell^2$ and satisfy

$$\phi = \psi \circ \mathcal{U}_{\sigma}, \qquad \omega = h \circ \mathcal{U}_{\sigma}. \tag{6.10}$$

Remark. 1) The mapping U_1 from this theorem is iso-spectral for the periodic eigenvalues, but not for the Dirichlet and Neumann eigenvalues, see Lemma 2.5. **Proof.** i) Let $q \in \mathcal{H}$. Due to (2.14) 2-periodic eigenvalues $\{\lambda_n^+, \lambda_n^\pm, n \ge 1\}$ are invariant under U_1 and (2.18) gives $\mu_n(q) = \nu_n(q^{(1)})$ for all $n \in \mathbb{N}$, where $q^{(1)} = U_1 q$. This yields

$$p_{c,n}(q) = \left(\frac{\lambda_n^- + \lambda_n^+}{2} - \mu_n\right)(q) = \left(\frac{\lambda_n^- + \lambda_n^+}{2} - \nu_n\right)(q^{(1)}) = \mathfrak{p}_{c,n}(q^{(1)}),$$

$$p_{s,n}(q) = \left(\left||p_n|^2 - (p_{c,n})^2\right|^{\frac{1}{2}} \operatorname{sign} h_{s,n}\right)(q) = \left(\left||\mathfrak{p}_n|^2 - (\mathfrak{p}_{c,n})^2\right|^{\frac{1}{2}} \operatorname{sign} \mathfrak{h}_{s,n}\right)(q^{(1)}) = \mathfrak{p}_{s,n}(q^{(1)}),$$

since (2.22) gives $h_{s,n}(q) = \mathfrak{h}_{s,n}(q^{(1)})$. Then $p = \mathfrak{p} \circ U_1$ and from Theorem 6.1 we deduce that \mathfrak{p} is a RAB between \mathcal{H} and $\ell^2 \oplus \ell^2$, since due to Proposition 1.1 U_1 is a RAB of \mathcal{L} onto itself. The proof of ii) is similar to the case i).

iii) Let $q^{\bullet} = \mathcal{U}_{\sigma}(q)$, where $q \in \mathcal{H}$. Let $n = 2j - 1 \geqslant 1$ for all $j \in \mathbb{N}$ and $\sigma_n = -1$. Due to (2.15) under the mapping \mathcal{U}_{σ} all mixed eigenvalues satisfy

$$\tau_n(q) = \varrho_n(q^{\bullet}), \quad \varrho_n(q) = \tau_n(q^{\bullet}),$$
(6.11)

and then the identities (2.13) imply

$$\varphi'(1,\cdot,q) = \vartheta(1,\cdot,q^{\bullet}), \quad \vartheta(1,\cdot,q) = \varphi'(1,\cdot,q^{\bullet}), \tag{6.12}$$

which jointly with (1.6) yields $h_{s,n}(q) = h_{s,n}(q^{\bullet})$ for all $n \ge 1$.

Let n=2j for all $j\in\mathbb{N}$. Then (2.16), (2.15) yields

$$\begin{cases} \mu_{j}(q) = \mu_{j}(q^{\bullet}), & \nu_{j}(q) = \nu_{j}(q^{\bullet}), & if \ \sigma_{2j} = 1\\ \mu_{j}(q) = \nu_{j}(q^{\bullet}), & \nu_{j}(q) = \mu_{j}(q^{\bullet}), & if \ \sigma_{2j} = -1 \end{cases}$$
(6.13)

Thus we have two cases:

A) If $\sigma_{2j} = 1$, then from (6.13), (6.12) we have $\phi_j(q^{\bullet}) = p_j(q^{\bullet})$, where

$$h_{s,j}(q^{\bullet}) = \ln |\varphi'(1,\mu_j(q^{\bullet}),q^{\bullet})| = \ln |\varphi'(1,\mu_j(q),q)| = h_{s,j}(q),$$

$$p_{cj}(q^{\bullet}) = \left(\frac{\lambda_n^- + \lambda_n^+}{2} - \mu_n\right)(q^{\bullet}) = \left(\frac{\lambda_n^- + \lambda_n^+}{2} - \mu_n\right)(q) = p_{cj}(q),$$

$$p_{sj}(q^{\bullet}) = \left(\left| |p_j|^2 - (p_{c,j})^2 \right|^{\frac{1}{2}} \operatorname{sign} h_{s,j} \right) (q^{\bullet}) = \left(\left| |p_j|^2 - (p_{c,j})^2 \right|^{\frac{1}{2}} \operatorname{sign} h_{s,j} \right) (q) = p_{s,j}(q).$$

B) If $\sigma_{2j} = -1$, then from (6.13), (6.12) we have $\phi_j(q^{\bullet}) = \mathfrak{p}_j(q^{\bullet})$, where

$$\mathfrak{h}_{s,j}(q^{\bullet}) = \ln |\vartheta(1,\nu_j(q^{\bullet}),q^{\bullet})| = \ln |\varphi'(1,\mu_j(q),q)| = h_{s,j}(q),$$

$$\mathfrak{p}_{cj}(q^{\bullet}) = \left(\frac{\lambda_n^- + \lambda_n^+}{2} - \nu_n\right)(q^{\bullet}) = \left(\frac{\lambda_n^- + \lambda_n^+}{2} - \mu_n\right)(q) = p_{cj}(q),$$

$$\mathfrak{p}_{sj}(q^{\bullet}) = \left(\left| |\mathfrak{p}_{j}|^{2} - (\mathfrak{p}_{c,j})^{2} \right|^{\frac{1}{2}} \operatorname{sign} \mathfrak{h}_{s,j} \right) (q^{\bullet}) = \left(\left| |p_{j}|^{2} - (p_{c,j})^{2} \right|^{\frac{1}{2}} \operatorname{sign} h_{s,j} \right) (q) = p_{s,j}(q).$$

From A and B we obtain $p = \phi \circ \mathcal{U}_{\sigma}$ and ϕ is a RAB between \mathcal{H} and $\ell^2 \oplus \ell^2$, since due to Proposition 1.1 U_{σ} is a RAB of \mathcal{L} onto itself. The proof for mapping ω is similar.

6.2. Smooth periodic potentials. We discuss isomorphic inverse problems for the case of smooth periodic potentials from Sobolev spaces \mathcal{H}_k defined by

$$\mathcal{H}_k = \{q, q^{(k)} \in \mathcal{H}\}, \quad k \geqslant 0.$$

Theorem 6.3. i) The mappings $\psi : \mathcal{H}_k \to \ell_k^2 \oplus \ell_k^2$ and $\mathfrak{p} : \mathcal{H}_k \to \ell_k^2 \oplus \ell_k^2$ given by (6.1), (6.2) are isomorphic and are bijections between \mathcal{H}_k and $\ell_k^2 \oplus \ell_k^2$. They satisfy $\mathfrak{p} = \psi \circ U_1$ and

$$\|\psi\|_{k} \leqslant C_{1} \|q\|_{(k)} (1 + \|q\|_{(k)})^{2s+1},$$

$$\|q\|_{(k)} \leqslant C_{2} \|\psi\|_{k} (1 + \|\psi\|_{k})^{2m(1+s)+s},$$

(6.14)

where $||q||_{(k)}^2 = \int_0^1 |q^{(k)}|^2 dx$ and $||\psi||_k^2 = \sum_{n \ge 1} (2\pi n)^{2k} |\psi_n|^2$, $||\psi_n|| = \frac{\lambda_n^+ - \lambda_n^-}{2}$ and $s = \frac{2k+1}{3}$, $m = \frac{k+1}{3} (1 + tk + t^2 k(k-1) + \dots + t^{k+1} k!)$, $t = \frac{2}{3}$. (6.15)

ii) The mappings $h: \mathcal{H}_k \to \ell_k^2 \oplus \ell_k^2$ and $\mathfrak{h}: \mathcal{H}_k \to \ell_k^2 \oplus \ell_k^2$ given by (6.1), (6.2) are isomorphic and are bijections between \mathcal{H}_k and $\ell_k^2 \oplus \ell_k^2$. They satisfy $\mathfrak{h} = h \circ U_1$ and

$$||h||_{k} \leq C_{3}||q||_{(k)} (1 + ||q||_{(k)})^{2s+1},$$

$$||q||_{(k)} \leq C_{4}||h||_{k} (1 + ||h||_{k})^{m} (1 + \sup_{j \in \mathbb{N}} |h_{j}|)^{(k+1)(1+m)}.$$
(6.16)

The constants $C_1, ..., C_4$ depend on k only and $||h||_k^2 = \sum_{n \geqslant 1} (2\pi n)^{2k} |2\pi n h_n|^2$.

Remark. The mapping $h: \mathcal{H}_k \to \ell_k^2 \oplus \ell_k^2, k \in \mathbb{N}$ is a bijection, see [32]. All estimates (6.14), (6.16) are new. There is an open problem about their sharpness, even for potentials $q \in \mathcal{H}$. **Proof.** i) If $q \in \mathcal{H}_k$, then estimates (6.14) yield that $\psi(q) \in \ell_k^2 \oplus \ell_k^2$. The mapping $\psi: \mathcal{H}_k \to \ell_k^2 \oplus \ell_k^2$ is an injection, since due to Theorem 6.1, ii) the mapping $\psi: \mathcal{H} \to \ell^2 \oplus \ell^2$ is a bijection. Let $b \in \ell_k^2 \oplus \ell_k^2$. Then due to Theorem 6.1, ii) there exists a unique $q \in \mathcal{H}_0$, such that $\psi(q) = b$. Thus using the following results: if $q \in \mathcal{H}$ and $k \in \mathbb{N}$, then

$$(|\psi_n|)_1^{\infty} \in \ell_k^2 \quad \text{or} \quad h \in \ell_k^2 \qquad \Leftrightarrow \qquad q \in \mathcal{H}_k.$$
 (6.17)

see e.g., Corollary 3.4 in [32] or [19], we deduce that $q \in \mathcal{H}_k$.

We show (6.14). We need estimates (see Theorem 2.3 and Theorem 2.6 from [19])

$$Q_k \leqslant c_1^2 \|\psi\|_k^2 (1 + \|\psi\|_k^{2s}), \tag{6.18}$$

$$\|\gamma\|_k^2 \leqslant c_2^2 Q_k (1 + Q_k),$$
 (6.19)

where $Q_k \geqslant 0$ is some non-linear functional of $q, q', ..., q^{(k)}$ and

$$Q_k \leqslant c_3^2 \|q^{(k)}\|^2 (1 + \|q^{(k)}\|)^{2s},$$
 (6.20)

$$||q^{(k)}|| \le c_4^2 Q_k^{\frac{1}{2}} (1 + Q_k^m),$$
 (6.21)

for some constants $c_1, ..., c_4$ depending on k only. Let $A = Q_k^{\frac{1}{2}} \ge 0$. Due to an inequality $(1+x)^{\frac{1}{2}} \le 1+x^{\frac{1}{2}}$ for $x \ge 0$ we rewrite these estimates in the form:

$$A \le c_1 \|\psi\|_k (1 + \|\psi\|_k)^s, \tag{6.22}$$

$$\|\psi\|_k \leqslant c_2 A(1+A),\tag{6.23}$$

and

$$A \leqslant c_3 \|q^{(k)}\| \left(1 + \|q^{(k)}\|\right)^s, \tag{6.24}$$

$$||q^{(k)}|| \le c_4 A (1 + A^{2m}),$$
 (6.25)

• Let A < 1. Then (6.23), (6.25) and (6.22) yield

$$\|\psi\|_{k} \leq 2c_{2}A \leq 2c_{2}c_{3}\|q^{(k)}\|\left(1+\|q^{(k)}\|\right)^{s},$$

$$\|q^{(k)}\| \leq 2c_{4}A \leq 2c_{4}c_{1}\|\psi\|_{k}\left(1+\|\psi\|_{k}\right)^{s}.$$
 (6.26)

• Let $A \ge 1$. Then (6.23), (6.25) and (6.22) yield

$$\|\psi\|_{k} \leqslant 2c_{2}A^{2} \leqslant 2c_{2}c_{3}^{2}\|q^{(k)}\|^{2} (1 + \|q^{(k)}\|)^{2s} \leqslant 2c_{2}c_{3}^{2}\|q^{(k)}\| (1 + \|q^{(k)}\|)^{2s+1},$$

$$\|q^{(k)}\| \leqslant 2c_{4}A^{1+2m} \leqslant 2c_{4}c_{1}^{1+2m}\|\psi\|_{k} (1 + \|\psi\|_{k})^{s(1+2m)+2m}.$$

$$(6.27)$$

Estimates (6.26), (6.27) imply (6.14).

ii) The proof for mappings h and \mathfrak{h} is similar, we need only to show (6.16). Let $q \in \mathcal{H}_k, k \in \mathbb{N}$. We need the following estimates (see Theorem 2.1 from [19]):

$$||h||_k^2 \le b^2 Q_k (1 + Q_k), \quad b^2 = 4^{8k+11},$$

$$Q_k \le \frac{1}{\pi (k+1)} h_+^{2(k+1)} ||h||_k^2, \qquad h_+ = \sup_{j \in \mathbb{N}} |h_j|.$$
(6.28)

Due to an inequality $(1+A^2)^{\frac{1}{2}} \leqslant 1+A$ for $A=Q_k^{\frac{1}{2}} \geqslant 0$ we rewrite (6.28) in the form:

$$||h||_{k} \leqslant bA(1+A),$$

$$A \leqslant \sigma_{*}||h||_{k}h_{+}^{k+1}, \qquad \sigma_{*} = (\pi(k+1))^{-\frac{1}{2}}.$$
(6.29)

• Let A < 1. Then (6.29), (6.24) and (6.25) yield

$$||h||_{k} \leqslant 2bA \leqslant 2bc_{3}||q^{(k)}|| (1 + ||q^{(k)}||)^{s},$$

$$||q^{(k)}|| \leqslant 2c_{4}A \leqslant 2c_{4}\sigma_{*}||h||_{k}h_{+}^{k+1}.$$
 (6.30)

• Let $A \ge 1$. Then (6.29), (6.24) and (6.25) yield

$$||h||_{k} \leqslant 2bA^{2} \leqslant 2bc_{3}^{2}||q^{(k)}||^{2} (1 + ||q^{(k)}||)^{2s} \leqslant 2bc_{3}^{2}||q^{(k)}|| (1 + ||q^{(k)}||)^{1+2s},$$

$$||q^{(k)}|| \leqslant 2c_{4}A^{1+m} \leqslant 2c_{4}\sigma_{*}^{1+m}||h||_{k}^{1+m}h_{+}^{(k+1)(1+m)}.$$

$$(6.31)$$

Estimates (6.30), (6.31) imply (6.16).

7. Appendix: Asymptotics and trace formulas

7.1. **Asymptotics.** We discuss asymptotics of fundamental solutions.

Proposition 7.1. Let $q \in \mathcal{L}$. Then following asymptotics as $n \to \infty$ hold true:

$$\ln\left[(-1)^n 2\mu_n^o \dot{\varphi}(1,\mu_n)\right] = \ell_1^2(n),\tag{7.1}$$

$$\ln\left[(-1)^{n} 2\sqrt{\tau_{n}^{o}} \,\dot{\varphi}'(1,\tau_{n})\right] = \ell_{1}^{2}(n),\tag{7.2}$$

$$\ln\left[(-1)^n 2\sqrt{\varrho_n^o} \ \dot{\vartheta}(1,\varrho_n)\right] = \ell_1^2(n). \tag{7.3}$$

Proof. Asymptotics (7.1) was proved in [2]. We show (7.2). Recall that the sequence $\hat{\tau}_n = \tau_n - \tau_n^o, n \geqslant 1$ belongs to ℓ^2 . Using (2.13) and $\dot{\varphi}'_o(1, \tau_n^o) = \frac{(-1)^n}{2\sqrt{\tau_n^o}}$ at q = 0 we obtain

$$\dot{\varphi}'(1,\tau_n) = \frac{-1}{\tau_n^o} \prod_{j \neq n}^{+\infty} \frac{\tau_j - \tau_n}{\tau_j^o} = \frac{(-1)^n}{2\sqrt{\tau_n^o}} \prod_{j \neq n}^{+\infty} \frac{\tau_j - \tau_n}{\tau_j^o - \tau_n^o} = \frac{(-1)^n}{2\sqrt{\tau_n^o}} \prod_{j \neq n}^{+\infty} \left[1 + \frac{\hat{\tau}_j - \hat{\tau}_n}{\tau_j^o - \tau_n^o} \right]. \tag{7.4}$$

Then

$$\log(-1)^{n} 2\sqrt{\tau_{n}^{o}} \dot{\varphi}'(1, \tau_{n}) = \log \prod_{j \neq n}^{+\infty} \left[1 + \frac{\hat{\tau}_{j} - \hat{\tau}_{n}}{\tau_{j}^{o} - \tau_{n}^{o}} \right] = \sum_{j \neq n}^{+\infty} \left[\frac{\hat{\tau}_{j} - \hat{\tau}_{n}}{\tau_{j}^{o} - \tau_{n}^{o}} + \frac{O(1)}{(\tau_{j}^{o} - \tau_{n}^{o})^{2}} \right]$$

$$= \sum_{j \neq n}^{+\infty} \frac{\hat{\tau}_{j} - \hat{\tau}_{n}}{\tau_{j}^{o} - \tau_{n}^{o}} + \frac{O(1)}{n^{2}} = \sum_{j \neq n}^{+\infty} \frac{\hat{\tau}_{j}}{\tau_{j}^{o} - \tau_{n}^{o}} - \frac{\hat{\tau}_{n}}{4\tau_{n}^{o}} + \frac{O(1)}{n^{2}} = \sum_{j \neq n}^{+\infty} \frac{\hat{\tau}_{j}}{\tau_{j}^{o} - \tau_{n}^{o}} + \frac{O(1)}{n^{2}},$$

$$(7.5)$$

since due to $\frac{1}{\tau_i^o - \tau_n^o} = \frac{1}{2\pi\sqrt{\tau_n^o}} \left(\frac{1}{j-n} - \frac{1}{j+n-1}\right)$ and $\frac{1}{2n-1} = \frac{\pi}{2\sqrt{\tau_n^o}}$ we have

$$\sum_{j\neq n}^{+\infty} \frac{1}{\tau_j^o - \tau_n^o} = \lim_{m \to \infty} \sum_{j\neq n}^m \frac{1}{\tau_j^o - \tau_n^o} = \frac{1}{2\pi\sqrt{\tau_n^o}} \lim_{m \to \infty} A_m, \quad A_m = \sum_{j=1, j\neq n}^m \left(\frac{1}{j-n} - \frac{1}{j+n-1}\right),$$

$$A_m = \frac{1}{2n-1} + \sum_{j=-m, j\neq n}^m \frac{1}{j-n} = \frac{\pi}{2\sqrt{\tau_n^o}} + o(1) \quad \text{as} \quad m \to \infty.$$
(7.6)

Let $\hat{\tau} = (\hat{\tau}_j)_{j \in \mathbb{Z}}$, where $\hat{\tau}_{\pm j} = \hat{\tau}_j$ for $j \in \mathbb{N}$ and $\hat{\tau}_0 = 0$. The last sum in (7.5) satisfies

$$\sum_{j\neq n}^{+\infty} \frac{\hat{\tau}_j}{\tau_j^o - \tau_n^o} = \frac{1}{2\pi\sqrt{\tau_n^o}} \sum_{j\neq n}^{+\infty} \left(\frac{\hat{\tau}_j}{j-n} - \frac{\hat{\tau}_j}{j+n-1}\right)$$

$$= \frac{1}{2\pi\sqrt{\tau_n^o}} \left[\frac{\pi\hat{\tau}_n}{2\sqrt{\tau_n^o}} + \sum_{j\in\mathbb{Z}\setminus\{n\}} \frac{\hat{\tau}_j}{j-n}\right] = \frac{\hat{\tau}_n}{4\tau_n^o} + \frac{1}{2\pi\sqrt{\tau_n^o}} (\mathcal{F}\hat{\tau})_n$$

$$(7.7)$$

where \mathcal{F} is the linear operator on $\ell^2(\mathbb{Z})$ and given by $(\mathcal{F}\hat{\tau})_n = \sum_{j \in \mathbb{Z} \setminus \{n\}} \frac{\hat{\tau}_j}{j-n}$. Due to the identity $\sum_{j \neq 0} \frac{e^{i2\pi jt}}{j} = -i(t-\frac{1}{2}), t \in (0,1)$ and the Fourier transform we deduce that \mathcal{F} is the bounded operator in $\ell^2(\mathbb{Z})$. Thus jointly (7.5), (7.7) it gives (7.2).

We show (7.3). Let $q^o = U_o(q)$. From (2.29) we obtain $\vartheta(1,\cdot,q) = \varphi'(1,\cdot,q^o)$. Then (2.28) gives $\vartheta(1,\varrho_n(q),q) = \varphi'(1,\tau_n(q^o),q^o)$ and Theorem 2.1 iii) implies (7.3).

In order to determined asymptotics of norming constants we need asymptotics of fundamental solutions. Recall that a solution of $-y'' + qy = z^2y, z > 1$ has the form

$$y(x,z) = e^{izx}Y(x,z), \quad Y(x,z) = 1 + \sum_{1}^{k} \frac{u_j(x)}{\varsigma^j} + \frac{\tilde{u}_k(x,z)}{\varsigma^{k+1}}, \quad \varsigma = i2z,$$
 (7.8)

see Sect. 3 in [32], where $u_1(x) = \int_0^x q(t)dt, \dots$ and

$$\widetilde{u}_k \in \mathcal{L}_2, \quad \widetilde{u}_k(0, z) = \widetilde{u}'_k(0, z) = u_j(0) = 0, \quad u_j \in \mathcal{L}_2, \quad j \in \mathbb{N}_{k+1},$$

$$(7.9)$$

$$\widetilde{u}_k(1,z) = u_{k+1}(1) - q_*^k(z) + \frac{1}{\varsigma} (C_q + c_q e^{-i2z} + \widehat{K}(z)), \quad c_q, C_q \in \mathbb{R},$$

$$q_*^k(z) = (-1)^k e^{-i2z} \int_0^1 e^{i2zt} q^{(k)}(t) dt, \quad \widehat{K}(z) = \int_0^1 e^{-i2zt} g(t) dt, \quad K \in L^2(0,1),$$

$$(7.10)$$

We can rewrite φ, ϑ in terms of $y(x, \pm z)$. For example, we compute the identities

$$\vartheta(1,z) = \frac{1}{w(z)} (y_0'(z)y_1(-z) - y_0'(-z)y_1(z)), \qquad \varphi(1,z) = \frac{1}{w(z)} (y_1(z) - y_1(-z)),$$
where $y_0'(z) = y'(0,z), \quad y_1(z) = y(1,z), \quad w(z) = y_0'(z) - y_0'(-z).$ (7.11)

We rewrite the function $y'_0(z)$, the Wronskian w(z) due to (7.8), (7.9) in terms of $1/\varsigma$:

$$y_0'(z) = iz + \sum_{1}^{k} \frac{u_j'(0)}{\varsigma^j} = \varsigma y_{\cdot}(z), \quad y_{\cdot}(\varsigma) = \frac{1}{2} + \sum_{1}^{k} \frac{2u_j'(0)}{\varsigma^{j+1}},$$

$$w(z) = \varsigma + \sum_{1}^{k} \frac{u_j'(0)}{\varsigma^j} \left(1 - (-1)^j\right) = \varsigma w_{\cdot}(\varsigma), \quad w_{\cdot}(\varsigma) = 1 + \sum_{2 \leqslant 2j \leqslant k+1} \frac{2u_{2j-1}'(0)}{\varsigma^{2j}}.$$

$$(7.12)$$

Lemma 7.2. Let $q \in \mathcal{L}_k, k \in \mathbb{N}$. Then the norming constants $h_{s,n} = \ln |\varphi'(1, \mu_n, q)|$ satisfy

$$h_{s,n} = \sum_{1 \le j \le d} \widetilde{\varphi}_j \varepsilon^{2j} + E_n \varepsilon^{k+1} + \ell_{k+2}^2(n) \quad \text{as} \quad n \to \infty,$$
 (7.13)

where $\varepsilon = \frac{1}{2\pi n}$, the coefficients $\widetilde{\varphi}_j \in \mathbb{R}$, $j \in \mathbb{N}_d$, $d = \left[\frac{k+1}{2}\right]$ depend on q and

$$E_n = (-1)^{1+\left[\frac{k}{2}\right]} \int_0^1 q^{(k)}(x) \mathcal{G}_k(nx) dx, \quad \mathcal{G}_k(nx) = \begin{cases} \sin[2\pi nx], & k \in 2\mathbb{N} \\ \cos[2\pi nx], & k+1 \in 2\mathbb{N} \end{cases}$$
 (7.14)

Proof. From (5.1) we deduce that the eigenvalues $z_n = \sqrt{\mu_n}$ have asymptotics

$$z_{n} = \pi n + r_{n}, \qquad r_{n} = \sum_{1 \leq j \leq d} a_{j} \varepsilon^{2j+1} + \varepsilon^{k+1} \widetilde{r}_{n}, \qquad (\widetilde{r}_{n})_{1}^{\infty} \in \ell^{2},$$

$$z_{n} = \pi n \mathbf{v}_{n}, \qquad \mathbf{v}_{n} = 1 + 2r_{n} \varepsilon = 1 + \sum_{1 \leq j \leq d} 2a_{j} \varepsilon^{2j+2} + \ell_{k+2}^{2}(n),$$

$$(7.15)$$

where $(a_j)_1^d \in \mathbb{R}^d$. Moreover, the function $\mathbf{v}_n = 1 + 2r_n \varepsilon$ and $1/(i2z_n)$ satisfy

$$\frac{1}{\mathbf{v}_{n}^{s}} = \frac{1}{(1+2r_{n}\varepsilon)^{s}} = 1 + \sum_{1 \leq j \leq d} c_{j}(s)\varepsilon^{2j+2} + \ell_{k+2}^{2}(n),$$

$$\frac{1}{(i2z_{n})^{s}} = \frac{(-i)^{s}\varepsilon^{s}}{\mathbf{v}_{n}^{s}} = (-i)^{s} \sum_{1 \leq j \leq d} C_{j}(s)\varepsilon^{2j+2+s} + \ell_{k+1+s}^{2}(n), \quad \forall s \in \mathbb{N},$$
(7.16)

where $C_j(s), c_j(s)$ are polynomial of a_j . The asymptotics of $(i2z_n)^{-s}$ has two terms. The first is $\varepsilon^s \Psi$, where Ψ is a polynomial in ε with even power $2j + 2 \le d + 2$. The second term is a remainder $\ell_{k+1+s}^2(n)$. It is a crucial fact in our proof. Using (7.12), (7.15),(7.16) we obtain

$$w_{\cdot}(z_{n}) = 1 + \sum_{1 \leq j \leq d} \widetilde{w}_{j} \varepsilon^{2j} + \ell_{k+3}^{2}(n), \qquad y_{\cdot}(z_{n}) = \frac{1}{2} + \sum_{1 \leq j \leq 2k} \widetilde{y}_{j} \varepsilon^{j} + \ell_{k+3}^{2}(n),$$

$$\frac{1}{w_{\cdot}(z_{n})} = 1 + \sum_{1 \leq j \leq d} \widehat{w}_{j} \varepsilon^{2j} + \ell_{k+3}^{2}(n),$$
(7.17)

for some coefficients \widetilde{w}_j , \widetilde{y}_j , \widehat{w}_j and thus (7.8)-(7.10) imply

$$Y(1, z_n) = 1 + \sum_{1}^{k} \frac{u_j(1)}{(2iz_n)^j} + \frac{\widetilde{u}_k(1, z)}{(2iz_n)^{k+1}} = 1 + \sum_{1}^{k+2} \mathfrak{p}_j \varepsilon^j + \varepsilon^{k+1} q_k^{\bullet}(n) + \ell_{k+2}^2(n), \tag{7.18}$$

for some coefficients \mathfrak{p}_i depending on q.

From the Wronskian we get $\vartheta(1, \mu_n)\varphi'(1, \mu_n) = 1$. Then in order to get (7.13) we can study the asymptotics of $e^{h_{s,n}} = (-1)^n \vartheta(1, \mu_n)$ given by (7.11). From (7.11), (7.8), (7.18) we have

$$e^{h_{s,n}} = \frac{y'_0(z)e^{-iz}Y_1(-z) - y'_0(-z)e^{iz}Y_1(z)}{(-1)^n w(z)} \Big|_{z_n} = \frac{y_{\cdot}(z)e^{-ir_n}Y_1(-z) + y_{\cdot}(-z)e^{ir_n}Y_1(z)}{w_{\cdot}(z)} \Big|_{z_n}$$

$$= \sum_{m=0}^{2k} \frac{i^m}{m!} A_m(z_n) + O(r_n^{2k+1}) = \sum_{m=0}^{2k} \frac{i^m}{m!} A_m(z_n) + O(\varepsilon^{6k+3}),$$
(7.19)

since due to (7.15) we have $r_n = O(\varepsilon^3)$ and where

$$A_m(z_n) = \frac{r_n^m}{w_{\cdot}(z_n)} B_m(z_n), \qquad B_m(z) = (-1)^m y_{\cdot}(z) Y_1(-z) + y_{\cdot}(-z) Y_1(z). \tag{7.20}$$

Consider the main term $A_0(z)$. Since $B_0(z)$ is even in z, then (7.17), (7.18) give

$$B_{0}(z_{n}) = y_{\cdot}(z_{n})Y_{1}(-z_{n}) + y_{\cdot}(-z_{n})Y_{1}(z_{n}) = 1 + \sum_{1 \leq j \leq d} b_{j}\varepsilon^{2j} + E_{n}\varepsilon^{k+1} + \ell_{k+2}^{2}(n),$$

$$A_{0}(z_{n}) = \frac{B_{0}(z_{n})}{w_{\cdot}(z_{n})} = 1 + \sum_{1 \leq j \leq d} \widetilde{b}_{j}\varepsilon^{2j} + E_{n}\varepsilon^{k+1} + \ell_{k+2}^{2}(n),$$

$$(7.21)$$

where b_j, \widetilde{b}_j polynomials from $a_j, u_j'(0), u_j(1), j \in \mathbb{N}_k$ and $E_n = -\frac{q_k^{\bullet}(n) + (-1)^{k+1} q_k^{\bullet}(-n)}{i^{k+1} 2}$ satisfies

if
$$\frac{k}{2} \in \mathbb{N} \implies E_n = \frac{\int_0^1 q^{(k)}(t)(e^{i2\pi nt} - e^{-i2\pi nt})dt}{(-1)^{1+\frac{k}{2}}2i} = (-1)^{1+\frac{k}{2}} \int_0^1 q^{(k)}(t)\sin[2\pi nt]dt,$$
 (7.22)

which yields (7.14) for even k. Similar arguments imply (7.14) for odd k.

Consider $A_m(z)$ with even $m \ge 2$. Then the asymptotics (7.21), (7.15) imply

$$A_m(z_n) = r^m A_0(z) = r^m \left(1 + \sum_{1 \le j \le d} \widetilde{b}_j \varepsilon^{2j} + \ell_{k+1}^2(n) \right) = \sum_{3 \le j \le 1 + d + 3m} b_j' \varepsilon^{2j} + \ell_{k+1+3m}^2(n). \tag{7.23}$$

Thus $A_m(z_n)$ is an even polynomial of ε with order $\leq 2(1+d+3m)$ plus a reminder $\ell_{k+1+3m}^2(n)$. Let m be odd. Then $B_m(z) = y_*(-z)Y_1(z) - y_*(z)Y_1(-z)$ is odd in z. Then (7.18), (7.17) give

$$B_{1}(z_{n}) = y_{\cdot}(-z)Y_{1}(z_{n}) - y_{\cdot}(z_{n})Y_{1}(-z_{n}) = \sum_{1 \leqslant 2j+1 \leqslant d} t_{j} \varepsilon^{2j+1} + \ell_{k+1}^{2}(n),$$

$$\frac{B_{1}(z_{n})}{w_{\cdot}(z_{n})} = \frac{1}{w_{\cdot}(z_{n})} \left(\sum_{1 \leqslant 2j+1 \leqslant d} t_{j} \varepsilon^{2j+1} + \ell_{k+1}^{2}(n) \right) = \sum_{1 \leqslant 2j+1 \leqslant d} \widetilde{t}_{j} \varepsilon^{2j+1} + \ell_{k+1}^{2}(n),$$

where the constants t_j, \widetilde{t}_j are polynomials from $a_j, u'_j(0), u_j(1), j \in \mathbb{N}_k$, and using (7.15) we obtain

$$A_m(z_n) = r_n^m \frac{B_1(z)}{w(z)} = r_n^m \left(\sum_{1 \le i \le d} \widetilde{t}_i \varepsilon^{2j+1} + \ell_{k+1}^2(n) \right) = \sum_{3 \le i \le d+3m} t_i' \varepsilon^{2j} + \ell_{k+2}^2(n), \tag{7.24}$$

for some constants t'_{j} . Collecting estimates (7.21), (7.23), (7.24) we obtain

$$e^{h_{s,n}} = 1 + P(\varepsilon) + \varepsilon^{k+1} E_n + \ell_{k+2}^2(n),$$

where $P(\varepsilon)$ is a polynomial in ε with even power $2j+2 \le d+2$, which yields (7.13).

Lemma 7.3. Let $q \in \mathcal{L}_k, k \in \mathbb{N}$. Then the norming constants $\mathfrak{t}_n = \ln |\varphi(1, \tau_n) \sqrt{\tau_n^o}|$ satisfy

$$\mathfrak{t}_n = \sum_{1 \leq j \leq d} \mathfrak{t}_{j,n} \delta^{2j} + E_n \varepsilon^{k+1} + \ell_{k+2}^2(n) \quad \text{as} \quad n \to \infty, \tag{7.25}$$

where $\delta = \frac{1}{2\pi(n-\frac{1}{2})}$, the coefficients $\mathfrak{t}_{j,n} \in \mathbb{R}$, $j \in \mathbb{N}_d$, $d = \left[\frac{k+1}{2}\right]$ depend on q and

$$E_n = (-1)^{1+\left[\frac{k}{2}\right]} \int_0^1 q^{(k)}(x) \mathcal{G}_k(n'x) dx, \quad \mathcal{G}_k(nx) = \begin{cases} \sin[2\pi nx], & k \in 2\mathbb{N} \\ \cos[2\pi nx], & k+1 \in 2\mathbb{N} \end{cases}$$
 (7.26)

Proof. From (5.3) we deduce that the eigenvalues $z_n = \sqrt{\tau_n}$ have asymptotics

$$z_{n} = \pi n' + r_{n}, \quad n' = n - \frac{1}{2}, \quad r_{n} = \sum_{1 \le j \le d} a_{j} \delta^{2j+1} + \delta^{k+1} \widetilde{r}_{n}, \qquad (\widetilde{r}_{n})_{1}^{\infty} \in \ell^{2},$$

$$z_{n} = \pi n' \mathbf{v}_{n}, \qquad \mathbf{v}_{n} = 1 + 2r_{n} \delta = 1 + \sum_{1 \le i \le d} 2a_{j} \delta^{2j+2} + \ell_{k+2}^{2}(n),$$

$$(7.27)$$

where $(a_j)_1^d \in \mathbb{R}^d$. Moreover, the function $\mathbf{v}_n = 1 + 2r_n\delta$ and $\varsigma = i2z$ satisfies

$$\frac{1}{\mathbf{v}_{n}^{s}} = \frac{1}{(1+2r_{n}\delta)^{s}} = 1 + \sum_{1 \leq j \leq d} c_{j}(s)\delta^{2j+2} + \ell_{k+2}^{2}(n),$$

$$\frac{1}{(i2z_{n})^{s}} = \frac{(-i)^{s}\delta^{s}}{\mathbf{v}_{n}^{s}} = (-i)^{s} \sum_{1 \leq j \leq d} C_{j}(s)\delta^{2j+2+s} + \ell_{k+1+s}^{2}(n), \quad \forall \ s \in \mathbb{N},$$
(7.28)

where $C_j(s), c_j(s)$ are polynomial of a_j . The asymptotics of z_n^{-s} has two terms. The first is $\varepsilon^s \mathcal{P}$, where \mathcal{P} a polynomial in ε with even power $2j+2 \leqslant d+2$. The second term is a remainder $\ell_{k+1+s}^2(n)$. It is a crucial fact in our proof.

Recall that due to (7.8) a solution to $-y'' + qy = z^2y, z > 1$ has the form

$$y(x,z) = e^{izx}Y(x,z), \quad Y(x,z) = 1 + \sum_{j=1}^{k} \frac{u_j(x)}{\varsigma^j} + \frac{\tilde{u}_k(x,z)}{\varsigma^{k+1}}, \quad \varsigma = i2z.$$
 (7.29)

We need also the Wronskian $w(z) = \{y(\cdot, -z), y(\cdot, z)\} = y_0'(z) - y_0'(-z)$, where the function $y_0'(z) = y'(0, z)$. The function φ has the form

$$\varphi(1,z) = \frac{1}{w(z)}(y_1(z) - y_1(-z)), \quad \text{where } y_1(z) = y(1,z).$$
 (7.30)

Using (7.12), (7.15), (7.16) we obtain

$$w_{\cdot}(z) = 1 + \sum_{1 \leq j \leq d} \widetilde{w}_{j} \delta^{2j} + \ell_{k+3}^{2}(n), \qquad y_{\cdot}(z) = \frac{1}{2} + \sum_{1 \leq j \leq 2k} \widetilde{y}_{j} \delta^{j} + \ell_{k+3}^{2}(n),$$

$$\frac{1}{w_{\cdot}(z)} = 1 + \sum_{1 \leq j \leq d} \widehat{w}_{j} \delta^{2j} + \ell_{k+3}^{2}(n),$$
(7.31)

for some coefficients \widetilde{w}_j , \widetilde{y}_j , \widehat{w}_j and (7.8)-(7.10) imply

$$Y(1, z_n) = 1 + \sum_{1}^{k} \frac{u_j(1)}{(2iz_n)^j} + \frac{\widetilde{u}_k(1, z)}{(2iz_n)^{k+1}} = 1 + \sum_{1}^{k+1} \mathfrak{p}_j \delta^j + \delta^{k+1} q_k^{\bullet}(n) + \ell_{k+2}^2(n),$$

$$q_*^k(z) = (-1)^{k+1} \int_0^1 e^{i2\pi n'x} q^{(k)}(x) dx$$
(7.32)

for some coefficients \mathfrak{p}_j depending on q. From (7.15), (7.30), (7.32) we have that the norming constant $e^{\mathfrak{t}_n} = (-1)^n (\pi n') \varphi(1, z_n)$ satisfies

$$e^{\mathfrak{t}_{n}} = (\pi n') \frac{e^{-iz_{n}} Y_{1}(-z_{n}) - e^{iz_{n}} Y_{1}(z_{n})}{(-1)^{n} (2iz_{n}) w_{.}(z_{n})} = \frac{e^{-ir_{n}} Y_{1}(-z_{n}) + e^{ir_{n}} Y_{1}(z_{n})}{2 \mathbf{v}_{n} w_{.}(z_{n})}$$

$$= \sum_{m=0}^{2k} \frac{i^{m}}{m!} A_{m}(z_{n}) + O(r_{n}^{2k+1}) = \sum_{m=0}^{2k} \frac{i^{m}}{m!} A_{m}(z_{n}) + O(\delta^{6k+3}),$$
(7.33)

since due to (7.27) we have $r_n = O(\delta^3)$ and where

$$A_m(z_n) = \frac{r^m}{\mathbf{v}_n w_{\cdot}(z_n)} B_m(z_n), \qquad B_m(z_n) = \frac{(-1)^m Y_1(-z_n) + Y_1(z_n)}{2}. \tag{7.34}$$

Consider the main term $A_0(z_n)$. Since $B_0(z)$ is even in z, then (7.17), (7.32) give

$$B_{0}(z_{n}) = \frac{Y_{1}(-z_{n}) + Y_{1}(z_{n})}{2} = 1 + \sum_{1 \leq j \leq d} b_{j} \delta^{2j} + E_{n} \delta^{k+1} + \ell_{k+2}^{2}(n),$$

$$A_{0}(z_{n}) = \frac{B_{0}(z)}{\mathbf{v}_{n} w_{\cdot}(z)} = 1 + \sum_{1 \leq j \leq d} \widetilde{b}_{j} \delta^{2j} + E_{n} \delta^{k+1} + \ell_{k+2}^{2}(n),$$

$$(7.35)$$

where b_j, \widetilde{b}_j polynomials from $a_j, u_j'(0), u_j(1), j \in \mathbb{N}_k$ and $E_n = -\frac{q_k^{\bullet}(n) + (-1)^{k+1} q_k^{\bullet}(-n)}{i^{k+1}2}$ satisfies

if
$$\frac{k}{2} \in \mathbb{N} \implies E_n = \frac{\int_0^1 (e^{i2\pi nt} - e^{-i2\pi nt})q^{(k)}(t)dt}{(-1)^{1+\frac{k}{2}}i2} = (-1)^{1+\frac{k}{2}} \int_0^1 \sin[2\pi nt]q^{(k)}(t)dt,$$
 (7.36)

which yields (7.14) for even k. Similar arguments imply (7.14) for odd k.

Consider $A_m(z)$ with even $m \ge 2$. Then the asymptotics (7.21), (7.15) imply

$$A_m(z_n) = r^m A_0(z) = r^m \left(1 + \sum_{1 \le j \le d} \widetilde{b}_j \delta^{2j} + \ell_{k+1}^2(n) \right) = \sum_{3 \le j \le 1 + d + 3m} b_j' \delta^{2j} + \ell_{k+1+3m}^2(n). \tag{7.37}$$

Thus $A_m(z_n)$ is an even polynomial of ε with order $\leq 2(1+d+3m)$ plus a reminder $\ell_{k+1+3m}^2(n)$. Let $m \in \mathbb{N}$ be odd. Then $B_1(z) = y_{\cdot}(-z)Y_1(z) - y_{\cdot}(z)Y_1(-z)$ is odd in z and (7.28), (7.32), (7.31) give

$$B_{1}(z_{n}) = y.(-z)Y_{1}(z_{n}) - y.(z_{n})Y_{1}(-z_{n}) = \sum_{1 \leq 2j+1 \leq d} t_{j}\delta^{2j+1} + \ell_{k+1}^{2}(n),$$

$$\frac{B_{1}(z_{n})}{\mathbf{v}_{n}w.(z_{n})} = \frac{1}{\mathbf{v}_{n}w.(z_{n})} \left(\sum_{1 \leq 2j+1 \leq d} t_{j}\delta^{2j+1} + \ell_{k+1}^{2}(n) \right) = \sum_{1 \leq 2j+1 \leq d} \widetilde{t}_{j}\delta^{2j+1} + \ell_{k+1}^{2}(n),$$

where the constants t_j, \widetilde{t}_j are polynomials from $a_j, u'_j(0), u_j(1), j \in \mathbb{N}_k$, and using (7.27) we obtain

$$A_m(z_n) = r_n^m \frac{B_1(z_n)}{\mathbf{v}_n w_{\cdot}(z)} = r_n^m \left(\sum_{1 \le i \le d} \widetilde{t}_j \delta^{2j+1} + \ell_{k+1}^2(n) \right) = \sum_{3 \le i \le d+3m} t_j' \delta^{2j} + \ell_{k+2}^2(n), \tag{7.38}$$

for some constants t'_{i} . Collecting estimates (7.35), (7.37), (7.38) we obtain

$$e^{h_{s,n}} = 1 + P(\varepsilon) + \varepsilon^{k+1} E_n + \ell_{k+2}^2(n),$$

where $P(\varepsilon)$ is a polynomial in ε^2 with even power $2j + 2 \leq d + 2$, which yields (7.25).

- 7.2. Trace formulas. We discuss trace formulas for Sturm-Liouvill problem for $q \in \mathcal{L}_1$.
- Gel'fand and Levitan determined trace formulas [10] for Dirichlet and Neumann eigenvalues

$$\frac{q(0)+q(1)}{4} = \sum_{n\geqslant 1} (\mu_n^o - \mu_n),
\frac{q(0)+q(1)}{4} = \sum_{n\geqslant 0} (\nu_n - \nu_n^o),$$
(7.39)

and mixed eigenvalues

$$\frac{q(0)-q(1)}{4} = \sum_{n\geqslant 1} (\tau_n^o - \tau_n), \tag{7.40}$$

$$\frac{q(0)-q(1)}{4} = \sum_{n\geqslant 1} (\varrho_n - \tau_n^o). \tag{7.41}$$

• Magnus and Winkler [30] determined trace formulas for periodic case:

$$0 = \sum_{n \in \mathbb{N}} (\lambda_{2n-1}^+ + \lambda_{2n-1}^- - 2\mu_{2n-1}^o),$$

$$\lambda_0^+ = -\sum_{n \geqslant 1} (\lambda_{2n}^+ + \lambda_{2n}^- - 2\mu_{2n}^o).$$
(7.42)

• Another trace formula for periodic case was obtained (see e.g., [14], [37])

$$q(0) = \lambda_0^+ + \sum_{n \ge 1} (\lambda_n^+ + \lambda_n^- - 2\mu_n), \quad q \in \mathcal{H}_1.$$
 (7.43)

In fact summing (7.39) for Dirichlet eigenvalues and (7.42) we obtain (7.43). Similar arguments imply the trace formula for periodic case via the Neumann eigenvalues in (7.39):

$$q(0) = 2\nu_0 - \lambda_0^+ - \sum_{n \ge 1} (\lambda_n^+ + \lambda_n^- - 2\nu_n). \tag{7.44}$$

We recall the known facts from the theory of Fourier series.

Lemma 7.4. Let $q = q_{od} + q_{ev}$ for some $q \in \mathcal{L}_1$, where

$$q_{od}(x) = \sum_{n \ge 1} 2q_{c,2n-1} \cos \pi (2n-1)x, \quad q_{ev}(x) = \sum_{n \ge 1} 2q_{c,2n} \cos \pi 2nx, \quad x \in (0,1), \quad (7.45)$$

and $q_{c,n} = \int_0^1 q(x) \cos \pi nx dx$. Then the following formulas hold true:

$$q_{ev}(0) = \frac{q(0)+q(1)}{2}, \qquad q_{od}(0) = \frac{q(0)-q(1)}{2}.$$
 (7.46)

Proof. There is a Fourier series $q(x) = \sum_{n \geq 1} 2q_{c,n} \cos \pi nx$, where due to $q \in \mathcal{L}_1$ a sequence $(q_{c,n})_1^{\infty} \in \ell_1^2(\mathbb{N})$. Then we obtain $q(0) = q_{od}(0) + q_{ev}(0)$, since here all functions are continuous on [0,1]. The first identity in (7.46) is well known. Thus these two facts imply (7.46).

We discuss trace formulas associated with the mappings $\tau \star \mu$ and \mathfrak{f} , defined by (1.11). Here also we show (7.40), (7.41), since we can not find a reference for them.

Proposition 7.5. Let $q \in \mathcal{L}_1$. Then the trace formulas (7.40), (7.41) and

$$q(0) = \nu_0 + \sum_{n \ge 1} \left((\nu_n - \mu_n) + (\varrho_n - \tau_n) \right),$$

$$q(1) = \nu_0 + \sum_{n \ge 1} \left((\nu_n - \mu_n) - (\varrho_n - \tau_n) \right),$$
(7.47)

and

$$q(0) = 2\sum_{n\geqslant 1} \left((\mu_n^o - \mu_n) + (\tau_n^o - \tau_n) \right) = 2\nu_0 + 2\sum_{n\geqslant 1} \left((\nu_n - \nu_n^o) + (\varrho_n - \varrho_n^o) \right),$$

$$q(1) = 2\sum_{n\geqslant 1} \left((\mu_n^o - \mu_n) - (\tau_n^o - \tau_n) \right) = 2\nu_0 + 2\sum_{n\geqslant 1} \left((\nu_n - \nu_n^o) - (\varrho_n - \varrho_n^o) \right).$$
(7.48)

hold true, where all series converge absolutely.

Proof. Firstly we discuss the trace formulas for mixed eigenvalues τ_n , since I can find their proofs. We show (7.40) shortly repeating well-known arguments [10], [7]. The standard arguments from the papers of Gel'fand and Levitan [10] and Dikiy [7] (see also a book [29]) imply the identity

$$\sum_{n\geq 1} \left((\tau_n^o - \tau_n) + 2 \int_0^1 q(x) \sin^2 k_n x dx \right) = 0, \qquad k_n = \sqrt{\tau_n^o} > 0, \tag{7.49}$$

where $\sqrt{2}\sin k_n x$ is the unperturbed eigenfunction corresponding to the unperturbed eigenvalue τ_n^o . The identity (7.46) gives

$$-\sum_{n\geq 1} \int_0^1 q(x) 2\sin^2 k_n x dx = \sum_{n\geq 1} \int_0^1 q\cos 2k_n x dx = \frac{q(0) - q(1)}{4},$$

which jointly with (7.49) yields (7.40). The proof of (7.41) for ϱ_n is similar.

Summing trace formulas in (7.39), (7.40), (7.41) we have the first identity in (7.47).

Summing trace formulas in (7.39), summing trace formulas in (7.40), (7.41) and take their deferens we obtain the second identity in (7.47).

Summing the first trace formula in (7.39) and (7.40) and take the deferens we have the first trace formula in (7.48).

Summing the second trace formula in (7.39) and (7.41) and take the deferens we have the second trace formula in (7.48).

• Summing trace formulas in (7.40), (7.41) and take the deferens we have

$$\sum_{n \ge 1} (\varrho_n - \tau_n) = \frac{q(0) - q(1)}{2}, \qquad \sum_{n \ge 1} (\varrho_n + \tau_n - 2\tau_n^o) = 0.$$
 (7.50)

Similar arguments and identities (7.47), (7.50) imply

$$\nu_0 + \sum_{n\geqslant 1} (\nu_n - \mu_n) = \frac{q(0)+q(1)}{2}, \qquad \nu_0 + \sum_{n\geqslant 1} (\nu_n + \mu_n - 2\mu_n^o) = 0.$$
 (7.51)

Acknowledgments. I thank Segei Kuksin (Paris) for usefull discusion about inverse problems.

REFERENCES

- [1] Chelkak, D.; Kargaev, P.; Korotyaev, E. Inverse problem for harmonic oscillator perturbed by potential, characterization. Comm. Math. Phys. 249 (2004), no. 1, 133–196.
- [2] Chelkak, D.; Korotyaev, E. Weyl-Titchmarsh functions of vector-valued Sturm-Liouville operators on the unit interval. Journal of Functional Analysis, 257 (2009), 1546–1588.
- [3] Clay, M.; Margalit, D. Office Hours with a Geometric Group Theorist. Princeton Univ. Press, 2017.
- [4] Coleman C.; McLaughlin, J. Solution of the inverse problem for an impedance with integrable derivative, I Commun. Pure Appl. Math. 46 (1993), 145–184.
- [5] Coleman C.; McLaughlin, J. Solution of the inverse problem for an impedance with integrable derivative II, Commun. Pure Appl. Math. 46(1993), 185–212.
- [6] Dahlberg, B.; Trubowitz, E. The inverse Sturm-Liouville problem. III. Comm. Pure Appl. Math. 37(1984), no. 2, 255–267.
- [7] Dikii, L. A. On a formula of Gel'fand-Levitan. (Russian) Uspehi Matem. Nauk (N.S.) 8, (1953). no. 2(54), 119–123.
- [8] Garnett, J., Trubowitz, E. Gaps and bands of one dimensional periodic Schrödinger operators II. Comment. Math. Helv. 62(1987), 18–37.
- [9] Gel'fand, I. M.; Levitan, B. M. On the determination of a differential equation from its spectral function. (Russian) Izvestiya Akad. Nauk SSSR. Ser. Mat. 15, (1951). 309–360. English Translation: Amer. Math. Soc. Transl. (2) 1 (1955), 253–304.

- [10] Gel'fand, I. M.; Levitan, B. M. On a simple identity for the characteristic values of a differential operator of the second order. (Russian) Doklady Akad. Nauk SSSR (N.S.) 88, (1953). 593–596.
- [11] Guillot J.G.; Ralston J.V. Inverse spectral theory for a singular Sturm-Liouville operator on [0, 1], J. Diff. Eq., 76 (1988), 353–373.
- [12] Isaacson, E. L.; Trubowitz, E. The inverse Sturm-Liouville problem. I. Comm. Pure Appl. Math. 36 (1983), no. 6, 767–783.
- [13] Isaacson, E. L.; McKean, H. P.; Trubowitz, E. The inverse Sturm-Liouville problem. II. Comm. Pure Appl. Math. 37 (1984), no. 1, 1–11.
- [14] Its, A. R.; Matveev, V. B. Schrödinger operators with the finite-band spectrum and the N-soliton solutions of the Korteweg-de Vries equation. (Russian) Teoret. Mat. Fiz. 23 (1975), no. 1, 51–68.
- [15] Kappeler, T.; Pöschel, J. KdV & KAM. Springer, 2003.
- [16] Kargaev, P.; Korotyaev E. The inverse problem for the Hill operator, direct approach. Invent. Math. 129(1997), 567–593.
- [17] Klein, M.; Korotyaev, E. Parametrization of periodic weighted operators in terms of gap lengths. Inverse Problems 16 (2000), no. 6, 1839–1860.
- [18] Korotyaev, E. Inverse Sturm-Liouville problems for non-Borg conditions, Journal of Inverse and Ill-Posed Problems, 27(2019), no 3, 445–452.
- [19] Korotyaev, E. Estimates for the Hill operator. II. J. Differential Equations 223 (2006), no. 2, 229–260.
- [20] Korotyaev, E. Estimates for the Hill operator. I. J. Differential Equations 162 (2000), no. 1, 1–26.
- [21] Korotyaev, E Inverse problem for periodic "weighted" operators. J. Funct. Anal. 170 (2000), no. 1, 188–218.
- [22] Korotyaev, E. The inverse problem and trace formula for the Hill operator, II. Math. Z. 231(1999), 345–368.
- [23] Korotyaev, E. Estimates of periodic potentials in terms of gap lengths. Comm. Math. Phys. 197 (1998), no. 3, 521–526.
- [24] Korotyaev, E. The inverse problem for the Hill operator. I. Int. Math. Res. Notices 1997, no. 3, 113–125.
- [25] Korotyaev, E. The estimates of periodic potentials in terms of effective masses. Comm. Math. Phys. 183 (1997), no. 2, 383–400.
- [26] Korotyaev, E.; Chelkak, D. The inverse Sturm-Liouville problem with mixed boundary conditions, St. Petersburg Math. Journal. 21(2009), no 5, 114–137.
- [27] Krein, M. G. Solution of the inverse Sturm-Liouville problem. (Russian) Doklady Akad. Nauk SSSR (N.S.) 76, (1951). 21–24.
- [28] Krein, M. G. On a method of effective solution of an inverse boundary problem. (Russian) Doklady Akad. Nauk SSSR (N.S.) 94, (1954). 987–990.
- [29] Levitan, B. Inverse Sturm-Liouville problems. Utrecht: VNU Science Press, 1987.
- [30] W. Magnus, S. Winkler. Hill's equation, Interscience Tracts in Pure and Applied Mathematics, No. 20 Interscience Publishers John Wiley & Sons, New York-London-Sydney, 1966.
- [31] Marchenko, V. A. Concerning the theory of a differential operator of the second order. (Russian) Doklady Akad. Nauk SSSR. (N.S.) 72, (1950). 457–460.
- [32] Marchenko, V.; Ostrovski, I. A characterization of the spectrum of the Hill operator. Mat. Sb. 97(139), (1975), 540–606.
- [33] Marchenko, V. Sturm-Liouville operator and applications. Basel, Birkhäuser 1986.
- [34] Pöschel, P., Trubowitz E. Inverse Spectral Theory. Boston, Academic Press, 1987.
- [35] Savchuk, A. M.; Shkalikov, A. A. On the properties of mappings associated with inverse Sturm-Liouville problems. Proc. Steklov Inst. Math. 260 (2008), no. 1, 218–237.
- [36] Savchuk, A. M.; Shkalikov, A. A. Inverse Problems for Sturm-Liouville Operators with Potentials in Sobolev Spaces: Uniform Stability, Funct. Anal. Appl., 44:4 (2010), 270–285.
- [37] Trubowitz, E. The inverse problem for periodic potentials. Comm. Pure Appl. Math. 30 (1977), no. 3, 321–337.

Department of Math. Analysis, Saint-Petersburg State University, Universitetskaya nab. 7/9, St. Petersburg, 199034, Russia, and HSE University, 3A Kantemirovskaya ulitsa, St. Petersburg, 194100, Russia korotyaev@gmail.com, e.korotyaev@spbu.ru