

A CHARACTERIZATION OF ZONOIDS

YOSSI LONKE

1. INTRODUCTION

A function defined on the n -dimensional sphere \mathbb{S}^{n-1} will be called *zonal* if its value at a point $x \in \mathbb{S}^{n-1}$ depends only on the angle between x and a fixed axis. Thus if u is a unit vector in the direction of the fixed axis, then a function is zonal with respect to u if its value at x depends only on the standard inner product $\langle x, u \rangle$. A natural way to generate zonal functions is as follows. Let $f \in C(\mathbb{S}^{n-1})$, and fix a direction $u \in \mathbb{S}^{n-1}$. Let $(O(u), m)$ denote the subgroup of orthogonal transformations which keep the point u fixed, equipped with the normalized Haar measure, m . Define

$$(S_u f)(x) = \int_{O(u)} f(Tx) dm(T)$$

Clearly, the function $S_u f$ is zonal. Applying this procedure to support functions leads to the following definition.

Definition 1.1. Suppose $u \in \mathbb{S}^{n-1}$ and K is a centrally symmetric convex body, with support function h_K . The u -spin of K is the convex body K_u whose support function is $S_u h_K$.

For example, computing the e_n -spin of the unit cube in \mathbb{R}^n , gives:

$$(S_{e_n} h_{B_n^\infty})(x) = \int_{O(e_n)} \|Tx\|_1 dm(T) = c_n \left(\sum_{i=1}^n x_i^2 \right)^{1/2} + |x_n|,$$

(c_n is a positive number depending on n), which is the support function of a cylinder. This illustrates the choice of the word 'spin'.

Thus, with each body there is associated a system of rotation-bodies, one for each possible direction. Since $S_u f(u) = f(u)$ for each $u \in \mathbb{S}^{n-1}$ and every function f , a body is seen to be uniquely defined by its spins. The main result of this note is the following.

Theorem 1.2. *A centrally symmetric convex body is a zonoid if and only if all its spins are zonoids.*

Date: February 2020.

2. PRELIMINARIES

$C^\infty(\mathbb{S}^{n-1})$ is the Frechet space of functions on \mathbb{S}^{n-1} that have derivatives of every order. Elements of this space are called *test functions*. Its dual space, denoted $\mathcal{D}(\mathbb{S}^{n-1})$, is the space of distributions on \mathbb{S}^{n-1} . If a subscript e is added to any of the above spaces, it is to designate the subspace of even objects.

The *cosine transform* is the operator $\mathcal{C} : C_e^\infty(\mathbb{S}^{n-1}) \rightarrow C_e^\infty(\mathbb{S}^{n-1})$ defined by:

$$(\mathcal{C}f)(x) = \int_{\mathbb{S}^{n-1}} |\langle x, y \rangle| f(y) d\sigma_{n-1}(y),$$

where σ_{n-1} is the normalized rotation-invariant measure on the sphere. It is well known that \mathcal{C} is a continuous bijection of $C_e^\infty(\mathbb{S}^{n-1})$ onto itself, and that it can be extended by duality to a bi-continuous bijection of the dual space $\mathcal{D}_e(\mathbb{S}^{n-1})$. Hence, if ρ is a distribution and $f \in C_e^\infty(\mathbb{S}^{n-1})$ is a test function, then

$$\langle \mathcal{C}\rho, f \rangle = \langle \rho, \mathcal{C}f \rangle$$

Since $C_e^\infty(\mathbb{S}^{n-1})$ and its dual space, the even measures, are both naturally embedded in $\mathcal{D}_e(\mathbb{S}^{n-1})$, it makes sense to speak about the cosine transform of a measure, or of a continuous function. A fundamental connection between distributions and centrally symmetric convex bodies was discovered by Weil, in [1]. Weil proved that for every centrally symmetric convex body $K \subset \mathbb{R}^n$ there corresponds a unique distribution ρ_K (called the *generating distribution* of K) such that $\mathcal{C}\rho_K = h_K$, where h_K is the support function of K . Suppose $f \in C^\infty(\mathbb{S}^{n-1})$. Then for every direction $u \in \mathbb{S}^{n-1}$, the function $S_u f$ also belongs to $C^\infty(\mathbb{S}^{n-1})$. Therefore, S_u can be defined to act on distributions by duality:

$$\langle S_u \rho, f \rangle = \langle \rho, S_u f \rangle, \quad \rho \in \mathcal{D}(\mathbb{S}^{n-1}), \quad f \in C^\infty(\mathbb{S}^{n-1})$$

A routine verification shows that the transforms S_u and \mathcal{C} commute on test functions, and therefore also as transforms of distributions.

3. PROOF OF THEOREM 1.1

If K is a zonoid, then $h_K = \mathcal{C}\mu$ for some positive measure μ , and for every $u \in \mathbb{S}^{n-1}$,

$$S_u h_K = S_u(\mathcal{C}(\mu)) = \mathcal{C}(S_u \mu)$$

Since $S_u \mu$ is a positive measure for every μ , every spin of K is a zonoid. This proves the easy part of the theorem.

Suppose K is a centrally symmetric convex body every spin of which is a zonoid. That is, for each direction $u \in \mathbb{S}^{n-1}$ there exists a positive measure μ_u such that $S_u h_K = \mathcal{C}\mu_u$. There exists a distribution ρ such that $h_K = \mathcal{C}\rho$, and so the assumption reduces to $\mathcal{C}(S_u \rho) = \mathcal{C}\mu_u$, where the commuting of S_u and \mathcal{C} was used. Since \mathcal{C} is one-to-one, the distribution ρ is seen to satisfy $S_u \rho = \mu_u$ for every $u \in \mathbb{S}^{n-1}$. It therefore remains to prove:

Lemma 3.1. *A distribution is positive if and only if $S_u \rho$ is positive for every $u \in \mathbb{S}^{n-1}$.*

Proof. The "only if" part is obvious. Suppose $\rho \in \mathcal{D}(\mathbb{S}^{n-1})$ is a distribution such that $S_u \rho$ is positive for every $u \in \mathbb{S}^{n-1}$. Then $\langle \rho, f \rangle \geq 0$ for every positive zonal test function f . Therefore if $g = \sum_1^m a_i f_i$, where $a_i \geq 0$ and f_i are positive zonal test functions, then also $\langle \rho, g \rangle \geq 0$.

Choose a positive test function $f \in C^\infty(\mathbb{S}^{n-1})$ and write its Poisson integral:

$$P_r f(x) = \int_{\mathbb{S}^{n-1}} \frac{1-r^2}{\|x-ry\|^n} f(y) d\sigma_{n-1}(y)$$

It is well known that $\lim_{r \rightarrow 1^-} P_r f = f$ in the topology of $C^\infty(\mathbb{S}^{n-1})$. Therefore, if $\langle \rho, P_r f \rangle \geq 0$ for every $0 < r < 1$ and every positive test function f , then ρ is positive, by continuity. Fix a sequence of convex combinations of Dirac measures of the form

$$(1) \quad \nu_N = \sum_{i \geq 1} \lambda_{iN} \delta_{y_i}, \quad y_i \in \mathbb{S}^{n-1}$$

such that $\nu_N \rightarrow \sigma_{n-1}$ in the w^* topology of measures. If φ is a test function, then

$$(2) \quad \int_{\mathbb{S}^{n-1}} \frac{1-r^2}{\|x-ry\|^n} \varphi(y) d\nu_N(y) \rightarrow \int_{\mathbb{S}^{n-1}} \frac{1-r^2}{\|x-ry\|^n} \varphi(y) d\sigma_{n-1}(y) = P_r \varphi(x)$$

If $F(x, y)$ is a continuous function on $\mathbb{S}^{n-1} \times \mathbb{S}^{n-1}$ then for every test function φ , the integrals $\langle F(x, y)\varphi(y), \nu_N \rangle$ converge to the integral $\langle F(x, y)\varphi(y), \sigma_{n-1} \rangle$ uniformly in x . Therefore, the sequence of functions of the variable x in the l.h.s of (2) converge to the r.h.s in the topology of $C^\infty(\mathbb{S}^{n-1})$, because the function $\|x-ry\|^{-n}$ for fixed y and $0 < r < 1$ is C^∞ with respect to x . Moreover, the distribution ρ is positive on every term of the l.h.s of (2), because for each y_i appearing in (1) one has

$$\|x-ry_i\|^{-n} = \int_{O(y_i)} \|Tx-ry_i\|^{-n} dm(T), \quad \forall x$$

so

$$\langle \rho, \|x-ry_i\|^{-n} \rangle = \langle \rho, S_{y_i}(\|x-ry_i\|^{-n}) \rangle = \langle S_{y_i} \rho, \|x-ry_i\|^{-n} \rangle \geq 0$$

Hence by continuity ρ is also positive of the r.h.s of (2) as well. It follows that ρ is a positive distribution. \square

It is well known that a positive distribution is in fact a positive measure, that is, it satisfies $|\rho(f)| \leq \rho(1)\|f\|_\infty$ for every test function, where 1 is the constant function 1. Consequently, there is a unique extension of ρ to a bounded linear functional on $C_c(\mathbb{S}^{n-1})$, and so in this sense it represents a measure. This completes the proof of the theorem.

REFERENCES

- [1] Weil, W, Centrally symmetric convex bodies and distributions, Israel J. Math., 24, 352–367 (1976)