

AUTOMORPHISMS OF GENERALIZED FERMAT MANIFOLDS

RUBÉN A. HIDALGO, HENRY F. HUGHES, AND MAXIMILIANO LEYTON-ÁLVAREZ

ABSTRACT. A group H of holomorphic automorphisms of a compact d -dimensional complex manifold M is called a generalized Fermat group of type $(d; k, n)$ if $H \cong \mathbb{Z}_k^n$ and there is a Galois holomorphic branched covering $\pi : M \rightarrow \mathbb{P}^d$, with deck group H , whose branch divisor consists of $n + 1$ hyperplanes in general position (each one of branch order k). We prove that generalized Fermat groups of type $(d; k, n)$ are unique for $(d; k, n) \notin \{(2; 2, 5), (2; 4, 3)\}$.

1. INTRODUCTION

In what follows, M will denote a compact d -dimensional complex manifold and $\text{Aut}(M)$ will be its group of holomorphic automorphisms. It is known that $\text{Aut}(M)$ is a complex Lie group (see page 77 of [19]) and, if $\text{Aut}^0(M)$ is its connected component of the identity, then $\text{Aut}(M)/\text{Aut}(M)^0$ is a countable discrete group.

If $H \leq \text{Aut}(M)$ is a finite abelian group, then the quotient $\mathcal{O}_{M,H} := M/H$ is a compact d -dimensional complex orbifold and there is a (branched) cover map $\pi : M \rightarrow \mathcal{O}_{M,H}$ with deck group H . The π -image of those points with non-trivial H -stabilizer is the branch locus of π . If the H -stabilizer of $x \in M$ is generated by involutions, each one with a co-dimension one locus of fixed points, then the branch value $\pi(x)$ is a smooth point of $\mathcal{O}_{M,H}$. If $\mathcal{O}_{M,H} = \mathbb{P}^d$, then Serre's cohomological criterion for ampleness, asserts that M is projective. In [23], Pardini constructed a building data which describes (up to isomorphisms) these abelian (branched) covers. In [9], Gao constructed cyclic equations of the manifold M .

We say that $H \leq \text{Aut}(M)$ is *unique* (or *rigid*) if for $G \leq \text{Aut}(M)$, $G \cong H$, such that $\mathcal{O}_{M,H}$ and $\mathcal{O}_{M,G}$ are topologically equivalent orbifolds (i.e., they are homeomorphic as topological spaces, but the homeomorphism respect the branching data), then $G = H$. In general, checking if H is unique is a difficult task.

If $d = 1$ (i.e., M a closed Riemann surface), then the uniqueness is known, for instance, in the following situations: (i) M is hyperelliptic and $H \cong \mathbb{Z}_2$ is generated by the hyperelliptic involution; (ii) $H \cong \mathbb{Z}_p$, where $p \geq 3$ is a prime integer such that $g > (p - 1)^2$ and M/H has genus zero [4]; (iii) $H \cong \mathbb{Z}_k^n$, where $k, n \geq 2$ with $(n - 1)(k - 1) > 2$, and M/H of genus zero with exactly $n + 1$ cone points of order k [15] (these Riemann surfaces are called generalized Fermat curves of type (k, n) and algebraic equations for them are provided as a fiber product of $n - 1$ classical Fermat curves of degree k).

If $d \geq 2$, following the case of generalized Fermat curves, we study the uniqueness for $H \cong \mathbb{Z}_k^n$, where $n, k \geq 2$, and M/H is the orbifold whose underlying space is the d -dimensional projective space \mathbb{P}^d and whose branch locus is the complete intersection of $n + 1$ hyperplanes in general position, each one with branch order k . In this case, we say that M (respectively, H and (M, H)) is a *generalized Fermat manifold* (respectively,

2010 *Mathematics Subject Classification.* 14J50; 32Q40; 53C15.

Partially supported by the projects Fondecyt 1230001, 1220261, and 1221535.

a *generalized Fermat group* and a *generalized Fermat pair* of type $(d; k, n)$. We note that necessarily $n \geq d$ (as M is assumed to be smooth). The case $n = d$ corresponds to $M = \mathbb{P}^d$. The main result of this paper is the following.

Theorem (A). *Let (M, H) be a generalized Fermat pair of type $(d; k, n)$, where $n \geq d+1$, $k, d \geq 2$. If $(d; k, n) \notin \{(2; 2, 5), (2; 4, 3)\}$, then H is the unique generalized Fermat group of type $(d; k, n)$ of M .*

Generalized Fermat manifolds (without that name) have been already considered in the literature. In [3], Bin studied some of these examples (case $k = 2$ and $n = 4$) and their corresponding Pardini's building data. For $d = 2$, in [16] (see also [17]), Hirzebruch studied the arrangement of $n+1$ lines in \mathbb{P}^2 (which are not necessarily in general position). Such an arrangement produces a singular surface X (with normal singularities) with an action of a \mathbb{Z}_k^n producing a branched cover over \mathbb{P}^2 with branching data being the given lines and each one with branch order k . If the lines are in general position, then X is a generalized Fermat manifold of type $(2; k, n)$ (if the lines are not in a general position, then one may produce a desingularization of X to obtain a generalized Fermat manifold of type $(2; k, n)$). In [16, Section (2.3)], it is remarked (without proof) that generalized Fermat manifolds of type $(2; k, n)$ are a complete intersection of $(n-2)$ Fermat surfaces of degree k in \mathbb{P}^n .

As a consequence of Pardini's building data, one may observe that if (M_1, H_1) and (M_2, H_2) are two generalized Fermat manifolds of type $(d; k, n)$, then there is an isomorphism between M_1 and M_2 that conjugates H_1 to H_2 if and only if the branch locus of M_1/H_1 is sent to the branch locus of M_2/H_2 by some $A \in \text{PGL}_{d+1}(\mathbb{C})$. This fact permits us to construct explicit algebraic models of generalized Fermat manifolds of type $(d; k, n)$ (see Section 2) as a complete intersection of $(n-d)$ Fermat hypersurfaces of dimension d and of degree k in \mathbb{P}^n (as was noted by Hirzebruch in the case $d = 2$). As a matter of completeness, in Appendix B (see Theorem 7), we provide an argument, based on basic covering space theory, to obtain these algebraic models.

The strategy of the proof of Theorem (A) is as follows. Let (M, H) be a generalized Fermat pair of type $(d; k, n)$, described algebraically as in Section 2. In this section, we also describe the locus of fixed points of each element of H .

We first observe that (see Lemma 1 in Section 3.1.1) $\text{Lin}(M)$ consists of matrices such that only an element in each row and each column is non-zero. Then, we use such an information (see Lemma 2) to prove that inside $\text{Lin}(M)$ the generalized Fermat group H is unique. Next, as the generalized Fermat manifold M is a non-singular complete intersection, its Picard group is torsion-free. So, if either (i) $d \geq 3$ or (ii) $d = 2$ with non-trivial canonical sheaf, then $\text{Aut}(M) = \text{Lin}(M)$ (see [22]). We verify that the canonical sheaf is not trivial if $(d; k, n) \notin \{(2; 2, 5), (2; 4, 3)\}$. Note that the two exceptional tuples correspond to the unique K3 surfaces in the family of generalized Fermat surfaces.

Notations:

- (1) If $n \geq 1$ and $k \geq 2$ are integers, then $\mathbb{Z}_k^n := \mathbb{Z}_k \times \cdots \times \mathbb{Z}_k$, where \mathbb{Z}_k denotes the cyclic group of order k .
- (2) We denote by \mathbb{P}^d the d -dimensional complex projective space.
- (3) The divisor D given by the union of $n+1$ hyperplanes $L_1, \dots, L_{n+1} \subset \mathbb{P}^d$, where $n \geq 2$, which are in general position, is called a strict normal crossing divisor.
- (4) We denote by M_{d+1}^k the degree k Fermat hypersurface in \mathbb{P}^d . For $d = 1$ (resp., $d = 2$) we also use the notation C_2^k (resp., S_3^k).

- (5) For $n \geq d + 1$ and a strict normal crossing divisor $D = L_1 + \cdots + L_{n+1}$ (where, for $d = 1$, D consists of $n + 1$ different points of \mathbb{P}^1), we will use the notation $M_n^k(D)$ for a generalized Fermat manifold of type $(d; k, n)$ associated to D . In the special case $d = 1$ (resp., $d = 2$), we will also use the notation $C_n^k(D)$ (resp., $S_n^k(D)$) for a generalized Fermat manifold of type $(1; k, n)$ (resp., $(2; k, n)$). In [10], it is proven that $C_n^k(D)$ is uniquely determined by the isomorphism class of D (class defined by the automorphisms of \mathbb{P}^1).
- (6) Let $X \subset \mathbb{P}^n$ be a smooth complex projective variety. In this article, by an automorphism of X we will mean a regular automorphism, so it is a holomorphic automorphism of the underlying complex manifold structure of X . Its group of automorphisms (respectively, group of linear automorphisms) will be denoted by $\text{Aut}(X)$ (respectively, $\text{Lin}(X)$).

2. ALGEBRAIC MODELS OF GENERALIZED FERMAT MANIFOLDS

We describe (algebraic) models of generalized Fermat manifolds of type $(d; k, n)$. These models are obtained as a consequence of Pardini's results [23] (but in Appendix 6, we provide another arguments to obtain them).

2.1. Hyperplanes in general position. There is a natural one-to-one correspondence between hyperplanes and points in the complex projective space \mathbb{P}^d . This correspondence comes from the fact that each hyperplane $L \subset \mathbb{P}^d$ (with projective coordinates $[t_1 : \cdots : t_{d+1}]$) has the form $L_q := \{\rho_1 t_1 + \cdots + \rho_{d+1} t_{d+1} = 0\}$, $q := [\rho_1 : \cdots : \rho_{d+1}] \in \mathbb{P}^d$.

A collection $L_{q_1}, \dots, L_{q_{n+1}} \subset \mathbb{P}^d$, where $n \geq d + 1$, of hyperplanes are in *general position* if the corresponding (pairwise different) points $q_1, \dots, q_{n+1} \in \mathbb{P}^d$ are in general position. This means that, for $d \geq 2$, any subset of $3 \leq s \leq d + 1$ of these points spans a $(s - 1)$ -plane $\Sigma \subset \mathbb{P}^d$. In this above situation, the divisor D , formed by the hyperplanes $L_{q_1}, \dots, L_{q_{n+1}}$, is a strict normal crossing divisor. If $q_j = [\rho_{1,j} : \cdots : \rho_{d+1,j}] \in \mathbb{P}^d$, for $j = 1, \dots, n + 1$, then the hyperplanes $L_{q_1}, \dots, L_{q_{n+1}}$ are in general position if and only if, for every all $(d + 1 \times d + 1)$ -minors of the matrix

$$M(q_1, \dots, q_{n+1}) := \begin{pmatrix} \rho_{1,1} & \cdots & \rho_{1,n+1} \\ \vdots & \vdots & \vdots \\ \rho_{d+1,1} & \cdots & \rho_{d+1,n+1} \end{pmatrix} \in M_{(d+1) \times (n+1)}(\mathbb{C})$$

are nonzero. For example, if $e_1 = [1 : 0 : \cdots : 0]$, $e_2 = [0 : 1 : 0 : \cdots : 0]$, \dots , $e_{d+1} = [0 : \cdots : 0 : 1]$ and $e_{d+2} = [1 : \cdots : 1]$, then the hyperplanes $L_{e_1}, \dots, L_{e_{d+2}}$ are in a general position.

2.2. The parameter spaces $X_{n,d}$ and $\overline{X}_{n,d}$. Let $n \geq d + 1$. Two ordered tuples $(L_{q_1}, \dots, L_{q_{n+1}})$ and $(L_{p_1}, \dots, L_{p_{n+1}})$, of hyperplanes in \mathbb{P}^d in general position, are equivalent if there is a $T \in \text{PGL}_2(\mathbb{C})$ such that $L_{p_j} = T(L_{q_j})$, for $j = 1, \dots, n + 1$. We denote by $X_{n,d}$ the set of such equivalence classes.

If $(L_{q_1}, \dots, L_{q_{n+1}})$ is one of such tuples, then there is a unique $T \in \text{PGL}_{d+1}(\mathbb{C})$ such that $T(L_{q_j}) = L_{e_j}$, for $j = 1, \dots, d + 2$. Then, for each $j = d + 3, \dots, n + 1$, there is some unique $\Lambda_i := [\lambda_{i,1} : \cdots : \lambda_{i,d} : 1] \in \mathbb{P}^d$, $i = 1, \dots, n - d - 1$, such that $T(L_{q_j}) = L_{\Lambda_{j-d-2}}$, for $j = d + 3, \dots, n + 1$. In this case, we set $\Lambda := (\lambda_1, \dots, \lambda_d) \in \mathbb{C}^{d(n-d-1)}$, where $\lambda_j := (\lambda_{1,j}, \dots, \lambda_{n-d-1,j}) \in \mathbb{C}^{n-d-1}$, for $j = 1, \dots, d$. So, if we set

$$\begin{aligned} L_j(\Lambda) &:= L_{e_j} \subset \mathbb{P}^d, \quad j = 1, \dots, d + 2, \\ L_j(\Lambda) &:= L_{\Lambda_{j-d-2}} \subset \mathbb{P}^d, \quad j = d + 3, \dots, n + 1, \end{aligned}$$

then $(L_1(\Lambda), \dots, L_{n+1}(\Lambda))$ is equivalent to $(L_{q_1}, \dots, L_{q_{n+1}})$. We call the tuple Λ a standard parameter.

This observation permits to make the identification (i) $X_{d+1,d} := \{(1, \cdot^d, 1)\}$ and (ii) for $n \geq d + 2$, $X_{n,d} \subset \mathbb{C}^{d(n-d-1)}$ as the set formed by the tuples $\Lambda \in \mathbb{C}^{d(n-d-1)}$ such that the $n + 1$ hyperplanes $L_1(\Lambda), \dots, L_{n+1}(\Lambda)$ are in a general position. We observe that, in this case, $X_{n,d}$ is a (non-empty) open set.

The symmetric group \mathfrak{S}_{n+1} acts by permuting the $n + 1$ coordinates of an ordered tuple of $n + 1$ hyperplanes in general position. Such an action induces an action of a group $\mathbb{G}_{n,d}$ of holomorphic automorphisms on $X_{n,d}$. The quotient complex orbifold $\overline{X}_{n,d} = X_{n,d}/\mathbb{G}_{n,d}$ provides a parameter space of (unordered) collection of $n + 1$ hyperplanes in general position. More details on this can be found in Appendix 5.

2.3. Algebraic models of generalized Fermat pairs, $n \geq d + 1$. We assume $n \geq d + 1$, $k \geq 2$ and $d \geq 1$. Let $\Lambda := (\lambda_1, \dots, \lambda_d) \in X_{n,d}$, where $\lambda_j := (\lambda_{1,j}, \dots, \lambda_{n-d-1,j}) \in \mathbb{C}^{n-d-1}$. Let us consider the complex projective algebraic set

$$(1) \quad M_n^k(\Lambda) := \left\{ \begin{array}{l} x_1^k + \dots + x_d^k + x_{d+1}^k + x_{d+2}^k = 0 \\ \lambda_{1,1}x_1^k + \dots + \lambda_{1,d}x_d^k + x_{d+1}^k + x_{d+3}^k = 0 \\ \vdots \\ \lambda_{n-d-1,1}x_1^k + \dots + \lambda_{n-d-1,d}x_d^k + x_{d+1}^k + x_{n+1}^k = 0 \end{array} \right\} \subset \mathbb{P}^n,$$

So, for $d = 1$ (i.e., $\Lambda = \lambda_1 \in \mathbb{C}^{n-2}$), the above is the generalized Fermat curve of type (k, n) [10]

$$(2) \quad C_n^k(\lambda_1) := \left\{ \begin{array}{l} x_1^k + x_2^k + x_3^k = 0 \\ \lambda_{1,1}x_1^k + x_2^k + x_4^k = 0 \\ \vdots \\ \lambda_{n-2,1}x_1^k + x_2^k + x_{n+1}^k = 0 \end{array} \right\} \subset \mathbb{P}^n,$$

and, for $d = 2$ (i.e., $\Lambda = (\lambda_1, \lambda_2) \in \mathbb{C}^{2(n-3)}$), is the algebraic surface

$$(3) \quad S_n^k(\lambda_1, \lambda_2) := \left\{ \begin{array}{l} x_1^k + x_2^k + x_3^k + x_4^k = 0 \\ \lambda_{1,1}x_1^k + \lambda_{1,2}x_2^k + x_3^k + x_5^k = 0 \\ \vdots \\ \lambda_{n-3,1}x_1^k + \lambda_{n-3,2}x_2^k + x_3^k + x_{n+1}^k = 0 \end{array} \right\} \subset \mathbb{P}^n.$$

Remark 1. If $n = d + 1$, then we will use the notations $C_2^k = \{x_1^k + x_2^k + x_3^k = 0\} \subset \mathbb{P}^2$, $S_3^k := \{x_1^k + x_2^k + x_3^k + x_4^k = 0\} \subset \mathbb{P}^3$ and $M_{d+1}^k := \{x_1^k + x_2^k + \dots + x_{d+2}^k = 0\} \subset \mathbb{P}^{d+1}$.

Theorem 1. *If $\Lambda \in X_{n,d}$, then $M_n^k(\Lambda)$ is an irreducible nonsingular complete intersection.*

Proof. Let us observe that the matrix of coefficients of (1) has rank $(m + 1) \times (m + 1)$ and all of its $(m + 1) \times (m + 1)$ -minors are different from zero (this is the general position condition of the $n + 1$ hyperplanes). The result follows from [32, Proposition 3.1.2]. For explicitness, let us work out the case $d = 2$ (the case $d = 1$ was noted in [10]).

Set $\lambda_{0,1} = \lambda_{0,2} = 1$ and consider the following degree k homogeneous polynomials $f_i := \lambda_{i,1}x_1^k + \lambda_{i,2}x_2^k + x_3^k + x_{4+i}^k \in \mathbb{C}[x_1, \dots, x_{n+1}]$, where $i \in \{0, 1, \dots, n - 3\}$. Let $V_i \subset \mathbb{P}^n$ be the hypersurface given as the zero locus of f_i .

The algebraic set $S_n^k(\lambda_1, \lambda_2)$ is the intersection of the $(n-2)$ hypersurfaces V_0, \dots, V_{n-3} . We consider the matrix of ∇f_i written as rows.

$$(4) \quad \begin{pmatrix} kx_1^{k-1} & kx_2^{k-1} & kx_3^{k-1} & kx_4^{k-1} & 0 & \dots & 0 \\ \lambda_{1,1}kx_1^{k-1} & \lambda_{1,2}kx_2^{k-1} & kx_3^{k-1} & 0 & kx_5^{k-1} & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \lambda_{n-3,1}kx_1^{k-1} & \lambda_{n-3,2}kx_2^{k-1} & kx_3^{k-1} & 0 & \dots & 0 & kx_{n+1}^{k-1} \end{pmatrix}.$$

By the defining equations of the curve, and as $(\lambda_1, \lambda_2) \in X_n$, we see that a point which has three variables $x_i = x_j = x_l = 0$ for $i \neq j \neq l \neq i$ and $1 \leq i, j, l \leq n+1$ has also $x_t = 0$ for $t = 1, \dots, n+1$. Therefore the above matrix has the maximal rank $n-2$ at all points of the surface. So the defining hypersurfaces are intersecting transversally and the corresponding algebraic surface they define is a nonsingular complete intersection.

Next, we proceed to see that the ideal $I_{k,n}$, defined by the $n-2$ equations defining $S_n^k(\lambda_1, \lambda_2) \subset \mathbb{P}^n$ is prime. This follows similarly as in [22, sec. 3.2.1]. Observe first that the defining equations f_0, \dots, f_{n-3} form a regular sequence, that $\mathbb{C}[x_1, \dots, x_{n+1}]$ is a Cohen-Macaulay ring and that the ideal $I_{k,n}$ they define is of codimension $n-2$. The ideal $I_{k,n}$ is prime as a consequence of the Jacobian Criterion [8, Th. 18.15], [22, Th. 3.1] and that the fact that the matrix (4) is of maximal rank $n-2$ on $S_n^k(\lambda_1, \lambda_2)$. In remark [22, 3.4] it is pointed out that an ideal I is prime if the singular locus of the algebraic set defined by I has big enough codimension (in our case, the singular set is the empty set). \square

Let $H_0 = \langle \varphi_1, \dots, \varphi_{n+1} \rangle$, where

$$\varphi_j([x_1 : \dots : x_j : \dots : x_{n+1}]) := [x_1 : \dots : w_k x_j : \dots : x_{n+1}],$$

and w_k is a primitive k -th root of unity. Let $\text{Fix}(\varphi_j) \subset M_n^k(\Lambda)$ be the set of fixed points of φ_j and $F(H_0) := \cup_{j=1}^{n+1} \text{Fix}(\varphi_j)$. The following facts can be deduced from the above.

- (I) $H_0 \cong \mathbb{Z}_k^n$.
- (II) $\varphi_1 \varphi_2 \dots \varphi_{n+1} = 1$.
- (III) $H_0 < \text{Aut}(M_n^k(\Lambda)) < \text{PGL}_{n+1}(\mathbb{C})$.
- (IV) The only non-trivial elements of H_0 with fixed set points being of maximal dimension $d-1$ are the non-trivial powers of the generators $\varphi_1, \dots, \varphi_{n+1}$ (see Proposition 2). Moreover, $\text{Fix}(\varphi_j) := \{x_j = 0\} \cap M_n^k(\Lambda)$, which is isomorphic to a generalized Fermat manifold of type $(d-1; k, n-1)$ for $d \geq 2$ and a collection of k^{n-1} points for $d = 1$. We call the set $\varphi_1, \dots, \varphi_{n+1}$ a canonical generators of H_0 (canonical generators are not unique, but any other set of them is of the form $\varphi_1^r, \dots, \varphi_{n+1}^r$, where $r \in \{1, \dots, k-1\}$ is relatively prime to k).

Remark 2. If we set the subgroups $H_j := \langle \varphi_1, \dots, \varphi_{j-1}, \varphi_{j+1}, \dots, \varphi_n \rangle$, $1 \leq j \leq n$, and $H_{n+1} := \langle \varphi_1, \dots, \varphi_{n-2}, \varphi_{n+1} \rangle$, then $H_j \cong \mathbb{Z}_k^{n-1}$. For $d \geq 2$, $H_j < \text{Aut}(F_j)$ and there is a holomorphic map $\pi_j : F_j \rightarrow L_j(\Lambda) = \mathbb{P}^{d-1}$ (given by the restriction of π_0), which is a Galois holomorphic branched cover with deck group H_j , such that its branch locus is given by the collection of the n intersections $\pi(F_j) \cap \pi(F_i)$, $i \neq j$, which are copies of \mathbb{P}^{d-2} in general position. In particular, (F_j, H_j) is a generalized Fermat pair of type $(d-1; k, n-1)$. This permits to study generalized Fermat manifolds from an inductive point of view.

Remark 3. In [32], T. Terasoma studied a family (depending on $n+1$ parameters) $X_{r,n,k} \subset \mathbb{P}^n$, being a complete intersection of $r+1$ Fermat hypersurfaces of degree $k \geq 2$, where $r+1 < n$, and obtained a decomposition of its Hodge structure $H_{\text{prim}}^{n-r-1}(X_{r,n,k}, \mathbb{Q})$. These algebraic varieties admit a group of automorphisms $H \cong \mathbb{Z}_k^n$ such that X/H is \mathbb{P}^{n-r-1} ,

and its branch locus is given by exactly $n + 1$ linear hyperplanes in general position. It follows that $X_{r,n,k}$ is an example of a generalized Fermat manifold of type $(n - r - 1; k, n)$. Now, for instance, if we take $r = n - 3$, the surface $X_{n-3,n,k}$ is an example of a generalized Fermat surface of type $(2; k, n)$. As generalized Fermat surfaces of type $(2; k, n)$ have moduli dimension $2n - 6$, for $n > 7$ Terasoma's examples form a strict subfamily.

Theorem 2. *($M_n^k(\Lambda), H_0$) is a generalized Fermat pair of type $(d; k, n)$. In particular, every generalized Fermat manifold of type $(d; k, n)$ whose associated branching divisor is $\text{PGL}_{d+1}(\mathbb{C})$ equivalent to the divisor associated to Λ is isomorphic to $M_n^k(\Lambda)$.*

Proof. The map

$$(5) \quad \pi_0 : M_n^k(\Lambda) \rightarrow \mathbb{P}^d : [x_1 : \cdots : x_{n+1}] \mapsto [x_1^k : \cdots : x_{d+1}^k]$$

is a regular branched cover with deck group H_0 and whose branch set is the union of the previous hyperplanes $L_1(\Lambda), \dots, L_{n+1}(\Lambda)$, each one of order k . In other words, the pair $(M_n^k(\Lambda), H_0)$ is a generalized Fermat pair of type $(d; k, n)$. The last part is a consequence of Pardini's building data (or the arguments provided in Appendix 6). \square

Remark 4. If $d \geq 2$, then (as $M_n^k(\Lambda)$ is an irreducible nonsingular complete intersection projective variety of dimension d) it follows that $M_n^k(\Lambda)$ is simply connected (this result is attributed to Lefschetz. For more information, see [12]).

Remark 5 (A fiber product of classical Fermat hypersurfaces). If $\Lambda = (\lambda_1, \dots, \lambda_d) \in X_{n,d}$, and setting $\lambda_{0,i} = 1$, for $i = 1, \dots, d$ and, for each $j \in \{0, 1, \dots, n - d - 1\}$, we consider the classical degree k Fermat hypersurface $F_j = \{\lambda_{j,1}x_1^k + \cdots + \lambda_{j,d}x_d^k + x_{d+1}^k + x_{d+2+j}^k = 0\} \subset \mathbb{P}^{d+1}$ and the rational map $\pi_j : F_j \rightarrow \mathbb{P}^d$ defined by $\pi_j([x_1 : \cdots : x_{d+1} : x_{d+2+j}]) = [x_1^k : \cdots : x_{d+1}^k]$. The branch values of π_j are given by the hyperplanes $L_i(\Lambda)$, for $i = 1, \dots, d + 1$, and $L_{d+2+j}(\Lambda)$. If we consider the fiber product of all these pairs $(F_0, \pi_0), \dots, (F_{n-d-1}, \pi_{n-d-1})$, then we obtain a reducible projective algebraic variety with the action of the group $\mathbb{Z}_k^{n(n-d)}$ and quotient \mathbb{P}^d with branching divisor given by the previous $n + 1$ hyperplanes. This fiber product has $k^{(n-d)(d+1)-n}$ irreducible components, each of them isomorphic to $M_n^k(\Lambda)$.

2.4. Some cohomological information. The fact that $M := M_n^k(\Lambda)$ is a complete intersection variety allows us to compute the cohomology groups of the twisting sheaf $\mathcal{O}_M(r)$ in a relatively direct way, and in particular, to obtain the following.

Proposition 1. *Let $\Lambda \in X_{n,d}$, $n \geq d + 1$, and $M := M_n^k(\Lambda)$. Then*

(1) *The plurigenera $P_m(M)$ of M satisfies*

$$P_m(M) = \frac{k^{n-d}((n-d)k - n - 1)^d}{d!} m^d + O(m^{d-1}).$$

(2) *The arithmetic genus $p_a(M)$ and the geometric genus $p_g(M)$ are given by*

$$p_a(M) = p_g(M) = \begin{cases} 0 & \text{if } r_1 < 0 \\ \binom{r_1+n}{n} & \text{if } 0 \leq r_1 < k \\ \sum_{j \in \Delta_{r_1}} \binom{r_1 - \bar{j} + d}{d} & \text{if } r_1 \geq k \end{cases}$$

(3) *If $(n-d)k - n - 1 = 0$, then M is a Calabi-Yau variety.*

(4) *If $d = 2$, then M is a general type surface except for the rational varieties cases $(k, n) \in \{(2, 3), (3, 3), (2, 4)\}$ and the K3 varieties $(k, n) \in \{(4, 3), (2, 5)\}$.*

Proof. Let $\mathbb{C}[x_1, \dots, x_m]_l$ be the homogeneous polynomials of degree l .

- (a) We first proceed to describe the cohomology groups of the twisting sheaf $\mathcal{O}_M(r)$, $r \in \mathbb{Z}$.
- (a1) Let $\Delta_r := \{(j_1, \dots, j_{n-d}) \in \mathbb{Z}^{n-d} : 0 \leq j_i \leq k-1, 0 \leq i \leq n-d, \text{ and } \bar{j} := j_1 + j_2 + \dots + j_{n-d} \leq r\}$. Then

$$H^0(M, \mathcal{O}_M(r)) := \begin{cases} 0 & \text{if } r < 0 \\ \mathbb{C}[x_1, \dots, x_{n+1}]_r & \text{if } 0 \leq r < k \\ \bigoplus_{j \in \Delta_r} Q_j & \text{if } r \geq k \end{cases}$$

where $Q_j := \mathbb{C}[x_1, \dots, x_{d+1}]_{(r-\bar{j})} x_{d+2}^{j_1} x_{d+3}^{j_2} \dots x_{n+1}^{j_{n-d}}$, $j := (j_1, \dots, j_{n-d})$.

- (a2) By Grothendieck's vanishing theorem we have that:

$$H^i(M, \mathcal{O}_M(r)) = 0 \text{ for } i > d \text{ and } r \in \mathbb{Z},$$

- (a3) and as M is a complete intersection variety we obtain that:

$$H^i(M, \mathcal{O}_M(r)) = 0 \text{ for } 0 < i < d \text{ and } r \in \mathbb{Z},$$

(see page 231 of [13]).

- (a4) Finally, using the Serre duality, we obtain that:

$$H^d(M, \mathcal{O}_M(r)) \cong H^0(M, \mathcal{O}_M(r_1 - r))$$

Remember that $\omega_M \cong \mathcal{O}_M(r_1)$, $r_1 = (n-d)k - n - 1$, (see page 188 of [13]).

- (b) With the former we can calculate the plurigenus of M

$$P_m(M) = \dim_{\mathbb{C}} H^0(M, \omega_M^{\otimes m}) = \dim_{\mathbb{C}} H^0(M, \mathcal{O}_M(r_m))$$

where $r_m := mr_1 = m((n-d)k - n - 1)$.

- (b1) If $(n-d)k - n - 1 < 0$, we obtain that $P_m(M) = 0$. This implies that the Kodaira dimension of M is $\kappa(M) = -\infty$.
- (b2) If $(n-d)k - n - 1 = 0$, we obtain that $P_m(M) = 1$. This implies that the Kodaira dimension of M is $\kappa(M) = 0$.
- (b3) If $(n-d)k - n - 1 > 0$, the canonical sheaf is very ample and

$$P_m(M) = \begin{cases} \binom{r_m + n}{n} & \text{if } 0 \leq r_m < k \\ \sum_{j \in \Delta_{r_m}} \binom{r_m - \bar{j} + d}{d} & \text{if } r_m \geq k \end{cases}$$

In particular, if $r_m \geq \max\{k, (n-d)(k-1)\}$, we obtain that

$$P_m(M) = \frac{k^{n-d}((n-d)k - n - 1)^d}{d!} m^d + O(m^{d-1})$$

This implies that the Kodaira dimension of M is $\kappa(M) = d$.

- (c) The former also permits us to determine the arithmetic genus and geometric genus of M . As seen from the above,

$$p_a(M) = p_g(M) = \dim_{\mathbb{C}} H^d(M, \mathcal{O}_M) = \dim_{\mathbb{C}} H^0(M, \mathcal{O}_M(r_1)),$$

so

$$p_a(M) = p_g(M) = \begin{cases} 0 & \text{if } r_1 < 0 \\ \binom{r_1 + n}{n} & \text{if } 0 \leq r_1 < k \\ \sum_{j \in \Delta_{r_1}} \binom{r_1 - \bar{j} + d}{d} & \text{if } r_1 \geq k \end{cases}$$

□

2.5. On the locus of fixed points of elements of H_0 . The diagonal presentation of the generators $\varphi_j \in H_0$, for $j = 1, \dots, n+1$, permits us to obtain the locus of fixed points of every element of H_0 (see Proposition 2 below). Let us start with some definitions and notations. For each tuple $(m_1, \dots, m_{n+1}) \in \{0, 1, \dots, k-1\}^{n+1}$ and each $l \in \{0, 1, \dots, k-1\}$, we set

$$L_l(m_1, \dots, m_{n+1}) := \{j \in \{1, \dots, n+1\} : m_j = l\}$$

and the (possibly empty) algebraic sets

$$\tilde{F}_l(m_1, \dots, m_{n+1}) = \{[x_1 : \dots : x_{n+1}] \in \mathbb{P}^n : x_i = 0, \forall i \notin L_l(m_1, \dots, m_{n+1})\}.$$

and

$$F_l(m_1, \dots, m_{n+1}) := \tilde{F}_l(m_1, \dots, m_{n+1}) \cap M_n^k(\Lambda)$$

Example 1. Let us take $n = k = 3$ and $d = 2$. In this case, S_3^3 is just the Fermat hypersurface $\{x_1^3 + x_2^3 + x_3^3 + x_4^3 = 0\} \subset \mathbb{P}^3$. If $\varphi = \varphi_1 \varphi_2 \varphi_3^2$, then $(m_1, m_2, m_3, m_4) = (1, 1, 2, 0)$ and $L_0(1, 1, 2, 0) = \{4\}$, $L_1(1, 1, 2, 0) = \{1, 2\}$, $L_2(1, 1, 2, 0) = \{3\}$. The locus of fixed points (in \mathbb{P}^3) of φ is given by

$$\begin{aligned} & \tilde{F}_0(1, 1, 2, 0) \cup \tilde{F}_1(1, 1, 2, 0) \cup \tilde{F}_2(1, 1, 2, 0) = \\ & \{[0 : 0 : 0 : 1]\} \cup \{[x_1 : x_2 : 0 : 0] \in \mathbb{P}^3\} \cup \{[0 : 0 : 1 : 0]\}. \end{aligned}$$

As the cardinalities of $L_0(1, 1, 2, 0)$ and $L_2(1, 1, 2, 0)$ are equal to $1 \leq n - d$, these two do not introduce fixed points of φ on S_3^3 (as also can be seen directly). The set $L_1(1, 1, 2, 0)$ has cardinality $2 \geq n - d + 1$, so it produces a zero-dimensional set of fixed points consisting of the three points $[1 : -1 : 0]$, $[1 : \omega_6 : 0]$ and $[1 : \omega_6^{-1} : 0]$, where $\omega_6 = e^{\pi i/3}$.

In the general case, we have:

Proposition 2. *Let $d \geq 1$, $n \geq d + 1$, $k \geq 2$, $\Lambda \in X_{n,d}$, $M_n^k(\Lambda)$, $H_0 \cong \mathbb{Z}_k^n$ and $\varphi_1, \dots, \varphi_{n+1} \in H_0$ as before. Then:*

- (1) $F_l(m_1, \dots, m_{n+1}) \neq \emptyset$ if and only if $\#L_l(m_1, \dots, m_{n+1}) \geq n + 1 - d$.
- (2) If $F_l(m_1, \dots, m_{n+1}) \neq \emptyset$, then it is a generalized Fermat manifold of dimension $\#L_l(m_1, \dots, m_{n+1}) + d - n - 1$.
- (3) If $m_1, \dots, m_{n+1} \in \{0, 1, \dots, k-1\}$, then the locus of fixed points of $\varphi := \varphi_1^{m_1} \dots \varphi_{n+1}^{m_{n+1}}$ is the disjoint union of the sets $F_l(m_1, \dots, m_{n+1})$, where $l \in \{1, \dots, k-1\}$. In particular, the number of (non-empty) connected components of its locus of fixed points (if non-empty) equals the number of exponents l appearing in φ at least $n + 1 - d$ times.

Proof. The locus of fixed points, in \mathbb{P}^n , of φ is the disjoint union of the algebraic sets

$$\tilde{F}_l(m_1, \dots, m_{n+1}) = \{[x_1 : \dots : x_{n+1}] \in \mathbb{P}^n : x_i = 0, \forall i \notin L_l(m_1, \dots, m_{n+1})\},$$

where the union is taking on all those $l \in \{1, \dots, k-1\}$ such that $\#L_l(m_1, \dots, m_{n+1}) \geq 1$. Note that each $\tilde{F}_l(m_1, \dots, m_{n+1})$ is: (i) just a point if $\#L_l(m_1, \dots, m_{n+1}) = 1$, and (ii) a projective space of dimension $\#L_l(m_1, \dots, m_{n+1}) - 1$ if $\#L_l(m_1, \dots, m_{n+1}) > 1$.

The locus of fixed points of φ on $M_n^k(\Lambda)$ is then given as the disjoint union of the sets $F_l(m_1, \dots, m_{n+1}) = \tilde{F}_l(m_1, \dots, m_{n+1}) \cap M_n^k(\Lambda)$.

But on $M_n^k(\Lambda)$ we cannot have points $[x_1 : \dots : x_{n+1}]$ with at least $d + 1$ coordinates equal to zero. This fact asserts that for $\#L_l(m_1, \dots, m_{n+1}) \leq n - d$ one has that $F_l(m_1, \dots, m_{n+1}) = \emptyset$.

Also, for $\#L_l(m_1, \dots, m_{n+1}) \geq n + 1 - d$, we obtain that $F_l(m_1, \dots, m_{n+1}) \neq \emptyset$ is a generalized Fermat manifold of dimension $\#L_l(m_1, \dots, m_{n+1}) + d - n - 1$. \square

Remark 6. If $k = 2$, then the above proposition asserts that $\varphi_1^{m_1} \cdots \varphi_{n+1}^{m_{n+1}}$ has no fixed points if and only if $\#L_0(m_1, \dots, m_{n+1}), \#L_1(m_1, \dots, m_{n+1}) \leq n - d$. As both cardinalities add to $n + 1$, the previous will only be possible if $n \geq 1 + 2d$. We will see later (see Example 2), for $d \geq 2$, that if $H \cong \mathbb{Z}_p^n$, where $p \geq 2$ is a prime integer, admits subgroups $K \cong \mathbb{Z}_p^{n-r}$ acting freely, then necessarily $n + 1 \leq (p^r - 1)/(p - 1)$. In particular, the case $r = 1$ is not possible and, for $p = 2$, neither $r = 2$. Note, for $p = 2$, that this is in contrast to the case $d = 1$ where these type of subgroups exists and are related to hyperelliptic Riemann surfaces.

Example 2 (Case $k = p$ prime). If $k = 2$, then $m_1, \dots, m_{n+1} \in \{0, 1\}$. The above proposition asserts that $\varphi := \varphi_1^{m_1} \cdots \varphi_{n+1}^{m_{n+1}}$ has no fixed points on $M_n^2(\Lambda)$ if and only if $\#L_0(m_1, \dots, m_{n+1}), \#L_1(m_1, \dots, m_{n+1}) \leq n - d$. As these two cardinalities must add to $n + 1$, this is only possible for $n \geq 1 + 2d$.

Let $d = 1, n \geq 4$ and $C := C_n^2(\Lambda)$ be a generalized Fermat curve of type $(1; 2, n)$ with corresponding generalized Fermat group $H_0 \cong \mathbb{Z}_2^n$. If $n \geq 5$ odd, then there is a (unique) subgroup $K \cong \mathbb{Z}_2^{n-1}$ of H_0 acting freely on C . For $n \geq 4$ even, the above is not true, but there are subgroups $K \cong \mathbb{Z}_2^{n-2}$ acting freely.

In the case $d \geq 2$ the above situation does not hold. In fact, let $k = p \geq 2$ be a prime integer and let $n \geq d + 1$ and $M = M_n^p(\Lambda)$ be a generalized Fermat manifold of type $(d; p, n)$ with associated generalized Fermat group $H_0 \cong \mathbb{Z}_p^n$. Assume $K \cong \mathbb{Z}_p^{n-r}$ is a subgroup of H_0 acting freely on M . Let $F_j \subset M, j = 1, \dots, n + 1$, be the locus of fixed points of the canonical generator φ_j . As H_0 is an abelian group, each F_j is invariant under K and acts freely on it. Let $N = M/K$ (which is a compact complex manifold of dimension d) and $X_j = F_j/K$ (a connected complex submanifold of N). The $(n + 1)$ connected sets X_j are the locus of fixed points of the induced holomorphic automorphism by φ_j . As each two different F_i and F_j always intersect transversely, it follows that the same happens for X_i and X_j . As the locus of fixed points of (finite) holomorphic automorphisms are smooth, it follows that different X_i and X_j are the fixed points of different cyclic groups of $A = H_0/K \cong \mathbb{Z}_p^r$. This in particular asserts that $n + 1 \leq (p^r - 1)/(p - 1)$. So, for instance, the case $r = 1$ is not possible and, the case $r = 2$ and $p = 2$ is neither possible.

2.6. The exceptional cases $2 \leq n \leq d$. In the above, we have assumed the condition $n \geq d + 1$. In this one, we consider the complementary case $2 \leq n \leq d$. The following result asserts that the only situation holds for $n = d$.

Theorem 3. *If (M, H) is a generalized Fermat pair of type $(d; k, n)$, where $d \geq 2$ and $2 \leq n \leq d$, then $n = d$ and, up to isomorphisms, $M = \mathbb{P}^d$ and $H = \langle \varrho_1, \dots, \varrho_d \rangle$, where $\varrho_j([x_1 : \cdots : x_{d+1}]) = [x_1 : \cdots : x_{j-1} : \omega_k x_j : x_{j+1} : \cdots : x_{d+1}]$.*

Proof. Let us consider a Galois branched covering $\pi : M \rightarrow \mathbb{P}^d$, with deck group H , and whose branch divisor locus is given by the hyperplanes L_1, \dots, L_{n+1} . Up to post-composition of π by a suitable projective linear automorphisms, we may assume $L_j = \{t_j = 0\}$. Now, we add the hyperplanes L_{n+2}, \dots, L_{d+2} , where, for $j \leq d + 1$, $L_j = \{t_j = 0\}$ and $L_{d+2} = \{t_1 + \cdots + t_{d+1} = 0\}$. Next, consider the orbifold M^{orb} , whose underlying complex manifold is M and whose orbifold divisor is obtained by lifting the hyperplanes L_{n+2}, \dots, L_{d+2} and giving them the branching order equal to k . By construction, the group H preserves that branch divisor of M^{orb} .

Let $M_{d+1}^k \subset \mathbb{P}^{d+1}$ be the generalized Fermat manifold of type $(d; k, d + 1)$ (the Fermat hypersurface of degree k) together its generalized Fermat group $H_0 = \langle \varphi_1, \dots, \varphi_{d+1} \rangle$ and the corresponding Galois branched covering $\pi_0 : M_{d+1}^k \rightarrow \mathbb{P}^d$ with deck group H_0 .

Note that $M^{orb}/H = M_{d+1}^k/H_0$ (which is \mathbb{P}^d with branch divisor $L_1 + \cdots + L_{d+2}$, each one with order k). As the universal covering of this last orbifold is $M_{d+1}^k \subset \mathbb{P}^{d+1}$, the branched covering π_0 it must factor through π , that is, there is some $\mathbb{Z}_k^{d+1-n} \cong K < H_0$ such that $M = M_{d+1}^k/K$. Because of the choices of the hyperplanes L_j , we must have

$$K = \langle \varphi_{n+2}, \dots, \varphi_{d+2} \rangle,$$

where $\varphi_1, \dots, \varphi_{d+2} \in H_0$ are the corresponding standard set of generators.

Let us first assume $n = d$. In this case, $K = \langle \varphi_{d+2} \rangle$. We may consider the (branched) Galois cover, with deck group K ,

$$\pi_K : M_{d+1}^k \rightarrow \mathbb{P}^d : [x_1 : \cdots : x_{d+2}] \mapsto [x_1 : \cdots : x_{d+1}] = [y_1 : \cdots : y_{d+1}].$$

In this way, $M \cong \mathbb{P}^d$ and $H = \langle \varrho_1, \dots, \varrho_d \rangle$, where ϱ_j is as described in the theorem. The corresponding Galois branched cover, with deck group H , is $\pi([y_1 : \cdots : y_{d+1}]) = [y_1^k : \cdots : y_{d+1}^k]$ whose branch divisor is $L_1 + \cdots + L_{d+1}$.

Now, if $2 \leq n < d$, then $M = M_{d+1}^k/K = \mathbb{P}^d/\langle \varrho_{n+2}, \dots, \varrho_{d+1} \rangle$. But, the point $[0 : \cdots : 0 : 1]$ (which is an isolated fixed point of φ_{d+2}) produces a singular point of the quotient $\mathbb{P}^d/\langle \varrho_{n+2}, \dots, \varrho_{d+1} \rangle$ as a consequence of the Chevalley-Shephard-Todd theorem [5]. \square

3. THE GROUP OF AUTOMORPHISMS OF GENERALIZED FERMAT MANIFOLDS

3.1. Uniqueness of the generalized Fermat group. In this section, we provide the proof of our main theorem.

Theorem 4. *Let $d \geq 2$, $k \geq 2$, $n \geq d + 1$ be integers, $\Lambda \in X_{n,d}$ and set $M = M_n^k(\Lambda)$. Then:*

- (1) *The elements in $\text{Lin}(M)$ consist of matrices such that only an element in each row and each column is non-zero.*
- (2) *If $(d; k, n) \notin \{(2; 2, 5), (2; 4, 3)\}$, then $\text{Aut}(M) = \text{Lin}(M)$ has a unique generalized Fermat group H of type $(d; k, n)$. If $(d; k, n) \in \{(2; 2, 5), (2; 4, 3)\}$, then the same uniqueness result holds for $\text{Lin}(M)$.*

Remark 7. If $n = d + 1$, then $M = M_{d+1}^k$ is the Fermat hypersurface of degree k , for which can be directly observed the uniqueness of H_0 as in this case $\text{Lin}(M) = H_0 \rtimes \mathfrak{S}_{n+1}$, where the permutation part acts as permutation of coordinates.

Remark 8. If $(d; k, n) \in \{(2; 2, 5), (2; 4, 3)\}$, then $M := M_n^k(\Lambda)$ is a K3 surface (see Section 2.4). If $(d; k, n) = (2; 4, 3)$, then M corresponds to the classical Fermat hypersurface of degree 4 in \mathbb{P}^3 for which $\text{Lin}(M) \cong \mathbb{Z}_4^3 \rtimes \mathfrak{S}_4$ and $\text{Aut}(M)$ infinite. If $(d; k, n) = (2; 2, 5)$, then $\text{Lin}(M)$ is a finite extension of \mathbb{Z}_2^5 (generically a trivial extension) and $\text{Aut}(M)$ is infinite by results due to Shioda and Inose in [31, Thm 5] (in [33] Vinberg computed it for a particular case). In these cases, the generalized Fermat group H (which is unique in $\text{Lin}(M)$) cannot be a normal subgroup of $\text{Aut}(M)$. Otherwise, every element of $\text{Aut}(M)$ will induce an automorphism of \mathbb{P}^2 (permuting the $n + 1$ branched lines), so a finite group of linear automorphism. This will ensure that $\text{Aut}(M)$ is a finite extension of H , so a finite group, a contradiction.

3.1.1. Proof of Theorem 4. If either (i) $d \geq 3$ or (ii) $d = 2$ and $(k, n) \neq (2, 5), (4, 3)$, then it follows from [22] that $\text{Aut}(M) = \text{Lin}(M)$. So, we are left to prove uniqueness of H_0 inside $\text{Lin}(M)$ and that the elements of $\text{Lin}(M)$ have the desired structure.

Let $\sigma \in L = \text{Lin}(M)$. As it is linear, it is given in terms of an $(n+1) \times (n+1)$ matrix $A_\sigma = (a_{i,j}) \in GL_{n+1}(\mathbb{C})$:

$$(6) \quad \sigma([x_1 : \cdots : x_{n+1}]) = \left[\sum_{j=1}^{n+1} a_{1,j} x_j : \cdots : \sum_{j=1}^{n+1} a_{n+1,j} x_j \right].$$

We start by proving the structure of σ .

Lemma 1. *There is only one non-zero element in each column and row of $A_\sigma = (a_{i,j})$.*

Proof. The arguments are independent of $d \geq 1$. In [15] it was done the case $d = 1$. Below, we indicate its simple modification for $d = 2$ (the general case is similar). Set $\lambda = (\lambda_1, \dots, \lambda_{n-3}), \mu = (\mu_1, \dots, \mu_{n-3})$ such that $(\lambda, \mu) \in X_{n,2}$ and $M = S_n^k(\lambda, \mu)$. If we set (by setting $\lambda_0 = \mu_0 = 1$)

$$f_j : \lambda_j x_1^k + \mu_j x_2^k + x_3^k + x_{4+j}^k, \quad j = 0, \dots, n-3,$$

then $S_n^k(\lambda, \mu) = V(f_0, \dots, f_{n-3})$. In this way, for every point $P \in S_n^k(\lambda, \mu)$, then $\sigma(P) \in S_n^k(\lambda, \mu)$, that is, $f_i \circ \sigma = \sigma^*(f_i) \in \langle f_0, \dots, f_{n-3} \rangle$, i.e.

$$(7) \quad f_i \circ \sigma = \sum_{j=0}^{n-3} g_{j,i} f_j,$$

for some constants $g_{j,i} \in \mathbb{C}$.

Let us consider $\sigma^*(\nabla f_i) = \nabla f_i \circ \sigma$. For every point $P \in S_n^k(\lambda, \mu)$ we have

$$(8) \quad \nabla(f_i \circ \sigma)(P) = \sum_{j=0}^{n-3} g_{j,i} \nabla f_j(P),$$

so,

$$(9) \quad \sigma^*(\nabla f_i) A_\sigma = \nabla(f_i \circ \sigma) = \sum_{j=0}^{n-3} g_{j,i} \nabla f_j,$$

that is,

$$(10) \quad \sigma^*(\nabla f_i) = \left(\sum_{j=0}^{n-3} g_{j,i} \nabla f_j \right) A_\sigma^{-1}.$$

Recall that $f_j = \lambda_j x_1^k + \mu_j x_2^k + x_3^k + x_{4+j}^k$ for $0 \leq j \leq n-3$, where $\lambda_0 = \mu_0 = 1$, and

$$Y_j := \nabla f_j = (k\lambda_j x_1^{k-1}, k\mu_j x_2^{k-1}, kx_3^{k-1}, 0, \dots, 0, kx_{j+4}^{k-1}, 0, \dots, 0).$$

The above asserts that

$$\sigma^*(Y_j) = \left(\lambda_j \left(\sum_{i=1}^{n+1} a_{1,i} x_i \right)^{k-1}, \mu_j \left(\sum_{i=1}^{n+1} a_{2,i} x_i \right)^{k-1}, \left(\sum_{i=1}^{n+1} a_{3,i} x_i \right)^{k-1}, 0, \dots, 0, \left(\sum_{i=1}^{n+1} a_{j+4,i} x_i \right)^{k-1}, 0, \dots, 0 \right),$$

where the fourth nonzero element is at the $j+4$ position.

Now, for $t \in \{1, 2, 3, j+4\}$, the t -th coordinate of $\sigma^*(Y_j)$ involves all combinations of the terms

$$\binom{k-1}{\nu_1, \dots, \nu_{n+1}} (a_{t,1}^{\nu_1} \cdots a_{t,n+1}^{\nu_{n+1}}) \cdot (x_1^{\nu_1} \cdots x_{n+1}^{\nu_{n+1}}), \quad \text{for } \nu_1 + \cdots + \nu_{n+1} = k-1,$$

where

$$\binom{k-1}{\nu_1, \dots, \nu_{n+1}} = \frac{(k-1)!}{\nu_1! \cdot \nu_2! \cdots \nu_{n+1}!}$$

Equation (10) asserts that each $\sigma^*(Y_j)$ is a linear combination of Y_0, \dots, Y_{n-3} . So, it only involves combinations of the monomials x_i^{k-1} . The above asserts that in the tuple $\bar{\nu} = (\nu_1, \dots, \nu_{n+1})$ there is only one coordinate $\nu_i = k-1$ and all the others are equal to zero. We set

$$\begin{aligned} \mathbf{x}^{\bar{\nu}} &= x_1^{\nu_1} \cdots x_{n+1}^{\nu_{n+1}}, \\ A_{l, \bar{\nu}} &= a_{l,1}^{\nu_1} \cdots a_{l,n+1}^{\nu_{n+1}}. \end{aligned}$$

In this notation, we have that if the tuple $\bar{\nu}$ has at least two non-zero coordinates, then $\mathbf{x}^{\bar{\nu}}$ cannot appear in the linear combination of Y_i . In this case, by using eq. (10), we must have

$$(A_{1, \bar{\nu}}, \dots, A_{n+1, \bar{\nu}}) \cdot A_\sigma = 0,$$

and, since A_σ is an invertible matrix, the above implies that

$$A_{l, \bar{\nu}} = 0, \quad l = 1, \dots, n+1.$$

Let us assume that, for some j and some $1 \leq l_1 < l_2 \leq n+1$, we have $a_{j, l_1} \neq 0 \neq a_{j, l_2}$. In this case, we consider the tuple $\bar{\nu} = (0, \dots, 0, t, 0, \dots, 0, k-1-t, 0, \dots, 0)$, where $t \in \{1, \dots, k-1\}$ (respectively, $k-1-t$) is in the position l_1 (respectively, in the position l_2). As $\mathbf{x}^{\bar{\nu}}$ cannot appear in the linear combination of Y_j , we must have that $a_{j, l_1}^t a_{j, l_2}^{k-1-t} = A_{j, \bar{\nu}} = 0$, a contradiction. \square

The above lemma permits us to note that H_0 is a normal subgroup of L . Next, we proceed to see that H_0 is the only generalized Fermat group of type $(d; k, n)$ in L . Which ends the proof of our theorem.

Lemma 2. *H_0 is the only generalized Fermat group of type $(d; k, n)$ in L .*

Proof. Let us assume that $H < L$ is another generalized Fermat group of M of type $(d; k, n)$ and let $\varphi_1^*, \dots, \varphi_{n+1}^*$ be an standard set of generators of H . Recall that the only non-trivial elements of H with a set of fixed points of the maximal dimension $d-1$ are the non-trivial powers of these standard generators. If φ_j^* is given by a diagonal matrix, then it is an element of H_0 . Let us assume that one of the standard generators say φ_1^* , is non-diagonal. By Lemma 1, the matrix representation of φ_1^* has exactly one non-zero element in each of its arrows and each of its columns. This means that the locus of fixed points \widehat{F} of φ_1^* in \mathbb{P}^n consists of a finite collection of linear spaces, each one of dimension at most $n-2$. As M is not contained inside a hyperplane, it follows that the locus of fixed points of φ_1^* in M , $F = \widehat{F} \cap M$, must be a finite collection of components each one of dimension at most $d-2$, a contradiction. It follows then that $H = H_0$ are required. \square

3.1.2. Remark: Another approach for uniqueness. Let us assume $n \geq d+2$ and that $H < L$ is another generalized Fermat group of M of type $(d; k, n)$. Let $\varphi_1, \dots, \varphi_{n+1}$ be an standard set of generators of H_0 and $\varphi_1^*, \dots, \varphi_{n+1}^*$ be an standard set of generators for H . Following in a similar way as done in the proof of Claim 1 of Section 5 in [15], we may assume that there are $i, j \in \{1, \dots, n\}$ such that $\langle \varphi_i \rangle = \langle \varphi_j^* \rangle$. Up to a permutation of indices, we may assume $i = j = 1$. So, let $\langle \varphi_1 \rangle = \langle \varphi_1^* \rangle$ and set $F_1 \subset M$ the locus of fixed points of φ_1 , which is a generalized Fermat manifold of type $(d-1; k, n-1)$, there are two generalized Fermat groups of that type, these being $H_0/\langle \varphi_1 \rangle$ and $H/\langle \varphi_1^* \rangle$. So, if we have the uniqueness for dimension $d-1$, then we will be done by an induction process. The starting case is $d = 2$, for which F_1 is a generalized Fermat curve of type

$(k, n - 1)$. The uniqueness, in this case, works if $(k - 1)(n - 2) > 2$ [15]. As we are assuming $n \geq d + 2 = 4$, the uniqueness will hold if either: (i) $n = 4$ and $k \geq 3$ or (ii) $n \geq 5$. So, in these cases, $H_0 = H$.

Let us now consider the left case $n = 4, k = 2$, and, by contradiction, assume $H \neq H_0$. In this case F_1 is a genus one Riemann surface. On it we have the group $J = H_0/\langle \varphi_1 \rangle \cong \mathbb{Z}_2^3$ of conformal automorphisms. If $j \neq 1$, then F_1 is tangent to each F_j (the locus of fixed points of φ_j) at four points (these are the fixed points of the action of φ_j on F_1). The torus F_1 corresponds to the plane generalized Fermat curve of genus one

$$F_1 = \left\{ \begin{array}{l} y_1^2 + y_2^2 + y_3^2 = 0 \\ \mu y_1^2 + y_2^2 + y_4^2 = 0 \end{array} \right\} \subset \mathbb{P}^3.$$

In the above model, the group J is generated by $a_1([y_1 : y_2 : y_3 : y_4]) = [-y_1 : y_2 : y_3 : y_4]$, $a_2([y_1 : y_2 : y_3 : y_4]) = [y_1 : -y_2 : y_3 : y_4]$ and $a_3([y_1 : y_2 : y_3 : y_4]) = [y_1 : y_2 : -y_3 : y_4]$, and the corresponding Galois branched covering is $\pi([y_1 : y_2 : y_3 : y_4]) = -(y_2/y_1)^2 = z$ whose branched values are $\infty, 0, 1$ and μ . Set $a_4 = a_1 a_2 a_3$. As we have assumed above that $\varphi_1 = \varphi_1^*$, then F_1 also admits the group $J' = H/\langle \varphi_1^* \rangle \cong \mathbb{Z}_2^3$ of conformal automorphisms; and it normalizes J . So, J' induces (under π) a group of conformal automorphisms of the Riemann sphere, that permutes the branch values of π , and isomorphic to \mathbb{Z}_2^t , for some $t > 0$ (as we are assuming $H \neq H'$). There are only two possibilities: $t \in \{1, 2\}$ (as the only finite abelian subgroups of Möbius transformations are either cyclic or \mathbb{Z}_2^2). It follows that $J \cap J'$ can be either \mathbb{Z}_2 or \mathbb{Z}_2^2 . Since $H \neq H_0$, one of the standard generators of H is different from the standard generators of H_0 , say φ_2^* . This generator induces an involution α with four fixed points on F_1 different from those in J . We may assume that it induces the involution $a(z) = \mu/z$, in which case $\alpha([y_1 : y_2 : y_3 : y_4]) = [y_2 : \epsilon_1 \sqrt{\mu} y_1 : \epsilon_2 y_4 : \epsilon_3 \sqrt{\mu} y_3]$, where $\sqrt{\mu}$ is a fixed square root of μ , $\epsilon_j \in \{\pm 1\}$ and $\epsilon_1 = \epsilon_2 \epsilon_3$. As $\alpha a_1 \alpha = a_2$ and $\alpha a_3 \alpha = a_4$, there is no subgroup isomorphic to \mathbb{Z}_2^2 inside J such that the group generated by it and α is \mathbb{Z}_2^3 . It follows that $J \cap J'$ cannot be \mathbb{Z}_2^2 . If $J \cap J' = \mathbb{Z}_2$, then J' must induce \mathbb{Z}_2^2 , one generator being a and the other being $b(z) = (z - \mu)/(z - 1)$, this last one induced by some involution $\beta \in J' \setminus J$. It can be seen that $\beta([y_1 : y_2 : y_3 : y_4]) = [y_3 : \epsilon_1 i y_4 : \epsilon_2 \sqrt{1 - \mu} y_1 : \epsilon_3 i \sqrt{1 - \mu} y_2]$, where $\sqrt{1 - \mu}$ is a fixed square root of $1 - \mu$, $\epsilon_j \in \{\pm 1\}$ and $\epsilon_2 = -\epsilon_1 \epsilon_3$. Similarly as for α , one may see that $\beta a_1 \beta = a_3$ and $\beta a_2 \beta = a_4$. It can be checked that there is no involution of J such that together α and β generate \mathbb{Z}_2^3 .

All the above produces a contradiction under the assumption that $H \neq H_0$.

3.2. Automorphisms of generalized Fermat manifolds. Let $d \geq 2, k \geq 2, n \geq d + 1$ be integers, $(d; k, n) \notin \{(2; 2, 5), (2; 4, 3)\}$, and M be a generalized Fermat manifold of type $(d; k, n)$. Then, by Theorem 4, M admits a unique generalized Fermat group H of type $(d; k, n)$. Let $\pi : M \rightarrow \mathbb{P}^d$ be a Galois branched covering, with H as its deck group, and let $\{L_1, \dots, L_{n+1}\}$ be its set of branch hyperplanes. Let G_0 be the $\text{PGL}_{d+1}(\mathbb{C})$ -stabilizer of these $n + 1$ branch hyperplanes.

As H is a normal subgroup of $\text{Aut}(M)$, it follows the existence of a homomorphism $\theta : \text{Aut}(M) \rightarrow G_0$, with kernel H . The universal property of M (see 7) assures that every element Q of G_0 lifts to a holomorphic automorphism \widehat{Q} of M . Then there is a short exact sequence

$$1 \rightarrow H \rightarrow \text{Aut}(M) \rightarrow G_0 \rightarrow 1.$$

In particular, $|\text{Aut}(M)| = |G_0|k^n$. Also, by the Schur-Zassenhaus theorem [7], in the case that the order of G_0 is relatively prime with k , then $\text{Aut}(M) \cong H \rtimes G_0$. We summarize all the above in the following.

Corollary 1. *Let $d \geq 2$, $k \geq 2$, $n \geq d + 1$ be integers and $(d; k, n) \notin \{(2; 2, 5), (2; 4, 3)\}$. Let M be a generalized Fermat manifold of type $(d; k, n)$ and let H be its unique generalized Fermat group of type $(d; k, n)$. If G_0 is the $\mathrm{PGL}_{d+1}(\mathbb{C})$ -stabilizer of the $n + 1$ branch hyperplanes of $M/H = \mathbb{P}^d$, then $|\mathrm{Aut}(M)| = |G_0|k^n$ and, if the order of G_0 is relatively prime with k , then $\mathrm{Aut}(M) \cong H \rtimes G_0$.*

Corollary 2. *Let $d \geq 2$, $k \geq 2$. If G_0 be a finite subgroup of $\mathrm{PGL}_{d+1}(\mathbb{C})$, then there exists a generalized Fermat pair (M, H) of type $(d; k, n)$, for some $n \geq d + 1$, such that $\mathrm{Aut}(M/H) \cong G_0$. In fact, for $|G_0| \leq d + 1$ we may assume $n = d + 1$ and, for $|G_0| \geq d + 2$, we may assume $n = |G_0| - 1$.*

Proof. If $|G_0| \leq d + 1$, then take $n = d + 1$ and note that for the classical Fermat hypersurface $M_n^k \subset \mathbb{P}^n$ of degree k one has that $\mathrm{Aut}(M_n^k)/H$ contains the permutation group of $d + 1$ letters. Let us assume $|G_0| \geq d + 2$. The linear group G_0 induces a linear action on the space $\mathbb{P}_{\mathrm{hyper}}^d$ of hyperplanes of \mathbb{P}^d . As G_0 is finite, we may find (generically) a point $q \in \mathbb{P}_{\mathrm{hyper}}^d$ whose G_0 -orbit is a generic set of points. Such an orbit determines a collection of $|G_0|$ lines in general position in \mathbb{P}^d . Let us observe that, by the generic choice, we may even assume the above set of points to have $\mathrm{PGL}_{d+1}(\mathbb{C})$ -stabilizer exactly G_0 , so the same situation for our collection of hyperplanes. Now the results follow from Corollary 1. \square

3.3. Equivalence of generalized Fermat manifolds. Let us remember that $X_{n,d}$ is the parameter space of the isomorphism classes of strict normal crossing divisors formed by an ordered collection of $n + 1$ hyperplanes. By abuse of notation, given a standard parameter $\Lambda \in X_{n,d}$, we also note Λ the ordered divisor $(L_1(\Lambda), L_2(\Lambda), \dots, L_{n+1}(\Lambda))$.

If $\Lambda := (\lambda_1, \dots, \lambda_d) \in X_{n,d}$, for $n \geq d + 1$, then we have associated a generalized Fermat manifold $M_n^k(\Lambda)$ with its unique generalized Fermat group H of type $(d; k, n)$. We also have a natural Galois branched covering $\pi_0 : M_n^k(\Lambda) \rightarrow \mathbb{P}^d$ defined by $\pi_0([x_1 : \dots : x_{n+1}]) = [x_1^k : \dots : x_{d+1}^k]$. Its branch locus is given by ordered divisor Λ .

As there is only one isomorphism class of generalized Fermat manifold of type $(d; k, d + 1)$, we assume that $n \geq d + 2$.

Lemma 3. *Let $\Lambda, \widehat{\Lambda} \in X_{n,d}$, $n \geq d + 2$, $k \geq 2$ and $(d; k, n) \notin \{(2; 2, 5), (2; 4, 3)\}$. The two generalized Fermat manifolds $M_n^k(\Lambda), M_n^k(\widehat{\Lambda})$ are isomorphic if and only if there is some $T \in \mathrm{PGL}_{d+1}(\mathbb{C})$ sending Λ onto $\widehat{\Lambda}$.*

Proof. If $\phi : M_n^k(\Lambda) \rightarrow M_n^k(\widehat{\Lambda})$ is an isomorphism, then the uniqueness of the generalized Fermat group H ensures the existence of a $T_\phi \in \mathrm{PGL}_{d+1}(\mathbb{C})$ such that $\pi_0 \circ \phi = T_\phi \circ \pi_0$, so sending Λ onto $\widehat{\Lambda}$. Conversely, if we have some $T \in \mathrm{PGL}_{d+1}(\mathbb{C})$ sending Λ onto $\widehat{\Lambda}$, Then $T \circ \pi_0 : M_n^k(\Lambda) \rightarrow \mathbb{P}^d$ is a regular branched cover where the branch locus is $\widehat{\Lambda}$. Then by the Theorem 7 we obtain that there exists a biholomorphism between $M_n^k(\Lambda)$ and $M_n^k(\widehat{\Lambda})$. \square

Now, by Lemma 3, we observe that two points in $X_{n,d}$ are in the same $\mathbb{G}_{n,d}$ -orbit if and only if they define isomorphic generalized Fermat manifolds of the same type $(d; k, n)$. Moreover, if we consider the homomorphism $R : \mathbb{S}_{n+1} \rightarrow \mathbb{G}_{n,d}$ of Appendix 5, we may observe that the preimage $R^{-1}(\mathrm{Stab}_{\mathbb{G}_{n,d}}(\Lambda))$, where $\mathrm{Stab}_{\mathbb{G}_{n,d}}(\Lambda)$ denotes the $\mathbb{G}_{n,d}$ -stabilizer of $\Lambda \in X_{n,d}$, is isomorphic to the reduced group $\mathrm{Aut}(M_n^k(\Lambda))/H_0$.

We summarize all the above in the following.

Theorem 5. *Let $d \geq 2$, $k \geq 2$ and $n \geq d + 2$ such that $(d; k, n) \notin \{(2; 2, 5), (2; 4, 3)\}$. Keeping with the above notation, we have the following.*

- (1) If $\Lambda, \widehat{\Lambda} \in X_{n,d}$, then $M_n^k(\Lambda)$ and $M_n^k(\widehat{\Lambda})$ are isomorphic if and only if the points Λ and $\widehat{\Lambda}$ are in the same \mathbb{G}_n -orbit
- (2) If $\Lambda \in X_{n,d}$, then $\text{Aut}(M_n^k(\Lambda))/H_0$ is isomorphic to an extension of the kernel of R (which is trivial if $(n, d) \neq (3, 1)$) by the $\mathbb{G}_{n,d}$ -stabilizer of Λ .
- (3) A model of the (coarse) moduli space of generalized Fermat manifolds of type $(d; k, n)$ is given by the complex orbifold $\overline{X}_{n,d} = X_{n,d}/\mathbb{G}_{n,d}$. The orbifold points are given by the equivalence classes of those points with non-trivial $\mathbb{G}_{n,d}$ -stabilizer.

3.4. A remark on the field of moduli. If $Z \subset \mathbb{P}^n$ be an algebraic variety, say defined as the zero of the homogeneous polynomials P_1, \dots, P_r . For each $\sigma \in \text{Gal}(\mathbb{C})$ (the group of field automorphisms of the complex numbers), let Z^σ be the algebraic variety defined by the polynomials $P_1^\sigma, \dots, P_r^\sigma$ (where P_j^σ is obtained from P_j by applying σ to its coefficients). Associated to Z is the group $G_Z = \{\sigma \in \text{Gal}(\mathbb{C}) : Z^\sigma \cong Z\}$. The field of moduli of Z is defined as the fixed subfield of \mathbb{C} under G_Z . The field of moduli was introduced by Shimura in [28] and it coincides with the intersection of all the definition fields of Z [21], but in general, it is not a field of definition. As a consequence of results due to Weil [34], if $\text{Aut}(Z)$ is trivial, then the field of moduli is a field of definition.

As M_{d+1}^k is just the classical Fermat hypersurface, its field of moduli is \mathbb{Q} , which is a field of definition. Now, assume $n \geq d + 2$ and $k \geq 2$ such that $(d; k, n) \notin \{(2; 2, 5), (2; 4, 3)\}$. Let $\Lambda \in X_{n,d}$. If $\sigma \in \text{Gal}(\mathbb{C})$, then $\Lambda^\sigma \in X_{n,d}$ and $M_n^k(\Lambda)^\sigma = M_n^k(\Lambda^\sigma)$. In particular, $G_\Lambda = \{\sigma \in \text{Gal}(\mathbb{C}) : M_n^k(\Lambda)^\sigma \cong M_n^k(\Lambda)\} = \{\sigma \in \text{Gal}(\mathbb{C}) : \Lambda^\sigma \equiv_{\mathbb{G}_{n,d}} \Lambda\}$ and the field of moduli of $M_n^k(\Lambda)$ is $\mathbb{Q}(I(\Lambda))$, where $I(\Lambda)$ is a (finite) set of invariants for the action of $\mathbb{G}_{n,d}$ on $X_{n,d}$.

In a forthcoming paper, we will deal with the explicit computation of the field of moduli, to describe conditions for it to be a field of definition and, if that is not the case, to obtain some (sharp) upper bound for the degree extension of a field of definition over the field of moduli (for $d = 1$ it is known that the field of moduli of a generalized Fermat curve is either a field of definition or there is a field of definition which is a quadratic extension of it [14]).

4. EXAMPLES

4.1. Jacobians of genus two surfaces. In [18, 29] it was observed that the desingularized Kummer surface of the Jacobian variety JC of the genus two hyperelliptic curve $C : y^2 = (x - \alpha_1) \cdots (x - \alpha_6)$, where $\alpha_1, \dots, \alpha_6 \in \mathbb{C}$ are different points, is isomorphic to the following complete intersection

$$S = \left\{ \begin{array}{l} x_1^2 + \cdots + x_6^2 = 0 \\ \alpha_1 x_1^2 + \cdots + \alpha_6 x_6^2 = 0 \\ \alpha_1^2 x_1^2 + \cdots + \alpha_6^2 x_6^2 = 0 \end{array} \right\} \subset \mathbb{P}^5.$$

The surface S admits the group $H \cong \mathbb{Z}_2^5$ of automorphisms (generated by those elements that multiply each coordinate by -1) and the quotient S/H is \mathbb{P}^2 branched at six lines in general position. This, in particular, asserts that S is a generalized Fermat surface of type $(2; 2, 5)$. Below, we proceed to find an equation of it in the form $S_5^2(\lambda_1, \lambda_2)$. If we set

$t_1 = (\alpha_1 - \alpha_5)(\alpha_1 - \alpha_6)x_1^2, t_2 = (\alpha_2 - \alpha_5)(\alpha_2 - \alpha_6)x_2^2, t_3 = (\alpha_3 - \alpha_5)(\alpha_3 - \alpha_6)x_3^2$, then $P : S \rightarrow \mathbb{P}^2 : [x_1 : \cdots : x_6] \mapsto [t_1 : t_2 : t_3]$ defines a Galois branched covering with deck group H . In this case, the six branched lines are

$$L_1 = \{t_1 = 0\}, L_2 = \{t_2 = 0\}, L_3 = \{t_3 = 0\}, L_4 = \{t_1 + t_2 + t_3 = 0\},$$

$$L_5 = \left\{ \frac{(\alpha_1 - \alpha_4)(\alpha_3 - \alpha_5)}{(\alpha_1 - \alpha_5)(\alpha_3 - \alpha_4)} t_1 + \frac{(\alpha_2 - \alpha_4)(\alpha_3 - \alpha_5)}{(\alpha_2 - \alpha_5)(\alpha_3 - \alpha_4)} t_2 + t_3 = 0 \right\},$$

$$L_6 = \left\{ \frac{(\alpha_1 - \alpha_4)(\alpha_3 - \alpha_6)}{(\alpha_1 - \alpha_6)(\alpha_3 - \alpha_4)} t_1 + \frac{(\alpha_2 - \alpha_4)(\alpha_3 - \alpha_6)}{(\alpha_2 - \alpha_6)(\alpha_3 - \alpha_4)} t_2 + t_3 = 0 \right\}.$$

It follows that $S \cong S_5^2(\lambda_1, \lambda_2)$, where

$$\lambda_1 = \left(\frac{(\alpha_1 - \alpha_4)(\alpha_3 - \alpha_5)}{(\alpha_1 - \alpha_5)(\alpha_3 - \alpha_4)}, \frac{(\alpha_1 - \alpha_4)(\alpha_3 - \alpha_6)}{(\alpha_1 - \alpha_6)(\alpha_3 - \alpha_4)} \right),$$

$$\lambda_2 = \left(\frac{(\alpha_2 - \alpha_4)(\alpha_3 - \alpha_5)}{(\alpha_2 - \alpha_5)(\alpha_3 - \alpha_4)}, \frac{(\alpha_2 - \alpha_4)(\alpha_3 - \alpha_6)}{(\alpha_2 - \alpha_6)(\alpha_3 - \alpha_4)} \right).$$

4.2. A connection between generalized Fermat surfaces and curves of the same type.

Let $(\lambda, \mu) \in X_{n,2}$, where $n \geq 4$, $\lambda = (\lambda_1, \dots, \lambda_{n-3})$, $\mu = (\mu_1, \dots, \mu_{n-3}) \in \mathbb{C}^{n-3}$. Associated with this pair, is the generalized Fermat surface $S_n^k(\lambda, \mu)$, $k \geq 2$, together its (unique) generalized Fermat group $H = H_0 \cong \mathbb{Z}_k^n$ and the Galois branched holomorphic map $\pi : S_n^k(\lambda, \mu) \rightarrow \mathbb{P}^2$ where $\pi[x_1 : \dots : x_{n+1}] = [t_1 = x_1^k : t_2 = x_2^k : t_3 = x_3^k]$ (whose deck group is H). Its branch locus set is given by the union of the lines $L_j(\lambda, \mu)$, $j = 1, \dots, n+1$, as previously defined.

Let us consider a line $L = \{\rho_1 t_1 + \rho_2 t_2 + \rho_3 t_3 = 0\} \subset \mathbb{P}^2$, $[\rho_1 : \rho_2 : \rho_3] \in \mathbb{P}^2$, such that the collection of $n+2$ lines $L_1(\lambda, \mu), \dots, L_{n+1}(\lambda, \mu), L$ are in general position. On L we have exactly $n+1$ intersection points with the branched lines $L_1(\lambda, \mu), \dots, L_{n+1}(\lambda, \mu)$. These points are given by

$$p_1 = [0 : 1], \quad p_2 = [1 : 0], \quad p_3 = \left[\frac{-\rho_2}{\rho_1} : 1 \right], \quad p_4 = \left[\frac{\rho_3 - \rho_2}{\rho_1 - \rho_3} : 1 \right],$$

$$p_j = \left[\frac{\mu_{j-4}\rho_3 - \rho_2}{\rho_1 - \lambda_{j-4}\rho_3} : 1 \right], \quad j = 5, \dots, n+1.$$

The preimage $\pi^{-1}(L) \subset S_n^k(\lambda, \mu)$ provides the following curve

$$\pi^{-1}(L) := \left\{ \begin{array}{l} \rho_1 x_1^k + \rho_2 x_2^k + \rho_3 x_3^k = 0 \\ x_1^k + x_2^k + x_3^k + x_4^k = 0 \\ \lambda_1 x_1^k + \mu_1 x_2^k + x_3^k + x_4^k = 0 \\ \vdots \\ \lambda_{n-3} x_1^k + \mu_{n-3} x_2^k + x_{n+1}^k = 0 \end{array} \right\} \subset \mathbb{P}^n.$$

By taking

$$y_1^k = \rho_1 x_1^k, \quad y_2^k = \rho_2 x_2^k, \quad y_3^k = \rho_3 x_3^k, \quad y_4^k = \frac{\rho_2 \rho_3}{\rho_3 - \rho_2} x_4^k,$$

$$y_j^k = \frac{\rho_2 \rho_3}{\mu_{4-j}\rho_3 - \rho_2} x_j^k, \quad j = 5, \dots, n+1,$$

we observe that the above is isomorphic to the following generalized Fermat curve of type (k, n)

$$C_n^k(\eta_{L;\lambda,\mu}) = \left\{ \begin{array}{l} y_1^k + y_2^k + y_3^k = 0 \\ \eta_1 y_1^k + y_2^k + y_4^k = 0 \\ \vdots \\ \eta_{n-2} y_1^k + y_2^k + y_{n+1}^k = 0 \end{array} \right\} \subset \mathbb{P}^n,$$

where $\eta_{L;\lambda,\mu} = (\eta_1, \dots, \eta_{n-2})$ and

$$\eta_1 := \frac{\rho_2(\rho_3 - \rho_1)}{\rho_1(\rho_3 - \rho_2)},$$

$$\eta_2 := \frac{\rho_2(\lambda_1\rho_3 - \rho_1)}{\rho_1(\mu_1\rho_3 - \rho_2)}, \dots, \eta_{n-2} := \frac{\rho_2(\lambda_{n-3}\rho_3 - \rho_1)}{\rho_1(\mu_{n-3}\rho_3 - \rho_2)}.$$

By varying L (and including the non-general ones as above), we obtain a two-dimensional family of such generalized Fermat curves for the fixed parameter (λ, μ) . This process provides the 3-fold

$$X(\lambda, \mu) \subset \mathbb{P}^2 \times \mathbb{P}^n$$

defined by those tuples $([\rho_1 : \rho_2 : \rho_3], [x_1 : \dots : x_{n+1}]) \in \mathbb{P}^2 \times \mathbb{P}^n$ satisfying the following equations

$$\left\{ \begin{array}{l} y_1^k + y_2^k + y_3^k = 0 \\ \rho_2(\rho_3 - \rho_1)y_1^k + \rho_1(\rho_3 - \rho_2)y_2^k + \rho_1(\rho_3 - \rho_1)y_3^k = 0 \\ \rho_2(\lambda_1\rho_3 - \rho_1)y_1^k + \rho_1(\mu_1\rho_3 - \rho_2)y_2^k + \rho_1(\mu_1\rho_3 - \rho_2)y_3^k = 0 \\ \vdots \\ \rho_2(\lambda_{n-3}\rho_3 - \rho_1)y_1^k + \rho_1(\mu_{n-3}\rho_3 - \rho_2)y_2^k + \rho_1(\mu_{n-3}\rho_3 - \rho_2)y_{n+1}^k = 0 \end{array} \right\},$$

together the morphism

$$\pi_{\lambda,\mu} : X(\lambda, \mu) \rightarrow \mathbb{P}^2 : ([\rho_1 : \rho_2 : \rho_3], [y_1 : \dots : y_{n+1}]) \mapsto [\rho_1 : \rho_2 : \rho_3],$$

such that $C_n^k(\eta_{L;\lambda,\mu}) \cong \pi_{\lambda,\mu}^{-1}([\rho_1 : \rho_2 : \rho_3])$.

Remark 9. If $\rho := [\rho_1 : \rho_2 : \rho_3] \in \mathbb{P}^2$ corresponds to a line L such that the collection of lines $L, L_1(\lambda, \mu), \dots, L_{n+1}(\lambda, \mu)$ is not in general position, then $C_n^k(\eta_{L;\lambda,\mu}) \cong \pi_{\lambda,\mu}^{-1}(\rho) \subset X(\lambda, \mu)$ is a singular curve (a singular generalized Fermat curve of type (k, n)).

The (possible singular) generalized Fermat curve $\pi_{\lambda,\mu}^{-1}(\rho)$ is invariant under the group $H \cong \mathbb{Z}_k^n = \langle b_1, \dots, b_n \rangle$ of automorphisms of $X(\lambda, \mu)$, where b_j is amplification of y_j by $e^{2\pi i/k}$, and such a restriction to $\pi_{\lambda,\mu}^{-1}(\rho)$ is its generalized Fermat group of type (k, n) .

Remark 10. The above provides a nice relation between the surface $S_n^k(\lambda, \mu)$ and the 3-fold $X(\lambda, \mu)$. The last one parametrizes all the (singular) generalized Fermat curves of type (k, n) inside the surface, each one being invariant under the generalized Fermat group H of $S_n^k(\lambda, \mu)$.

4.3. Generalized Fermat surfaces and smooth quadrics. In this example, let us consider a (smooth) quadric $Q \subset \mathbb{P}^2$ which is tangent to the lines $L_1 = L_{[1:0:0]}$, $L_2 = L_{[0:1:0]}$, $L_3 = L_{[0:0:1]}$ and $L_4 = L_{[1:1:1]}$.

Following the same notations as above, we proceed to describe a one-to-one correspondence between the collection of generalized Fermat curves of type (k, n) and that of the generalized Fermat surfaces of type $(2; k, n)$ whose associated lines $L_j(\lambda, \mu)$ are each one tangent to Q .

Lemma 4. *The family of (smooth) quadrics in \mathbb{P}^2 being tangent to the lines L_1, L_2, L_3 and L_4 is provided by the following ones:*

$$Q_\alpha := \{4t_1^2 + \alpha^2 t_2^2 + (4 + \alpha^2)t_3^2 + 4\alpha t_1 t_2 + 4(2 - \alpha)t_1 t_3 - 2\alpha(2 - \alpha)t_2 t_3\},$$

where $\alpha \notin \{0, 2\}$.

Proof. The family all quadrics in \mathbb{P}^2 is given by

$$Q = \{\alpha_1 t_1^2 + \alpha_2 t_2^2 + \alpha_3 t_3^2 + \alpha_4 t_1 t_2 + \alpha_5 t_1 t_3 + \alpha_6 t_2 t_3 = 0\}.$$

The condition for Q to be tangent to the given lines asserts that

$$\alpha_1 \alpha_2 \alpha_3 (\alpha_2 + \alpha_3 - \alpha_6) (\alpha_1 + \alpha_3 - \alpha_5) (\alpha_1 + \alpha_2 - \alpha_4) \neq 0.$$

Tangency of Q at these lines obligates that the point in $L_i \cap L_j$, where $i \neq j$, cannot belong to Q . These points are given by $[1 : 0 : 0]$, $[0 : 1 : 0]$, $[0 : 0 : 1]$, $[0 : 1 : -1]$, $[1 : 0 : -1]$ and $[1 : -1 : 0]$. Now, $[1 : 0 : 0] \in Q$ if and only if $\alpha_1 = 0$, $[0 : 1 : 0] \in Q$ if and only if $\alpha_2 = 0$, $[0 : 0 : 1] \in Q$ if and only if $\alpha_3 = 0$, $[0 : 1 : -1] \in Q$ if and only if $\alpha_2 + \alpha_3 - \alpha_6 = 0$, $[1 : 0 : -1] \in Q$ if and only if $\alpha_1 + \alpha_3 - \alpha_5 = 0$, and $[1 : -1 : 0] \in Q$ if and only if $\alpha_1 + \alpha_2 - \alpha_4 = 0$.

In this way, we may assume that $\alpha_1 = 1$. The intersection locus $Q \cap L_1$ is given by the points $[0 : t_2 : t_3]$ such that $\alpha_2 t_2^2 + \alpha_3 t_3^2 + \alpha_6 t_2 t_3 = 0$. As $\alpha_2 \neq 0$, then $t_3 \neq 0$, so we may assume $t_3 = 1$. In this case, the above equation is $\alpha_2 t_2^2 + \alpha_3 + \alpha_6 t_2 = 0$ and the tangency condition is equivalent to $\alpha_6^2 = 4\alpha_2 \alpha_3$. Working in a similar fashion for L_2 and L_3 , respectively, provide the equalities $\alpha_5^2 = 4\alpha_3$ and $\alpha_4^2 = 4\alpha_2$. In particular, the above equalities asserts that $\alpha_4 \alpha_5 \neq 0$ and, as $4\alpha_6^2 = 16\alpha_2 \alpha_3 = (4\alpha_2)(4\alpha_3) = \alpha_4^2 \alpha_5^2$, that $2\alpha_6 = \pm \alpha_4 \alpha_5$.

The tangency condition of Q and L_4 provides the equality

$$(*) \quad -4(1 + \alpha_3 - \alpha_5)(\alpha_2 + \alpha_3 - \alpha_6) + (-2\alpha_3 - \alpha_4 + \alpha_5 + \alpha_6)^2 = 0.$$

By replacing α_2 by $\alpha_4^2/4$ and α_3 by $\alpha_5^2/4$ in $(*)$ we obtain the equality

$$(**) \quad (\alpha_4 \alpha_5 - 2\alpha_6)(8 - 4\alpha_4 - 4\alpha_5 + \alpha_4 \alpha_5 + 2\alpha_6) = 0.$$

If $2\alpha_6 = \alpha_4 \alpha_5$, then equality $(**)$ holds. But in this case, Q corresponds to the equation

$$4t_1^2 + \alpha^2 t_2^2 + \beta^2 t_3^2 + 4\alpha t_1 t_2 + 4\beta t_1 t_3 + 2\alpha\beta t_2 t_3 = 0,$$

which is equivalent to

$$(2t_1 + \alpha_4 t_2 + \alpha_5 t_3)^2 = 0,$$

which is not a smooth conic.

If $2\alpha_6 = -\alpha_4 \alpha_5$, then equality $(**)$ is given as $\alpha_4 \alpha_5 (\alpha_4 + \alpha_5 - 2) = 0$, that is, $\alpha_5 = 2 - \alpha_4$. In this case, the conic Q is given by the equation

$$4t_1^2 + \alpha_4^2 t_2^2 + (4 + \alpha_4^2) t_3^2 + 4\alpha_4 t_1 t_2 + 4(2 - \alpha_4) t_1 t_3 - 2\alpha_4(2 - \alpha_4) t_2 t_3 = 0.$$

□

Let us fix any $\alpha \notin \{0, 2\}$ and let $\mathscr{W}_\alpha \subset X_{n,2}$ be the collection of those tuples (λ, μ) such that all the lines $L_j(\lambda, \mu)$, for $5 \leq j \leq n+1$, are tangent to the quadric Q_α . The condition for the tangency provides a polynomial $E_\alpha \in \mathbb{C}[x, y]$ such that $E_\alpha(\lambda_{j-4}, \mu_{j-4}) = 0$, for all values of j . (So \mathscr{W}_α is just the locus of zeroes of a set of $n-3$ polynomials).

Each $(\lambda, \mu) \in \mathscr{W}_\alpha$ determines $n+1$ points on Q_α , these being the tangency points of it with the $n+1$ lines $L_j(\lambda, \mu)$, $j = 1, \dots, n+1$. By identifying Q_α with $\widehat{\mathbb{C}}$, these $n+1$ tangency points can be identified with $n+1$ different points on $\widehat{\mathbb{C}}$, which, up to a Möbius transformation, can be assumed to be $\infty, 0, 1, \eta_1 := \eta_1(\alpha; \lambda, \mu), \dots, \eta_{n-2} := \eta_{n-2}(\alpha; \lambda, \mu)$. These points produce a a generalized Fermat curve of type (k, n) , with

equations

$$C_n^k(\eta_{\alpha}; \lambda, \mu) = \left\{ \begin{array}{l} y_1^k + y_2^k + y_3^k = 0 \\ \eta_1 y_1^k + y_2^k + y_4^k = 0 \\ \vdots \\ \eta_{n-2} y_1^k + y_2^k + y_{n+1}^k = 0 \end{array} \right\} \subset \mathbb{P}^n,$$

where $\eta_{\alpha}; \lambda, \mu = (\eta_1, \dots, \eta_{n-2})$.

Remark 11. Conversely, given $n + 1$ different points on Q_{α} , where four of them are the intersections of it with the lines $L_j = L_j(\lambda, \mu)$, for $j = 1, 2, 3, 4$, we may obtain a tuple $(\lambda, \mu) \in \mathcal{W}_{\alpha}$ such that the corresponding tangencies are these given points. This means that the above process can be reversed.

In this way, we obtain a one-to-one correspondence between generalized Fermat curves of type (k, n) and generalized Fermat surfaces of type $(2; k, n)$ parametrized by the tuples in \mathcal{W}_{α} .

Remark 12. Let $\alpha \notin \{0, 2\}$ and $(\lambda, \mu) \in \mathcal{W}_{\alpha}$. Note that there is a two-fold branched cover $P_{\alpha} : \mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^2$, whose deck group is $\langle J(x, y) = (y, x) \rangle$ and with branch locus $Q_{\alpha} = P_{\alpha}(\Delta)$, where $\Delta = \{(x, x) \in \mathbb{P}^1 \times \mathbb{P}^1\} \cong \mathbb{P}^1$. Under P_{α} , the tangency point of Q_{α} with $L_j(\lambda, \mu)$ lifts to a point $(q_j, q_j) \in \Delta$ and $L_j(\lambda, \mu)$ lifts to the lines $L_j^+(\alpha; \lambda, \mu) = \{q_1\} \times \mathbb{P}^1$ and $L_j^-(\alpha; \lambda, \mu) = \mathbb{P}^1 \times \{q_1\}$. The points $q_1, \dots, q_{n+1} \in \mathbb{P}^1$ determines a generalized Fermat curve of type (k, n) , isomorphic to $C_n^k(\eta_{\alpha}; \lambda, \mu)$. Now, we may consider the complex surface $W = C_n^k(\eta_{\alpha}; \lambda, \mu) \times C_n^k(\eta_{\alpha}; \lambda, \mu) \subset \mathbb{P}^n \times \mathbb{P}^n$. On this surface, we have the action of the group $H \times H \cong \mathbb{Z}_k^n \times \mathbb{Z}_k^n$, where each factor H acts on the corresponding factor as its (unique) generalized Fermat group. We also have the action of $\mathbb{Z}_2 = \langle \tilde{J}(x, y) = (y, x) \rangle$, which permutes both factors in $H \times H$. The locus of fixed points of \tilde{J} is the diagonal, this being a copy of $C_n^k(\eta_{\alpha}; \lambda, \mu)$. The quotient $Z = W / \langle \tilde{J} \rangle$ turns out to be a surface (this situation is similar as for the quotient $(\mathbb{P}^1 \times \mathbb{P}^1) / \langle J \rangle = \mathbb{P}^2$) admitting the action of H such that Z/H is \mathbb{P}^2 and its branch locus (by such an action) being the lines $L_j(\lambda, \mu)$, $j = 1, \dots, n + 1$. In this way, we may observe that Z is a copy of $S_n^k(\lambda, \mu)$ in this case. In the particular case $k = 2$ and n odd, this was observed in [25].

5. APPENDIX: ON THE HOMOMORPHISM $R : \mathfrak{S}_{n+1} \rightarrow \mathbb{G}_{n,d}$

Let $\Lambda \in X_{n,d}$. If $\hat{\Lambda} \in X_{n,d}$ is, up to permutation, equivalent to Λ . Then (by the definition) there exists $T \in \text{PGL}_{d+1}(\mathbb{C})$ and there is a permutation $\eta \in \mathfrak{S}_{n+1}$ of the set $\{1, \dots, n + 1\}$ defined by $T(L_j(\Lambda)) = L_{\eta(j)}(\hat{\Lambda})$. Conversely, given $\eta \in \mathfrak{S}_{n+1}$, consider the reordered collection $(L_{\eta^{-1}(1)}(\Lambda), \dots, L_{\eta^{-1}(n+1)}(\Lambda))$. There is a unique $\hat{\Lambda} \in X_{n,d}$ such that there exists $T_{\eta} \in \text{PGL}_{d+1}(\mathbb{C})$ satisfying that $T_{\eta}(L_{\eta^{-1}(j)}(\Lambda)) = L_j(\hat{\Lambda})$. It follows that $\hat{\Lambda} \in X_{n,d}$ is, up to permutations, equivalent to Λ . In particular, the above asserts that, for every $\eta \in \mathfrak{S}_{n+1}$, there is a (unique) bijection $R_{\eta} : X_{n,d} \rightarrow X_{n,d}$ such that $R_{\eta}(\Lambda)$ and $\hat{\Lambda}$ are standard parameters for the same divisor.

If $\text{Sym}(X_{n,d})$ denotes the group of bijective maps of $X_{n,d}$, then the above induces a function $R : \mathfrak{S}_{n+1} \rightarrow \text{Sym}(X_{n,d})$, defined by $R(\eta) = R_{\eta}$. As \mathfrak{S}_{n+1} is generated by the permutations $\sigma_1 = (1, 2)$ and $\sigma_2 = (1, 2, \dots, n, n + 1)$, we only need to describe R_{σ_1} and R_{σ_2} . These are given as follows:

$$R_{\sigma_1}(\lambda_1, \dots, \lambda_d) = (\lambda_2, \lambda_1, \lambda_3, \dots, \lambda_d), \quad R_{\sigma_2}(\lambda_1, \dots, \lambda_d) = (\hat{\lambda}_1, \dots, \hat{\lambda}_d),$$

where

$$\begin{aligned}\widehat{\lambda}_{1,1} &= \frac{\lambda_{n-d-1,d}}{\lambda_{n-d-1,d}-1}, \\ \widehat{\lambda}_{1,j} &= \frac{\lambda_{n-d-1,d}(\lambda_{n-d-1,j-1}-1)}{\lambda_{n-d-1,j-1}(\lambda_{n-d-1,d}-1)}, \quad j = 2, \dots, d, \\ \widehat{\lambda}_{i+1,1} &= \frac{\lambda_{n-d-1,d}}{\lambda_{n-d-1,d}-\lambda_{i,d}}, \quad i = 1, \dots, n-d-2, \\ \widehat{\lambda}_{i+1,j} &= \frac{\lambda_{n-d-1,d}(\lambda_{n-d-1,j-1}-\lambda_{i,j-1})}{\lambda_{n-d-1,j-1}(\lambda_{n-d-1,d}-\lambda_{i,d})}, \quad j = 2, \dots, d, \quad i = 1, \dots, n-d-2.\end{aligned}$$

The above explicit forms for R_{σ_1} and R_{σ_2} permits to see that these two are holomorphic automorphisms of $X_{n,d}$, which generate a group $\mathbb{G}_{n,d}$. Now, using the fact that R is a homomorphism of groups, it follows that for every $\eta \in \mathfrak{S}_{n+1}$ one has that $R_\eta \in \text{Aut}(X_{n,d})$. In this way, we obtain a homomorphism (we consider multiplication at the left in the permutation group)

$$R : \mathfrak{S}_{n+1} \rightarrow \mathbb{G}_{n,d} \leq \text{Aut}(X_{n,d}) : \eta \mapsto R_\eta.$$

That R is a homomorphism of groups can be seen as follows. Each hyperplane L of \mathbb{P}^d corresponds to a point $Q = [q_1 : \dots : q_{d+1}] \in \mathbb{P}^d$; the equation of L is given by $\sum_{j=1}^{g+1} q_j x_j = 0$. This identification is represented by the horizontal arrows in the following commutative diagrams.

Let $\eta, \rho \in \mathfrak{S}_{n+1}$ and $\Lambda \in X_{n,d}$. We identify the hyperplane $L_j(\Lambda)$ with the point $Q_j(\Lambda)$ as above. We then have the following two commutative diagrams. To simplify the notation, in the diagrams the symbol f_\bullet represents the $n+1$ -tuple $(f_1, f_2, \dots, f_{n+1})$. This is to say replace \bullet by the numbers $1, 2, \dots, n+1$ to obtain the respective tuple.

$$\begin{array}{ccc} L_\bullet(\Lambda) & \longrightarrow & Q_\bullet(\Lambda) \\ \downarrow \eta^{-1} & & \downarrow \eta^{-1} \\ L_{\eta^{-1}(\bullet)}(\Lambda) & \longrightarrow & Q_{\eta^{-1}(\bullet)}(\Lambda) \\ \downarrow T_\eta \in \text{PGL}_{d+1}(\mathbb{C}) & & \downarrow T_\eta \in \text{PGL}_{d+1}(\mathbb{C}) \\ T_\eta(L_{\eta^{-1}(\bullet)}(\Lambda)) & \longrightarrow & Q_{\eta^{-1}(\bullet)}(\Lambda)T_\eta^{-1} \\ \downarrow = & & \downarrow = \\ L_\bullet(R_\eta(\Lambda)) & \longrightarrow & Q_\bullet(R_\eta(\Lambda)) \\ \downarrow \rho^{-1} & & \downarrow \rho^{-1} \\ L_{\rho^{-1}(\bullet)}(R_\eta(\Lambda)) & \longrightarrow & Q_{\rho^{-1}(\bullet)}(R_\eta(\Lambda)) \\ \downarrow T_\rho \in \text{PGL}_{d+1}(\mathbb{C}) & & \downarrow T_\rho \in \text{PGL}_{d+1}(\mathbb{C}) \\ T_\rho(L_{\rho^{-1}(\bullet)}(R_\eta(\Lambda))) & \longrightarrow & Q_{\rho^{-1}(\bullet)}(R_\eta(\Lambda))T_\rho^{-1} \\ \downarrow = & & \downarrow = \\ L_\bullet(\Lambda^*) & \longrightarrow & Q_\bullet(\Lambda^*) \end{array}$$

and

$$\begin{array}{ccc}
L_{\bullet}(\Lambda) & \longrightarrow & Q_{\bullet}(\Lambda) \\
\downarrow (\eta\rho)^{-1} & & \downarrow (\eta\rho)^{-1} \\
L_{(\eta\rho)^{-1}(\bullet)}(\Lambda) & \longrightarrow & Q_{(\eta\rho)^{-1}(\bullet)}(\Lambda) \\
\downarrow T_{\eta\rho} \in \mathrm{PGL}_{d+1}(\mathbb{C}) & & \downarrow T_{\eta\rho} \in \mathrm{PGL}_{d+1}(\mathbb{C}) \\
T_{\eta\rho}(L_{(\eta\rho)^{-1}(\bullet)}(\Lambda)) & \longrightarrow & Q_{(\eta\rho)^{-1}(\bullet)}(\Lambda)T_{\eta\rho}^{-1} \\
\downarrow = & & \downarrow = \\
L_{\bullet}(R_{\eta\rho}(\Lambda)) & \longrightarrow & Q_{\bullet}(R_{\eta\rho}(\Lambda))
\end{array}$$

Now, from the first commutative diagram, we notice that the equalities

$$Q_j(\Lambda^*) = Q_{\rho^{-1}(j)}(R_{\eta}(\Lambda))T_{\rho}^{-1} = Q_{\rho^{-1}(\eta^{-1}(j))}(\Lambda)T_{\eta}^{-1}T_{\rho}^{-1} = Q_{(\eta\rho)^{-1}(j)}(\Lambda)T_{\eta}^{-1}T_{\rho}^{-1}$$

and from the second commutative diagram

$$Q_j(R_{\eta\rho}(\Lambda)) = Q_{(\eta\rho)^{-1}(j)}(\Lambda)T_{\eta\rho}^{-1}.$$

It can be checked, by direct inspection of the definitions of the linear transformations $T_{\rho}, T_{\eta}, T_{\eta\rho}$ that $T_{\eta\rho} = T_{\rho}T_{\eta}$. So, the above asserts that

$$Q_j(R_{\eta\rho}(\Lambda)) = Q_j(\Lambda^*)$$

from which it can be seen that $R_{\eta\rho}(\Lambda) = \Lambda^* = R_{\rho}R_{\eta}(\Lambda)$.

Example 3. If $d = 1$ and $n = 3$, then the kernel of R is given by the Klein group $\{e, (1, 2)(3, 4), (1, 3)(2, 4), (1, 4)(2, 3)\} \cong \mathbb{Z}_2^2$ (where e denotes the identity permutation). In this case, $\mathbb{G}_{3,1} \cong \mathfrak{S}_3$. Also, for $n \geq 4$, it can be seen that $\mathbb{G}_{n,1} \cong \mathfrak{S}_{n+1}$.

6. APPENDIX: ALGEBRAIC MODELS OF GENERALIZED FERMAT MANIFOLDS

As mentioned before, Pardini's building data permits to obtain algebraic models of a generalized Fermat manifold, given as a complete intersection of Fermat hypersurfaces. This section provides another argument (based on basic covering theory) to obtain such models. Let us observe that if M is a compact d -dimensional complex manifold and $K < \mathrm{Aut}(M)$ is a finite group, then its locus of fixed points $\mathrm{Fix}(K) = \{p \in M : a(p) = p, \forall a \in K\}$ is a closed submanifold of M (it might not be connected and, if it is not connected, then its connected components might have different dimensions) [24, Corollary 4.6].

Remark 13. Let $H < \mathrm{Aut}(M)$ be a finite group. Assume that there is a branched Galois holomorphic covering $\pi : M \rightarrow \mathbb{P}^d$, with deck group H , and whose branch locus D consists of the union of $n + 1$ hyperplanes $L_1, \dots, L_{n+1} \subset \mathbb{P}^d$, where $n \geq 2$, which are in general position (in other words, D is a strict normal crossing divisor). If $W = \mathbb{P}^d \setminus (L_1 \cup \dots \cup L_{n+1})$, then $\pi : \pi^{-1}(W) \rightarrow W$ is a Galois (unbranched) covering map, whose deck group still H . If $P : \widetilde{W} \rightarrow W$ is a universal covering, then there is a Galois covering $Q : \widetilde{W} \rightarrow \pi^{-1}(W)$ such that $P = \pi \circ Q$. In [35] (for $d = 2$) and [26, 27] (for $d \geq 3$) (see also [1]) it is noted that $\pi_1(W) \cong \mathbb{Z}^n$. In particular, for $d \geq 2$, H must be an abelian group, a quotient of \mathbb{Z}^n . Let us assume $H \cong \mathbb{Z}_k^n$, for some integer $k \geq 2$. If for every $j \in \{1, \dots, n + 1\}$ and every connected component F of $\pi^{-1}(L_j)$ (which is a submanifold of dimension $d - 1$), the isotropy group $H_F = \{a \in H : F \subset \mathrm{Fix}(a)\} \cong \mathbb{Z}_k$, then H a generalized Fermat group of type $(d; k, n)$. Note that $F \subset \mathrm{Fix}(H_F)$, but under the assumption that $n \geq d + 1$, $F = \mathrm{Fix}(H_F)$ and, if moreover $d \geq 2$, that $\pi^{-1}(L) = F$.

6.1. The Riemann extension theorem. Let Ω be a domain in \mathbb{C}^m and $A \subset \Omega$. It is said that A is *thin* if for every $z \in \Omega$ there is an open polydisk $D(z; r) \subset \Omega$ and a non-zero holomorphic map $f : D(z; r) \rightarrow \mathbb{C}$ such that f is identically zero in $A \cap D(z; r)$. Let $h : \Omega \setminus A \rightarrow \mathbb{C}$ be a function. It is said that h is *locally bounded in Ω* if for every $z \in \Omega$ there exists an open polydisc $D(z; r) \subset \Omega$ such that h is bounded in $D(z; r) \cap (\Omega \setminus A)$.

Theorem 6 (Riemann extension theorem). [11] *Let Ω be a domain in \mathbb{C}^m and let $A \subset \Omega$ be a thin subset. Let $f : \Omega \setminus A \rightarrow \mathbb{C}$ be a locally bounded holomorphic map on Ω . Then there is a (unique) holomorphic map $F : \Omega \rightarrow \mathbb{C}$ such that $F(z) = f(z)$ for every $z \in \Omega \setminus A$.*

6.2. The Algebraic model. Let $D \subset \mathbb{P}^d$, $d \geq 1$, be a strict normal crossing divisor formed by a collection of $n \geq d + 1$ hyperplanes. Let $\mathcal{F}(k; D)$ be the collection of pairs (N, K) which satisfy the following properties:

- (1) N is a compact complex manifold of dimension d and $K < \text{Aut}(N)$ is an abelian finite group.
- (2) There is a Galois branched holomorphic covering $\pi_N : N \rightarrow \mathbb{P}^d$, with K as its deck group, whose branch locus is the divisor D .
- (3) Given one of the hyperplanes L which make up D and F any connected component of $\pi^{-1}(L)$, the subgroup $\{a \in K : F \subset \text{Fix}(a)\}$ is isomorphic to group \mathbb{Z}_k .

Let us consider two pairs $(N, K), (N', K') \in \mathcal{F}(k; D)$. By a morphism $\varphi : (N, K) \rightarrow (N', K')$ of these two pairs we mean a morphism of complex manifolds, $\varphi : N \rightarrow N'$ such that $\pi_N = \pi_{N'} \circ \varphi$. If φ is a biholomorphism, we will say that the pairs (N, K) and (N', K') are equivalent in $\mathcal{F}(k; D)$.

Remark 14. If (M, H) is a generalized Fermat pair of type $(d; k, n)$, where $d \geq 1$, $k \geq 2$ and $n \geq d + 1$, and $\pi : M \rightarrow \mathbb{P}^d$ is a Galois branched covering, with deck group is H , whose branch locus is given by the strict normal crossing divisor $D \subset \mathbb{P}^d$, then $(M, H) \in \mathcal{F}(k; D)$.

Fixing an order over the hyperplanes of D , let $\Lambda := (\lambda_1, \dots, \lambda_d) \in X_{n,d}$ be the standard parameter of the ordered divisor D . By composition, by a suitable linear projective automorphism of \mathbb{P}^d at the left of π , we may assume these branch locus to be given by the ordered divisor $(L_1(\Lambda), \dots, L_{n+1}(\Lambda))$ (by abuse of notation, we note by Λ this divisor). Let's remember that $\pi_0 : M_n^k(\Lambda) \rightarrow \mathbb{P}^d$ is the regular branched cover with deck group H_0 , whose branch set is Λ .

Theorem 7. *The pair $(M_n^k(\Lambda), H_0) \in \mathcal{F}(k; D)$ satisfies the following universal property: If $(N, K) \in \mathcal{F}(k; D)$, then there exists a unique morphism $\varphi : (M_n^k(\Lambda), H_0) \rightarrow (N, K)$. What's more, each generalized Fermat pair $(M, H) \in \mathcal{F}(k; D)$ is equivalent to $(M_n^k(\Lambda), H_0)$.*

Proof. The case $d = 1$ was proved in [10], so we only need to consider the case $d \geq 2$. The quotient orbifold $\mathcal{O} = M_n^k(\Lambda)/H_0$ can be identified with \mathbb{P}^d and whose branch locus consists of exactly $(n + 1)$ hyperplanes in general position, each one with branch order k . Let $\pi_1(\mathcal{O})$ be the orbifold fundamental group of \mathcal{O} . For generalities on orbifold fundamental groups and their coverings, the reader may consult the reference [6]. By Remark 4 $M_n^k(\Lambda)$ is simply connected. So $M_n^k(\Lambda)$ is the universal cover of \mathcal{O} and the orbifold fundamental group $\pi_1(\mathcal{O}) \cong \mathbb{Z}_k^n$. Let's remember that N is a compact complex manifold admitting an abelian group K of automorphisms such that $N/K = \mathcal{O}$. Then there exists a pair $(N', K') \in \mathcal{F}(k; \Lambda)$ which satisfies the following properties:

- (1) There exists a subgroup J in H_0 , acting freely on $M_n^k(\Lambda)$, such that $N' := M_n^k(\Lambda)/J$ and $K' := H/J$.

(2) There exists a homeomorphism $\varphi : N' \rightarrow N$ such that $\pi_N \circ \varphi = \varphi_{N'}$.

Outside of the branch locus Λ the holomorphisms $\pi_N, \pi_{N'}$ are locally invertible, i.e., φ is holomorphic in a dense open set Ω of N' and the closed set $N' \setminus \Omega$ is locally thin. It follows from Theorem 6 that φ is a biholomorphism. If (M, H) is generalized Fermat pairs, then $J = 1$. Which finishes the proof of the theorem. \square

Remark 15. In Proposition 2 we have provided the conditions for any element of H to act freely on M . This can be used to construct subgroups K of H acting freely on M and, in particular, those manifolds N and groups A as above.

REFERENCES

- [1] W. Arbola. The fundamental group of the complement of an arrangement of complex hyperplanes. *Topology* **31** (4) (1992), 757–765.
- [2] W. P. Barth, K. Hulek, C. A. M. Peters, and A. Van de Ven. *Compact complex surfaces*, Vol. 4 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, second edition, 2004.
- [3] Nguyen Bin. Some examples of algebraic surfaces with canonical map of degree 20. *Comptes Rendus* **359** (2021), 1145–1153.
- [4] G. Castelnuovo. Sulle serie algebriche di gruppi di punti appartenenti ad una curve algebrica. *Rend. R. Accad. Lincei Ser. (5)* **XV** (1906) (Memorie scelte p. 509).
- [5] C. Chevalley. Invariants of finite groups generated by reflections. *Amer. J. Math.* **77** (1955), 778–782.
- [6] Daryl Cooper, Craig D. Hodgson and Steven P. Kerckhoff. *Three-dimensional Orbifolds and Cone-Manifolds*. Mathematical Society of Japan Memoirs **5**, 2000.
- [7] D. S. Dummit and R. M. Foote. *Abstract Algebra*. Wiley (2003). ISBN 978-0-471-43334-7.
- [8] D. Eisenbud. *Commutative algebra. With a view toward algebraic geometry*. Springer-Verlag, Graduate texts in Mathematics **150**, 1995.
- [9] Yun Gao. A note on finite abelian covers. *Science China. Mathematics* **54** No. 7 (2011), 1333–1342. doi: 10.1007/s11425-011-4201-1
- [10] G. González-Diez, R. A. Hidalgo, and M. Leyton-Álvarez. Generalized Fermat curves. *J. Algebra* **321** (6) (2009), 1643–1660.
- [11] H. Grauert and R. Remmert. RIEMANN Extension Theorem and Analytic Coverings. In: *Coherent Analytic Sheaves. Grundlehren der mathematischen Wissenschaften (A Series of Comprehensive Studies in Mathematics)*, vol **265**. Springer, Berlin, Heidelberg (1984).
- [12] R. Hartshorne. Varieties of small codimension in projective space. *Bull. Amer. Math. Soc.*, 80:1017–1032, 1974.
- [13] R. Hartshorne. *Algebraic geometry*. Springer-Verlag, Graduate Texts in Mathematics **52** (1977).
- [14] R. A. Hidalgo. A remark on the field of moduli of Riemann surfaces. *Archiv der Mathematik* **114** (2020), 515–526.
- [15] R. A. Hidalgo, A. Kontogeorgis, M. Leyton-Álvarez and P. Paramantzoglou. Automorphisms of generalized Fermat curves. *Journal of Pure and Applied Algebra* **221** (2017), 2312–23337.
- [16] F. Hirzebruch. Arrangements of lines and algebraic surfaces. In: *Arithmetic and Geometry*, vol. II. Progr Math, 36. Birkhäuser, Boston: Mass, 1983, 113–140.
- [17] F. Hirzebruch. Some examples of algebraic surfaces. *Contemporary Mathematics* **9** (1982), 55–71.
- [18] V. F. Klein. Zur Theorie der Linencomplexe des ersten und zweiten Grades. *Math. Ann.* **2** (1870), 198–226.
- [19] Shoshichi Kobayashi. *Transformation groups in differential geometry*. Classics in Mathematics. Springer-Verlag, Berlin, 1995. Reprint of the 1972 edition.
- [20] K. Kodaira. On the structure of compact complex surfaces. I. *Amer. J. Math.* **86** (1964), 751–798.
- [21] S. Koizumi. The fields of moduli for polarized abelian varieties and for curves. *Nagoya Math. J.* **48** (1972), 37–55.
- [22] A. Kontogeorgis. Automorphisms of Fermat-like varieties. *Manuscripta Math.* **107** (2) (2002), 187–205.
- [23] R. Pardini. Abelian covers of algebraic variety. *J. Reine Angew. Math.* **417** (1991), 191–213.
- [24] Y. Prokhorov and C. Shramov. Automorphism groups of compact complex surfaces. [arXiv:1708.03566v5](https://arxiv.org/abs/1708.03566) [math.AG] 3 Jul 2019.
- [25] R. Quezada. Sobre superficies de Büchi generalizadas. Mathematical Master thesis, Universidad de Concepción, 2020.

- [26] R. Randell. The fundamental group of the complement of a union of complex hyperplanes. *Inventiones math.* **69** (1982), 103–108.
- [27] R. Randell. The fundamental group of the complement of a union of complex hyperplanes: correction. *Inventiones math.* **80** (1985), 467–468.
- [28] G. Shimura. On the field of rationality for an abelian variety. *Nagoya Math. J.* **45** (1972), 167–178.
- [29] T. Shioda. Some results on unirationality of algebraic surface. *Math. Ann.* **230** (1977), 153–168.
- [30] T. Shioda. Arithmetic and geometry of Fermat curves. In *Algebraic Geometry Seminar (Singapore, 1987)*, pages 95–102. World Sci. Publishing, Singapore, 1988.
- [31] T. Shioda and H. Inose. On singular K3 surfaces. In *Complex analysis and algebraic geometry*, pages 119–136. Iwanami Shoten, Tokyo, 1977.
- [32] T. Terasoma. Complete intersections of hypersurfaces. *Japan J. Math.* **14** No. 2 (1988), 309–384.
- [33] È. B. Vinberg. The Two Most Algebraic Algebraic K3 Surfaces. *Math. Ann.* 265 (1983), 1–21.
- [34] A. Weil. The field of definition of a variety. *Am. J. Math.* **78** (1956), 509–524.
- [35] O. Zariski. On the Problem of Existence of Algebraic Functions of Two Variables Possessing a Given Branch Curve. *American Journal of Mathematics* **51** (2) (1929), 305–328.

DEPARTAMENTO DE MATEMÁTICA Y ESTADÍSTICA, UNIVERSIDAD DE LA FRONTERA, CASILLA 54-D,
4780000 TEMUCO, CHILE

Email address: ruben.hidalgo@ufrontera.cl

DEPARTAMENTO DE MATEMÁTICA Y ESTADÍSTICA, UNIVERSIDAD DE LA FRONTERA, CASILLA 54-D,
4780000 TEMUCO, CHILE

Email address: h.hughes01@ufromail.cl

INSTITUTO DE MATEMÁTICA Y FÍSICA, UNIVERSIDAD DE TALCA, 3460000 TALCA, CHILE

Email address: leyton@inst-mat.otalca.cl